We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Recent Trends in Microbiological Decontamination of Aflatoxins in Foodstuffs

Carlos Augusto Fernandes Oliveira, Fernanda Bovo,
Carlos Humberto Corassin,
Alessandra Vincenzi Jager and
Kasa Ravindranadha Reddy

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/51120>

1. Introduction

Nowadays, about 100,000 fungi have already been identified. From these, more than 400 may be considered potentially toxigenic, and about 5% are known to produce toxic compounds or classes of compounds that cause adverse effects in animals and humans in several parts of the world [1]. These compounds, called mycotoxins, are secondary metabolites of low molecular weight produced by mycelia or spores of filamentous fungi [2]. It is suggested that mycotoxin production is generally limited to a relatively small number of mold species, and that toxin may be produced by the whole species or just one specific strain [3]. The more complex the synthesis pathway of a mycotoxin, the lesser the number of mold species that produce it.

The term “mycotoxin” originates from the Greek word “Mykes”, meaning fungus, and from the Latin word “Toxicum”, meaning poison or toxin [2]. Mycotoxins are classified as the most important chronic and noninfectious foodborne risk factor, more important than synthetic contaminants, plant toxins, food additives, and pesticide residues. Both humans and animals may show acute or chronic intoxication caused by mycotoxin ingestion, and the pathological condition that results from this ingestion is called mycotoxicosis [4]. Some factors affect the magnitude of toxicity in humans or animals, including the animal species, mechanism of action, metabolism and defense mechanisms [5].

About 400 types of mycotoxins have been already discovered, and they are generally divided into groups based on structural similarities and most important toxic effects [6]. From all

mycotoxins that have been isolated, aflatoxin is one of the most well-known and widely distributed in foodstuffs, with proven and marked toxic properties. Aflatoxins are predominantly produced by *Aspergillus flavus* and *A. parasiticus*, but may also be produced by other strains, such as *A. nomius*, *A. tamari*, and *A. pseudotamarii* [7]. Contamination of foodstuff with aflatoxigenic fungi may occur at any moment during production, harvesting, processing, transportation, and storage [8]. The most different kinds of foods may be affected, such as corn, peanuts, cotton seeds, rice, pistachio, almonds, chestnuts, Brazil nuts, and pumpkin seeds, as well as other oily seeds, such as sunflower and coconut [9].

Aflatoxins are distributed worldwide. *Aspergillus* species are able to grow in a wide variety of substrates and under different environmental conditions. Toxin formation in agricultural products occurs in hot and humid weather, and in inadequate or deficient storage facilities. The most important factors that influence growth and aflatoxin production are relative humidity, ranging from 88 to 95% in most of the cases [8], and temperature, ranging from 36 to 38 °C for mold growth, and 25 to 27 °C for maximum toxin production [10].

Other factors may also influence aflatoxin production: substrate composition, water activity, pH, atmosphere (concentration of oxygen and carbon dioxide), microbial competition, mechanical damage to the seeds, mold lineage, strain specificity and variation, instability of toxigenic properties, plant stress, insect infestation, and use of fungicides or fertilizers [2, 5, 11]. It is important to remember that aflatoxin contamination is cumulative, and the moment of harvesting and drying, and storage conditions may also play an important role in aflatoxin production [12].

Concerns related to the negative impacts of aflatoxins on health led to the study of strategies to prevent toxin formation in foodstuffs, as well as to eliminate, inactivate or reduce toxin bioavailability in contaminated products [13]. Contamination may be prevented by improved agricultural practices, antifungal agents, genetic engineering, and control of storage conditions [2]. Bioavailability may be reduced by enterosorption, which is done by adding nutritionally inert adsorbent compounds to the diet. These compounds are mycotoxin sequestrants, and prevent the toxin from being absorbed in the gastrointestinal tract of the animals, making its distribution to the target organs impossible [14]. This method has limited practical use, due to the safety of the adsorbent agents used, and the difficulty in applying them to human foods [15]. Elimination or inactivation, that is, decontamination, may be achieved by physical, chemical, and biological methods, which have to present the following characteristics: complete inactivation; destruction or removal of the toxin; no production or toxic residues in foods or no remainders of them; preservation of nutritional value and palatability of the food; destruction of fungal spores and mycelia to prevent production or re-appearance of the toxin; no significant changes in the physical properties of the food; low cost and ease of use [1,11].

Physical methods for mycotoxin decontamination involve procedures such as thermal inactivation, ultraviolet light, ionizing radiation, or extraction with solvents. Chemical methods are based on agents that break mycotoxin structure, such as chlorine treatment (sodium hypochlorite or chlorine gas), oxidizing agents (hydrogen peroxide, ozone and sodium disulfide), or hydrolytic agents (acids, alkalis and ammonia). However, both chemical and

physical methods have disadvantages, either because removal is not efficient, or because of high costs or nutritional losses to the product [16,17]. Biological methods are based on the action of microorganisms on mycotoxins. These microorganisms may be yeasts, filamentous fungi, bacteria, algae, among others, and their mechanisms of action is based on competition by nutrients and space, interactions, and antibiosis, among others [18].

Biodegradation of aflatoxins by microorganisms offers an attractive alternative for the control or elimination of aflatoxins in foods and animal feed, preserving their quality and safety [19]. Besides, their use have a more "natural" appeal, given the ever-growing resistance of the consumer to chemical treatments [1]. Biological decontamination methods are being widely studied and may be a very promising choice, provided they show to be efficient, specific, cost-effective, and are environmentally friendly [20]. Among the types of microorganisms available and that may be used to remove aflatoxins from a contaminated medium, lactic acid bacteria (LAB) and yeasts are the most studied ones, showing the most promising results.

Therefore, the objective of this chapter was to present results of studies on microbiological methods for aflatoxin decontamination, more specifically on the ability of LAB and yeasts to degrade or sequester this mycotoxin.

2. Toxicological Properties of Aflatoxins

Nowadays, there are 18 similar compounds called aflatoxins. However, the most important types in terms of health and medical interest are identified based on their fluorescence under ultraviolet light (B = Blue and G = Green), such as aflatoxin B₁ (AFB₁), B₂ (AFB₂), G₁ (AFG₁) and G₂ (AFG₂). From these compounds, AFB₁ is the most prevalent and toxic one [21]. When AFB₁ is ingested by domestic animals in contaminated feed or foodstuffs, such as by dairy cows, the toxin undergoes liver biotransformation and is converted into aflatoxin M₁ (AFM₁), becoming the hydroxylated form of AFB₁, which is excreted in milk, tissues and biological fluids of these animals [22-24]. It was reported that of all AFB₁ ingested in feed, about 0.3% to 6.2% is transformed in AFM₁ in milk and that there is a linear relationship between the concentration of AFM₁ in milk and the concentration of AFB₁ in contaminated feeds consumed by the animals [25,26].

Chronic exposure to low levels of aflatoxins represents a serious risk to economy, and mainly to health [21]. Economic losses are related to decreased efficiency in industrial or agricultural production, with loss in quality, lower yield, and defective product [27]. It was also reported that in some states of the USA, economic losses to agriculture amount to 100 million dollars [19]. On the other hand, these losses caused by mold contamination and mycotoxins are greater than 1.6 billion dollars in the US, and African feeds lose about 670 billion dollars a year due to barriers to the trade of aflatoxin-contaminated foodstuffs [28].

As for human and animal health, biological effects of aflatoxins may be carcinogenic, mutagenic, teratogenic, hepatotoxic, and immunosuppressive [29]. The International Agency for

Research on Cancer classifies AFB₁ and AFM₁ as Group 1 human carcinogens, even though AFM₁ is about 10 times less carcinogenic than AFB₁ [30]. All these aflatoxin effects are influenced by variations according to the animal species, sex, age, nutritional status, and effects of other chemical products, besides the dose of toxin and the length of exposure of the organism to it [31].

Aflatoxicosis is the poisoning caused by the ingestion of moderate to high levels of aflatoxin in contaminated foods. Acute aflatoxicosis causes quick and progressive jaundice, edema of the limbs, pain, vomiting, necrosis, cirrhosis and, in severe cases, acute liver failure and death, caused by the ingestion of about 10 to 20 mg of aflatoxin in adults. Aflatoxin LD50 shows the following order of toxicity: AFB₁ > AFM₁ > AFG₁ > AFB₂ > AFG₂ [4, 32]. Chronic aflatoxicosis causes cancer, immunosuppression and other pathological conditions, having the liver as the primary target organ [4].

The greatest risk presented by aflatoxins for human beings is chronic exposure causing hepatocellular carcinoma, which may be made worse by hepatitis A virus [5]. It was also reported that aflatoxins were found in the tissues of children affected by Reye syndrome (encephalopathy with serious lesions in liver and kidneys after influenza or chickenpox), and Kwashiorkor (protein-energy malnutrition). Aflatoxicosis is considered, then, a contributing factor to these diseases.

AFB₁ is metabolized in the liver by the cytochrome P450 system, generating its most carcinogenic metabolite, AFB₁-8,9-epoxide (AFBO), or other less mutagenic forms, such as AFM₁, Q₁ or P₁. There are several pathways for AFBO after it is metabolized, with one of them leading to cancer, another to toxicity and another one, to excretion. AFBO exo-form easily binds to cell macromolecules, including genetic material such as DNA proteins, producing adducts. Formation of these DNA adducts leads to genetic mutations and cancer, and their excretion in the urine of infected people is not only a proof that humans have the necessary biochemical pathways for carcinogenesis, but also offers a reliable biomarker for AFB₁ exposure [24].

Potential risk to human health caused by aflatoxins has led to surveillance programs for the toxin in different raw materials, as well as regulations determined by almost every country in the world [9]. A study carried out by the Food and Agriculture Organization of the United Nations (FAO) in 2002 pointed out that about 100 countries had specific regulations for the presence of aflatoxin in foods, dairy products and animal feed, and that the total population of these countries amounted to 90% of the world population. The same study showed that regulations for aflatoxin are getting more diverse and detailed, including sampling methods and methods of analysis [33].

In countries where a regulation for aflatoxin exists, tolerance levels for the total aflatoxin (sum of aflatoxins B₁, B₂, G₁ and G₂) ranges from 1 to 35 µg/kg for foods, with an average of 10 µg/kg; and from zero to 50 µg/kg for animal feed, with an average of 20 µg/kg. For AFM₁ in milk, tolerance levels are between 0.05 and 0.5 µg/kg, with most countries adopting a threshold of 0.05 µg/kg [10].

3. Decontamination of Aflatoxins by Lactic Acid Bacteria

LAB is a large group of genetically different bacteria that, besides producing lactic acid as the main product of their metabolism, have similar characteristics: they are all gram-positive, non-sporoformers, non-motile, and catalase, and oxidase negative. They are, therefore, aerotolerant anaerobes. Besides, they mandatorily ferment sugars and tend to be nutritionally fastidious, frequently requiring specific amino acids and B-complex vitamins as growth factors [34]. Several LAB genera, such as *Lactobacillus*, *Bifidobacterium* and *Lactococcus* are known for they ability to act as preserving agents in fermented foods, such as vegetables, cereals, dairy and meat products, actively inhibiting spoilage and growth of pathogenic bacteria, besides increasing shelf life and sensory properties of these foods [23].

Fermentation enables longer shelf life and improves sensory and nutritional properties of the product, as sugar fermentation lowers pH and inhibits growth of spoilage and pathogenic microorganisms. Fermentation is also responsible for other reactions, such as proteins hydrolysis, improving texture and flavor; synthesis of aromatic components and texturizers, affecting the consistency of the product; and production of inhibitory components [35,36]. This inhibition is, in part, caused by the final products of fermentation, such as lactic acid, diacetyl, acetaldehyde and acetic acid, which may accumulate in inhibitory concentrations in certain foods and drinks. In other cases, inhibition may also be caused by secondary by-products of metabolism, such as hydrogen peroxide or bacteriocins [37].

Therefore, two aspects may be considered when LAB are used: fermentation and antibiosis ability. In the first case, the starter culture added to the food acts on the substrate, causing advantages to the food. In the second case, the starter culture has to inhibit the development of undesirable microorganisms that may spoil the product or be hazardous to human health. In reference [38], authors state that one of the effects that were identified in LAB was protection against toxins found in foods, such as heterocyclic amines, polycyclic aromatic hydrocarbons, reactive oxygen species, and mycotoxins. In the latter case, studies have demonstrated that LAB have the ability to inhibit aflatoxin biosynthesis, or that they have the ability to remove mycotoxins from the medium, reducing their effects.

It should be emphasized that with increased interest in probiotic food production all over the world, selection of LAB cultures with probiotic characteristics and greater ability to remove mycotoxins may help to reduce risk of exposure to these toxins in foodstuffs, which is a very promising line of research in mycotoxicology. Yeast and LAB strains have great ability to remove mycotoxins, and may be used as part of starter cultures in the fermentation of foods and drinks [39]. These microorganisms have, thus, ability to ferment and decontaminate the medium, and purified components of these strains may be used in small amounts as food additives without compromising the characteristics of the final product.

One of the first studies in this area was carried out in the 1960s, when these authors evaluated the ability of about 1,000 types of microorganisms to degrade aflatoxins [40]. Yeasts, filamentous fungi, bacteria, actinomycetes, algae, and fungal spores were among the organisms studied. From these, only the bacterium *Flavobacterium aurantiacum* B-184 (known today as *Nocardia corynebacterioides*) was able to irreversibly remove aflatoxins from the solution.

After this study, many others followed. However, the most significant ones started to appear after the 1990s. Table 1 presents the most relevant studies carried out with bacteria for aflatoxin decontamination. The action of 7 different types of bacteria on AFB₁ was evaluated and it was found that some strains of *Lactobacillus* (*L. rhamnosus* GG and *L. rhamnosus* LC-705) were able to efficiently remove most mycotoxin from the medium, up to about 80% [17]. In reference [27] authors analyzed 9 strains of *Lactobacillus* and achieved the same result, that *L. rhamnosus* GG and *L. rhamnosus* LC-705 were the most efficient strains in removing AFB₁, with removal rates of 78.9% and 76.5%, respectively. Fifteen types of LAB, among them *Lactobacillus* and *Lactococcus*, and 5 types of bifidobacteria, were studied and it was observed that removal of AFB₁ ranged from 5.6% to 59.7% [23]. Strains of *Lactobacillus amylovorus* (CSCC 5160 and CSCC 5197) and *L. rhamnosus* LC 1/3 showed the best results: 59.7%, 57.8%, and 54.6%, respectively. It was also observed that different strains of bifidobacteria removed from 37% to 46% AFB₁, and that *Staphylococcus aureus* and *Escherichia coli* removed 46% and 37%, respectively [22]. It may be observed that among a given genus, and even a given species, not all the strains show equivalent toxin removal rates. On the contrary, the ability to remove aflatoxin is a characteristic of specific lineages, and efficiency varies widely [41].

Most assays on aflatoxin removal in the studies cited above were carried out in phosphate-buffered saline (PBS). In reference [42], besides testing the ability of 27 strains of *Lactococcus* spp. and 15 strains of *Streptococcus* spp. isolated from yogurt, raw milk, and Karish cheese to remove AFB₁ in buffered solution, observed that *Lactococcus L. lactis* and *Streptococcus thermophilus* presented the greatest rates of toxin removal (54.85% and 81.0%, respectively). They also tested the ability of viable and non-viable cells to remove AFB₁ in different vegetable oils, and observed that viable *L. lactis* cells removed from 71% to 86.7% AFB₁, whereas non-viable cells removed 100% of the toxin in all the oils. Moreover, viable *S. thermophilus* cells removed from 66.5% to 91.5% of the toxin, and non-viable ones, from 81.7% to 96.8%.

AFB₁ was added to yogurt and acidified milk in concentrations ranging from 1,000 to 1,400 g/kg, and a reduction of AFB₁ in yogurt (pH 4.0), ranging from 97.8% to 90% was obtained [43]. Maximum decrease in AFB₁ was observed during milk fermentation. As for milk acidified with citric, lactic, and acetic acid (pH 4.0) AFB₁ reduction (concentration of 1,000 µg/Kg) was 90%, 84% and 73%, respectively. The ability of probiotic bacteria (*L. paracasei*, *L. casei*, *L. brevis* and *L. plantarum*) and the yeast *Saccharomyces cerevisiae* to remove a sum of aflatoxins (B₁, B₂, G₁ and G₂) during fermentation of dough made up of 50% barley flour, 45% wheat flour, and 5% corn flour was evaluated [44]. They observed that after 6 hours of fermentation, the amount of aflatoxin had decreased 18% and 33% for dough added of 4 and 40 µg of aflatoxin, respectively, and after 24 hours, the amount of aflatoxin decreased 27% and 50%, respectively.

Toxin polarity has an important role in the binding mechanism. The percentage of aflatoxin removed by LAB decreases in the following order: AFB₁ > AFB₂ > AFG₁ > AFG₂. This observation correlates with the decrease in the polarity of these toxins, and is consistent with hydrophobic reactions, which may also have a role in the binding mechanism [45]. AFM₁ is less efficiently removed than AFB₁. However, scientific literature has few studies on the ability of LAB to remove AFM₁.

In reference [46], authors examined the ability of 4 strains of *Lactobacillus* spp. and 2 strains of *Bifidobacterium* spp. to remove AFM₁ in PBS and reconstituted skim milk. In PBS, viable cells of 6 strains were able to remove from 10.22 to 26.65% AFM₁ in solution, depending on the level of contamination and the length of incubation, whereas non-viable cells removed from 14.04 to 28.97% of the toxin. In reconstituted skim milk incubated for 4 hours, 7.85 to 25.94% AFM₁ were removed by viable cells, and 12.85 to 27.31% for cells rendered non-viable by heat treatment. These researchers concluded that the removal process was fast, with no differences between 0, 4, and 24 hours of contact, different from what was observed in [47] for strains of *Lactobacillus* spp., *Lactobacillus* spp. and *Bifidobacterium* spp., which showed removal rates ranging from 0 to 14.6% after 24 hours of contact, and from 4.5 to 73.1% after 96 hours of contact.

The ability of *L. rhamnosus* GG to remove AFM₁ from reconstituted skim and whole milk was investigated and it was observed rates of 18.8% and 26.0%, respectively [29]. The authors concluded that the decrease in removal efficiency may be explained by the fact that AFM₁ is possibly not accessible in milk, that is, it is associated with casein, and the interference of proteins in toxin removal may be the greatest responsible factor for the difference between skim milk and whole milk (approximately 10% lower), once powdered skim milk used in the study contained 37g of protein / 100 g, whereas protein content in powdered whole milk was 25g /100g. In the same study, AFM₁ removal in buffered solution (50.7%) was compared with AFB₁ removal by the same bacterial strain in the same solution (75.3%). It was concluded that AFM₁ removal was less effective possibly due to the presence of an -OH group in the molecule, increasing its polarity and making it less hydrophilic, what increases the tendency of the molecule to be retained in aqueous solutions.

Some physical, chemical, and enzymatic treatments may increase the ability of LAB to bind to aflatoxin in the medium. In reference [48] authors studied the ability of *L. rhamnosus* GG to bind to AFB₁, observing little difference between aflatoxin removal by heat-treated and acid-treated cells (85% and 91%, respectively), compared with viable bacterial cells (86%). The use of physical and chemical treatments (chloric acid, and heat treatment in autoclave or boiling at 100 °C) on *L. rhamnosus* GG and LC-705 caused a significant increase in AFB₁ removal, showing that metabolic degradation caused by viable bacterial cells may be ruled out as a possible mechanism of action [15-17].

Comparing the ability of viable and heat-treated bifidobacteria cells, it was observed that viable cells removed 4 to 56% AFB₁ from the medium, whereas non-viable cells removed 12 to 82% [23]. Evaluating the influence of the inactivation treatment on the ability of 4 types of *Lactobacillus* spp. to remove AFB₁, it was observed that acid treatment (58.6 to 87.0%) and heat treatment (33.5 to 71.9%) increased the ability to remove the toxin, compared with viable cells in PBS (16.3 to 56.6%) [49]. On the other hand, alkali treatment (8.3 to 27.4%) and ethanol treatment (15.9 to 46.5%) decreased the amount of aflatoxin removed from the medium.

Removal of AFM₁ with 8 LAB strains showed that heat-treated cells bound more efficiently (25.5 to 61.5%) to the toxin than viable bacterial cells (18.1 to 53.8%) [29]. In reference [50] it was observed that heat-treated cells removed greater percentages of AFM₁ (12.4% to 45.7%) in PBS compared with viable cells (5.6% to 33.5%), with no significant differences between 15

minutes or 24 hours of contact. Similar results were found in [51], because viable cells of *Lactobacillus delbrueckii* spp. *bulgaricus* CH-2 removed 29.42% AFM₁ in PBS after 4 hours of contact at 37 °C. These authors also analyzed the ability of *Streptococcus thermophilus* ST-36, observing that 18.70% AFM₁ was removed from the medium. Until today, only one bacterium, *Flavobacterium aurantiacum* NRRL B-184, was able to remove 100% of AFM₁ from contaminated liquid medium, at a cell concentration of 5×10^{10} CFU/mL and 4 hours of contact [52].

In [53] authors observed that *B. subtilis* UTBSP1 presented significant removal of AFB₁ from a medium contaminated with 2.5 µg/g (52.67% and 80.53%, after 24 and 48 hours, respectively). After 72 and 96 hours, there was no significant increase in the amount of toxin removed from the medium. Strains of *B. subtilis* were analyzed and it was concluded that strain ANSB060 was the one that best removed AFB₁, AFM₁, and AFG₁ from the medium (81.5%, 60%, and 80.7%, respectively) [54]. Results of this study also demonstrated that aflatoxin degradation is mainly observed in the supernatant culture, compared with cells or cell extracts. Besides, in assays that simulated the gastrointestinal environment (pH 2.0, and 0.3% of biliary salts), viable cells of the same strain were able to survive for 24 hours of incubation, and presented antimicrobial activity against *E. coli*, *S. typhimurium*, and *S. aureus*.

These examples show that both viable and non-viable cells are able to remove aflatoxin from aqueous solutions. As non-viable cells are also able to remove the toxin, it is supposed that cells are physically bound to the toxin, that is, components of the bacterial cell wall adhere to it, mainly polysaccharides and peptidoglycans, taking into account the possibility of a covalent bond or degradation caused by bacterial metabolism [1, 55, 56].

Both polysaccharides and peptidoglycans of the bacterial cell wall may be extremely affected by heat and acid treatment, once heat may denature proteins or form Maillard reaction products. Besides, acid treatment may break glycosidic bonds of polysaccharides, releasing monomers that may be further broken into aldehydes, also degrading proteins to smaller components, such as peptides and amino acids. Thus, acid treatment may break the peptidoglycan structure, compromising its structural integrity, that is, decreasing the thickness of this layer, reducing cross links and increasing the size of the pores. These changes caused by the treatments cited above enable AFB₁ to bind to the bacterial cell wall and to the components of the plasmatic membrane that were not available when the bacterial cell was intact [27].

In reference [57] authors explained that the integrity of the bacterial cell wall is important in the process of toxin removal by both viable and non-viable cells. In their study of AFB₁, they observed that both the bacterial cell wall and its purified fragments were able to remove aflatoxin from the medium. However, when the cell wall was lost or destroyed (totally or partially) by enzymatic treatment, there was a significant decrease in the ability to remove the toxin. It was observed, using atomic force microscopy, that the bond between AFB₁ and *Lactobacillus casei* Shirota produced structural changes that modified the surface of the bacterial cell [58]. Before the toxin was bound to it, the surface was well-defined, smooth and homogenous, and after AFB₁ adsorption, there were changes in shape. These changes were probably caused by the bond between the toxin and the surface of the cell wall, which became very irregular and rough, with undefined edges. The authors suggest that changes in the shape of teichoic acids are responsible for these alterations, once these molecules are

found inside the cell wall in such a way that they produce no differences in the texture of the surface before the toxin was bound to it.

The ability of *L. rhamnosus* GG to bind to AFB₁ was studied, observing that the addition of urea - an anti-hydrophobic agent - to the medium, significantly decreased removal of the toxin by non-viable cells, from 85-91% to 50-60%, showing that hydrophobic interactions have a relevant role in the process [48]. Besides, addition of different concentrations of NaCl and CaCl₂ (from 0.01 to 1 M), and pH variations from 2.5 to 8.5 had practically no effect on AFB₁ removal by the bacterium, suggesting that hydrogen bonds and electrostatic interactions are not important in this process.

In the use of pronase E, lipase and periodate, treatment with periodate led to significant reduction in the ability to remove the toxin, both by viable and non-viable cells, once it oxidizes the -OH cis groups in aldehyde and carboxylic acid groups, suggesting that the bonds involve predominantly bacterial polysaccharides. Treatment with pronase E caused the same significant reduction in AFB₁ removal, evidencing that proteins may also be involved in the process. Thus, the fact that pronase E and periodate both have a significant reduction on AFB₁ removal indicates that binding sites are made of protein. Treatment with lipase, on its turn, did not cause any significant reduction in AFB₁, showing that lipids, such as lyso-teichoic acid probably do not have a role in the process. Although the treatments decreased AFB₁ removal, it was still substantial in all cases, possibly showing the involvement of multiple components in the bond with mycotoxin [48].

However, not only the type of bacterial strain and the inactivation treatment used may influence formation and stability of the LAB/aflatoxin complex, but also of other factors, such as bacterial counts, specificity of the bacteria, pH, incubation temperature, addition of nutrients, and the solvents used, among others [23, 27, 48].

As for the number of bacterial cells in the medium, it has been concluded that there was a significant decrease in the amount of AFB₁ removed when cell counts changed from 10⁷ CFU/mL (0 to 5.02%) to 10⁸ CFU/mL (10.22 to 26.65%), indicating that bacterial counts are critical factors in the removal of AFB₁ by LAB [46]. In reference [59] authors observed that no less than 5 × 10⁹ CFU/mL of *Lactobacillus acidophilus* or *Bifidobacterium longum* are necessary to remove only 13% AFB₁ in about one hour.

In reference [17] authors reported that, for *Lactobacillus rhamnosus* (strains GG and LC705), minimum counts of 2 × 10⁹ CFU/mL were required to remove 50% AFB₁, and greater removal rates were obtained when LAB concentration was increased to 10¹⁰ CFU/mL. In this same study, the authors observed that the process depended on the temperature, once the efficiency in aflatoxin removal was greater at 37 °C than at 4 and 25 °C. Besides, the authors observed that Gram-positive bacteria are better aflatoxin sequestrants than Gram-negative bacteria, with removal rates of 80% and 20%, respectively, suggesting the ability to remove the toxin depends on the structure of the cell wall. It has also been stated that aflatoxin concentration in the medium also influences adsorption rates, leading to the conclusion that the greater its concentration in the medium, the greater the removal rate, both for viable and non-viable cells [60].

Assays with AFB₁ and *L. rhamnosus* GG and LC-705 at different incubation temperatures was also carried out, but it was not observed significant differences in the stability of the LAB/AFB₁ complex formed in the temperatures range between 4 °C and 37 °C [27]. When pH of the medium was changed from 2 to 10, a range that includes the pH switch in the gastrointestinal tract, only 10% AFB₁ removed was released back into the solution, different from what happened when organic solvents were used. In this case, almost all AFB₁ that was removed by the bacterial strains was released back into the medium, providing extra evidence that the process is based on a non-covalent bond. In this study, the release efficiency by solvents presented the following order: methanol < acetonitrile = benzene < chloroform, which does not coincide with the order of decreasing polarity. This may be explained by the fact that AFB₁ hydrophobicity is similar to that of the chloroform molecule. These results show once more that hydrophobic interactions have an important role in the binding mechanism between LAB and the toxin.

The effect of washing on the stability of the LAB/aflatoxin complex was analyzed [47]. They observed that after the first washing of bacterial pellets with PBS, the proportion of AFM₁ released by the bacteria was 87.3% for *Lactobacillus* spp. strains; 85.7% for *Lactococcus* spp. strains, and 85.7% for strains of *Bifidobacterium* spp. They also observed that after the third washing, practically all bacteria had released adsorbed AFM₁ back into the medium (92.0 a 100%). In reference [46] they concluded that AFM₁ removal by bacteria was reversible, and that small amounts of toxin were released back to the PBS solution (5.62 to 8.54%). This finding is consistent with those observations of reference [27], who reported that *L. rhamnosus* GG, *L. rhamnosus* LC-705, and *Lactobacillus casei* Shirota released, respectively, 3.7%, 3.0% and 2.4% AFB₁ back into the solution. Differently, in [23] authors showed that release of AFB₁ back into the solution in the first washing was 48.6%, 30.7% and 26.5% for *L. amylovorus* (strains CSCC 5160 and CSCC 5197) and *L. rhamnosus* Lc 1/3, respectively. After 5 washings, AFB₁ adsorbed by *L. amylovorus* CSCC 5160 was almost completely released (94.4%), whereas *L. amylovorus* CSCC 5197 and *L. rhamnosus* Lc 1/3 retained, respectively, only 17.4% and 32.2% AFB₁ found in the original solution.

Thus, the LAB/aflatoxin complex seems to be unstable, once part of the aflatoxin, both for AFB₁ and AFM₁, is released from the complex after washing, and gradually returns to the aqueous solution. Therefore, the greater the number of washings, the greater the amount of aflatoxin released back into the solution. This shows that the bond is not a strong one, suggesting it is a weak non-covalent bond and an association with hydrophobic sites on the surface of the bacteria [23, 48].

Different from this hypothesis, in reference [61], performing the same washings on a complex between *Flavobacterium aurantiacum* and AFB₁, authors observed that aflatoxin was not released into the aqueous solution. Analyzing the stability of the complex formed between AFB₁ and 8 strains of *Lactobacillus casei* after the washings, it was demonstrated that the amount of aflatoxin released ranged from practically zero and 9.2% [13]. Possible explanations for this variation in aflatoxin release include the differences in binding sites found in the different strains, or more probably, that these binding sites are similar, but that they present minimal differences depending on the strain. Authors explained that lower rate of toxin release into

the medium after the washings may be attributed to the interactions between aflatoxin molecules retained on the cell wall of a bacterium and molecules retained on the cell wall of the adjacent bacterium, forming a kind of reticulated matrix that prevents aflatoxin release. It has also been suggested that the greater the number of molecules that are removed by the bacterial cells, the longer these molecules remain adsorbed on the cell surface [60].

The stability of the LAB/aflatoxin complex in a wide range of pH is an important factor in the use of these microorganisms to remove aflatoxin from foods, once gastric release of the toxin would have negative health implications. Therefore, the complex formed has to resist environmental stress caused by the gastrointestinal tract, such as low pH and presence of bile. When the influence of the presence of bile on the LAB/aflatoxin complex was analyzed, it was observed that *Lactobacillus casei* removed more AFB₁ when exposed to bile, suggesting that this exposure causes changes in the structure and composition of the bacterial cell wall, probably inducing the formation of new binding sites for aflatoxin, or increasing the size of the sites available [13].

The ability of *L. rhamnosus* (strains GG and LC705) and *Propionibacterium freudenreichii* spp. *shermanii* JS to remove AFB₁ from intestinal liquid medium extracted from the duodenum of chicks was investigated, and it was observed that AFB₁ concentration was reduced in 54% in only 1 minute in the presence of *L. rhamnosus* GG, whereas it was reduced in only 44% in the presence of *L. rhamnosus* LC705, and 36% in the presence of *P. freudenreichii* spp. *shermanii* JS [62]. The authors observed that the accumulation of AFB₁ in the intestinal tissue was reduced in 74%, 63%, and 37%, respectively, for *L. rhamnosus* (strains GG and LC705) and *P. freudenreichii* spp. *shermanii* JS, showing that these bacteria may affect aflatoxin bioavailability and be used to reduce its toxicity to humans and animals.

Rats treated with feed added of aflatoxin (3 mg/kg of feed) presented a significant decrease in the feed intake compared with the control group, different from the animals fed diets containing *Lactobacillus casei* and *Lactobacillus reuteri* (10 mL/kg of feed, with 1×10^{11} CFU/mL) and aflatoxin [63]. The second group did not show reduced feed intake. Consequently, animals treated with the diet containing only aflatoxin presented lower body weight, significant increase in serum levels of transaminase, alkaline phosphatase, cholesterol, triglycerides, total lipids, creatinine, uric acid, and nitric oxide; and in lipid peroxidation in the liver and kidneys, followed by a significant decrease in total antioxidant capacity. Treatment with bacteria was able to induce a significant improvement in all biochemical parameters and in the histological condition of the liver, with *L. reuteri* being more efficient than *L. casei*.

In Egypt, a pilot study investigated the effect of the addition of *L. rhamnosus* LC-705 and *P. freudenreichii* spp. *shermanii* JS in human diet on the levels of aflatoxin in feces samples. It was observed that from 11 of 20 volunteers, AFB₁ ranged from 1.8 to 6 µg AFB₁/kg feces, and after two weeks of supplementation with probiotic bacteria, there was a significant reduction in the excretion rate, showing that these strains have the ability to influence the concentration of AFB₁ in feces [64].

Microorganism	AF	Bound (%)	Conditions	Ref.
<i>L. rhamnosus</i> GG	B1			
Viable cells	5 µg/mL	78.4	2 x 10 ¹⁰ cfu/mL, 0h, 37 °C, PBS	[17]
Freeze-dried cells		65	4h, 37 °C, PBS	
Heat-treated cells		81	4h, 37 °C, PBS	
<i>L. rhamnosus</i> LC-705				
Viable cells		78.8	2 x 10 ¹⁰ cfu/mL, 0h, 37 °C, PBS	
Freeze-dried cells		50	4h, 37 °C, PBS	
Heat-treated cells		82	4h, 37 °C, PBS	
<i>L. gasseri</i>		58.1	2 x 10 ¹⁰ cfu/mL, 0h, 37 °C, PBS	
<i>L. acidophilus</i>		67.4	7 x 10 ⁹ cfu/mL, 0h, 37 °C, PBS	
<i>L. casei</i> Shirota		33.2	1 x 10 ¹⁰ cfu/mL, 0h, 37 °C, PBS	
<i>E. coli</i>		16.3	5 x 10 ¹⁰ cfu/mL, 0h, 37 °C, PBS	
<i>L. paracasei</i> , <i>L. casei</i> , <i>L. brevis</i> , <i>L. plantarum</i> and <i>S. cerevisiae</i>	B ₁ , B ₂ , G ₁ , G ₂ 4 or 40 µg/kg	18-33 27-50	6h, 37 °C, barley flour (50%), wheat flour (45%) and corn flour (5%) mixed with water in 1:1.5 proportion 24h, 37 °C, barley flour (50%), wheat flour (45%) and corn flour (5%) mixed with water in 1:1.5 proportion	[44]
<i>Lc. lactis</i> ssp. <i>cremoris</i>	B ₁	5.6	1 x 10 ¹⁰ cfu/mL, 24h, 37 °C, PBS	[23]
<i>Lactobacillus delbrueckii</i>	5 µg/mL	17.3		
<i>Lb. acidophilus</i>		18.2		
<i>Lb. rhamnosus</i>		22.7		
<i>Lb. plantarum</i>		28.4		
<i>Lc. lactis</i> ssp. <i>lactis</i>		31.6		
<i>Bifidobacterium lactis</i>		18.0		
<i>Lb. helveticus</i>		34.2		
<i>Lc. lactis</i> ssp. <i>cremoris</i>		41.1		
<i>Lb. rhamnosus</i> Lc		54.6		
<i>Lb. acidophilus</i>		20.7		
<i>Lb. fermentum</i>		22.6		
<i>Lb. johnsonii</i>		30.1		
<i>Lb. rhamnosus</i>		33.1		
<i>Lb. amylovorus</i>		57.8		
<i>Lb. amylovorus</i>		59.7		
<i>Bb. lactis</i>		34.7		
<i>Bb. longum</i>		37.5		
<i>Bb. animalis</i>		45.7		
<i>Bb. lactis</i>		48.7		

Microorganism	AF	Bound (%)	Conditions	Ref.
<i>L. rhamnosus</i> GG	B1			[27]
Viable cells	5 µg/mL	78.9	1 x 10 ¹⁰ cfu/mL, 1h, 37 °C, PBS	
Heat-treated cells		84.1		
Acid-treated cells		86.7		
<i>L. rhamnosus</i> LC-705				
Viable cells		76.5	1 x 10 ¹⁰ cfu/mL, 1h, 37°C, PBS	
Heat-treated cells		87.8		
Acid-treated cells		88.3		
<i>L. acidophilus</i> LC1				
Viable cells		59.7	1 x 10 ¹⁰ cfu/mL, 1h, 37 °C, PBS	
Heat-treated cells		74.7		
Acid-treated cells		84.2		
<i>L. lactis</i> subsp. <i>lactis</i>				
Viable cells		59.0	1 x 10 ¹⁰ cfu/mL, 1h, 37 °C, PBS	
Heat-treated cells		58.1		
Acid-treated cells		69.5		
<i>L. acidophilus</i> ATCC 4356				
Viable cells		48.3	1 x 10 ¹⁰ cfu/mL, 1h, 37 °C, PBS	
Heat-treated cells		69.7		
Acid-treated cells		81.3		
<i>L. plantarum</i>				
Viable cells		29.9	1 x 10 ¹⁰ cfu/mL, 1h, 37 °C, PBS	
Heat-treated cells		35.5		
Acid-treated cells		62.7		
<i>L. casei</i> Shirota				
Viable cells		21.8	1 x 10 ¹⁰ cfu/mL, 1h, 37 °C, PBS	
Heat-treated cells		41.5		
Acid-treated cells		32.3		
<i>L. delbrueckii</i> subsp. <i>bulgaricus</i>			1 x 10 ¹⁰ cfu/mL, 1h, 37 °C, PBS	
Viable cells		15.6		
Heat-treated cells		33.7		
Acid-treated cells		75.8		
<i>L. helveticus</i>				
Viable cells		17.5	1 x 10 ¹⁰ cfu/mL, 1h, 37 °C, PBS	
Heat-treated cells		29.8		
Acid-treated cells		58.1		
<i>P. freudenreichii</i> subsp. <i>shermanii</i> JS				
Viable cells		22.3	1 x 10 ¹⁰ cfu/mL, 1h, 37 °C, PBS	
Heat-treated cells		67.3		

Microorganism	AF	Bound (%)	Conditions	Ref.
Acid-treated cells		82.5		
<i>Lc. lactis</i> subsp. <i>cremoris</i>				
Viable cells		26.9	1 x 10 ¹⁰ cfu/mL, 1h, 37 °C, PBS	
Heat-treated cells		40.1		
Acid-treated cells		43.7		
<i>S. thermophilus</i>				
Viable cells		32.7	1 x 10 ¹⁰ cfu/mL, 1h, 37 °C, PBS	
Heat-treated cells		42.0		
Acid-treated cells		63.8		
<i>E. Coli</i>	B1	37	30 min, 37 °C, PBS	[22]
<i>L. rhamnosus</i> GG	2 µg/mL	37		
<i>S. aureus</i>		46		
<i>Bifidobacterium</i> sp. Bf6		25		
<i>B. adolescentis</i> 14		31		
<i>B. bifidum</i> BGN4		46		
<i>Bifidobacterium</i> sp. CH4		37		
<i>B. longum</i> JR20		37		
<i>Bifidobacterium</i> sp. JO3		41		
<i>Lc. lactis</i>	B1		10 ⁷ -10 ⁸ cfu/mL,30 min, 37 °C, in:	[42]
Living cells	0.5 µg/mL	54.8	PBS	
		86.7	maize oil	
		82.3	sunflower oil	
		71.0	soybean oil	
Dead cells by boiling		81.0	PBS	
		100	maize, sunflower or soybean oil	
Dead cells by autoclaving		80.0	PBS	
<i>S. thermophilus</i>				
Living cells		81.0	PBS	
		91.5	maize oil	
		90.7	sunflower oil	
		66.5	soybean oil	
Dead cells by boiling		100.0	PBS	
		96.8	maize oil	
		81.7	sunflower oil	
		96.0	soybean oil	
Dead cells by autoclaving		83.0	PBS	

Microorganism	AF	Bound (%)	Conditions	Ref.
Yoghurt Culture	B1			[43]
	0.6mg/kg	97	42 °C/3h, pH 4.0, overnight, milk	
	1 mg/kg	91		
	1.4mg/kg	90		
	1 mg/kg	90	milk acidified with citric acid	
		84	milk acidified with latic acid	
		73	milk acidified with acetic acid	
<i>L. acidophilus</i> NCC12	M1			[46]
Living cells	5, 10 and 20	14.9-20.2 10 ⁸ cfu/mL	0, 4, 24 h, 37 °C, PBS	
		14.4-15.4 10 ⁸ cfu/mL	4h, 37 °C, milk	
Heated cells	ng/mL	17.0-24.9	0, 4, 24 h, 37 °C, PBS	
		16.6-19.0	4h, 37 °C, milk	
<i>L. acidophilus</i> NCC36				
Living cells		20.4-25.3 10 ⁸ cfu/mL	0, 4, 24 h, 37 °C, PBS	
		21.8-22.7 10 ⁸ cfu/mL	4h, 37 °C, milk	
Heated cells		22.1-26.8	0, 4, 24 h, 37 °C, PBS	
		23.7-25.1	4h, 37 °C, milk	
<i>L. acidophilus</i> NCC68				
Living cells		10.2-16.0 10 ⁸ cfu/mL	0, 4, 24 h, 37 °C, PBS	
		7.8-10.5 10 ⁸ cfu/mL	4h, 37 °C, milk	
Heated cells		14.0-21.8	0, 4, 24 h, 37 °C, PBS	
		12.8-15.9	4h, 37 °C, milk	
<i>B. bifidum</i> Bb13				
Living cells		23.5-26.6 10 ⁸ cfu/mL	0, 4, 24 h, 37 °C, PBS	
		24.0-25.9 10 ⁸ cfu/mL	4h, 37 °C, milk	
Heated cells		24.3-28.9	0, 4, 24 h, 37 °C, PBS	
		25.4-27.4	4h, 37 °C, milk	
<i>B. bifidum</i> NCC 381				
Living cells		16.6-22.1 10 ⁸ cfu/mL	0, 4, 24 h, 37 °C, PBS	
		15.5-18.3 10 ⁸ cfu/mL	4h, 37 °C, milk	
Heated cells		17.4-23.5	0, 4 and 24 h, 37 °C, PBS	
		17.1-22.2	4h, 37 °C, milk	
<i>L. rhamnosus</i>				
Living cells		20.1-24.0 10 ⁸ cfu/mL	0, 4, 24 h, 37 °C, PBS	
		20.4-22.2 10 ⁸ cfu/mL	4h, 37 °C, milk	
Heated cells		23.4-27.8	0, 4 and 24 h, 37 °C, PBS	
		22.9-26.3	4h, 37 °C, milk	
<i>Lactobacillus</i> strains	AFM1	9.4-73.1	96 h, 37 °C, PBS	[47]

Microorganism	AF	Bound (%)	Conditions	Ref.
<i>Lactococcus</i> strains		4.5-38.3	96 h, 37 °C , PBS	
<i>Bifidobacterium</i> strains		7.8-41.6	96 h, 37 °C , PBS	
<i>L. plantarum</i>		73	96 h, 37 °C , PBS	
<i>B. adolescentes</i>		41.6	96 h, 37 °C , PBS	
<i>Lactobacillus</i> strains		64-80.5	96 h, 37 °C ,milk	
<i>Lactococcus</i> strains		46.0-68.5	96 h, 37 °C , milk	
<i>Bifidobacterium</i> strains		67.0-72.5	96 h, 37 °C , milk	
<i>L. bulgaricus</i>		80.5	96 h, 37 °C , milk	
<i>B. adolescentes</i>		73	96 h, 37 °C , milk	
<i>L. rhamnosus</i> strain GG (pre-cultured)	M1 0.15		5.3 x 10 ⁸ , 15 - 16h, , 37 °C, in:	[29]
Viable cells	µg/ml	50.7	PBS	
		18.8	skim milk	
		26.0	full cream milk	
Heat-killed cells		57.8	PBS	
		26.6	skim milk	
		36.6	full cream milk	
<i>L. rhamnosus</i> strain LC-705 (pre-cultured)				
Viable cells		46.3	PBS	
		69.6	skim milk	
		27.4	full cream milk	
Heat-killed cells		51.6	PBS	
		63.6	skim milk	
		30.1	full cream milk	
<i>L. rhamnosus</i> strain GG (lyophilized)				
Viable cells		53.8	1.0 x 10 ¹⁰ , 15-16h, 37 °C, PBS	
Heat-killed cells		56.2		
<i>L. rhamnosus</i> strain LC-705 (lyophilized)				
Viable cells		45.7	1.0 x 10 ¹⁰ , 15-16h, 37 °C, PBS	
Heat-killed cells		57.4		
<i>L. lactis</i> ssp. <i>cremoris</i> strain ARH74				
Viable cells		40.4	2.9 x 10 ⁸ , 15-16h, , 37 °C, PBS	

Microorganism	AF	Bound (%)	Conditions	Ref.
Heat-killed cells <i>L. gasseri</i> (ATCC 33323)		38.9		
Viable cells		30.8	3.9 x 10 ⁸ , 15-16h, , 37 °C, PBS	
Heat-killed cells <i>L. acidophilus</i> strain LA1		61,5		
Viable cells		18,3	1.7 x 10 ⁹ , 15-16h, , 37 °C, PBS	
Heat-killed cells <i>L. rhamnosus</i> strain 1/3		25,5		
Viable cells		18,1	3.9 x 10 ⁸ , 15-16h, , 37 °C, PBS	
Heat-killed cells		39,9		
<i>L. rhamnosus</i> strain GG	B ₁			[48]
Pre-treatment:	5 µg/mL			
Pronase E				
Viable cells		66	1h, 37 °C, 5% CO ₂ , PBS	
Heat-treated cells		72	Boiled for 1h, PBS	
Acid-treated cells		85	2 mol/L HCl, 1h, 37 °C, 5% CO ₂	
Lipase				
Viable cells		76	1h, 37 °C, 5% CO ₂ , PBS	
Heat-treated cells		74	Boiled for 1h, PBS	
Acid-treated cells		89	2 mol/L HCl, 1h, 37 °C, 5% CO ₂	
Phosphate Buffer				
Viable cells		86	1h, 37 °C, 5% CO ₂ , PBS	
Heat-treated cells		85	Boiled for 1h, PBS	
Acid-treated cells		91	2 mol/L HCl, 1h, 37 °C, 5% CO ₂	
m-Periodater				
Viable cells		60	1h, 37 °C, 5% CO ₂ , PBS	
Heat-treated cells		49	Boiled for 1h, PBS	
Acid-treated cells		36	2 mol/L HCl, 1h, 37 °C, 5% CO ₂	
Iodate				
Viable cells		83	1h, 37 °C, 5% CO ₂ , PBS	
Heat-treated cells		84	Boiled for 1h, PBS	
Acid-treated cells		80	2 mol/L HCl, 1h, 37 °C, 5% CO ₂	
Urea				
Viable cells		64	1h, 37 °C, 5% CO ₂ , PBS	
Heat-treated cells		60	Boiled for 1h, PBS	
Acid-treated cells		50	2 mol/L HCl, 1h, 37 °C, 5% CO ₂	
Water (Milli Q)				
Viable cells		76	1h, 37 °C, 5% CO ₂ , PBS	
Heat-treated cells		83	Boiled for 1h, PBS	
Acid-treated cells		84	2 mol/L HCl, 1h, 37 °C, 5% CO ₂	

Microorganism	AF	Bound (%)	Conditions	Ref.
<i>L. acidophilus</i>	B1			[49]
Pre-treatment:	5 µg/mL			
None		56.6	4h, 37 °C, PBS	
Heat		71.9		
Ethanol		46.5		
Acid		87.0		
Alkaline		27.4		
<i>L. casi</i>				
None		22.4		
Heat		41.8		
Ethanol		21.8		
Acid		43.1		
Alkaline		12.0		
<i>L. helveticus</i>				
None		17.8		
Heat		28.5		
Ethanol		18.0		
Acid		56.3		
Alkaline		9.1		
<i>L. bulgaricus</i>				
None		16.3		
Heat		33.5		
Ethanol		15.9		
Acid		586		
Alkaline		8.3		
<i>L. plantarum</i>	M1		10 ¹⁰ cfu/mL, 15 min, 37°C, in:	[50]
Viable cells	0.15	5.6	PBS	
Heat-killed cells	µg/mL	8.1	PBS	
<i>E. avium</i>	PBS			
Viable cells	solution	7.4	PBS	
Heat-killed cells	0.5	6.6	PBS	
<i>P. pentosaceus</i>	µg/mL			
Viable cells	skimmed	8.7	PBS	
Heat-killed cells	milk	7.8	PBS	
<i>L. gasseri</i>				
Viable cells		21.4	PBS	
Heat-killed cells		22.8	PBS	
<i>L. bulgaricus</i>				
Viable cells		30.2	PBS	
Heat-killed cells		33.5	PBS	

Microorganism	AF	Bound (%)	Conditions	Ref.
		33.5	skimmed milk	
<i>L. rhamnosus</i>				
Viable cells		17.1		
Heat-killed cells		27.8	PBS	
		24.5	PBS	
			skimmed milk	
<i>B. lactis</i>				
Viable cells		16.9	PBS	
Heat-killed cells		23.6	PBS	
		32.5	skimmed milk	
<i>L. delbrueckii</i> subsp.	M1	18.7	4h, 37 °C, PBS	[51]
<i>bulgaricus</i> CH-2	10 ng/mL	27.6	4h, 42 °C, milk	
<i>S. thermophilus</i> ST-36		29.4	4h, 37 °C, PBS	
		39.2	4h, 42 °C, milk	
		14.8	Yoghurt	
<i>F. aurantiacum</i> NRRL	M1	100	5 x 10 ¹⁰ cfu/mL, 30 °C, 4h, PBS	[52]
B-184	10 µg/mL		and milk	
<i>B. subtilis</i> UTBSP1	B1			[53]
Viable cells	2.5	85.7	96h, 30 °C, nutrient broth culture	
Cell Free Supernatant	µg/mL	95	10 ⁸ cfu/ml, 120 h, 30 °C, pistachio nuts	
			120 h, 35 °C, nutrient broth culture	
		78.4		
<i>B. subtilis</i> ANSB060	B ₁			[54]
"Inocula" suspension	G ₁	81.5	72h, 37 °C, Luria-Bertani medium	
	M1	80.7		
	(0.5 µg/mL)	60		
Cell	B1	10.5	72h, 37 °C, PBS	
Cell extract		9.6		
Culture Supernatant		78.7		
<i>L. rhamnosus</i> strain GG	B1		10 ¹⁰ cfu/mL:	[62]
"In vivo"	3 µg/mL	51	1 min, duodenum of chicks	
		92		
"In vitro"		80	1 h, duodenum of chicks	
<i>L. rhamnosus</i> strain LC-705			37 °C, 1h , pH 7.3	

Microorganism	AF	Bound (%)	Conditions	Ref.
"In vivo"		36	1 min, duodenum of chicks	
		71	1 h, duodenum of chicks	
"In vitro"		77	37 °C, 1h , pH 7.3	
<i>P. freudenreichii</i> subsp.				
<i>shermanii</i> JS				
"In vivo"		37	1 min, duodenum of chicks	
		82	1 h, duodenum of chicks	
"In vitro"		22	37 °C, 1h , pH 7.3	

Table 1. Aflatoxin binding / absorption by microorganisms. Note: PBS, Phosphate-Buffered Saline; cfu, colony formingunit.

4. Decontamination of Aflatoxins by Yeasts

Yeasts are non-photosynthetic organisms with a separate nucleus and complex life cycle. They are larger than bacteria, normally spherical, non-motile, and reproduce by budding. Although their main function is alcoholic fermentation, these organisms are also capable of producing enzymes and vitamins. The primary substrates for yeasts are fermentable sugars, which are mainly transformed in ethanol, carbon dioxide, and biomass under oxygen-limited conditions. Under adequate oxygen supply, yeast produces carbon dioxide, water, and biomass [65]. *Saccharomyces cerevisiae* (SC) is the most well-known and commercially important species of yeast, and SC strains are widely used in the production of alcoholic drinks and in the baking industry.

As it occurs with LAB, SC cells have been studied to evaluate their ability to remove aflatoxins from contaminated media. The most important results obtained until now are summarized in Table 2. Products based on SC (cell wall from baker and brewer yeasts, inactivated baker yeast, or alcohol yeast) was studied, and it was observed that in pH 3, 37 °C and 15 minutes of contact, AFB₁ removal ranged from 2.5% to 49.3%, depending on the concentration of the toxin in the medium, and on the yeast-based products used [66]. These authors also observed a decrease in toxin adsorption as the initial concentration increased, and concluded that adsorption is not a linear phenomenon. Similar results with a SC strain and AFB₁ concentration ranging from 1 to 20 µg/mL was also reported [56]. At the 1 µg/mL concentration, 69.1% AFB₁ was removed; at 5 g/mL, removal rate was 41%; and at 20 µg/mL, 34%. *S. cerevisiae* strains were isolated from animal feed, feces and intestines, and tested for their ability to tolerate gastrointestinal conditions and remove AFB₁ from a contaminated medium [67]. These researchers observed that all strains isolated were able to survive in gastrointestinal conditions, and that the percentage of toxin removed ranged among SC strains (10⁷ CFU/mL), and with AFB₁ concentration used (16.4% to 82% of adsorption for 50 ng/mL AFB₁; 21.3% to 48.7% for 100 ng/mL AFB₁; and 20.2% to 65.5% for 500 ng/mL AFB₁).

The ability of SC (0.1%, 0.2%, and 0.3%) to adsorb AFB₁ in contaminated corn (150, 300, 450 and 800 µg/kg corn was analyzed [68]. The adsorption process showed an inversely proportional relationship with the concentration, that is, the greater the AFB₁ concentration in the medium, the lower the efficiency of AFB₁ removal by SC (16% to 66% for 800 µg/kg AFB₁ vs. 40% to 93% for 150 µg/kg AFB₁). The authors concluded, using densitogram analysis, that the adsorption process did not change the molecular structure of the mycotoxin, and that the decreased AFB₁ adsorption rates observed as the toxin concentration increased may possibly be caused by saturation of the adsorption sites on the SC cell. Other factors, such as length of incubation, pH, method of biomass purification, and methods of analysis, may also influence this process.

Immobilized SC cells (ATTC 9763) was investigated for their ability to remove AFB₁ from pistachio seeds, and it was observed that the amount of toxin removed was dependent on its concentration in the medium (40% and 70% of removal for concentrations of 10 ng/mL and 20 ng/mL AFB₁, respectively) [69]. The authors also concluded that this ability to remove the toxin was greater in SC exponential growth phase, and that the process was a quick one, being saturated after 3 hours of contact. Besides, the ability of SC cells to remove toxin was increased after treatment with acid (60% and 73% for 10 ng/mL and 20 ng/mL AFB₁, respectively) and heat (55% and 75%, respectively). In another study, authors also concluded that the treatment of SC cells with heat at 60 °C and 120 °C, and with chloric acid (2 mol/L) increased their ability to remove AFB₁ from the medium to 68.8%, 79.3%, and 72.1%, respectively, against 38.7% when viable yeast cells were used [56].

Heat treatment may increase the permeability of the external layer of the cell wall due to the suspension of some mannanes on the cell surface, leading to increased availability of previously hidden binding sites. Besides, countless physical-chemical changes take place on the cell wall during heat treatment, leading to more exposed binding sites. On the other hand, acid conditions may affect polysaccharides by releasing monomers, which are further fragmented in aldehydes after glycosidic bonds are broken. Continuous removal of aflatoxin, even after use of acid and heat treatments, confirms that yeast cell viability is not a significant factor for the removal of aflatoxin from the medium [69].

During the fermentation of broiler feed using LAB (3 strains of *Lactobacillus*) and SC strains resistant to gastric juices and bile, 55% AFB₁ was removed when AFB₁ concentration in the medium was 1 mg/kg, and 39% when concentration was 5 mg/kg AFB₁, after 6 hours [70]. This tendency for removal was maintained as incubation continued, and after 24 hours, the amount of AFB₁ removed was 73% and 53%, respectively, for the two concentrations of the toxin. The authors considered that, from a practical point of view, the most important factor was the 6-hour fermentation period, once the passage of feed through the gastrointestinal tract of broilers lasts from 4 to 8 hours. In reference [71], authors analyzed the ability of SC to remove AFB₁ from a contaminated medium at different pH values (3.0, 6.0, and 8.0), and observed that the three strains analyzed showed great ability to remove the toxin (41.6% to 94.5%), and that after washing, only a small amount of AFB₁ was released back into the medium. *In vitro* studies are not always good indications of the *in vivo* responses, as *in vivo*

studies are affected by physiological parameters, such as pH, peristaltic movements, and gastric and intestinal secretions.

In vivo studies using SC are not as rare as those with LAB, mainly in poultry science. Generally, SC is added to the feed as a growth promoter. However, the addition of yeasts has also presented beneficial effects against the exposure to AFB₁. It was observed that the addition of 1% SC to feed contaminated with 5 g/g of AFB₁ prevented loss of weight; liver and heart hyperplasia; and decreased serum albumin and total protein concentrations in broilers [72]. The addition of SC in feed containing aflatoxin decreased the deleterious effects on feed intake, weight gain, and feed conversion in Japanese quails [73]. Compared with control animals, weight gain was 37% lower in birds fed a diet added only of aflatoxin, and was 15% greater than the control in the group that received feed containing aflatoxin and SC. The authors concluded that the diet containing with only SC significantly improved all growth parameters investigated (about 40%), compared with the control group.

In a study with mice, it was observed that the addition of AFB₁ to the diet (0.4 and 0.8 mg/kg) caused a significant reduction in weight gain, and an increase of 85% (0.8 mg/kg) in the rate of micronucleated normochromatic erythrocytes (MNE) after 3 weeks of ingestion, compared with the control group [68]. When diets containing AFB₁ and SC (0.3%) were administered, weight gain was twice greater than in diets that contained only the toxin, and the rate of MNE increased only 46% (0.8 mg/kg). The authors stated that reduced body weight is one of the most common consequences of AFB₁ ingestion, because the toxin alters the activity of several digestive enzymes, giving rise to a malabsorption syndrome characterized by steatorrhea, hypovitaminosis A and a decrease in the levels of bile, pancreatic lipase, trypsin, and amylase. Besides, biotransformation of AFB₁ gives rise to several metabolites, particularly AFB₁-8,9-epoxide, which may bind covalently to DNA and proteins, changing enzymatic processes such as gluconeogenesis, Krebs cycle, and fatty acid synthesis [74]. MNE rate is used to determine the genotoxicity of AFB₁, because it quantifies broken chromosomes and whole chromosomes that are abnormally distributed to daughter cells, showing thus, that AFB₁ is a potent mutagenic agent.

A diet containing 5 g/g of aflatoxin (82.06% AFB₁, 12.98% AFB₂, 2.84% AFG₁, and 1.12% AFG₂) by female quails (49 to 84 days of age) led to decreased egg production, feed intake, and feed conversion (31%, 28%, and 47%, respectively) [75]. However, addition of SC (2 g/kg) significantly increased these parameters (16%, 4%, and 14%, respectively). They also observed that the diet with aflatoxins caused a marked decrease in weight gain and egg weight, besides increasing animal mortality (39%, 7%, and 50%, respectively), whereas addition of SC reverted the negative effect on these parameters (65%, 8%, and 50%, respectively). The authors stated that these negative effects of aflatoxins in egg production, feed intake, and feed conversion may have been caused by anorexia, apathy, and inhibition of protein synthesis and lipogenesis. Besides, affected liver function and mechanisms of use of protein and lipids may have affected performance criteria and the general health of the animals. In reference [76] authors reported that the components of the cells wall of SC are able to adsorb mycotoxins, stimulate the immune system, and compete for binding sites in the enterocytes, inhibiting intestinal colonization by pathogens.

SC cell wall is mainly made up of polysaccharides (80-90%), and its mechanical resistance is due to an inside layer composed of β -D-glucans, which are formed by a complex network of highly polymerized β -(1,3)-D-glucans, branched off as β -(1,6)-D-glucans, that have a low level of polymerization. This inside layer is firmly bound to the plasmatic membrane by linear chains of chitin, which have a significant role in the insolubility of the overall structure and packing of the branched β -D-glucans. Both chitin chains and β -D-glucans affect the plasticity of the cell wall. The external layer of the yeast cell wall is formed by mannoproteins, which have an important role in the exchanges with the external environment. This whole structure is highly dynamic and may vary according to the yeast strain, phase of the cell cycle, and culture conditions, such as pH, temperature, oxygenation rate, nature of the medium, concentration and nature of the carbon source. Thus, these differences in the composition of the cell wall among yeast strains are related with their ability to bind to the mycotoxin [77].

Studies have shown that the components of SC cell wall, called oligomannanes, after esterification, are able to bind more than 95% AFB₁ [78]. Addition of 0.05% glucomannanes in the basal diet improved broiler performance [79].

The possible binding mechanisms between yeast cell wall and mycotoxins were studied, and authors suggested that β -D-glucans are the components of the cell wall that are responsible for forming the complex with the toxin, and that the reticular organization of β -D-glucans and their distribution in β -(1,3)-D-glucans and β -(1,6)-D-glucans have an important role in the efficiency of the bond [77]. Besides, studies have shown that weak hydrogen and van der Waals bonds are involved in the complex chemical connection between the mycotoxins and β -D-glucans, a chemical interaction that is much more "adsorption" than "bond". As for AFB₁, they observed that the aromatic ring, the lactone and ketone groups of the polar form of AFB₁, or chemical bonds with glucose units in the single helix of the β -D-glucans, are what keep the toxin bound to the glucans.

It was demonstrated that yeast strains isolated from environments where animals are raised are able to bind to AFB₁ in saline solution (PBS, pH 7) [67]. These strains presented other properties that were beneficial to the host, such as the inhibition of pathogenic bacteria. Therefore, SC strains acted both as probiotics (co-aggregation and inhibition of pathogenic bacteria), and as mycotoxin adsorbents.

In reference [72], SC was able to reduce the deleterious effects of AFB₁ in the diet of broilers and in [68] authors replicated these findings in rats. Protective effect against aflatoxins produced by yeasts was confirmed in rats. However, when yeast cells were inactivated by heat, they were inefficient [80] but when glucomannanes extracted from the cell wall of yeasts were used, there was an increase in the efficiency of the bond with AFB₁, OTA and T-2 toxin [81-84], individually or in combination [75, 79, 85, 86]. The addition of SC in the diet reduced AFB₁ toxic effects in chickens [72, 87]. The ability of SC to reduce AFB₁ toxic effects in quails was demonstrated, and this effect was apparently more efficient with the increase in inclusion rates [88].

In [89] authors obtained a significant reduction in AFB₁ concentration during beer production, probably due to the bond between mycotoxins and SC cell. This hypothesis was supported by other studies [39, 90]. A 19% reduction in AFB₁ during dough fermentation in bread production was observed [91].

Microorganism	AF	Bound (%)	Conditions	Ref.
<i>S. cerevisiae</i>	B ₁		15 min, 37 °C:	[66]
	0.0058-	7.6-49.3	YCW from brewer's yeast	
	6.35 µg/mL	7.6-29	YCW from brewer's yeast	
		10-24	Inactivated baker's yeast	
		4-29	YCW from baker's yeast	
		17-44	Inactivated baker's yeast	
		3-44	YCW from baker's yeast	
		23-35	YCW from baker's yeast	
		27-44	Alcohol yeast	
<i>S. cerevisiae</i>	B1			[56]
Strain A18	1 µg/mL	69.1	3h, 25 C, PBS	
	5 µg/mL	41		
	10 µg/mL	33		
	20 µg/mL	34.2		
Strain 26.1.11	1 µg/mL	65.1	3h, 25 C, PBS	
	5 µg/mL	37.2		
	10 µg/mL	31		
	20 µg/mL	32.6		
Pre-treatment:				
Heated cells 52°C	5 µg/mL	58.8	3h, 25 C, PBS	
Strain A18		56.5		
Strain 26.1.11				
Heated cells 55 °C		64.5		
Strain A18		64		
Strain 26.1.11				
Heated cells at 60 °C		68.8		
Strain A18		67		
Strain 26.1.11				
Heat cells at 120 °C		79.3		
Strain A18		77.7		
Strain 26.1.11				
2 mol/L HCl / 1h		72.1		
Strain A18		69.3		
Strain 26.1.11				
<i>S. cerevisiae</i>	B ₁ (ng/mL)			[67]

Microorganism	AF	Bound (%)	Conditions	Ref.
Strain RC008	50 ng/mL		10 ⁷ cells/mL, 1h, 37 °C, PBS	
	100	67.6		
	500	43.5		
	50	38.2		
Strain RC009	100	16.4		
	500	21.3		
	50	31.8		
Strain RC012	100	29.6		
	500	20.6		
	50	20.2		
Strain RC016	100	82.0		
	500	48.7		
		65.5		
<i>S. cerevisiae</i>	B1			[68]
Yeast concentration:	(µg/kg)			
0.1 %	150	40	37 °C, 24 h, corn	
	300	25		
	450	17		
	800	16		
0.2 %	150	88	37 °C, 24 h, corn	
	300	76		
	450	64		
	800	51		
0.3 %	150	93	37 °C, 24 h, corn	
	300	86		
	450	81		
	800	66		
<i>S. cerevisiae</i> ATTC 9763	B1			[69]
Pre-treatment:	(ng/mL)			
None	10	40	3 h, 25 °C, pistachio nuts	
	20	70		
Acid treated cells	10	60		
(2 mol/L / 90 min)	20	73		
Heat-treated cells	10	55		
(120 °C / 20 min)	20	75		
<i>L. paracasei</i> LOCK 0920,	B ₁			[70]
<i>L. brevis</i> LOCK 0944, <i>L.</i>	1 mg/kg	55	37 °C, 6h fermentation in broiler feed	
<i>plantarum</i> LOCK 0945,	5 mg/kg	39		
and <i>S. cerevisiae</i> LOCK 0140				

Microorganism	AF	Bound (%)	Conditions	Ref.
<i>S. cerevisiae</i>	B ₁			[89]
	1 µg /g	86	12 °C, 8 days, brewing process	
	10 µg /g	72		

Table 2. Aflatoxin binding by yeasts. YCW, Yeast Cell Wall

5. Concluding Remarks

Considering the data from several studies carried out until now, it may be observed that microorganisms, among them lactic acid bacteria and yeasts, have a huge potential application in aflatoxin degradation in foodstuffs. However, new studies are necessary to identify bacterial species with greater binding potential with aflatoxins, once there are differences in sensitivity and selectivity, besides the influence of factors that are intrinsic and extrinsic to the bacteria in the decontamination process. After this step of choosing species with greater efficiency has been overcome, new production technologies that are economically viable to be applied to human and animal foods may be developed.

Several studies have demonstrated that the cell wall of SC and LAB and their components are responsible for binding with aflatoxins. However, the mechanisms by which this bond occurs remain unclear. Cell walls with glucomannanes and manno-oligosaccharides have been pointed out as the responsible elements for AFB₁ bond with yeasts. The great advantage in the commercial use of these microorganisms as binding agents is that these strains are approved and already used in a wide range of fermented food products, being recognized as safe. However, aflatoxin may be released from the cell-aflatoxin complex with changes in the pH and temperature conditions. Therefore, further studies are necessary to determine the behavior of yeasts in the different environmental conditions before they are used commercially.

Acknowledgements

The authors wish to acknowledge the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) and the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) of Brazil for financial support and fellowships.

Author details

Carlos Augusto Fernandes Oliveira^{1*}, Fernanda Bovo¹, Carlos Humberto Corassin¹, Alessandra Vincenzi Jager¹ and Kasa Ravindranadha Reddy²

*Address all correspondence to: carlosaf@usp.br

1 Faculty of Animal Science and Food Engineering, University of São Paulo, Brazil

2 Research and Development Department, SRI Biotech Laboratories Ltd, India

References

- [1] Bata, A., & Lasztity, R. (1999). Detoxification of mycotoxin-contaminated food and feed by microorganisms. *Trends in Food Science & Technology*, 10-223.
- [2] Gonçalves, E., Pinto, M. M., & Felicio, J. D. (2001). Análise de micotoxinas no Instituto Biológico de 1989 a 1999. *Biológico*, 63(1/2), 15-19.
- [3] D'Mello, J. P. F., & Mac, Donald. A. M. C. (1997). Mycotoxins. *Animal Feed Science and Technology*, 69-155.
- [4] Nierman, W. C., Cleveland, T. E., Payne, G. A., Keller, N. P., Campbell, B. C., Bennett, J. W., Guo, B., Yu, J., & Robens, J. F. (2008). Mycotoxin Production and Prevention of Aflatoxin Contamination in Food and Feed. In: Goldman GH, Osmani SA. *The Aspergilli: Genomics, Medical Aspects, Biotechnology, and Research Methods*. Boca Raton: CRC Press , 457-472.
- [5] Hussein, H. S., & Brasel, J. M. (2001). Toxicity, metabolism, and impact of mycotoxins on humans and animals. *Toxicology*, 167-101.
- [6] Bhat, R., Rai, R. V., & Karim, A. A. (2010). Mycotoxins in Food and Feed: Present Status and Future Concerns. *Comprehensive Reviews in Food Science and Food Safety*, 9-57.
- [7] Alberts, J. F., Engelbrecht, Y., Steyn, P. S., Holzapfel, W. H., & Van Zyl, W. H. (2006). Biological degradation of aflatoxin B1 by *Rhodococcus erythropolis* cultures. *International Journal of Food Microbiology*, 109-121.
- [8] Park, D. L., & Liang, B. (1993). Perspectives on aflatoxin control for human food and animal feed. *Trends in Food Science & Technology*, 41-334.
- [9] Chu, F. S. (1991). Mycotoxins: food contamination, mechanism, carcinogenic potential and preventive measures. *Mutation Research*, 259-291.
- [10] Abbas, H. K. (2005). Aflatoxin and food safety. Boca Raton, CRC Press.
- [11] Magan, N., & Olsen, M. (2006). Mycotoxins in food: Detection and Control. Boca Raton, CRC Press.
- [12] Prandini, A., Tansini, G., Sigolo, S., Filippi, L., Laporta, M., & Piva, G. (2009). On the occurrence of aflatoxin M₁ in milk and dairy products. *Food and Chemical Toxicology*, 47-984.

- [13] Hernandez-Mendoza, A., Garcia, H. S., & Steele, J. L. (2009). Screening of *Lactobacillus casei* strains for their ability to bind aflatoxin B₁. *Food and Chemical Toxicology*, 47-1064.
- [14] Gratz, S., Mykkänen, H., & El-Nezami, H. (2005). Aflatoxin B1 binding by a mixture of *Lactobacillus* and *Propionibacterium*: in vitro versus ex vivo. *Journal of Food Protection*, 68(11), 2470-2474.
- [15] El-Nezami, H., Kankaanpää, P., Salminen, S., & Ahokas, J. (1998). Physicochemical alterations enhance the ability of dairy strains of lactic acid bacteria to remove aflatoxin from contaminated media. *Journal of Food Protection*, 61(4), 466-468.
- [16] Line, J. E., & Brackett, R. E. (1995). Factors affecting aflatoxin B1 removal by *Flavobacterium aurantiacum*. *Journal of Food Protection*, 58(1), 91-94.
- [17] El-Nezami, H., Kankaanpää, P., Salminen, S., & Ahokas, J. (1998). Ability of dairy strains of lactic acid bacteria to bind a common food carcinogen, aflatoxin B₁. *Food and Chemical Toxicology*, 36-321.
- [18] Fazeli, M. R., Hajimohammadali, M., Moshkani, A., Samadi, N., Jamalifar, H., Khoshayand, M. R., et al. (2009). Aflatoxin B1 binding capacity of autochthonous strains of lactic acid bacteria. *Journal of Food Protection*, 72(1), 189-192.
- [19] Alberts, J. F., Gelderblomb, W. C. A., Botha, A., & Van Zyl, W. H. (2009). Degradation of aflatoxin B1 by fungal laccase enzymes. *International Journal of Food Microbiology*, 135-47.
- [20] Wu, Q., Jezkova, A., Yuan, Z., Pavlikova, L., Dohnal, V., & Kuca, K. (2009). Biological degradation of aflatoxins. *Drug Metabolism Reviews*, 41(1), 1-7.
- [21] Oliveira, C. A. F., & Germano, P. M. L. (1997). Aflatoxinas: conceitos sobre mecanismos de toxicidade e seu envolvimento na etiologia do câncer hepático celular. *Revista de Saúde Pública*, 31(4), 417-424.
- [22] Oatley, J. T., Rarick, M. D., Ji, G. E., & Linz, J. E. (2000). Binding of aflatoxin B1 to bifidobacteria in vitro. *Journal of Food Protection*, 63(8), 1133-36.
- [23] Peltonen, K., El-Nezami, H., Haskard, C., Ahokas, J., & Salminen, S. (2001). Aflatoxin B1 binding by dairy strains of lactic acid bacteria and bifidobacteria. *Journal of Dairy Science*, 84, 2152-2156.
- [24] Murphy, P. A., Hendrich, S., Landgren, C., & Bryant, C. M. (2006). Food Mycotoxins: An Update. *Journal of Food Science*, 71(5), 51-65.
- [25] Creppy, E. E. (2002). Update of survey, regulation and toxic effects of mycotoxins in Europe. *Toxicology Letters*, 127-19.
- [26] Bakirci, I. (2001). A study on the occurrence of aflatoxin M1 in milk and milk products produced in Van province of Turkey. *Food Control*, 12-47.

- [27] Haskard, C. A., El -Nezami, H. S., Kankaanpaa, P. E., Salminen, S., & Ahokas, J. T. (2001). Surface binding of aflatoxin B1 by lactic acid bacteria. *Applied and Environmental Microbiology*, 67(7), 3086-3091.
- [28] Oluwafemi, F., & Silva, F. A. (2009). Removal of aflatoxins by viable and heat-killed *Lactobacillus* species isolated from fermented maize. *Journal of Applied Biosciences.*, 16-871.
- [29] Pierides, M., El -Nezami, H., Peltonem, K., Salminen, S., & Ahokas, J. (2000). Ability of dairy strains of lactic acid bacteria to bind aflatoxin M1 in a food model. *Journal of Food Protection*, 63(5), 645-650.
- [30] IARC International Agency for Research on Cancer- World Health Organization. (2002). *IARC Monograph on the Evaluation of Carcinogenic Risk to Humans.*, 82, 171.
- [31] Mishra, H. N., & Das, C. (2003). A Review on biological control and metabolism of aflatoxin. *Critical Reviews in Food Science and Nutrition*, 43(3), 245-264.
- [32] Laciaková, A., Cicoðová, P., Máté, D., & Laciak, V. (2008). Aflatoxins and possibilities for their biological detoxification. *Medycyna Weterynaryjna*, 64(3), 276-279.
- [33] Van Egmond, H. P., & Jonker, M. A. (2004). Worldwide regulations on aflatoxins- The situation in 2002. *Toxin Reviews*, 23, 273-293.
- [34] Walstra, P., Wouters, J. T. M., & Geurts, T. J. (2006). *Dairy Science and Technology*. Boca Raton, CRC Press.
- [35] Salminen, S., Von, Wright. A., & Ouwehand, A. C. (2004). *Lactic acid bacteria: microbiological and functional aspects*. 3.ed. New York Marcel Dekker, Inc.
- [36] Sybesma, W. F., & Hugenholtz, J. (2004). Food fermentation by lactic acid bacteria for the prevention of cardiovascular disease. In: Arnoldi A. *Functional Foods, Cardiovascular Disease and Diabetes*. Boca Raton CRC Press
- [37] Onilude, A. A., Fagade, O. E., Bello, M. M., & Fadahunsi, I. F. (2005). Inhibition of aflatoxin-producing aspergilli by lactic acid bacteria isolates from indigenously fermented cereal gruels. *African Journal of Biotechnology*, 4(12), 1404-1408.
- [38] Fuchs, S., Sontag, G., Stidl, R., Ehrlich, V., Kundi, M., & Knasmüller, S. (2008). Detoxification of patulin and ochratoxin A, two abundant mycotoxins, by lactic acid bacteria. *Food and Chemical Toxicology*, 46-1398.
- [39] Shetty, P. H., & Jespersen, L. (2006). *Saccharomyces cerevisiae* and lactic acid bacteria as potential mycotoxin decontaminating agents. *Trends in Food Science & Technology*, 17-48.
- [40] Ciegler, A., Lillehoj, E. B., Peterson, R. E., & Hall, H. H. (1966). Microbial detoxification of aflatoxin. *Applied Environmental Microbiology*, 14(6), 934-939.
- [41] El -Nezami, H., Mykkänen, H., Haskard, C., Salminen, S., & Salminen, E. (2004). Lactic acid bacteria as a tool for enhancing food safety by removal of dietary toxins. In:

- Salminen S, Von Wright A, Ouwehand A. Lactic Acid Bacteria: Microbiological and Functional Aspects. 3.ed. Nova Iorque Marcel Dekker , 397-406.
- [42] Shahin, A. A. M. (2007). Removal of aflatoxin B₁ from contaminated liquid media by dairy lactic acid bacteria. *International Journal of Agriculture and Biology*, 9(1), 71-75.
- [43] Rasic, J. L., Skrinjar, M., & Markov, S. (1991). Decrease of aflatoxins B₁ in yoghurt and acidified milks. *Mycopathologia*, 113-117.
- [44] Biernasiak, J., Piotrowska, M., & Libudzisz, Z. (2006). Detoxification of mycotoxins by probiotic preparation for broiler chickens. *Mycotoxin Research*; , 22(4), 230-235.
- [45] Kabak, B., & Dobson, A. D. W. (2009). Biological strategies to counteract the effects of mycotoxins. *Journal of Food Protection*, 72(9), 2006-2016.
- [46] Kabak, B., & Var, I. (2008). Factors affecting the removal of aflatoxin M₁ from food model by Lactobacillus and Bifidobacterium strains. *Journal of Environmental Science and Health, Part B*, 43-617.
- [47] Elgerbi, A. M., Aidoo, K. E., Candlish, A. A. G., & Williams, A. G. (2006). Effects of lactic acid bacteria and bifidobacteria on levels of aflatoxin M₁ in milk and phosphate buffer. *Milchwissenschaft*, 61(2), 197-199.
- [48] Haskard, C., Binnion, C., & Ahokas, J. (2000). Factors affecting the sequestration of aflatoxin by Lactobacillus rhamnosus strain GG. *Chemico-Biological Interactions*, 128-39.
- [49] Azab, R. M., Tawakkol, W. M., Hamad, A. R. M., Abou-Elmagd, M. K., El -Agrab, H. M., & Refai, M. K. (2005). Detection and estimation of aflatoxin B₁ in feeds and its bio-degradation by bacteria and fungi. *Egypt Journal of Natural Toxins*, 2-39.
- [50] Bovo, F., Corassin, C. H., Rosim, R. E., & Oliveira, C. A. F. (2012). Efficiency of lactic acid bacteria strains for decontamination of aflatoxin M₁ in phosphate buffer saline solution and in skimmed milk. *Food and Bioprocess Technology*.
- [51] Sarimehmetoglu, B., & Küplülü, Ö. (2004). Binding ability of aflatoxin M₁ to yoghurt bacteria. *Ankara Üniversitesi Veteriner Fakültesi Dergisi*; , 51-195.
- [52] Lillehoj, E. B., Stubblefield, R. D., Shannon, G. M., & Shotwell, O. L. (1971). Aflatoxin M₁ removal from aqueous solutions by Flavobacterium aurantiacum. *Mycopathologia et mycologia applicata*., 45-259.
- [53] Farzaneh, M., Shi, Z. Q., Ghassempour, A., Sedaghat, N., Ahmadzadeh, M., Mirabol-fathy, M., & Javan-Nikkhah, M. (2012). Aflatoxin B₁ degradation by Bacillus subtilis UTBSP1 isolated from pistachio nuts of Iran. *Food Control*, 23-100.
- [54] Gao, X., Ma, Q., Zhao, L., Lei, Y., Shan, Y., & Ji, C. (2011). Isolation of Bacillus subtilis: screening for aflatoxins B₁, M₁ and G₁ detoxification. *European Food Research and Technology*, 232-957.

- [55] Lahtinen, S. J., Haskard, C. A., Ouwehand, A. C., Salminen, S. J., & Ahokas, J. T. (2004). Binding of aflatoxin B₁ to cell wall components of *Lactobacillus rhamnosus* strain GG. *Food Additives and Contaminants*, 21(2), 158-164.
- [56] Shetty, P. H., Hald, B., & Jespersen, L. (2007). Surface binding of aflatoxin B₁ by *Saccharomyces cerevisiae* strains with potential decontaminating abilities in indigenous fermented foods. *International Journal of Food Microbiology*, 113-41.
- [57] Hernandez-Mendoza, A., Guzman-de-Peña, D., & Garcia, H. S. (2009). Key role of teichoic acids on aflatoxin B₁ binding by probiotic bacteria. *Journal of Applied Microbiology*, 107-395.
- [58] Hernandez-Mendoza, A., Guzman-de-Peña, D., González-Córdova, A. F., Vallejo-Córdova, B., & Garcia, H. S. (2010). In vivo assessment of the potential protective effect of *Lactobacillus casei* Shirota against aflatoxin B₁. *Dairy Science and Technology*, 90-729.
- [59] Bolognani, F., Rummey, C. J., & Rowland, I. R. (1997). Influence of carcinogen binding by lactic acid-producing bacteria on tissue distribution and in vivo mutagenicity of dietary carcinogens. *Food and Chemical Toxicology*, 35-535.
- [60] Lee, Y. K., El -Nezami, H., Haskard, C. A., Gratz, S., Puong, K. Y., Salminen, S., & Mykkänen, H. (2003). Kinetics of adsorption and desorption of aflatoxin B₁ by viable and nonviable bacteria. *Journal of Food Protection*, 66(3), 426-430.
- [61] D'Souza, D. H., & Brackett, R. E. (2001). Aflatoxin B₁ degradation by *Flavobacterium aurantiacum* in the presence of reducing conditions and seryl and sulfhydryl group inhibitors. *Journal of Food Protection*, 64(2), 268-271.
- [62] El -Nezami, H., Mykkänen, H., Kankaanpää, P., Salminen, S., & Ahokas, J. (2000). Ability of *Lactobacillus* and *Propionibacterium* strains to remove aflatoxin B₁ from the chicken duodenum. *Journal of Food Protection*, 63(4), 549-552.
- [63] Hathout, A. S., Mohamed, S. R., El -Nekeety, A. A., Hassan, N. S., Aly, S. E., & Abdel-Wahhab, M. A. (2011). Ability of *Lactobacillus casei* and *Lactobacillus reuteri* to protect against oxidative stress in rats fed aflatoxins-contaminated diet. *Toxicon*, 58-179.
- [64] El -Nezami, H. S., Mykkänen, H., Kankaanpää, P., Suomalainen, T., Ahokas, J. T., & Salminen, S. (2000). The ability of a mixture of *Lactobacillus* and *Propionibacterium* to influence the faecal recovery of aflatoxins in healthy Egyptian volunteers: a pilot clinical study. *Bioscience and Microflora*, 19-41.
- [65] Mc Kinney, R. E. (2004). Environmental pollution control microbiology: a fifty-year perspective. Boca Raton, CRC Press.
- [66] Joannis-Cassan, C., Tozlovanu, M., Hadjeba-Medjdoub, K., Ballet, N., & Pfohl-Lesch, A. (2011). Binding of zearalenone, aflatoxin B₁, and ochratoxin A by yeast-based products: a method for quantification of adsorption performance. *Journal of Food Protection*, 74(7), 1175-1185.

- [67] Armando, M. R., Dogi, C. A., Pizzolitto, R. P., Escobar, F., Peirano, M. S., Salvano, M. A., Sabini, L. I., Combina, M., Dalcerro, A. M., & Cavaglieri, L. R. (2011). *Saccharomyces cerevisiae* strains from animal environment with in vitro aflatoxin B₁ binding ability and anti-pathogenic bacterial influence. *World Mycotoxin Journal*, 4(1), 59-68.
- [68] Madrigal-Santillán, E., Madrigal-Bujaidar, E., Márquez-Márquez, R., & Reyes, A. (2006). Antigenotoxic effect of *Saccharomyces cerevisiae* on the damage produced in mice fed with aflatoxin B₁ contaminated corn. *Food and Chemical Toxicology*, 44, 2058-2063.
- [69] Rahaie, S., Emam-Djomeh, Z., Razavi, S. H., & Mazaheri, M. (2010). Immobilized *Saccharomyces cerevisiae* as a potential aflatoxin decontaminating agent in pistachio nuts. *Brazilian Journal of Microbiology*, 41-82.
- [70] Slizewska, K., & Smulikowska, S. (2011). Detoxification of aflatoxin B₁ and change in microflora pattern by probiotic in vitro fermentation of broiler feed. *Journal of Animal and Feed Sciences*, 20-300.
- [71] Dogi, C. A., Armando, R., Ludueña, R., Le Blanc, A. M., Rosa, C. A. R., Dalcerro, A., & Cavaglieri, L. (2011). *Saccharomyces cerevisiae* strains retain their viability and aflatoxin B₁ binding ability under gastrointestinal conditions and improve ruminal fermentation. *Food Additives and Contaminantes, Part A*, 28(12), 1705-1711.
- [72] Stanley, V. G., Ojo, R., Woldeesenbet, S., Hutchinson, D. H., & Kubena, L. F. (1993). The use of *Saccharomyces cerevisiae* to suppress the effects of aflatoxicosis in broiler chicks. *Poultry Science*, 72(10), 1867-1872.
- [73] Parlat, S. S., Özcan, M., & Oguz, M. (2001). Biological suppression of aflatoxicosis in Japanese quail (*Coturnix coturnix japonica*) by dietary addition of yeast (*Saccharomyces cerevisiae*). *Research in Veterinary Science*, 7-207.
- [74] Lesson, S., Diaz, G., & Summers, J. (1995). Poultry metabolic disorders and mycotoxins. Montreal (Canada), University Books.
- [75] Yildiz, A. O., Parlat, S. S., & Yildirim, I. (2004). Effect of dietary addition of live yeast (*Saccharomyces cerevisiae*) on some performance parameters of adult Japanese quail (*Coturnix coturnix japonica*) induced by aflatoxicosis. *Revue Médecine Vétérinaire*, 155(1), 38-41.
- [76] Raju, M. V. L. N., & Devegowda, G. (2000). Influence of esterified-glucomannan on performance and organ morphology, serum biochemistry and haematology in broilers exposed to individual and combined mycotoxicosis (aflatoxin, ochratoxin and T-2 toxin). *British Poultry Science*, 41(5), 640-650.
- [77] Jouany, J. P., Yiannikouris, A., & Bertin, G. (2005). The chemical bonds between mycotoxins and cell wall components of *Saccharomyces cerevisiae* have been identified. *Archiva Zootechnica*, 8-26.

- [78] Devegowda, G., Aravind, B. I. R., & Morton, M. G. (1996). Saccharomyces cerevisiae and mannanoligosaccharides to counteract aflatoxicoses in broilers. *Proceedings of Australian Poultry Science Symposium*, , 103-106.
- [79] Aravind, K. L., Patil, V. S., Devegowda, G., Umakhanta, B., & Ganpule, S. P. (2003). Efficacy of esterified glucomannan to counteract mycotoxicosis in naturally contaminated feed on performance and serum biochemical, hematological parameters in broilers. *Poultry Science*, 82-571.
- [80] Baptista, A. S., Horii, J., Calori-domingues, M. A., Gloria, E. M., Salgado, J. M., & Vizzioli, M. R. (2002). Thermolysed and active yeast to reduce the toxicity of aflatoxin. *ScientiaAgricola*,, 59(2), 257-260.
- [81] Bejaoui, H., Mathieu, F., Taillandier, P., & Lebrihi, A. (2004). Ochratoxin A removal in synthetic and natural grape juices by selected oenological Saccharomyces strains. *Journal of Applied Microbiology*,, 97-1038.
- [82] Cecchini, F., Morassut, M., Moruno, E., & Di Stefano, R. (2007). Influence of yeast strain on ochratoxin A content during fermentation of white and red must. *Food Microbiology*, 23-411.
- [83] Angioni, A., Caboni, P., Garau, A., Farris, A., Orro, D., Budroni, M., & Cabras, P. (2007). In vitro interaction between ochratoxin A and different strains of Saccharomyces cerevisiae and Kloeckera apiculata. *Journal of Agricultural and Food Chemistry*, 55, 2043-2048.
- [84] Freimund, S., Sauter, M., & Rys, P. (2003). Efficient adsorption of the mycotoxins zearalenone and T-2 toxin on a modified yeast glucan. *Journal of Environmental Science and Health Part B*, 38-243.
- [85] Yiannikouris, A., François, J., Poughon, L., Dussap, C. G., Bertin, G., Jeminet, G., & Jouany, J. P. (2004). Alkali extraction of β -D-glucans from Saccharomyces cerevisiae cell wall and study of their adsorptive properties toward zearalenone. *Journal of Agriculture and Food Chemistry*, 52-3666.
- [86] Karaman, M., Basmacioglu, H., Ortatatli, M., & Oguz, H. (2005). Evaluation of the detoxifying effect of yeast glucomannan on aflatoxicosis in broilers as assessed by gross examination and histopathology. *British Poultry Science*, 46-394.
- [87] Santin, E., Paulillo, A. C., Nakagui, L. S. O., Alessi, A. C., Polveiro, W. J. C., & Maioraka, A. (2003). Evaluation of the efficacy of Saccharomyces cerevisiae cell wall to ameliorate the toxic effects of aflatoxin in broilers. *International Journal of Poultry Science*, 2(6), 465-468.
- [88] Eshak, M. G., Khalil, W. K. B., Hegazy, E. M., Ibrahim, M. F., Fadel, M., & Stino, K. R. D. (2010). Effect of SCE on reduction of aflatoxicosis, enhancement of growth performance and expression of neural and gonadal genes in Japanese quail. *Journal of American Science*, 6-824.

- [89] Chu, F. S., Chang, C. C., Ashoor, S. H., & Prentice, N. (1975). Stability of Aflatoxin B₁ and Ochratoxin A in Brewing. *Applied Microbiology*, 29(3), 313-316.
- [90] Celyk, K., Denly, M., & Savas, T. (2003). Reduction of toxic effects of aflatoxin by using baker yeast (*Saccharomyces cerevisiae*): in growing broiler chicken diets. *Revista Brasileira de Zootecnia*, 32-615.
- [91] El -Banna, A. A., Lau, P. Y., & Scott, P. M. (1983). Fate of mycotoxins during processing of foodstuffs. II- Deoxynivalenol (vomitoxin) during making of Egyptian bread. *Journal of Food Protection*, 46-484.