We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Starvation Conditions Effects on Carbohydrate Metabolism of Marine Bacteria

Monia El Bour

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/50877>

1. Introduction

In the coastal shorelines terrestrial organic materials transported by river runoff represent an important material source to the ocean and is estimated that 0.4×10^{15} g yr⁻¹ organic carbon is discharged to ocean by land flows or rivers (Meybeck, 1982, He et al, 2010). This amount of riverine organic carbon is sufficient to support the entire organic carbon turnover in the ocean (Williams and Druffel, 1987) even though, little terrestrial signal by carbon isotopic ratio ($\delta^{13}\text{C}$) of the bulk DOC in ocean was shown, suggesting that terrestrial organic carbon undergo rapid removal and decomposition within estuarine mixing (Druffel et al, 1992; Hedges et al, 1997).

Carbohydrates are, among others, the major components of identified organic matter in ocean and account for 3% to 30% of the bulk Dissolved Organic Carbon (DOC) (Gueuen et al, 2006; Hung et al, 2003; Pakulski and Benner, 1994), whereas, in estuarine and marine surface waters they are considered the most labile fractions of bulk organic matter and may play key roles in the geochemical cycles as reported by several studies (Benner et al, 1992; Burdige and Zheng, 1998; Middelboe et al, 1995; Murrell and Hollibaugh, 2000).

In complex macro-aggregates mixtures, the carbohydrates are dominant with proteins and lipids, with evidence that in mucilage samples carbohydrates are relevant components of the organic carbon (17–45% on dry weight basis depending on the age of the aggregates) with heavy colonization by several heterotrophic bacteria and autotrophic organisms which embedded the organic matrix (Simon et al, 2002; Urbani and Sist, 2003). Thus high density of micro organisms enriches the matrix in organic and inorganic nutrients in comparison with the surrounding water and such macro flocks are important hot-spots for bacterial growth and carbon cycling (Kaltenbock and Herndl, 1992; Alldredge, 2000 and Azam and Long, 2001) and then significance of bacteria for the formation and decomposition of aggregates appears to be higher than previously estimated (Simon et al, 2002). .

2. Importance of carbohydrates availability for marine bacteria

Marine micro-organisms have important established roles which may reflect large-scale changes within inter tidal systems and heterotrophic bacteria, for example, are crucial to transformations and re-mineralization of organic carbon, nitrogen and other nutrients throughout oceans (Azam, 1998; Azam et al, 1993).

Bacterial utilization of carbohydrates depends heavily on their chemical composition (Aluwihare and Repeta, 1999; Arnosti, 2000, Zoppini et al, 2010) and slow hydrolysis may provide the pathway for the accumulation in aquatic environments (Cowie and Heges, 1984; Cowie et al, 1995; Benner et al, 1992).

Else, bacteria play an important role in regulation of the rate of organic matter mineralization, nutrient cycling, and energy transfer in aquatic environments and demineralization or the turnover of carbohydrates has been used to evaluate the efficiency of the microbial community and the liability of the organic carbon in coastal waters (Azam and Worden, 2004; Lennon, 2007).

Considering that total number of bacteria on Earth, the largest proportion of bacterial cells presumably reside in oceanic subsurface rather than in terrestrial areas (3.5×10^{30} and $0.25\text{--}2.5 \times 10^{30}$ respectively) and therefore bacterial cells are estimated to contain, in total, 350–550 Pg of carbon, up to 60–100% of the total carbon found in plants, as well as large amounts of nitrogen and phosphorous (Whitman *et al* 1998). Thus, despite their modest size as individuals, as a group these organisms not only contribute to the flow of nutrients worldwide, but may also constitute a significant proportion of the nutrients in living biomass (Horner-Devine et al, 2009).

In estuarine areas, biodegradation by heterotrophic bacteria for Organic composition changed rapidly along the estuary, showing a selected removal of carbohydrates and amino acids within the DOC pool in the upper reach and mixing zone, and an autotrophic source of Particular Carbohydrates (PCHO) in the lower estuary, which gave an insight into the DOC estuarine process (He et al, 2010).

For hydrothermal marine compartment, heterotrophic thermophiles in culture far exceeds the number of their autotrophic counterparts, and many are known to metabolize carbohydrates *via* fermentation or respiration nevertheless, the energetic associated with these high-temperature microbial processes have been all but ignored (Amend et al, 2001).

In Coastal marine regions, sediments receive organic matter from a variety of sources, and its reactivity towards anaerobic fermentation and respiration is determined by the relative content of e.g. carbohydrates, lipids and proteins (Kristensen et al, 1995; Kristensen and Holmer, 2001). Proteins and labile carbohydrates derived from algae usually constitute the main organic sources for marine sediments, but coastal areas may also receive detritus rich in structural carbohydrates (e.g. cellulose) from vascular plants (Cowie and Hedges, 1992; Leeuw and Largeau, 1993; Kristensen, 1994). Furthermore, sediments of anthropogenic point sources, such as marine fish farms, may concentrate high loads of protein and lipid rich materials of animal origin (Ackefors and Enell, 1994). Proteins and simple carbohydrates are

generally labile towards anaerobic degradation (Arnosti and Repeta, 1994; Holmer and Kristensen, 1994; Arnosti and Holmer, 1999), whereas anaerobic degradation of lipids and structural carbohydrates occur at much lower rates (Boetius and Lochte, 1996; Canuel and Martens, 1996). This difference is most likely caused by differential efficiency of extra-cellular enzymes produced by anaerobic fermenting bacteria (Valdemarsen et al, 2010)

2.1. Bio mineralization of carbohydrates by marine bacteria and production of carbohydrases

Bacteria are the most abundant and most important biological component involved in the transformation and mineralization of organic matter in the biosphere (Cho et al, 1988; , Pomeroy et al, 1991). Heterotrophic bacteria contribute to the cycles of nutrients and carbon in two major ways: by the production of new bacterial biomass (secondary production) and by the re mineralization of organic carbon and nutrients. Understanding this dual character of planktonic bacteria in aquatic ecosystems is a central paradigm of contemporary microbial ecology (Billen et al, 1984; Ducklow et al, 1992 and Del Giorgio and Cole, 1998).

Further, bacteria are capable of out competing, other organisms for organic compounds at low concentrations since they possess high substrate affinities, surface-to volume ratios, and metabolic rates. Natural DOM sources are heterogeneous mixtures of compounds, and bacteria can utilize several molecules simultaneously (Bott et Kaplan, 1985).

Extracellular enzymes, term generally utilised to define enzymes located outside the cytoplasmic membrane, are important catalyst in the decomposition of particulate organic matter (POM) and dissolved organic matter (DOM) including marine snow (Cho and Azam, 1988; Smith et al, 1992) and only small molecules, around 600 Da (Weiss et al, 1991), can be transported across the bacterial membrane and thus bacteria secrete enzymes to hydrolyze high-molecular-weight organic matter. Moreover extra cellular enzymes play an important role in nutrient cycling as they may be produced in order to acquire the limiting nutrient (Hoppe, 1983; Zoppini et al, 2010).

In the aggregates, the activity of the extracellular enzymes transforms non utilizable POM in DOM, thus bringing small nutrient molecules into solution (Smith et al, 1992; Simon et al, 2002). The metabolic activity of colonizing bacteria drives sinking particulate organic matter into non sinking DOM (Azam and Long, 2001; Smith et al, 1992) fueling free-living bacteria (Kiørboe and Jackson, 2001). This bun coupled solubilisation causes rapid POMYDOM transition with important biogeochemical implications for carbon flux in the oceans (Smith et al, 1992; Simon et al, 2002).

Further, almost heterotrophic bacteria have developed several mechanisms which allow the preferred utilization of the most efficiently metabolised carbohydrates when are exposed to a mixture of carbon sources. Interestingly, similar mechanisms are used by some pathogens to control various steps of their infection process. The efficient metabolism of a carbon source might serve as signal for proper fitness. Alternatively, the presence of a specific carbon source might indicate to bacterial cells that they thrive in infection-related organs,

tissues or cells and that specific virulence genes should be turned on or switched off. Frequently, virulence gene regulators are affected by changes in carbon source availability. Thus, the activity of PrfA, the major virulence regulator in *Listeria monocytogenes*, seems to be controlled by the phosphorylation state of phosphotransferase system (PTS) components. In *Vibrio cholerae* synthesis of HapR, which regulates the expression of genes required for motility, is controlled via the Crp/cAMP CCR mechanism, whereas synthesis of *Salmonella enterica* HilE, which represses genes in a pathogenicity island, is regulated by the carbohydrate responsive, PTS-controlled Mlc (Poncet et al, 2009).

2.2. Use of carbohydrates mineralization for marine bacteria identification

The metabolic diversity of bacteria is perhaps as remarkable as their taxonomic and evolutionary diversity. Although culture-based studies are limited in their ability to estimate bacterial diversity, and several studies demonstrated varieties of modes of energy conversion of bacteria and wide ranges of substrate uses and metabolic pathways.). The flexibility of bacterial metabolism is similarly best illustrated by their abilities to degrade xenobiotic compounds such as malathion (an insecticide) and 2,4,5-trichlorophenoxyacetic acid (a herbicide), which are toxic to many other organisms (Brock et al, 1987 and Horner et al, 2003).

Metabolic assays mainly based on carbohydrates degradation on either aerobic or anaerobic conditions still effectively used to identify environmental bacteria besides molecular methods. Among these methods, the most widely used commercial identification systems are the API, the Biolog systems and more recently Microgen systems (Awong-Taylor et al, 2007).

The API 20 system included profiles of approximately 40 groups and species of bacteria in addition to the *Enterobacteriaceae* group in the API data set segregated mainly on assimilation of several carbohydrates with varied number depending on the type of API test kit used. (MacDonell et al, 1982; Bertone et al, 1996).

The Biolog identification system (Biolog, Inc., Hayward, CA, USA) is a bacterial identification method that establishes identifications based on the exchange of electrons generated during respiration, leading subsequently to tetrazolium-based color changes. This system tests the ability of micro organisms to oxidize a panel of 95 different carbon sources (Truu et al, 1999).

3. Effects of starvation conditions in carbohydrates metabolism

In their natural environment especially in marine ecosystems, bacteria are frequently exposed to major changes in growth conditions. These changes can include temperature, salt concentrations, essential nutrients, oxygen supply, pressure...etc. In order to sustain these changes, bacteria have developed mechanisms which allow them to adapt to drastic environmental alterations. Mainly, response to carbon source availability seems to be of special importance, because many bacteria dispose of more than one mechanism to adapt Carbon Metabolism and Virulence to changes in carbohydrate composition. These mechanisms include induction of specific carbohydrate transport and utilization system by

the presence of the corresponding carbon source and their repression when a more efficiently utilizable carbohydrate is present in addition. The latter phenomenon is called carbon catabolite repression (CCR) and is often mediated by more than one signal transduction pathway, which can include regulation of transcriptional activators or repressors, anti terminators, carbohydrate transporters or metabolic enzymes (Poncet et al, 2009).

3.1. *In vivo* starvation effect on catabolic carbohydrates profiles of *E.coli*

Escherichia coli, a fecal coliform, was found to survive for longer periods of time in non sterile natural seawater when sediment material was present than in seawater without sediments, and growth was observed to occur by one occasion (Gerba et al, 1976). This enteric bacterium was found to increase rapidly in number in autoclaved natural seawater and autoclaved sediment taken from areas receiving domestic wastes, even when the seawater had salinities as high as 34 g/kg. Else, longer survival of *E. coli* in sediment is revealed and attributed to greater content of organic matter present in sediment than the seawater.

Experimental, mainly *in vitro*, studies have been performed on the survival of different enteric bacterial species in seawater and results showed solar radiation is the most adverse factor for enteric bacteria. Its effect was found to be wavelength-dependent (most inactivation caused by UVB solar spectrum with a range of dose from 0.98 to 4 kJ/m²) and thus restricted to shallow (45 to 90 cm) or clear waters (Rozen and Belkin, 2001; Sinton et al, 1994; Villarino et al, 2003; Whitman et al, 2004). Moreover, in cases of its release from wastewater into marine environment, enteric bacteria are subject to osmotic stress with effects depending on water salinity (a high loss of viability for *E. coli* occurred for salinities between 15 and 30 g/L) (Anderson et al, 1979; Pereira and Alcantara, 1993). Most coastal waters provide low concentrations of nutrients essential to the growth and survival of enteric bacteria (Barcina et al, 1997). These stresses induce different resistance mechanisms for at least part of the cellular population (Martin et al 1998; Rozen et al 2002). As a consequence, different cellular states occur in the stressed population, including the viable but nonculturable state (VBNC). The ability of such cells to recover their growth capacity is still under debate (Arana et al, 2007).

Thus, we investigated survival and virulence of *Escherichia coli* strains (*E.coli* O126:B16 and *E.coli* O55:B5) exposed to natural conditions in brackish water by incubation in water microcosms in the Bizerte lagoon in Northern Tunisia and exposed for 12 days to natural sunlight in June (231 to 386 W/m², 26.61 uC, 30 g/L) and in April (227 to 330 W/m², 17.61 uC, 27 g/L) or maintained in darkness for 21 days (17.61 uC, 27 g/L). The results revealed sunlight as the most significant inactivating factor (decrease of 3 Ulog within 48 hours for the two strains) compared to salinity and temperature (in darkness). Survival time of the strains was prolonged as they were maintained in darkness. Local strain (*E. coli* O55:B5) showed better survival capacity (T₉₀ 552 hours) than *E. coli* O126:B16 (T₉₀ 511 h).

For both, modifications were noted only for some metabolic activities of carbohydrates hydrolysis (Table 1). Thus, *E. coli* O126:B16 culturable cells lost the ability to assimilate

amygdalin, and *E. coli* O55:B5 lost ability to assimilate melibiose, amygdalin, rhamnose, and saccharose.

Although, in previous data revealed exposure increase to sunlight, particularly UV radiation, alleviates some of the negative effects of altered organic matter quality through selective photolytic degradation. Both ultraviolet radiation (UV-B, 280–320 nm and UV-A, 320–400 nm) and visible light (photo synthetically active radiation, PAR, 400–700 nm) and induce major photolytic changes to complex organic molecules and generate large quantities of readily utilizable substrates for bacterial metabolism (Lindell et al 1995, Espeland & Wetzel 2001, Tietjen & Wetzel 2003). Thus, rates of carbon and nutrient cycling are often accelerated, stimulating ecosystem productivity. Interestingly, this enhanced microbial respiration may lead to increased CO₂ production and evasion to the atmosphere, creating a potential positive feedback to atmospheric CO₂ concentrations (Wetzel 2001; Lingo et al, 2007)

Strains	ONPG	ADH	LDC	ODC	CIT	H ₂ S	URE	TDA	IND	VP	GEL	GLU	MAN	INO	SOR	RHA	SAC	MEL	AMY	ARA	OX	CAT
Sunlight incubation (April and June) (n=2)																						
<i>E.coli</i> O126B16 ^a	+	+	+	+	-	-	-	+	+	-	-	+	+	-	+	-	-	-	+	+	-	+
after 3 days	+	+	+	+	-	-	-	+	+	-	-	+	+	-	+	-	-	-	+	+	-	+
<i>E.coli</i> O55B5 ^a	+	+	+	+	-	-	-	+	+	-	-	+	+	-	+	+	+	+	+	+	-	+
after 12 days	+	+	+	+	-	-	-	+	+	-	-	+	+	-	+	+	+	+	+	+	-	+
Darkness incubation (n=2)																						
<i>E.coli</i> O126B16 ^a	+	+	+	+	-	-	-	+	+	-	-	+	+	-	+	-	-	-	+	+	-	+
after 2 days	+	+	+	+	-	-	-	+	+	-	-	+	+	-	+	-	-	-	(-)	+	-	+
after 21 days	+	+	+	+	-	-	-	+	+	-	-	+	+	-	+	-	-	-	(-)	+	-	+
<i>E.coli</i> O55B5 ^a	+	+	+	+	-	-	-	+	+	-	-	+	+	-	+	+	+	+	+	+	-	+
after 2 days	+	+	+	+	-	-	-	+	+	-	-	+	+	-	+	+	+	(-)	(-)	+	-	+
after 16 days	+	+	+	+	-	-	-	+	+	-	-	+	+	-	+	(-)	(-)	(-)	(-)	+	-	+
after 21 days	+	+	+	+	-	-	-	+	+	-	-	+	+	-	+	(-)	(-)	(-)	(-)	+	-	+

(+) positive reaction; (-) : negative reaction; *n*: number of colony tested. ONPG: β-galactosidase, ADH: arginine dihydrolase, ODC: ornithine decarboxylase, CIT: use of citrate, URE: urease, TDA: tryptophane desaminase, IND: indole production, VP: acetoin production, GEL: gelatinase, GLU: glucose assimilation, MAN: mannitol, SOR: sorbitol, RHA: rhamnose, SAC: saccharose, MEL: melibiose, ARA: arabinose

Table 1. Characteristics of *E.coli* O55B5 and *E.coli* O126B16 before and during seawater incubation

3.2. *In vitro* starvation effect on catabolic carbohydrates profiles of *E.coli*

Depending on the nature of aquatic environmental receptor (seawater, wastewater or brackishwater), different physiological, metabolic and pathogenic modifications related to stress conditions have been reported by much studies (Gauthier *et al*, 1993; Dupray and Derrien, 1995; Baleux *et al*, 1998; Troussellier *et al*, 1998; Monfort and Baleux, 1999). And different studies showed that adaptation forms of enterobacteria to drastic conditions

occurring gradually during discharging in wastewater (Dupray and Derrien, 1995), or during the transfer from a previous culture to a saline medium or to mixed wastewater/seawater medium (Munro *et al*, 1987), allowing better survival in seawater conditions. In this report, *E. coli* was chosen as a representative strain of Thermotolerant Coliform (TTC), used as fecal contamination indicator. Most of *E. coli* strains are normal intestinal flora components; however, certain pathogenic strains are important disease factors. Those pathogenic strains are classified into at least six distinct groups: enteropathogenic *E.coli* (EPEC), enterotoxigenic *E.coli* (ETEC), enteroinvasive *E.coli* (EIEC), diffuse-adhering *E.coli* (DAEC), enteroaggregative *E.coli* (EAEC), enterohemorrhagic *E.coli* (EHEC) (Schroeder *et al*, 2004). Thus, we studied the behavior of these 6 different types of *E.coli* when discharged in seawater with or without a previous incubation in wastewater microcosms. Else, we examined their metabolic, antibiotic and virulence resistance patterns after exposure to salinity gradient.

Therefore, the studies were carried out using membrane chambers with 120ml capacity, constituted by assembly of glass tubes allowing dissolved matter to penetrate, through 0.2 µm- pore- size, 25 mm-diameter filters, fixed between two Teflon joints. For each *E.coli* strains two types of microcosms were prepared: one type was inoculated by seawater and the second with wastewater.

For microcosm inoculation, the seawater used was brought from Bizerte lagoon (northern part in Tunisia of Tunis). Wastewater was brought from sewage treatment station (near Tunis town).

The results obtained (Table 2) showed most important modifications noted for the strains previously inoculated in wastewater and O126B16 *E.coli* strain lost ability of fermenting glucose and mannitol, producing indole and to use citrate, but it acquired urease and gelatinase activities. All the other pathogenic *E.coli* strains maintained their major metabolic characteristics except for the ability to use citrate and to produce indole.

In further study, we compared the behavior of *E.coli* (ATCC 14948), *Vibrio paraheamolyticus* and *Salmonella Typhymurium* (ATCC 17802) when discharged in seawater with or without previous incubation in wastewater microcosms and we examined mainly their metabolic, patterns after exposure to salinity gradient. The results obtained showed metabolic profiles changes both for *E.coli* and *Salmonella Typhymurium* and loss for production of several carbolases mainly glucose, mannose, rhamnase, sorbitol, sucrose, melibiose and arabinose hydrolases whereas for *Vibrio paraheamolyticus* no changes were observed.

Similar results were found previously by Ben Kahla –Nakbi *et al* (2007) for *V. alginolyticus* strains incubated in seawater microcosms for long period of starvation and for what the result of metabolic carbohydrates profiles tested by results of API system tests, showed no modification. These findings were explained by preparation of cells to enter into VBNC state characterized by low metabolic rate, reduction in size, morphological changes or synthesis of specific proteins. This adaptation strategy developed by bacteria in aquatic environments allows them to survive during long period of time. Modifications of cellular morphology should be related to modifications with carbohydrates metabolites contained in parietal membranes for these species.

Strains	API 20E*															
	ONPG	ADH	ODC	CIT	URE	TDA	IND	VP	GEL	GLU	MAN	SOR	RHA	SAC	MEL	ARA
O126B16	+	+	-	+	-	+	+	-	-	+	+	+	+	+	+	+
S.W.	-	-	-	-	-	+	+	-	-	+	+	+	-	-	+	+
W.W.	-	-	-	+	+	+	-	+	+	-	-	-	-	+	-	-
ECEAgg	+	-	+	+	-	+	+	-	-	+	+	+	+	+	+	+
S.W.	+	-	+	-	-	+	-	-	-	+	+	+	+	+	+	+
W.W.	+	+	+	-	-	+	-	-	-	+	+	+	+	+	+	+
ECEI	+	-	+	+	-	+	+	-	-	+	+	+	+	-	+	+
S.W.	+	-	+	-	-	+	+	-	-	+	+	+	+	-	+	+
W.W.	+	+	+	+	-	-	-	-	-	+	+	+	-	-	-	-
ECEH	+	-	+	+	-	+	+	-	-	+	+	+	+	-	+	+
S.W.	+	-	-	-	-	+	+	+	-	+	+	+	+	-	-	+
W.W.	+	+	-	+	-	+	-	-	-	+	+	+	+	-	-	+
ECEP	+	-	+	+	-	+	+	-	-	+	+	+	+	-	+	+
S.W.	+	-	-	-	-	+	+	+	-	+	+	+	+	-	-	+
W.W.	+	-	-	+	-	+	-	-	-	+	+	+	+	-	-	+
ECET	+	-	+	+	-	+	+	-	-	+	+	+	+	+	+	+
S.W.	+	-	+	-	-	+	+	+	-	+	+	+	+	+	+	+
W.W.	+	+	+	+	-	+	-	-	-	+	+	+	-	+	-	+

*ONPG: β -galactosidase, ADH: arginine dihydrolase, ODC: ornithine decarboxylase, CIT: use of citrate, URE: urease, TDA: tryptophane desaminase, IND: indole production, VP: acetoin production, GEL: gelatinase, GLU: glucose, MAN: mannitol, SOR: sorbitol, RHA: rhamnose, SAC: saccharose, MEL: melibiose, ARA: arabinose.

Table 2. Biochemical patterns. Modifications of some biochemical characters on APi 20E strips of different *E.coli* strains after survival in seawater with previous incubation in seawater (SW) or in wastewater (WW). O126B16 (reference *E.coli* strain), EaggEC (Enteraggregative *E.coli* strain), EIEC (Enteroinvasive *E.coli* strain), EHEC (Enterohemorrhagic *E.coli* strain), EPEC (Enteropathogenic *E.coli* strain), ETEC (Enterotoxigenic *E.coli* strain).

Keymer et al (2007) described similar results for *V.cholerae* and pointed that environmental parameters, measured *in situ* during sample collection, should correlated to the presence of specific dispensable genes and metabolic capabilities, including utilization of mannose, sialic acid, citrate, and chitosan oligosaccharides. Thus, gene content identified and metabolic pathways that are likely selected for in certain coastal environments and may influence *V. cholerae* population structure in aquatic environments.

Our previous data described *A.hydrophila* cultured for 30 days in marine water microcosms which maintained the ability to metabolize all the carbohydrates and continued to produce several enzymes with differences of its homologous cultured in waste water previous to incubation in marine water microcosm which failed to produce numerous enzymes and metabolize carbohydrates (El Mejri et al, 2008). Thus, for marine heterotrophic bacterial populations as *Vibrio sp* and *Aeromonas sp* halotolerant conserve major potential of their carbohydrate metabolism.

For high salinity or halotolerant bacteria, many strains that grew fermentatively on carbohydrates in the presence of air (determined by the acidification of the medium) grew especially well on glucose, totally by substrate- level phosphorylation in the absence of air. Tomlinson and Hochstein (1976) demonstrated that O₂ consumption by *Halobacterium saccharovororum* in presence of glucose was 18% of the theoretical amount required for its

complete oxidation it was for 83% in presence of galactose or fructose which like galactose, requires more oxygen for its oxidation and therefore was generally a poorer substrate for fermentation (Javor et al, 1984). The measurement of O₂ consumption, growth rates, and fermentation products in the presence of glycerol, pyruvate, and acetate by strains that demonstrated relatively good anaerobic growth on these substrates would provide further evidence of the importance of fermentative metabolism in extreme halophiles.

In the water column, biofilm microenvironments in suspended flocks may form a stabilizing refuge that enhances the survival and propagation of pathogenic (i.e., disease-causing) bacteria entering coastal waters from terrestrial and freshwater sources. The EPS matrix offers microbial cells a tremendous potential for resiliency during periods of stress, and may enhance the overall physiological activities of bacteria as it's emphasized by Giller et al (1994) that influences small-scale microbial biofilms must be addressed in understanding larger-scale processes within inter tidal systems. Intertidal systems are a key interface of the ocean, atmosphere, and terrestrial environments, and as such, are characterized by frequent fluctuations in temperature, ion concentration, desiccation, UV-irradiation, and wave action. The relative frequency of these fluctuations poses both physical and biochemical challenges to micro organisms which inhabit this environments such estuarine areas. The characteristics and intensities of such stresses may vary substantially (Decho et al, 2000).

Therefore, in estuarine region with with salinity of 35 ppt it is been reported that the growth rate of lingo-cellulose-degrading populations (50%) is inhibited by NaCl (Liu and Boone, 1991). According to Park et al (2006) the high salinity seems to significantly reduce ecto enzyme activities. Thus, it should be possible that type of bacterial population found at particular station can be influenced by salinity changes. Also, it has been reported in most of the aquatic environments there is a significant relationship between extracellular enzyme activities, their corresponding substrates (polymers) and their hydrolysis products (monomers) (Münster et al, 1992). Thus, presence of cellulolytic and hemi-cellulolytic bacteria in one particular region and their absence in other region should indicated presence of wide-ranging organic matter in each region (Khanderparker et al, 2011). Previous research has also put forth a widely accepted concept that hydrolytic enzymes are induced by presence of polymeric substrates (Chrost, 1991; Vetter and Deming, 1999). It was observed that the genus *Bacillus* and *Vibrio* were the dominant hemicellulase and cellulase producers in both the estuaries. According to Ruger (1989) the genus *Bacillus* comprised phylogenetically and phenotypically diverse species, which are ubiquitous in terrestrial and fresh water habitats and are also widely distributed in seawater ecosystems. While the family of *Vibrionaceae* represent the most important bacterial autochthonous groups in marine environments. Members of this family often predominate in the bacterial flora of seawater, plankton, and fish. In the West Pacific Ocean, , *Vibriosis* accounted for nearly 80% of the bacterial population in surface seawater (Simidu et al, 1980) and *Vibrio* phylogenetic diversity of their culturable forms in Bulgarian hot springs as described along with their abilities to metabolize carbohydrates (Derekova et al, 2008)..

For Moari et al, (2011), marine models tested showed, prokaryotic abundance in superficial marine sediments controlled by organic trophic resources, while in sub-surface sediments, prokaryotic activities and abundance were driven by environmental factors and predatory pressure, suggesting that the shift in prokaryotic community structure could be coupled to a change in life-style of microbial assemblages.

Previously, Bouvier et al, (2002), discussed the relationship between the compositional succession and changes in single-cell metabolic activities in the Choptank River, and suggested that profound phylogenetic shifts were linked to cell stress, loss of activity, and death (Del Giorgio and Bouvier, 2002). Similarly both types of succession, i.e., activation/inactivation and replacement, occurred simultaneously, and understanding the relative importance of such processes in determining bacterial succession and community composition remain to be clarified considering environmental conditions that trigger bacterial succession observed within the mixing areas. .

4. Conclusion

Carbohydrates metabolism still basic to marine microbiota mainly for autochthonous communities as *Vibrio*, *Aeromonas* and *Bacillus* genus with high potential of degradation and considerable supply of big amounts of different carbolases and pathways which should take more attention in order to be clarified.

Increase in significant amounts of organic materials are being exchanged between the land and marine ecosystems with higher amounts of dissolved carbohydrates provided especially the polymeric substrates which represent the most abundant molecules used by marine bacteria as structural and /or storage compounds under extreme variations in environmental parametres and biological marine habitats.

The intrinsic ability of bacterial cells to enter in to viable but non culturable state (VBNC) characterized by low metabolic rate and morphological modifications (reduction in cell size and membrane alterations) should be related with carbohydrates metabolism and change observed experimentally. Therefore, investigations in carbohydrates metabolites which continue going on will certainly elucidate more molecular adaptations forms of bacteria (for mainly enteric species) in variable marine ecosystems .

Author details

Monia El Bour

National Institute of Sea Sciences and Technologies, Salammbô, Tunisia

5. References

- Ackefors H. & Enell M. (1994) The release of nutrients and organic matter from aquaculture systems in Nordic countries. *Journal of Applied Ichthyology*. Vol.10, pp 225–241. Ackefors and Enell, 1994

- Allredge AL.(2000). Interstitial dissolved organic carbon (DOC) concentrations within sinking marine aggregates and their potential contribution to carbon flux. *Limnologie and Oceanography*. Vol45, pp1245–1253
- Aluwihare, L.I., Repeta, D.J. & Chen, R.F., (1997). A major biopolymeric component to dissolved organic carbon in surface sea water. *Nature*. Vol387, pp166–169.
- Amend J.P. & Plyasunov A.V. (2001). Carbohydrates in thermophile metabolism: Calculation of the standard molal thermodynamic properties of aqueous pentoses and hexoses at elevated temperatures and pressures. *Cosmochimica Acta*. Vol 65, pp. 3901-3917.
- Anderson, I.C., Rhodes, M.W., Kator, H.I., (1979). Sublethal stress in *Escherichia coli*: a function of salinity. *Applied and Environmental Microbiology* Vol.38, pp1147-1152. Pereira and Alcantara, 1993
- Arnosti C. and Repeta D. J. (1994). Oligosaccharide degradation by anaerobic marine bacteria : characterization of an experimental system to study polymer degradation in sediments. *Limnology and Oceanography*. Vol 39, pp1865–1877.
- Arnosti C. and Holmer M. (1999). Carbohydrate dynamics and contributions to the carbon budget of an organic-rich coastal sediment. *Geochimistry and. Cosmochimistry*. Acta. Vol. 63, pp 393–403.
- Arnosti C.(2000). Substrate specificity in polysaccharide hydrolysis: contrast between bottom water and sediments. *Limnologie and Oceanography*. Vol.45, pp1112– 9.
- Azam F, & Long R. (2001). Sea snow microcosm. *Nature*. Vol. 8, pp414:495..
- Azam, F. & Worden, A.Z., (2004). Microbes, molecules, and marine ecosystems. *Science*. Vol. 303, pp 1622-1624.
- Baleux, B., Caro, A., Lesne, J., Got P., Binard, S., Delpeuch, B., 1998. Survie et maintien de la virulence de *Salmonella Typhimurium* VNC exposée simultanément à trois facteurs stressants expérimentaux. *Océanologica Acta* Vol.21, pp939-950.
- Barcina I., LeBaron P., & Vives-Rego J. (1997). Survival of allochthonous bacteria in aquatic systems: A biological approach. *FEMS Microbiology and Ecology*. Vol.23, pp 1–9.
- Ben Kahla-Nakbi A, Besbes A, Chaieb K, Rouabhia M, & Bakhrouf A. (2007). *Marine Environmental Research*, Vol. 64,pp 469
- Benner, R., Pakulski, J.D., McCarty, M., Hedges, J.I. & Hatcher, P.G. (1992). Bulk chemical characteristics of dissolved organic matter in the ocean. *Science* Vol. 255, pp 1561-1564.
- Billen G. (1984). Heterotrophic utilization and regeneration of nitrogen. In *Heterotrophic Activity in the Sea*, ed. JE Hobbie, P.JleB Williams, pp. 313–55. New York: Plenum
- Boetius, A., Ravensschlag, K., Schubert, C.J., Rickert, D., Widdel, F., Gieseke, A., Amann, R., Jørgensen, B.B., Witte, U. & Pfannkuche, O. (2000). A marine microbial consortium apparently mediating anaerobic oxidation of methane. *Nature*. Vol. 407, pp 623-626. .
- Bott T.L.& Kaplan L.(1985). Bacterial biomass, metabolic state and activity in stream sediments: relation to environmental variables and multiple assay comparisons. *Applied and Environmental Microbiology*. Vol. 50, pp 508-522.
- Bouvier T.C. & Del Giorgio P.A.(2002). Compositional changes in free-living bacterial communities along a salinity gradient in two temperate estuaries. *Limnologie and Oceanography*. Vol. 2002, pp. 453-470.

- Brock, T. D. (1987). The study of microorganisms *in situ*: progress and problems. *Symposium Of Society Of. General Microbiology* Vol.. 41, pp1–17.
- Burdige, D.J. & Zheng, S.L.(1998). The biogeochemical cycling of dissolved organic nitrogen in estuarine sediments. *Limnology and Oceanography*. Vol.43, pp1796–1813.
- Canuel E. A. and Martens C. S. (1996). Reactivity of recently deposited organic matter: degradation of lipid compounds near the sediment–water interface. *Geochimistry and Cosmochimistry Acta* 60, pp 1793–1806.
- Cho BC & Azam F.(1988) Major role of bacteria in biogeochemical fluxes in the ocean's interior. *Nature*. Vol.332, pp441–3.
- Chrost, R.J., 1991. Environmental control of the synthesis and activity of aquatic microbial ectoenzymes. In: *Microbial Enzymes in Aquatic Environments*. Springer, pp. 29-59.
- Cowie G. L. and Hedges J. I. (1992) Sources and reactivities of amino-acids in a coastal marine environment. *Limnologie Oceanography*. Vol.37, pp703–724.
- Cowie GL, Heges JI, Prahl FG & De Lange GJ.(1995) Elemental and major biochemical changes across an oxidation front in a relict turbidite: an oxygen effect. *Geochimistry Cosmochim Acta*. Vol.59. pp33– 46..
- Del Giorgio P.A. & Cole J.J.(1998). Bacterial growth efficiency in natural aquatic systems. *Annual Review of Ecology*. Vol. 29, pp 503-541.
- Decho A.W.,(2000). Microbial biofilms in intertidal systems: an overview. *Continental Shelf Research*. Vol.20, pp. 1257-1273.
- Derekova, A., Mandeva, R., Kambourova, M., (2008). Phylogenetic diversity of thermophilic carbohydrate degrading Bacilli from Bulgarian hot springs. *World Journal of Microbiology and Biotechnology*. Vol. 24, pp 1697-1702..
- Druffel, E.R.M., Williams, P.M., Bauer, J.E. & Ertel, J.R. (1992). Cycling of dissolved and particulate organic matter in the open ocean. *Journal of Geophysical Research*. Vol.97, pp15639–15659.
- Ducklow HW & Carlson CA. (1992). Oceanic bacterial production *Advances in Microbiol Ecology*. Vol.12, pp113–81
- Dupray E., Derrien A. (1995). Influence du passage de *Salmonella* spp. Et *Escherichia coli* en eaux usées sur leur survie ulterieure en eau de mer. *Water Research*. Vol. 29, pp1005-1011.
- Espeland, E. M. & Wetzel, R. G., (2001). Complexation, stabilization and UV photolysis of extracellular and surface-bound glucosidase and alkaline phosphatase: implications for biofilm microbiota. – *Microbiology and Ecology*. Vol. 42, pp 572–585.
- Gauthier M.J., Munro P.M., Flatau G.N., Clément R.L. & Breittmayer V.A. (1993). Nouvelles perspectives sur l'adaptation des entérobactéries dans le milieu marin. *Marine Life*, Vol.3, pp1-18.
- Gerba C.P. & McLeod J.S. (1976).Effect of Sediments on the Survival of *Escherichia coli* in Marine Waters. *Applied and Environmental Microbiology*. Vol.32, pp 114-120.
- Giller, P.S., Hildrew, A.G., Ralaelli, D.G. (Eds). (1994). *Aquatic Ecology: Scale, Pattern and Processes*. Blackwell Publishers, Oxford, 649 pp..
- Gueuen, C., Guo, L., Wang, D., Tanaka, N., Hung, C.C. (2006). Chemical characteristics and origin of dissolved organic matter in the Yukon River. *Biochemistry* Vol. 77, pp139–155.

- Holmer M. & Kristensen E. (1994) Organic-matter mineralization in an organic-rich sediment – experimental stimulation of sulfate reduction by fish food pellets. *FEMS Microbiology and Ecology*. Vol.14, pp 33–44.
- Hedges, J.I., Keil, R.G., Benner, R., 1997. What happens to terrestrial organic matter in the ocean? *Organic Geochemistry*. Vol. 27, pp195–212.
- Hoppe HG. (1983). Significance of exoenzymatic activities in the ecology of brackish water: measurements by means of methylumbelliferylsubstrates. *Marine Ecological Progressive Series* Vol.11, pp 299–308.
- He B., Dai M., Zhai W., Wng L., Wang L., Chen J., Lin J., Han A. & Xu Y. (2010). Distribution, degradation and dynamics of dissolved organic carbon and its major compound classes in the Pearl River estuary, China. *Marine Chemistry*. Vol. 119, pp 52–64.
- Horner-Devine M.C., Carney K.M. & Bohannan .M.J. (2004). An ecological perspective on bacterial biodiversity. *Proceedings of the Royal Society of London*. Vol.271, 113-122
- Hung, C.-C., Guo, L., Santschi, P.H., Alvarado-Quiroz, N. & Haye, J.M. (2003). Distributions of carbohydrate species in the Gulf of Mexico. *Marine Chemistry*. Vol. 81, pp119–135.
- Javor B.J.,(1984).Growth Potential of Halophilic Bacteria Isolated from Solar Salt Environments: Carbon Sources and Salt Requirements. *Applied and Environmental Microbiology*. Vol.48, pp. 352-360.
- Kaltenbock E & Herndl GJ.(1992). Ecology of amorphous aggregates (marine snow) in the northern Adriatic Sea IV Dissolved nutrients and autotrophic community associated with marine snow. *Marine Ecological Progressive Series* Vol.87, pp147–59.
- Keymer D.P., Miller M.C., Schoolnik G.K, Boehm A.B. (2007). Genomic and phenotypic diversity of costal *Vibrio cholerae* strains is linked to environmental factors. *Applied and Environmental Microbiology*. Vol 73, pp 3705-3714.
- Khanderparker R.,Verna P., Meena R.M. & Deobagkar D.D. (2011). Phylogenetic diversity of carbohydrate degrading culturable bacteria from Mandovi and Zuari estuaries, Goa, west coast of India. *Estuarine, Coastal and Shelf Science*. Vol. 95, pp 359-366.
- Kiørboe T. & Jackson GA.(2001). Marine snow, organic solute plumes, and optimal chemosensory behaviour of bacteria. *Limnology Oceanography*. Vol.46, pp1309– 18.
- Kristensen E. (1994). Decomposition of macroalgae, vascular plants and sediment detritus in seawater: use of step wise thermogravimetry. *Biogeochemistry* Vol.26, pp1–24.
- Kristensen E., Ahmed S. I. and Devol A. H. (1995) Aerobic and anaerobic decomposition of organic matter in marine sediment: which is fastest? *Limnology and Oceanography*. Vol 40, pp1430–1437.
- Kristensen E. and Holmer M. (2001) Decomposition of plant materials in marine sediment exposed to different electron acceptors (O₂, NO₃, and SO₄²⁻), with emphasis on substrate origin, degradation kinetics, and the role of sulfate reduction. *Geochimistry and Cosmochimistry*. Acta Vol.65, pp419–433.
- Lango Z., Leech D.M., Wetzel R.G., (2007). Indirect effects of elevated atmospheric CO₂ and solar radiation on the growth of culturable bacteria. *Fundamental and Applied Limnology Archiv Für Hydrobiologie*, Vol. 168, pp. 327-333..
- Lennon, J.T., 2007. Diversity and metabolism of marine bacteria cultivated on dissolved DNA. *Applied and Environmental Microbiology* Vol. 73, pp 2799-2805.

- Leeuw J. W. D. and Largeau C. (1993). A review of macromolecular organic compounds that comprise living organisms and their role in kerogen, coal and petroleum formation. In Organic (eds. M. H. Engel and S. A. Macko). Plenum Press, New York.
- Lindell, M. J., Granéli, H. W. & Tranvik, L. J., (1995). Enhanced bacterial growth in response to photochemical transformation of dissolved organic matter. *Limnology and Oceanography*, Vol. 40, pp 195–199.
- Liu, Y., Boone, D.R., 1991. Effects of salinity on methanogenic decomposition. *Bioresource Technology*. Vol. 35, pp 271-274.
- Martin, Y., Troussellier, M., Bonnefont, J-L., 1998. Adaptive responses of *Escherichia coli* to marine environmental stresses: a modelling approach based on viability and dormancy concepts. *Oceanologica Acta* 21, 951-964.
- Meybeck, M., (1982). Carbon, nitrogen, and phosphorus transport by world rivers. *American Journal of Science*. Vol.282, pp401–450.
- Middelboe, M., Borch, N.H., Kirchman, D.L., 1995. Bacterial utilization of dissolved free amino acids, dissolved combined amino acids and ammonium in the Delaware Bay estuary: effects of carbon and nitrogen limitation. *Marine Ecology Progress Series*. Vol.28, pp109–120.
- Molari M., Giovannelli D., D’Errico G. & Manini E. (2011). Factors influencing prokaryotic community structure composition in sub-surface coastal sediments. *Estuarine, Coastal and Shelf Science*. Vol. xxx, pp1-8.
- Monfort P., Baleux B. (1999). Bactéries viables non cultivables: réalité et conséquences. *Bulletin de la Société Française de Microbiologie*, Vol.14, pp 201-207.
- Murrell, M.C. & Hollibaugh, J.T. (2000). Distribution and composition of dissolved and particulate organic carbon in northern San Francisco Bay during low flow conditions. *Estuarine, Coastal and Shelf Science*. Vol. 51, pp75–90.
- Munro P., Gauthier J., Laumond F. (1987). Changes in *Escherichia coli* cells starved in seawater or grown in seawater-wastewater mixtures. *Applied Environmental Microbiology*, Vol.53, pp1476-1481.
- Münster, U., Einiö, P., Nurminen, J. & Overbeck, J. (1992). Extracellular enzymes in a polyhumic lake: important regulators in detritus processing. *Hydrobiologia*. Vol. 229, pp. 225-238.
- Pakulski, J.D. & Benner, R. (1994). Abundance and distribution of carbohydrates in the ocean. *Limnology and Oceanography*. Vol. 39, pp930–940..
- Park, J.S., Choi, D.H., Hwang, G.J., & Park, B.C. (2006). Seasonal study on ectoenzyme activities, carbohydrate concentrations, prokaryotic abundance and production in a solar saltern in Korea. *Aquatic Microbial Ecology*. Vol. 43, pp 153-163.
- Pereira, M.G., Alcantara, F., 1993. Culturability of *Escherichia coli* and *Streptococcus faecalis* in batch culture and “in situ” in estuarine water (Portugal). *Water Research* Vol.27, pp.1351-1360
- Pomeroy LR, Wiebe WJ, Deibel D, Thompson RJ & Rowe GT,(1991). Bacterial responses to temperature and substrate concentration during the Newfoundland spring bloom. *Marine Ecology Progressive Series*. Vol 75, pp143–59

- Poncet S., Milohanic E., Mazé A., Nait Abdallah J., Aké F., Larribe M., Deghmane A.E., Taha M.K., Dozot M., De Bolle X., Letesson. J.J & Deutscher J.(2009). Correlations between Carbon Metabolism and Virulence in Bacteria. In Contributions to Microbiology (Mattias Collins & Raymond Scush (Editors). Vol. 16, pp.88-102.
- Rozen, Y. & Belkin, S., 2001. Survival of enteric bacteria in seawater. *Microbiology Review* 25, 513-529.
- Rozen, Y., LaRossa, R.A., Templeton, L.J., Smulski, D.R. & Belkin, S. (2002). Gene expression analysis of the response by *Escherichia coli* to seawater. *Antonie van Leeuwenhoek*. Vol.81, pp15-25.
- Ruger, H.J., 1989. Benthic studies of the northwest African upwelling region psychrophilic bacterial communities from areas with different upwelling intensities. *Marine Ecological Progress Series*. Vol. 57, pp. 45-52.
- Simidu, U., Taga, N., Cohvell, R.R., Schwartz, J.R., (1980). Heterotrophic bacterial flora of the seawater from the Nansei Shoto (Ryukyu Retto) area. *Bulletine Japanese Society of Science and Fisheries*. Vol. 46, pp.505-510.
- Simon M, Grossart HP, Schweitzer B & Ploug H.(2002).Microbial ecology of organic aggregates in aquatic ecosystems. *Aquat Microb Ecol*. Vol.28, pp175– 211.
- Smith, D.C., Simon, M., Alldredge, A.L. & Azam, F., (1992). Intense hydrolytic enzyme activity on marine aggregates and implications for rapid particle dissolution. *Nature* Vol. 359, pp. 139-142.
- Sinton, L.W., Davies-Colley, R.J., Bell, R., 1994. Inactivation of enterococci and fecal coliforms from sewage and meatworks effluents in seawater chambers, *Applied and Environmental Microbiology*. Vol60, pp2040-2048.
- Tietjen, T. & Wetzel, R. G., (2003) Extracellular enzyme-clay mineral complexes: enzyme adsorption, alteration of enzym activity, and protection from photodegradation. *Aquatic Ecology*, Vol. 37, pp 331–339.
- Tomlinson G. A., & Hochstein L.I. (1976). *Halobacterium saccharovororum* sp. nov., a carbohydrate-metabolizing, extremely halophilic bacterium. *Canadian Journal of Microbiologie*. Vol. 22, pp.587-591.
- Troussellier, M., Got, P., Bouvy, M., M'Boup, M., Arfi, R., Lebihan, F., Monfort, P., Corbin, D. & Bernard, C., (2004). Water quality and health status of the Senegal River estuary. *Marine Pollution Bulletin*. Vol. 48, pp 852-862.
- Truu J., Talpsep E., Heimrn E., Stottmeister U., Wand H., Hermarn A. (1999). Comparison system of Api 20NE and Biolog GN identification systems assessed by techniques of multivariate analyses. *Journal of Microbiological Methods*. Vol.36, pp 193-201.
- Urbani, R., Sist, P., 2003. Studio di caratterizzazione chimica delle componenti polisaccaridiche e del loro ruolo nel meccanismo di formazione degli aggregati gelatinosi. In: Istituto per la Ricerca Scientifica e Tecnologica Applicata al Mare, Ministero dell'Ambiente e della Tutela del Territorio, editors. Programma di monitoraggio e studio sui processi di formazione delle mucillagini nell'Adriatico e nel Tirreno — MAT. Final Report, June 2003. pp.135–70.

- Valdemarsen T., & Kristensen E.(2010). Degradation of dissolved organic monomers and short-chain fatty acids in sandy marine sediment by fermentation and sulfate reduction. *Geochimica et Cosmochimica Acta*, Vol. 74, pp1593–1605
- Vetter, Y.A. & Deming, J.W. (1999). Growth rates of marine bacterial isolates on particulate organic substrates solubilized by freely released extracellular enzymes. *Microbial Ecology*, Vol. 37, pp 86-94..
- Weiss MS, Abele J, Weckesser W, Welte W, Schultz E, Schulz GE. (1991) Molecular architecture and electrostatic properties of a bacterial porin. *Science*. Vol.254, pp1627– 30.
- Wetzel, R. G. & Tuchman, N. C. (2005) Effects of atmospheric CO₂ enrichment and sunlight on degradation of plant particulate and dissolved organic matter and microbial utilization – *Archives of. Hydrobiology*, Vol. 162, pp 287–308.
- Williams PJeB. (1981). Microbial contribution to overall marine plankton metabolism: direct measurements of respiration. *Oceanologica. Acta* Vol.4, pp359–64
- Williams, P.M. & Druffel, E.R.M. (1987). Radiocarbon in dissolved organic matter in the Central North Pacific Ocean. *Nature*. Vol. 330, pp246–248.
- Whitman, W. B., Coleman, D. C. & Wiebe, W. J. (1998). Prokaryotes: the unseen majority. *Proceedings of the. Natural Academic of Sciences USA*, Vol. 95, pp 6578–6583.
- Zoppini A., Puddu A., Fagi S., Rosati M. & Sist P. (2005). Extracellular enzyme activity and dynamics of bacterial community in mucilaginous aggregates of the northern Adriatic Sea. *Science of the Total Environment*. Vol.353, pp 270– 286