We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Role of Inflammation and Angiogenic Growth Factors in Malignant Mesothelioma

Loredana Albonici, Camilla Palumbo and Vittorio Manzari

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/47951>

1. Introduction

Malignant mesothelioma (MM) is a highly aggressive tumor which arises from the mesothelial cell lining of the serosal surfaces, most cases (>90%) being of pleural origin (Attanoos & Gibbs, 1997; Robinson & Lake, 2005). The pathogenesis of MM has been mainly associated with previous asbestos exposure (Berman & Crump, 2008), with a latency period of up to 40 years, although other agents such as Simian virus 40 (SV40) or genetic susceptibility factors have been linked to the development of this tumor (Carbone et al., 2002; Pisick & Salgia, 2005). Indeed, human mesothelial cells are highly susceptible to SV40-mediated transformation *in vitro* and SV40 DNA sequences and large T antigen (Tag) have been detected in human MM cells (Bocchetta et al., 2000; Carbone et al., 2012; Gazdar et al., 2003).

MM is largely unresponsive to conventional chemotherapy or radiotherapy and, despite its low metastatic efficiency, it is highly invasive to surrounding tissues so that its extensive growth leads to the failure of the organs underlying the serosal membranes (Astoul, 1999). In fact, the primary cause of fatality in MM is related to the propensity of the tumor cells to invade locally, even though MM metastasis are more common after surgery and, at the autopsy, metastatic diffusion is observed in 50% of patients (Astoul, 1999). At present, the median survival from diagnosis of MM is less than two years (Palumbo et al., 2008).

The mesothelium is not just a passive protective surface, but a highly dynamic membrane (Mutsaers, 2004). It consists of a single layer of elongated, flattened, squamous-like cells of mesodermal origin, characterized by dual epithelial/mesenchymal features. Cuboidal mesothelial cells can also be found at various locations in physiological conditions. Further, mesothelial cells can adopt a cuboidal morphology, which reflects a metabolically activated state, after injury or stimulation of the serosal surface (Mutsaers, 2004). Indeed, mesothelial cells are sentinel cells that can sense and respond to a variety of signals within their

microenvironment. They participate in serosal inflammation by secreting both pro- and anti-inflammatory as well as immunomodulatory mediators. Besides, these cells can act as antigen presenting cells for T lymphocytes (Hausmann et al., 2000), regulate tissue repair, control fibrin deposition and breakdown, and modulate adhesion, growth and dissemination of tumor cells metastasizing to the serosal membranes (Mutsaers, 2002). In particular, in response to different types of stimuli, including cytokines and asbestos fibers, mesothelial cells have been reported to release prostaglandins, chemokines, reactive oxygen and nitrogen species and growth factors which represent key effectors in the modulation of inflammatory reactions that occur in response to pleural injury (Fleury-Feith et al., 2003; Mutsaers, 2002).

2. Asbestos-induced carcinogenesis as an inflammation-driven process

The association between exposure to asbestos fibers and development of lung cancer and mesothelioma is well established in both humans and animals models (Greillier & Astoul, 2008; Huang et al., 2011; Mossman & Churg, 1998; Yarborough, 2007). A variety of mediators, either generated directly from asbestos fibers or elaborated intracellularly or extracellularly by cells exposed to asbestos, are implicated in the initiation and promotion of mesothelial cell transformation.

The mechanisms underlying asbestos-induced carcinogenesis involve mutagenic and non-mutagenic pathways, the latter including inflammation, enhanced mitogenesis, cell signaling alterations, and cytotoxic apoptosis/necrosis. Neither of these two mechanisms alone fully accounts for the complex biological abnormalities produced by asbestos fibers, even though in MM asbestos appears to act as a complete carcinogen (Dong et al., 1994; Huang et al., 2011). Still, the chronic inflammatory response induced by asbestos inhalation seems to play a critical role in mesothelial cell transformation.

Asbestos exposure induces an inflammatory reaction with a large component of mononuclear phagocytes (Antony et al., 1993; Branchaud et al., 1993; Carbone et al., 2012; Choe et al., 1997). Upon differentiation into macrophages, these cells phagocytize asbestos fibers and, in response, release numerous cytokines and reactive oxygen species with mutagenic properties (Robledo & Mossman, 1999). Thus, many of the pathological consequences occurring in the lung following exposure to asbestos fibers are believed to arise from an inflammatory cascade involving both autocrine and paracrine events (Hillegass et al., 2010). Persistent pulmonary inflammation is observed in animal models of asbestosis that can be correlated with fibroproliferative responses (Mossman & Churg, 1998).

Experimental models, as well as *in vitro* studies, have shown that mesothelial cells are particularly susceptible to the cytotoxic effects of asbestos (Baldys et al., 2007; Bérubé et al., 1996; Broaddus et al., 1996). Asbestos does not induce transformation of primary human mesothelial cells *in vitro*, instead, it is very cytotoxic to this cell type, causing extensive cell death. This finding raised an apparent paradoxical issue of how asbestos causes MM if human mesothelial cells exposed to this mineral die (Liu et al., 2000). This apparent paradox is reconciled by the current hypothesis that the chronic inflammation induced by asbestos

leads to the persistent activation of the nuclear factor kappa B (NF- κ B) transcription factor, which in turn mediates the activation of prosurvival genes and prevents apoptosis of the damaged mesothelial cells (Mantovani et al., 2008; Micheau & Tschopp, 2003; Philip et al., 2004). This allows mesothelial cells with asbestos-induced DNA damage to survive and divide rather than die and, if sufficient genetic damage accumulates, to eventually develop into a MM (Miura et al., 2006; Nymark, 2007). In fact, apoptosis is an important mechanism by which cells with DNA damage are eliminated without eliciting an inflammatory response (Ullrich et al., 2008; Yoshida et al., 2010). However, failure of apoptosis in cells with unrepaired DNA and chromosomal damage after chronic exposure to asbestos may lead to permanent genetic alterations and trigger the development of a clone of cancerous cells (Roos & Kaina, 2006; Wu, 2006). Consistently, MM cells are found to be apoptosis-resistant as compared to primary cultured mesothelial cells (Fennel & Rudd, 2004; Villanova et al., 2008).

2.1. Tumor Necrosis Factor- α and other pro-inflammatory cytokines

Tumor Necrosis Factor- α (TNF- α) is probably the most studied candidate for initiating inflammatory and fibrotic events linked to lung diseases such as asbestosis. Asbestos fibers cause the accumulation of macrophages in the pleura and lung. When these macrophages encounter asbestos, they release TNF- α . At the same time, asbestos induces the secretion of TNF- α and the expression of TNF- α receptor I (TNF-RI) in mesothelial cells (Yang et al., 2006). Remarkably, treatment of mesothelial cells with TNF- α significantly reduced asbestos cytotoxicity. Indeed, TNF- α activates NF- κ B, which in turn promotes mesothelial cell survival and resistance to the cytotoxic effects of asbestos. Thus, TNF- α signaling through NF- κ B-dependent mechanisms increases the percentage of mesothelial cells that survive asbestos exposure, thereby increasing the pool of asbestos-damaged cells susceptible to malignant transformation (Haegens et al., 2007; Janssen-Heininger et al., 1999; Yang et al., 2006).

It has been reported that rats receiving a single intratracheal instillation of fibrogenic chrysotile asbestos developed lung chronic inflammatory reactions characterized by the accumulation of alveolar macrophages producing elevated levels of both Interleukin (IL)-1 and IL-6 (Lemaire & Ouellet, 1996). An increased production and/or release of these cytokines triggers inflammatory cell recruitment, thus amplifying and sustaining local inflammation. It has also been demonstrated that crocidolite asbestos and TNF- α can stimulate a dose-dependent increase in IL-6 expression and secretion from cultured, transformed and normal, human alveolar type II epithelial cells that is dependent upon intracellular redox potential (Simeonova et al., 1997). Interestingly, although MM cells appear to express low levels of IL-6 receptor (IL-6R), IL-6 can act as a growth factor for these cells through a trans-signaling mechanism involving the interaction of macromolecular complexes of IL-6 and soluble IL-6R (sIL-6R) with the transmembrane glycoprotein gp130 expressed on the surface of MM cells (Adachi et al., 2006; Rose-John et al., 2007). High levels of both IL-6 and sIL-6R are typical of several chronic inflammatory conditions (Rose-John et al., 2007).

Thus, inflammatory cytokines such as TNF- α and IL-6 appear to play a dual role in MM pathogenesis: they induce and sustain pleural inflammation and at the same time can act as survival or mitogenic factors for normal and transformed mesothelial cells, respectively.

2.2. Reactive Oxygen and Nitrogen Species (ROS/RNS)

The mechanisms of injury and disease development caused by asbestos fibers are presumed to be related to their greater fibrogenic and carcinogenic properties in comparison to other minerals. Asbestos-induced mutagenicity is mediated through both direct and indirect pathways. Asbestos fibers may induce mutagenicity and genotoxicity directly through physical interaction with the mitotic machinery after being phagocytized by the target cells, or indirectly as a result of DNA and chromosome damage caused by asbestos-induced reactive oxygen (ROS) and nitrogen species (RNS) (Kamp & Weitzman, 1999; Shukla et al., 2003a, 2003b). ROS and RNS can be generated primarily by asbestos fibers or secondarily through fiber-induced inflammation (Aust et al., 2011; Gulumian, 2005; Hoidal, 2001). Free radicals generated from asbestos fibers plus the direct damage induced by the fibers are linked to cell signaling, inflammation, and a plethora of other responses (mutagenesis, proliferation, etc.) associated with the pathogenesis of asbestos-associated diseases (Heinz et al., 2010; Manning et al., 2002; Shukla et al., 2003a, 2003b).

Several evidences indicate that a main factor in determining the surface and biological reactivity of different types of asbestos fibers is their ability to participate in redox reactions that generate free radicals (Kamp & Weitzman, 1999; Shukla et al., 2003a). Although the nature of the free radical-generating surface sites on asbestos fibers is not yet clear, asbestos fibers have an intrinsic redox activity and contain ferrous iron, which catalyzes reactions generating active oxygen intermediates on the fiber surface. Within the tissues several asbestos fiber types can produce reactive oxygen free radicals from hydrogen peroxide, a common product of intermediary tissue metabolism. Epidemiological studies have identified crocidolite as one of the most potent forms of asbestos associated with the induction of MM (Heintz et al., 2010). Crocidolite has a greater surface-area and a higher ferrous iron content compared to other fiber types such as chrysotile, and it is more biologically active in the generation of free radicals (Toyokuni, 2009). However, the ability of asbestos fibers to elicit these effects is not related to total iron content, suggesting the presence of specific iron active sites at the fibers' surface (Shukla et al., 2003a).

Cells exposed to asbestos have also been reported to produce a higher amount of nitric oxide (NO). In this regard, it has been reported that in human mesothelial cells crocidolite increases the expression of the inducible NO synthase (NOS) isoform (iNOS), the activity of the constitutive endothelial NOS (eNOS), and the synthesis of NO via NF- κ B and Akt activation (Riganti et al., 2007). Thus, the asbestos-induced upregulation of iNOS or NO in the lungs, as well as the induction of inflammation by fibers, may contribute along with ROS, to the pathogenesis of lung and pleural injury (Hussain et al., 2003; Tanaka et al., 1998). Indeed, ROS and RNS can cause breakage of DNA, lipid peroxidation, release of inflammatory cytokines such as TNF- α , and the modification of cellular proteins including phosphatases involved in cell signaling cascades (Gossart et al., 1996; Hussain et al., 2003), so that their increased synthesis by various cell types may have multiple roles in cellular events critical to the establishment of lung and pleural inflammation and uncontrolled cell proliferation.

Finally, in mesothelial and lung epithelial cells asbestos fibers, as opposed to nonpathogenic minerals, cause a persistent induction of the redox-sensitive transcription factors NF- κ B and Activator Protein-1 (AP-1), which is accompanied by chronic alterations in gene expression (Heintz et al., 1993; Janssen et al., 1995). As mentioned above, the aberrant activation of the NF- κ B pathway is regarded as a critical event for mesothelial cell transformation (Toyooka et al., 2008).

2.3. Transcription factors

2.3.1. NF- κ B

NF- κ B proteins are dimeric transcription factors composed of five different subunits, namely p65 (RelA), RelB, c-Rel, NF- κ B1 p50 and NF- κ B2 p52, which regulate gene expression events that impact on cell survival and differentiation. Moreover, since activation of NF- κ B is critical in up-regulating the expression of many genes linked to proliferation, apoptosis resistance, and chemokine/cytokine production, this is undoubtedly a critical transcription factor in inflammatory responses occurring in target cells of asbestos-related diseases (Janssen et al., 1995, 1997).

In unstimulated cells, the NF- κ B transcription dimers are retained in the cytoplasm in an inactive state through the interaction with a family of inhibitors called I κ Bs (Inhibitors of κ B) or with the p50 and p52 precursor proteins, p105 and p100, respectively (Hayden & Ghosh, 2008; Scheidereit, 2006). Indeed, p50 and p52 are translated as precursor proteins containing an I κ B-like C-terminal portion (Sun, 2011).

Two different NF- κ B-activation pathways exist: the classical and the alternative NF- κ B pathway. The classical NF- κ B pathway is initiated by signals elicited by diverse receptors, including TNF receptors type 1/2, Toll/IL-1 receptor, T-cell and B-cell receptors and EGF receptor, and also by cellular stresses and DNA damage (Hayden & Ghosh 2004; Le Page et al., 2005). These signals induce the activation of the I κ B kinase (IKK) complex, which is composed by the catalytic subunits IKK α and IKK β and by the regulatory subunit IKK γ /NEMO (Hayden & Ghosh, 2008; Scheidereit, 2006; Sun, 2011). The activated IKK complex phosphorylates I κ B proteins, thereby triggering their proteasomal degradation. As a consequence, NF- κ B dimers are released and can translocate into the nucleus. This pathway mainly leads to the activation of p50:RelA dimers (Sun, 2011). Conversely, the alternative NF- κ B pathway predominantly targets activation of RelB:p52 complexes. This pathway relies on the inducible processing of p100 triggered by signaling from TNF receptor family members via the NF- κ B-inducing kinase (NIK): NIK activates IKK α , which, in turn, phosphorylates p100 and triggers its processing to p52. This event results in the conversion of p100-inhibited NF- κ B complexes into p52-containing NF- κ B dimers, capable of translocating into the nucleus (Hayden & Ghosh, 2008; Scheidereit, 2006; Sun, 2011).

NF- κ B-regulated genes have distinct requirements for NF- κ B dimers. For instance, the NF- κ B binding site of the IL-2 gene has been reported to bind preferentially c-Rel homodimers

and p50:c-Rel, while that of the gene encoding IL-8 has been found to selectively bind Rel A (Hoffman et al., 2003, 2006). On the other hand, several genes are redundantly induced by more than one dimer (Hoffman et al., 2003, 2006; Sacconi et al., 2003).

A number of studies have shown that nuclear retention and DNA binding of NF- κ B protein complexes are increased following exposure of various cell types to a variety of extracellular stimuli that include oxidative stress (Bowie & O'Neill, 2000), hypoxia (Jung et al., 2003; Royds et al., 1998) and inflammatory cytokines (Mantovani et al., 2008). These observations are consistent with the hypothesis that persistent activation of NF- κ B can contribute to the induction of multiple genes that are critical to the pathogenesis of asbestos-associated diseases, since oxidants, local hypoxia and inflammatory cytokines are all components involved in the effects induced by asbestos exposure.

It is noteworthy that among various carcinogenic and non-carcinogenic fibers studied for their effect on nuclear translocation of NF- κ B, only carcinogenic fibers were found to cause a dose-dependent translocation of this transcription factor to the nucleus, and this effect was reported to be oxidative stress-dependent (Brown et al., 1999). In lung macrophages, the asbestos-induced expression and secretion of TNF- α are mediated by iron-catalyzed ROS products (Simeonova & Luster, 1995) through a process that involves NF- κ B activation (Cheng et al., 1999). In rat alveolar type 2 cells, the crocidolite-induced activation of NF- κ B as well as the expression of the macrophage inflammatory protein-2 (MIP-2) gene have also been shown to be dependent on mitochondrial-derived oxidative stress (Driscoll et al., 1998).

2.3.2. AP-1

AP-1 is a homo- or heterodimeric transcription factor composed by proteins encoded by the *fos* and *jun* early response proto-oncogenes. This family of proteins includes c-Fos, FosB, FosL1 (Fra-1), FosL2 (Fra-2), c-Jun, JunB and JunD (Milde-Langosch, 2005). Whereas Jun members are capable of forming homodimers able to bind DNA and regulate transcription, all Fos members must form heterodimers with Jun family members to bind DNA.

AP-1 is a redox-sensitive transcription factor typically associated with cell proliferation and tumor promotion (Eferl & Wagner, 2003). The first evidence showing that asbestos exerts regulatory effects linked to aberrant transcriptional responses, cell proliferation and cell transformation derives from studies in which asbestos fibers caused induction of *c-fos* and *c-jun* proto-oncogene mRNAs in pleural mesothelial cells and tracheo-bronchial epithelial cells in a dose-response fashion (Heintz et al., 1993).

The persistent induction of AP-1 by asbestos suggests a model of asbestos-induced carcinogenesis involving chronic stimulation of cell proliferation through activation of early response genes (Schonthaler et al., 2011). Of note, early response genes are a set of genes whose transcription is rapidly induced in response to growth factors. Furthermore, AP-1 activity is induced by growth factors, pro-inflammatory cytokines and genotoxic stress (Jochum et al., 2001; Shaulian & Karin, 2002). These stimuli activate mitogen-activated protein kinase (MAPK) cascades through the phosphorylation of distinct substrates such as

ERK, JNK and p38 MAPK (Chang & Karin, 2001). Indeed, the MAPK signal transduction pathway uses AP-1 as a converging point not only to regulate the expression of various genes but also to autoregulate AP-1 gene transcription (Reuter et al., 2010).

Several genes, which play very important roles in injury, repair, and differentiation, contain binding site(s) for AP-1 in their promoter and/or enhancer regions (Chang & Karin, 2001). These genes include extracellular matrix metalloproteinases (MMPs), antioxidant enzymes, growth factors and their receptors, differentiation markers, cytokines, chemokines and other transcription factors (Shaulian & Karin, 2001).

2.3.3. Nuclear Factor of Activated T Cells (NFAT)

The Nuclear Factor of Activated T cells (NFAT) family of transcription factors consists of five proteins that are evolutionarily related to the Rel/NF- κ B family. NFAT can be present in both the cytoplasm and the nucleus. In the cytoplasm NFAT is in a highly phosphorylated, inactive state. Cell stimuli leading to the elevation of intracellular Ca^{2+} levels induce the activation of the phosphatase PP2B/Calcineurin which dephosphorylates NFAT. This results in its nuclear relocation and transcriptional activation. Interestingly, NFAT family members can act synergistically with AP-1 on composite DNA elements which contain adjacent NFAT and AP-1 binding sites (Macián et al., 2001). A functional cooperation has also been reported to occur between NFAT and NF- κ B (Jash et al., 2012).

Initially, NFAT was identified in lymphocytes and was reported to be expressed in activated but not resting T cells (Macián et al., 2005; Shaw et al., 1988). NFAT regulates not only T cell activation and differentiation but also the function of other immune cells, including dendritic cells (DCs), B cells and megakaryocytes. In addition, NFAT has crucial roles in numerous developmental programs in vertebrates.

Dysregulation of NFAT signalling is now known to be associated with malignant transformation and the development of cancer (Mancini & Toker, 2009; Müller & Rao, 2010). The observation that NFAT can be activated by asbestos-induced oxidative stress suggests that this transcription factor may play multiple roles in asbestos-induced inflammation and carcinogenesis (Li et al., 2002). Indeed, NFAT mediates the expression of several inflammatory cytokines, including TNF- α , and is involved in cell transformation, proliferation, invasive migration, tumor cell survival and tumor angiogenesis (Mancini & Toker, 2009).

3. Multifaceted role of angiogenic growth factors in MM

Angiogenesis is a common feature of solid tumors. Indeed, the development of a clinically observable tumor requires the neoformation of a vascular network sufficient to sustain tumor growth (Ribatti et al., 2007). Tumor angiogenesis is stimulated by the secretion of angiogenic molecules which induce endothelial cells from nearby vessels to switch from a quiescent to an activated state. Further, upon the stimulation of angiogenic growth factors, activated endothelial cells disrupt the extracellular matrix, proliferate and migrate (Ribatti et

al., 2007). Angiogenic growth factors include, among the others, Vascular Endothelial Growth Factor (VEGF), Placenta Growth Factor (PlGF), Platelet-Derived Growth Factor (PDGF) and acidic and basic Fibroblast Growth Factors (FGF-1 and -2, respectively). VEGF is regarded as the most important player in angiogenesis (Ono, 2008).

The link between angiogenesis and tumor progression is provided by the negative prognostic value of intratumoral microvascular density (IMD) (Folkman, 2006; Kerbel, 2008). In MM the IMD has an independent prognostic value (Kumar-Singh et al., 1997). MM demonstrates a higher IMD than colon and breast tumors and, consistently, presents with minimal central necrosis despite its huge size (Gasparini & Harris, 1995; Kumar-Singh et al., 1997).

On the other hand, the involvement of angiogenic growth factors in MM goes beyond the stimulation of angiogenesis. Indeed, as discussed below, MM cells express receptors for several angiogenic factors which, accordingly, can directly modulate MM cell behavior.

3.1. Angiogenic growth factors of the VEGF family

The human VEGF family consists of five members: VEGF (VEGF-A), VEGF-B, VEGF-C, VEGF-D and PlGF. These growth factors are secreted as dimers and their biological effects are mediated by binding to three tyrosine kinase receptors, *i.e.* VEGF-R1/Flt-1, VEGF-R2/KDR (whose murine homologue is known as Flk-1) and VEGF-R3/Flt-4, and two non-enzymatic co-receptors known as neuropilin-1 and -2 (Ferrara et al., 2003; Koch et al., 2011; Roskoski, 2007).

3.1.1. VEGF

VEGF is regarded as the major mediator of tumor angiogenesis. It is expressed in the majority of cancers and has a central role in tumor growth and metastasis. In fact, this growth factor is essential for the mobilization of bone-marrow-derived endothelial precursors in neovascularization (Asahara et al., 1999), and stimulates vascular endothelial cells mobility, proliferation and survival (Waltenberger et al., 1994).

High levels of VEGF are present both in malignant and non-malignant pleural effusions leading to increased vascular permeability. On the other hand, VEGF levels in serum or pleural effusions of MM patients are higher than those found in patients with non-malignant pleuritis or lung cancer involving malignant pleural effusions. Further, in MM patients elevated serum or pleural effusion levels of VEGF correlate with a worse prognosis and may also contribute to increase resistance to chemotherapy (Hirayama et al., 2011; Yasumitsu et al., 2010; Zebrowski et al., 1999). In fact, VEGF status has proved to be of value in predicting the effectiveness of radiotherapy and chemotherapy on different cancers (Choi et al., 2008; Kumar et al., 2009; Toi et al., 2001).

In addition to its role in tumor vascularization, VEGF can directly affect the behavior of cancer cells in an autocrine or paracrine manner. Indeed, many tumor cell types express VEGF receptors. VEGF has been found to promote the growth of transformed cell lines *in vitro* (Masood et al., 2001) and to act as a survival factor for tumor cells by enhancing the

expression of the antiapoptotic factors bcl-2 (Harmey & Bouchier-Hayes, 2002) and survivin (Kanwar et al., 2011). In this context, MM cells have been shown to express high amounts of VEGF, VEGF receptors and co-receptors both *in vitro* and *in vivo*, and VEGF has been demonstrated to act as an autocrine growth factor for this tumor cell type (Albonici et al., 2009; Ohta et al., 1999; Pompeo et al., 2009; Strizzi et al., 2001a).

VEGF-R1 participates in cell migration; it has an important role in monocyte chemotaxis and promotes recruitment of circulating endothelial precursor cells from bone marrow (Hattori et al., 2002). Its expression is increased in various tumors, correlates with disease progression and can predict poor prognosis, metastasis and recurrent disease in humans (Dawson et al., 2009; Fischer et al., 2008; Kerber et al., 2008). This receptor is also expressed by MM cells *in vitro* and *in vivo*, where it appears to mediate proliferative and cell survival responses (Albonici et al., 2009; Strizzi et al., 2001a).

VEGF-R2 is the main mediator of VEGF-stimulated endothelial cell migration, proliferation, survival and enhanced vascular permeability (Olsson et al., 2006; Shibuya, 2006). VEGF-R2 expression is induced in conjunction with active angiogenesis, such as during the reparative process, and in pathological conditions associated with neovascularization, such as cancer (Plate et al., 1993). VEGF-R2 is overexpressed in MM cells and specimens, and VEGF-R2 silencing by small interfering RNA has been shown to induce cell death in MM or immortalized mesothelial cells *in vitro* (Albonici et al., 2009; Catalano et al., 2009; Pompeo et al., 2009; Strizzi et al., 2001a). Interestingly, it has been reported that in MM cells this receptor can be activated also via the semaphorin-6D receptor Plexin-A1, triggering a prosurvival program that promotes anchorage-independent growth through a NF- κ B-dependent pathway (Catalano et al., 2009). Remarkably, the expression of plexin-A1 is induced by asbestos fibers and overexpression of plexin-A1 in non-malignant mesothelial cells inhibits cell death after asbestos exposure, thus suggesting a role for this receptor not only in MM promotion and progression but also in asbestos-induced mesothelial carcinogenesis (Catalano et al., 2009).

In vitro studies have shown that transfection of normal mesothelial cells with SV40 Tag potently increases VEGF protein and mRNA levels (Cacciotti et al., 2002) as well as mesothelial cell proliferation (Catalano et al., 2002). These data indicate that VEGF regulation by SV40 transforming proteins can also represent a key event in MM onset and progression.

3.1.2. PlGF

PlGF, originally identified in the placenta during the early embryonic development (Khaliq et al., 1996; Maglione et al., 1991), is expressed in several other organs including the heart, lung, thyroid, skeletal muscle and adipose tissue (Persico et al., 1999) but not normal mesothelium (Albonici et al., 2009).

Although the role exerted by PlGF in tumor growth is controversial yet, PlGF can stimulate vessel growth and maturation directly by affecting endothelial and mural cells, as well as indirectly by recruiting pro-angiogenic cell types (Barillari et al., 1998; Carmeliet, 2003). It

also promotes the recruitment and maturation of angiogenesis-competent myeloid progenitors to growing sprouts and collateral vessels (Hattori et al., 2002; Luttun et al., 2002; Rafii et al., 2003). Further, PlGF is able to protect endothelial cells from apoptosis, in a similar manner as VEGF, by inducing the expression of antiapoptotic genes such as survivin (Adini et al., 2002).

Under pathological conditions, PlGF abundance is elevated in various cell types and tissues, including vascular endothelial cells, and many different tumor cells (Albonici et al., 2009; Cao et al., 1996; Fischer et al., 2007; Oura et al., 2003). PlGF expression is switched on in hyperplastic/reactive mesothelium and in MM cells (Albonici et al., 2009). Moreover, in MM as well as in different types of cancer, including melanoma, gastric, colorectal and breast carcinomas, PlGF plasma levels and intratumoral expression have been found to correlate with tumor stage, vascularity, recurrence, metastasis and survival (Chen et al., 2004; Marcellini et al., 2006; Parr et al. 2005; Pompeo et al., 2009; Wei et al., 2005).

In vitro studies have shown that administration of recombinant PlGF to MM cells triggers the activation of Akt but does not elicit a significant stimulation of cell growth. Conversely, the administration of PlGF-neutralizing antibodies causes a significant reduction of MM cell viability, demonstrating the PlGF acts as a survival factor for MM cells (Albonici et al., 2009).

PlGF binds VEGF-R1 and the co-receptors neuropilin-1 and -2, but, unlike VEGF, it does not bind VEGF-R2. Accordingly, it can act independently of VEGF in cells which primarily express VEGF-R1 (Fischer et al., 2007). Worthy of note, even though VEGF and PlGF both bind VEGF-R1, PlGF was reported to stimulate the phosphorylation of specific VEGF-R1 tyrosine residues and the expression of distinct downstream target genes as compared to VEGF (Autiero et al., 2003). On the other hand, PlGF can also sustain VEGF activity through different mechanisms involving both VEGF-R1 and VEGF-R2. One of these mechanisms relies on the formation of PlGF:VEGF heterodimers. Indeed, PlGF:VEGF heterodimers have been isolated from cells producing both factors and shown to bind VEGF-R1:VEGF-R2 receptor complexes, thus inducing receptor cross-talk and activation of VEGF-R2, the major mediator of VEGF activities (Autiero et al., 2003; Cao et al., 1996). In addition, the activation of VEGF-R1 by PlGF homodimers may induce the intermolecular transphosphorylation and activation of VEGF-R2 (Carmeliet et al., 2001).

It is noteworthy that *in vivo* anti-PlGF treatment was reported to inhibit tumor growth without affecting healthy vessels, thus reducing tumor infiltration by angiogenic macrophages and severe tumor hypoxia, and preventing the switch on of the angiogenic rescue program leading to the enhanced release different angiogenic factors responsible for resistance to VEGF receptors inhibitors (Fischer et al., 2007).

3.2. PDGF

PDGFs comprise a family of dimeric growth factors structurally and functionally related to VEGFs (Andrae et al., 2008). PDGF homodimers are formed by four different chains, *i.e.* PDGF-A, PDGF-B, PDGF-C and PDGF-D. In addition, PDGF-A and -B chains can form the

heterodimeric PDGF-AB. The biological effects of PDGF are mediated by two tyrosine kinase receptors, namely the PDGF receptor alpha (PDGFR α), which binds PDGF-A, -B, and -C chains, and the PDGF receptor beta (PDGFR β), which binds PDGF-B and -D. Accordingly, upon ligand binding different receptor dimers may form depending on ligand configuration and the pattern of receptor expression. Cellular responses to PDGF signaling include stimulation of cell growth, differentiation, migration and inhibition of apoptosis (Andrae et al., 2008).

An increased PDGF activity has been linked with tumors, vascular and fibrotic diseases (Andrae et al., 2008). Autocrine PDGF signaling leading to enhanced proliferation of tumor cells occurs in several types of cancer (Ostman, 2004). In addition, PDGF secretion by cancer cells and activated endothelial cells promotes the formation of both fibrous and vascular tumor stroma. In particular, PDGF-BB participates in tumor angiogenesis by stimulating endothelial cell motility and pericyte recruitment to neoformed vessels, thus leading to vessel stabilization, tumor cell survival and growth. Instead, both PDGF-AA and PDGF-BB appear involved in tumor recruitment of PDGFR-positive fibroblasts which, in turn, can be activated by PDGFs to produce VEGF and other tumor-promoting growth factors (Andrae et al., 2008; Cao et al., 2008; Homsí & Daud, 2007).

Either high PDGF-AB serum levels or a strong expression of PDGFR signaling effectors in MM tissues have been associated with a lower survival in MM patients (Filiberti et al., 2005; Kothmaier et al., 2008). In fact, several evidence support a role for PDGF in MM promotion and progression through both autocrine and paracrine mechanisms.

While PDGFR α expression levels are lower in MM than in normal mesothelial cells, PDGFR β , PDGF-A and PDGF-B are overexpressed in MM cells as compared to their non-transformed counterparts (Langerak et al., 1996a, 1996b; Metheny-Barlow et al., 2001). Functional studies have shown that transduction of MM cells with a hammerhead ribozyme against PDGFR β mRNA reduced both PDGFR β expression and MM cell proliferation, demonstrating the involvement of a PDGF-BB autocrine loop in MM cell growth (Dorai et al., 1994). Conversely, the role of PDGF-A in MM cell proliferation is controversial. Indeed, the transfection of MM cells with antisense oligonucleotides to PDGF-A has been reported to either inhibit or stimulate MM cell growth *in vitro* (Garlepp & Leong, 1995; Metheny-Barlow et al., 2001). On the other hand, PDGF-A appears to play an important role in sustaining MM cell growth *in vivo* through paracrine mechanisms. Indeed, PDGF-A overexpression in MM cells inoculated in nude mice was found to increase tumor incidence, tumor growth rate and to decrease the latency period to tumor formation (Metheny-Barlow et al., 2001). In this regard, it has been suggested that PDGF-A participates in a malignant cytokine network through which MM cells instigates tumor-associated fibroblasts to produce growth factors, such as hepatocyte growth factor (HGF), with tumor-promoting activities (Li et al., 2011).

3.3. FGF

The FGF family encompasses 22 structurally related ligands in mammals. The effects of most FGF family members, including FGF-1 and -2, are mediated by binding to a family of

tyrosine kinase receptors designated FGF receptors (FGFR1 to FGFR5), whereas a smaller number of FGF isoforms does not bind FGFRs but interacts with voltage-gated sodium channels (Knights & Cook, 2010).

FGFs regulate cell proliferation, differentiation, survival, wound healing and angiogenesis. In cancer, FGF signaling is frequently de-regulated, resulting in mitogenic, anti-apoptotic and angiogenic responses (Knights & Cook, 2010). FGF-1 and -2, but also other less-studied FGF isoforms, exert pro-angiogenic effects by modulating proliferation and migration of endothelial cells and by stimulating the production of proteases (Lieu et al., 2011; Saylor et al., 2012). Worthy of note, it has been demonstrated that FGF-2 can synergize with both VEGF and PDGF-BB in stimulating neovascularization, this synergism relying on multiple mechanisms. For instance, FGF-2 promotes hypoxia-induced VEGF release by cancer cells and the expression of both VEGF and VEGFRs in endothelial cells, whereas VEGF, in turn, upregulates the expression of FGF-2 (Lieu et al., 2011; Saylor et al., 2012). Moreover, FGF-2 upregulates PDGFRs expression and increases the responsiveness to PDGF-BB in endothelial cells, whereas PDGF-BB enhances FGFR1 expression and FGF-2 responsiveness in vascular smooth muscle cells (Cao et al., 2008; Liu et al., 2011). Remarkably, FGFs are thought to play a critical role in the resistance to anti-VEGF therapy (Lieu et al., 2011; Saylor et al., 2005). Besides, both FGF-1 and -2 may also be involved in tumor cell growth through cell-autonomous, autocrine mechanisms (Kumar-Singh et al., 1999).

FGF-1 and -2 are expressed in the majority of MMs *in vivo* and high levels of FGF-2 in tumor tissues, serum or pleural effusions are associated with a worse prognosis in MM patients (Davidson et al., 2004; Kumar-Singh et al., 1999; Strizzi et al., 2001b). Furthermore, the combined expression levels of FGF-1, FGF-2, VEGF and Transforming Growth Factor beta (TGF β) in MM tissues correlates with both IMD and a poorer prognosis (Kumar-Singh et al., 1999). In addition to their role in tumor angiogenesis, FGFs act as autocrine growth factors for MM cells. Indeed, MM cells express FGFs and FGF receptors and the transfection with short interfering RNAs to FGF-1 and FGF-2 reduces MM cell proliferation (Kumar-Singh et al., 1999; Liu & Klominek, 2003; Stapelberg et al., 2005). It has also been reported that treatment of MM cells with exogenous FGF-2 stimulates the secretion of matrix metalloproteinases involved in tumor invasion and angiogenesis (Liu & Klominek, 2003).

4. Cross-talk between inflammation and angiogenic growth factors

Experimental and epidemiological evidences indicate that chronic inflammation is associated with most, if not all, tumors and supports their progression (Coussens & Werb 2002; Mantovani et al., 2008; Mantovani et al., 2010; Porta et al., 2009). Chronic inflammation appears to have a versatile function in tumor onset and progression. Indeed, as discussed above, a long-lasting inflammation can contribute to cancer initiation through the production ROS and RNS with DNA-damaging properties. On the other hand, it can also participate in cancer promotion and progression by increasing the availability of mediators (growth factors, cytokines, chemokines, prostaglandins) which contribute to the growth of initiated cells and to neoangiogenesis (Mantovani, 2010). Besides, once a tumor is

established, cancer cells promote a constant influx of myelomonocytic cells that express inflammatory mediators supporting pro-tumoral functions. In this regard, myelomonocytic cells are key orchestrators of cancer-related inflammatory processes supporting proliferation and survival of malignant cells, subversion of adaptive immune responses, stromal remodeling and angiogenesis (David Dong et al., 2009; Loges et al., 2009; Porta et al., 2009).

Tissue infiltration by macrophages is a dramatic and common feature of inflammation, angiogenesis and cancer (Pollard, 2004; Sica, 2010). High densities of tumor-infiltrating macrophages are associated with poor survival in patients with MM (Burt et al., 2011). In fact, the recruitment and infiltration of macrophages in the tumor microenvironment can activate them to support the malignant progression of cancer cells. These macrophages are called tumor-associated macrophages (TAMs) (Lawrence, 2011; Sica, 2010). Cancer cells co-cultured with macrophages and incubated with inflammatory cytokines are synergistically stimulated to produce various angiogenesis-related factors (Izzi et al., 2009; Ono, 2008). This inflammatory angiogenesis is mediated, in part, by activation of NF- κ B and AP-1 (Angelo & Kurzrock, 2007; Huang et al., 2000; Ono, 2008). In fact, treatment of both vascular endothelial cells and cancer cells with IL-1 α/β , TNF- α and ROS *in vitro* results in a marked induction of VEGF and FGF-2, through the transcriptional activation of NF- κ B, Specificity protein 1 (Sp-1), AP-1 and hypoxia response elements.

In addition to macrophages, other tumor-infiltrating immune cells including T cells, B cells, natural killer cells and neutrophils can release cytokines, such as IL-1 α/β , TNF- α and IL-6, able to sustain the synthesis of angiogenic growth factors (Angelo & Kurzrock, 2007). As for, IL-6, this pro-inflammatory cytokine has been reported to play a critical role in the stimulation of VEGF synthesis by different cell types, including MM cells (Adachi et al., 2006; Angelo & Kurzrock, 2007). Of note, MMs usually produce high levels of IL-6 but express low levels of IL-6R, so that the presence of sIL-6Rs, which may be provided by inflammatory cells recruited to the tumor region, is essential for the IL-6-dependent stimulation of VEGF expression by MM cells (Adachi et al., 2006). Inflammation can also induce the expression of receptors for angiogenic growth factors. In this regard, the expression of PDGFRs is known to be induced by inflammatory cytokines such as TNF- α and IL-1 (Andrae et al., 2008). Besides, inflammatory cells themselves can directly release angiogenic factors such as VEGF, PlGF, FGF-2 and PDGF, among many others, which exert mitogenic and migratory effects on surrounding cells (Sica 2010, Ono 2008). Inflammatory cells recruited in the tumor microenvironment can also produce matrix metalloproteinases which promote the formation of new vessels by degrading the basement membrane and by releasing angiogenic growth factors, such as VEGF, PlGF-2 and FGF-2, stored in the extracellular matrix (Barillari et al., 1998; Cao et al., 2008; Lieu et al., 2011).

The high amount of chemokines/cytokines, growth factors, proteolytic enzymes, proteoglycans, lipid mediators and prostaglandins which is typically found in the tumor microenvironment sustains and exacerbates both inflammation and angiogenesis (Costa et al., 2007; Lin & Karin, 2007; Ono, 2008). In this context, the cross-talk between inflammation and angiogenesis is further corroborated by the evidence that, if on one hand inflammatory mediators have significant effects on angiogenesis, on the other hand angiogenic factors can

effectively promote inflammation. As a matter of fact, in addition to their angiogenic role, VEGF and PlGF appear to act as direct proinflammatory mediators in the pathogenesis of different inflammatory conditions (Angelo & Kurzrock, 2007; Yoo et al., 2008). In this regard, VEGF was found to increase the production of TNF- α and IL-6 by human peripheral blood mononuclear cells and macrophages (Yoo et al., 2008). Moreover, VEGF stimulates monocyte recruitment to tumor areas (Barleon et al., 1996). An additional link between inflammatory and angiogenic growth factors has been provided with the demonstration that in myelomonocytic cells TNF- α is upregulated by PlGF in a NFAT1-dependent manner and, in turn, contributes to PlGF-induced myelomonocytic cell recruitment (Ding et al., 2010). PlGF can also contribute to inflammation by acting as survival factor for monocytes and macrophages (Adini et al., 2002).

5. Cooperation between asbestos and angiogenic growth factors in MM onset and progression

As reported above, asbestos stimulates the expression of *c-fos* and *c-jun* mRNA in mesothelial cells in a dose-dependent fashion (Heintz et al., 1993; Ramos-Nino et al., 2002). One of the mechanisms by which VEGF and PlGF elicit biological responses is the induction of Fos-B and c-Fos expression in endothelial cells and monocytes (Holmes & Zachary, 2004). The coexistence of different stimuli, such as asbestos fibers and angiogenic growth factors, concurring to the activation of early response genes might lead to the persistent induction of AP-1 in mesothelial cells and to the chronic stimulation of mesothelial cell proliferation, thus favoring cell transformation.

Further, asbestos and angiogenic growth factors can cooperate in inducing an immunosuppressive tumor microenvironment. Indeed, asbestos has been found to possess immunosuppressive properties. For example, chrysotile fibers have been shown to depress the *in vitro* proliferation of phytohemagglutinin-stimulated peripheral blood lymphocytes and to suppress natural killer activity. Moreover, asbestos significantly reduces the generation and activity of lymphokine-activated killer (LAK) cells, which are immune effectors with a strong lytic activity against MM cells (Manning et al., 1991; Valle et al., 1998).

Immunosuppressive properties have been reported for angiogenic growth factors as well (Ohm et al., 2001; Ziogas et al., 2012). Impaired antigen-presenting function in DCs as a result of abnormal differentiation is an important mechanism of tumor escape from immune control. It has been demonstrated that VEGF can inhibit the maturation of DCs induced by lipopolysaccharide (Takahashi et al., 2004). VEGF can also affect the ability of hematopoietic progenitor cells (HPCs) to differentiate into functional DCs during the early stages of hematopoiesis *in vivo* (Gabrilovich et al., 1996; Oyama et al., 1998). In this regard, it has been shown that VEGF binds to specific receptors on the surface of HPCs and this binding appears to involve VEGF-R1. Interestingly, the number of binding sites available for VEGF decreased with DC maturation and correlated with decreased levels of VEGF-R1 mRNA expression in the late-stage cells (Gabrilovich et al., 1996). PlGF was also found to inhibit the

activation and maturation of human DCs effectively and rapidly through the NF- κ B pathway (Lin et al., 2007). The results of this study further indicate that by modulating the function of DCs, PlGF can down-regulate T helper immune responses (Lin et al., 2007). In addition, both VEGF and PlGF are also involved in the recruitment of macrophages with immunosuppressive, tumor-promoting roles to the tumor stroma.

On the whole, these findings suggest mechanisms by which tumor-derived soluble factors such as VEGF or PlGF may synergize with asbestos to down-regulate immune responses to MM antigens.

6. Conclusions

Collectively, the reported findings demonstrate that a complex network involving asbestos, inflammation and angiogenic factors upregulation is involved in the pathogenesis of MM. In particular, the abnormal expression of angiogenic factors appears to play multiple roles in MM: it stimulates tumor neovascularization, increases pleural effusion formation by increasing vascular permeability, supports autocrine tumor cell growth and finally, in synergism with asbestos fibers, can sustain inflammation and bias host immune responses. Accordingly, the upregulation of angiogenic growth factors appears to be a crucial event in mesothelial cell transformation and MM progression.

Given the involvement of multiple angiogenic growth factors in the formation of tumor vessels, in tumor inflammation and MM cell growth and survival, the therapeutic development of antiangiogenic agents for the treatment of this tumor should be aimed at blocking multiple growth factor signaling pathways and their complex interactive network (Cao et al., 2008; Ikuta et al., 2009; Homsí & Daud, 2007; Lieu et al., 2011).

Author details

Loredana Albonici, Camilla Palumbo and Vittorio Manzari

Department of Experimental Medicine and Biochemical Sciences, University of Rome "Tor Vergata", Rome, Italy

7. References

- Adachi, Y., Aoki, C., Yoshio-Hoshino, N., Takayama, K., Curiel, D. T. & Nishimoto N. (2006). Interleukin-6 induces both cell growth and VEGF production in malignant mesotheliomas. *Int. J. Cancer*, Vol. 119, No. 6, (September 2006), pp. 1303-1311, ISSN 0020-7136
- Adini, A., Kornaga, T., Firoozbakht, F. & Benjamin, L. E. (2002). Placenta growth factor is a survival factor for human endothelial cells and macrophages. *Cancer Res.*, Vol. 62, No. 10, (May 2002), pp. 2749-2752, ISSN 0008-5472
- Albonici, L., Doldo, E., Palumbo, C., Orlandi, A., Bei, R., Pompeo, E., Mineo, T. C., Modesti, A. & Manzari, V. (2009). Placenta growth factor is a survival factor for human

- malignant mesotelioma cells. *Int. J. Immunopathol. Pharmacol.*, Vol. 22, No. 2, (April-June 2009), pp. 389-401, ISSN 0394-6320
- Andrae, J., Gallini, R. & Betsholtz, C. (2008). Role of platelet-derived growth factors in physiology and medicine. *Genes Dev.*, Vol. 22, No. 10, (May 2008), pp. 1276-1312, ISSN 0890-9369
- Angelo, L. S. & Kurzrock, R. (2007). Vascular endothelial growth factor and its relationship to inflammatory mediators. *Clin. Cancer Res.*, Vol. 13, No. 10, (May 2007), pp. 2825-2830, ISSN 1078-0432
- Antony, V. B., Godbey, S. W., Kunkel, S. L., Hott, J. W., Hartman, D. L., Burdick, M. D. & Strieter, R. M. (1993). Recruitment of inflammatory cells to the pleural space. Chemotactic cytokines, IL-8, and monocyte chemotactic peptide-1 in human pleural fluids. *J. Immunol.*, Vol. 151, No. 12, (December 1993), pp. 7216-7223, ISSN 0022-1767
- Asahara, T., Takahashi, T., Masuda, H., Kalka, C., Chen, D., Iwaguro, H., Inai, Y., Silver, M. & Isner, J. M. (1999). VEGF contributes to postnatal neovascularization by mobilizing bone marrow-derived endothelial progenitor cells. *EMBO J.*, Vol. 18, No. 14, (July 1999), pp. 3964-3972, ISSN 0261-4189
- Astoul, P. (1999). Pleural mesothelioma. *Curr. Opin. Pulm. Med.*, Vol. 5, No. 4, (July 1999), pp. 259-268, ISSN 1070-5287
- Attanoos, R. L. & Gibbs, A. R. (1997). Pathology of malignant mesothelioma. *Histopathology*, Vol. 30, No. 5, (May 1997), pp. 403-418, ISSN 0309-0167
- Aust, A. E., Cook, P.M. & Dodson, R. F. (2011). Morphological and chemical mechanisms of elongated mineral particle toxicities. *J. Toxicol. Environ. Health. B Crit. Rev.*, Vol. 14, No. 1-4, pp. 40-75, ISSN 1093-7404
- Autiero, M., Waltenberger, J., Communi, D., Kranz, A., Moons, L., Lambrechts, D., Kroll, J., Plaisance, S., De Mol, M., Bono, F., Kliche, S., Fellbrich, G., Ballmer-Hofer, K., Maglione, D., Mayr-Beyrle, U., Dewersch, M., Dombrowski, S., Stanimirovic, D., Van Hummelen, P., Dehio, C., Hicklin, D. J., Persico, G., Herbert, J. M., Communi, D., Shibuya, M., Collen, D., Conway, E. M. & Carmeliet, P. (2003). Role of PlGF in the intra- and intermolecular cross-talk between the VEGF receptors Flt-1 and Flk-1. *Nat. Med.*, Vol. 9, No. 7, (July 2003), pp. 936-943, ISSN 1078-8956
- Baldys, A., Pande, P., Mosleh, T., Park, S. H. & Aust, A. E. (2007). Apoptosis induced by crocidolite asbestos in human lung epithelial cells involves inactivation of Akt and MAPK pathways. *Apoptosis*, Vol. 12, No. 2, (February 2007), pp. 433-447, ISSN 1360-8185
- Barillari, G., Albonici, L., Franzese, O., Modesti, A., Liberati, F., Barillari, P., Ensoli, B., Manzari, V. & Santeusano, G. (1998). The basic residues of placenta growth factor type 2 retrieve sequestered angiogenic factors into soluble form. Implication for tumor angiogenesis. *Am. J. Pathol.*, Vol. 152, No. 5, (May 1998), pp. 1161-1166, ISSN 0002-9440
- Barleon, B., Sozzani, S., Zhou, D., Weich, H. A., Mantovani, A. & Marmé, D. (1996). Migration of human monocytes in response to vascular endothelial growth factor (VEGF) is mediated via the VEGF receptor flt-1. *Blood*, Vol. 87, No. 8, (April 1996), pp. 3336-3343, ISSN 0006-4971

- Berman, D. W. & Crump, K. S. (2008). A meta-analysis of asbestos-related cancer risk that addresses fiber size and mineral type. *Crit. Rev. Toxicol.*, Vol. 38, Suppl. 1, pp. 49–73, ISSN 1040-8444
- BéruBé, K. A., Quinlan, T. R., Fung, H., Magae, J., Vacek, P., Taatjes, D. J. & Mossman, B. T. (1996). Apoptosis is observed in mesothelial cells after exposure to crocidolite asbestos. *Am. J. Respir. Cell. Mol. Biol.*, Vol. 15, No. 1, (July 1996), pp. 141-147, ISSN 1044-1549
- Bocchetta, M., Di Resta, I., Powers, A., Fresco, R., Tosolini, A., Testa, J. R., Pass, H.I., Rizzo, P. & Carbone, M. (2000). Human mesothelial cells are unusually susceptible to simian virus 40-mediated transformation and asbestos cocarcinogenicity. *Proc. Natl. Acad. Sci. USA*, Vol. 97, No. 18, (August 2000), pp. 10214–10219, ISSN 0027-8424
- Bowie, A. & O'Neill, L. A. (2000). Oxidative stress and nuclear factor-kappaB activation: a reassessment of the evidence in the light of recent discoveries. *Biochem. Pharmacol.*, Vol. 59, No. 1, (January 2000), pp. 13-23, ISSN 0006-2952
- Branchaud, R. M., Garant, L. J. & Kane A. B. (1993). Pathogenesis of mesothelial reactions to asbestos fibers. Monocyte recruitment and macrophage activation. *Pathobiology*, Vol. 61, No. 3-4, pp. 154–163, ISSN 1015-2008
- Broadbush, V. C., Yang, L., Scavo, L. M., Ernst, J. D. & Boylan, A. M. (1996). Asbestos induces apoptosis of human and rabbit pleural mesothelial cells via reactive oxygen species. *J. Clin. Invest.*, Vol. 98, No. 9, (November 1996), pp. 2050–2059, ISSN 0021-9738
- Brown, D., Beswick, P. & Donaldson, K. (1999). Induction of nuclear translocation of NF- κ B in epithelial cells by respirable mineral fibres. *J. Pathol.*, Vol. 189, No. 2, (October 1999), pp. 258–264, ISSN 0022-3417
- Burt, B. M., Rodig, S. J., tilleman, T. R., Elbardissi, A. W., Bueno, R., Sugarbaker, D. J. (2011). Circulating and tumor-infiltrating myeloid cells predict survival in human pleural mesothelioma. *Cancer*, Vol. 117, No. 22, (November 2011), pp. 5234-5244, ISSN 0008-543X
- Cacciotti, P., Strizzi, L., Vianale, G., Iaccheri, L., Libener, R., Porta, C., Tognon, M., Gaudino, G. & Mutti, L. (2002). The presence of simian-virus 40 sequences in mesothelioma and mesothelial cells is associated with high levels of vascular growth factor. *Am. J. Respir. Cell. Mol. Biol.*, Vol. 26, No. 2, (February 2002), pp. 189-193, ISSN 1044-1549
- Cao, Y., Cao, R. & Hedlund, E. M. (2008). R Regulation of tumor angiogenesis and metastasis by FGF and PDGF signaling pathways. *J Mol. Med. (Berl.)*, Vol. 86, No. 7, (July 2008), pp. 785-789, ISSN 0946-2716
- Cao, Y., Chen, H., Zhou, L., Chiang, M. K., Anand-Apte, B., Weatherbee, J. A., Wang, Y., Fang, F., Flanagan, J. G. & Tsang, M. L. (1996). Heterodimers of placenta growth factor/vascular endothelial growth factor. Endothelial activity, tumor cell expression, and high affinity binding to Flk-1/KDR. *J. Biol. Chem.*, Vol. 271, No. 6, (February 1996), pp. 3154-3162, ISSN 0021-9258
- Carbone, M., Kratzke, R. A. & Testa, J. R. (2002). The pathogenesis of mesothelioma. *Semin. Oncol.*, Vol. 29, No. 1, (February 2002), pp. 2-17, ISSN 0093-7754
- Carbone, M., Ly, B. H., Dodson, R. F., Pagano, I., Morris, P. T., Dogan, U. A., Gazdar, A. F., Pass, H. I. & Yang, H. (2012). Malignant mesothelioma: Facts, myths and hypotheses. *J. Cell. Physiol.*, Vol. 227, No. 1 (January 2012), pp. 44-58, ISSN 0021-9541

- Carmeliet, P. (2003). Angiogenesis in health and disease. *Nat. Med.*, Vol. 9, No. 6, (June 2003), pp. 653-660, ISSN 1078-8956
- Carmeliet, P., Moons, L., Luttun, A., Vincenti, V., Compernelle, V., De Mol, M., Wu, Y., Bono, F., Devy, L., Beck, H., Scholz, D., Acker, T., DiPalma, T., Dewerchin, M., Noel, A., Stalmans, I., Barra, A., Blacher, S., Vandendriessche, T., Ponten, A., Eriksson, U., Plate, K. H., Foidart, J. M., Schaper, W., Charnock-Jones, D. S., Hicklin, D. J., Herbert, J. M., Collen, D. & Persico, M. G. (2001). Synergism between vascular endothelial growth factor and placental growth factor contributes to angiogenesis and plasma extravasation in pathological conditions. *Nat. Med.*, Vol. 7, No. 5, (May 2001), pp. 575-583, ISSN 1078-8956
- Catalano, A., Lazzarini, R., Di Nuzzo, S., Orciari, S. & Procopio A. (2009). The plexin-A1 receptor activates vascular endothelial growth factor-receptor 2 and nuclear factor-kappaB to mediate survival and anchorage-independent growth of malignant mesothelioma cells. *Cancer Res.*, Vol. 69, No. 4, (February 2009), pp. 1485-1493, ISSN 0008-5472
- Catalano, A., Romano, M., Martinotti, S. & Procopio, A. (2002). Enhanced expression of vascular endothelial growth factor (VEGF) plays a critical role in the tumor progression potential induced by simian virus 40 large T antigen. *Oncogene*, Vol. 21, No. 18 (April 2002), pp. 2896-2900, ISSN 0950-9232
- Chang, L. & Karin, M. (2001). Mammalian MAP kinase signalling cascades. *Nature*, Vol. 410, No. 6824, (March 2001), pp. 37-40, ISSN 0028-0836
- Chen, C. N., Hsieh, F. J., Cheng, Y. M., Cheng, W. F., Su, Y. N., Chang, K. J. & Lee, P. H. (2004). The significance of placenta growth factor in angiogenesis and clinical outcome of human gastric cancer. *Cancer Lett.*, Vol. 213, No. 1, (September 2004), pp. 73-82, ISSN 0304-3835
- Cheng, N., Shi, X., Ye, J., Castranova, V., Chen, F., Leonard, S. S., Vallyathan, V. & Rojanasakul, Y. (1999). Role of transcription factor NF-kappaB in asbestos-induced TNFalpha response from macrophages. *Exp. Mol. Pathol.*, Vol. 66, No. 3, (August 1999), pp. 201-210, ISSN 0014-4800
- Choe, N., Tanaka, S., Xia, W., Hemenway, D. R., Roggli, V. L. & Kagan, E. (1997). Pleural macrophage recruitment and activation in asbestos-induced pleural injury. *Environ. Health Perspect.*, Vol. 105, Suppl. 5, (September 1997), pp. 1257-1260, ISSN 0091-6765
- Choi, C. H., Song, S. Y., Choi, J. J., Park, Y. A., Kang, H., Kim, T. J., Lee, J. W., Kim, B. G., Lee, J. H. & Bae, D. S. (2008). Prognostic significance of VEGF expression in patients with bulky cervical carcinoma undergoing neoadjuvant chemotherapy. *BMC Cancer*, Vol. 8, (October 2008), p. 295, ISSN 1471-2407
- Costa, C., Incio, J. & Soares, R. (2007). Angiogenesis and chronic inflammation: cause or consequence? *Angiogenesis*, Vol. 10, No. 3, pp. 149-166, ISSN 0969-6970
- Coussens, L. M. & Werb, Z. (2002). Inflammation and cancer. *Nature*, Vol. 420, No. 6917, (December 2002), pp. 860-867, ISSN 0028-0836
- David Dong, Z. M., Aplin, A. C. & Nicosia, R. F. (2009). Regulation of angiogenesis by macrophages, dendritic cells, and circulating myelomonocytic cells. *Curr. Pharm. Des.*, Vol. 15, No. 4, pp. 365-379, ISSN 1381-6128

- Davidson, B., Vintman, L., Zcharia, E., Bedrossian, C., Berner, A., Nielsen, S., Ilan, N., Vlodavsky, I & Reich, R. (2004). Heparanase and basic fibroblast growth factor are co-expressed in malignant mesothelioma. *Clin. Exp. Metastasis*, Vol. 21, No. 5, pp. 469-476, ISSN 0262-0898
- Dawson, M. R., Duda, D. G., Fukumura, D. & Jain, R. K. (2009). VEGFR1-activity-independent metastasis formation. *Nature*, Vol. 461, No. 7262, (September 2009), pp. E4-E5, ISSN 0028-0836.
- Ding, Y., Huang, Y., Song, N. Gao, X., Yuan, S., Wang, X., Cai, H., Fu, Y. & Luo, Y. (2010). NFAT1 mediates placental growth factor-induced myelomonocytic cell recruitment via the induction of TNF-alpha. *J. Immunol.*, Vol. 184, No. 5, (March 2010), pp. 2593-2601, ISSN 0022-1767
- Dong, H. Y., Buard, A., Renier, A., Levy, F., Saint-Etienne, L. & Jaurand, M. C. (1994). Role of oxygen derivatives in the cytotoxicity and DNA damage produced by asbestos on rat pleural mesothelial cells in vitro. *Carcinogenesis*, Vol. 15, No. 6, (June 1994), pp. 1251-1255, ISSN 0143-3334
- Dorai, T., Kobayashi, H.; Holland, J. F. & Onhuma, T. (1994). Modulation of platelet-derived growth factor-beta mRNA expression and cell growth in a human mesothelioma cell line by a hammerhead ribozyme. *Mol. Pharmacol.*, Vol. 46, No. 3, (September 1994, pp. 437-444, ISSN 0026-895X
- Driscoll, K., Carter, J., Howard, B., Hassenbein, D., Janssen, Y. & Mossman, B. T. (1998). Crocidolite activates NF- κ B and MIP-2 gene expression in rat alveolar epithelial cells. Role of mitochondrial-derived oxidants. *Environ. Health Perspect.*, Vol. 106, Suppl. 5, (October 1998), pp. 1171-1174, ISSN 0091-6765
- Eferl, R., & Wagner, E. F.(2003). AP1: a double-edged sword in tumorigenesis. *Nature Rev. Cancer*, Vol. 3, No. 11, (November 2003), pp. 859-868, ISSN 1474-175X
- Fennell, D. A. & Rudd, R. M. (2004). Defective core-apoptosis signaling in diffuse malignant pleural mesothelioma: opportunities for effective drug development. *Lancet Oncol.*, Vol. 5, No. 6, (June 2004), pp. 354-362, ISSN 1470-2045
- Ferrara, N., Gerber, H. P. & LeCouter, J. (2003). The biology of VEGF and its receptors. *Nat. Med.*, Vol. 9, No. 6, (June 2003), pp. 669-676, ISSN 1078-8956
- Filiberti, R., Marroni, P., Neri, M., Ardizzoni, A., Betta P. G., Cafferata, M. A., Canessa, P. A., Puntoni, R., Ivaldi, G. P., & Paganuzzi, M. (2005). Serum PDGF-AB in pleural mesotelioma. *Tumour Biol.*, Vol. 26, No. 5, (September-October 2005), pp. 221-226, ISSN 1010-4283
- Fischer, C., Jonckx, B., Mazzone, M., Zacchigna, S., Loges, S., Pattarini, L., Chorianopoulos, E., Liesenborghs, L., Kock M., De Mol, M., Autiero, M., Wyns, S., Plaisance, S., Moons, L., van Rooijen, N., Giacca, M., Stassen J. M., Dewerchin, M., Collen, D. & Carmeliet, P. (2007). Anti-PlGF inhibits growth of VEGF(R)-inhibitor-resistant tumors without affecting healthy vessels. *Cell*, Vol. 131, No. 3, (November 2007), pp. 463-475, ISSN 0092-8674
- Fischer, C., Mazzone, M., Jonckx, B. & Carmeliet, P. (2008). FLT1 and its ligands VEGFB and PlGF: drug targets for anti-angiogenic therapy? *Nat. Rev. Cancer.*, Vol. 8, No. 12, (December 2008), pp. 942-956, ISSN 1474-175X

- Fleury-Feith, J., Pilatte, Y. & Jaurand, M. C. (2003). Cells in the pleural cavity, In: *Textbook of pleural diseases*, Light, R. W. & Lee, Y. C. G., pp. 17-34, Arnold Publishers, ISBN 9780340807941, London.
- Folkman, J. (2006). Angiogenesis. *Annu. Rev. Med.*, Vol. 57, pp. 1-18, ISSN 0066-4219
- Gabrilovich, D. I., Chen, H. L., Girgis, K. R., Cunningham, H. T., Meny, G. M., Nadaf, S., Kavanaugh, D. & Carbone, D. P. (1996). Production of vascular endothelial growth factor by human tumors inhibits the functional maturation of dendritic cells. *Nat. Med.*, Vol. 2, No. 10, (October 1996), pp. 1096-1103, ISSN 1078-8956
- Garlepp, M. J & Leong, C. C. (1995). Biological and immunological aspects of malignant mesothelioma. *Eur. Respir. J.*, Vol. 8, No. 4, (April 1995), pp. 643-650, ISSN 0903-1936
- Gasparini, G. & Harris, A. L. (1995). Clinical importance of the determination of tumor angiogenesis in breast carcinoma: much more than a new prognostic tool. *J. Clin. Oncol.*, Vol. 13, No. 3, (March 1995), pp. 765-782, ISSN 0732-183X
- Gazdar, A. F. & Carbone, M. (2003). Molecular pathogenesis of malignant mesothelioma and its relationship to simian virus 40. *Clin. Lung Cancer*, Vol. 5, No. 3, (November 2003), pp. 177-181, ISSN 1525-7304
- Gossart, S., Cambon, C., Orfila, C., Séquélas, M. H., Lepert, J. C., Rami, J., Carrè, P. & Pipy, B. (1996). Reactive oxygen intermediates as regulators of TNF- α production in rat lung inflammation induced by silica. *J. Immunol.*, Vol. 156, No. 4, (February 1996), pp. 1540-1548, ISSN 0022-1767
- Greillier, L. & Astoul, P. (2008). Mesothelioma and asbestos-related pleural diseases. *Respiration*, Vol. 76, No. 1, pp. 1-15, ISSN 0025-7931
- Gulumian, M. (2005). An update on the detoxification processes for silica particles and asbestos fibers: success and limitations. *J. Toxicol. Environ. Health. B Crit. Rev.*, Vol. 8, No. 6, (November-December 2005), pp. 453-483, ISSN 1093-7404
- Haegens, A., Barrett, T. F., Gell, J., Shukla, A., Macpherson, M., Vacek, P., Poynter, M. E., Butnor, K. J., Janssen-Heininger, Y. M., Steele, C. & Mossman, B. T. (2007). Airway epithelial NF- κ B activation modulates asbestos-induced inflammation and mucin production in vivo. *J. Immunol.*, Vol. 178, No. 3, (February 2007), pp. 1800-1808, ISSN 0022-1767
- Harmey, J. H. & Bouchier-Hayes, D. (2002). Vascular endothelial growth factor (VEGF), a survival factor for tumour cells: implications for anti-angiogenic therapy. *Bioessays*, Vol. 24, No. 3, (March 2002), pp. 280-283, ISSN 0265-9247
- Hattori, K., Heissig, B., Wu, Y., Dias, S., Tejada, R., Ferris, B., Hicklin, D. J., Zhu, Z., Bohlen, P., Witte, L., Hendriks, J., Hackett, N. R., Crystal, R. G., Moore, M. A., Werb, Z., Lyden, D. & Rafii, S. (2002). Placental growth factor reconstitutes hematopoiesis by recruiting VEGFR1(+) stem cells from bone-marrow microenvironment. *Nat. Med.*, Vol. 8, No. 8, (August 2002), pp. 841-849, ISSN 1078-8956
- Hausmann, M. J., Rogachev, B., Weiler, M., Chaimovitz, C. & Douvdevani, A. (2000). Accessory role of human peritoneal mesothelial cells in antigen presentation and T-cell growth. *Kidney Int.*, Vol. 57, No. 2, (February 2000), pp. 476-86, ISSN 0085-2538
- Hayden, M. S. & Gosh, S. (2004). Signaling to NF- κ B. *Genes Dev.*, Vol. 18, No. 18, (September 2004), pp. 2195-2224, ISSN 0890-9369

- Hayden, M. S. & Ghosh, S. (2008). Shared principles in NF-kappaB signaling. *Cell*, Vol. 132, No. 3, (February 2008), pp. 344–362, ISSN 0092-8674
- Heintz, N. H., Janssen, Y. M. & Mossman, B. T. (1993). Persistent induction of c-fos and c-jun expression by asbestos. *Proc. Natl. Acad. Sci. USA*, Vol. 90, No. 8, (April 1993), pp. 3299–3303, ISSN 0027-8424
- Heintz, N. H., Janssen-Heininger, Y. M. & Mossman, B. T. (2010). Asbestos, lung cancers, and mesotheliomas: from molecular approaches to targeting tumor survival pathways. *Am. J. Respir. Cell Mol. Biol.*, Vol. 42, No. 2, (February 2010), pp. 133–139, ISSN 1044-1549
- Hillegass, J. M., Shukla, A., Lathrop, S. A., MacPherson, M. B., Beuschel, S. L., Butnor, K. J., Testa, J. R., Pass, H. I., Carbone, M., Steele, C. & Mossman, B. T. (2010). Inflammation precedes the development of human malignant mesotheliomas in a SCID mouse xenograft model. *Ann. N. Y. Acad. Sci.*, Vol. 1203, (August 2010), pp. 7-14, ISSN 0077-8923
- Hirayama, N., Tabata, C., Tabata, R., Maeda, R., Yasumitsu, A., Yamada, S., Kuribayashi, K., Fukuoka, K. & Nakano T. (2011). Pleural effusion VEGF levels as a prognostic factor of malignant pleural mesothelioma. *Respir. Med.*, Vol. 105, No. 1 (January 2011), pp. 137-142, ISSN 0954-6111
- Hoffmann, A., Natoli, G. & Baltimore, D. (2003). Genetic analysis of NF-kappaB/Rel transcription factor defines functional specificities. *EMBO J.*, Vol. 22, No. 20, (October 2003), pp. 5530-5539, ISSN 0261-4189
- Hoffmann, A., Natoli, G. & Ghosh, G. (2006). Transcriptional regulation via the NF-kappaB signaling module. *Oncogene*, Vol. 25, No. 51, (October 2006), pp. 6706-6716, ISSN 0950-9232
- Hoidal, J. R. (2001). Reactive oxygen species and cell signaling. *Am. J. Respir. Cell Mol. Biol.*, Vol. 25, No. 6, (December 2001), pp. 661–663, ISSN 1044-1549
- Holmes, D. I. & Zachary, I. (2004). Placental growth factor induces FosB and c-Fos gene expression via Flt-1 receptors. *FEBS Lett.*, Vol. 557, No. 1-3, (January 2004), pp. 93-98, ISSN 0014-5793
- Homsy, J. & Daud, A. I. (2007). Spectrum of activity and mechanism of action of VEGF/PDGF inhibitors. *Cancer Control*, vol. 14, No. 3, (July 2007), pp. 285-294, ISSN 1073-2748
- Huang, S., Robinson, J. B., Deguzman, A., Bucana, C. D. & Fidler, I. J. (2000). Blockade of nuclear factor-kappaB signaling inhibits angiogenesis and tumorigenicity of human ovarian cancer cells by suppressing expression of vascular endothelial growth factor and interleukin 8. *Cancer Res.*, Vol. 60, No. 19, (October 2000), pp. 5334–5539, ISSN 0008-5472
- Huang, S. X., Jaurand, M. C., Kamp, D. W., Whysner, J. & Hei, T. K. (2011). Role of mutagenicity in asbestos fiber-induced carcinogenicity and other diseases. *J. Toxicol. Environ. Health B. Crit. Rev.*, Vol. 14, No. 1-4, pp. 179-245, ISSN 1093-7404
- Hussain, S. P., Hofseth, L. J. & Harris, C. C. (2003). Radical causes of cancer. *Nat. Rev. Cancer.*, Vol. 3, No. 4, (April 2003), pp. 276-285, ISSN 1474-175X
- Ikuta, K., Yano, S., Trung, V. T., Hanibuchi, M., Goto, H., Li, Q., Wang, W., Yamada, T., Ogino, H., Kakiuchi, S., Uehara, H., Sekido, Y., Uenaka, T., Nishioka, Y & Sone, S. (2009). E7080, a multi-tyrosine kinase inhibitor, suppresses the progression of malignant

- pleural mesothelioma with different proangiogenic cytokine production profiles. *Clin. Cancer Res.*, Vol. 15, No. 23, (December 2009), pp. 7229-7237, ISSN 1078-0432
- Izzi, V., Chiurchiù, V., D'Aquilio, F., Palumbo, C., Tresoldi, I., Modesti, A. & Baldini, P. M. (2009). Differential effects of malignant mesothelioma cells on THP-1 monocytes and macrophages. *Int. J. Oncol.*, Vol. 34, No. 2, (February 2009), pp. 543-550, ISSN 1019-6439
- Janssen-Heininger, Y. M., Macara, I. & Mossman, B. T. (1999). Cooperativity between oxidants and tumor necrosis factor in the activation of nuclear factor (NF)-kappaB: requirement of Ras/mitogen-activated protein kinases in the activation of NF-kappaB by oxidants. *Am. J. Respir. Cell. Mol. Biol.*, Vol. 20, No. 5, (May 1999), pp. 942-952, ISSN 1044-1549
- Janssen, Y. M., Barchowsky, A., Treadwell, M., Driscoll, K. E. & Mossman, B. T. (1995). Asbestos induces nuclear factor- κ B (NF- κ B) DNA-binding activity and NF- κ B-dependent gene expression in tracheal epithelial cells. *Proc. Natl. Acad. Sci. USA*, Vol. 92, No. 18, (August 1995), pp. 8458-8462, ISSN 0027-8424
- Janssen, Y. M., Driscoll, K. E., Howard, B., Quinlan, T. R., Treadwell, M., Barchowsky, A. & Mossman, B. T. (1997). Asbestos causes translocation of p65 protein and increases NF-kappa B DNA binding activity in rat lung epithelial and pleural mesothelial cells. *Am. J. Pathol.*, Vol. 151, No. 2, (August 1997), pp. 389-401, ISSN 0002-9440
- Jash, A., Sahoo, A., Kim, G. C., Chae, C. S., Hwang, J. S., Kim, J. E. & Im, S. H. (2012). Nuclear factor of activated T cells 1 (NFAT1) induced permissive chromatin modification facilitates nuclear Factor- κ B (NF- κ B) mediated interleukin-9 (IL-9) transactivation. *J. Biol. Chem.*, (March 2012), Epub ahead of print, ISSN 1083-351X
- Jochum, W., Passequé, E. & Wagner, E. F. (2001). AP-1 in mouse development and tumorigenesis. *Oncogene*, Vol. 20, No. 19, (April 2001), pp. 2401-2412, ISSN 0950-9232
- Jung, Y. J., Isaacs, J. S., Lee, S., Trepel, J. & Neckers, L. (2003). IL-1beta-mediated up-regulation of HIF-1alpha via an NF-kappaB/COX-2 pathway identifies HIF-1 as a critical link between inflammation and oncogenesis. *FASEB J.*, Vol. 17, No. 14, (November 2003), pp. 2115-2117, ISSN 0892-6638
- Kamp, D. W. & Weitzman, S. A. (1999). The molecular basis of asbestos induced lung injury. *Thorax*, Vol. 54, No. 7, (July 1999), pp. 638-652, ISSN 0040-6376
- Kanwar, J. R., Kamalapuram, S. K. & Kanwar, R. K. (2011). Targeting surviving in cancer: the cell-signalling perspective. *Drug Discov. Today*, Vol. 16, No. 11-12, (June 2011), pp. 485-494, ISSN 1359-6446
- Kerbel, R. S. (2008). Tumor angiogenesis. *N. Engl. J. Med.*, Vol. 358, No. 19, (May 2008), pp. 2039-2049, ISSN 0028-4793
- Kerber, M., Reiss, Y., Wickersheim, A., Jugold, M., Kiessling, F., Heil, M., Tchaikovski, V., Waltenberger, J., Shibuya, M., Plate, K. H. & Machein, M. R. (2008). Flt-1 signaling in macrophages promotes glioma growth in vivo. *Cancer Res.*, Vol. 68, No. 18, (September 2008), pp. 7342-7351, ISSN 0008-5472
- Khaliq, A., Li, X. F., Shams, M., Sisi, P., Acevedo, C. A., Whittle, M. J., Weich, H. & Ahmed, A. (1996). Localization of placenta growth factor (PlGF) in human term placenta. *Growth Factors*, Vol. 13, No. 3-4, pp. 243-250, ISSN 0897-7194

- Knights, V. & Cook, S. J. (2010). De-regulated FGF receptors as therapeutic targets in cancer. *Pharmacol. Ther.*, Vol. 125, No. 1, (January 2010), pp. 105-117, ISSN 0163-7258
- Koch, S., Tugues, S., Li, X., Gualandi, L. & Claesson-Welsh, L. (2011). Signal transduction by vascular endothelial growth factor receptors. *Biochem J.*, Vol. 437, No. 2, (July 2011), pp. 169-183, ISSN 0264, 6021
- Kothmaier, H., Quehenberger, F., Halbewedl, I., Morbini, P., Demirag, F., Zeren, H., Comin, C. E., Murer, B., Cagle, P. T., Attanoos, R., Gibbs, A. R., Galateau-Salle, F. & Popper, H. H. (2008). EGFR and PDGFR differentially promote growth in malignant epithelioid mesothelioma of short and long term survivors. *Thorax*, Vol. 63, No. 4, (April 2008), pp. 345-351, ISSN 0040-6376
- Kumar, S., Guleria, R., Singh, V., Bharti, A. C., Mohan, A. & Das, B. C. (2009). Efficacy of plasma vascular endothelial growth factor in monitoring first-line chemotherapy in patients with advanced non-small cell lung cancer. *BMC Cancer*, Vol. 9, (December 2009), p. 421, ISSN 1471-2407
- Kumar-Singh, S., Vermuelen, P. B., Weyler, J., Segers, K., Wejn, B., Van Daele, A., Dirix, L. Y., Van Oosterom, A. T. & Van Mark, E. (1997). Evaluation of tumor angiogenesis as a prognostic marker in malignant mesothelioma. *J. Pathol.*, Vol. 182, No. 2, (June 1997), pp. 211-216, ISSN 0022-3417
- Kumar-Singh, S., Weyler, J., Martin, M. J., Vermeulen, P. B. & Van Marck, E. (1999). Angiogenic cytokines in mesothelioma: a study of VEGF, FGF-1 and -2, and TGF beta expression. *J. Pathol.*, Vol. 189, No. 1, (September 1999), pp. 72-82, ISSN 0022-3417
- Langerak, A. W., van der Linden-van Beurden, C. A. & Versnel, M. A. (1996a). Regulation of differential expression of platelet-derived growth factor alpha- and beta-receptor mRNA in normal and malignant human mesothelial cell lines. *Biochim. Biophys. Acta.*, Vol. 1305, No. 1-2, (February 1996), pp. 63-70, ISSN 0006-3002
- Langerak, A. W., De Laat, P. A., Van Der Linden-Van Beurden, C. A., Delahaye, M., Van Der Kwast, T. H., Hoogsteden, H. C., Benner, R. & Versnel, M. A. (1996b). Expression of platelet-derived growth factor (PDGF) and PDGF receptors in human malignant mesothelioma in vitro and in vivo. *J. Pathol.*, Vol. 178, No. 2, (February 1996), pp. 151-160, ISSN 0022-3417
- Lawrence, T. (2011). Macrophages and NF- κ B in cancer. *Curr. Top. Microbiol. Immunol.*, Vol. 349, pp. 171-184, ISSN 0070-217X
- Lemaire, I. & Ouellet, S. (1996). Distinctive profile of alveolar macrophage-derived cytokine release induced by fibrogenic and nonfibrogenic mineral dusts. *J. Toxicol. Environ. Health.*, Vol. 47, No. 5, (April 1996), pp. 465-478, ISSN 0098-4108
- Le Page, C., Koumakpayi, I. H., Lessard, L., Mes-Masson, A. M. & Saad, F. (2005). EGFR and Her-2 regulate the constitutive activation of NF-kappaB in PC-3 prostate cancer cells. *Prostate*, Vol. 65, No. 2, (October 2005), pp. 130-140, ISSN 0270-4137
- Li, J., Huang, B., Shi, X., Castranova, V., Vallyathan, V. & Huang, C. (2002). Involvement of hydrogen peroxide in asbestos-induced NFAT activation. *Mol. Cell. Biochem.*, Vol. 234-235, No. 1-2, (May-June 2002), pp. 161-168, ISSN 0300-8177
- Li, Q., Wang, W., Yamada, T., Matsumoto, K., Sakai, K., Bando, Y., Uehara, H., Nishioka, Y., Sone, S., Iwakiri, S., Itoi, K., Utsugi, T., Yasumoto, K. & Yano, S. (2011). Pleural

- mesothelioma instigates tumor-associated fibroblasts to promote progression via a malignant cytokine network. *Am. J. Pathol.*, Vol. 179, No. 3, (September 2011), pp. 1483-1493, ISSN 0002-9440
- Lieu, C., Heymach, J., Overman, M., Tran, H. & Kopetz, S. (2011). Beyond VEGF: inhibition of the fibroblast growth factor pathway and antiangiogenesis. *Clin. Cancer Res.*, Vol. 17, No. 19, (October 2011), pp. 6130-6139, ISSN 1078-0432
- Lin, Y. L., Liang, Y. C. & Chiang, B. L. (2007). Placental growth factor down-regulates type 1 T helper immune response by modulating the function of dendritic cells. *J. Leukoc. Biol.*, Vol. 82, No. 6, (December 2007), pp. 1473-1480, ISSN 0741-5400
- Lin, W. W. & Karin, M. (2007). A cytokine-mediated link between innate immunity, inflammation, and cancer. *J. Clin. Invest.*, Vol. 117, No. 5, (May 2007), pp. 1175-1182, ISSN 0021-9738
- Liu, W., Ernst, J. D. & Broaddus, V. C. (2000). Phagocytosis of crocidolite asbestos induces oxidative stress, DNA damage, and apoptosis in mesothelial cells. *Am. J. Respir. Cell. Mol. Biol.*, vol. 23, No. 3, (September 2000), pp. 371-378, ISSN 1044-1549
- Liu, Z. & Klominek, J. (2003). Regulation of matrix metalloprotease activity in malignant mesothelioma cell lines by growth factors. *Thorax*, Vol. 58, No. 3, (March 2003), pp. 198-203, ISSN 0040-6376
- Loges, S., Schmidt, T. & Carmeliet, P. (2009). "Antimyoangiogenic" therapy for cancer by inhibiting PlGF. *Clin. Cancer Res.*, Vol. 15, No. 11, (June 2009), pp. 3648-3653, ISSN 1078-0432
- Luttun, A., Tjwa, M. & Carmeliet, P. (2002). Placenta growth factor (PlGF) and its receptor Flt-1 (VEGFR-1): novel therapeutic targets for angiogenic disorders. *Ann. N. Y. Acad. Sci.*, Vol. 979, (December 2002), pp. 80-93, ISSN 0077-8923
- Macian, F. (2005). NFAT proteins: key regulators of T-cell development and function. *Nature Rev. Immunol.*, Vol. 5, No. 6, (June 2005), pp. 472-484, ISSN 1474-1733.
- Macián, F., López-Rodríguez, C., & Rao, A. (2001). Partners in transcription: NFAT and AP-1. *Oncogene*, Vol. 20, No. 19, (April 2001), pp. 2476-2489, ISSN 0950-9232
- Maglione, D., Guerriero, V., Viglietto, G., Delli-Bovi, P. & Persico, M. G. (1991). Isolation of a human placenta cDNA coding for a protein related to the vascular permeability factor. *Proc. Natl. Acad. Sci. USA*, Vol. 88, No. 20, (October 1991), pp. 9267-9271, ISSN 0027-8424
- Mancini, M. & Toker, A. (2009). NFAT proteins: emerging roles in cancer progression. *Nature Rev. Cancer.*, Vol. 9, No. 11, (November 2009), pp. 810-820, ISSN 1474-175X
- Manning, C. B., Vallyathan, V. & Mossman, B. T. (2002). Diseases caused by asbestos: mechanisms of injury and disease development. *Int. Immunopharmacol.*, Vol. 2, No. 2-3, (February 2002), pp. 191-200, ISSN 1567-5769
- Manning, L. S., Davis, M. R. & Robinson, B. W. (1991). Asbestos fibres inhibit the in vitro activity of lymphokine-activated killer (LAK) cells from healthy individuals and patients with malignant mesothelioma. *Clin. Exp. Immunol.*, Vol. 83, No. 1, (January 1991), pp. 85-91, ISSN 0009-9104
- Mantovani, A. (2010). Molecular pathways linking inflammation and cancer. *Curr. Mol. Med.*, Vol. 10, No. 4, (June 2010), pp. 369-373, ISSN 1566-5240

- Mantovani, A., Allavena, P., Sica, A., & Balkwill, F. (2008). Cancer-related inflammation. *Nature*, Vol. 454, No. 7203, (July 2008), pp. 436-444, ISSN 0028-0836
- Marcellini, M., De Luca, N., Riccioni, T., Ciucci, A., Orecchia, A., Lacal, P. M., Ruffini, F., Pesce, M., Cianfarani, F., Zambruno, G., Orlandi, A. & Failla, M. C. (2006). Increased melanoma growth and metastasis spreading in mice overexpressing placenta growth factor. *Am. J. Pathol.*, Vol. 169, No. 2, (August 2006), pp. 643-654, ISSN 0002-9440
- Masood, R., Cai, J., Zheng, T., Smith, D. L., Hinton, D. R. & Gill, P. S. (2001). Vascular endothelial growth factor (VEGF) is an autocrine growth factor for VEGF receptor-positive human tumors. *Blood*, Vol. 98, No. 6, (September 2001), pp. 1904-1913, ISSN 0006-4971
- Metheny-Barlow, L. J., Flynn, B., van Gijssel, H.E., Marrogi, A. & Gerwin, B. I. (2001). Paradoxical effects of platelet-derived growth factor-A overexpression in malignant mesothelioma. Antiproliferative effects in vitro and tumorigenic stimulation in vivo. *Am. J. Respir. Cell Mol. Biol.*, Vol. 24, No. 6, (June 2001), pp. 694-702, ISSN 1044-1549
- Micheau, O. & Tschopp, J. (2003). Induction of TNF receptor I-mediated apoptosis via two sequential signaling complexes. *Cell*, Vol. 114, No. 2, (July 2003), pp. 181-190, ISSN 0092-8674
- Milde-Langosch, K. (2005). The Fos family of transcription factors and their role in tumourigenesis. *Eur. J. Cancer.*, Vol. 41, No. 16, (November 2005), pp. 2449-2461, ISSN 0014-2964
- Miura, Y., Nishimura, Y., Katsuyama, H., Maeda, M., Hayashi, H., Dong, M., Hyodoh, F., Tomita, M., Matsuo, Y., Uesaka, A., Kuribayashi, K., Nakano, T., Kishimoto, T. & Otsuki T. (2006). Involvement of IL-10 and Bcl-2 in resistance against an asbestos-induced apoptosis of T cells. *Apoptosis*, Vol. 11, No. 10, (October 2006), pp. 1825-1835, ISSN 1360-8185
- Mossman, B. T. & Churg, A. (1998). Mechanisms in the pathogenesis of asbestosis and silicosis. *Am. J. Respir. Crit. Care Med.*, Vol. 157, No. 5, (May 1998), pp. 1666-1680, ISSN 1073-449X
- Müller, M. R. & Rao, A. (2010). NFAT, immunity and cancer: a transcription factor comes of age. *Nat. Rev. Immunol.*, Vol. 10, No. 9, (September 2010), pp. 645-656, ISSN 1474-1733
- Mutsaers, S.E. (2002). Mesothelial cells: Their structure, function and role in serosal repair. *Respirology*, Vol. 7, No. 3, (September 2002), pp. 171-191, ISSN 1323-7799
- Mutsaers, S. E. (2004). The mesothelial cell. *Int. J. Biochem. Cell Biol.*, Vol. 36, No. 1, (January 2004), pp. 9-16, ISSN 1357-2725
- Nymark, P., Lindholm, P. M., Korpela, M. V., Lahti, L., Ruosaari, S., Kaski, S., Hollmén, J., Anttila, S., Kinnula, V. L. & Knuutila, S. (2007). Gene expression profiles in asbestos-exposed epithelial and mesothelial lung cell lines. *BMC Genomics*, Vol. 8, (March 2007), p. 62, ISSN 1471-2164
- Ohm, J. E. & Carbone, D. P. (2001). VEGF as a mediator of tumor-associated immunodeficiency. *Immunol. Res.*, Vol. 23, No. 2-3, pp. 263-272, ISSN 0257-277X
- Ohta, Y., Shridhar, V., Bright, R. K., Kalemkerian, G. P., Du, W., Carbone, M., Watanabe, Y. & Pass, H. I. (1999). VEGF and VEGF type C play an important role in angiogenesis and

- lymphangiogenesis in human malignant mesothelioma tumours. *Br. J. Cancer.*, Vol. 81, No. 1, (September 1999), pp. 54–61, ISSN 0007-0920
- Olsson, A. K., Dimberg, A., Kreuger, J. & Claesson-Welsh, L. (2006). VEGF receptor signaling – in control of vascular function. *Nat. Rev. Mol. Cell. Biol.*, Vol. 7, No. 5, (May 2006), pp. 359-371, ISSN 1471-0072
- Ono, M. (2008). Molecular links between tumor angiogenesis and inflammation: inflammatory stimuli of macrophages and cancer cells as targets for therapeutic strategy. *Cancer Sci.*, Vol. 99, No. 8, (August 2008), pp. 1501–1506, ISSN 1347-9032
- Ostman, A. (2004). PDGF receptors-mediators of autocrine tumor growth and regulators of tumor vasculature and stroma. *Cytokine growth Factor Rev.*, Vol. 15, No. 4, (August 2004), pp. 275-286, ISSN 1359-6101
- Oura, H., Bertocini, J., Velasco, P., Brown, L. F., Carmeliet, P. & Detmar, M. (2003). A critical role of placenta growth factor in the induction of inflammation and edema formation. *Blood*, Vol. 101, No. 2, (January 2003), pp. 560-567, ISSN 0006-4971
- Oyama, T., Ran, S., Ishida, T., Nadaf, S., Kerr, L., Carbone, D. P. & Gabrilovich, D. I. (1998). Vascular endothelial growth factor affects dendritic cell maturation through the inhibition of nuclear factor-kappa B activation in hemopoietic progenitor cells. *J Immunol.*, Vol. 160, No. 3, (February 1998), pp. 1224-1232, ISSN 0022-1767
- Palumbo, C., Bei, R., Procopio, A. & Modesti A. (2008). Molecular targets and targeted therapies for malignant mesothelioma. *Curr. Med. Chem.*, Vol. 15, No. 9, pp. 855-867, ISSN 0929-8673
- Parr, C., Watkins, G., Boulton, M., Cai, J. & Jiang, W. G. (2005). Placenta growth factor is over-expressed and has prognostic value in human breast cancer. *Eur. J. Cancer*, Vol. 41, No. 18, (December 2005), pp. 2819-2827, ISSN 0014-2964
- Persico, M. G., Vincenti, V. & DiPalma, T. (1999). Structure, expression and receptor-binding properties of placenta growth factor (PlGF). *Curr. Top. Microbiol. Immunol.*, Vol. 237, pp. 31-40, ISSN 0070-217X
- Philip, M., Rowley, D. A. & Schreiber, H. (2004). Inflammation as a tumor promoter in cancer induction. *Semin. Cancer Biol.*, Vol. 14, No. 6, (December 2004), pp. 433-439, ISSN 1044-579X
- Pisick, E. & Salgia, R. (2005). Molecular biology of malignant mesothelioma: a review. *Hematol. Oncol. Clin. North Am.*, Vol. 19, No. 6, (December 2005), pp. 997-1023, ISSN 0889-8588
- Plate, K. H., Breier, G., Millauer, B., Ullrich, A. & Risau, W. (1993). Up-regulation of vascular endothelial growth factor and its cognate receptors in a rat glioma model of tumor angiogenesis. *Cancer Res.*, Vol. 53, No. 23, (December 1993), pp. 5822-5827, ISSN 0008-5472
- Pollard, J. W. (2004). Tumor-educated macrophages promote tumor progression and metastasis. *Nat. Rev. Cancer*, Vol. 4, No. 1, (January 2004), pp. 71–78, ISSN 1474, 175X
- Pompeo, E., Albonici, L., Doldo, E., Orlandi, A., Manzari, V., Modesti, A. & Mineo, T. C. (2009). Placenta growth factor expression has prognostic value in malignant pleural mesothelioma. *Ann. Thorac. Surg.*, Vol. 88, No. 2, pp. 426-431, ISSN 0003-4975

- Porta, C., Larghi, P., Rimoldi, M., Totaro, M. G., Allavena, P., Mantovani, A. & Sica, A. (2009). Cellular and molecular pathways linking inflammation and cancer. *Immunobiology*, Vol. 214, No. 9-10, pp. 761-777, ISSN 0171-2985
- Rafii, S., Avezilla, S., Shmelkov, S., Shido, K., Tejada, R., Moore, M. A., Heissig, B. & Hattori, K. (2003). Angiogenic factors reconstitute hematopoiesis by recruiting stem cells from bone marrow microenvironment. *Ann. N. Y. Acad. Sci.*, Vol. 996, (May 2003), pp. 49-60, ISSN 0077-8923
- Ramos-Ninos, M., Timblin, C. & Mossman, B. T. (2002). Mesothelial cell transformation requires increased AP-1 binding activity and ERK-dependent Fra-1 expression. *Cancer Res.*, Vol. 62, No. 21, (November 2002), pp. 6065-6069, ISSN 0008-5472
- Reuter, S., Gupta, S. C., Chaturvedi, M. M. & Aggarwal, B. B. Oxidative stress, inflammation, and cancer: how are they linked? *Free Radic. Biol. Med.*, Vol. 49, No. 11, (December 2010), pp. 1603-1616, ISSN 0891-5849
- Ribatti, D., Nico, B., Crivellato, E., Roccaro, A. M. & Vacca, A. (2007). The history of the angiogenic switch concept. *Leukemia*, Vol. 21, No. 1, (January 2007), pp. 44-52, ISSN 0887-6924
- Riganti, C., Orecchia, S., Silvagno, F., Pescarmona, G., Betta, P. G., Gazzano, E., Aldieri, E., Ghigo, D. & Bosia A. (2007). Asbestos induces nitric oxide synthesis in mesothelioma cells via Rho signaling inhibition. *Am. J. Respir. Cell. Mol. Biol.*, Vol. 36, No. 6, (June 2007), pp. 746-756, ISSN 1044-1549
- Robinson, B. W. & Lake, R. A. (2005). Advances in malignant mesothelioma. *N. Engl. J. Med.*, Vol. 353, No. 15, (October 2005), pp. 1591-1603, ISSN 0028-4793
- Robledo, R. & Mossman, B. (1999). Cellular and molecular mechanisms of asbestos-induced fibrosis. *J. Cell. Physiol.*, Vol. 180, No. 2, (August 1999), pp. 158-166, ISSN 0021-9541
- Roos, W. P. & Kaina, B. (2006). DNA damage-induced cell death by apoptosis. *Trends Mol. Med.*, Vol. 12, No. 9, (September 2006), pp. 440-450, ISSN 1471-4914
- Rose-John, S., Waetzig, G. H., Scheller, J., Grötzinger, J. & Seeger, D. (2007). The IL-6/sIL-6R complex as a novel target for therapeutic approaches. *Expert Opin. Ther. Targets*, Vol. 11, No. 5, (May 2007), pp. 613-624, ISSN 1472-8222
- Roskoski, R. Jr. (2007). Vascular endothelial growth factor (VEGF) signaling in tumor progression. *Crit. Rev. Oncol. Hematol.*, Vol. 62, No. 3, (June 2007), pp. 179-213, ISSN 1040-8428
- Royds, J. A., Dower, S. K., Qwarnstrom, E. E. & Lewis, C. E. (1998). Response of tumour cells to hypoxia: role of p53 and NFkB. *Mol. Pathol.*, Vol. 51, No. 2, (April 1998), pp. 55-61, ISSN 1366-8714
- Saccani, S., Pantano, S. & Natoli, G. (2003) Modulation of NF-kappaB activity by Exchange of dimers. *Mol. Cell*, Vol. 11, No. 6, (June 2003), pp. 1563-1574, ISSN 1097-2765
- Saylor, P. J., Escudier, B. & Michaelson, M. D. (2012). Importance of Fibroblast Growth Factor Receptor in Neovascularization and Tumor Escape from Antiangiogenic Therapy. *Clin. Genitourin. Cancer*, (February 2012), Epub ahead of print, ISSN 1558-7673
- Scheidereit, C. (2006). IkappaB kinase complexes: gateways to NF-kappaB activation and transcription. *Oncogene*, Vol. 25, No. 51, (2006), (October 2006), pp. 6685-6705, ISSN 0950-9232

- Shaulian, E. & Karin, M. (2001). AP-1 in cell proliferation and survival. *Oncogene*, Vol. 20, No. 19, (April 2001), pp. 2390-2400, ISSN 0950-9232
- Shaulian, E. & Karin, M. (2002). AP-1 as a regulator of cell life and death. *Nat. Cell Biol.*, Vol. 4, No. 5, (May 2002), pp. E131-E136, ISSN 1465-7392
- Shaw, J. P., Utz, P. J., Durand, D. B., Toole, J. J., Emmel, E. A. & Crabtree, G. R. (1988). Identification of a putative regulator of early T cell activation genes. *Science*, Vol. 241, No. 4862, (July 1988), pp. 202-205, ISSN 0036-8075
- Shibuya, M. (2006). Differential roles of vascular endothelial growth factor receptor-1 and receptor-2 in angiogenesis. *J. Biochem. Mol. Biol.*, Vol. 39, No. 5, (September 2006), pp. 469-478, ISSN 1225-8687
- Schonthaler, H. B., Guinea-Viniegra, J. & Wagner, E. F. (2011). Targeting inflammation by modulating the Jun/AP-1 pathway. *Ann. Rheum. Dis.*, Vol. 70, Suppl. 1, (March 2011), pp. i109-i112, ISSN 0003-4967
- Shukla, A., Gulumian, M., Hei, T. K., Kamp, D., Rahman, Q. & Mossman, B. T. (2003a). Multiple roles of oxidants in the pathogenesis of asbestos-induced diseases. *Free Radical Biol. Med.*, Vol. 34, No. 9, (May 2003), pp. 1117-1129, ISSN 0891-5849
- Shukla, A., Jung, M., Stern, M., Fukagawa, N. K., Taatjes, D. J., Sawyer, D., Van Houten, B. & Mossman, B. T. (2003b). Asbestos induces mitochondrial DNA damage and dysfunction linked to the development of apoptosis. *Am. J. Physiol. Lung Cell. Mol. Physiol.*, Vol. 285, No. 5, (November 2003), pp. L1018-L1025, ISSN 1040-0605
- Sica, A. (2010). Role of tumour-associated macrophages in cancer-related inflammation. *Exp. Oncol.*, Vol. 32, No. 3, (September 2010), pp. 153-158, ISSN 1812-9269
- Simeonova, P. & Luster, M. (1995). Iron and reactive oxygen species in the asbestos-induced tumor necrosis factor- α response from alveolar macrophages. *Am. J. Respir. Cell. Mol. Biol.*, Vol. 12, No. 6, (June 1995), pp. 676-683, ISSN 1044-1549
- Simeonova, P. P., Toriumi, W., Kommineni, C., Erkan, M., Munson, A. E., Rom, W. N. & Luster, M. I. (1997). Molecular regulation of IL-6 activation by asbestos in lung epithelial cells: role of reactive oxygen species. *J. Immunol.*, Vol. 159, No. 8, (October 1997), pp. 3921-3928, ISSN 0022-1767
- Stapelberg, M., Gellert, N., Swettenham, E., Tomasetti, M., Witting, P. K., Procopio, A. & Neuzil, J. (2005). Alpha-tocopheryl succinate inhibits malignant mesothelioma by disrupting the fibroblast growth factor autocrine loop: mechanism and the role of oxidative stress. *J. Biol. Chem.*, Vol. 280, No. 27, (July 2005), pp. 25369-25376, ISSN 0021-9258
- Strizzi, L., Catalano, A., Vianale, G., Orecchia, S., Casalini, A., Tassi, G., Puntoni, R., Mutti, L. & Procopio, A. (2001a). Vascular endothelial growth factor is an autocrine growth factor in human malignant mesothelioma. *J. Pathol.*, Vol. 193, No. 4, (April 2001), pp. 468-475, ISSN 0022-3417
- Strizzi, L., Vianale, G., Catalano, A., Muraro, R., Mutti, L. & Procopio, A. (2001b). Basic fibroblast growth factor in mesothelioma pleural effusions: correlation with patient survival and angiogenesis. *Int. J. Oncol.*, Vol. 18, No. 5, (May 2001), pp. 1093-1098, ISSN 1019-6439

- Sun, S. C. (2011). Non-canonical NF- κ B signaling pathway. *Cell Res.*, Vol. 21, No. 1, (January 2011), pp. 71-85, ISSN 1001-0602
- Takahashi, A., Kono, K., Ichihara, F., Sugai, H., Fuji, H. & Matsumoto, Y. (2004). Vascular endothelial growth factor inhibits maturation of dendritic cells induced by lipopolysaccharide, but not by proinflammatory cytokines. *Cancer Immunol., Immunother.*, Vol. 53, No. 6, (June 2004), pp. 543-550, ISSN 0340-7004
- Tanaka, S., Choe, N., Hemenway, D. R., Zhu, S., Matalon, S. & Kagan, E. (1998). Asbestos inhalation induces reactive nitrogen species and nitrotyrosine formation in the lungs and pleura of the rat. *J. Clin. Invest.*, Vol. 102, No. 2, (July 1998), pp. 445-454, ISSN 0021-9738
- Toi, M., Matsumoto, T. & Bando, H. (2001). Vascular endothelial growth factor: its prognostic, predictive, and therapeutic implications. *Lancet Oncol.*, Vol. 2, No. 11, (November 2001), pp. 667-673, ISSN 1470-2045
- Toyokuni, S. (2009). Mechanisms of asbestos-induced carcinogenesis. *Nagoya J. Med. Sci.*, Vol. 71, No. 1-2, (February 2009), pp. 1-10, ISSN 0027-7622
- Toyooka, S., Kishimoto, T. & Date, H. (2008). Advances in the molecular biology of malignant mesothelioma. *Acta Med., Okayama*, Vol. 62, No. 1, (February 2008), pp. 1-7, ISSN 0386-300X
- Ullrich, E., Bonmort, M., Mignot, G., Kroemer, G. & Zitvogel, L. (2008). Tumor stress, cell death and the ensuing immune response. *Cell Death Differ.*, Vol. 15, No. 1, (January 2008), pp. 21-28, ISSN 1350-9047
- Valle, M. T., Castagneto, B., Procopio, A., Carbone, M., Giordano, A. & Mutti, L. (1998). Immunobiology and immune defense mechanisms of mesothelioma cells. *Monaldi Arch. Chest Dis.*, Vol. 53, No. 2, (April 1998), pp. 219-227, ISSN 1122-0643
- Villanova, F., Procopio, A. & Rippo, M. R. (2008). Malignant mesothelioma resistance to apoptosis: recent discoveries and their implication for effective therapeutic strategies. *Curr. Med. Chem.*, Vol. 15, No. 7, pp. 631-641, ISSN 0929-8673
- Waltenberger, J., Claesson-Welsh, L., Siegbahn, A., Shibuya, M. & Heldin, C. H. (1994). Different signal transduction properties of KDR and Flt1, two receptors for vascular endothelial growth factor. *J. Biol. Chem.*, Vol. 269, No. 43, (October 1994), pp. 26988-26995, ISSN 0021-9258
- Wei, S. C., Tsao, P. N., Yu, S. C., Shun, C. T., Tsai-Wu, J. J., Wu, C. H., Su, Y. N., Hsieh, F. J. & Wong, J. M. (2005). Placenta growth factor expression is correlated with survival of patients with colorectal cancer. *Gut*, Vol. 54, No. 5, (May 2005), pp. 666-672, ISSN 0017-5749
- Wu, W. S. (2006). The signaling mechanism of ROS in tumor progression. *Cancer Metastasis Rev.*, Vol. 25, No. 4, (December 2006), pp. 695-705, ISSN 0167-7659
- Yang, H., Bocchetta, M., Kroczyńska, B., Elmishad, A. G., Chen, Y., Liu, Z., Bubici, C., Mossman, B. T., Pass, H. I., Testa, J. R., Franzoso, G. & Carbone, M. (2006). TNF- α inhibits asbestos-induced cytotoxicity via a NF- κ B-dependent pathway, a possible mechanism for asbestos-induced oncogenesis. *Proc. Natl. Acad. Sci. USA.*, Vol. 103, No. 27, (July 2006), pp. 10397-10402, ISSN 0027-8424

- Yarborough, C.M. (2007). The risk of mesothelioma from exposure to chrysotile asbestos. *Curr. Opin. Pulm. Med.*, Vol. 13, No. 4, (July 2007), pp. 334-338, ISSN 1070-5287
- Yasumitsu, A., Tabata, C., Tabata, R., Hirayama, N., Murakami, A., Yamada, S., Terada, T., Iida, S., Tamura, K., Fukuoka, K., Kuribayashi, K. & Nakano, T. (2010). Clinical significance of serum vascular endothelial growth factor in malignant pleural mesothelioma. *J. Thorac. Oncol.*, Vol. 5, No. 4, (April 2010), pp. 479-483, ISSN 1556-0864
- Yoo, S. A., Kwok, S. K. & Kim W. U. (2008). Proinflammatory role of vascular endothelial growth factor in the pathogenesis of rheumatoid arthritis: prospect for therapeutic intervention. *Mediators Inflamm.*, Vol. 2008, Article ID 129873, ISSN 0962-9351
- Yoshida, K. & Miki, Y. (2010). The cell death machinery governed by the p53 tumor suppressor in response to DNA damage. *Cancer Sci.*, Vol. 101, No. 4, (April 2010), pp. 831-835, ISSN 1347-9032
- Zebrowski, B. K., Yano, S., Liu, W., Shaheen, R. M., Hicklin, D. J., Putnam, J. B. & Ellis, L. M. (1999). Vascular endothelial growth factor levels and induction of permeability in malignant pleural effusions. *Clin. Cancer Res.*, Vol. 5, No. 11, (November 1999), pp. 3364-3368, ISSN 1078-0432
- Ziogas, A. C., Gavalas, N. G., Tsiatas, M., Tsitsilonis, O., Politi, E., Terpos, E., Rodolakis, A., Vlahos, G., Thomakos, N., Haidopoulos, D., Antsaklis, A., Dimopoulos, M. A. & Bamias, A. (2012). VEGF directly suppresses activation of T cells from ovarian cancer patients and healthy individuals via VEGF receptor Type 2. *Int. J. Cancer.*, Vol. 130, No. 4, (February 2012), pp. 857-864, ISSN 0020-7136