We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

The Cardioprotection of Silymarin in Coronary Artery Bypass Grafting Surgery

D. Tagreed Altaei, D. Imad A. Jamal and
D. Diyar Dilshad

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/55125>

1. Introduction

Coronary artery bypass grafting surgery (CABG) is one of the most effective therapies for coronary artery disease (CAD). CABG is conventionally performed with the use of cardiopulmonary bypass (CPB), which has been associated with an increased frequency of complications [1-3]. A variety of risk factors has been described to help delineate risk assessment of patients undergoing CABG, including preoperative left ventricular ejection fraction (LVEF) [4, 5] and postoperative increase in creatine kinase myocardial band (CK-MB) levels [6-11]. In patients undergoing coronary artery bypass grafting (CABG), the use of cardiopulmonary bypass (CPB) in combination with aortic clamping during coronary artery bypass grafting (CABG) elicits ischemic myocardial injury [12]. The vascular endothelium is a complex synthetic organ subject to injury from numerous potential insults, including oxidative stress [13, 14] modified lipoproteins [15], and hemodynamic forces [16]. Injured endothelial cells initiate a largely stereotyped, initially protective response. The concurrent uptake of low-density lipoproteins (LDLs) by monocyte-derived macrophages transforms them into the lipid laden foam cells that constitute a key element of the fatty streak, the first recognizable progenitor of the advanced atherosclerotic lesion [17, 18].

Silymarin, a flavonolignan from 'milk thistle' (*Silybum marianum*) plant, is used from ancient times as a hepatoprotective, antioxidant, anti-lipid peroxidative, antifibrotic, anti-inflammatory, immunomodulatory and liver regenerating. Silymarin has cardioprotective activity against ischemia-reperfusion induced myocardial infarction in rats [19].

Protective efficacy of Silymarin treatment confirmed by anti-inflammatory and antioxidant actions against reperfusion injury and inflammation during CABG surgery [20].

This may represent a novel cardioprotective agent to be used pre CABG, to our knowledge; this is the first study of pretreatment Silymarin as cardioprotective agent in patients undergoing CABG.

2. Patients and methods

The local ethics committee approved the investigation, and informed written consent was obtained from all patients entering the study. 140 patients admitted to the hospital for the first time for elective coronary artery bypass surgery were invited to take part. They were randomized into three groups (G); G I: Administered Silymarin (Legalon® tablet), 140 mg×3; 1 day before surgery. G II: Administered Silymarin (Legalon® tablet), 140 mg×3; 3 days before surgery. G III: Control (no treatment). Patients receiving corticosteroids were deemed not eligible. Any drugs were withheld on the morning of surgery.

Surgical procedure: Specifications on the extracorporeal circulation circuit, cardiopulmonary bypass procedures and surgical procedures have been described previously [20].

At baseline, demographic data (age, sex, weight, BSA, BMI), and history of conventional vascular risk factors (hypertension, diabetes mellitus, hyperlipidemia, smoking habit, alcohol abuse) were obtained.

Routine laboratory investigations were performed the first day after admission to the hospital after overnight fasting and later before discharged. It included levels of WBCs; differential counts (Neutrophils, Monocytes, Lymphocytes), RBCs, ESR, total cholesterol, LDL, vLDL, HDL, triglycerides, SGOT, SGPT, B.urea, S.creatinin, alkaline phosphatase, serum bilirubin, blood sugar, HbA1c, & ESR. Blood samples for troponin I (T), & creatine kinase; (CK-MB) measurement were obtained within 24 hours before surgery, & 2 hr, 24 hr post CABG. All laboratory tests were performed according to companys' procedures. Laboratory staffs were blind to the tested drug and groups.

Statistical analysis

The present Data was analyzed using Student 't' test, one-way analysis of covariance. $P < 0.05$ was considered to be significant. All data were analyzed using the statistical package SPSS (version 10.0) Continuous data are presented as mean $\pm$ SD and categorical data as absolute numbers, or mean.

3. Results

One hundred and forty patients [105 (75%) males, and 35 (25%) females] with a mean age of 64.5 years were included in the study. All underwent on-pump CABG.

No significant differences were noted between the groups in age, body surface area, BMI, and operation data. The demographic data on the 140 patients completing the study are presented

in Table 1. Clinical & operative characteristics (Age, sex, body weight, smoking, left ventricular ejection fraction, usual administered drugs, other diseases, family history of coronary artery disease) were largely independent of silymarin treatment with no significance ($p>0.05$).

Variables	Silymarin (SM) Group (No. = 90)		Control Group (No. = 50)
	G I (n=40)	G II (n=50)	G III
Male/Female (No.)	68/22		37/13
Age (years) ^a	65		64
Body surface area (m ²) ^b	1.6 ± 0.2		1.7 ± 0.1
BMI (kg/m ²) ^b	42.3 ± 0.2		44.7 ± 0.2
Ejection fraction (%) ^b	58 ± 1		60 ± 1
Operative time (min) ^b	201 ± 16		200 ± 18
No. of grafts ^a	3		3.5
Hospitalization (days) ^a	4		7

Table 1. Clinical and Patients’ data in Silymarin treated and control groups.

There was a significant decrease in post operative values of; WBCs counts, Neutrophil, monocytes, lymphocytes, RBCs, ESR (Figure1), total cholesterol, LDL, vLDL, & triglycerides (Figure2), SGOT, SGPT, alkaline phosphatase (47%), showed in (Figure 3), B.urea, S.creatinine, serum bilirubin (48%) (Figure4), blood sugar, HbA1c in diabetic patients (43%) as showed in figure 5, in SM treated group compared to baseline and control group; ($P=0.002$), while there was an elevation of postoperative HDL values in patients treated with SM compared to control, ($p=0.001$).

Figure 1. The mean values of Neutrophils, Monocytes, Lymphocytes, total WBCs, RBCs, and ESR for Silymarin treated and control groups pre and post CABG.

Figure 2. The mean values (mg/dL) of total cholesterol, LDL, vLDL, HDL, and triglycerides for Silymarin treated and control groups pre and post CABG.

Figure 3. Mean values of SGOT, SGPT, & Alkaline phosphatase for both groups compared pre and post CABG.

Figure 4. The mean values of blood urea & serum creatinin, & serum bilirubin in tested groups pre & post operation.

Figure 5. Mean values of fasting blood sugar, & glycosylated heamoglobin pre & post CABG for both groups.

Surgery was associated with a significant increase in troponin I (T1), CK-MB (CK1) in both groups, but after 24 hr post CABG there was a significant reduction of Troponin I (T2) values in SM treated group compared to baseline (T0), after 2 hour (T1), and control group, [p=0.001]. SM-treated patients released significantly less creatine kinase (CK)-MB than the control subjects postoperatively (CK1) after 2 hours, then back to normal levels after 24 hours (CK 2) [p = 0.004]; indicating less myocardial injury in patients receiving SM when compared to the control subjects (no significant change $p > 0.05$), as showed in figure 6.

Figure 6. Mean Troponin I, & CK-MB values at different times in Silymarin (SM) treated patients and control groups. [T0=pre CABG, T1=2 hours after CABG, T2= 24 hours after CABG, CK-MB 0=pre CABG, CK-MB 1= 2 hours after CABG, CK-MB 2= 24 hours after CABG].

Patients treated with Silymarin before 3 days better than those treated before 1 day, but there was no statistical significant differences between them (G I & G II), $p>0.05$.

4. Discussion

Myocardial ischemia and reperfusion is a common occurrence in CABG patients. Reintroduction of oxygen to previously ischemic myocardium can result in irreversible tissue injury, and Ischemic myocardial damage is associated with inflammation [21, 22].

SM has been shown to have a potential positive effect on immune function by its ability to enhance neutrophil activity [23]. Silymarin & Silibinin by interacting with the lipid component of cell membranes can influence their chemical & physical properties. Studies in erythrocytes, mast cells, leucocytes, macrophages & hepatocytes have shown that SM renders cell membranes more resistant to lesions [24].

Studies have shown that silymarin exerts a number of effects, including inhibition of neutrophil migration [25, 24]. The inhibitory effects of silymarin on neutrophil function prevent post _ ischemic mucosal injury [26]. Activated neutrophils are thought to play a major role in ischemia-reperfusion injury [27]. This study agrees with that, SM treated groups showed a significant reduction of WBCs, neutrophils count post CABG, & ESR also. According to this, silymarin may prevent reperfusion injury so it may have a beneficial effect during CABG.

Milk thistle was able to inhibit the biosynthesis of cholesterol in the liver and reduce LDL cholesterol oxidation, one of the primary mechanisms of atherosclerosis [28,25]. This study agrees with it, SM treatment showed a significant reduction of total cholesterol, LDL, & vLDL, while HDL was elevated significantly.

SM interact directly with the cell membrane components to prevent any abnormalities in the content of lipid fraction responsible for maintaining normal fluidity [29].

Silymarin appears to act as an antioxidant not only because it acts as a scavenger of the free radicals that induce lipid peroxidation, but also because it influences enzyme systems associated with glutathione & superoxide dismutase [30]. Also, prevent damage to rat heart membrane primarily through a free radical scavenging mechanism [31]. This study agrees with previous mentioned studies, the antioxidant activity of Silymarin prevents vascular endothelium injury during CABG.

In the present study, all patients showed significantly higher plasma levels of markers of peri-operative myocardial tissue injury early after the start of reperfusion. Silymarin treated group showed a significant reduction of all measured parameters compared to baseline and control.

Alkaline phosphatase, & serum bilirubin was elevated preoperatively in 47%, and 48% of enrolled patients, respectively, pretreatment with Silymarin showed a significant reduction post operatively because SM have the ability to prevent injury from different causes.

Blood sugar & HbA1c in diabetic patients (43%) were diminished significantly in those treated by SM.

SM counteracted the increase in the cardiac enzymes and cTnI concentration induced by cisplatin, toward near normal levels. Rao and Viswanath 2007 [19] reported that the administration of SM before ischemia-reperfusion-induced myocardial infarction maintained the levels of marker enzymes (LDH, CK and CK-MB) compared to isoproterenol-injected rats. A possible explanation is that silymarin, via its anti-lipid peroxidation activity, causes stabilization of cardiac membranes and prevents the leakage of cardiac enzymes [32]. Silibinin induced cardiac myocyte expression of Bcl-2 protein, which prevented permeability transition pore opening, and, therefore, cytochrome c release decreased. These events might be one of the mechanisms of silymarin-mediated stabilization of the mitochondrial membrane [33]. This study agrees with the above study in which Troponin I, and CK-MB diminished significantly in SM treated group compared to baseline and control.

The anti-inflammatory activity (significant reduction of cytokines post operatively), & antioxidant effects of SM during CABG [20], in addition to that, the highly significant reduction in serum levels of troponin I, & CK-MB, confirm the cardioprotection activity of SM during CABG. This mechanism of action may reduce ischemia-reperfusion injury, and protect the myocardium.

The results of this study indicate that SM pretreatment induces potent endogenous protection against subsequent ischemic stress in the human myocardium, and reduced the damage caused by reperfusion injury. SM has the ability to regulate membrane permeability and to increase membrane stability in the presence of xenobiotic damage, capacity to regulate nuclear expression by means of a steroid-like effect.

There was no statistically significant difference between the two treated groups, even the results of patients treated with SM before 3 days better than those treated before 1 day of CABG surgery. The data of this study needs further large-scale, randomized, double blind technique studies for other cardiovascular diseases to confirm the present study before clinical use of Silymarin.

5. Conclusion

The authors conclude that CABG surgery need using cardioprotective agent pre operation and suggest using other pharmaceutical preparation I.V. dosage form of SM and studying the pharmacokinetics during CABG surgery. The pre-operative administration of Silymarin may reduce perioperative morbidity and myocardial injury during CABG surgery. The authors suggest a large-scale, randomized, double blind study with cardiovascular events before Silymarin could be considered for clinical use.

Acknowledgements

The authors wish to thanks all patients who so kindly took part, and all staff.

Author details

D. Tagreed Altaei*, D. Imad A. Jamal and D. Diyar Dilshad

*Address all correspondence to: tagreedaltaei@yahoo.com

Dep. of Pharmacology and Toxicology, College of Pharmacy, Hawler Medical University, Erbil, Iraq

References

- [1] Czerny, M, Baumer, H, Kilo, J, et al. Inflammatory response and myocardial injury following coronary artery bypass grafting with or without cardiopulmonary bypass. *Eur J Cardiothorac Surg.* (2000). , 17, 737-42.
- [2] Wan, I. Y, Arifi, A. A, Wan, S, et al. Beating heart revascularization with or without cardiopulmonary bypass: evaluation of inflammatory response in a prospective randomized trial. *J Thorac Cardiovasc Surg.* (2004). , 127, 1624-31.
- [3] Schulze, C, Conrad, N, Schutz, A, et al. Reduced expression of systemic proinflammatory cytokines after off-pump versus conventional coronary artery bypass grafting. *Thorac Cardiovasc Surg.* (2000). , 48, 364-9.
- [4] Magovern, J. A, Sakert, T, Magovern, G. J, et al. A model that predicts morbidity and mortality after coronary artery bypass graft surgery. *J Am Coll Cardiol.* (1996). , 28, 1147-53.

- [5] Provenchere, S, Berroeta, C, Reynaud, C, et al. Plasma brain natriuretic peptide and cardiac troponin I concentrations after adult cardiac surgery: association with postoperative cardiac dysfunction and 1-year mortality. *Crit Care Med.* (2006). , 34, 995-1000.
- [6] Costa, M. A, Carere, R. G, Lichtenstein, S. V, et al. Incidence, predictors, and significance of abnormal cardiac enzyme rise in patients treated with bypass surgery in the arterial revascularization study (ARTS). *Circulation.* (2001). , 104, 2689-93.
- [7] Klatte, K, Chaitman, B. R, Theroux, P, et al. Guardian Investigators (The GUARD during Ischemia Against Necrosis). Increased mortality after coronary artery bypass graft surgery is associated with increased levels of postoperative creatine kinase-myocardial band isoenzyme release: results from the guardian trial. *J Am Coll Cardiol.* (2001). , 38, 1070-7.
- [8] Steuer, J, Horte, L. G, Lindahl, B, & Stahle, E. Impact of perioperative myocardial injury on early and long-term outcome after coronary artery bypass grafting. *Eur Heart J.* (2002). , 23, 1219-27.
- [9] Brener, S. J, Lytle, B. W, Schneider, J. P, et al. Association between CKMB elevation after percutaneous or surgical revascularization and three-year mortality. *J Am Coll Cardiol.* (2002). , 40, 1961-7.
- [10] Engoren, M. C, Habib, R. H, Zacharias, A, et al. The association of elevated creatine kinase-myocardial band on mortality after coronary artery bypass grafting surgery is time and magnitude limited. *Eur J Cardiothorac Surg.* (2005). , 28, 114-9.
- [11] Ramsay, J, Shernan, S, Fitch, J, et al. Increased creatine kinase MK level predicts postoperative mortality after cardiac surgery independent of new Q waves. *J Thorac Cardiovasc Surg.* (2005). , 129, 300-6.
- [12] Fransen, E. J, Maessen, J. G, & Hermens, W. T. Glatz JFC. Demonstration of ischemia-reperfusion injury separate from postoperative infarction in Coronary Artery Bypass Graft Patients. *Ann Thorac Surg.* (1998). , 65, 48-53.
- [13] Gong, K. W, Zhu, G. Y, & Wang, L. H. Effect of active oxygen species on intimal proliferation in rat aorta after arterial injury. *J Vasc Res.* (1996). , 42-46.
- [14] Liao, F, Andalibi, A, & Qiao, J. H. Genetic evidence for a common pathway mediating oxidative stress, inflammatory gene induction, and aortic fatty streak formation in mice. *J Clin Invest.* (1994). , 94, 877-84.
- [15] Steinberg, D. Antioxidants and atherosclerosis: A current assessment. *Circulation.* (1991). , 84, 1420-25.
- [16] Glagov, S, Zarins, C, & Giddens, D. P. Hemodynamics and atherosclerosis: Insights and perspectives gained from studies of human arteries. *Arch Pathol lab Med.* (1988). , 112, 1018-31.

- [17] Aqel, N. M, Ball, R. Y, & Waldmann, H. Monocytic origin of foam cells in human atherosclerotic plaques. *Atherosclerosis*. (1984). , 53, 265-71.
- [18] Stary, H. C. Evolution and progression of atherosclerotic lesions in coronary arteries of children and young adults, *Arteriosclerosis*. (1989). , 9, 119-32.
- [19] Rao, P. R, & Viswanath, R. K. Cardioprotective activity of silymarin in ischemia-reperfusion-induced myocardial infarction in albino rats. *Exp Clin Cardiol*. (2007). , 12(4), 179-187.
- [20] Altaei, Tagreed. Protective effect of Silymarin during coronary artery bypass grafting surgery. *Exp Clin Cardiol*. (2011). , 17(1), 34-38.
- [21] Hawkins, H. K, Entman, M. L, Zhu, J. Y, et al. Acute inflammatory reaction after myocardial ischemic injury and reperfusion. Development and use of a neutrophil-specific antibody. *Am J Pathol*.(1996). , 148, 1957-1969.
- [22] Entman, M. L, Michael, L, Rossen, R. D, et al. Inflammation in the course of early myocardial ischemia. *FASEB J*. (1991). , 5, 2529-2537.
- [23] Kalmar, L. Silibinin (Legalon-70) enhances the motility of human neutrophils immobilizes by calcium ionophore, lymphokine, and normal human serum. *Agents Actions*. (1990).
- [24] De La Puerta, R, Martinez, E, Bravo, L, et al. Effect of silymarin on different acute inflammation models and leukocyte migration. *J Pharm Pharmacol*. (1996).
- [25] Skottova, N, & Krecman, V. Silymarin as a potential hypocholesterolaemic drug. *Physiol Res*. (1998). , 47, 1-7.
- [26] Alarcon de la Lastra ACMartin MJ, Motilva V, Jimenez M, La Casa C, Lopez A. Gas-troprotection induced by silymarin, the hepatoprotective principle of *Silybum marianum* in ischemia-reperfusion mucosal injury: role of neutrophils. *Planta Med*. (1995).
- [27] Weiss, S. J. Tissue destruction by neutrophils. *N Engl J Med*. (1989). , 320, 365-376.
- [28] Bialecka, M. The effect of bioflavonoids and lecithin on the course of experimental atherosclerosis. *Acad Med Stetin*. (1997).
- [29] Muriel, P, & Mourelle, M. Prevention by silymarin of membrane alterations in acute CCl₄ liver damage. *J Appl Toxicol*. (1990). , 10, 275-9.
- [30] Letteron, P, Labbe, G, Degott, C, et al. Mechanism for the protective effects of silymarin against carbon tetrachloride-induced lipid peroxidation and hepatotoxicity in mice. *Biochem Pharmacol*. (1990).
- [31] Psotova, J, Chlopckikova, S, Grambal, F, & Simanek, V. Ulrichova J; Influence of silymarin and its flavolignans on doxorubicin-iron induced lipid peroxidation in rat heart microsomes and mitochondria in comparison with quercetin: *Phytother Res*. (2002). Suppl 1: S63_7.

- [32] Abu Ghadeer A.R., Ali, S.E., Osman, S.A., Abu Bedair, F.A., Abbady, M.M., El-Kady, M.R. Antagonistic role of silymarin against cardiotoxicity and impaired antioxidation induced by adriamycin and/or radiation exposure in albino rats. *Pakistan J. Biol. Sci.* (2001). , 4, 604-607.
- [33] Zhou, B, Wu, L, Tashiro, S, Onodera, S, Uchiumi, F, & Ikejima, T. Silibinin protect rat cardiac myocyte from isoproterenol-induced DNA damage independent on regulation of cell cycle. *Biol. Pharm. Bull.* (2006 a). , 29, 1900-1905.