We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Metabolism of Plasma Membrane Lipids in Mycobacteria and Corynebacteria

Paul K. Crellin, Chu-Yuan Luo and Yasu S. Morita

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/52781>

1. Introduction

Bacteria of the Corynebacterineae, a suborder of the Actinobacteria, comprise *Mycobacterium*, *Corynebacterium*, *Nocardia*, *Rhodococcus* and other genera. This suborder of high GC gram-positive bacteria includes a number of important human pathogens, such as *Mycobacterium tuberculosis*, *Mycobacterium leprae* and *Corynebacterium diphtheriae*, the causative agents of tuberculosis, leprosy and diphtheria, respectively. *M. tuberculosis* is the most medically significant species, a devastating human pathogen infecting around one-third of the entire human population and responsible for more than 1 million deaths annually. The Corynebacterineae also includes non-pathogenic species such as *Mycobacterium smegmatis*, a saprophytic species, and *Corynebacterium glutamicum*, an industrial workhorse for the production of amino acids and other useful compounds. These relatively fast-growing species serve as useful models to study metabolic processes essential to the growth and survival of the slow-growing pathogens.

All these bacteria share a common feature, a distinctive multilaminate cell wall composed of peptidoglycan, complex polysaccharides, and both covalently linked lipids and free lipids/lipoglycans (Fig. 1). Among them, mycolic acids are the hallmark of these species. These long chain α -branched, β -hydroxylated fatty acids are covalently linked to the arabinogalactan polysaccharide layer. This mycolic acid layer is complemented by a glycolipid layer to form an outer “mycomembrane” analogous to the outer membrane of Gram-negative bacteria. [1, 2]. The outer leaflet of the mycomembrane is composed of a variety of lipids including trehalose dimycolates (TDMs), glycopeptidolipids (GPLs), phthiocerol dimycocerosates (PDIMs), sulfolipids, phenolic glycolipids (PGLs), and lipooligosaccharides. Some of these lipids are widely distributed while others are restricted to particular species. For example, TDMs and their structural equivalents are found in both mycobacteria and corynebacteria, while PDIMs and PGLs are restricted to a subset of

mycobacteria. The structure and hydrophobic properties of the cell wall make it a potent permeability barrier that is responsible for intrinsic resistance of mycobacteria to an array of host microbiocidal processes, many antibiotics and sterilization conditions [3, 4]. Many of the cell wall components of pathogenic mycobacterial species are essential for pathogenesis and *in vitro* growth, hampering efforts to characterize the function of individual proteins in their assembly. In contrast, some non-pathogenic species such as *C. glutamicum* can tolerate the loss of major cell wall components, making them useful model systems for delineating processes involved in the assembly of core cell wall structures.

Figure 1. Mycobacterial plasma membrane and cell wall with flow of key metabolic pathways. Some of the metabolites are exported to the mycomembrane. SLD, small lipid droplet; LD, lipid droplet; FA-CoA, fatty acyl-CoA. See text for other abbreviations used in the figure.

Studies on mycobacteria and corynebacteria provide a unique opportunity to illustrate the complexity and diversity of lipid metabolic pathways in bacteria. They have a significantly higher lipid content than other bacteria with cell wall lipids comprising ~40% of the dry cell mass. *M. tuberculosis* produces a diversity of lipids unparalleled in bacteria, from simple fatty acids to highly complex long chain structures such as mycolic acids. It has devoted a significant proportion of its coding capacity to lipid metabolism and produces about 250 enzymes dedicated to fatty acid metabolism, which is around five times the number produced by *Escherichia coli* [5]. Lipid biosynthesis places a significant metabolic burden on the organism but is ultimately advantageous, allowing *M. tuberculosis* to survive and replicate in the inhospitable environment of host macrophages. While capable of *de novo* synthesis, these bacteria also scavenge and degrade host cell membrane lipids to acetyl-CoA, via broad families of β -oxidation and other catabolic enzymes, for incorporation into their own metabolic pathways and to fuel cellular processes.

The plasma membrane provides the platform for lipid metabolism. While some lipid metabolic reactions take place in the cytoplasm or cell wall, the plasma membrane is the

pivotal site for the metabolism of lipids. At the same time, this membrane must perform many other functions associated with energy production, nutrient uptake, protein export, and various sensing/signaling reactions. Studies on how these metabolic and cellular processes might be organized within bacterial plasma membranes are in their infancy. Understanding the homeostasis of the plasma membrane is particularly important in Corynebacterineae organisms because this structure must support the high biosynthetic demands of sustaining such a lipid-rich cell wall. In this chapter, we focus our discussion on processes of lipid metabolism that are critical for the biogenesis and maintenance of the plasma membrane, and illustrate the recent progress on our understanding of plasma membrane biogenesis in mycobacteria and corynebacteria.

2. Functions of plasma membrane lipids in mycobacteria and corynebacteria

In this section we will describe the functions of plasma membrane lipids. First, we will describe the functions of major structural phospholipids. We will then describe quantitatively minor lipids, which have important metabolic/physiological functions. Lastly, we will discuss the functions of neutral lipids because their biosynthesis is closely linked to phospholipid metabolism and neutral lipid storage is a critical part of plasma membrane homeostasis.

2.1. Structural lipids

Major structural components of the mycobacterial plasma membrane are phospholipids such as cardiolipin (CL), phosphatidylethanolamine (PE), phosphatidylinositol (PI), and glycosylated PIs (*i.e.* phosphatidylinositol mannosides (PIM), lipomannans (LM) and lipoarabinomannans (LAM), see below). The ratio of these phospholipids may vary depending on the species and growth conditions [6-8]. For example, one study indicated that CL, PE, and PI/PIMs represent about 37, 32, and 28%, respectively, of the total phospholipids in the plasma membrane in *M. smegmatis* [9], while another reported the ratio in *Mycobacterium phlei* to be about 50, 10, and 40% [10]. Phosphatidylglycerol (PG), which is abundant in other bacteria, is a relatively minor species in mycobacteria. Deletion of the PI biosynthetic gene has been shown to be lethal in *M. smegmatis* [9], indicating that PI or glycosylated PIs are essential for mycobacterial viability. In *M. tuberculosis*, putative PI synthetase (*Rv2612c*) and PGP synthetase (*Rv2746c*, involved in CL synthesis) genes are predicted to be essential [11], while the PS synthetase gene (*Rv0436c*, involved in PE synthesis) is not [12]. In corynebacteria, major species of phospholipids are PI, PG, CL, and acylphosphatidylglycerol (APG) [13], and PE appears to be absent.

CL is widely found in both prokaryotes and eukaryotes. It forms aggregates within the membrane bilayer. Nonyl acridine orange (NAO) is a fluorescent dye which is proposed to bind the hydrophobic surface created by the CL cluster [14], allowing microscopic visualization of CL domains. Indeed, using NAO, CLs were found to be enriched in septa and poles of actively dividing *M. tuberculosis* and *M. smegmatis* cells [15, 16]. CL has a non-

bilayer structure [17, 18], and carries a small partially immobilized head group that is more exposed to the aqueous environment than those of other glycerophospholipids [19]. Although the physiological function of CL is unclear, its physical properties may indicate that it provides a platform for membrane-protein interactions. Indeed, some mycobacterial enzymes require CL for activity [20-22], although the molecular basis for these observations has not been clarified. Recent fractionation studies in *C. glutamicum* revealed that CL (as well as other phospholipids) is enriched in the plasma membrane [23, 24]. However, a large proportion of CL is also found to be associated with the outer membrane [24], suggesting that some of these phospholipids are exported to the outer membrane in corynebacteria. Similarly, CL is released from *M. bovis* bacillus Calmette-Guerin residing in host phagosomes, and converted to lyso-CL by a host phospholipase A₂ [25]. It has been suggested that lyso-CL may influence host immune responses during infection.

PE is another major class of glycerophospholipids in mycobacteria. Although PE is generally found in all organisms, it is particularly abundant in bacterial plasma membranes [26]. Mycobacteria are no exception [20], but corynebacteria apparently lack the capacity to synthesize PE [27]. Indeed, PE biosynthetic enzymes, such as PS synthetase and PS decarboxylase, appear to be absent in corynebacterial genomes. *Corynebacterium aquaticum* has been reported to possess PE [28], but this species was later reclassified as *Leifsonia aquatica* [29], which belongs to the suborder Micrococccineae of the order Actinomycetales. The functions of PE remain elusive at the molecular level, but it appears to play important roles as a component of the plasma membrane. For example, TBsmr, a small multidrug resistance family protein from *M. tuberculosis*, shows enhanced catalytic activities when PE is supplemented in a reconstituted liposome [30].

PIs are an important class of phospholipids, and are known to be further modified by extensive glycosylation. The resultant lipoglycans, termed PIMs, LM, and LAM, are essential structural components of mycobacterial and corynebacterial cell walls. Furthermore, in pathogenic species, they have been suggested to perform additional roles in the modulation of host immune responses in favor of the pathogen through myriad effects on macrophages including cytokine production, inhibition of phagosome maturation and apoptosis [31-34]. PIMs are oligo-mannosylated PIs carrying up to 6 mannose residues while LM/LAM carry much longer mannose polymers with arabinan modifications. It remains controversial if these glycolipids are embedded in the plasma membrane or exported to the outer membrane. A recent study suggests that LM/LAM appear to be anchored to both the plasma membrane and outer membrane [35]. In *C. glutamicum*, the outer membrane and plasma membrane were fractionated on sucrose gradients upon cell lysis, and the analysis of these membrane sub-fractions demonstrated that PIMs, LM and LAM are all enriched in the plasma membrane fraction [23]. Another recent study also suggested that PI/PIMs are major components of the plasma membrane of *C. glutamicum* [24]. In the latter study, however, substantial amounts of PI/PIMs were detected in the outer membrane as well. The functional significance of these subcellular localizations, as well as the physiological roles of LM/LAM in each of these locations, remain important questions. The structural importance of PIMs remains unclear as well. For example, a *pimE*-deletion mutant that cannot produce mature

PIM6 species (see below) is viable, but shows severe plasma membrane abnormalities [36], suggesting that higher order PIMs may be involved in the maintenance of plasma membrane integrity.

It is notable that some unusual phospholipids have been identified in corynebacteria. APG is an acylated form of PG which is widespread in corynebacteria [37-40], and is a major phospholipid species in *Corynebacterium amycolatum*. Another interesting phospholipid from *C. amycolatum* is acyl-phosphatidylinositol (API), which was identified by electrospray ionization mass spectroscopy [41]. *C. amycolatum* lacks a mycolic acid-based outer membrane, and does not appear to have a fracture plane other than the plasma membrane [42]. Therefore, APG and API are likely to be components of the plasma membrane, and are suggested to play structural roles. Very little is known about their biosynthesis, and acyltransferases responsible for their synthesis remain to be identified for both lipid species.

2.2. Functional lipids

There are some examples of lipids that appear to play no structural roles in the plasma membrane. They often exist in low quantities but play important functional roles. Among these, polyprenol-phospho-sugars function as sugar donors. Two well-studied examples are polyprenol phosphomannose (PPM) and decaprenol phosphoarabinose (DPA). These molecules are the donors of mannose and arabinose, respectively, and their biosynthesis will be discussed in a later section.

PI 3-phosphate, recently identified in both *M. smegmatis* and *C. glutamicum* [43], may prove to be another interesting example of a functional lipid. It accumulates only transiently upon stimulation by high concentrations of salt, and behaves as if it is involved in a signaling cascade. However, whether PI 3-phosphate represents a mediator of stress responses remains to be addressed. More recently, lysinylated PG was identified as a minor phospholipid species in *M. tuberculosis* [44]. The synthesis of lysinylated PG is mediated by LysX and a *lysX* deletion mutant showed altered phospholipid metabolism and membrane integrity [16, 44], suggesting a regulatory role of lysinylated PG in plasma membrane homeostasis.

Carotenoids are photo-protective pigments and serve to scavenge free radicals or harvest light [45]. Several mycobacterial species are known to produce carotenoids with the notable exception of *M. tuberculosis*, despite the presence of a carotenoid oxidase in the human pathogen [46]. These hydrophobic pigments are thought to be present in the plasma membrane but whether they play structural roles in addition to a photo-protective role remains to be elucidated.

2.3. Lipid storage for energy and carbon

Neutral lipids are an important reservoir of stored energy and carbon, and their metabolism is closely linked to plasma membrane phospholipid metabolism. Unlike many other bacteria which use polyhydroxyalkanoates as a lipid storage material [47], Actinobacteria use

triacylglycerides (TAGs) as a major form of lipid storage, and the presence of TAGs has been reported in *Mycobacterium*, *Streptomyces*, *Rhodococcus* and *Nocardia* [48-52]. Interestingly, corynebacteria seem to lack the capacity to synthesize TAG, indicating that some lineages of Actinobacteria have eliminated this capacity at some point in their evolution. Recent evidence suggests that *M. tuberculosis* accumulates TAG-based lipid droplets while residing in macrophages using fatty acids released from host TAGs, and this process is critical for acquiring a dormancy phenotype [53]. Nevertheless, a mutant defective in accumulating TAG remained viable under *in vitro* dormancy-inducing conditions [54]. These somewhat contradictory observations suggest that our understanding of TAG metabolism in mycobacteria is far from complete. As we illustrate later, there appear to be several redundant genes involved in the final step of TAG synthesis, suggesting that it is an important regulatory step of lipid metabolism in these bacteria.

Cholesterol has recently been suggested to be an alternative form of lipid storage in mycobacteria. Neither mycobacteria nor corynebacteria have the capacity to synthesize cholesterol. However, cholesterol is taken up by *M. tuberculosis* cells residing in the host, and components of the *mce4* operon have been shown to be involved in cholesterol import [55]. Cholesterol catabolism is critical in the chronic phase of animal infection, and a fully functional catabolic pathway is encoded by the *M. tuberculosis* genome [56]. Furthermore, cholesterol appears to accumulate in the mycobacterial cell envelope, and this might represent a potential form of lipid storage for *M. tuberculosis* during animal infection [57, 58]. Although the authors of this study suggested that cholesterol accumulates in the outer membrane, it remains possible that the plasma membrane is the true site of accumulation. Therefore, in addition to acting as a lipid storage molecule, cholesterol may play roles in plasma membrane structure and function, and these possibilities await further exploration.

Catabolism of cholesterol, amino acids and odd-chain-length/methyl branched fatty acids produces propionyl-coenzyme A (CoA). Propionate accumulation has been shown to be toxic in various organisms [59-61], and *M. tuberculosis* has multiple pathways to metabolize propionyl-CoA [62]. Metabolized propionyl-CoA is in part incorporated into TAG [63], and it has been suggested that TAG functions as a sink for reducing equivalents in addition to being a source of carbon and energy.

3. Structure and metabolism of plasma membrane lipids in mycobacteria and corynebacteria

In this section, we will describe the structure and metabolism of various lipids found in the plasma membrane of mycobacteria and corynebacteria in more detail. Lipids are categorized into the following four classes based on their key structural features.

3.1. Fatty acids

M. tuberculosis devotes a large proportion of its coding capacity to genes involved in fatty acid metabolism [5], highlighting the importance of lipids to the organism. Fatty acid

metabolism is essential for intracellular survival of the pathogen since it forms the precursors of key membrane components such as plasma membrane phospholipids and outer membrane glycolipids. In particular, mycolic acids, which are very long chain α -alkyl β -hydroxy fatty acids, form the hydrophobic, protective mycomembrane described earlier. *M. tuberculosis* encodes two distinct enzyme systems for biosynthesis of fatty acids, designated FAS (fatty acid synthase) I and II (Fig. 2). Studies on fatty acid synthesis date back to the 1970s when *M. smegmatis* was shown to contain both type I fatty acid synthetase (FAS-I), involving a large multifunctional polypeptide, and type II fatty acid synthetase (FAS-II), consisting of a series of distinct enzymes [64]. The key elongation unit is malonyl-CoA, which is produced by acetyl-CoA carboxylase (ACCase) and the *M. tuberculosis* genome encodes several such enzymes (AccA1-3 and AccD1-6). The resultant malonyl-CoA is incorporated into fatty acids by the two FAS systems.

Figure 2. Fatty acid biosynthesis pathways in mycobacteria. Point of inhibition by the front-line tuberculosis drug isoniazid is indicated. Product profile of FAS-I is bimodal, and C₁₆-C₁₈-CoA and C₂₄-C₂₆-CoA are produced. Dashed lines indicate that some of the fatty acid products are further utilized for mycolic acid production.

3.1.1.1. De novo synthesis by FAS-I

Surprisingly, members of the Corynebacterineae use a eukaryote-like FAS-I system for *de novo* fatty acid synthesis. The single, essential [11], 9.2kb *fas* gene encodes a 326 kDa protein containing all seven domains necessary to perform the iterative series of reactions: acyl transferase, enoyl reductase, β -hydroxyacyl dehydratase, malonyl transferase, acyl carrier protein, β -ketoacyl reductase, and β -ketoacyl synthase [65, 66]. This very large protein elongates acetyl groups by 2-carbon (acetate) units using acetyl-CoA and malonyl-CoA. Early rounds of elongation yield C₁₆ to C₁₈-CoA products that are used for synthesis of membrane phospholipids or to feed into the FAS-II system. More extensive elongation yields C₂₄-C₂₆ products that ultimately form the α -branch of mycolic acids. Unlike *M. tuberculosis*, *C. glutamicum* encodes two *fas* genes (*fasA* and *fasB*) with FasA taking the

dominant role [67]. The presence of two Fas proteins may compensate for the lack of a FAS-II system in this organism.

3.1.2. Elongation by FAS-II

The FAS-II system is commonly found in bacteria and plants and, unlike FAS-I, is composed of a series of separate enzymes, each performing one step in the pathway. FAS-II elongates medium chain fatty acids derived from FAS-I using malonyl-CoA, producing C₁₈-C₃₀ fatty acids [68]. FAS-II has been extensively studied in *E. coli* [69] and orthologs of the *fab* genes have been identified in mycobacteria. AcpM is a mycobacterial acyl carrier protein (ACP) and plays a key role in transferring acyl groups between the various enzyme components [70]. The seven genes are located in two clusters on the *M. tuberculosis* chromosome [5], comprising *mtfabD-acpM-kasA-kasB-accD6* and *mabA-inhA*. Initially, the malonate group is transferred from malonyl-CoA to AcpM by the MtFabD protein. Then MtFabH performs a Claisen condensation of malonyl-ACP with acyl-CoA to form β -ketoacyl-ACP. A four-step cycle is then initiated [64] in which:

1. β -ketoacyl-ACP reductase MabA reduces the β -keto group with concomitant oxidation of NADPH
2. β -hydroxyacyl-ACP dehydratase dehydrates the β -hydroxyl to enoyl-ACP
3. enoyl-ACP reductase InhA, a target of the first-line anti-tuberculosis drug isoniazid (INH) [71], reduces enoyl-ACP to acyl-ACP with concomitant oxidation of NADPH
4. β -ketoacyl-ACP synthase KasA/B elongates acyl-ACP by 2 carbon units, forming β -ketoacyl-ACP, which can feed back into step 1.

In this way, the hydrocarbon chain increases by 2 carbons each cycle. Further elongation and processing of the products of FAS-II produces the precursors of the long meromycolate chains that are condensed with the α -branches derived from FAS-I by the large polyketide synthase Pks13 [72]. Reduction of the β -keto group by CmrA forms the mature C₆₀-C₉₀ mycolic acid [73].

3.2. Glycerolipids

Glycerolipids include both nonpolar lipids and polar phospholipids. Their biosynthesis is overlapping and 1,2-diacyl-*sn*-glycerol 3-phosphate, commonly known as phosphatidic acid (PA), is an important intermediate at the branch point (Fig. 3) [74]. In this section, we focus our discussion on the biosynthesis of PA and its conversion to non-polar lipids. Non-polar lipids are generally divided into three different classes depending on the number of fatty acids attached to glycerol: monoacylglycerol (MAG), diacylglycerol (DAG) and TAG. TAG is a glycerol carrying three fatty acyl chains, and its biosynthesis diverges from phospholipid synthesis after the synthesis of PA. TAG is a major component of lipid droplets, which accumulate in the cytoplasm. How TAG is made in the plasma membrane and incorporated into lipid droplets remains largely unclear. Here, we provide an overview of the TAG metabolic pathway.

Figure 3. Glycerolipid/phospholipid biosynthesis pathways. Some pathways such as TAG and PE biosynthesis (shown as green arrows) do not occur in corynebacteria while some others (shown as blue arrows) are known to occur only in corynebacteria. PG is abundant in corynebacteria, but is a minor species in mycobacteria.

3.2.1. Biosynthesis of PA

The first step of PA biosynthesis is mediated by glycerol phosphate acyltransferase (GPAT) transferring an acyl chain from acyl-CoA to glycerol-3-phosphate, forming acyl-glycerol 3-phosphate. In general, this reaction produces 1-acyl-*sn*-glycerol 3-phosphate. However, mycobacteria are unusual in that 2-acyl-*sn*-glycerol 3-phosphate is used as the main intermediate for the production of PA [75]. Another unusual feature is that oleic acid, an unsaturated fatty acid often found at the *sn*-2 positions of glycerolipids, is found at the *sn*-1 position in mycobacteria. Instead, palmitic acid, a saturated fatty acid, is the preferred fatty acid attached to the *sn*-2 position in mycobacteria [75, 76]. In the second step, acylglycerol phosphate acyltransferase (AGPAT) further transfers a fatty acid from acyl-CoA to 2-acyl-*sn*-glycerol 3-phosphate, producing PA. PA can be diverted to TAG synthesis, or activated

to form cytidine diphosphate-diacylglycerol (CDP-DAG), which is the precursor for the synthesis of phospholipids. Therefore, PA represents an important branch point for the synthesis of TAG and phospholipids [74]. An alternative pathway for PA synthesis is phosphorylation of DAG by DAG kinase, and Rv2252 has been suggested to be involved in this reaction [77]. Disruption of this enzyme results in altered PIM biosynthesis, but precise functions of this metabolic pathway remain unclear.

3.2.2. TAG Biosynthesis

TAG is *de novo* synthesized by two steps. First, PA is dephosphorylated to become DAG, and this reaction is mediated by phosphatidic acid phosphatase (PAP). PAP was discovered from animal tissues in 1957 by the group of Eugene Kennedy [78], and the gene encoding this activity was recently identified in *Saccharomyces cerevisiae* [79]. Nothing is known about this enzyme in mycobacteria or corynebacteria. In the second step, diacylglycerol acyltransferase (DGAT) catalyzes the addition of a fatty acyl-CoA to DAG to form TAG. Until recently, little was known about the genes involved in this final step of TAG synthesis in mycobacteria. Analysis of this final step is complicated because there are multiple genes encoding TAG synthetase in mycobacteria and corynebacteria. For example, the *M. tuberculosis* genome encodes 15 putative TAG synthetase genes [48, 80]. Despite the redundancies, recent studies reported that some of these *tgs* genes are critical for TAG synthesis in *M. tuberculosis* [48, 54]. Specifically, TAG synthetases encoded by *Rv3130c* (*tgs1*), *Rv3734c* (*tgs2*), *Rv3234c* (*tgs3*), and *Rv3088* (*tgs4*) have been shown to have TAG synthetase activities [53]. Furthermore, Tgs1 has been demonstrated to be the main contributor to TAG synthesis and lipid droplet formation in *M. tuberculosis* [53]. More recently, Ag85A, which is known as a mycolyltransferase involved in TDM biosynthesis, was shown to possess DGAT activity [81]. Ag85A is not homologous to other *tgs* genes, and may represent a novel class of TAG biosynthetic enzymes. TAG not only forms a lipid droplet in the cytoplasm, but also accumulates in the cell wall of mycobacteria [82]. Therefore, Ag85A located in the cell wall might be involved in the production of surface-exposed TAGs.

3.2.3. Utilization of TAG

Under starvation conditions where stored TAG needs to be mobilized for energy production, TAG is catabolized by lipases. In 1977, TAG lipase was purified from stationary phase *M. phlei* and predicted to have a molecular weight of about 40 kDa [83]. More recently, LipY, encoded by the *M. tuberculosis* *Rv3097c* gene, was identified as a TAG lipase [84]. LipY appears to play a critical role in TAG catabolism because a *M. tuberculosis* *lipY* deletion mutant cannot utilize accumulated TAG under starvation conditions. Another recent study demonstrated that LipY has a dual localization pattern [85]: while a fraction of LipY was found in the cytoplasm, consistent with its role in the catabolism of intracellular TAG, a significant fraction of LipY was also localized to the outer membrane of the cell wall, indicating that it may be involved in the breakdown of exogenously available TAGs. Indeed, it has been long known that *M. tuberculosis* depends on fatty

acids as a preferred energy source during infection [86], and LipY may well be a critical enzyme for the utilization of host lipids during an *M. tuberculosis* infection. Another lipase encoded by *Rv0183* shows preference for MAG over DAG and TAG, and is localized to the cell wall [87], suggesting its involvement in subsequent reactions of TAG breakdown. However, whether it is involved in degradation of host-derived TAG or intracellular TAG remains to be determined.

3.2.4. Lipid droplet formation

In eukaryotes, lipid droplets form in between the two leaflets of the endoplasmic reticulum membrane [88]. In bacteria, a distinct mechanism of lipid body formation has been proposed. For example, in rhodococci, TAG is formed in the cytoplasmic surface of the plasma membrane. Small lipid droplets are then fused to each other, coated by a monolayer of phospholipids, and released from the surface of the plasma membrane into the cytoplasm as mature lipid droplets [89]. Although no endogenous proteins have been found to associate with lipid droplets in rhodococci or mycobacteria, heterologous expression of known lipid droplet-associated proteins resulted in correct targeting of these proteins to lipid droplets in both *R. opacus* and *M. smegmatis* [90, 91], allowing visualization of lipid droplets in these organisms.

3.3. Phospholipids

3.3.1. CDP-DAG

In both eukaryotic and prokaryotic cells, PA is activated by CTP to form CDP-DAG, and this reaction is mediated by CDP-DAG synthase [92]. The synthesis of CDP-DAG commits the pathway to phospholipid biosynthesis, and CDP-DAG is a common precursor for the biosynthesis of all glycerophospholipids in mycobacteria and corynebacteria. The activity of CDP-DAG synthetase is associated with plasma membrane in *M. smegmatis*, and is possibly encoded by the *cdsA* (*Rv2881c*) gene in *M. tuberculosis* H37Rv [93].

3.3.2. CL

CL is composed of four acyl chains, three glycerols and two phosphates, and is structured in a 1,3-diphosphatidylglycerol configuration [94]. It is a common phospholipid in bacteria, and is one of the abundant phospholipids in mycobacteria and corynebacteria. To initiate CL synthesis, PG phosphate synthase first produces PG phosphate (PGP) using CDP-DAG and glycerol 3-phosphate as substrates. An *M. smegmatis* strain engineered to overexpress *M. tuberculosis* PgsA3 (encoded by *Rv2746c*) was shown to overproduce PG, suggesting that PgsA3 is the PGP synthase [9]. PGP is then converted into PG via PGP phosphatase. Three phosphatases, PgpA, PgpB, and PgpC, have been identified as PGP phosphatases in *E. coli* [95-97]. Furthermore, Gep4 and PTPMT1 have been identified as PGP phosphatases in yeast and mammals, respectively [98, 99]. Some homologs exist in the genomes of mycobacteria and corynebacteria, but experimental verification of these genes remains to be performed.

Typically, the final step of CL synthesis in prokaryotes is mediated by a reaction that utilizes two PG molecules, producing one molecule of CL and one molecule of glycerol. However, in mycobacteria, the eukaryote-like reaction, which utilizes PG and CDP-DAG to produce CL, has been shown to occur [100], and Jackson and colleagues have suggested that PgsA2 might be the enzyme responsible for this reaction [9].

3.3.3. PE

The precise structure of PE was recently reported as 1-*O*-tuberculostearoyl-2-*O*-palmitoyl-*sn*-glycero-3-phosphoethanolamine in *M. tuberculosis* using tandem mass spectrometry [101]. In the initial step of PE synthesis, PS synthetase transfers serine to CDP-DAG and produces PS. In the second step, PE is produced by decarboxylation of PS mediated by PS decarboxylase. Genes encoding putative PS synthetase (*pssA*, *Rv0436c*) and PS decarboxylase (*psd*, *Rv0437c*) are found in tandem in the *M. tuberculosis* genome [9]. However, there is no experimental evidence demonstrating the identities of the genes. In *M. smegmatis*, PS synthetase and PS decarboxylase activities are enriched in different membrane fractions, which can be distinguished by sucrose gradient sedimentation [102]. However, the significance of differential membrane localization remains to be clarified.

3.3.4. PI

PI is a major phospholipid in both mycobacteria and corynebacteria and forms the anchor for the PIMs, which are substrates for heavy mannosylation to form LMs and additional arabinosylation to produce LAMs. PI is formed by the PI synthase PgsA (*Rv2612c*) from CDP-DAG and *myo*-inositol [9, 103]. Inositol is not a common metabolite in bacteria. It is therefore surprising that mycobacteria produce copious amounts of inositol through pathways shared with eukaryotes for incorporation into a range of metabolic pathways (recently reviewed in [104]). The enzyme D-*myo*-inositol 3-phosphate synthase (*Ino1*) converts glucose-6-phosphate to D-*myo*-inositol 3-phosphate, which is dephosphorylated by one of several inositol monophosphatases to produce *myo*-inositol [105].

3.3.5. PIMs

All Corynebacterineae synthesize PIMs that are important components of the cell envelope. Polar PIM species can also serve as membrane anchors for LM and LAM. Many of the steps of PIM/LM/LAM biosynthesis have now been elucidated [106]. Extensive genetic and biochemical studies have demonstrated that the synthesis of PIMs occurs linearly in mycobacteria with PI as the starting substrate (reviewed in [107]) (Fig. 4A). Early steps of the pathway occur on the cytoplasmic face of the plasma membrane. A PIM biosynthetic membrane, enriched in the early steps, has been purified by sucrose gradient fractionation as a membrane subdomain termed PMf, which is distinct from the bulk plasma membrane [102]. Mannosyltransferases performing the early steps utilize the water-soluble mannose donor, GDP-Man, which can be produced from exogenously acquired mannose or via *de novo* synthesis from the glycolytic pathway when fructose-6-phosphate is transformed by

the enzymes ManA (Rv3255c), ManB (Rv3257c) and ManC (Rv3264c) [108-111], with a degree of redundancy reported at the ManC (NCgl0710) step in *C. glutamicum* [112]. The first step of PIM synthesis involves mannosylation of the C2-position of the inositol ring of PI by the enzyme PimA (Rv2610c) to form PIM1 [113, 114]. PimA is an essential enzyme in mycobacteria [11, 113] and absent in humans, making it a current target for drug development by several groups. The crystal structure of PimA from *M. smegmatis* has been solved in complex with GDP and GDP-Man at resolutions of 2.4Å and 2.6Å, respectively [115, 116]. PimA has a typical GT-B fold of glycosyltransferases consisting of two Rossmann-fold domains with a deep fissure at the interface, which contains the active site. Close to the GDP-Man binding site, the N-terminal domain displays a deep pocket containing highly conserved hydrophobic residues. This pocket is proposed to bind the acyl moieties of the acceptor substrate PI [116].

Next, O-6 mannosylation of the *myo*-inositol ring is performed by the cytoplasmic α -mannosyltransferase PimB' (Rv2188c) [117, 118] resulting in the formation of PIM2. While the mycobacterial enzyme Rv0557 was originally assigned this function [119], later studies showed that Rv2188c was the true PimB (designated PimB') [118], with Rv0557 being renamed MgtA and included in the LM-B pathway [120, 121] (see below). PimB' is an essential enzyme in mycobacteria [11, 117] and the crystal structure of the equivalent corynebacterial enzyme has been solved at high resolution complexed with nucleotide [122]. In corynebacteria, PIM2 is detected mainly in its mono-acylated form, but in mycobacteria PIM2 accumulates in the cell envelope as monoacyl (AcPIM2) and diacyl (Ac₂PIM2) forms, the former produced by the acyltransferase Rv2611c [123] which acts optimally on the product of the PimB' reaction [117].

The next enzyme in the pathway, PimC, has been identified in *M. tuberculosis* strain CDC1551 and could produce trimannosylated PIMs [124]. However, the absence of this enzyme in other strains indicates redundancy at this step and the enzymes involved remain to be identified, as does the putative "PimD" protein. Flipping of PIM intermediates from the cytoplasmic face of the membrane to the periplasm is thought to occur at this point of the pathway, but the precise intermediate and transporter involved are also undefined.

AcPIM4 species can be further mannosylated to form more polar PIMs in reactions thought to take place on the periplasmic side of the cytoplasmic membrane. These reactions are performed by glycosyltransferases that require a lipid sugar donor in the form of PPM, since these reactions are amphotycin-sensitive [36, 125-128]. In mycobacteria, AcPIM4 is proposed to be a branch point for synthesis of polar PIM end products and LM/LAM. PimE (Rv1159) has been shown to elongate AcPIM4 with one or more α 1-2 linked mannoses to form AcPIM6 [36]. This polytopic membrane protein has sequence similarities with eukaryotic PIG-M mannosyltransferases and localizes to a cell wall-associated plasma membrane subdomain in *M. smegmatis*, termed PM-CW, to which enzymatic activities of AcPIM4-6 synthesis are enriched [102]. Its catalytic activity was successfully mapped to a conserved aspartate residue in the first outer loop of the protein. Whether PimE also forms AcPIM6 is unknown. Interestingly, no PimE orthologue is present in *C. glutamicum* and

Ac/Ac₂PIM6 do not accumulate in this species, so the formation of Ac/Ac₂PIM5/6 appears to be a mycobacterium-specific side-branch of the pathway.

Surprisingly, studies in *M. smegmatis* have revealed a role for a lipoprotein in PIM/LAM synthesis, since *lpqW* mutants produce reduced levels of LM/LAM [129]. As the only non-enzymatic component of the pathway identified to date, LpqW has been proposed to have a regulatory role at the bifurcation point of the pathways, as well as a functional connection with PimE, since mutations in the *pimE* gene can bypass the requirement for LpqW [130]. Very recent studies have implicated the corynebacterial ortholog of LpqW in the LM-B pathway as well (Rainczuk *et al*, submitted). Structural studies on the *M. smegmatis* LpqW revealed a scaffold similar to substrate binding proteins associated with ABC transporters, which has evolved to fulfill a new role in the regulation of PIM/LAM biosynthesis [131].

3.3.6. LM/LAM

A subpopulation of PIMs (AcPIM4 in mycobacteria [128, 129, 132] and AcPIM2 in corynebacteria [133]) can be extended with chains of α 1-6 linked mannose to form LM that is further modified with a number of single α 1-2 mannose side chains [134-136]. MptB is a PPM-dependent mannosyltransferase involved in extending AcPIM2 to form the proximal α 1-6 mannan backbone of LM in *C. glutamicum* but is redundant in *M. smegmatis* [133], indicating additional complexity at this step in mycobacteria. Further elongation is performed by MptA (Rv2174) [135, 136], with MptC (Rv2181) required for addition of α 1-2 linked Man side chains [106, 134, 137, 138]. Further additions of arabinose units by EmbC (Rv3793), AftC (Rv2673), AftD (Rv0236c) and at present unidentified α 1-5 arabinofuranosyltransferases, result in the formation of mature LAM [139-142]. In *M. tuberculosis* and other pathogenic mycobacteria, additional mannose capping is present [143], synthesized by the enzymes MptC [137] and Rv1635c [144]. Alternatively, *M. smegmatis* LAM is capped with inositol phosphate [145].

While the general PI→PIM→LM→LAM pathway is conserved in corynebacteria, a second pathway of lipoglycan biosynthesis exists in which a sub-population of LM lipoglycans is assembled on a glucopyranosyluronic acid diacylglycerol (Gl-A, GlcADAG) glycolipid anchor [118, 121, 133]. In this pathway (Fig. 4B), Gl-A is first mannosylated by MgtA (NCgl0452 in *C. glutamicum* and previously termed PimB, see above) forming mannosyl-glucuronic acid diacylglycerol (Gl-X, ManGlcADAG), which is subjected to further α 1-6 (backbone) and α 1-2 (side unit) mannosylation resulting in LM [121]. This pathway shares some PPM-dependent enzymes with the PI-based LM pathway including MptB, since an *mptB* (NCgl1505) mutant of *C. glutamicum* fails to produce intermediates beyond Gl-X or AcPIM2 [133]. For clarity, the PI-based LM pool has been designated LM-A while this second pathway produces LM-B [121, 146], the major LM pool in *C. glutamicum* [118]. While *C. glutamicum* produces LAMs via LM-A using a similar pathway to mycobacteria, its LAMs are smaller and structurally distinct, with more extensive mannosylation and singular Araf capping [147].

Figure 4. Summary of Lipoglycan Biosynthesis Pathways in Corynebacterineae. A) The mycobacterial pathway and structures for LM-A/LAM are shown, although several steps are inferred from studies in *C. glutamicum*. A progression from PIM1 to AcPIM2 via AcPIM1 can also occur but is sub-optimal in mycobacteria. AcPIM3 is shown as the substrate for the unidentified flippase but AcPIM2 may be the preferred substrate: AcPIM2 appears to be the flipped intermediate in *C. glutamicum*. Reactions associated with the PMf and PM-CW subfractions of *M. smegmatis* are shaded orange and grey, respectively. B) LM-B pathway in *C. glutamicum*. Its presence in mycobacteria remains to be determined. Intermediates representing Gl-Y and Gl-Z have been detected in very recent studies (Rainczuk et al, submitted). C) PPM serves as the lipid-linked donor of mannose for both pathways. The mechanism of flipping is undetermined. PM, plasma membrane. Other abbreviations are as defined in the text.

3.4. Prenol lipids

Polyprenol phosphate (Pol-P) is a key carrier lipid in synthesis of the core structures of the mycobacterial cell wall, including peptidoglycan and arabinogalactan. Unlike most bacteria, mycobacteria contain multiple types of Pol-P. For example, *M. smegmatis* produces decaprenol phosphate (C₅₀, Dec-P, [148]) and heptaprenol phosphate (C₃₅, Hep-P, [149]). Polyprenols are thought to be synthesized via the condensation of two C₅ lipids, isopentenyl diphosphate [150-152] and dimethylallyl diphosphate derived from the mevalonate-independent methylerythritol 4-phosphate (MEP) pathway. These reactions are catalyzed by prenil diphosphate synthases and two *M. tuberculosis* proteins, Rv2361c and Rv1086, acting

sequentially, are thought to fulfill this role [148, 149, 153-155]. Finally, the C₅₀ decaprenol diphosphate is dephosphorylated to produce Dec-P by an unknown phosphatase.

3.4.1. Polyprenol-phospho-sugars

3.4.1.1. PPM

PPM, a β -D-mannosyl-1-monophosphoryldecaprenol, is utilized by periplasmic mannosyltransferases for synthesis of polar PIM species and LMs [106]. C₃₅/C₅₀-P-Man_p is formed by the PPM synthase Ppm1 (Rv2051c) from GDP-Man_p and Pol-P [126, 127] (Fig. 4C). In *M. tuberculosis* Ppm1 consists of two domains: a C-terminal catalytic domain and a N-terminal membrane anchor with 6 transmembrane helices. However, the equivalent domains of the *M. smegmatis* PPM synthase are two distinct, but interacting, proteins [156]. The importance of PPM in cell wall synthesis has been highlighted by analysis of a *ppm1* mutant in *C. glutamicum* that failed to produce PPM, resulting in severe defects in lipoglycan biosynthesis. While the mutant could synthesize PIM2 species, all downstream products (LM, LAM) were absent, indicating their reliance on the PPM donor [127]. These findings were consistent with others showing that amphomycin, an antibiotic specific for PPM-dependent polymerases, blocked the PIM pathway at the PIM2 or PIM3 stage [125, 128]. The product of the *Rv3779* gene has also been implicated in PPM synthesis but its role remains unclear [157, 158].

3.4.1.2. DPA

DPA is the only known donor of arabinose (Ara) for mycobacterial cell wall synthesis, contributing Ara_f units to arabinogalactan and LAM [106] with concomitant release of Pol-P. The Ara_f portion is derived from the pentose-phosphate pathway [159-161]. 5-phosphoribose 1-diphosphate is transferred to Dec-P by the *Rv3806c* gene product [162] and the resultant Dec-P- β -D-5-phosphoribose is dephosphorylated to form Dec-P- β -D-ribose. Oxidation of the 2' hydroxyl is followed by a reduction reaction to form DPA. This two-step epimerization reaction is catalyzed by the combined activities of DprE1 (Rv3790) and DprE2 (Rv3791) [163]. Since DPA is the sole donor of Ara_f residues for mycobacterial cell wall synthesis, this pathway is of interest for drug development. Indeed, DprE1 is the target of dinitrobenzamide derivatives (DNBs) [164] and a set of nitro-compounds related to DNBs, the nitro-benzothiazinones (BTZ), a class of compounds with nanomolar anti-*M. tuberculosis* activities but minimal host-cell toxicity [165-167]. The essential nature of DprE1 in species beyond *M. tuberculosis* [168] reinforces it as a "magic" drug target [169]. The crystal structure of *M. tuberculosis* DprE1 complexed with BTZ inhibitors has been reported very recently, revealing the mode of inhibitor binding [170].

3.4.2. Carotenoids

Carotenoids are isoprenoid pigments widely distributed in biology and mostly based on C₄₀-polyene. Synthesis of these pigments has been poorly studied in mycobacteria but they have proven useful for taxonomic and identification purposes. Mycobacterial pigments are

generally yellow or orange and most have been confirmed as carotenoids [171]. While carotenoid genetics has been best studied in plants, the key enzymes of the pathway have been identified in bacteria, including mycobacteria [172-174]. There are two classes of carotenoids in bacteria, carotenes and xanthophylls, the latter of which contain oxygen. Both classes are composed of eight isoprenoid units with a long central chain of double bonded carbons. A consensus pathway for carotenoid biosynthesis in bacteria has been elucidated with orthologues of key enzymes identified in several species of mycobacteria [175]. As described above for prenol lipids, the carotenoid pathway begins with isopentenyl diphosphate and dimethylallyl diphosphate derived from the MEP pathway [176]. Head-to-tail condensation of these terpenes produces geranylgeranyl pyrophosphate (GGPP) due to the activity of GGPP synthase (CrtE). Condensation of two GGPP molecules [177] is followed by desaturation to phytoene by phytoene synthase (CrtB). Phytoene desaturase (CrtI) converts phytoene to lycoprene followed by cyclization to β -carotene by lycoprene cyclase (CrtY) [178].

4. Concluding remarks

Lipid metabolism in mycobacteria and corynebacteria is a highly complex network of catabolic and anabolic reactions. While the metabolic pathways and many of the enzymes involved have been actively elucidated over the past decade, substantial efforts are still needed to draw a comprehensive map of lipid metabolism in these organisms. In particular, our understanding of regulatory mechanisms of lipid metabolism is currently at an early stage. In addition, there are very few studies describing the interactions between multiple metabolic pathways of lipid biosynthesis. One promising approach for the comprehensive understanding of lipid metabolism is lipidomics, which is the study of lipid biosynthetic and catabolic pathways at a global level [179]. In the past, metabolic pathways have generally been examined in isolation without consideration of how different pathways might interact with, and influence, one another. Since the plasma membrane is a shared platform for most lipid biosynthetic pathways, and some donors are shared between different pathways (see above), it seems unlikely that the various pathways are truly independent. Recent advances in mass spectrometry (*e.g.* MALDI-MS, ESI-MS), nuclear magnetic resonance spectroscopy and associated computational methods have fuelled the development of this field [180]. Members of the Corynebacterineae, with their extensive lipid repertoires and complex metabolic pathways, would seem to be ideal targets to assess the true potential of lipidomics technologies. Recently, appropriate databases and methods for detection and identification of all major lipid classes of *M. tuberculosis* from a single crude extract have been developed [181, 182]. Very recently, a lipidomics profiling platform has been reported that uses high-performance liquid chromatography/mass spectrometry to resolve more than 12,000 molecules from *M. tuberculosis* [183]. These exciting advances provide a basis for future studies on the regulation of lipid metabolism and may allow, for the first time, a true appreciation of the interactive lipid networks of mycobacteria and corynebacteria.

Author details

Paul K. Crellin

Australian Research Council Centre of Excellence in Structural and Functional Microbial Genomics, Department of Microbiology, Monash University, Australia

Chu-Yuan Luo and Yasu S. Morita*

Department of Microbiology, University of Massachusetts, Amherst, USA

5. References

- [1] 1. Hoffmann C, Leis A, Niederweis M, Plitzko JM, Engelhardt H. Disclosure of the mycobacterial outer membrane: cryo-electron tomography and vitreous sections reveal the lipid bilayer structure. *Proc Natl Acad Sci U S A*. 2008;105:3963-7.
- [2] Zuber B, Chami M, Houssin C, Dubochet J, Griffiths G, Daffe M. Direct visualization of the outer membrane of mycobacteria and corynebacteria in their native state. *J Bacteriol*. 2008;190:5672-80.
- [3] Barry CE, 3rd, Mdluli K. Drug sensitivity and environmental adaptation of mycobacterial cell wall components. *Trends Microbiol*. 1996;4:275-81.
- [4] Daffe M, Draper P. The envelope layers of mycobacteria with reference to their pathogenicity. *Adv Microb Physiol*. 1998;39:131-203.
- [5] Cole ST, Brosch R, Parkhill J, Garnier T, Churcher C, Harris D, et al. Deciphering the biology of *Mycobacterium tuberculosis* from the complete genome sequence. *Nature*. 1998;393:537-44.
- [6] Nandedkar AK. Comparative study of the lipid composition of particular pathogenic and nonpathogenic species of *Mycobacterium*. *J Natl Med Assoc*. 1983;75:69-74.
- [7] Subramoniam A, Subrahmanyam D. Light-induced changes in the phospholipid composition of *Mycobacterium smegmatis* ATCC 607. *J Gen Microbiol*. 1982;128:419-21.
- [8] Khuller GK, Taneja R, Nath N. Effect of fatty acid supplementation on the lipid composition of *Mycobacterium smegmatis* ATCC 607, grown at 27 degrees and 37 degrees C. *J Appl Bacteriol*. 1983;54:63-8.
- [9] Jackson M, Crick DC, Brennan PJ. Phosphatidylinositol is an essential phospholipid of mycobacteria. *J Biol Chem*. 2000;275:30092-9.
- [10] Akamatsu Y, Nojima S. Separation and analyses of the individual phospholipids of mycobacteria. *J Biochem*. 1965;57:430-9.
- [11] Sassetti CM, Boyd DH, Rubin EJ. Genes required for mycobacterial growth defined by high density mutagenesis. *Mol Microbiol*. 2003;48:77-84.
- [12] Lamichhane G, Zignol M, Blades NJ, Geiman DE, Dougherty A, Grosset J, et al. A postgenomic method for predicting essential genes at subsaturation levels of mutagenesis: application to *Mycobacterium tuberculosis*. *Proc Natl Acad Sci U S A*. 2003;100:7213-8.

* Corresponding Author

- [13] Yague G, Segovia M, Valero-Guillen PL. Phospholipid composition of several clinically relevant *Corynebacterium* species as determined by mass spectrometry: an unusual fatty acyl moiety is present in inositol-containing phospholipids of *Corynebacterium urealyticum*. *Microbiology*. 2003;149:1675-85.
- [14] Mileykovskaya E, Dowhan W, Birke RL, Zheng D, Lutterodt L, Haines TH. Cardiolipin binds nonyl acridine orange by aggregating the dye at exposed hydrophobic domains on bilayer surfaces. *FEBS Lett*. 2001;507:187-90.
- [15] Maloney E, Madiraju SC, Rajagopalan M, Madiraju M. Localization of acidic phospholipid cardiolipin and DnaA in mycobacteria. *Tuberculosis (Edinb)*. 2011;91 Suppl 1:S150-5.
- [16] Maloney E, Lun S, Stankowska D, Guo H, Rajagoapalan M, Bishai WR, et al. Alterations in phospholipid catabolism in *Mycobacterium tuberculosis* lysX mutant. *Front Microbiol*. 2011;2:19.
- [17] Dahlberg M. Polymorphic phase behavior of cardiolipin derivatives studied by coarse-grained molecular dynamics. *J Phys Chem B*. 2007;111:7194-200.
- [18] Powell GL, Marsh D. Polymorphic phase behavior of cardiolipin derivatives studied by ³¹P NMR and X-ray diffraction. *Biochemistry*. 1985;24:2902-8.
- [19] Lewis RN, Zweytick D, Pabst G, Lohner K, McElhaney RN. Calorimetric, x-ray diffraction, and spectroscopic studies of the thermotropic phase behavior and organization of tetramyristoyl cardiolipin membranes. *Biophys J*. 2007;92:3166-77.
- [20] Imai T, Kageyama Y, Tobari J. *Mycobacterium smegmatis* malate dehydrogenase: activation of the lipid-depleted enzyme by anionic phospholipids and phosphatidylethanolamine. *Biochim Biophys Acta*. 1995;1246:189-96.
- [21] Kimura T, Tobari J. Participation of flavin-adenine dinucleotide in the activity of malate dehydrogenase from *Mycobacterium avium*. *Biochim Biophys Acta*. 1963;73:399-405.
- [22] Tobari J. Requirement of flavin adenine dinucleotide and phospholipid for the activity of malate dehydrogenase from *Mycobacterium avium*. *Biochem Biophys Res Commun*. 1964;15:50-4.
- [23] Marchand CH, Salmeron C, Bou Raad R, Meniche X, Chami M, Masi M, et al. Biochemical disclosure of the mycolate outer membrane of *Corynebacterium glutamicum*. *J Bacteriol*. 2012;194:587-97.
- [24] Bansal-Mutalik R, Nikaido H. Quantitative lipid composition of cell envelopes of *Corynebacterium glutamicum* elucidated through reverse micelle extraction. *Proc Natl Acad Sci U S A*. 2011;108:15360-5.
- [25] Fischer K, Chatterjee D, Torrelles J, Brennan PJ, Kaufmann SH, Schaible UE. Mycobacterial lysocardiolipin is exported from phagosomes upon cleavage of cardiolipin by a macrophage-derived lysosomal phospholipase A2. *J Immunol*. 2001;167:2187-92.
- [26] Murzyn K, Rog T, Pasenkiewicz-Gierula M. Phosphatidylethanolamine-phosphatidylglycerol bilayer as a model of the inner bacterial membrane. *Biophys J*. 2005;88:1091-103.
- [27] Brennan PJ, Lehane DP. The phospholipids of corynebacteria. *Lipids*. 1971;6:401-9.

- [28] Khuller GK, Brennan PJ. Further studies on the lipids of corynebacteria. The mannolipids of *Corynebacterium aquaticum*. *Biochem J.* 1972;127:369-73.
- [29] Suzuki KI, Suzuki M, Sasaki J, Park YH, Komagata KK. Leifsonia gen. nov., a genus for 2,4-diaminobutyric acid-containing actinomycetes to accommodate "*Corynebacterium aquaticum*" Leifson 1962 and *Clavibacter xyli* subsp. *cynodontis* Davis et al. 1984. *J Gen Appl Microbiol.* 1999;45:253-62.
- [30] Charalambous K, Miller D, Curnow P, Booth PJ. Lipid bilayer composition influences small multidrug transporters. *BMC Biochem.* 2008;9:31.
- [31] Chatterjee D, Khoo KH. Mycobacterial lipoarabinomannan: an extraordinary lipoheteroglycan with profound physiological effects. *Glycobiology.* 1998;8:113-20.
- [32] Nigou J, Gilleron M, Rojas M, Garcia LF, Thurnher M, Puzo G. Mycobacterial lipoarabinomannans: modulators of dendritic cell function and the apoptotic response. *Microbes Infect.* 2002;4:945-53.
- [33] Strohmeier GR, Fenton MJ. Roles of lipoarabinomannan in the pathogenesis of tuberculosis. *Microbes Infect.* 1999;1:709-17.
- [34] Vercellone A, Nigou J, Puzo G. Relationships between the structure and the roles of lipoarabinomannans and related glycoconjugates in tuberculosis pathogenesis. *Front Biosci.* 1998;3:e149-63.
- [35] Pitarque S, Larrouy-Maumus G, Payre B, Jackson M, Puzo G, Nigou J. The immunomodulatory lipoglycans, lipoarabinomannan and lipomannan, are exposed at the mycobacterial cell surface. *Tuberculosis (Edinb).* 2008;88:560-5.
- [36] Morita YS, Sena CB, Waller RF, Kurokawa K, Sernee MF, Nakatani F, et al. PimE is a polyprenol-phosphate-mannose-dependent mannosyltransferase that transfers the fifth mannose of phosphatidylinositol mannoside in mycobacteria. *J Biol Chem.* 2006;281:25143-55.
- [37] Lechevalier MP, Debievre C, Lechevalier H. Chemotaxonomy of aerobic actinomycetes - phospholipid composition. *Biochem Syst Ecol.* 1977;5:249-60.
- [38] Yague G, Segovia M, Valero-Guillen PL. Acyl phosphatidylglycerol: a major phospholipid of *Corynebacterium amycolatum*. *FEMS Microbiol Lett.* 1997;151:125-30.
- [39] Niepel T, Meyer H, Wray V, Abraham WR. Intraspecific variation of unusual phospholipids from *Corynebacterium* spp. containing a novel fatty acid. *J Bacteriol.* 1998;180:4650-7.
- [40] Mazzella N, Molinet J, Syakti AD, Dodi A, Doumenq P, Artaud J, et al. Bacterial phospholipid molecular species analysis by ion-pair reversed-phase HPLC/ESI/MS. *J Lipid Res.* 2004;45:1355-63.
- [41] Valero-Guillen PL, Yague G, Segovia M. Characterization of acyl-phosphatidylinositol from the opportunistic pathogen *Corynebacterium amycolatum*. *Chem Phys Lipids.* 2005;133:17-26.
- [42] Puech V, Chami M, Lemassu A, Laneelle MA, Schiffler B, Gounon P, et al. Structure of the cell envelope of corynebacteria: importance of the non-covalently bound lipids in the formation of the cell wall permeability barrier and fracture plane. *Microbiology.* 2001;147:1365-82.

- [43] Morita YS, Yamaryo-Botte Y, Miyanagi K, Callaghan JM, Patterson JH, Crellin PK, et al. Stress-induced synthesis of phosphatidylinositol 3-phosphate in mycobacteria. *J Biol Chem.* 2010;285:16643-50.
- [44] Maloney E, Stankowska D, Zhang J, Fol M, Cheng QJ, Lun S, et al. The two-domain LysX protein of *Mycobacterium tuberculosis* is required for production of lysinylated phosphatidylglycerol and resistance to cationic antimicrobial peptides. *PLoS Pathog.* 2009;5:e1000534.
- [45] Fraser PD, Bramley PM. The biosynthesis and nutritional uses of carotenoids. *Prog Lipid Res.* 2004;43:228-65.
- [46] Scherzinger D, Scheffer E, Bar C, Ernst H, Al-Babili S. The *Mycobacterium tuberculosis* ORF *Rv0654* encodes a carotenoid oxygenase mediating central and excentric cleavage of conventional and aromatic carotenoids. *FEBS J.* 2010;277:4662-73.
- [47] Steinbuchel A, Valentin HE. Diversity of bacterial polyhydroxyalkanoic acids. *FEMS Microbiol Lett.* 1995;128:219-28.
- [48] Daniel J, Deb C, Dubey VS, Sirakova TD, Abomoelak B, Morbidoni HR, et al. Induction of a novel class of diacylglycerol acyltransferases and triacylglycerol accumulation in *Mycobacterium tuberculosis* as it goes into a dormancy-like state in culture. *J Bacteriol.* 2004;186:5017-30.
- [49] Alvarez HM, Steinbuchel A. Triacylglycerols in prokaryotic microorganisms. *Appl Microbiol Biotechnol.* 2002;60:367-76.
- [50] Alvarez HM, Kalscheuer R, Steinbuchel A. Accumulation and mobilization of storage lipids by *Rhodococcus opacus* PD630 and *Rhodococcus ruber* NCIMB 40126. *Appl Microbiol Biotechnol.* 2000;54:218-23.
- [51] Alvarez HM, Luftmann H, Silva RA, Cesari AC, Viale A, Waltermann M, et al. Identification of phenyldecanoic acid as a constituent of triacylglycerols and wax ester produced by *Rhodococcus opacus* PD630. *Microbiology.* 2002;148:1407-12.
- [52] Olukoshi ER, Packter NM. Importance of stored triacylglycerols in *Streptomyces*: possible carbon source for antibiotics. *Microbiology.* 1994;140 (Pt 4):931-43.
- [53] Daniel J, Maamar H, Deb C, Sirakova TD, Kolattukudy PE. *Mycobacterium tuberculosis* uses host triacylglycerol to accumulate lipid droplets and acquires a dormancy-like phenotype in lipid-loaded macrophages. *PLoS Pathog.* 2011;7:e1002093.
- [54] Sirakova TD, Dubey VS, Deb C, Daniel J, Korotkova TA, Abomoelak B, et al. Identification of a diacylglycerol acyltransferase gene involved in accumulation of triacylglycerol in *Mycobacterium tuberculosis* under stress. *Microbiology.* 2006;152:2717-25.
- [55] Pandey AK, Sassetti CM. Mycobacterial persistence requires the utilization of host cholesterol. *Proc Natl Acad Sci U S A.* 2008;105:4376-80.
- [56] Griffin JE, Gawronski JD, Dejesus MA, Ioerger TR, Akerley BJ, Sassetti CM. High-resolution phenotypic profiling defines genes essential for mycobacterial growth and cholesterol catabolism. *PLoS Pathog.* 2011;7:e1002251.
- [57] Brzostek A, Pawelczyk J, Rumijowska-Galewicz A, Dziadek B, Dziadek J. *Mycobacterium tuberculosis* is able to accumulate and utilize cholesterol. *J Bacteriol.* 2009;191:6584-91.

- [58] Av-Gay Y, Sobouti R. Cholesterol is accumulated by mycobacteria but its degradation is limited to non-pathogenic fast-growing mycobacteria. *Can J Microbiol.* 2000;46:826-31.
- [59] Wright LD, Skeggs HR. Reversal of sodium propionate inhibition of *Escherichia coli* with beta-alanine. *Arch Biochem.* 1946;10:383-6.
- [60] Brock M, Buckel W. On the mechanism of action of the antifungal agent propionate. *Eur J Biochem.* 2004;271:3227-41.
- [61] Maruyama K, Kitamura H. Mechanisms of growth inhibition by propionate and restoration of the growth by sodium bicarbonate or acetate in *Rhodopseudomonas sphaeroides* S. *J Biochem.* 1985;98:819-24.
- [62] Savvi S, Warner DF, Kana BD, McKinney JD, Mizrahi V, Dawes SS. Functional characterization of a vitamin B12-dependent methylmalonyl pathway in *Mycobacterium tuberculosis*: implications for propionate metabolism during growth on fatty acids. *J Bacteriol.* 2008;190:3886-95.
- [63] Singh A, Crossman DK, Mai D, Guidry L, Voskuil MI, Renfrow MB, et al. *Mycobacterium tuberculosis* WhiB3 maintains redox homeostasis by regulating virulence lipid anabolism to modulate macrophage response. *PLoS Pathog.* 2009;5:e1000545.
- [64] Bloch K. Control mechanisms for fatty acid synthesis in *Mycobacterium smegmatis*. *Adv Enzymol Relat Areas Mol Biol.* 1977;45:1-84.
- [65] Fernandes ND, Kolattukudy PE. Cloning, sequencing and characterization of a fatty acid synthase-encoding gene from *Mycobacterium tuberculosis* var. bovis BCG. *Gene.* 1996;170:95-9.
- [66] Bloch K, Vance D. Control mechanisms in the synthesis of saturated fatty acids. *Annu Rev Biochem.* 1977;46:263-98.
- [67] Radmacher E, Alderwick LJ, Besra GS, Brown AK, Gibson KJ, Sahm H, et al. Two functional FAS-I type fatty acid synthases in *Corynebacterium glutamicum*. *Microbiology.* 2005;151:2421-7.
- [68] Odriozola JM, Ramos JA, Bloch K. Fatty acid synthetase activity in *Mycobacterium smegmatis*. Characterization of the acyl carrier protein-dependent elongating system. *Biochim Biophys Acta.* 1977;488:207-17.
- [69] Rock CO, Cronan JE. *Escherichia coli* as a model for the regulation of dissociable (type II) fatty acid biosynthesis. *Biochim Biophys Acta.* 1996;1302:1-16.
- [70] Kremer L, Nampoothiri KM, Lesjean S, Dover LG, Graham S, Betts J, et al. Biochemical characterization of acyl carrier protein (AcpM) and malonyl-CoA:AcpM transacylase (mtFabD), two major components of *Mycobacterium tuberculosis* fatty acid synthase II. *J Biol Chem.* 2001;276:27967-74.
- [71] Marrakchi H, Laneelle G, Quemard A. InhA, a target of the antituberculous drug isoniazid, is involved in a mycobacterial fatty acid elongation system, FAS-II. *Microbiology.* 2000;146:289-96.
- [72] Portevin D, De Sousa-D'Auria C, Houssin C, Grimaldi C, Chami M, Daffe M, et al. A polyketide synthase catalyzes the last condensation step of mycolic acid biosynthesis in mycobacteria and related organisms. *Proc Natl Acad Sci U S A.* 2004;101:314-9.

- [73] Lea-Smith DJ, Pyke JS, Tull D, McConville MJ, Coppel RL, Crellin PK. The reductase that catalyzes mycolic motif synthesis is required for efficient attachment of mycolic acids to arabinogalactan. *J Biol Chem.* 2007;282:11000-8.
- [74] Athenstaedt K, Daum G. Phosphatidic acid, a key intermediate in lipid metabolism. *Eur J Biochem.* 1999;266:1-16.
- [75] Okuyama H, Kameyama Y, Fujikawa M, Yamada K, Ikezawa H. Mechanism of diacylglycerophosphate synthesis in mycobacteria. *J Biol Chem.* 1977;252:6682-6.
- [76] Okuyama H, Kankura T, Nojima S. Positional distribution of fatty acids in phospholipids from mycobacteria. *J Biochem.* 1967;61:732-7.
- [77] Owens RM, Hsu FF, VanderVen BC, Purdy GE, Hestande E, Giannakas P, et al. *M. tuberculosis* Rv2252 encodes a diacylglycerol kinase involved in the biosynthesis of phosphatidylinositol mannosides (PIMs). *Mol Microbiol.* 2006;60:1152-63.
- [78] Smith SW, Weiss SB, Kennedy EP. The enzymatic dephosphorylation of phosphatidic acids. *J Biol Chem.* 1957;228:915-22.
- [79] Han GS, Wu WI, Carman GM. The *Saccharomyces cerevisiae* Lipin homolog is a Mg^{2+} -dependent phosphatidate phosphatase enzyme. *J Biol Chem.* 2006;281:9210-8.
- [80] Kalscheuer R, Steinbuchel A. A novel bifunctional wax ester synthase/acyl-CoA:diacylglycerol acyltransferase mediates wax ester and triacylglycerol biosynthesis in *Acinetobacter calcoaceticus* ADP1. *J Biol Chem.* 2003;278:8075-82.
- [81] Elamin AA, Stehr M, Spallek R, Rohde M, Singh M. The *Mycobacterium tuberculosis* Ag85A is a novel diacylglycerol acyltransferase involved in lipid body formation. *Mol Microbiol.* 2011;81:1577-92.
- [82] Ortalo-Magne A, Lemassu A, Laneelle MA, Bardou F, Silve G, Gounon P, et al. Identification of the surface-exposed lipids on the cell envelopes of *Mycobacterium tuberculosis* and other mycobacterial species. *J Bacteriol.* 1996;178:456-61.
- [83] Paznokas JL, Kaplan A. Purification and properties of a triacylglycerol lipase from *Mycobacterium phlei*. *Biochim Biophys Acta.* 1977;487:405-21.
- [84] Deb C, Daniel J, Sirakova TD, Abomoelak B, Dubey VS, Kolattukudy PE. A novel lipase belonging to the hormone-sensitive lipase family induced under starvation to utilize stored triacylglycerol in *Mycobacterium tuberculosis*. *J Biol Chem.* 2006;281:3866-75.
- [85] Mishra KC, de Chastellier C, Narayana Y, Bifani P, Brown AK, Besra GS, et al. Functional role of the PE domain and immunogenicity of the *Mycobacterium tuberculosis* triacylglycerol hydrolase LipY. *Infect Immun.* 2008;76:127-40.
- [86] Munoz-Elias EJ, McKinney JD. Carbon metabolism of intracellular bacteria. *Cell Microbiol.* 2006;8:10-22.
- [87] Cotes K, Dhouib R, Douchet I, Chahinian H, de Caro A, Carriere F, et al. Characterization of an exported monoglyceride lipase from *Mycobacterium tuberculosis* possibly involved in the metabolism of host cell membrane lipids. *Biochem J.* 2007;408:417-27.
- [88] Waltermann M, Steinbuchel A. Neutral lipid bodies in prokaryotes: recent insights into structure, formation, and relationship to eukaryotic lipid depots. *J Bacteriol.* 2005;187:3607-19.

- [89] Waltermann M, Hinz A, Robenek H, Troyer D, Reichelt R, Malkus U, et al. Mechanism of lipid-body formation in prokaryotes: how bacteria fatten up. *Mol Microbiol.* 2005;55:750-63.
- [90] Hanisch J, Waltermann M, Robenek H, Steinbuchel A. Eukaryotic lipid body proteins in oleogenous actinomycetes and their targeting to intracellular triacylglycerol inclusions: Impact on models of lipid body biogenesis. *Appl Environ Microbiol.* 2006;72:6743-50.
- [91] Hanisch J, Waltermann M, Robenek H, Steinbuchel A. The *Ralstonia eutropha* H16 phasin PhaP1 is targeted to intracellular triacylglycerol inclusions in *Rhodococcus opacus* PD630 and *Mycobacterium smegmatis* mc²155, and provides an anchor to target other proteins. *Microbiology.* 2006;152:3271-80.
- [92] Dowhan W. CDP-diacylglycerol synthase of microorganisms. *Biochim Biophys Acta.* 1997;1348:157-65.
- [93] Nigou J, Besra GS. Cytidine diphosphate-diacylglycerol synthesis in *Mycobacterium smegmatis*. *Biochem J.* 2002;367:157-62.
- [94] Lecocq J, Ballou CE. On the structure of cardiolipin. *Biochemistry.* 1964;3:976-80.
- [95] Icho T, Raetz CR. Multiple genes for membrane-bound phosphatases in *Escherichia coli* and their action on phospholipid precursors. *J Bacteriol.* 1983;153:722-30.
- [96] Funk CR, Zimniak L, Dowhan W. The *pgpA* and *pgpB* genes of *Escherichia coli* are not essential: evidence for a third phosphatidylglycerophosphate phosphatase. *J Bacteriol.* 1992;174:205-13.
- [97] Lu YH, Guan Z, Zhao J, Raetz CR. Three phosphatidylglycerol-phosphate phosphatases in the inner membrane of *Escherichia coli*. *J Biol Chem.* 2011;286:5506-18.
- [98] Osman C, Haag M, Wieland FT, Brugger B, Langer T. A mitochondrial phosphatase required for cardiolipin biosynthesis: the PGP phosphatase Gep4. *EMBO J.* 2010;29:1976-87.
- [99] Zhang J, Guan Z, Murphy AN, Wiley SE, Perkins GA, Worby CA, et al. Mitochondrial phosphatase PTPMT1 is essential for cardiolipin biosynthesis. *Cell Metab.* 2011;13:690-700.
- [100] Mathur AK, Murthy PS, Saharia GS, Venkitasubramanian TA. Studies on cardiolipin biosynthesis in *Mycobacterium smegmatis*. *Can J Microbiol.* 1976;22:354-8.
- [101] Ter Horst B, Seshadri C, Sweet L, Young DC, Feringa BL, Moody DB, et al. Asymmetric synthesis and structure elucidation of a glycerophospholipid from *Mycobacterium tuberculosis*. *J Lipid Res.* 2010;51:1017-22.
- [102] Morita YS, Velasquez R, Taig E, Waller RF, Patterson JH, Tull D, et al. Compartmentalization of lipid biosynthesis in mycobacteria. *J Biol Chem.* 2005;280:21645-52.
- [103] Salman M, Lonsdale JT, Besra GS, Brennan PJ. Phosphatidylinositol synthesis in mycobacteria. *Biochim Biophys Acta.* 1999;1436:437-50.
- [104] Morita YS, Fukuda T, Sena CB, Yamaryo-Botte Y, McConville MJ, Kinoshita T. Inositol lipid metabolism in mycobacteria: biosynthesis and regulatory mechanisms. *Biochim Biophys Acta.* 2011;1810:630-41.
- [105] Nigou J, Besra GS. Characterization and regulation of inositol monophosphatase activity in *Mycobacterium smegmatis*. *Biochem J.* 2002;361:385-90.

- [106] Mishra AK, Driessen NN, Appelmelk BJ, Besra GS. Lipoarabinomannan and related glycoconjugates: structure, biogenesis and role in *Mycobacterium tuberculosis* physiology and host-pathogen interaction. FEMS Microbiol Rev. 2011;35:1126-57.
- [107] Guerin ME, Kordulakova J, Alzari PM, Brennan PJ, Jackson M. Molecular basis of phosphatidyl-*myo*-inositol mannoside biosynthesis and regulation in mycobacteria. J Biol Chem. 2010;285:33577-83.
- [108] Ma Y, Stern RJ, Scherman MS, Vissa VD, Yan W, Jones VC, et al. Drug targeting *Mycobacterium tuberculosis* cell wall synthesis: genetics of dTDP-rhamnose synthetic enzymes and development of a microtiter plate-based screen for inhibitors of conversion of dTDP-glucose to dTDP-rhamnose. Antimicrob Agents Chemother. 2001;45:1407-16.
- [109] McCarthy TR, Torrelles JB, MacFarlane AS, Katawczik M, Kutzbach B, Desjardin LE, et al. Overexpression of *Mycobacterium tuberculosis* manB, a phosphomannomutase that increases phosphatidylinositol mannoside biosynthesis in *Mycobacterium smegmatis* and mycobacterial association with human macrophages. Mol Microbiol. 2005;58:774-90.
- [110] Ning B, Elbein AD. Purification and properties of mycobacterial GDP-mannose pyrophosphorylase. Arch Biochem Biophys. 1999;362:339-45.
- [111] Patterson JH, Waller RF, Jeevarajah D, Billman-Jacobe H, McConville MJ. Mannose metabolism is required for mycobacterial growth. Biochem J. 2003;372:77-86.
- [112] Mishra AK, Krumbach K, Rittmann D, Batt SM, Lee OY, De S, et al. Deletion of manC in *Corynebacterium glutamicum* results in a phospho-*myo*-inositol mannoside- and lipoglycan-deficient mutant. Microbiology. 2012;158:1908-17.
- [113] Kordulakova J, Gilleron M, Mikusova K, Puzo G, Brennan PJ, Gicquel B, et al. Definition of the first mannosylation step in phosphatidylinositol mannoside synthesis. PimA is essential for growth of mycobacteria. J Biol Chem. 2002;277:31335-44.
- [114] Gu X, Chen M, Wang Q, Zhang M, Wang B, Wang H. Expression and purification of a functionally active recombinant GDP-mannosyltransferase (PimA) from *Mycobacterium tuberculosis* H37Rv. Protein Expr Purif. 2005;42:47-53.
- [115] Guerin ME, Buschiazzi A, Kordulakova J, Jackson M, Alzari PM. Crystallization and preliminary crystallographic analysis of PimA, an essential mannosyltransferase from *Mycobacterium smegmatis*. Acta Crystallograph Sect F Struct Biol Cryst Commun. 2005;61:518-20.
- [116] Guerin ME, Kordulakova J, Schaeffer F, Svetlikova Z, Buschiazzi A, Giganti D, et al. Molecular recognition and interfacial catalysis by the essential phosphatidylinositol mannosyltransferase PimA from mycobacteria. J Biol Chem. 2007;282:20705-14.
- [117] Guerin ME, Kaur D, Somashekar BS, Gibbs S, Gest P, Chatterjee D, et al. New insights into the early steps of phosphatidylinositol mannoside biosynthesis in mycobacteria: PimB' is an essential enzyme of *Mycobacterium smegmatis*. J Biol Chem. 2009;284:25687-96.
- [118] Lea-Smith DJ, Martin KL, Pyke JS, Tull D, McConville MJ, Coppel RL, et al. Analysis of a new mannosyltransferase required for the synthesis of phosphatidylinositol

- mannosides and lipoarabinomannan reveals two lipomannan pools in *Corynebacterineae*. J Biol Chem. 2008;283:6773-82.
- [119] Schaeffer ML, Khoo KH, Besra GS, Chatterjee D, Brennan PJ, Belisle JT, et al. The *pimB* gene of *Mycobacterium tuberculosis* encodes a mannosyltransferase involved in lipoarabinomannan biosynthesis. J Biol Chem. 1999;274:31625-31.
- [120] Mishra AK, Batt S, Krumbach K, Eggeling L, Besra GS. Characterization of the *Corynebacterium glutamicum* *deltapimB'* *deltamtgA* double deletion mutant and the role of *Mycobacterium tuberculosis* orthologues Rv2188c and Rv0557 in glycolipid biosynthesis. J Bacteriol. 2009;191:4465-72.
- [121] Tatituri RV, Illarionov PA, Dover LG, Nigou J, Gilleron M, Hitchen P, et al. Inactivation of *Corynebacterium glutamicum* NCgl0452 and the role of MgtA in the biosynthesis of a novel mannosylated glycolipid involved in lipomannan biosynthesis. J Biol Chem. 2007;282:4561-72.
- [122] Batt SM, Jabeen T, Mishra AK, Veerapen N, Krumbach K, Eggeling L, et al. Acceptor substrate discrimination in phosphatidyl-myo-inositol mannoside synthesis: structural and mutational analysis of mannosyltransferase *Corynebacterium glutamicum* PimB'. J Biol Chem. 2010;285:37741-52.
- [123] Kordulakova J, Gilleron M, Puzo G, Brennan PJ, Gicquel B, Mikusova K, et al. Identification of the required acyltransferase step in the biosynthesis of the phosphatidylinositol mannosides of *Mycobacterium* species. J Biol Chem. 2003;278:36285-95.
- [124] Kremer L, Gurucha SS, Bifani P, Hitchen PG, Baulard A, Morris HR, et al. Characterization of a putative alpha-mannosyltransferase involved in phosphatidylinositol trimannoside biosynthesis in *Mycobacterium tuberculosis*. Biochem J. 2002;363:437-47.
- [125] Besra GS, Morehouse CB, Rittner CM, Waechter CJ, Brennan PJ. Biosynthesis of mycobacterial lipoarabinomannan. J Biol Chem. 1997;272:18460-6.
- [126] Gurucha SS, Baulard AR, Kremer L, Loch C, Moody DB, Muhlecker W, et al. Ppm1, a novel polyprenol monophosphomannose synthase from *Mycobacterium tuberculosis*. Biochem J. 2002;365:441-50.
- [127] Gibson KJ, Eggeling L, Maughan WN, Krumbach K, Gurucha SS, Nigou J, et al. Disruption of Cg-Ppm1, a polyprenyl monophosphomannose synthase, and the generation of lipoglycan-less mutants in *Corynebacterium glutamicum*. J Biol Chem. 2003;278:40842-50.
- [128] Morita YS, Patterson JH, Billman-Jacobe H, McConville MJ. Biosynthesis of mycobacterial phosphatidylinositol mannosides. Biochem J. 2004;378:589-97.
- [129] Kovacevic S, Anderson D, Morita YS, Patterson J, Haites R, McMillan BN, et al. Identification of a novel protein with a role in lipoarabinomannan biosynthesis in mycobacteria. J Biol Chem. 2006;281:9011-7.
- [130] Crellin PK, Kovacevic S, Martin KL, Brammananth R, Morita YS, Billman-Jacobe H, et al. Mutations in *pimE* restore lipoarabinomannan synthesis and growth in a *Mycobacterium smegmatis* *lpqW* mutant. J Bacteriol. 2008;190:3690-9.

- [131] Marland Z, Beddoe T, Zaker-Tabrizi L, Lucet IS, Brammananth R, Whisstock JC, et al. Hijacking of a substrate-binding protein scaffold for use in mycobacterial cell wall biosynthesis. *J Mol Biol.* 2006;359:983-97.
- [132] Besra GS, Brennan PJ. The mycobacterial cell wall: biosynthesis of arabinogalactan and lipoarabinomannan. *Biochem Soc Trans.* 1997;25:845-50.
- [133] Mishra AK, Alderwick LJ, Rittmann D, Wang C, Bhatt A, Jacobs WR, Jr., et al. Identification of a novel $\alpha(1\rightarrow6)$ mannopyranosyltransferase MptB from *Corynebacterium glutamicum* by deletion of a conserved gene, *NCgl1505*, affords a lipomannan- and lipoarabinomannan-deficient mutant. *Mol Microbiol.* 2008;68:1595-613.
- [134] Kaur D, Berg S, Dinadayala P, Gicquel B, Chatterjee D, McNeil MR, et al. Biosynthesis of mycobacterial lipoarabinomannan: role of a branching mannosyltransferase. *Proc Natl Acad Sci U S A.* 2006;103:13664-9.
- [135] Kaur D, McNeil MR, Khoo KH, Chatterjee D, Crick DC, Jackson M, et al. New insights into the biosynthesis of mycobacterial lipomannan arising from deletion of a conserved gene. *J Biol Chem.* 2007;282:27133-40.
- [136] Mishra AK, Alderwick LJ, Rittmann D, Tatituri RV, Nigou J, Gilleron M, et al. Identification of an $\alpha(1\rightarrow6)$ mannopyranosyltransferase (MptA), involved in *Corynebacterium glutamicum* lipomannan biosynthesis, and identification of its orthologue in *Mycobacterium tuberculosis*. *Mol Microbiol.* 2007;65:1503-17.
- [137] Kaur D, Obregon-Henao A, Pham H, Chatterjee D, Brennan PJ, Jackson M. Lipoarabinomannan of *Mycobacterium*: mannose capping by a multifunctional terminal mannosyltransferase. *Proc Natl Acad Sci U S A.* 2008;105:17973-7.
- [138] Sena CB, Fukuda T, Miyanagi K, Matsumoto S, Kobayashi K, Murakami Y, et al. Controlled expression of branch-forming mannosyltransferase is critical for mycobacterial lipoarabinomannan biosynthesis. *J Biol Chem.* 2010;285:13326-36.
- [139] Birch HL, Alderwick LJ, Bhatt A, Rittmann D, Krumbach K, Singh A, et al. Biosynthesis of mycobacterial arabinogalactan: identification of a novel $\alpha(1\rightarrow3)$ arabinofuranosyltransferase. *Mol Microbiol.* 2008;69:1191-206.
- [140] Alderwick LJ, Lloyd GS, Ghadbane H, May JW, Bhatt A, Eggeling L, et al. The C-terminal domain of the arabinosyltransferase *Mycobacterium tuberculosis* EmbC is a lectin-like carbohydrate binding module. *PLoS Pathog.* 2011;7:e1001299.
- [141] Skovierova H, Larrouy-Maumus G, Zhang J, Kaur D, Barilone N, Kordulakova J, et al. AftD, a novel essential arabinofuranosyltransferase from mycobacteria. *Glycobiology.* 2009;19:1235-47.
- [142] Zhang N, Torrelles JB, McNeil MR, Escuyer VE, Khoo KH, Brennan PJ, et al. The Emb proteins of mycobacteria direct arabinosylation of lipoarabinomannan and arabinogalactan via an N-terminal recognition region and a C-terminal synthetic region. *Mol Microbiol.* 2003;50:69-76.
- [143] Chatterjee D, Hunter SW, McNeil M, Brennan PJ. Lipoarabinomannan. Multiglycosylated form of the mycobacterial mannosylphosphatidylinositols. *J Biol Chem.* 1992;267:6228-33.

- [144] Appelmelk BJ, den Dunnen J, Driessen NN, Ummels R, Pak M, Nigou J, et al. The mannose cap of mycobacterial lipoarabinomannan does not dominate the *Mycobacterium*-host interaction. *Cell Microbiol.* 2008;10:930-44.
- [145] Khoo KH, Dell A, Morris HR, Brennan PJ, Chatterjee D. Inositol phosphate capping of the nonreducing termini of lipoarabinomannan from rapidly growing strains of *Mycobacterium*. *J Biol Chem.* 1995;270:12380-9.
- [146] Mishra AK, Klein C, Gurcha SS, Alderwick LJ, Babu P, Hitchen PG, et al. Structural characterization and functional properties of a novel lipomannan variant isolated from a *Corynebacterium glutamicum* pimB' mutant. *Antonie Van Leeuwenhoek.* 2008;94:277-87.
- [147] Tatituri RV, Alderwick LJ, Mishra AK, Nigou J, Gilleron M, Krumbach K, et al. Structural characterization of a partially arabinosylated lipoarabinomannan variant isolated from a *Corynebacterium glutamicum* ubiA mutant. *Microbiology.* 2007;153:2621-9.
- [148] Wolucka BA, McNeil MR, de Hoffmann E, Chojnacki T, Brennan PJ. Recognition of the lipid intermediate for arabinogalactan/arabinomannan biosynthesis and its relation to the mode of action of ethambutol on mycobacteria. *J Biol Chem.* 1994;269:23328-35.
- [149] Takayama K, Schnoes HK, Semmler EJ. Characterization of the alkali-stable mannophospholipids of *Mycobacterium smegmatis*. *Biochim Biophys Acta.* 1973;316:212-21.
- [150] Bailey AM, Mahapatra S, Brennan PJ, Crick DC. Identification, cloning, purification, and enzymatic characterization of *Mycobacterium tuberculosis* 1-deoxy-D-xylulose 5-phosphate synthase. *Glycobiology.* 2002;12:813-20.
- [151] Dhiman RK, Schaeffer ML, Bailey AM, Testa CA, Scherman H, Crick DC. 1- Deoxy-D-xylulose 5-phosphate reductoisomerase (IspC) from *Mycobacterium tuberculosis*: towards understanding mycobacterial resistance to fosmidomycin. *J Bacteriol.* 2005;187:8395-402.
- [152] Argyrou A, Blanchard JS. Kinetic and chemical mechanism of *Mycobacterium tuberculosis* 1-deoxy-deoxy-D-xylulose -5-phosphate isomeroreductase. *Biochemistry.* 2004;43:4375-84.
- [153] Takayama K, Goldman DS. Enzymatic synthesis of mannosyl-1-phosphoryl-decaprenol by a cell-free system of *Mycobacterium tuberculosis*. *J Biol Chem.* 1970;245:6251-7.
- [154] Wolucka BA, de Hoffmann E. The presence of beta-D-ribosyl-1-monophosphodecaprenol in mycobacteria. *J Biol Chem.* 1995;270:20151-5.
- [155] Wolucka BA, de Hoffmann E. Isolation and characterization of the major form of polyprenyl-phospho-mannose from *Mycobacterium smegmatis*. *Glycobiology.* 1998;8:955-62.
- [156] Baulard AR, Gurcha SS, Engohang-Ndong J, Gouffi K, Locht C, Besra GS. *In vivo* interaction between the polyprenol phosphate mannose synthase Ppm1 and the integral membrane protein Ppm2 from *Mycobacterium smegmatis* revealed by a bacterial two-hybrid system. *J Biol Chem.* 2003;278:2242-8.

- [157] Scherman H, Kaur D, Pham H, Skovierova H, Jackson M, Brennan PJ. Identification of a polyprenylphosphomannosyl synthase involved in the synthesis of mycobacterial mannosides. *J Bacteriol.* 2009;191:6769-72.
- [158] Skovierova H, Larrouy-Maumus G, Pham H, Belanova M, Barilone N, Dasgupta A, et al. Biosynthetic origin of the galactosamine substituent of arabinogalactan in *Mycobacterium tuberculosis*. *J Biol Chem.* 2010;285:41348-55.
- [159] Klutts JS, Hatanaka K, Pan YT, Elbein AD. Biosynthesis of D-arabinose in *Mycobacterium smegmatis*: specific labeling from D-glucose. *Arch Biochem Biophys.* 2002;398:229-39.
- [160] Scherman M, Weston A, Duncan K, Whittington A, Upton R, Deng L, et al. Biosynthetic origin of mycobacterial cell wall arabinosyl residues. *J Bacteriol.* 1995;177:7125-30.
- [161] Scherman MS, Kalbe-Bournonville L, Bush D, Xin Y, Deng L, McNeil M. Polyprenylphosphate-pentoses in mycobacteria are synthesized from 5-phosphoribose pyrophosphate. *J Biol Chem.* 1996;271:29652-8.
- [162] Huang H, Scherman MS, D'Haeze W, Vereecke D, Holsters M, Crick DC, et al. Identification and active expression of the *Mycobacterium tuberculosis* gene encoding 5-phospho-D-ribose-1-diphosphate: decaprenyl-phosphate 5-phosphoribosyltransferase, the first enzyme committed to decaprenylphosphoryl-d-arabinose synthesis. *J Biol Chem.* 2005;280:24539-43.
- [163] Mikusova K, Huang H, Yagi T, Holsters M, Vereecke D, D'Haeze W, et al. Decaprenylphosphoryl arabinofuranose, the donor of the D-arabinofuranosyl residues of mycobacterial arabinan, is formed via a two-step epimerization of decaprenylphosphoryl ribose. *J Bacteriol.* 2005;187:8020-5.
- [164] Christophe T, Jackson M, Jeon HK, Fenistein D, Contreras-Dominguez M, Kim J, et al. High content screening identifies decaprenyl-phosphoribose 2' epimerase as a target for intracellular antimycobacterial inhibitors. *PLoS Pathog.* 2009;5:e1000645.
- [165] Makarov V, Manina G, Mikusova K, Mollmann U, Ryabova O, Saint-Joanis B, et al. Benzothiazinones kill *Mycobacterium tuberculosis* by blocking arabinan synthesis. *Science.* 2009;324:801-4.
- [166] Makarov V, Riabova OB, Yuschenko A, Urlyapova N, Daudova A, Zipfel PF, et al. Synthesis and antileprosy activity of some dialkyldithiocarbamates. *J Antimicrob Chemother.* 2006;57:1134-8.
- [167] Pasca MR, Degiacomi G, Ribeiro AL, Zara F, De Mori P, Heym B, et al. Clinical isolates of *Mycobacterium tuberculosis* in four European hospitals are uniformly susceptible to benzothiazinones. *Antimicrob Agents Chemother.* 2010;54:1616-8.
- [168] Crellin PK, Brammananth R, Coppel RL. Decaprenylphosphoryl-beta- D-ribose 2'-epimerase, the target of benzothiazinones and dinitrobenzamides, is an essential enzyme in *Mycobacterium smegmatis*. *PLoS One.* 2011;6:e16869.
- [169] Manina G, Pasca MR, Buroni S, De Rossi E, Riccardi G. Decaprenylphosphoryl-beta- D-ribose 2'-epimerase from *Mycobacterium tuberculosis* is a magic drug target. *Curr Med Chem.* 2010;17:3099-108.

- [170] Batt SM, Jabeen T, Bhowruth V, Quill L, Lund PA, Eggeling L, et al. Structural basis of inhibition of *Mycobacterium tuberculosis* DprE1 by benzothiazinone inhibitors. *Proc Natl Acad Sci U S A*. 2012;109:11354-9.
- [171] Tarnok I, Tarnok Z. Carotenes and xanthophylls in mycobacteria. II. Lycopene, alpha- and beta-carotene and xanthophyll in mycobacterial pigments. *Tubercle*. 1971;52:127-35.
- [172] Houssaini-Iraqi M, Lazraq MH, Clavel-Seres S, Rastogi N, David HL. Cloning and expression of *Mycobacterium aurum* carotenogenesis genes in *Mycobacterium smegmatis*. *FEMS Microbiol Lett*. 1992;69:239-44.
- [173] Ramakrishnan L, Tran HT, Federspiel NA, Falkow S. A crtB homolog essential for photochromogenicity in *Mycobacterium marinum*: isolation, characterization, and gene disruption via homologous recombination. *J Bacteriol*. 1997;179:5862-8.
- [174] Viveiros M, Krubasik P, Sandmann G, Houssaini-Iraqi M. Structural and functional analysis of the gene cluster encoding carotenoid biosynthesis in *Mycobacterium aurum* A+. *FEMS Microbiol Lett*. 2000;187:95-101.
- [175] Robledo JA, Murillo AM, Rouzaud F. Physiological role and potential clinical interest of mycobacterial pigments. *IUBMB Life*. 2011;63:71-8.
- [176] Kim SW, Keasling JD. Metabolic engineering of the nonmevalonate isopentenyl diphosphate synthesis pathway in *Escherichia coli* enhances lycopene production. *Biotechnol Bioeng*. 2001;72:408-15.
- [177] Ito M, Yamano Y, Tode C, Wada A. Carotenoid synthesis: retrospect and recent progress. *Arch Biochem Biophys*. 2009;483:224-8.
- [178] Misawa N, Satomi Y, Kondo K, Yokoyama A, Kajiwara S, Saito T, et al. Structure and functional analysis of a marine bacterial carotenoid biosynthesis gene cluster and astaxanthin biosynthetic pathway proposed at the gene level. *J Bacteriol*. 1995;177:6575-84.
- [179] Wenk MR. The emerging field of lipidomics. *Nat Rev Drug Discov*. 2005;4:594-610.
- [180] Wenk MR. Lipidomics: new tools and applications. *Cell*. 2010;143:888-95.
- [181] Layre E, Sweet L, Hong S, Madigan CA, Desjardins D, Young DC, et al. A comparative lipidomics platform for chemotaxonomic analysis of *Mycobacterium tuberculosis*. *Chem Biol*. 2011;18:1537-49.
- [182] Sartain MJ, Dick DL, Rithner CD, Crick DC, Belisle JT. Lipidomic analyses of *Mycobacterium tuberculosis* based on accurate mass measurements and the novel "Mtb LipidDB". *J Lipid Res*. 2011;52:861-72.
- [183] Madigan CA, Cheng TY, Layre E, Young DC, McConnell MJ, Debono CA, et al. Lipidomic discovery of deoxysiderophores reveals a revised mycobactin biosynthesis pathway in *Mycobacterium tuberculosis*. *Proc Natl Acad Sci U S A*. 2012;109:1257-62.