We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Non-Coding RNAs and Cancer

Gianpiero Di Leva and Michela Garofalo

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/54997>

1. Introduction

The question of which regions of the human genome constitute its functional elements—those expressed as genes or serving as regulatory elements—has long been a central topic in biology. In the 1970s and 1980s, early cloning-based methods revealed the presence of more than 7000 genes in human genome [1], and large-scale analyses of expressed sequence tags (ESTs) in the 1990s suggested that the estimated number of human genes range from 35,000 to 100,000 [2]. The completion of the human genome project narrowed the focus considerably by highlighting the surprisingly small number of protein-coding genes, which is now conventionally cited as less than 25,000 [3]. While the number of protein-coding genes (20,000–25,000) has maintained broad consensus, recent studies of the human transcriptome have revealed an astounding number of non-coding RNAs (ncRNAs) [4–6]. In fact, the increased sensitivity of genome tiling arrays provides an even more detailed view, revealing that the extent of non-coding sequence transcription is at least four times greater than coding sequence, and that the abundance of non-coding transcripts had been previously overlooked. The RNA world hypothesis proposes that early life was based on RNAs, which subsequently devolved the storage of information to more stable DNA, and catalytic functions to more versatile proteins. Consequently, despite crucial roles in the ancient processes of translation and splicing, RNA is assumed to have been largely relegated to an intermediate between gene and protein, encapsulated in the central dogma ‘DNA makes RNA makes protein’ [7]. However, the finding that most of the genome in complex organisms is transcribed and the discovery of new classes of regulatory non-coding RNAs (ncRNAs) challenges this assumption and suggests that RNAs have continued to evolve and expand alongside proteins and DNA.

ncRNAs are considered as RNA transcripts that do not encode for a protein. In the past decade, a great diversity of ncRNAs has been observed. Depending on the type of ncRNA, transcription can occur by any of the three RNA polymerases (RNA Pol I, RNA Pol II, or RNA Pol III). General conventions divide ncRNAs into two main categories: small ncRNAs

less than 200 bp and long ncRNAs greater than 200 bps [8]. Within these two categories, there are also many individual classes of ncRNAs (**Table1**), although the degree of biological and experimental support for each class ranges substantially and should be evaluated individually. The relevance of ncRNAs in gene regulation has been rapidly unveiling during the last decade. However, the functional elements in the primary sequence of noncoding genes that determine their role as RNA molecules remain unknown. Protein-coding genes have a defined language with a set of grammatical rules: three nucleotides forms a codon that translates into a specific amino acid [9]. Aberrations in codons of a protein-coding gene can be interpreted in terms of the amino acids they encode. We can recognize a mutation in a codon and determine its contribution to a given disease. In contrast to the genetic code for protein synthesis, ‘the ncRNA alphabet’ – a specific set of RNA sequences or structural motifs important for ncRNA function – remains to be largely elucidated. However, it has become increasingly apparent that the ncRNAs are of crucial functional importance for normal development, physiology and disease [10]. The functional relevance of the ncRNAs is particularly evident for a class of small non-coding RNAs called microRNAs (miRNAs) [11-12]. In human diseases, particularly cancer, it has been shown that epigenetic and genetic defects in miRNAs and their processing machinery are a common hallmark of disease [13-16]. However, miRNAs are just the tip of the iceberg, and other ncRNAs such as small nucleolar RNAs (snoRNAs), PIWI-interacting RNAs (piRNAs), large intergenic non-coding RNAs (lincRNAs) and, overall, the heterogeneous group of long non-coding RNAs (lncRNAs), might also contribute to the development of many different human disorders. Here we discuss the most recent genetic studies on ncRNAs and their related proteins in the context of cancer and we will analyze the new regulatory elements of the noncoding language to interpret their contribution to the pathogenesis of cancer.

2. MicroRNAs

In 1993, Victor Ambros and colleagues discovered a gene, *lin-4*, that affected development in *Caenorhabditis elegans* and found that its product was a small nonprotein-coding RNA [31]. The number of known small RNAs in different organisms such as *Caenorhabditis elegans*, *Drosophila melanogaster*, plants, and mammals—including humans—has since expanded substantially, mainly as a result of the cloning and sequencing of size-fractionated RNAs. MiRNAs are single stranded RNAs (ssRNAs) of 19–25 nucleotides in length that are generated from endogenous hairpin transcripts [32]. They play an important role in the negative regulation of gene expression by base-pairing to partially complementary sites on the target messenger RNAs (mRNAs), usually in the 3′ untranslated region (UTR). Binding of a miRNA to the target mRNA typically leads to translational repression and exonucleolytic mRNA decay, although highly complementary targets can be cleaved endonucleolytically. A genomic analysis of miRNAs has revealed that more than 50% of mammalian miRNAs are located within the intronic regions of annotated protein-coding or non-protein-coding genes [33]. These miRNAs could therefore use their host gene transcripts as carriers, although it remains possible that some are actually transcribed separately from internal promoters. Other miRNAs, located in intergenic regions, apparently have their own transcriptional regulatory elements and thus constitute

Category	Name	Supporting data	Function	Role in cancer	Refs.
Housekeeping RNAs	Ribosomal RNAs (rRNA)	high	ribosome structure	no	17-18
	Transfer RNAs (tRNA)	high	protein translation	no	17-18
	Small nuclear RNAs (snRNA)	high	splicing	no	17-18
	Small nucleolar RNAs (snoRNA)	high	post-translational modification	yes	17-18
Short non coding RNAs (above 200nt in size)	MicroRNAs	high	translational repression	yes	20,21
	Tiny transcription initiation RNAs	high	may regulate gene expression	not known	18
	Repeat associated small interfering RNAs	high	gene regulation, transposon control and viral defence	not known	18
	Promoter-associated short RNAs	high	may regulate gene expression at chromatin level	not known	18, 22,24
	Termini-associated short RNAs	high	may regulate gene expression at chromatin level	not known	18, 22,24
	Antisense termini associated short	high	may regulate gene expression at chromatin level	not known	17, 22,24
	Piwi-interacting RNAs	high	regulate transposon activity and chromatin state	yes	23
	Transcription start site antisense RNAs	moderate	may regulate transcription	not known	17
	Retrotranspon-derived RNAs	high	may regulate transcription	not known	24
	3'UTR-derived RNAs	moderate	may regulate transcription	not known	17
	Splice-site RNAs	poor	not known	not known	18
Long non coding RNAs (over 200nt in size)	Long or large intergenic ncRNAs	high	epigenetic regulation, protein complex subcellular compartments or localization	yes	25,26
	Transcribed ultraconserved regions	high	not known	yes	27
	Pseudogenes	high	competitive endogenous RNAs	yes	25, 28
	Enhancer RNAs	high	not known	yes	29
	Long intronic ncRNAs	moderate	not known	not known	17,18
	Repeat associated ncRNAs	high	not known	not known	23
	Antisense RNAs	high	gene expression	not known	28
	Promoter associated long RNAs	moderate	may regulate gene expression at chromatin level	not known	22, 30
	Long stress-induced non-coding transcripts	moderate	epigenetic regulation, protein complex subcellular compartments or localization	yes	17, 18

Table 1. Non coding RNA in human genome.

independent transcription units. Animal miRNAs are processed from longer primary transcripts (pri-miRNAs) that can contain multiple miRNAs [34,35]. Few pri-miRNA transcripts have been studied in detail, but in general miRNAs are regulated and transcribed similar to protein encoding genes by (Pol) II with the exception of the rapidly evolving RNA polymerase (Pol) III transcribed miRNA cluster [36]. MiRNA processing occurs in three essential steps (**Figure 1**). First, the nuclear endoribonuclease protein Drosha recognizes the miRNA hairpins in the primary transcript and cleaves each hairpin ~11 nt from its base [37-38]. It has been proposed that Drosha may recognize the pri-miRNA through the stem-loop structure and then cleave the stem at a fixed distance from the loop to liberate the pre-miRNA. How is the Drosha enzyme able to discriminate the pri-miRNA stem-loop structure from the other stem-loop cellular RNAs? Both cell culture experiments and in vitro Drosha cleavage assays have shown that proteins associated with Drosha confer specificity to this process. In fact, Drosha has been found to be part of a large, ~650-kDa protein complex known as the Microprocessor [39], where Drosha interacts with its cofactor DGCR8 (the DiGeorge syndrome critical region gene 8 protein) in the human and interacts with Pasha in *Drosophila melanogaster* [40]. The next step in miRNA biogenesis is recognition of the ~60 nt pre-miRNA by exportin-5 and export into the cytoplasm in a ran-guanine-GTP-dependent manner [41-43]. The Exp5/Ran-GTP complex has a high affinity for pre-miRNAs,

Figure 1. miRNA biogenesis and function. The primary miRNA (pri-miRNA) is transcribed by RNA pol II from its genomic location and cleaved by the microprocessor complex, which comprises Drosha and DGCR8. The resulting pre-miRNA is actively transported to the cytoplasm by exportin 5 (Expt.5), where the pre-miRNA undergoes further processing into the mature miRNA by Dicer and its co-factors, protein activator of interferon-induced protein kinase (PACT) and TAR RNA binding protein (TRBP). Normally, one strand of this duplex is degraded (miRNA star), whereas the other strand accumulates as a mature miRNA. From the miRNA-miRNA duplex, only the miRNA enters preferentially in the protein effector complex, formed by the RNA-induced silencing complex (RISC) and miRgonaute and binds with partial complementarity to the 3' untranslated region (UTR) of target messenger RNAs (mRNAs) to mediate translational repression.

protecting them from the moment they are generated in the nucleus until they are ready for the next cleavage step in the cytoplasm, where GTP is hydrolyzed to guanosine diphosphate (GDP); at that point, the Exp5/Ran-GDP complex releases its cargo. Third, the endoribonuclease protein Dicer cleaves the pre-miRNA into ~22 nt duplexes and, with the help of cofactors such as TAR RNA binding protein (TRBP) and protein activator of the interferon-induced protein kinase (PACT), preferentially incorporates one of the duplex strands into the RNA induced-silencing complex (RISC) [44-50]. The final product is a miRNA-miRNA duplex that needs to be unwound to act as a single-stranded guide in the RISC to recognize its target mRNAs. It was originally proposed that an ATP-dependent helicase (known as unwindase) separates the two small RNA strands, after which the resulting single-stranded guide is loaded into Ago proteins. However, it was later shown that *Drosophila* Ago2 [51], as well as human Ago2 [52], directly receive double-stranded small RNA from the RISC-loading complex. Ago2 then cleaves the passenger strand, thereby liberating the single-stranded guide to form mature Ago2-RISC. In mammals, miRNAs guide the RISC to complementary target sites in mRNAs, where endonucleolytically active Ago proteins cleave the RNA [53] (**Figure 1**). Finally, RISC can cleave [54-55] degrade [56-57] or suppress translation [58-59] of target mRNAs depending on the complementarity between miRNA and mRNA. Imperfect base pairing between small RNAs and their target mRNAs leads to repression of translation and/or deadenylation (removal of the polyA tail of the target), followed by destabilization of the target [60], whereas perfect base pairing usually leads to mRNA degradation.

3. MicroRNAs and cancer

Cancer is a multistep process in which normal cells experience genetic changes that progress them through a series of pre-malignant states (initiation) into invasive cancer (progression) that can spread throughout the body (metastasis). The dysregulation of genes involved in cell proliferation, differentiation and/or apoptosis is associated with cancer initiation and progression. Genes linked with cancer development are characterized as oncogenes and tumor suppressors. Recently, the definition of oncogenes and tumor suppressors has been expanded from the classical protein coding genes to include miRNAs [61-62]. MiRNAs have been found to regulate more than 60% of mRNAs and have roles in fundamental processes, such as development [63], differentiation [64], cell proliferation [65], apoptosis [66], and stress responses [67]. Over the past few years, many miRNAs have been implicated in various human cancers. The first evidence that miRNAs are involved in cancer comes from the finding that miR-15 and miR-16 are downregulated or deleted in most patients with chronic lymphocytic leukemia [68]. This discovery has projected miRNAs to the center stage of molecular oncology and, in the past few years, a myriad of genome-wide miRNA expression profiling analyses have shown a general dysregulation of miRNA expression in all tumors (**Table 2**) [69]. Surprisingly, the use of miRNA profiles is newly becoming highly preferred to the traditional mRNA signature for a variety of reasons. First, the remarkable stability of miRNAs, due to their short length, has allowed scientists to perform analyses also in samples considered to be technically challenging, such as formalin fixed specimens. High sensitive and refined miRNA detection technique provide high reliability in the use of miRNAs as a diagnostic tools. Finally, miRNA fingerprints have demonstrated the ability to

identify the tissue of origin for cancer that have already spread in multiple metastatic sites, thereby reducing patient's psychological burden and overall procedure costs. To date, over 1000 miRNAs have been reported in humans (miRbase: 1527 at November 2011), and both loss and gain of miRNA functions contribute to cancer development through a range of different mechanisms that we will discuss in the following sections.

Cancer	Authors	Samples	Main findings	miRNA signature
Breast Cancer	Iorio et al. 2005	10 normal 76 tumors	First miRNA signature of breast cancer. 15 miRNAs predict the nature of the sample analyzed with 100% accuracy.	miR-21 , -155 were up-regulated while miR-10b , -125b , -145 were down-regulated.
	Mattie et al. 2006	20 tumors	Unique sets of miRNAs are associated with ErbB2 and ER/PR status.	ErbB2 status (let-7f , let-7g , miR-107 , -10b , -126 , -154 , -195) ER/PR status (miR-142-5p , -200a , -205 , -25).
	Blenkiron et al. 2007	93 tumors	Striking differences in miRNA expression between breast tumor molecular subtypes (luminal A, luminal B, basal-like, HER2+ and normal breast-like).	-
	Sempere et al. 2007	> 100 pairs tumor-normal	In situ hybridization method to reveal the spatial distribution of miRNA expression in archived formalin-fixed breast tumors	20 miRNAs differentiate matched normal/tumor. miR-21 up-regulated in all tumor specimens, miR-145 and miR-451 were the only miRs high expressed in normal specimens and low expressed in tumor specimens. let-7a , miR-21 , -141 , -214 preferentially expressed in luminal cells, miR-145 and -205 in myoepithelial cells.
	Foekens et al. 2008	299 tumors	Identification of miRNAs associated with metastatic capability.	miR-7 , -128a , -210 , -516-3p associated with tumor aggressiveness in ER-positive patients with lymph node-negative disease. miR-210 associated with early relapse in ER-negative patients with lymph node-negative disease and with poor outcome.
	Volinia et al. 2012	80 IDC 8 DCIS 6 normal	Identification of miRNA dysregulation during the transition from ductal carcinoma in situ to invasive ductal carcinoma	Let-7d , miR-210 , and -221 down-regulated in DCIS and up-regulated in IDC. miR-210 , -21 , -106b* , -197 , and let-7i associated with prognosis. Only miR-210 involved in the invasive transition.
Lung Cancer	Yanaihara et al. 2006	104 pairs tumor-normal	miRNA expression profiles are diagnostic and prognostic markers of lung cancer.	43 miRNAs discriminate between normal ad tumors. let-7a-2 , miR-155 , -17-3p , -145 , -21 were associated with adenocarcinoma patients survival.
	Raponi et al. 2009	10 normal 61 squamous	miRNAs may have greater clinical utility in predicting the prognosis of patients with squamous cell lung carcinomas than mRNA-based signatures.	15 miRNAs that were differentially expressed including miR-17/92 cluster, miR-155 , let-7 . miR-146b has the strongest prediction accuracy for stratifying prognostic groups.
	Landi et al. 2010	290 tumors 165AC-125S Q	miRNA expression profiles can distinguish adenocarcinoma (AC) from squamous carcinoma (SQ) and predict survival.	let-7g and miR-26 were the most down-regulated in SQ versus AC. miR-25 , -34c-5p , -34a , -191 and let-7e strongly predicted SQ survival for the 107 male smokers with early-stage SQ tumors. miR-21 is overexpressed in AC and may be a marker of tumor progression in adenocarcinoma.
	Tan et al. 2011	34 pairs tumors-normal	miRNA expression profiles of squamous carcinoma vs normal tissues.	miR-210 , -182 , -486-5p , -30a , -140-3p discriminate between cancerous and normal lung tissues. miR-31 was associated with poor survival in squamous cell carcinoma.
	Lu et al. 2012	527 tumors	Identification of miRNA signatures that predict prognosis of stage I NSCLC.	Two miRNA signatures that are highly predictive of recurrence/relapse free survival were identified. The first contained 34 miRNAs derived from 357 stage I NSCLC patients independent of cancer subtype, whereas the second containing 27 miRNAs was adenocarcinoma specific.
	Guan et al. 2012	14 different dataset	A meta-analysis reviews 14 microRNA expression profiling studies that compared the microRNAs expression profiles in lung cancer tissues with those in normal lung tissues.	184 miRNAs discriminate lung cancer tissues from normal tissues; 61 miRNAs were reported in at least two studies. Up-regulated miRNA: miR-210 was reported in 9 studies and miR-21 in 7 studies. Down-regulated miRNA: miR-126 was reported in 10 studies and miR-30a in 8 studies.
Thyroid Cancer	Pallante et al. 2006	10 normals 30 tumors	Identification of the miRNA expression profile of papillary carcinomas.	A significant increase in miR-221 , -222 and -181b was detected in papillary carcinomas in comparison with normal thyroid tissue.
	Visone et al. 2007	10 normals 76 tumors	Identification of the miRNA expression profile of anaplastic carcinomas from normal thyroid tissues.	A significant decrease in miR-30d , -125b , -26a , -30a-5p was detected in anaplastic carcinomas compared to normal thyroid tissue.
	Nikiforova et al. 2008	60 normals 60 tumors	Identification of miRNA signature for the different thyroid tumors subtypes: Oncocytic, conventional follicular, papillary and medullary carcinomas.	miR-187 , -221 , -222 , -146b , -155 , -224 , -197 were the most differentially overexpressed in thyroid tumors vs. hyperplastic nodules and combination of them classified the different subtypes.
	Yip et al. 2011	32 tumors	Identification of a specific signature in aggressive (17) compared with nonaggressive papillary carcinomas (15).	Upregulation of miR-146b , -221 , -222 , -155 , -31 and downregulation of miR-1 , -34b , -130b , -138 downregulation. miR-146 overexpression was associated with aggressive behavior in BRAF-positive tumors.
	Kitano et al. 2011	47 tumors	miRNAs expression signature for samples representing difficult to diagnose histologic subtypes of thyroid neoplasm (21 benign, 26 malignant).	Out of 34 differentially expressed miRNA, miR-126 and miR-7 had high diagnostic accuracy and could be helpful in the classification of benign and malignant thyroid tumors.

Cancer	Authors	Samples	Main findings	miRNA signature
Colon Cancer	Schetter et al. 2008	84 pairs tumor-normal	miRNA expression profiles and clinical correlation for colon cancer.	37 miRNAs differentially expressed in tumors: high miR-21 was in adenomas and tumors with more advanced TNM staging. High miR-21 expression was also associated with poor survival and therapeutic outcome.
	Scheepeler et al. 2008	10 normals 49 tumors	miRNA expression profiles of colon cancer with different microsatellite status and prognosis.	miR-145 showing the lowest expression in cancer relative to normal tissue. miR-142-3p , -212 , -151 , -144 were associated with tumor microsatellite status. High expression of miR-320 or -498 associated with good prognosis.
	Arndt et al. 2009	8 cell lines 4 normals 45 tumors	miRNA expression profiles for colon cancer.	37 miRNAs discriminate between colorectal cancer and normal tissues. 22 miRNAs were differentially expressed between normal and early stage cancer including increases in miR-21 and -224 and decreases in miR-133a and -145 . A differential expression in miR-31 , -7 , -99b , -378* , -133a , -125a discriminates between early and late stage.
Prostate Cancer	Ambs et al. 2008	16 normals 60 tumors	First miRNA expression profiles for prostate cancer.	Up-regulated miRs: miR-32 , -182 , -31 , -26a , -200c , -196a , -106b/25 cluster; down-regulated miRNAs included miR-520h , -494 , -490 , -1/133a cluster.
	Tong et al. 2009	40 pairs normal-tumor	miRNA expression profiles of paired microdissected malignant and non-involved areas from stage T2a/b, early relapse and non-relapse cancer patients.	miR-23b , -100 , -145 , -221 , -222 were significantly downregulated in malignant tissues. Patients with post-surgery elevation of prostate-specific antigen (chemical relapse) displayed a distinct profile of 16 miRNAs, as compared with those with non-relapse disease.
	Schaefer et al. 2010	76 pairs normal-tumor	miRNA expression profiles for prostate cancer and clinical correlation.	miR-16 , -31 , -125b , -145 , -149 , -181b , -184 , -205 , -221 , -222 were downregulated and miR-96 , -182 , -182 , -183 , -375 were upregulated. Expression of 5 miRNAs correlated with Gleason score or pathological tumor stage.
Liver Cancer	Murakami et al. 2006	25 pairs normal-tumor	miRNA expression profiles in hepatocellular carcinoma and non-tumorous tissues.	miR-18 , precursor miR-18 , -224 , -199a* , -195 , -199a , -200a , -125a differentially expressed between cancer and normal tissues; miR-92 , -20 , -18 and precursor miR-18 were significantly higher in poorly differentiated tumors. In contrast, miR-99a exhibited a positive correlation between expression levels and degree of tumor differentiation.
	Budhu et al. 2008	241 pairs normal-tumor	miRNA expression profiles in liver cancer predict metastasis.	A unique 20-miRNA metastasis signature was identified that could predict primary neoplastic tissues with venous metastases from metastasis-free solitary tumors. miR-219-1 , -207 , and -338 were most highly up-regulated, whereas miR-34a , -30c-1 , -148a were most highly down-regulated in metastasis cases.
	Wang et al. 2008	4 normals 46 tumors	miRNA expression profiles in hepatocellular carcinoma and non-tumorous tissues.	miR-224 overexpression identified in all tumors and miR-200c , -200 , -21 , -224 , -10b , -222 specific deregulation in benign or malignant tumors. miR-96 was overexpressed in HBV tumors, and miR-126* was down-regulated in alcohol-related hepatocellular carcinoma. Down-regulations of miR-107 and miR-375 were specifically associated with HNF1alpha and beta-catenin gene mutations, respectively.
	Ji et al. 2009	241 pairs normal-tumor	miRNA expression profiles in hepatocellular carcinoma and non-tumorous tissues and significant correlation to survival.	Reduced miR-26 in tumors and the expression of miR-26a and miR-26b in nontumor liver tissue was higher in women than in men. Low miR-26 is associated to a short overall survival but a better response to interferon therapy.
	Toffanin et al. 2011	89 tumors	miRNA expression profiles in hepatocellular carcinoma and non-tumorous tissues: identification of tumor subtypes and new oncomiRs.	3 main clusters of hepatocellular carcinoma: beta-catenin gene mutated tumors (36%), interferon-response-related genes (33%), and tumors with abnormal activation of IGF and mTOR-(PI)3K pathways (31%). A subset of tumors in last subclass (9%) overexpressed a family of miRNAs from chr19q13.42: miR-517a and miR-520c (from chr19q13.42) increased proliferation, migration, and invasion of HCC cells in vitro.

Cancer	Authors	Samples	Main findings	miRNA signature
Ovarian Cancer	Iorio et al. 2007	5 cell lines 15 normals 69 tumors	miRNA expression profiles of ovarian cancer vs normal and tumor subtype-specific miRNA signature.	29 miRNAs differentially expressed between normal and tumor with a classification rate of 89%. 4 up-modulated: miR-200a, -200b, -200c, -141; 25 down-modulated: miR-199a, -140, -145, -125b-1 among the most significant.
	Yang et al. 2008	10 normal cells 10 tumors	miRNA expression profiles of ovarian cancer vs normal epithelial cells.	Up-regulation of miR-214, -199a*, -200a and down-regulation of miR-100; alterations of the first three miRNAs is associated with late-stage and high-grade ovarian tumors. miR-214 induces cell survival and cisplatin resistance by targeting the PTEN/Akt pathway.
	Wu et al. 2009	10 pairs normal-tumor	miRNA expression profiles of endometrioid ovarian cancer vs normal tissues.	17 up-regulated (miR-205, -449, -429) and 6 down-regulated (miR-204, -99b, -193b) miRNAs in endometrioid adenocarcinoma samples.
	Marchini et al. 2011	144 tumors	miRNA expression profiles of stage I epithelial ovarian cancer assess the existence of a miRNA signature associated with overall and progression-free survival.	34 miRNAs were associated with survival. Between them miR-200c, -199a-3p, -199a-5p were highly associated with overall and progression-free survival.
	Devor et al. 2011	4 normal 23 tumors	miRNA signature in ovarian cancer vs normal and in serous (9) and endometrioid (14) subtypes	7 miRNAs was down-regulated and 13 miRNAs up-regulated in both adenocarcinomas. miR-133b the most repressed miRNA in both adenocarcinomas; miR-205 the most expressed miRNA. miR-135b, -200a, -200b, -200c, -141, -429 significantly overexpressed in both types of endometrial cancers.
Gastric Cancer	Luo et al. 2009	2 cell lines 3 normals 24 tumors	miRNA expression profiles of gastric cancer vs normal.	19 miRNAs down-regulated and 7 miRNAs up-regulated. miR-433 and miR-9 were remarkably down-regulated in the carcinoma samples.
	Ueda et al. 2010	160 pairs normal-tumor	miRNA expression profiles of gastric cancer vs normal and miRNA signature in histological subtypes	22 miRNAs upregulated and 13 downregulated in gastric cancer. Diffuse-type and intestinal-type subtypes were discriminated by miRNA expression. miR-125b, -199a, -100 were the most important microRNAs involved progression signature. Low let-7g and miR-433 and high expression of miR-214 were associated with unfavourable outcome in overall survival independent of clinical covariates, including depth of invasion, lymph-node metastasis, and stage.
	Brenner et al. 2011	45 tumors	miRNA expression profiles of primary tumor of patients with recurrent and non-recurrent gastric cancer.	miR-451, -199a-3p, -195 differentially expressed in gastric tumors from patients with good prognosis vs bad prognosis. High expression of each miR was associated with poorer prognosis for both recurrence and survival. miR-451 showed a positive predictive value for non-recurrence of 100%.
	Kim et al. 2011	34 normals 90 tumors	miRNA signature distinguishes gastric cancer from normal stomach epithelium from healthy volunteers, and a chemoresistance miRNA signature that is correlated with time to progression after cisplatin/fluorouracil therapy	30 miRNAs inversely correlated with time to progression of disease after chemotherapy whereas 28 miRNAs positively correlated. Among the upregulated miRNAs associated with chemosensitivity: let-7g, miR-342, -16, -181, -1, -34 known to regulate apoptosis.
Esophageal Cancer	Guo et al. 2008	31 pairs normal-tumor	miRNA expression profiles of esophageal cancer vs normal tissues.	46 miRNAs differently expressed between the cancerous and adjacent normal tissues. A minimal set of 7 distinguishes malignant from normal esophageal tissues: miR-25, -424, -151 showed up-regulation and miR-100, -99a, -29c, -140* showed down-regulation. High miR-103/107 correlated with poor survival.
	Yang et al. 2009	32 pairs normal-tumor	miRNA expression profiles of esophageal cancer progression from Barrett's low to high grade dysplasia to adenocarcinoma.	111 miRNAs differentiated the adenocarcinoma tissues with 100% accuracy. 11 miRNAs may be important in the progression from low-grade to high-grade dysplasia. let-7b/a/c/f, miR-345, -494, -193a were modulated in the progression from high-grade dysplasia to adenocarcinoma, and all of them were down-regulated in esophageal adenocarcinoma.
	Mathe et al. 2009	170 tumors	Identification of miRNAs involved in major histologic types of esophageal carcinoma and significant associations with prognosis.	In adenocarcinoma patients: high expression of miR-21, -223, -192, -194 and low miR-203 levels. In squamous carcinoma: high expression of miR-21 and low miR-375 levels. High miR-21 in normal tissue of squamous carcinoma and low levels of miR-375 in cancerous tissue of adenocarcinoma patients with Barrett's were strongly associated with worse prognosis.
	Fassan et al. 2011	14 normals 23 dysplasias	miRNA expression profiles in Barrett's subtypes (7 low grade dysplasia, 5 high grade dysplasia, 11 Barrett's adenocarcinomas)	Up-regulation of miR-215, -560, -615-3p, -192, -326, -147 and down-regulation of miR-100, -23a, -605, -99a, -205, let-7c, -203.
	Feber et al. 2011	45 tumors	miRNA expression profiles provide prognostic utility in staging esophagus cancer patients and elucidate steps in the metastatic pathway and allow for development of targeted therapy.	Up-regulation of miR-143, -199a_3p, -199a_5p, -100, -99a predicted a worse survival. miR-99b, -199a_3p and _5p also associated with the presence of lymph node metastasis.

C

Table 2. miRNA profiling in cancer.

4. Oncogenic microRNAs

Although studies linking miRNA dysfunctions to human diseases are in their infancy, a great deal of data already exists, establishing an important role for miRNAs in the pathogenesis of cancer. Many miRNAs have been shown to function as oncogenes in the

majority of cancers profiled to date (**Table 3**). *MiR-21* displays a strong evolutionary conservation across a wide range of vertebrate species in mammalian, avian and fish clades [70]. It has been demonstrated that a primary transcript containing *miR-21* (i.e., *pri-miR-21*) is independently transcribed from a conserved promoter that is located within the intron of the overlapping protein-coding gene *TMEM49* [71]. Several studies suggest that this miRNA is oncogenic [72-74] and that it may act as an antiapoptotic factor. For example, Chan et al. have found that miR-21 is commonly and markedly up-regulated in human glioblastoma and that inhibiting miR-21 expression leads to caspase activation and associated apoptotic cell death [72]. Moreover, Zhu and collaborators provided the first evidence that *miR-21* regulates invasion and metastasis, at least in part, by targeting metastasis-related tumor suppressor genes such as TPM1, programmed cell death 4 (PDCD4) and maspin [73]. Furthermore, examination of human breast tumor specimens revealed an inverse correlation of *miR-21* with PDCD4 and maspin [74]. The final proof of miR-21 oncogenic activity came from the Slack laboratory where the first conditional knock-in of miR-21 overexpressing mice was generated. The mice developed a severe pre-B-cell lymphoma but when miR-21 was reduced to endogenous levels, the mouse tumors completely disappeared, defining the concept of “oncomiR addition” [75].

Another important oncogenic miRNA is represented by *miR-155*. Several groups have shown that *miR-155* is highly expressed in pediatric Burkitt’s lymphoma [76], Hodgkin’s disease [77], primary mediastinal non-Hodgkin’s lymphoma [77], chronic lymphocytic leukemia (CLL) [78], acute myelogenous leukemia (AML) [79], lung cancer [80], pancreatic cancer [81], and breast cancer [80]. Dr. Croce laboratory reported that *miR-155* transgenic mice develop acute lymphoblastic leukemia/high-grade lymphoma and that most of these leukemias start at approximately nine months, irrespective of the mouse strain, preceded by a polyclonal pre-B-cell proliferation [82].

Another example of “oncomiR” is represented by *miR-221&222* cluster that is highly upregulated in a variety of solid tumors, including thyroid cancer [83], hepatocarcinoma [84], estrogen receptor negative breast tumor [85], and melanoma [86]. Elevated *miR-221&222* expression has been causally linked to proliferation [85-87], apoptosis [88-89], and migration [89] of several cancer cell lines. We recently reported that the hepatocyte growth factor receptor (MET) oncogene, through c-Jun transcriptional activation, upregulates *miR-221&222* expression, which, in turn, by targeting *PTEN* and *TIMP3*, confers resistance to tumor necrosis factor-related apoptosis-inducing ligand (TRAIL) and enhances tumorigenicity of lung and liver cancer cells [89]. The results suggest that therapeutic intervention involving the use of miRNAs should not only sensitize tumor cells to drug-inducing apoptosis but also inhibit their survival, proliferation, and invasion [89].

The miR-106b-25 polycistron is composed of the highly conserved miR-106b, miR-93, and miR-25 that accumulate in different types of cancer, including gastric, prostate, and pancreatic neuroendocrine tumors, as well as neuroblastoma and multiple myeloma. Petrocca and collaborators [90] demonstrated that E2F1 regulates miR-106b, miR-93, and miR-25, inducing their accumulation in gastric tumors. Conversely, miR-106b and miR-93 control E2F1 expression, establishing a negative feedback loop that may be important in preventing E2F1 self-activation and apoptosis. On the other hand, miR-106b, miR-93, and

<i>miRNA</i>	<i>Target</i>	<i>Tumor</i>
<i>miR-21</i>	PTEN	cholangiocarcinoma
	TPM1	breast cancer
	PDCD4	breast cancer
	SPRY1	
	RECK, TIMP3	glioblastoma
	p63, JMY, TOPORS, TP53BP2, DAXX, HNRPK, TGFβRII	glioblastoma
	MARKS	prostate cancer
<i>miR-155</i>	ANP32A, SMARCA4	prostate cancer
	SOCS1	breast cancer
	CEBPB, PU.1, CUTL1, PICALM	AML
	BACH1, ZIC3	
	ETS1, MEIS1	human cord blood CD34+
	C-MAF	lymphocytes
	HGAL	diffuse large B-cell lymphoma
	JMJD1A	nasopharyngeal carcinoma
	WEE1	breast cancer
	TP53INP1	pancreatic cancer
	SMAD1, SMAD5, HIVEP2, CEBPB, RUNX2, MYO10	
	FOXO3A	breast cancer
	hMSH2, hMSH6, hMLH1	colon cancer
	SMAD5	diffuse large B-cell lymphoma
	p27(KIP1)	glioblastoma, prostate and thyroid carcinoma
	p57 (KIP2)	normal fibroblast
<i>miR-221/222</i>	PTEN, TIMP3	non small cell lung cancer and hepatocellular carcinoma
	FOXO3A	breast cancer
	KIT	Endotelial cells
	ESR1	breast cancer
	PUMA	glioblastoma
	TRSP1	breast cancer
	PTPμ	glioblastoma
	DICER	breast cancer
	APAF1	non small cell lung cancer
	BIM, p21	gastric cancer
<i>miR-106a~363</i>	E2F1	prostate cancer
<i>miR-106b~25</i>	PTEN	prostate cancer
<i>miR-17-92</i>	TSP-1, CTGF	colon
	E2F2, E2F3	prostate/Burkitt lymphoma/testis carcinoma/
	BIM PTEN	c-Myc induced lymphoma
	HIF1α	lung cancer
	PTPRO	cervix tumor cell line
	p63	myeloid cells
	BIM, PTEN, PRKAA1, PPP2R5e,	T-cell acute lymphoblastic leukaemia
	JAK1	endothelial cells
	HBP1	breast cancer
	p21(WAF1)	Ras induced senescent-fibroblasts
	TGFβII, SMAD4	glioblastoma
	MnSOD, GPX2, TRXR2	prostate
	HOXB1, HOXB3	pancreatic cancer
	HOXD10	breast cancer
	KLF4	esophageal cancer
<i>miR-10a/10b</i>	TIAM1	breast cancer
	NF1	Ewing's sarcoma

Table 3. *-oncomiRs*

miR-25 overexpression causes a decreased response of gastric cancer cells to TGF β by downregulating p21 and Bim, the two most downstream effectors of TGF β -dependent cell cycle arrest and apoptosis, respectively.

Another example of a miRNA locus with oncogenic properties is represented by the *miR-17-92* cluster, which consists of six miRNAs: miR-17-5p, -18, -19a, -19b, -20a, and -92-1. The miR-17-92 cluster is located in a region frequently amplified in several types of lymphoma and solid tumors [91-92]. It has been shown that mice deficient for miR-17-92 die shortly after birth with lung hypoplasia and a ventricular septal defect. This cluster is also essential for B cell development; its absence, in fact, leads to increased levels of the proapoptotic protein Bim and inhibits B cell development at the pro-B-to-pre-B transition [93]. All together these studies indicate that many miRNAs have oncogenic activity. Importantly, their knockdown through the use of antisense oligonucleotides, inhibits the development of cancer-associated phenotypes, laying the groundwork for the creation of miRNA-based therapies [94-96].

5. Tumor suppressor microRNAs

The first evidence that miRNAs are involved in cancer comes from the finding that miR-15 and miR-16 are downregulated or deleted in most patients with chronic lymphocytic leukemia (CLL) (**Table 4**) [68]. They are transcribed as a cluster (*miR-15a-miR-16-1*) that resides in the 13q14 chromosomal region. Deletions or point mutations in region 13q14 occur at high frequency in CLL, lymphoma, and several solid tumors [97]. Their expression is inversely correlated to *BCL2* expression in CLL [98]. The tumor suppressor function of *miR-15a/16-1* has also been addressed in vivo. In immunocompromised nude mice, ectopic expression of *miR-15a/16-1* was found to cause dramatic suppression of tumorigenicity of MEG-01 leukemic cells that exhibited a loss of endogenous expression of *miR-15a/16-1*. Furthermore, Klein et al. [99] generated transgenic mice with a deletion of the *miR-15a-miR-16-1* cluster, causing development of indolent B-cell-autonomous, clonal lymphoproliferative disorders, recapitulating the spectrum of CLL-associated phenotypes observed in humans. Recently, Bonci et al. reported that the *miR-15a-miR-16-1* cluster targets not only *BCL2* but also *CCND1* (encoding cyclin D1) and *WNT3A* mRNA, which promote several prostate tumorigenic features, including survival, proliferation, and invasion [100]. Together, these data suggest that *miR-15a/16-1* genes are natural antisense interactors of *BCL2* and probably other oncogenes and that they can be used to suppress tumor growth in therapeutic application for a variety of tumors [100].

In mammals, the miR-34 family comprises three processed miRNAs that are encoded by two different genes: miR-34a is encoded by its own transcript, whereas miR-34b and miR-34c share a common primary transcript. The miR-34 family has been shown to form part of the p53 tumor-suppressor network: their expression is directly induced by p53 in response to DNA damage or oncogenic stress [101-102]. He et al. identified different miR-34 targets such as cyclin E2 (*CCNE2*), CDK4, and MET. Silencing these selected miR-34 targets through the use of small interfering RNAs (siRNAs) led to a substantial cell cycle arrest in G1.

Moreover, ectopic miR-34 delivery caused a decrease in levels of phosphorylated retinoblastoma gene product (Rb), consistent with lowered activity of both CDK4 and CCNE2 complexes [102]. BCL2 and MYCN were also identified as miR-34a targets and likely mediators of the tumor suppressor phenotypic effect in neuroblastoma [103]. It has been also reported that p53 activation suppressed the EMT-inducing transcription factor SNAIL via induction of the miR-34a/b/c genes. In fact, suppression of miR-34a/b/c by anti-miRs caused up-regulation of SNAIL and cells displayed EMT markers, enhanced migration and invasion [104].

MicroRNA-122 (miR-122) is a liver-specific microRNA and is frequently downregulated in liver cancer [105]. Xu et al. reported that restoration of miR-122 in hepatocellular carcinoma cells could render cells sensitive to chemotherapeutic agents adriamycin or vincristine through downregulating antiapoptotic gene Bcl-w and cell cycle related gene cyclin B1 [106]. Another group found that over-expression of miR-122 inhibits hepatocellular carcinoma cell growth and promotes the cell apoptosis by affecting Wnt/ β -catenin signalling pathway [107]. Coulouarn et al. showed that miR-122 is specifically repressed in a subset of primary hepatocellular tumors that are characterized by poor prognosis [108]. They further reported that loss of miR-122 resulted in an increase of cell migration and invasion and that restoration of miR-122 reverses this phenotype [108]. The final understanding of the tumor suppressor role for mir-122 role in liver cancer came from a recent study where miR-122 knockout mice were studied. When miR-122 KO mice aged, hepatic inflammation ensued, preceding the progressive onset of fibrosis and, eventually, tumors resembling human liver cancer. These pathologic manifestations were associated with hyperactivity of oncogenic pathways and hepatic infiltration of inflammatory cells that produce pro-tumorigenic cytokines, including IL-6 and TNF [109].

<i>miRNA</i>	<i>Target</i>	<i>Tumor</i>
<i>miR-15/16</i>	BCL2	CLL
	COX-2	colon cancer
	CHEK1	follicular lymphoma
	CEBP β , CDC25a, CCNE1	fibroblast
	VEGF, VEGFR2, FGFR1	fibroblast
	FGF2, FGFR1	cancer associated fibroblast
	CCNE1	
	FGFR1, PI3KCa, MDM4, VEGFa	multiple myeloma
	WIP1	
	BMI-1	ovarian cancer
<i>miR-34</i>	CCND1, CCND2, CCNE1	lung cancer
	SIRT1	colon cancer
	BCL2, NOTCH, HMGA2	
	MYC	fibroblast
	AXL	lung cancer
	MET	ovarian cancer
	NANOG, SOX2, MYCN	embryonic fibroblast
	SNAIL	colon cancer

Table 4. Tumor suppressor *miRS*

6. MetastamiRs

Metastasis is the result of cancer cells detaching from a primary tumor, consequently adapting to distant tissues and organs, and forming a secondary tumor [110] and this ability of cancer cells to metastasize is a hallmark of malignant tumors [111-112]. To successfully metastasize, a tumor cell must complete a complex set of processes, including invasion, survival and arrest in the circulatory system, and colonization of foreign organs. Despite great advancements in knowledge of metastasis biology, the molecular mechanisms are still not completely understood. Several miRNAs have been shown to initiate invasion and metastasis by targeting multiple proteins that are major players in these cellular events, thus they have been denominated as metastamiRs (**Table 5**). It seems that these metastasis-associated miRNAs do not influence primary tumor either in development or initiation steps of tumorigenesis, but they regulate key steps in the metastatic program and processes, such as epithelial-mesenchymal transition (EMT), apoptosis, and angiogenesis. Ma et. al reported that miR-10b is highly expressed in metastatic breast cancer cells and positively regulates cell migration and invasion. Overexpression of miR-10b in otherwise non-metastatic breast tumors initiates robust invasion and metastasis [113]. The team led by Joan Massague found that miR-335, miR-126, and miR-206 are metastasis-suppressors in breast cancer [114]. MiR-126 and miR-206 restoration reduced overall tumor growth and proliferation, whereas miR-335 inhibits metastatic cell invasion through targeting of the progenitor cell transcription factor *SOX4* and extracellular matrix component tenascin C [114]. Others miRNAs with prominent roles in breast cancer metastasis have been reported. It has been reported that miR-31 inhibited multiple steps of metastasis including invasion, anoikis, and colonization leading to almost complete reduction of lung metastasis [115]. Clinically, miR-31 levels were lower in breast cancer patients with metastasis. In addition, miR-9, which is up-regulated in breast cancer cells, directly targets *CDH1*, the E-cadherin-encoding messenger RNA, leading to increased cell motility and invasiveness [116].

Another important aspect of the metastatic dissemination is represented by the epithelial-to-mesenchymal transition (EMT) that allow neoplastic cells to abandon their primary site and survive in the new tissue. During EMT, an epithelial neoplastic cell loses cell adhesion by repressing E-cadherin expression and thereby the cell increases its motility. Numerous studies have shown that different microRNAs are modulated during EMT and one of the best-studied example is represented by the miR-200 family. These miRs are commonly lost in aggressive tumors such as lung, prostate, and pancreatic cancer. It has been shown that miR-200 family members directly target *ZEB1* and *ZEB2*, transcription repressors of E-cadherin [117]. In fact, in the highly aggressive mouse lung cancer model where *KRAS* is constitutively activated and *p53* function is perturbed, miR-200 ectopic expression prevented metastasis by repressing *ZEB1* and *ZEB2* and preventing E-cadherin down-regulation [117]. However, overexpression of the miR-200 family is associated with an increased risk of metastasis in breast cancer and this overexpression promotes metastatic colonization in mouse models, phenotypes that cannot be explained by E-cadherin expression alone [118]. By using proteomic profiling of the targets of mesenchymal-to-epithelial (MET)-inducing miR-200, the authors discovered that miR-200 globally targets secreted proteins in breast cancer cells. Between the 38 modulated target genes, *Sec23a*,

which is involved in transporting protein cargo from the endoplasmic reticulum to the Golgi, shows a superior association with human metastatic breast cancer as compared to the currently recognized miR-200 targets ZEB1 and the EMT marker E-cadherin. EMT is first acquired in the onset of transmigration and then reversed in the new metastatic site. Korpál et al. have shown that the miR-200 status predicts predisposition of the cancer to successful metastasis [119].

<i>miRNA</i>	<i>Target</i>	<i>Tumor</i>
<i>miR-10a/10b</i>	HOXB1, HOXB3	pancreatic cancer
	HOXD10	breast cancer
	KLF4	esophageal cancer
	TIAM1	breast cancer
	NF1	Ewing's sarcoma
<i>miR-9</i>	PRDM1/BLIMP-1	lymphomas
	CDH1	breast cancer
	CAMTA	glioblastoma
<i>miR-31</i>	ITGA5, RDX, RHOA FZD3, M-RIP, MMP16	breast cancer
	SATB2	cancer associated fibroblast
<i>miR-200 family</i>	ZEB1, ZEB2	breast cancer
	ERRFI-1	bladder cancer
	ZEB1, CTNNB1	nasopharyngeal carcinoma
	BMI-1	pancreatic cancer
	PLC γ 1	breast cancer
	FAP1	
	SUZ12	breast cancer
	FLT1/VEGFR1	lung cancer
	JAG1, MALM2, MALM3	
	FN1, LEPR, NTRK2, ARHGAP19	breast and endometrial cancer
	p38 α	ovarian cancer

Table 5. *metastamiRS*

7. Other non-coding RNAs: Biology and implications in cancer

7.1. snoRNAs: From post-transcriptional modification to cancer

Small nucleolar RNAs (snoRNAs) have, for many years, been considered one of the best-characterized classes of non-coding RNAs (ncRNAs) [120-123] but despite the common assumption that snoRNAs only have cellular housekeeping functions, in the past few years, independent reports have converged in implicating snoRNAs in the control of cell fate and oncogenesis [124-130]. SnoRNAs are small RNAs of 60-300nt in length that specifically accumulate in the nucleolar compartment of the cell where are in charge of the 2'-O-ribose methylation and pseudouridylation of specific ribosomal RNA nucleotides, essential

modification for the efficient and accurate production of the ribosome [120-122]. The snoRNAs carry out their function in the form of small nucleolar ribonucleoproteins (snoRNPs), each of which consists of a box C/D or box H/ACA guide RNA, and four associated C/D or H/ACA snoRNP proteins (**Figure 2**). In both cases, snoRNAs hybridize specifically to the complementary sequence in the rRNAs, and the associated protein complexes then carry out the appropriate modification on the nucleotide that is identified by the snoRNAs. Biogenesis of vertebrate snoRNPs is remarkable and highly variable: in fact snoRNA gene organization ranges from independently transcribed genes, endowed with their own promoter elements, to intronic coding units lacking an independent promoter. In both yeast and animals, processing of intron-encoded snoRNAs is largely splicing-dependent; in contrast, the production of plant snoRNAs from introns seems to rely on a splicing-independent process [131]. Moreover, in both contexts (intergenic or intronic), genes can be either single or part of clusters. In the latter case, the generation of individual snoRNAs involves the enzymatic processing of polycistronic precursor RNAs. Such a processing, at least in yeast, appears to involve the same combination of endo- and exoribonucleases required for the maturation of monocistronic pre-snoRNAs [132-134]. The first indication that snoRNAs might have important roles in human disease was provided by the genetic studies on Prader-Willi syndrome (PWS), an inherited human disorder characterized by a complex phenotype, including mental retardation, decreased muscle tone and failure to thrive at birth, short stature, hypogonadism, sleep apnea, behavioral problems and hyperphagia (an insatiable appetite) that can lead to severe obesity [135]. The disease is caused by the genomic loss of the imprinted chromosomal 15q11-q13 locus which is normally only active on the paternal allele. The only characterized and conserved genes within this 121-kb-long genomic interval are the numerous HBII-85 snoRNA gene copies, thus suggesting that loss of expression of these repeated small C/D RNA genes might play a role in conferring some (or even all) phenotypes of the human disease and PWS-like phenotypes in mice (neonatal lethality, growth retardation and hypotonia). In fact, it has been shown that a site-specific deletion of the entire murine MBII-85 gene cluster led to post-natal growth retardation with low postnatal lethality (<15%) only seen in some genetic backgrounds, but no obesity [136]. Although all the imprinted C/D RNAs that have been tested accumulate within the nucleolus, none of them appear to act as RNA guides to modify rRNAs or spliceosomal U-snRNAs; they are called 'orphan C/D RNAs'. So far, the MBII-52 gene clusters have attracted much attention, given that the neuronal-specific MBII-52 small RNA is predicted to interfere (A-to-I RNA editing and/or alternative RNA splicing) with the post-transcriptional regulation of the pre-mRNA that encodes the 5-HT_{2c} (5-hydroxytryptamine 2C) receptor, playing a key role in regulating serotonergic signal transduction [137-138]. These observations raised the possibility that snoRNAs could have functions completely independent from their traditional activities and carry out other regulatory roles. The first insights into the potential roles of snoRNAs in cancer began with a study that identified C/D box snoRNA U50 and its host gene U50HG at the breakpoint in the t(3;6) (q21;q15) translocation in a diffuse large B cell lymphoma [139]. Moreover, snoRNAU50 gene has been found to undergo to a frequent copy number loss and a transcriptional downregulation in breast and prostate cancer samples [139,140]. In addition, a 2-bp deletion in U50 sequence also occurred both somatically and in germline, leading to increased incidence of homozygosity for the deletion in cancer cells [140].

Figure 2. snoRNAs. **A.** Boxed sequences C and D (named from conserved, nuclease-resistant sequences that were originally identified in snoRNA U3) are hallmarks of the C/D box snoRNAs; boxed sequences H (Hinge region) and ACA are hallmarks of the H/ACA box snoRNAs. These conserved boxed sequences are important for the associations with protein components that are required to form the functional small nucleolar ribonucleoprotein (snoRNP) complexes and for accumulation in the nucleolus. C/D box snoRNAs associate with several proteins, including fibrillarin, which is the methyl transferase that is involved in the 2'-O-methylation of particular ribonucleotides, and H/ACA box snoRNAs associate with proteins such as the pseudouridine synthase dyskerin. Antisense sequences within the C/D box and H/ACA box snoRNAs guide the snoRNP complex to the appropriate nucleotide within the target RNA (most often ribosomal RNA). In a minority of cases both C/D-associated and C'D'-associated antisense sequences within the same C/D box snoRNA can act as guides for 2'-O-methylation of the target RNA. The eukaryotic H/ACA box snoRNAs contain two hairpin domains with complementary regions flanking the uridine to be converted in the target rRNA, at a position 14–16 nucleotides upstream of the conserved H and/or ACA box. Most mammalian snoRNAs are encoded within the introns of genes producing 5' terminal oligopyrimidine (5'TOP) RNAs. **B.** Organization of snoRNA genes in representative eukaryotic genomes **C.** Small nucleolar RNAs (snoRNAs) in vertebrate are predominantly located in introns. Following splicing, debranching and trimming, mature snoRNAs are either exported, in which case they function in ribosomal RNA (rRNA) processing, or remain in the nucleus, where they are involved in alternative splicing and additional yet unknown functions.

SnoRNA42 (SNORA42) is located on chromosome 1q22 which is a commonly frequent amplified genomic region in lung cancer and overexpression of SNORA42 is frequently and remarkably found in NSCLC cells [141]. In addition, SNORA42 exhibited close correlations between its increases of copy number and expression level, suggesting that SNORA42

overexpression could be activated through its amplification. Importantly, engineered repression of SNORA42 caused marked repression of lung cancer growth *in vitro* and *in vivo* and it is associated with increased apoptosis by a p53-dependent pathway. Although not exhibiting apoptosis, p53 null and mutant p53 cancer cells with reduced levels of SNORA42 also show inhibited proliferation and growth, suggesting that SNORA42 knockdown can inhibit cell proliferation in p53-dependent or -independent manner. These independent studies on U50 and SNORA42 provide evidence for the functional importance of snoRNAs in cancer, and they show that snoRNAs can promote, as well as suppress, tumour development. In 2002, Wu and coworkers demonstrated that the expression of snoRNAs 5S was differentially displayed in different tissues and noticeably was highly expressed in normal brain, but its expression drastically decreased in meningioma [142]. Recently, genome-wide approaches identified six snoRNAs (SNORD33, SNORD66, SNORD73B, SNORD76, SNORD78, and SNORA42) that were statistically differently expressed between the non small cell lung cancer tumor and paired noncancerous samples [143]. Specifically, all these snoRNAs displayed a strong up-regulation in lung tumor specimens and the majority of them is located in commonly frequent genomic amplified regions in lung cancer: SNORD33 is located in chromosome 19q13.3 that contain potential oncogenes in lung cancer, while SNORD66 and SNORD76 are situated in chromosomal regions 3q27.1 and 1q25.1, respectively 3q27.1 and 1q25.1 are two of the most frequently amplified chromosomal segments in solid tumors, particularly NSCLC [143].

As well as the initial evidence that snoRNAs are involved in cancer development, there are some preliminary data showing that the genes that host snoRNAs might also contribute to the aetiology of this disease. A research screening for potential tumor-suppressor genes identified that Growth arrest-specific transcript 5 (gas5) gene as almost undetectable in actively growing cells but highly expressed in cells undergoing serum starvation or density arrest [144-145]. Gas5 is a multi-snoRNA host gene which encodes 9 (in mouse) or 10 (in human) snoRNAs and like all known snoRNA host genes exhibit characteristics which belong to the class of genes encoding 5' terminal oligopyrimidine (5'TOP) mRNAs [146]. The first and stronger evidence that GAS5 is related to cancer is the identification that GAS5 transcript levels are significantly reduced in breast cancer samples relative to adjacent unaffected normal breast epithelial tissues and some, but not all, GAS5 transcripts sensitize mammalian cells to apoptosis inducers [147]. Other studies have also showed that GAS5 reduced expression is associated with poor prognosis in both breast cancer and head and neck squamous cell carcinoma [148]. Of note, GAS5 has been also identified as a novel partner of the BCL6 in a patient with diffuse large B-cell lymphoma, harboring the t(1;3)(q25;q27) [149]. Another example of a mature spliced transcript that harbors C/D-box snoRNAs and can function independently of the snoRNAs is represented by the transcript Zfas1 [150]. This gene intronically hosts three C/D box snoRNAs (Snord12, Snord12b, and Snord12c) and has been identified as one of the most differentially expressed gene during mouse mammary development. siRNA-mediated downregulation of Zfas1 mRNA in a mouse mammary cell line increased proliferation and differentiation without substantially affecting the levels of the snoRNA hosted within its intron. The human homologue, ZFAS1 (also known as ZNF1-AS1), which is predicted to share secondary structural features with

mouse *Zfas1*, is expressed at high levels in the mammary gland and is downregulated in breast cancer. Taken together, these findings indicate that snoRNA host genes might have important functions in regulating cellular homeostasis and, potentially, cancer biology but more studies are needed to understand their involvement in molecular basis of disease and classify them as sources of potential biomarkers and therapeutic targets.

Another important aspect of the association between snoRNAs and tumorigenesis is represented by the involvement of their associated proteins in cancer. A point mutation in the *DKC1* gene is the cause of a rare X-linked recessive disease, the dyskeratosis congenita (DC) [151-152]. Individuals with DC display features of premature aging, as well as nail dystrophy, mucosal leukoplakia, interstitial fibrosis of the lung, and increased susceptibility to cancer. *DKC1* codes for dyskerin, a putative pseudouridine synthase, which carries out two separate functions, both fundamental for proliferating cells. One function is the pseudouridylation of ribosomal RNA (rRNA) molecules as a part of the H/ACA ribonucleoprotein complex, and the other is the stabilization of the telomerase RNA component necessary for telomerase activity. *Dkc1* mutant mice recapitulate the major features of DC, including an increased susceptibility to tumor formation. Early generation (G1 and G2) of *Dkc1* mutant mice showed a full spectrum of DC and presented alterations in rRNA modification, whereas defects in telomere length were not evident until G4 mice, suggesting that deregulated ribosome function is important for the initiation of DC and that impairment in telomerase activity in *Dkc1* mutant mice may modify and/or exacerbate the disease in later generations. To this regard, *DKC1* was identified as one of only seventy genes that, collectively, constitute a gene expression profile that strongly correlates with the development of aneuploidy and is associated with poor clinical prognosis in a variety of human cancers. Therefore, one hypothesis is that an alteration of physiologic dyskerin function, irrespective of the mechanism, may perturb mitosis and contribute to tumorigenesis but this idea will require more detailed investigation. Another possibility is related to the strong effect of dyskerin loss on H/ACA accumulation. Recent findings in fact have shown that some H/ACA box and C/D box can be processed to produce small RNAs, at least some of which can function like miRNAs [153]. Such processing may be of crucial importance, as miRNAs have important roles in the development of many cancers as previously discussed. To date, Xiao and colleagues have recently reported that an H/ACA box snoRNA-derived miRNA, miR-605, has a key role in stress-induced stabilization of the p53 tumour suppressor protein [154]. p53 transcriptionally activates its negative regulator, MDM2, in addition to miR-605. miR-605 counteracts MDM2 through post-transcriptional repression; under conditions of stress, this snoRNA-derived miRNA offsets the MDM2 negative-feedback loop, generating a positive-feedback loop to enable the rapid accumulation of p53. However, whether this regulation of p53 by miR-605 is relevant to cancer biology has not yet been addressed. Like dyskerin, NHP2 and NOP10 proteins, both components of the H/ACA snoRNPs, are also significantly up-regulated in sporadic cancers and high levels may be associated with poor clinical prognosis. Moreover, germline NHP2 and NOP10 mutations give rise to autosomal recessive forms of dyskeratosis congenita, and cancer susceptibility is also a feature of these genetic forms of the disease. Since the functions of several snoRNAs have not yet been identified (orphan snoRNAs), it is possible

that disruption of snoRNP biogenesis by any mechanism may affect an array of important cellular processes, and could potentiate cancer development and/or progression.

7.2. piRNAs: Guardians of the genome

Piwi-interacting RNAs (piRNAs) are germline-specific small silencing RNAs of 24–30 nt in length, that suppress transposable elements (TE) activity and maintain genome integrity during germline development, a role highly conserved across animal species [155–156]. TEs are genomic parasites that threaten the genomic integrity of the host genome: they are able to move to new sites by insertion or transposition and thereby disrupt genes and alter the genome [157]. In animals, endogenous siRNAs also silence TEs, but the piRNA pathway is at the forefront of defense against transposons in germ cells [158]. piRNAs specifically associate with PIWI proteins, which are germline-specific members of the AGO protein family, AGO3, Aubergine (Aub) and Piwi, and form a piRNA-induced silencing complex (piRISC) which will guide the TE silencing [159–162]. Any mutations in each of the three members of the PIWI family lead to transposon derepression in the germline, indicating that they act non-redundantly during TE silencing. Initial screening of piRNA sequences revealed that there are hundreds of thousands, if not millions, of individual piRNA sequences [163–165]. Furthermore, they are characterized by the absence of specific sequence motifs or secondary structures such as miRNA precursors. Despite their large diversity, most piRNAs can be mapped to a relatively small number of genomic regions called piRNA clusters. Each cluster extends from several to more than 200 kilobases, it contains multiple sequences that generate piRNAs and some piRNAs map to both genomic strands, suggesting bidirectional transcription [163–165]. Indeed, analysis of piRNA clusters in different *Drosophila* species has shown that, although the clusters locations are conserved, their sequence content has evolved very quickly suggesting adjustments in the piRNAs patrimony in order to suppress new active transposons invading the species. Therefore, piRNA clusters may be considered as repositories of information, enabling production of many mature piRNAs that target diverse TEs. Two main pathways, highly conserved in many animal species, have been discovered to be responsible for the biogenesis of the piRNAs: the primary pathways and the Ping-Pong amplification (**Figure 3**) [166–168]. First, the primary piRNA biogenesis pathway provides an initial pool of piRNAs that target multiple TEs. Next, the Ping-Pong cycle further shapes the piRNA population by amplifying sequences that target active transposons. It is currently unclear how primary piRNAs are produced from piRNA clusters but it is likely that piRNA precursors are single-stranded and therefore do not require Dicer for their processing. Interestingly, piRNAs that associate with each member of the PIWI protein family have a distinct size, suggesting that PIWI proteins can act as ‘rulers’ that define the size of mature piRNAs. Several additional proteins (e.s. Zucchini, Armitage and Yb) have also been identified that are involved in primary piRNA biogenesis and mutations in and/or depletion of any of these three proteins eliminates primary piRNAs associated with PIWI proteins. In some cell types, such as somatic follicle cells of the *D. melanogaster* ovary, primary piRNA biogenesis is the only mechanism that generates piRNAs. However, in germline cells of the *D. melanogaster* ovary and in the pre-meiotic spermatogonia in mice, there is another mechanism called the Ping-Pong cycle that amplifies specific sequences generated by the primary biogenesis pathway

[163,169]. Mainly the Ping-Pong pathway engages AGO3 and Aubergine, both of which are accumulated in perinuclear structures located at the cytoplasmic face of the nuclear envelope in animal germline cells, named “nuage”. The pathway depends on the endoribonuclease or Slicer activity of AGO3 and Aubergine, which act catalytically one after the other, leading to a cleavage of the target RNAs between their tenth and eleventh nucleotides relative to the ‘guide’ small RNAs. This process results in the generation of repeated rounds of piRNA production having exactly the same sequence of the original primary piRNA. The ping-pong pathway amplifies piRNAs in *D. melanogaster* testes, especially those originating from TEs. Non-TE-derived piRNAs seem to be barely amplified by the amplification loop. This two steps of piRNA biogenesis can be compared with the function of the adaptive immune system in protecting against pathogens. The primary piRNA biogenesis pathway resembles the initial generation of the hypervariable antibody repertoire, whereas the amplification loop is analogous to antigen-directed clonal expansion of antibody-producing lymphocytes during the acute immune response. An emerging number of studies highlight the role of piRNAs or PIWI proteins in the regulation of tumorigenesis. First examples of the piRNA involvement in cancer is represented by the up-regulation of HIWI, one of the four human Piwi homologues, in about 60 % of seminomas [170]. In fact, HIWI maps to a locus known as a germ cell tumor susceptibility locus (12aq24.33). HIWI overexpression has also been found in somatic cells such as soft-tissue sarcomas or ductal pancreas adenocarcinoma, and strongly correlates with bad prognosis and high incidence of tumor-related death, providing an example for a potential tumorigenic role of a piRNA-related protein in somatic cells [171,172]. In some cancers, PIWIL2 overexpression has been suggested to induced resistance in cells to cisplatin, which might arise because of increased chromatin condensation that prevents the normal process of DNA repair [173]. Furthermore, new high-throughput sequencing data revealed the presence of piRNAs in somatic cells, such as HeLa cells. These somatic piRNAs appear located in the nucleolus and in the cytoplasmic area surrounding the nuclear envelope and in contrast with the large population of known piRNAs in male germ cells, this population of piRNAs is dramatically smaller [174]. Another recent study demonstrated that the level of piR-651 is significantly higher in several cancer histotype including lung, mesothelium, breast, liver, and cervical cancer compared to non-cancerous adjacent tissues and inhibition of piR-651 induced block of gastric cancer cells at the G2/M phase [175,176]. Another example is represented by the downregulation of piR-823 in gastric cancer tissues; its enforced expression inhibited gastric cancer cell growth in vitro and in vivo, suggesting a tumor suppressive properties for piR-823 [177]. Interestingly, piRNAs not are only involved in direct regulation by degradation of TE but they have also been linked to DNA methylation of the retrotraspon regions, extending piRNA functions beyond post-transcriptional silencing. In fact, CpG DNA methylation, which is required for efficient transcriptional silencing of LINE and LTR retrotransposons in the genome, is decreased in the male germ line of mice with defective PIWI proteins. Specifically, mice with defective PIWI proteins fail to establish *de novo* methylation of TE sequences during spermatogenesis, leading to the hypothesis that the piRISC can also guide the *de novo* methylation machinery to TE loci. In this scenerio, piRNAs may present a perfect guide for discriminating TE sequences from normal protein-coding genes and marking them for DNA methylation; however, the biochemical details of how these two mechanisms of piRNA

action might be linked have not yet been revealed [178,179]. All together, these data revealed that PIWI-associated RNAs and PIWI pathway has a more profound function outside germline cells than was originally thought but many more studies are needed to clarify their specific role in tumorigenesis.

Figure 3. piRNAs. **A**, schematic representation of the *Drosophila* egg chamber. **B**, piRNAs (which are 24–32 nt in length) are processed from single-stranded RNA precursors that are transcribed largely from mono- or bidirectional intergenic repetitive elements known as piRNA clusters. Unlike miRNAs and siRNAs, piRNAs do not require Dicer for their processing. First, primary piRNAs are produced through the primary processing pathway and are amplified through the ping-pong pathway, which requires Slicer activity of PIWI proteins. Subsequently, additional piRNAs are produced through a PIWI-protein-catalysed amplification loop (called the 'ping-pong cycle') via sense and antisense intermediates. Primary piRNA processing and loading onto mouse PIWI proteins might occur in the cytoplasm. The PIWI ribonucleoprotein (piRISC) complex functions in transposon repression through target degradation and epigenetic silencing. **C**, total number of piRNA clusters in different animal species according to the piRNA Database (<http://pirnabank.ibab.ac.in/>).

8. The emergence of long non-coding RNAs

Over the last decade, advances in genome-wide analyses of the eukaryotic transcriptome have revealed that most of the human genome is transcribed, generating a large repertoire of (>200 nt) long non-coding RNAs (lncRNA or lincRNA, for long intergenic ncRNA) that

map to intronic and intergenic regions [181,181]. Given their unexpected abundance, lncRNAs were initially thought to be spurious transcriptional noise resulting from low RNA polymerase fidelity [182]. However, the restricted expression of many long ncRNAs to particular developmental contexts, the often exhibiting precise subcellular localization and the binding of transcription factors to non-coding loci, suggested that a significant portion of ncRNAs fulfills functional roles beyond transcriptional remodelling [183-187]. lncRNA typically refers to a polyadenylated long ncRNA that is transcribed by RNA polymerase II and is associated with epigenetic signatures common to protein-coding genes, such as trimethylation of histone 3 lysine 4 (H3K4me3) at the transcriptional start site (TSS) and trimethylation of histone 3 lysine 36 (H3K36me3) throughout the gene body [188-189]. lncRNAs also commonly exhibit splicing of multiple exons into a mature transcript, and their transcription occurs from an independent gene promoter and is not coupled to the transcription of a nearby or associated parental gene. RNA-Seq studies now suggest that several thousand uncharacterized lncRNAs are present in any given cell type [188-189], and that the human genome may harbor nearly as many lncRNAs as protein-coding genes (perhaps ~15,000 lncRNAs), although only a fraction is expressed in a given cell type. One main characteristic of the lncRNAs is their very low sequence conservation that had fueled the idea that they are not functional. This assertion needs to be carefully considered and takes in consideration several points. First, a recent study identified the presence of 1,600 lncRNAs that show a strong evolutionary conservation and function ranging from embryonic stem cell pluripotency to cell proliferation [189]. In contrast to the protein coding genes, long ncRNAs can exhibit shorter stretches of sequence that are conserved to maintain functional domains and structures. Indeed, many long ncRNAs with a known function, such as *Xist*, only exhibit high conservation over short sections of their length [190]. Third, rather than being indicative of non-functionality, low sequence conservation can also be explained by high rates of primary sequence evolution if long ncRNAs have, like promoters and other regulatory elements, more plastic structure–function constraints than proteins [190]. The diverse selection pressures acting on long ncRNAs probably reflect the wide range of their functions which can be regrouped in three major subclasses: chromatin remodeling, transcriptional modulation and nuclear architecture/subnuclear localization.

long ncRNAs can mediate epigenetic changes by recruiting chromatin remodelling complexes to specific genomic loci resolving the paradox of how a small repertoire of chromatin remodelling complexes are able to specify the large array of chromatin modifications without any apparent specificity for the genomic loci [191,192]. A recent study found that 20% of 3300 human long non coding RNAs are bound by Polycomb Repressive Complex 2 (PRC2) [193]. Although the specific molecular mechanisms are not defined, there are several examples that can illustrate the silencing potential of lncRNAs (**Figure 4**). The first most known example is represented by the X-chromosome inactivation which is carried out by a number of lncRNAs including *Xist* and *RepA*, which bind PRC2 complex, and the antagonist of *Xist*, *Tsix* [194]. In pre-X-inactivation cells, *Tsix* competes with *RepA* for the binding of PRC2 complex; when the X-inactivation starts *Tsix* is downregulated and PRC2 becomes available to *RepA* which can actively induced the transcription of *Xist*. The up-regulated *Xist* in turn preferentially binds to PRC2 and spreads across the chromosome X

inducing PRC2-mediated trimethylated histone H3 lysine27. Another important example is represented by the hundreds of long ncRNAs which are sequentially expressed along the temporal and spatial developmental axes of the human homeobox (Hox) loci, where they define chromatin domains of differential histone methylation and RNA polymerase accessibility [195]. One of these ncRNAs, Hox transcript antisense RNA (HOTAIR), originates from the HOXC locus and silences transcription across 40 kb of the HOXD locus in trans by inducing a repressive chromatin state, which is proposed to occur by recruitment of the Polycomb chromatin remodelling complex PRC2 by HOTAIR (**Figure 4**). Recently, it has been proposed that HOTAIR has the ability to bind other histone-modifying enzymes such as the demethylase LSD1 [196]. In fact, knockdown of HOTAIR induces a rapid loss of LSD1 or PRC2 at hundreds of gene loci with the corresponding increase in expression. This model fits other chromatin modifying complexes, such as Mll, PcG, and G9a methyltransferase, which can be similarly directed by their associated ncRNAs [196]. As modulator of epigenetic landmark, it has been shown that HOTAIR has a profound effect on tumorigenesis. In fact, HOTAIR is upregulated in breast carcinoma and colon cancer and its correlates with metastasis and poor prognosis [197]. Enforced expression of HOTAIR consistently changed the pattern of occupancy of Polycomb proteins from the typical epithelial mammary cells pattern to that of embryonic fibroblasts [198]. Another important effect of lncRNAs on chromatin modification that can highlight their impact on cancer is the relationship between the lncRNA ANRIL and the INK4b/ARF/INK4a locus, encoding for three tumor-suppressor genes highly deleted or silenced in a large cohort of tumors [199]. ANRIL, which is transcribed antisense to the protein coding genes of the locus, controls the epigenetic status of the locus by interacting with subunits of PRC1 and PRC2. High expression of ANRIL is found in some cancer tissues and is associated to a high levels of PCR-mediated trimethylated histone H3 lysine27. Inhibition of ANRIL releases PRC1 and PRC2 complexes from the locus, decreases the histone methylation status with the following increase of the protein coding gene transcription. Many other tumor suppressor genes that are frequently silenced by epigenetic mechanisms in cancer also have antisense partners, which can affect gene expression with different other mechanism. First, antisense ncRNAs can mask key cis-elements in mRNA by the formation of RNA duplexes, as in the case of the Zeb2 antisense RNA, which complements the 5' splice site of an intron of Zeb2 mRNA [200]. Expression of the ncRNA prevents the splicing of the intron that contains an internal ribosome entry site required for efficient translation and expression of the ZEB2 protein with a further efficient translation (**Figure 4**). In this context, it has been evaluated that the prevalence of lncRNAs are antisense to introns, hypothesizing their role in the regulation of splicing or capable of generating mRNA duplexes that fuel the RISC machinery to silence gene expression. One major emergent theme is the involvement of the lncRNAs in the assembly or activity of transcription factors functioning as a scaffold for the docking of many proteins, mimicking functional DNA elements or modulation of PolII itself. The first example is represented by the suppression of CCND1 mediated by the lncRNAs through the recruitment and integration of the RNA binding protein TLS into a transcriptional programme. DNA damage signals induce the expression of long ncRNAs associated with the cyclin D1 gene promoter, where they act cooperatively to recruit the RNA binding

Figure 4. lncRNAs. Schematic representation of the control operated on protein coding gene by the lncRNAs at the level of chromatin remodelling, transcriptional control and post-transcriptional processing. **A**, lncRNAs (Xist, HOTAIR, ANRIL, etc) can recruit chromatin modifying complexes to specific genomic loci to localize their catalytic activity. In this case, the lncRNA recruits the Polycomb complex by inducing trimethylation of the lysine 27 residues (me3K27) of histone H3 to produce heterochromatin formation and repress gene expression. **B**, **C**, **D**, lncRNAs can regulate the transcriptional process through a range of mechanisms. First, lncRNAs tethered to the promoter of the cyclin D1 gene recruit the RNA binding protein TLS to modulate the histone acetyltransferase activity of CREB binding protein (CBP) and p300 to repress gene transcription. Second, an ultraconserved enhancer is transcribed as a long ncRNA, Evf2, which subsequently acts as a co-activator to the transcription factor DLX2, to regulate the Dlx6 gene transcription. Third, a lncRNA transcribed from the DHFR minor promoter form a triplex at the major promoter to reduce the access of the general transcription factor TFIID, and thereby suppress DHFR gene expression. **E**, a lncRNA is antisense to Zeb2 mRNA and mask the 5' splice site resulting in intron retention. This retention results in an efficient Zeb2 translation related to the presence of an internal ribosome entry site (IRE) in the retained intron.

protein TLS. The modified and promoter-docked TLS inhibits the histone acetyltransferase activities of CREB binding protein and p300 inducing the silencing of cyclin D1 expression (Figure 4) [201]. A different co-activator activity mediated by lncRNAs is also evident in the regulation of Dlx genes, important modulators of neuronal development and patterning [202]. Dlx5-6 expression is regulated by two ultraconserved enhancers one of which is transcribed in a lncRNA, named Evf-2. Evf2 forms a stable complex with the homeodomain protein DLX-2 which in turn acts as a transcriptional enhancer of Dlx5-6 gene (Figure 4). In some cases, lncRNAs can also affect RNA polymerase activity by influencing the initiation complex in the choice of the promoter. For example, in humans, a ncRNA transcribed from an upstream region of the dihydrofolate reductase (DHFR) locus forms a triplex in the major

promoter of DHFR to prevent the binding of the transcriptional co-factor TFIID (Figure 4). This could be a widespread mechanism for controlling promoter usage as thousands of triplex structures exist in eukaryotic chromosomes. Recently, lncRNAs have also shown their tumorigenic potential by modulating the transcriptional program of p53 [203]. An 3kb lncRNAs, linc-RNA-p21, transcriptionally activated by p53, has been shown to collaborate with p53 in order to control the gene expression in response to DNA damage. Specifically, silencing of lincRNA-p21 derepresses the expression of hundred of genes which are also derepressed following p53 knockdown. It has also been discovered that lincRNA-p21 interacts with hnRNPK and this binding is essential for the modulation of p53 activity.

The final category of lncRNAs is represented by those molecules capable to generate the formation of compartmentalized nuclear organelles, subnuclear membraneless nuclear bodies whose function is relative unknown. One of them is represented by cell-cycle regulated nuclear foci, named paraspeckles. In addition to protein components, two lncRNAs, NEAT1 and Men epsilon, have been detected as essential part of the paraspeckles. While depletion of NEAT or Men epsilon disrupts the paraspeckles, their overexpression strongly increases their number. There is a number of different lncRNAs that localize to different nuclear regions [204]. Metastasis-associated lung adenocarcinoma transcript 1 (MALAT1) localizes to the splicing speckles, Xist and Kcnq1ot1 both, localize to the perinucleolar region during the S phase of the cell cycle, a class of repeat-associated lncRNAs (es SatIII) are associated to nuclear stress bodies which are produced on specific pericentromeric heterochromatic domains containing SatIII gene itself.

9. Conclusions

Alterations in microRNAs and other short or long non-coding RNA (ncRNA) are involved in the initiation, progression, and metastasis of human cancer. Over the last decade, a growing number of non-coding transcripts have been found to have roles in gene regulation and RNA processing. The most well known small non-coding RNAs are the microRNAs, but the network of long and short non-coding transcripts is complex and is likely to contain as yet unidentified classes of molecules that form transcriptional regulatory networks. The field of small and long non coding RNAs is rapidly advancing toward in vivo delivery for therapeutic purposes. Advanced molecular therapies aimed at downmodulating or upmodulating the level of a given miRNA in model organisms have been successfully established. RNA-based gene therapy can be used to treat cancer by using RNA or DNA molecules as therapy against the mRNA of genes involved in cancer pathogenesis or by directly targeting the ncRNAs that participate in pathogenesis. The use of miRNAs is still being evaluated preclinically; no clinical or toxicologic studies have been published but the future is promising. Kota and colleagues reported that systemic administration of this miRNA in a mouse model of HCC using adeno-associated virus (AAV) results in inhibition of cancer cell proliferation, induction of tumor-specific apoptosis, and dramatic protection from disease progression without toxicity (116). Recently, Pineau et al. (117) identified DNA damage-inducible transcript 4 (DDIT4), a modulator of the mTor pathway, as a bona fide target of miR-221. They introduced into liver cancer cells, by lipofection, LNA-modified oligonucleotides specifically designed for miR-221

(antimiR-221) and miR-222 (antimiR-222) knockdown. Treatment by antagomiRs, but not scrambled oligonucleotide, reduced cell growth in liver cancer cell lines that overexpressed miR-221 and miR-222 by 35% and 22%, respectively. Thus the use of synthetic inhibitors of miR-221 may prove to be a promising approach to liver cancer treatment (117). Despite recent progress in silencing of miRNAs in rodents, the development of effective and safe approaches for sequence-specific antagonism of miRNAs in vivo remains a significant scientific and therapeutic challenge. Recently, Elmen and collaborators (118) showed for the first time, that the simple systemic delivery of an unconjugated, PBS-formulated LNA-antimiR effectively antagonizes the liver-expressed miR-122 in nonhuman primates. Administration by intravenous injections of LNA-antimiR into African green monkeys resulted in the formation of stable heteroduplexes between the LNA-antimiR and miR-122, accompanied by depletion of mature miR-122 and dose-dependent lowering of plasma cholesterol. These findings demonstrate the utility of systemically administered LNA-antimiRs in exploring miRNA functions in primates and show the impressive potential of this strategy to overcome a major hurdle for clinical miRNA therapy. In conclusion, the discovery of small RNAs and their functions has revitalized the prospect of controlling expression of specific genes in vivo, with the ultimate hope of building a new class of gene-specific medical therapies. Just how significant are the ncRNAs? They appear to be doing something important and highly sophisticated; there are so many of them, their sequences are so highly conserved, their expression is tissue specific, and they have recognition sites on more than 30% of the entire transcriptome. It seems that ncRNAs were overlooked in the past simply because researchers were specifically looking for RNAs that code proteins. The above discussed data highlight that the complexity of genomic control operated by the ncRNAs is somewhat greater than previously imagined, and that they could represent a total new order of genomic control. In this scenario, understanding the precise roles of ncRNAs is a key challenge. The targeting of other ncRNAs, in addition to miRNAs, is still in its infancy, but new important developments are expected in this area. Therefore, small RNAs could become powerful therapeutic tools in the near future.

Author details

Gianpiero Di Leva and Michela Garofalo*

The Ohio State University, Department of Molecular Immunology, Virology and Medical Genetics. Columbus, OH, USA

10. References

- [1] Matsubara K, Okubo K. (1993). Identification of new genes by systematic analysis of cDNAs and database construction. *Curr Opin Biotechnol.* 4:672-677.
- [2] Liang F, Holt I, Pertea G, Karamycheva S, Salzberg SL, Quackenbush J. (2000). Gene index analysis of the human genome estimates approximately 120,000 genes. *Nat Genet.* 25:239-240.

* Corresponding Author

- [3] Lander ES, Linton LM, Birren B, Nusbaum C, Zody MC, Baldwin J, Devon K, Dewar K, Doyle M, FitzHugh W, Funke R, Gage D, Harris K, Heaford A, Howland J, Kann L, Lehoczky J, LeVine R, McEwan P, McKernan K, Meldrim J, Mesirov JP, Miranda C, Morris W, Naylor J, Raymond C, Rosetti M, Santos R, Sheridan A, Sougnez C, Stange-Thomann N, Stojanovic N, Subramanian A, Wyman D, Rogers J, Sulston J, Ainscough R, Beck S, Bentley D, Burton J, Clee C, Carter N, Coulson A, Deadman R, Deloukas P, Dunham A, Dunham I, Durbin R, French L, Grafham D, Gregory S, Hubbard T, Humphray S, Hunt A, Jones M, Lloyd C, McMurray A, Matthews L, Mercer S, Milne S, Mullikin JC, Mungall A, Plumb R, Ross M, Shownkeen R, Sims S, Waterston RH, Wilson RK, Hillier LW, McPherson JD, Marra MA, Mardis ER, Fulton LA, Chinwalla AT, Pepin KH, Gish WR, Chissoe SL, Wendl MC, Delehaunty KD, Miner TL, Delehaunty A, Kramer JB, Cook LL, Fulton RS, Johnson DL, Minx PJ, Clifton SW, Hawkins T, Branscomb E, Predki P, Richardson P, Wenning S, Slezak T, Doggett N, Cheng JF, Olsen A, Lucas S, Elkin C, Uberbacher E, Frazier M, Gibbs RA, Muzny DM, Scherer SE, Bouck JB, Sodergren EJ, Worley KC, Rives CM, Gorrell JH, Metzker ML, Naylor SL, Kucherlapati RS, Nelson DL, Weinstock GM, Sakaki Y, Fujiyama A, Hattori M, Yada T, Toyoda A, Itoh T, Kawagoe C, Watanabe H, Totoki Y, Taylor T, Weissenbach J, Heilig R, Saurin W, Artiguenave F, Brottier P, Bruls T, Pelletier E, Robert C, Wincker P, Smith DR, Doucette-Stamm L, Rubenfield M, Weinstock K, Lee HM, Dubois J, Rosenthal A, Platzer M, Nyakatura G, Taudien S, Rump A, Yang H, Yu J, Wang J, Huang G, Gu J, Hood L, Rowen L, Madan A, Qin S, Davis RW, Federspiel NA, Abola AP, Proctor MJ, Myers RM, Schmutz J, Dickson M, Grimwood J, Cox DR, Olson MV, Kaul R, Raymond C, Shimizu N, Kawasaki K, Minoshima S, Evans GA, Athanasiou M, Schultz R, Roe BA, Chen F, Pan H, Ramser J, Lehrach H, Reinhardt R, McCombie WR, de la Bastide M, Dedhia N, Blöcker H, Hornischer K, Nordsiek G, Agarwala R, Aravind L, Bailey JA, Bateman A, Batzoglu S, Birney E, Bork P, Brown DG, Burge CB, Cerutti L, Chen HC, Church D, Clamp M, Copley RR, Doerks T, Eddy SR, Eichler EE, Furey TS, Galagan J, Gilbert JG, Harmon C, Hayashizaki Y, Haussler D, Hermjakob H, Hokamp K, Jang W, Johnson LS, Jones TA, Kasif S, Kasprzyk A, Kennedy S, Kent WJ, Kitts P, Koonin EV, Korf I, Kulp D, Lancet D, Lowe TM, McLysaght A, Mikkelsen T, Moran JV, Mulder N, Pollara VJ, Ponting CP, Schuler G, Schultz J, Slater G, Smit AF, Stupka E, Szustakowski J, Thierry-Mieg D, Thierry-Mieg J, Wagner L, Wallis J, Wheeler R, Williams A, Wolf YI, Wolfe KH, Yang SP, Yeh RF, Collins F, Guyer MS, Peterson J, Felsenfeld A, Wetterstrand KA, Patrinos A, Morgan MJ, de Jong P, Catanese JJ, Osoegawa K, Shizuya H, Choi S, Chen YJ. (2001). International Human Genome Sequencing Consortium. Initial sequencing and analysis of the human genome. *Nature*. 409:860-921.
- [4] Guttman M, Amit I, Garber M, French C, Lin MF, Feldser D, Huarte M, Zuk O, Carey BW, Cassady JP, Cabili MN, Jaenisch R, Mikkelsen TS, Jacks T, Hacohen N, Bernstein BE, Kellis M, Regev A, Rinn JL, Lander ES. (2009). Chromatin signature reveals over a thousand highly conserved large non-coding RNAs in mammals. *Nature*. 458:223–227
- [5] Guttman M, Garber M, Levin JZ, Donaghey J, Robinson J, Adiconis X, Fan L, Koziol MJ, Gnirke A, Nusbaum C, Rinn JL, Lander ES, Regev A. (2010). Ab initio reconstruction of

- cell type-specific transcriptomes in mouse reveals the conserved multi-exonic structure of lincRNAs. *Nat. Biotechnol.* 28:503–510.
- [6] Marques AC, Ponting CP. (2009). Catalogues of mammalian long noncoding RNAs: modest conservation and incompleteness. *Genome Biol.* 10:R124.
 - [7] Cech TR. (2009). Crawling out of the RNA world. *Cell.* 136:599–602.
 - [8] Kapranov P, Cheng J, Dike S, Nix DA, Duttagupta R, Willingham AT, Stadler PF, Hertel J, Hackermüller J, Hofacker IL, Bell I, Cheung E, Drenkow J, Dumais E, Patel S, Helt G, Ganesh M, Ghosh S, Piccolboni A, Sementchenko V, Tammanna H, Gingeras TR. (2007). RNA maps reveal new RNA classes and a possible function for pervasive transcription. *Science.* 316:1484–1488.
 - [9] Crick FH, Barnett L, Brenner S, Watts-Tobin RJ. (1961). General nature of the genetic code for proteins. *Nature.* 192:1227–1232.
 - [10] Mercer TR, Dinger ME, Mattick JS. (2009) Long non-coding RNAs insight into functions. *Nature Rev. Genet.* 10:155–159.
 - [11] He L, Hannon GJ. (2004). MicroRNAs: small RNAs with a big role in gene regulation. *Nature Rev. Genet.* 5:522–531.
 - [12] Mendell JT. (2005). MicroRNAs: critical regulators of development, cellular physiology and malignancy. *Cell Cycle* 4:1179–1184.
 - [13] Esquela-Kerscher A, Slack FJ. (2006). OncomiRs — microRNAs with a role in cancer. *Nature Rev. Cancer* 6:259–269.
 - [14] Hammond SM. (2007). MicroRNAs as tumor suppressors. *Nature Genet.* 39:582–583.
 - [15] Croce CM. (2009). Causes and consequences of microRNA dysregulation in cancer. *Nature Rev. Genet.* 10:704–714.
 - [16] Nicoloso M S, Spizzo R, Shimizu M, Rossi S, Calin GA. (2009). MicroRNAs — the micro steering wheel of tumour metastases. *Nature Rev. Cancer.* 9:293–302.
 - [17] Carninci P, Kasukawa T, Katayama S, Gough J, Frith MC, Maeda N, Oyama R, Ravasi T, Lenhard B, Wells C, Kodzius R, Shimokawa K, Bajic VB, Brenner SE, Batalov S, Forrest AR, Zavolan M, Davis MJ, Wilming LG, Aidinis V, Allen JE, Ambesi-Impimombato A, Apweiler R, Aturaliya RN, Bailey TL, Bansal M, Baxter L, Beisel KW, Bersano T, Bono H, Chalk AM, Chiu KP, Choudhary V, Christoffels A, Clutterbuck DR, Crowe ML, Dalla E, Dalrymple BP, de Bono B, Della Gatta G, di Bernardo D, Down T, Engstrom P, Fagiolini M, Faulkner G, Fletcher CF, Fukushima T, Furuno M, Futaki S, Gariboldi M, Georgii-Hemming P, Gingeras TR, Gojobori T, Green RE, Gustincich S, Harbers M, Hayashi Y, Hensch TK, Hirokawa N, Hill D, Huminiecki L, Iacono M, Ikeo K, Iwama A, Ishikawa T, Jakt M, Kanapin A, Katoh M, Kawasawa Y, Kelso J, Kitamura H, Kitano H, Kollias G, Krishnan SP, Kruger A, Kummerfeld SK, Kurochkin IV, Lareau LF, Lazarevic D, Lipovich L, Liu J, Liuni S, McWilliam S, Madan Babu M, Madera M, Marchionni L, Matsuda H, Matsuzawa S, Miki H, Mignone F, Miyake S, Morris K, Mottagui-Tabar S, Mulder N, Nakano N, Nakauchi H, Ng P, Nilsson R, Nishiguchi S, Nishikawa S, Nori F, Ohara O, Okazaki Y, Orlando V, Pang KC, Pavan WJ, Pavesi G, Pesole G, Petrovsky N, Piazza S, Reed J, Reid JF, Ring BZ, Ringwald M, Rost B, Ruan Y, Salzberg SL, Sandelin A, Schneider C, Schönbach C, Sekiguchi K, Semple CA, Seno S, Sessa L, Sheng Y, Shibata Y, Shimada H, Shimada K, Silva D, Sinclair B, Sperling S, Stupka E, Sugiura K, Sultana R, Takenaka Y, Taki K, Tammoja K, Tan SL, Tang S, Taylor MS, Tegner J,

- Teichmann SA, Ueda HR, van Nimwegen E, Verardo R, Wei CL, Yagi K, Yamanishi H, Zabarovsky E, Zhu S, Zimmer A, Hide W, Bult C, Grimmond SM, Teasdale RD, Liu ET, Brusic V, Quackenbush J, Wahlestedt C, Mattick JS, Hume DA, Kai C, Sasaki D, Tomaru Y, Fukuda S, Kanamori-Katayama M, Suzuki M, Aoki J, Arakawa T, Iida J, Imamura K, Itoh M, Kato T, Kawaji H, Kawagashira N, Kawashima T, Kojima M, Kondo S, Konno H, Nakano K, Ninomiya N, Nishio T, Okada M, Plessy C, Shibata K, Shiraki T, Suzuki S, Tagami M, Waki K, Watahiki A, Okamura-Oho Y, Suzuki H, Kawai J, Hayashizaki Y; FANTOM Consortium; RIKEN Genome Exploration Research Group and Genome Science Group (Genome Network Project Core Group).(2005). The transcriptional landscape of the mammalian genome. *Science*. 309:1559–1563.
- [18] Gibb EA, Brown CJ, Lam WL. (2011). The functional role of long non-coding RNA in human carcinomas. *Mol Cancer* 10:38.
- [19] Taft RJ, Pang KC, Mercer TR, Dinger M, Mattick JS. (2010) Non-coding RNAs: regulators of disease. *J Pathol*.220:126-39.
- [20] Di Leva G, Croce CM. (2010). Roles of small RNAs in tumor formation. *Trends Mol Med*.16:257-67.
- [21] Garofalo M, Croce CM. (2011). microRNAs: Master regulators as potential therapeutics in cancer. *Annu Rev Pharmacol Toxicol*. 51:25-43.
- [22] ENCODE Project Consortium, Birney E, Stamatoyannopoulos JA, Dutta A, Guigó R, Gingeras TR, Margulies EH, Weng Z, Snyder M, Dermitzakis ET, Thurman RE, Kuehn MS, Taylor CM, Neph S, Koch CM, Asthana S, Malhotra A, Adzhubei I, Greenbaum JA, Andrews RM, Flicek P, Boyle PJ, Cao H, Carter NP, Clelland GK, Davis S, Day N, Dhami P, Dillon SC, Dorschner MO, Fiegler H, Giresi PG, Goldy J, Hawrylycz M, Haydock A, Humbert R, James KD, Johnson BE, Johnson EM, Frum TT, Rosenzweig ER, Karnani N, Lee K, Lefebvre GC, Navas PA, Neri F, Parker SC, Sabo PJ, Sandstrom R, Shafer A, Vetrie D, Weaver M, Wilcox S, Yu M, Collins FS, Dekker J, Lieb JD, Tullius TD, Crawford GE, Sunyaev S, Noble WS, Dunham I, Denoeud F, Reymond A, Kapranov P, Rozowsky J, Zheng D, Castelo R, Frankish A, Harrow J, Ghosh S, Sandelin A, Hofacker IL, Baertsch R, Keefe D, Dike S, Cheng J, Hirsch HA, Sekinger EA, Lagarde J, Abril JF, Shahab A, Flamm C, Fried C, Hackermüller J, Hertel J, Lindemeyer M, Missal K, Tanzer A, Washietl S, Korb J, Emanuelsson O, Pedersen JS, Holroyd N, Taylor R, Swarbreck D, Matthews N, Dickson MC, Thomas DJ, Weirauch MT, Gilbert J, Drenkow J, Bell I, Zhao X, Srinivasan KG, Sung WK, Ooi HS, Chiu KP, Foissac S, Alioto T, Brent M, Pachter L, Tress ML, Valencia A, Choo SW, Choo CY, Ucla C, Manzano C, Wyss C, Cheung E, Clark TG, Brown JB, Ganesh M, Patel S, Tammanna H, Chrast J, Henriksen CN, Kai C, Kawai J, Nagalakshmi U, Wu J, Lian Z, Lian J, Newburger P, Zhang X, Bickel P, Mattick JS, Carninci P, Hayashizaki Y, Weissman S, Hubbard T, Myers RM, Rogers J, Stadler PF, Lowe TM, Wei CL, Ruan Y, Struhl K, Gerstein M, Antonarakis SE, Fu Y, Green ED, Karaöz U, Siepel A, Taylor J, Liefer LA, Wetterstrand KA, Good PJ, Feingold EA, Guyer MS, Cooper GM, Asimenos G, Dewey CN, Hou M, Nikolaev S, Montoya-Burgos JI, Löytynoja A, Whelan S, Pardi F, Massingham T, Huang H, Zhang NR, Holmes I, Mullikin JC, Ureta-Vidal A, Paten B, Seringhaus M, Church D, Rosenbloom K, Kent WJ, Stone EA; NISC Comparative Sequencing Program; Baylor College of Medicine Human Genome Sequencing Center; Washington University

- Genome Sequencing Center; Broad Institute; Children's Hospital Oakland Research Institute, Batzoglou S, Goldman N, Hardison RC, Haussler D, Miller W, Sidow A, Trinklein ND, Zhang ZD, Barrera L, Stuart R, King DC, Ameur A, Enroth S, Bieda MC, Kim J, Bhinge AA, Jiang N, Liu J, Yao F, Vega VB, Lee CW, Ng P, Shahab A, Yang A, Moqtaderi Z, Zhu Z, Xu X, Squazzo S, Oberley MJ, Inman D, Singer MA, Richmond TA, Munn KJ, Rada-Iglesias A, Wallerman O, Komorowski J, Fowler JC, Couttet P, Bruce AW, Dovey OM, Ellis PD, Langford CF, Nix DA, Euskirchen G, Hartman S, Urban AE, Kraus P, Van Calcar S, Heintzman N, Kim TH, Wang K, Qu C, Hon G, Luna R, Glass CK, Rosenfeld MG, Aldred SF, Cooper SJ, Halees A, Lin JM, Shulha HP, Zhang X, Xu M, Haidar JN, Yu Y, Ruan Y, Iyer VR, Green RD, Wadelius C, Farnham PJ, Ren B, Harte RA, Hinrichs AS, Trumbower H, Clawson H, Hillman-Jackson J, Zweig AS, Smith K, Thakkapallayil A, Barber G, Kuhn RM, Karolchik D, Armengol L, Bird CP, de Bakker PI, Kern AD, Lopez-Bigas N, Martin JD, Stranger BE, Woodroffe A, Davydov E, Dimas A, Eyraas E, Hallgrimsdóttir IB, Huppert J, Zody MC, Abecasis GR, Estivill X, Bouffard GG, Guan X, Hansen NF, Idol JR, Maduro VV, Maskeri B, McDowell JC, Park M, Thomas PJ, Young AC, Blakesley RW, Muzny DM, Sodergren E, Wheeler DA, Worley KC, Jiang H, Weinstock GM, Gibbs RA, Graves T, Fulton R, Mardis ER, Wilson RK, Clamp M, Cuff J, Gnerre S, Jaffe DB, Chang JL, Lindblad-Toh K, Lander ES, Koriabine M, Nefedov M, Osoegawa K, Yoshinaga Y, Zhu B, de Jong PJ. Identification and analysis of functional elements in 1% of the human genome by the ENCODE pilot project.(2007). *Nature*.447:799-816.
- [23] Siomi MC, Sato K, Pezic D, Aravin AA. (2011). PIWI-interacting small RNAs: the vanguard of genome defence. *Nat Rev Mol Cell Biol*.12:246-58.
- [24] Faulkner GJ, Kimura Y, Daub CO, Wani S, Plessy C, Irvine KM, Schroder K, Cloonan N, Steptoe AL, Lassmann T, Waki K, Hornig N, Arakawa T, Takahashi H, Kawai J, Forrest AR, Suzuki H, Hayashizaki Y, Hume DA, Orlando V, Grimmond SM, Carninci P. (2009). The regulated retrotransposon transcriptome of mammalian cells. *Nat Genet*. 41:563-71.
- [25] Huarte M, Rinn JL. (2010). Large non-coding RNAs: missing links in cancer? *Hum Mol Genet*. 19:R152-61.
- [26] Pauli A, Rinn JL, Schier AF. (2011). Non-coding RNAs as regulators of embryogenesis. *Nat Rev Genet*.12:136–49.
- [27] Calin GA, Liu CG, Ferracin M, Hyslop T, Spizzo R, Sevignani C, Fabbri M, Cimmino A, Lee EJ, Wojcik SE, Shimizu M, Tili E, Rossi S, Taccioli C, Pichiorri F, Liu X, Zupo S, Herlea V, Gramantieri L, Lanza G, Alder H, Rassenti L, Volinia S, Schmittgen TD, Kipps TJ, Negrini M, Croce CM. (2007). Ultraconserved regions encoding ncRNAs are altered in human leukemias and carcinomas. *Cancer Cell*. 12:215-29.
- [28] Poliseno L, Salmena L, Zhang J, Carver B, Haveman WJ, Pandolfi PP. (2010). A coding-independent function of gene and pseudogene mRNAs regulates tumour biology. *Nature*. 465:1033-8.
- [29] He Y, Vogelstein B, Velculescu VE, Papadopoulos N, Kinzler KW. (2008). The antisense transcriptomes of human cells. *Science*. 322:1855-7.
- [30] Kim TK, Hemberg M, Gray JM, Costa AM, Bear DM, Wu J, Harmin DA, Laptewicz M, Barbara-Haley K, Kuersten S, Markenscoff-Papadimitriou E, Kuhl D, Bito H, Worley PF,

- Kreiman G, Greenberg ME. (2010). Widespread transcription at neuronal activity-regulated enhancers. *Nature*. 465:182-7.
- [31] Lee RC, Feinbaum RL, Ambros V. (1993). The *C. elegans* heterochronic gene *lin-4* encodes small RNAs with antisense complementarity to *lin-14*. *Cell*. 75:843-54.
- [32] Kim VN. 2005. MicroRNA biogenesis: coordinated cropping and dicing. *Nat. Rev. Mol. Cell Biol.* 6:376-85.
- [33] Rodriguez A, Griffiths-Jones S, Ashurst JL, Bradley A. (2004). Identification of mammalian microRNA host genes and transcription units. *Genome Res.* 14:1902-10.
- [34] Lagos-Quintana M, Rauhut R, Yalcin A, Meyer J, Lendeckel W, Tuschl T. (2002). Identification of tissue-specific microRNAs from mouse. *Curr Biol.* 12: 735-739.
- [35] Altuvia Y, Landgraf P, Lithwick G, Elefant N, Pfeffer S, Aravin A, Brownstein MJ, Tuschl T, Margalit H. (2005). Clustering and conservation patterns of human microRNAs. *Nucleic Acids Res* 33: 2697-2706.
- [36] Borchert GM, Lanier W, Davidson BL. (2006). RNA polymerase III transcribes human microRNAs. *Nat Struct Mol Biol.* 13: 1097-1101.
- [37] Han J, Lee Y, Yeom KH, Nam JW, Heo I, Rhee JK, Sohn SY, Cho Y, Zhang BT, Kim VN. (2006). Molecular basis for the recognition of primary microRNAs by the Drosha-DGCR8 complex. *Cell* 125: 887-901.
- [38] Saetrom P, Snove O, Nedland M, Grunfeld TB, Lin Y, Bass MB, Cannon JR. (2006). Conserved microRNA characteristics in mammals. *Oligonucleotides.* 16:115-44.
- [39] Gregory RI, Yan KP, Amuthan G, Chendrimada T, Doratotaj B, Cooch N, Shiekhattar R. (2004). The Microprocessor complex mediates the genesis of microRNAs. *Nature* 432:235-40.
- [40] Han J, Lee Y, Yeom KH, Kim YK, Jin H, Kim VN. (2004). The Drosha-DGCR8 complex in primary microRNA processing. *Genes Dev.* 18:3016-27.
- [41] Yi R, Qin Y, Macara IG, Cullen BR. (2003). Exportin-5 mediates the nuclear export of pre-microRNAs and short hairpin RNAs. *Genes Dev.* 17:3011-3016.
- [42] Lund E, Guttinger S, Calado A, Dahlberg JE, Kutay U. (2004). Nuclear export of microRNA precursors. *Science.* 303:95-98.
- [43] Bohnsack MT, Czaplinski K, Gorlich D. (2004). Exportin 5 is a RanGTP-dependent dsRNA-binding protein that mediates nuclear export of pre-miRNAs. *RNA.* 10:185-191.
- [44] Maniatakis E, Mourelatos Z. (2005). A human, ATP-independent, RISC assembly machine fueled by pre-miRNA. *Genes Dev.* 19:2979-2990.
- [45] Lee Y, Hur I, Park SY, Kim YK, Suh MR, Kim VN. (2006). The role of PACT in the RNA silencing pathway. *EMBO J.* 25:522-532.
- [46] Haase AD, Jaskiewicz L, Zhang H, Laine S, Sack R, Gatignol A, Filipowicz W. (2005). TRBP, a regulator of cellular PKR and HIV-1 virus expression, interacts with Dicer and functions in RNA silencing. *EMBO Rep* 6: 961-967.
- [47] Gregory RI, Chendrimada TP, Cooch N, Shiekhattar R. (2005). Human RISC couples microRNA biogenesis and posttranscriptional gene silencing. *Cell.* 123:631-640.
- [48] Forstemann K, Tomari Y, Du T, Vagin VV, Denli AM, Bratu DP, Klattenhoff C, Theurkauf WE, Zamore PD. (2005). Normal microRNA maturation and germ-line stem cell maintenance requires Loquacious, a double-stranded RNA-binding domain protein. *PLoS Biol.* 3:e236.

- [49] Chendrimada TP, Gregory RI, Kumaraswamy E, Norman J, Cooch N, Nishikura K, Nishikura K, Shiekhattar R. (2005). TRBP recruits the Dicer complex to Ago2 for microRNA processing and gene silencing. *Nature*. 436: 740–744.
- [50] Castanotto D, Sakurai K, Lingeman R, Li H, Shively L, Aagaard L, Soifer H, Gatignol A, Riggs A, Rossi JJ. (2007). Combinatorial delivery of small interfering RNAs reduces RNAi efficacy by selective incorporation into RISC. *Nucleic Acids Res*. 35:5154–64.
- [51] Rand TA, Petersen S, Du F, Wang X. (2005). Argonaute2 cleaves the anti-guide strand of siRNA during RISC activation. *Cell*. 123:621–29.
- [52] Leuschner PJ, Ameres SL, Kueng S, Martinez J. (2006). Cleavage of the siRNA passenger strand during RISC assembly in human cells. *EMBO Rep*. 7:314–20.
- [53] Martinez J, Patkaniowska A, Urlaub H, Luhrmann R, Tuschl T. (2002). Single-stranded antisense siRNAs guide target RNA cleavage in RNAi. *Cell*. 110:563–74.
- [54] Zamore PD, Tuschl T, Sharp PA, Bartel DP. (2000). RNAi: double-stranded RNA directs the ATP-dependent cleavage of mRNA at 21 to 23 nucleotide intervals. *Cell*. 101: 25–33.
- [55] Yekta S, Shih IH, Bartel DP. (2004). MicroRNA-directed cleavage of HOXB8 mRNA. *Science*. 304: 594–596.
- [56] Wu L, Fan J, Belasco JG. (2006). MicroRNAs direct rapid deadenylation of mRNA. *Proc Natl Acad Sci USA*. 103: 4034–4039.
- [57] Bagga S, Bracht J, Hunter S, Massirer K, Holtz J, Eachus R, Pasquinelli AE. (2005). Regulation by let-7 and lin-4 miRNAs results in target mRNA degradation. *Cell*. 122: 553–563.
- [58] Reinhart BJ, Slack FJ, Basson M, Pasquinelli AE, Bettinger JC, Rougvie AE, Horvitz HR, Ruvkun G. (2000). The 21-nucleotide let-7 RNA regulates developmental timing in *Caenorhabditis elegans*. *Nature* 403: 901–906.
- [59] Olsen PH, Ambros V. (1999). The lin-4 regulatory RNA controls developmental timing in *Caenorhabditis elegans* by blocking LIN-14 protein synthesis after the initiation of translation. *Dev Biol*. 216: 671–680.
- [60] Pillai RS, Bhattacharyya SN, Filipowicz W. (2007). Repression of protein synthesis by miRNAs: how many mechanisms? *Trends Cell Biol*. 17:118–26.
- [61] Garzon R, Fabbri M, Cimmino A, Calin GA, Croce CM. (2006). MicroRNA expression and function in cancer. *Trends Mol. Med*. 12:580–587.
- [62] Wu W, Sun M, Zou GM, Chen J. (2007). MicroRNA and cancer: Current status and prospective. *Int. J. Cancer*. 120: 953–960.
- [63] Boettger T, Braun T. (2012). A New Level of Complexity: The Role of MicroRNAs in Cardiovascular Development. *Circ Res*. 110:1000–1013.
- [64] Han R, Kan Q, Sun Y, Wang S, Zhang G, Peng T, Jia Y. (2012). MiR-9 promotes the neural differentiation of mouse bone marrow mesenchymal stem cells via targeting zinc finger protein 521. *Neurosci Lett*. Ahead of print.
- [65] D'Urso PI, D'Urso OF, Storelli C, Mallardo M, Gianfreda CD, Montinaro A, Cimmino A, Pietro C, Marsigliante S. (2012). miR-155 is up-regulated in primary and secondary glioblastoma and promotes tumour growth by inhibiting GABA receptors. *Int J Oncol*. doi: 10.3892/ijo.2012.1420.
- [66] Garofalo M, Romano G, Di Leva G, Nuovo G, Jeon YJ, Ngankea A, Sun J, Lovat F, Alder H, Condorelli G, Engelman JA, Ono M, Rho JK, Cascione L, Volinia S, Nephew KP,

- Croce CM. (2011). EGFR and MET receptor tyrosine kinase-altered microRNA expression induces tumorigenesis and gefitinib resistance in lung cancers. *Nat Med.*18:74-82.
- [67] Frank D, Gantenberg J, Boomgaarden I, Kuhn C, Will R, Jarr KU, Eden M, Kramer K, Luedde M, Mairbäurl H, Katus HA, Frey N. (2012). MicroRNA-20a inhibits stress-induced cardiomyocyte apoptosis involving its novel target Egl3/PHD3. *J Mol Cell Cardiol.* 52:711-717.
- [68] Calin GA, Dumitru CD, Shimizu M, Bichi R, Zupo S, Noch E, Aldler H, Rattan S, Keating M, Rai K, Rassenti L, Kipps T, Negrini M, Bullrich F, Croce CM. (2002). Frequent deletions and down-regulation of micro-RNA genes miR15 and miR16 at 13q14 in chronic lymphocytic leukemia. *Proc. Natl. Acad. Sci. USA.* 99:15524–29
- [69] Calin GA, Croce CM. (2006). MicroRNA signatures in human cancers. *Nat Rev Cancer.* 6:857–866.
- [70] Cai X, Hagedorn CH, Cullen BR. (2004). Human microRNAs are processed from capped, polyadenylated transcripts that can also function as mRNAs. *RNA.* 10:1957–1966.
- [71] Cai X, Hagedorn CH, Cullen BR. (2004). Human microRNAs are processed from capped, polyadenylated transcripts that can also function as mRNAs. *RNA.* 10:1957–1966.
- [72] Chan JA, Krichevsky AM, Kosik KS. 2005. MicroRNA-21 is an antiapoptotic factor in human glioblastoma cells. *Cancer Res.* 65:6029–6033.
- [73] Zhu S, Wu H, Wu F, Nie D, Sheng S, Mo YY. (2008). MicroRNA-21 targets tumor suppressor genes in invasion and metastasis. *Cell Res.* 18:350–359.
- [74] Zhu Q, Wang Z, Hu Y, Li J, Li X, Zhou L, Huang Y. (2012). miR-21 promotes migration and invasion by the miR-21-PDCD4-AP-1 feedback loop in human hepatocellular carcinoma. *Oncol Rep.*27:1660-1668.
- [75] Medina PP, Nolde M, Slack FJ. (2010). OncomiR addiction in an in vivo model of microRNA-21-induced pre-B-cell lymphoma. *Nature.* 467:86-90.
- [76] Metzler M, Wilda M, Busch K, Viehmann S, Borkhardt A. (2004). High expression of precursor microRNA-155/BIC RNA in children with Burkitt lymphoma. *Genes Chromosom. Cancer.* 39:167–169.
- [77] Kluiver J, Poppema S, de Jong D, Blokzijl T, Harms G, Jacobs S, Kroesen BJ, van den Berg A. (2005). BIC and miR-155 are highly expressed in Hodgkin, primary mediastinal and diffuse large B cell lymphomas. *J. Pathol.* 207:243–249.
- [78] Calin GA, Ferracin M, Cimmino A, Di Leva G, Shimizu M, Wojcik SE, Iorio MV, Visone R, Sever NI, Fabbri M, Iuliano R, Palumbo T, Pichiorri F, Roldo C, Garzon R, Sevignani C, Rassenti L, Alder H, Volinia S, Liu CG, Kipps TJ, Negrini M, Croce CM. (2005). A microRNA signature associated with prognosis and progression in chronic lymphocytic leukemia. *N. Engl. J. Med.* 353:1793–1801.
- [79] Garzon R, Volinia S, Liu CG, Fernandez-Cymering C, Palumbo T, Pichiorri F, Fabbri M, Coombes K, Alder H, Nakamura T, Flomenberg N, Marcucci G, Calin GA, Kornblau SM, Kantarjian H, Bloomfield CD, Andreeff M, Croce CM. (2008). MicroRNA signatures associated with cytogenetics and prognosis in acute myeloid leukemia. *Blood.* 111:3183–3189.

- [80] Volinia S, Calin GA, Liu CG, Ambs S, Cimmino A, Petrocca F, Visone R, Iorio M, Roldo C, Ferracin M, Prueitt RL, Yanaihara N, Lanza G, Scarpa A, Vecchione A, Negrini M, Harris CC, Croce CM. (2006). A microRNA expression signature of human solid tumors defines cancer gene targets. *Proc. Natl. Acad. Sci. USA.* 103:2257–2261.
- [81] Greither T, Grochola LF, Udelnow A, Lautenschlager C, Würl P, Taubert H. (2010). Elevated expression of microRNAs 155, 203, 210 and 222 in pancreatic tumors is associated with poorer survival. *Int. J. Cancer.* 126:73–80.
- [82] Costinean S, Zanesi N, Pekarsky Y, Tili E, Volinia S, Heerema N, Croce CM. (2006). Pre-B cell proliferation and lymphoblastic leukemia/high-grade lymphoma in E(mu)-miR155 transgenic mice. *Proc Natl Acad Sci U S A.* 103:7024–7029.
- [83] Pallante P, Visone R, Ferracin M, Ferraro A, Berlingieri MT, Troncone G, Chiappetta G, Liu CG, Santoro M, Negrini M, Croce CM, Fusco A. (2006). MicroRNA deregulation in human thyroid papillary carcinomas. *Endocr. Relat. Cancer.* 13:497–508.
- [84] Fornari F, Gramantieri L, Ferracin M, Veronese A, Sabbioni S, Calin GA, Grazi GL, Giovannini C, Croce CM, Bolondi L, Negrini M. (2008). MiR-221 control CDKN1C/p57 and CDKN1B/p27 expression in human hepatocellular carcinoma. *Oncogene.* 27:5651–5661.
- [85] Di Leva G, Gasparini P, Piovan C, Nganheu A, Garofalo M, Taccioli C, Iorio MV, Li M, Volinia S, Alder H, Nakamura T, Nuovo G, Liu Y, Nephew KP, Croce CM. (2010). MicroRNA cluster 221–222 and estrogen receptor α interactions in breast cancer. *J. Natl. Cancer Inst.* 102:706–721.
- [86] Felicetti F, Errico MC, Bottero L, Segnalini P, Stoppacciaro A, Biffoni M, Felli N, Mattia G, Petrini M, Colombo MP, Peschle C, Carè A. (2008). The promyelocytic leukemia zinc finger-microRNA-221/-222 pathway controls melanoma progression through multiple oncogenic mechanisms. *Cancer Res.* 68:2745–2754.
- [87] le Sage C, Nagel R, Egan DA, Schrier M, Mesman E, Mangiola A, Anile C, Maira G, Mercatelli N, Ciafrè SA, Farace MG, Agami R. (2007). Regulation of the p27Kip1 tumor suppressor by miR-221 and miR-222 promotes cancer cell proliferation. *EMBO J.* 26:3699–3708.
- [88] Garofalo M, Quintavalle C, Di Leva G, Zanca C, Romano G, Taccioli C, Liu CG, Croce CM, Condorelli G. (2008). MicroRNA signatures of TRAIL resistance in human non-small cell lung cancer. *Oncogene.* 27:3845–3855.
- [89] Garofalo M, Di Leva G, Romano G, Nuovo G, Suh SS, Nganheu A, Taccioli C, Pichiorri F, Alder H, Secchiero P, Gasparini P, Gonelli A, Costinean S, Acunzo M, Condorelli G, Croce CM. (2009). miR-221&222 regulate TRAIL resistance and enhance tumorigenicity through PTEN and TIMP3 downregulation. *Cancer Cell.* 16:498–509.
- [90] Petrocca F, Visone R, Onelli MR, Shah MH, Nicoloso MS, de Martino I, Iliopoulos D, Pillozzi E, Liu CG, Negrini M, Cavazzini L, Volinia S, Alder H, Ruco LP, Baldassarre G, Croce CM, Vecchione A. (2008). E2F1-regulated microRNAs impair TGF β -dependent cell-cycle arrest and apoptosis in gastric cancer. *Cancer Cell.* 13:272–286.
- [91] Ota A, Tagawa H, Karnan S, Tsuzuki S, Karpas A, Kira S, Yoshida Y, Seto M. (2004). Identification and characterization of a novel gene, C13orf25, as a target for 13q31-q32 amplification in malignant lymphoma. *Cancer Res.* 64:3087–3095.

- [92] Hayashita Y, Osada H, Tatematsu Y, Yamada H, Yanagisawa K, Tomida S, Yatabe Y, Kawahara K, Sekido Y, Takahashi T. (2005). A polycistronic microRNA cluster, miR-17-92, is overexpressed in human lung cancers and enhances cell proliferation. *Cancer Res.* 65:9628–9632.
- [93] Ventura A, Young AG, Winslow MM, Lintault L, Meissner A, Erkeland SJ, Newman J, Bronson RT, Crowley D, Stone JR, Jaenisch R, Sharp PA, Jacks T. (2008). Targeted deletion reveals essential and overlapping functions of the miR-17 through 92 family of miRNA clusters. *Cell.*132:875-886.
- [94] Park JK, Lee EJ, Esau C, Schmittgen TD. (2009). Antisense inhibition of microRNA-21 or -221 arrests cell cycle, induces apoptosis, and sensitizes the effects of gemcitabine in pancreatic adenocarcinoma. *Pancreas.* 38:e190–199.
- [95] Inomata M, Tagawa H, Guo YM, Kameoka Y, Takahashi N, Sawada K. (2009). MicroRNA-17-92 downregulates expression of distinct targets in different B-cell lymphoma subtypes. *Blood.* 113:396–402.
- [96] Wang PY, Li YJ, Zhang S, Li ZL, Yue Z, Xie N, Xie SY. (2010). Regulating A549 cells growth by ASO inhibiting miRNA expression. *Mol. Cell. Biochem.* 339:163–171.
- [97] Bandi N, Zbinden S, Gugger M, Arnold M, Kocher V, Hasan L, Kappeler A, Brunner T, Vassella E. (2009). miR-15a and miR-16 are implicated in cell cycle regulation in a Rb-dependent manner and are frequently deleted or down-regulated in non-small cell lung cancer. *Cancer Res.* 69:5553–5559.
- [98] Cimmino A, Calin GA, Fabbri M, Iorio MV, Ferracin M, Shimizu M, Wojcik SE, Aqeilan RI, Zupo S, Dono M, Rassenti L, Alder H, Volinia S, Liu CG, Kipps TJ, Negrini M, Croce CM. (2005). miR-15 and miR-16 induce apoptosis by targeting BCL2. *Proc. Natl. Acad. Sci. USA.*102:13944–13949.
- [99] Klein U, Lia M, Crespo M, Siegel R, Shen Q, Mo T, Ambesi-Impiombato A, Califano A, Migliazza A, Bhagat G, Dalla-Favera R. (2010). The DLEU2/miR-15a/16-1 cluster controls B cell proliferation and its deletion leads to chronic lymphocytic leukemia. *Cancer Cell* 17:28–40.
- [100] Bonci D, Coppola V, Musumeci M, Addario A, Giuffrida R, Memeo L, D'Urso L, Pagliuca A, Biffoni M, Labbaye C, Bartucci M, Muto G, Peschle C, De Maria R. (2008). The miR-15a–miR-16-1 cluster controls prostate cancer by targeting multiple oncogenic activities. *Nat. Med.* 14:1271–1277.
- [101] He X, He L, Hannon GJ. (2007). The guardian's little helper: microRNAs in the p53 tumor suppressor network. *Cancer Res.*67:11099-11101.
- [102] He L, He X, Lim LP, de Stanchina E, Xuan Z, Liang Y, Xue W, Zender L, Magnus J, Ridzon D, Jackson AL, Linsley PS, Chen C, Lowe SW, Cleary MA, Hannon GJ. (2007). A microRNA component of the p53 tumour suppressor network. *Nature.* 447:1130-1134.
- [103] Cole KA, Attiyeh EF, Mosse YP, Laquaglia MJ, Diskin SJ, Brodeur GM, Maris JM. (2008). A functional screen identifies miR-34a as a candidate neuroblastoma tumor suppressor gene. *Mol. Cancer Res.* 6:735–742.
- [104] Siemens H, Jackstadt R, Hüntten S, Kaller M, Menssen A, Götz U, Hermeking H. (2011). miR-34 and SNAIL form a double-negative feedback loop to regulate epithelial-mesenchymal transitions. *Cell Cycle.*10:4256-4271.

- [105] Tsai WC, Hsu PW, Lai TC, Chau GY, Lin CW, Chen CM, Lin CD, Liao YL, Wang JL, Chau YP, Hsu MT, Hsiao M, Huang HD, Tsou AP. (2009). MicroRNA-122, a tumor suppressor microRNA that regulates intrahepatic metastasis of hepatocellular carcinoma. *Hepatology*. 49:1571-1582.
- [106] Xu Y, Xia F, Ma L, Shan J, Shen J, Yang Z, Liu J, Cui Y, Bian X, Bie P, Qian C. (2011). MicroRNA-122 sensitizes HCC cancer cells to adriamycin and vincristine through modulating expression of MDR and inducing cell cycle arrest. *Cancer Lett*. 310:160-169.
- [107] Xu J, Zhu X, Wu L, Yang R, Yang Z, Wang Q, Wu F. (2012). MicroRNA-122 suppresses cell proliferation and induces cell apoptosis in hepatocellular carcinoma by directly targeting Wnt/ β -catenin pathway. *Liver Int*. 32:752-760.
- [108] Coulouarn C, Factor VM, Andersen JB, Durkin ME, Thorgeirsson SS. (2009). Loss of miR-122 expression in liver cancer correlates with suppression of the hepatic phenotype and gain of metastatic properties. *Oncogene*. 28:3526-3536.
- [109] Hsu SH, Wang B, Kota J, Yu J, Costinean S, Kutay H, Yu L, Bai S, La Perle K, Chivukula RR, Mao H, Wei M, Clark KR, Mendell JR, Caligiuri MA, Jacob ST, Mendell JT, Ghoshal K. (2012). Essential metabolic, anti-inflammatory, and anti-tumorigenic functions of miR-122 in liver. *J Clin Invest*. 122:2871-2883.
- [110] Nguyen DX, Bos PD, Massague J. (2009). Metastasis: From dissemination to organ-specific colonization. *Nat. Rev. Cancer*. 9:274–284.
- [111] Hanahan D, Weinberg RA. (2000). The hallmarks of cancer. *Cell*. 100:57–70.
- [112] Steeg PS. (2006). Tumor metastasis: Mechanistic insights and clinical challenges. *Nat. Med*. 12:895–904.
- [113] Ma L, Reinhardt F, Pan E, Soutschek J, Bhat B, Marcusson EG, Teruya-Feldstein J, Bell GW, Weinberg RA. (2010). Therapeutic silencing of miR-10b inhibits metastasis in a mouse mammary tumor model. *Nat. Biotechnol*. 28:341–347.
- [114] Tavazoie SF, Alarcon C, Oskarsson T, Padua D, Wang Q, Bos PD, Gerald WL, Massagué J. (2008). Endogenous human microRNAs that suppress breast cancer metastasis. *Nature*. 451:147–52.
- [115] Valastyan S, Reinhardt F, Benaich N, Calogrias D, Szász AM, Wang ZC, Brock JE, Richardson AL, Weinberg RA. (2009). A pleiotropically acting microRNA, miR-31, inhibits breast cancer metastasis. *Cell*. 137:1032–1046.
- [116] Ma L, Young J, Prabhala H, Pan E, Mestdagh P, Muth D, Teruya-Feldstein J, Reinhardt F, Onder TT, Valastyan S, Westermann F, Speleman F, Vandesompele J, Weinberg RA. (2010). MiR-9, a MYC/MYCN-activated microRNA, regulates E-cadherin and cancer metastasis. *Nat. Cell Biol*. 12: 247–256.
- [117] Gregory, P.A., et al., The miR-200 family and miR-205 regulate epithelial to mesenchymal transition by targeting ZEB1 and SIP1. *Nat Cell Biol*, 2008. 10(5): p. 593-601.
- [118] Gibbons, D.L., et al., Contextual extracellular cues promote tumor cell EMT and metastasis by regulating miR-200 family expression. *Genes Dev*, 2009. 23(18): p. 2140-51.
- [119] Korpel, M., et al., Direct targeting of Sec23a by miR-200s influences cancer cell secretome and promotes metastatic colonization. *Nat Med*, 2011. 17(9): p. 1101-8.

- [120] Kiss-Laszlo, Z., Henry, Y., Bachellerie, J., Caizergues- Ferrer, M. & Kiss, T. (1996). Site-specific ribose methylation of preribosomal RNA: a novel function for small nucleolar RNAs. *Cell*. 85:1077–1088.
- [121] Tollervey, D. & Kiss, T. (1997). Function and synthesis of small nucleolar RNAs. *Curr. Opin. Cell Biol.* 9:337–342.
- [122] Weinstein L. B., Steitz J. A. (1999). Guided tours: from precursor snoRNA to functional snoRNP. *Curr. Opin. Cell Biol.* 11: 378–384.
- [123] Williams GT, Hughes JP, Stoneman V, Anderson CL, McCarthy NJ, Mourtada-Maarabouni M, Pickard M, Hedge VL, Trayner I, Farzaneh F. (2006). Isolation of genes controlling apoptosis through their effects on cell survival. *Gene Ther. Mol. Biol.* 10B:255–261.
- [124] Mourtada-Maarabouni M, Hedge VL, Kirkham L, Farzaneh F, Williams GT. (2008). Growth arrest in human T-cells is controlled by the non-coding RNA growth-arrest-specific transcript 5 (GAS5). *J. Cell Sci.* 121:939–946.
- [125] Mourtada-Maarabouni M, Pickard MR, Hedge VL, Farzaneh F, Williams GT. (2009) GAS5, a non-protein-coding RNA, controls apoptosis and is downregulated in breast cancer. *Oncogene*. 28:195–208.
- [126] Dong XY, Guo P, Boyd J, Sun X, Li Q, Zhou W, Dong JT. (2009). Implication of snoRNA U50 in human breast cancer. *J. Genet. Genomics* 36:447–454.
- [127] Dong XY, Rodriguez C, Guo P, Sun X, Talbot JT, Zhou W, Petros J, Li Q, Vessella RL, Kibel AS, Stevens VL, Calle EE, Dong JT. (2008). SnoRNA U50 is a candidate tumor-suppressor gene at 6q14.3 with a mutation associated with clinically significant prostate cancer. *Hum. Mol. Genet.* 17:1031–1042.
- [128] Gee HE, Buffa FM, Camps C, Ramachandran A, Leek R, Taylor M, Patil M, Sheldon H, Betts G, Homer J, West C, Ragoussis J, Harris AL. (2011). The small-nucleolar RNAs commonly used for microRNA normalisation correlate with tumour pathology and prognosis. *Br. J. Cancer*. 104:1168–1177.
- [129] Liao J, Yu L, Mei Y, Guarnera M, Shen J, Li R, Liu Z, Jiang F. (2010). Small nucleolar RNA signatures as biomarkers for non-small-cell lung cancer. *Mol. Cancer*. 9:198.
- [130] Martens-Uzunova ES, Jalava SE, Dits NF, van Leenders GJ, Møller S, Trapman J, Bangma CH, Litman T, Visakorpi T, Jenster G. (2011). Diagnostic and prognostic signatures from the small non-coding RNA transcriptome in prostate cancer. *Oncogene*. 31:978–991.
- [131] Brown JW, Marshall DF, Echeverria M. (2008). Intronic noncoding RNAs and splicing. *Trends Plant Sci.* 13:335–342.
- [132] Chanfreau G, Legrain P, Jacquier A. (1998). Yeast RNase III as a key processing enzyme in small nucleolar RNAs metabolism. *J. Mol. Biol.* 284:975–988.
- [133] Petfalski E, Dandekar T, Henry Y, Tollervey D. (1998). Processing of the precursors to small nucleolar RNAs and rRNAs requires common components. *Mol. Cell. Biol.* 18:1181–1189.
- [134] Qu LH, Henras A, Lu YL, Zhou H, Zhou WX, Zhu YQ, Zhao J, Henry Y, Caizergues-Ferrer M, Bachellerie JP. (1999). Seven novel methylation guide small nucleolar RNAs are processed from a common polycistronic transcript by Rat1p and RNase III in yeast. *Mol. Cell. Biol.* 19:1144–1158.

- [135] Nicholls RD, Knepper JL. (2001). Genome organization, function, and imprinting in Prader–Willi and Angelman syndromes *Annu. Rev. Genomics Hum. Genet.* 2:153–175.
- [136] Skryabin BV, Gubar LV, Seeger B, Pfeiffer J, Handel S, Robeck T, Karpova E, Rozhdestvensky TS, Brosius J. (2007). Deletion of the MBII-85 snoRNA gene cluster in mice results in postnatal growth retardation. *PLoS Genet.* 3:e235.
- [137] Gurevich I, Englander MT, Adlersberg M, Siegal NB, Schmauss C. (2002). Modulation of serotonin 2C receptor editing by sustained changes in serotonergic neurotransmission *J. Neurosci.* 22:10529–1053.
- [138] Burns CM, Chu H, Rueter SM, Hutchinson LK, Canton H, Sanders-Bush E, Emeson RB. (1997). Regulation of serotonin-2C receptor G-protein coupling by RNA editing *Nature.* 387:303–308.
- [139] Tanaka R, Satoh H, Moriyama M, Satoh K, Morishita Y, Yoshida S, Watanabe T, Nakamura Y, Mori S. (2000). Intronic U50 small-nucleolar-RNA (snoRNA) host gene of no protein-coding potential is mapped at the chromosome breakpoint t(3;6) (q27;q15) of human B-cell lymphoma. *Genes Cells.* 5:277–287.
- [140] Dong XY, Rodriguez C, Guo P, Sun X, Talbot JT, Zhou W, Petros W, Li Q, Vessella RL, Kibel AS, Stevens VL, Calle EE, Dong DJ. (2008). SnoRNA U50 is a candidate tumor-suppressor gene at 6q14.3 with a mutation associated with clinically significant prostate cancer *Hum. Mol. Genet.* 17:1031–1042.
- [141] Mei YP, Liao JP, Shen JP, Yu L, Liu BL, Liu L, Li RY, Ji L, Dorsey SG, Jiang ZR, Katz RL, Wang JY, Jiang F. (2011). Small nucleolar RNA 42 acts as an oncogene in lung tumorigenesis. *Oncogene.* (doi:10.1038/onc.2011.449).
- [142] Chang LS, Lin S Y, Lieu AS, Wu TL. (2002). Differential expression of human 5S snoRNA genes. *Biochem. Biophys. Res. Commun.* 299:196–200.
- [143] Liao J, Yu L, Mei Y, Guarnera M, Shen J. (2010). Small nucleolar RNA signatures as biomarkers for non-small-cell lung cancer. *Mol. Cancer.* 9:198.
- [144] Li R, Wang H, Bekele BN, Yin Z, Caraway NP, Katz RL, Stass SA, Jiang F. (2006). Identification of putative oncogenes in lung adenocarcinoma by a comprehensive functional genomic approach. *Oncogene.* 18:2628–2635.
- [145] Schneider C, King RM, Philipson L. Genes specifically expressed at growth arrest of mammalian cells. *Cell.* 1988;54:787–793.
- [146] Amaldi F, Pierandrei-Amaldi P. (1997). TOP genes: a translationally controlled class of genes including those coding for ribosomal proteins. In: Jeanteur P, editor. *Progress in molecular and subcellular biology.* Vol. 18. Springer-Verlag; Berlin, Germany: pp. 1–17.
- [147] Mourtada-Maarabouni M, Pickard MR, Hedge VL, Farzaneh F, Williams GT. (2009). GAS5, a non-protein-coding RNA, controls apoptosis and is downregulated in breast cancer GAS5 regulates apoptosis. *Oncogene.* 28:195–208.
- [148] Gee HE, Buffa FM, Camps C, Ramachandran A, Leek R, Taylor M, Patil M, Sheldon H, Betts G, Homer J, West C, Ragoussis J, Harris AL. (2011). The small-nucleolar RNAs commonly used for microRNA normalisation correlate with tumour pathology and prognosis. *Br. J. Cancer.* 104, 1168–1177.
- [149] Nakamura Y, Takahashi N, Kakegawa E, Yoshida K, Ito Y, Kayano H, Niitsu N, Jinnai I, Bessho M. (2008). The GAS5 (growth arrest-specific transcript 5) gene fuses to BCL6 as

- a result of t(1;3)(q25;q27) in a patient with B-cell lymphoma. *Cancer Genetics and Cytogenetics*.182:144-149.
- [150] Askarian-Amiri ME, Crawford J, French JD, Smart CE, Smith MA, Clark MB, Ru K, Mercer TR, Thompson ER, Lakhani SR, Vargas AC, Campbell IG, Brown MA, Dinger ME, Mattick JS. (2011). SNORD-host RNA Zfas1 is a regulator of mammary development and a potential marker for breast cancer RNA. *17*:878-891.
- [151] Gupta V, Kumar A. (2010). Dyskeratosis congenita. *Adv Exp Med Biol*.685:215-219.
- [152] Ruggero D, Grisendi S, Piazza F, Rego E, Mari F, Rao PH, Cordon-Cardo C, Pandolfi PP. (2003). Dyskeratosis congenita and cancer in mice deficient in ribosomal RNA modification. *Science*. 299:259-62.
- [153] Ono M, Scott MS, Yamada K, Avolio F, Barton GJ, Lamond AI. (2011). Identification of human miRNA precursors that resemble box C/D snoRNAs. *Nucleic Acids Res*.39:3879-3891.
- [154] Xiao J, Lin H, Luo X, Luo X, Wang Z. (2011). miR-605 joins p53 network to form a p53:miR-605:Mdm2 positive feedback loop in response to stress. *EMBO J*.30:5021.
- [155] Siomi MC, Sato K, Pezic D, Aravin AA. (2011). PIWI-interacting small RNAs: the vanguard of genome defence. *Nat Rev Mol Cell Biol*.12:246-258.
- [156] Ishizu H, Nagao A, Siomi H. (2011). Gatekeepers for Piwi-piRNA complexes to enter the nucleus. *Curr Opin Genet Dev*. 21:484-490.
- [157] Kazazian HH Jr. (2004). Mobile elements: drivers of genome evolution. *Science* 303:1626–1632.
- [158] Saito K, Siomi MC. (2010). Small RNA-mediated quiescence of transposable elements in animals. *Dev. Cell*. 19:687–697.
- [159] Vagin VV, Klenov MS, Kalmykova AI, Stolyarenko AD, Kotelnikov RN, Gvozdev VA. (2004). The RNA interference proteins and vasa locus are involved in the silencing of retrotransposons in the female germline of *Drosophila melanogaster*. *RNA Biol*. 1:54–58.17.
- [160] Kalmykova AI, Klenov MS, Gvozdev VA. (2005). Argonaute protein PIWI controls mobilization of retrotransposons in the *Drosophila* male germline. *Nucleic Acids Res*. 33:2052–2059.
- [161] Savitsky M, Kwon D, Georgiev P, Kalmykova A, Gvozdev V. (2006). Telomere elongation is under the control of the RNAi-based mechanism in the *Drosophila* germline. *Genes Dev*. 20:345–354.
- [162] Li C, Vagin VV, Lee S, Xu J, Ma S, Xi H, Seitz H, Horwich MD, Syrzycka M, Honda BM, Kittler EL, Zapp ML, Klattenhoff C, Schulz N, Theurkauf WE, Weng Z, Zamore PD. (2009). Collapse of germline piRNAs in the absence of Argonaute3 reveals somatic piRNAs in flies. *Cell*. 137:509–521.
- [163] Aravin A, Gaidatzis D, Pfeffer S, Lagos-Quintana M, Landgraf P, Iovino N, Morris P, Brownstein MJ, Kuramochi-Miyagawa S, Nakano T, Chien M, Russo JJ, Ju J, Sheridan R, Sander C, Zavolan M, Tuschl T. (2006). A novel class of small RNAs bind to MILI protein in mouse testes. *Nature*. 442:203–207.
- [164] Girard A, Sachidanandam R, Hannon GJ, Carmell MA. (2006) A germline-specific class of small RNAs binds mammalian Piwi proteins. *Nature*. 442:199–202.

- [165] Brennecke J, Aravin AA, Stark A, Dus M, Kellis M, Sachidanandam R, Hannon GJ. (2007). Discrete small RNA-generating loci as master regulators of transposon activity in *Drosophila*. *Cell*. 128:1089–1103.
- [166] Lau NC, Seto AG, Kim J, Kuramochi-Miyagawa S, Nakano T, Bartel DP, Kingston RE. (2006). Characterization of the piRNA complex from rat testes. *Science* 313, 363–367
- [167] Houwing S, Berezikov E, Ketting RF. (2008). Zili is required for germ cell differentiation and meiosis in zebrafish. *EMBO J*. 27:2702–2711.
- [168] Robine N, Lau NC, Balla S, Jin Z, Okamura K, Kuramochi-Miyagawa S, Blower MD, Lai EC. (2009) A broadly conserved pathway generates 3'UTR-directed primary piRNAs. *Curr. Biol*. 19:2066–2076.
- [169] Aravin AA, Sachidanandam R, Bourc'his D, Schaefer C, Pezic D, Toth KF, Bestor T, Hannon GJ. (2008). A piRNA pathway primed by individual transposons is linked to de novo DNA methylation in mice. *Mol. Cell*. 31:785–799.
- [170] Qiao D, Zeeman AM, Deng W, Looijenga LH, Lin H. (2002). Molecular characterization of hiwi, a human member of the piwi gene family whose overexpression is correlated to seminomas. *Oncogene*.21:3988-3999.
- [171] Taubert H, Würfl P, Greither T, Kappler M, Bache M, Bartel F, Kehlen A, Lautenschläger C, Harris LC, Kaushal D, Füssel S, Meye A, Böhnke A, Schmidt H, Holzhausen HJ, Hauptmann S. (2007). Stem cell-associated genes are extremely poor prognostic factors for soft-tissue sarcoma patients. *Oncogene*. 26:7170-7174.
- [172] Taubert H, Greither T, Kaushal D, Würfl P, Bache M, Bartel F, Kehlen A, Lautenschläger C, Harris L, Kraemer K, Meye A, Kappler M, Schmidt H, Holzhausen HJ, Hauptmann S. (2007) Expression of the stem cell self-renewal gene Hiwi and risk of tumour-related death in patients with soft-tissue sarcoma. *Oncogene*.26:1098-1100.
- [173] Wang QE, Han C, Milum K, Wani AA. (2011). Stem cell protein Piwil2 modulates chromatin modifications upon cisplatin treatment. *Mutat Res*.708:59-68.
- [174] Lu Y, Li C, Zhang K, Sun H, Tao D, Liu Y, Zhang S, Ma Y. (2010). Identification of piRNAs in Hela cells by massive parallel sequencing. *BMB Rep*.43:635-41.
- [175] Cui L, Lou Y, Zhang X, Zhou H, Deng H, Song H, Yu X, Xiao B, Wang W, Guo J. (2011). Detection of circulating tumor cells in peripheral blood from patients with gastric cancer using piRNAs as markers. *Clin Biochem*. 44:1050-1057.
- [176] Cheng J, Guo JM, Xiao BX, Miao Y, Jiang Z, Zhou H, Li QN. (2011). piRNA, the new non-coding RNA, is aberrantly expressed in human cancer cells. *Clin Chim Acta*. 412:1621-1625.
- [177] Cheng J, Deng H, Xiao B, Zhou H, Zhou F, Shen Z, Guo J. (2012). piR-823, a novel non-coding small RNA, demonstrates in vitro and in vivo tumor suppressive activity in human gastric cancer cells. *Cancer Lett*. 315:12-17
- [178] Reuter M, Chuma S, Tanaka T, Franz T, Stark A, Pillai RS. (2009). Loss of the Mili-interacting Tudor domain-containing protein-1 activates transposons and alters the Mili-associated small RNA profile. *Nature Struct. Mol. Biol*. 16:639–646.
- [179] Shoji M, Tanaka T, Hosokawa M, Reuter M, Stark A, Kato Y, Kondoh G, Okawa K, Chujo T, Suzuki T, Hata K, Martin SL, Noce T, Kuramochi-Miyagawa S, Nakano T, Sasaki H, Pillai RS, Nakatsuji N, Chuma S. (2009) The TDRD9-MIWI2 complex is

- essential for piRNA-mediated retrotransposon silencing in the mouse male germline. *Dev. Cell* 17:775–787.
- [180] Struhl, K. (2007). Transcriptional noise and the fidelity of initiation by RNA polymerase II. *Nature Struct. Mol. Biol.* 14:103–105.
- [181] Amaral PP, Mattick JS. (2008). Noncoding RNA in development. *Mamm. Genome.* 19:454–492.
- [182] Dinger ME, Amaral PP, Mercer TR, Pang KC, Bruce SJ, Gardiner BB, Askarian-Amiri ME, Ru K, Soldà G, Simons C, Sunkin SM, Crowe ML, Grimmond SM, Perkins AC, Mattick JS. (2008). Long noncoding RNAs in mouse embryonic stem cell pluripotency and differentiation. *Genome Res.* 18:1433–1445.
- [183] Mercer TR, Dinger ME, Sunkin SM, Mehler MF, Mattick JS. (2008). Specific expression of long noncoding RNAs in the adult mouse brain. *Proc. Natl Acad. Sci. USA.* 105:716–721.
- [184] Cawley S, Bekiranov S, Ng HH, Kapranov P, Sekinger EA, Kampa D, Piccolboni A, Sementchenko V, Cheng J, Williams AJ, Wheeler R, Wong B, Drenkow J, Yamanaka M, Patel S, Brubaker S, Tammanna H, Helt G, Struhl K, Gingeras TR. (2004). Unbiased mapping of transcription factor binding sites along human chromosomes 21 and 22 points to widespread regulation of noncoding RNAs. *Cell.* 116:499–509.
- [185] Ponjavic J, Ponting CP, Lunter G. (2007). Functionality or transcriptional noise? Evidence for selection within long noncoding RNAs. *Genome Res.* 17:556–565.
- [186] Faulkner GJ, Kimura Y, Daub CO, Wani S, Plessy C, Irvine KM, Schroder K, Cloonan N, Steptoe AL, Lassmann T, Waki K, Hornig N, Arakawa T, Takahashi H, Kawai J, Forrest AR, Suzuki H, Hayashizaki Y, Hume DA, Orlando V, Grimmond SM, Carninci P. (2009). The regulated retrotransposon transcriptome of mammalian cells. *Nat Genet.* 41:563–71.
- [187] Guttman M, Amit I, Garber M, French C, Lin MF, Feldser D, Huarte M, Zuk O, Carey BW, Cassady JP, Cabili MN, Jaenisch R, Mikkelsen TS, Jacks T, Hacohen N, Bernstein BE, Kellis M, Regev A, Rinn JL, Lander ES. (2009). Chromatin signature reveals over a thousand highly conserved large non-coding RNAs in mammals. *Nature.* 458:223–227.
- [188] Pang KC, Frith MC, Mattick JS. (2006). Rapid evolution of noncoding RNAs: lack of conservation does not mean lack of function. *Trends Genet.* 22:1–5.
- [189] Cheng LL, Carmichael GG. (2010). Decoding the function of nuclear long non-coding RNAs. *Curr Opinion in Cell Biology.* 22:357–364.
- [190] Kugel JF, Goodrich JA. (2012). Non-coding RNAs: key regulators of mammalian transcription. *Trends in Biochemical Sciences.* 37:144–151.
- [191] Khalil AM, Guttman M, Huarte M, Garber M, Ray A, Rivea Morales D, Thomas K, Presser A. (2009). Many human large intergenic noncoding RNAs associated with chromatin-modifying complexes and affect gene expression. *Proc Natl Acad Sci USA.* 106:11667–11672.
- [192] Zhao J, Sun BK, Erwin JA, Song JJ, Lee JT. (2008). Polycomb proteins targeted by a short repeat RNA to the mouse X chromosome. *Science* 322:750–756.
- [193] Rinn JL, Kertesz M, Wang JK, Squazzo SL, Xu X, Brugmann SA, Goodnough LH, Helms JA, Farnham PJ, Segal E, Chang HY. (2007) Functional demarcation of active and silent chromatin domains in human HOX loci by noncoding RNAs. *Cell* 129, 1311–1323.

- [194] Tsai MC, Manor O, Wan Y, Mosammaparast N, Wang JK, Lan F, Shi Y, Segal E, Chang HY. (2010). Long noncoding RNA as modular scaffold of histone modification complexes. *Sciences*. 329:689-693.
- [195] Morris KV, Santoso S, Turner AM, Pastori C, Hawkins PG. (2008). Bidirectional transcription directs both transcriptional gene activation and suppression in human cells. *PLoS Genet*. 4:e1000258.
- [196] Nagano T, Mitchell JA, Sanz LA, Pauler FM, Ferguson-Smith AC, Feil R, Fraser P. (2008). The Air noncoding RNA epigenetically silences transcription by targeting G9a to chromatin. *Science*. 322:1717-1720.
- [197] Gupta RA, Shah N, Wang KC, Kim J, Horlings HM, Wong DJ, Tsai MC, Hung T, Argani P, Rinn JL, Wang Y, Brzoska P, Kong B, Li R, West RB, van de Vijver MJ, Sukumar S, Chang HY. (2010). Long non-coding RNA HOTAIR reprograms chromatin state to promote cancer metastasis. *Nature*. 464:1071-1076.
- [198] Kogo R, Shimamura T, Mimori K, Kawahara K, Imoto S, Sudo T, Tanaka F, Shibata K, Suzuki A, Komune S, Miyano S, Mori M. (2011). Long non-coding RNA HOTAIR regulates Polycomb-dependent chromatin modification and is associated with poor prognosis in colorectal cancers. *Cancer Res*. 71: 6320-6326.
- [199] Yap KL, Li S, Muñoz-Cabello AM, Raguz S, Zeng L, Mujtaba S, Gil J, Walsh MJ, Zhou MM. (2010). Molecular interplay of the non coding RNA ANRIL and methylated histone H3 lysine 27 by polycomb CBX7 in transcriptional silencing of INK4a. *Mol Cell*. 38:662-674.
- [200] Beltran M, Puig I, Peña C, García JM, Alvarez AB, Peña R, Bonilla F, de Herreros AG. (2008). A natural antisense transcript regulates Zeb2/Sip1 gene expression during Snail1-induced epithelial-mesenchymal transition. *Genes Dev*. 22:756-769.
- [201] Wang X, Arai S, Song X, Reichart D, Du K, Pascual G, Tempst P, Rosenfeld MG, Glass CK, Kurokawa R. (2008). Induced ncRNAs allosterically modify RNA-binding proteins in cis to inhibit transcription. *Nature*. 454:126-130.
- [202] Feng J, Bi C, Clark BS, Mady R, Shah P, Kohtz JD. (2006). The Evf-2 noncoding RNA is transcribed from the Dlx-5/6 ultraconserved region and functions as a Dlx-2 transcriptional coactivator. *Genes Dev*. 20:1470-1484.
- [203] Huarte M, Guttman M, Feldser D, Garber M, Koziol MJ, Kenzelmann-Broz D, Khalil AM, Zuk O, Amit I, Rabani M, Attardi LD, Regev A, Lander ES, Jacks T, Rinn JL. (2010). A large intergenic noncoding RNA induced by p53 mediates global gene repression in the p53 response. *Cell*. 142:409-419.
- [204] Bond CS, Fox AH. (2009). Paraspeckles: nuclear bodies built on long noncoding RNA. *J Cell Biol*. 186:637-644.