We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Endocannabinoids and Kisspeptins: Two Modulators in Fight for the Regulation of GnRH Activity

Rosaria Meccariello, Rosanna Chianese, Silvia Fasano and Riccardo Pierantoni

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/48443>

1. Introduction

The master system in the control of reproductive functions is the communication into the hypothalamus-pituitary-gonadal axis (HPG), whose main actor is the hypothalamic gonadotropin releasing hormone (GnRH). Such a decapeptide triggers the release of pituitary gonadotropins [Follicular Stimulating Hormone (FSH) and Luteinizing Hormone (LH)] which in turn reach the gonads, induce the biosynthesis of steroids (mainly testosterone in males and estradiol/progesterone in females) and of other non steroidal substances (i.e. activin, follistatin, inhibin) modulating the gametogenesis in both sexes. In the last decades, a significant upsurge of studies aimed to define several actors and mechanisms supporting reproductive activity. Ultra short, short and long feedback in HPG communication finely modulate reproduction. Nevertheless, this picture is still puzzling and the complete knowledge of the full process has to be unravelled.

“Only on the basis of an extensive comparative biology can authentic general biology emerge” (Bern 1967). Besides the importance of evolutionary track in the research of adaptive phenomena, comparative approaches provide a deep insight into the physiological mechanism in building general models. At present, 25 GnRH molecular forms have been detected in metazoan, also in species lacking pituitary gland. Up to 15 molecular forms have been detected in vertebrates; fish, amphibians, reptiles, birds and also humans possess two GnRH molecular forms (GnRH1 and GnRH2, formerly known as mammalian GnRH and chicken 2 GnRH, respectively) as well as one GnRH receptor (GnRHR), at least. A third GnRH molecular form, GnRH3, is often detected in fish telencephalon and peripheral tissues (Pierantoni et al., 2002). In this respect, current hypothesis postulates that GnRH action progressively evolved from the control of simple basic functions in early metazoan to an indirect way to check gonadal activity in

vertebrates, through a sophisticated network of finely tuned neurons (Chianese et al., 2011a; Kah et al., 2007; Kavanaugh et al., 2008; Pierantoni et al., 2002; White et al., 1998). Despite both GnRH1 and GnRH2 share the ability to trigger gonadotropin discharge (Pierantoni et al., 2002), the coexistence of multiple forms of GnRHs in the brain let to hypothesize the division of functional roles. In this respect, GnRH2 activity is also involved in processes other than gonadotropin discharge such as the control of sexual behaviour, or local action at gonadal level. Food intake and energy balance, stress and many other environmental cues deeply affect reproductive success *via* GnRH2. Hence, in such a complex scenario emerged: 1) the need to integrate and convey all information to GnRH neurons, the major hierarchical elements of the HPG and 2) the need to discover possible intermediary neuronal populations in this chain of events (Fernandez-Fernandez et al., 2006; Herbison & Pape, 2001).

Therefore, in this review, we focus on endocannabinoid (ECB) and kisspeptin systems, two modulators of GnRH activity. ECBs are lipidic mediators capable to inhibit the release of hypothalamic GnRH (Scorticati et al., 2004), affecting, as a consequence, both steroidogenesis and gonadal functions (Wang et al., 2004). While ECBs exert such a negative effect upon GnRH release, kisspeptins, the product of *kiss* gene, positively affect GnRH release. Impairment of kisspeptin system causes idiopathic hypogonadotropic hypogonadism and affects puberty onset (de Roux et al., 2003; Seminara et al., 2003). Thus, at hypothalamic level ECBs and kisspeptin modulate GnRH circuitry in opposite manner. Similarly to GnRH, ECBs and kisspeptin exert a direct effect upon gonadal activity, affecting steroidogenesis, spermatogenesis, spermatozoa functions, follicular development and oocyte maturation, suggesting the existence of a possible local crosstalk among these systems. Thus, in the next paragraphs the activity of ECBs and kisspeptin along the HPG will be properly discussed.

2. New modulators of GnRH/gonadotropin activity at central level

2.1. Endocannabinoid system

The endocannabinoid system (ECS) is an ancient, evolutionarily conserved system, well-characterized in mammalian and non-mammalian vertebrates (Buznikov et al., 2010; Fasano et al., 2009; McPartland et al., 2006). Such a system comprises ECBs, several ECB receptors (CBs), many enzymatic machineries responsible for ECB degradation and biosynthesis and ECB transporters (EMT) (Pierantoni et al., 2009). A schematic representation of ECS components is depicted in Figure 1.

In general, ECBs are amides, esters and ethers of long-chain polyunsaturated fatty acid, isolated from brain, peripheral tissues and reproductive fluids (Devane et al., 1992; Sugiura et al., 1995; Schuel et al., 2002); they mimic the effects of the phytocannabinoid Δ^9 -tetrahydrocannabinol (Δ^9 -THC), the psychoactive constituent of marijuana plant, *Cannabis sativa*. The main ECBs are the N-arachidonoyl-ethanolamine (AEA, anandamide), the first ECB discovered in porcine brain (Devane et al., 1992), and 2-arachidonoylglycerol (2-AG) (Sugiura et al., 1995). ECBs have the ability to activate a wide range of CBs: the most studied are CB1 and CB2, classical seven transmembrane spanning G coupled receptors (Matsuda et

al., 1990; Munro et al., 1993) widely distributed in both brain and peripheral tissues, gonads included (Galiegue et al., 1995, Shire et al., 1995, Brown et al., 2002). The orphan G coupled receptor GPR55 is currently accounted as the third CB (Lauckner et al., 2008). AEA, but not 2-AG, selectively acts as an intracellular ligand of the transient potential type1 vanilloid receptor (TRPV1) channel, a six transmembrane spanning receptor whose structure forms a ligand gated non selective cationic channel activated by capsaicin, one of red chilli pepper component (van der Stelt & Di Marzo, 2004, 2005). Lastly, direct nuclear action of ECBs has been postulated since many ECBs [for instance AEA, 2-AG, N-oleoyl-ethanolamine (OEA), N-palmitoyl-ethanolamine (PEA), noladin ether and virodhamine], the phytocannabinoid Δ^9 -THC, CB agonists (HU210, WIN55121-2) as well as cannabinoid metabolites have the ability to activate also PPAR (peroxisome-proliferator-activated receptor) family of nuclear receptors (O'Sullivan, 2007; Sun & Bennett, 2007). In order to activate PPAR receptors, cytoplasmic-nuclear translocation of ECBs requires the fatty acid binding proteins (FABPs) as intracellular carriers (Kaczocha et al., 2012).

ECS comprises ECBs, CBs, ECB biosynthetic and hydrolyzing enzymes, as well as membrane and intracellular carriers (See text for details). Figure modified from: Pierantoni et al., 2009.

Figure 1. Schematic representation of ECS components.

ECBs activity strongly depends on the balance between their biosynthetic and hydrolyzing pathways. AEA and 2-AG are usually released from membrane phospholipid precursors through the activation of N-acyl phosphatidylethanolamine phospholipase D (NAPE-PLD) and diacylglycerol lipase (DAGL), respectively (Bisogno et al., 2003; Okamoto et al., 2004).

Two fatty acid amide hydrolases (FAAH and FAAH-2) (Cravatt et al., 1996; Wei et al., 2006) as well as *N*-acylethanolamine-hydrolyzing acid amidase (NAAA) (Tsuboi et al., 2005; Ueda et al., 2010) release arachidonic acid and ethanolamine from AEA. Besides FAAH, 2-AG is cleaved into arachidonic acid and glycerol by a specific monoacylglycerol lipase (MAGL) (Dinh et al., 2002; Ho et al., 2002). Despite of their lipidic nature, data concerning the existence of a membrane carrier able to mediate ECBs transport is discussed. Recently, in neuronal cells a FAAH-like AEA transporter (FLAT) has been identified. Such a molecule is encoded by a splicing variant of *FAAH-1*, lacks the catalytic activity of FAAH but has the ability to bind AEA (Fu et al., 2011).

Nowadays, ECS elements have been identified in the central and peripheral nervous system as well as in gonads and gametes, demonstrating a deep involvement of the system in the control of reproductive functions, both at central and local level (Battista et al., 2012).

In marijuana smokers as well as in animal models, cannabinoids and ECBs interfere in the neuroendocrine control of reproductive function impairing GnRH and LH production, gonadic steroid production, spermatogenesis, ovulation, embryo development and implantation, as well as sexual behaviour (Murphy et al., 1998; Pagotto et al., 2006; Wang et al., 2006). In the brain, Δ^9 -THC and ECBs are well known retrograde signals that act at presynaptic level in order to inhibit the release of specific neurotransmitters [i.e. γ -aminobutyric acid (GABA)]. Current opinion postulates that ECB mediated LH, but not FSH, inhibition is the result of hypothalamic ECB activity. In fact, ECBs are well known inhibitors of GnRH release (Scorticati et al., 2004) and GnRH transcription (Chianese et al., 2011b; Meccariello et al., 2008). By contrast, direct or indirect action of ECBs upon GnRH secreting neurons is still controversial and under investigation. ECBs inhibit several neuronal systems, positively involved in GnRH circuitry (i.e. norepinephrine and glutamate); by contrast, they activate well known inhibitors of GnRH activity [i.e. dopamine, endogenous opioid peptides and corticotrophin-releasing hormone (Murphy et al., 1998)]. Current hypothesis postulates that ECBs interfere in the well known regulation of GnRH neurons by long loop gonadal steroid feedback through steroids receptor expressing afferents such as GABAergic neurons. For instance, there is a growing consensus that GABA can act through the GABAA receptor to exert both depolarizing and hyperpolarizing effects on GnRH neurons (Herbison & Moenter, 2011). In male mice, GnRH-secreting neurons tonically release 2-AG in presynaptic fissure, which in turn activates CB1 located on GABAergic afferents, in tight relationship with GnRH neurons. The activation of CB1 inhibits the spontaneous release of GABA (Figure 2). As a consequence, postsynaptic GABA receptors (GABAA and GABAB), located on GnRH-secreting neurons, are not activated and GnRH is not released (Farkas et al., 2010). ECB biosynthesis in GnRH secreting neurons might be induced by the activation of metabotropic glutamate receptor (mGluR) located on astrocytes. In fact, in female mice, a complementary hypothesis suggests that local GnRH-GABA circuits uses just the glia derived prostaglandins and/or ECBs in a steroid dependent fashion (Glanowska & Moenter, 2011). GnRH neurons interact with their afferent neurons using several mechanisms and these local circuits can be modified by both sex and steroid feedback. ECBs tone is certainly a key factor in ECBs activity. The inhibitory effect of AEA on GnRH-secreting neurons is reversed by estrogens (Scorticati et al., 2004), through the

inhibition of astrocyte mGluR. Such a process inhibits prostaglandin mediated release of ECBs from GnRH secreting neurons (Glanowska & Moenter, 2011). Alternatively, estradiol might directly prevent the ECB mediated inhibition of GABA neurons (Glanowska & Moenter, 2011). In the brain, functional relationships between CB1 and FAAH emerged, since they have a quite overlapping localization (Egertová et al., 1998). Since estradiol modulates the transcription of FAAH hydrolase, whose promoter contains an ERE element (Waleh et al., 2002), it is not excluded that estradiol might reverse the adverse activity of AEA on GnRH neurons, by means of FAAH upregulation and AEA degradation.

Conversely, it is not excluded that neuronal systems other than GABAergic, as for example kisspeptin neurons described in the next paragraphs, might modulate GnRH-secreting neurons activity *via* ECBs in an estradiol dependent fashion. A model for proposed circuits and possible mechanisms of GnRH neurons activity in males and females are, respectively, reported in Figures 2 and 3. The use of CB1 knockout mice (*CB1*^{-/-}) also contributed to elucidate the mechanism of ECB mediated LH inhibition. AEA decreases both LH and prolactin (PRL) in *CB1*^{-/-} mice whereas 2-AG is able to suppress LH in wild-type, but not in *CB1*^{-/-} mice (Oláh et al., 2008). Thus, receptors other than CB1 might be involved in such a signalling. In such a context, the main candidate is TRPV1 (Oláh et al., 2008), whose expression has been reported in the hypothalamus but not in pituitary gland.

The basal crosstalk between ECS and GnRH is evolutionarily conserved, since it has been described also in lower vertebrates (Chianese et al., 2008, 2011b; Cottone et al., 2008, Meccariello et al., 2008). In both amphibians and teleost fish, CB1 was detected in the forebrain, the encephalic macro-area containing the anterior preoptic area, the encephalic region mainly involved in GnRH activity and in the control of gonadotropin discharge (Cottone et al., 2003, 2005; Lam et al., 2006; Migliarini et al., 2006; Meccariello et al., 2008; Valenti et al., 2005). Besides an involvement in food intake, also in lower vertebrates ECS negatively modulates neuroendocrine machinery and reproduction. In fish forebrain, CB1 colocalizes with GnRH3, the GnRH molecular form mainly detected in the telencephalon of fish (Cottone et al., 2008). A CB1 mediated self modulation of GnRH secreting neurons emerged in the anuran amphibian *Rana esculenta* (Meccariello et al., 2008). In male frogs, GnRH1 and CB1 share the localization inside the basal telencephalon and septum (Cottone et al., 2008; Meccariello et al., 2008); most GnRH1 secreting neurons are close to CB1 expressing neurons whereas a subpopulation of GnRH1 secreting neurons - in the approximate order of 20% - coexpresses CB1 (Meccariello et al., 2008). Such a neuroanatomical observation finds a possible functional explanation at molecular level. In fact, mRNA and protein profiles of CB1 and GnRH1 are opposite in frog diencephalon during the annual sexual cycle (Chianese et al., 2008; Meccariello et al., 2008). Treatments of male diencephalons with buserelin, a long acting GnRH analogue, inhibit *GnRH1* transcription and upregulate *CB1* transcription; conversely, AEA treatments downregulate *GnRH1* expression. In this respect, GnRH secreting neurons might produce ECBs in order to properly suppress GnRH secreting activity (ultrashort feedback). This saga is becoming more intricate since in this amphibian a second GnRH molecular form - GnRH2, with a suggested hypophysiotropic role (Pierantoni et al., 2002) - and three GnRHRs (GnRHR1,

GnRHR2, GnRHR3) have been cloned (Chianese et al., 2011b). In different periods of the annual reproductive cycle, AEA also inhibits *GnRH2* expression and upregulates the expression of *GnRHR1*, *GnRHR2*, but not of *GnRHR3* (Chianese et al., 2011b). Also immortalized neuronal cell lines (GT1) are both target and source of ECBs; *in vitro* they have the ability to produce and secrete ECBs (2-AG and AEA), to uptake and degrade ECBs, and possess CBs (both CB1 and CB2); the activation of CBs inhibits the pulsatile release of GnRH (Gammon et al., 2005). Nevertheless, such observations did not find any confirmation *in vivo*, although GnRH secreting neurons are close to cannabinergic fibres and scantily express CB1 (Gammon et al., 2005). By contrast, micro-array analysis revealed CB2 expression in a subpopulation of GnRH secreting neurons (Todman et al., 2005).

ECS interferes in GnRH circuitry also modulating the activity of neuronal populations that usually converge environmental, stressors, metabolic and photoperiodic cues at different levels of HPG. Stress and food intake, well-known processes under ECBs control (Pagotto et al., 2006), interfere in GnRH secretion. It is interesting to include in this scenario the gonadotropin-inhibitory hormone (GnIH). GnIH belongs to the super-family of RFamide neuropeptides, but its role in the control of gonadotropin secretion is negative, thus proposing the existence of a balance between stimulatory and inhibitory systems in the control of reproduction. Interestingly, this concept does not represent a rule; in fact, in male Syrian hamster, the mammalian ortholog of avian GnIH, the RFRP-3, works on the gonadotropic axis as a stimulator, inducing LH, FSH and testosterone secretion, *via* GnRH neurons activation. Furthermore, this effect might not only vary across species, but also include sex-specific differences in the same species, due to the loss of RFRP-3 mediated stimulation in females (Ancel et al., 2012).

Besides the effect in the hypothalamus, still controversial is the direct activity of ECBs on the pituitary. ECB binding sites, as well as the expression of CBs, biosynthetic and hydrolyzing enzymes have been reported in pituitary *pars distalis* and in pituitary cell cultures (Gonzales et al., 1999, 2000; Lynn & Herkenham, 1994; Murphy et al., 1998; Wenger et al., 1999).

Nevertheless, the localization of CBs in the gonadotropes is confirmed in amphibians and mammals, but not in humans (Cesa et al., 2002; Wenger et al., 1999; Yasuo et al., 2010a). For instance, in rats, AEA exhibits differential effects on the *in vitro* secretion of LH, and pituitary hormones other than gonadotropins [i.e. PRL, corticotrophin (ACTH) and growth hormone (GH)] (Wenger et al., 2000). In a sex steroid dependent fashion, rats express CB1 in pituitary gland. In females, CB1 expression and AEA content depends on the phase of the ovarian cycle and, in general, pituitary AEA content is opposite to that observed in the hypothalamus (Gonzales et al., 2000). Recently, ECBs have been included among tuberulins, the messengers supposed to be secreted from the *pars tuberalis* - a brain area located between the median eminence, the pituitary portal vessels and the pituitary *pars distalis* - to target the pituitary *pars distalis* (Yasuo et al., 2010b). In both hamsters and humans, *pars tuberalis* produces high levels of 2-AG and low levels of other ECBs (AEA, PEA and OEA), while the pituitary *pars distalis* possesses CB1 (Yasuo et al., 2010a, 2010b). Such a crosstalk might be involved in gonadal response to photoperiodic changes.

GnRH neuron, through GPR54 receptor, directly receives ARC kisspeptin input and conveys it to pituitary and testis. Gonadal steroids inhibit GnRH secretion through ARC kisspeptin neurons. 2-AG synthesized from GnRH neuron - by means of the CB1 activation - blocks GABA release toward GnRH neuron. ECBs directly act upon GnRH neuron since it expresses CB1; possibly, ECBs may have a further action on kisspeptin neuron. Anyway, an autocrine regulation of GnRH neuron - through ECBs - may exist.

Figure 2. Model for possible regulatory network along the male HPG.

Gonadal steroids exert both inhibitory and stimulatory effects on GnRH neurons, through ARC and AVPV kisspeptin neurons, respectively. While CB1 expression and so ECBs action on GnRH neuron are sure, it remains doubtful the mediation of kisspeptin neurons in the inhibitory regulation of GnRH neuron from ECBs. GnRH neuron is able to produce ECBs; these, in turn, bind CB1 on the GABA-ergic afferent, thus to inhibit GABA release. A possible alternative mechanism for this regulation supposes possible glutamate release from GnRH neuron. Glutamate - through metabotropic glutamate receptor (mGluR) activation - stimulates astrocytes to produce prostaglandins (PGs). These in turn can stimulate GnRH neuron to produce ECBs or regulate CB1 trafficking on GABA-ergic axon. Estradiol blocks this circuit inhibiting mGluR or CB1 activity.

Figure 3. Model for possible regulatory network along the female HPG.

2.2. Kisspeptin system

Kisspeptins belong to the family of RFamide peptides, encoded by the *kiss1* gene, originally detected as a metastasis-suppressor gene in several malignancies (Lee et al., 1996). Kiss name just derives from its role as a suppressor sequence (ss); the letters “Ki” were added after in homage to the location of its discovery, Hershey, Pennsylvania, home of the famous “Hershey Chocolate Kiss”. The major kisspeptin product - known as kisspeptin-54 - is a 54 amino acid peptide (Ohtaki et al., 2001); for its instability, it is proteolitically cleaved into shorter peptides (kisspeptin-10, -13 and -14) (Kotani et al., 2001). In 2001, four independent groups showed that all kisspeptin forms bind and activate with similar affinity the orphan G protein-coupled membrane receptor, GPR54 (Clements et al., 2001; Kotani et al., 2001; Muir et al., 2001; Ohtaki et al., 2001).

Hypogonadotropic hypogonadism has been described in mice lacking a functional *kiss* gene or in human and mice with mutations/targeted deletions of *GPR54* genes (Oakley et al., 2009).

Multiple and intricate are the molecular pathways activated by the kisspeptin/GPR54 system to exert its functions in a cell specific way. Starting from G-protein $G_{q/11}$ stimulation, kisspeptins induce phospholipase C (PLC) activation and intracellular calcium mobilization; kisspeptins are also able to induce a strong, sustained stimulation of phosphorylation of the MAP kinases extracellular signal regulated kinases ERK1 and ERK2 and a weak stimulation of p38 MAPK phosphorylation, whereas no activation was observed for stress-activated protein kinase/c-Jun NH2-terminal kinase (SAPK/JNK) (Castaño et al., 2009 and references therein).

The “reproductive” facet of kisspeptins got its disclosure in 2003 when mutations in *kiss1* or *GPR54* genes were associated with idiopathic hypogonadotropic hypogonadism and impaired pubertal maturation (de Roux et al., 2003; Seminara et al., 2003).

Since, the demonstration that kisspeptins are able to stimulate LH and, to a lesser extent, FSH secretion in several species (Roa et al., 2009 and references therein). However, a more consistent hypogonadotropic phenotype characterizes *GPR54* knock-out (KO) mice, compared to *kiss*-KO, suggesting the existence of other possible endogenous ligands for the receptor.

By contrast, GPR54 seems to be the only receptor responsible for the effects of kisspeptins (Lapatto et al., 2007). The main experimental evidences supporting the involvement of kisspeptin in the control of GnRH secreting neuron activity and gonadotropin discharge are summarized in Table 1. The activation of GnRH neurons appears under the control of a kisspeptin tone, but also involves an increase of the kisspeptin neuron projections to GnRH neurons (Roa et al., 2008). Accordingly, in rodents, the detection of kisspeptin-immunoreactive fibres around GnRH neuron cell bodies only starts from postnatal day 25 onwards, and the number of kisspeptin fibres increases at puberty (Clarkson & Herbison, 2006), suggesting that GnRH neurons become more sensitive to kisspeptin just before the onset of puberty. Surprisingly, at least in rodents, electrophysiological studies have demonstrated that a subset of GnRH neurons - responsive to group I metabotropic glutamate receptor agonists - is insensitive to kisspeptins (Dumalska et al., 2008).

In rodents, two major kisspeptin neuronal populations are located at the arcuate nucleus (ARC) and the preoptic area (POA) or the anteroventral periventricular nucleus (AVPV) of the hypothalamus (Mikkelsen et al., 2009). Although in these encephalic districts the kisspeptin-immunoreactive neuron distribution is highly similar in both male and female, the number of cell bodies detected in the AVPV shows a substantial sexual dimorphism being higher in female than in male, at least in rodents (Clarkson & Herbison, 2006). In contrast, in sheep and primates, hypothalamic kisspeptin neurons are especially located in the ARC/infundibular region (Estrada et al., 2006). In this neuro-anatomical picture, although GPR54 and kisspeptin immunoreactivities are rather overlapping, there are some examples of receptor-ligand mismatch. In particular, the ARC is rich in kisspeptin fibres and devoid of GPR54 (Herbison et al., 2010). Over again, the existence of unidentified receptors or ligands appears an intriguing interpretation of this phenomenon.

Key aspect of kisspeptin populations is the high degree of anatomical heterogeneity that may underlie a functional discernable network in the control of GnRH secretion. In this view, in the ARC - but not in the POA/AVPV - kisspeptin neurons co-localize two other neuropeptides, neurokinin B (NKB) and dynorphin (DYN), in several species analyzed to date. For convenience, these cells have been termed KNDy (Kisspeptin, Neurokinin B, Dynorphin) (Cheng et al., 2010). They constitute a central node in the control of GnRH secretion, given also the recent observation that genetic inactivation of *TAC3* or *TACR3*, which encode NKB peptide and its receptor, respectively, causes hypogonadotropic hypogonadism (Topaloglu et al., 2009). KNDy cells establish an interconnected network, through extensive reciprocal connections, which might serve to promote auto-regulatory mechanisms; in addition, they have a suggested dual peptidergic as well as glutamatergic phenotype (Lehman et al., 2010a). Conversely, in the AVPV kisspeptin neurons co-localize galanin, a neuropeptide involved in female reproductive functions (Vida et al., 2009), and tyrosine hydroxylase, a marker for dopaminergic neurons (Kauffman et al., 2007). Starting from ARC and AVPV nuclei, both trans-synaptic (indirect) and direct kisspeptin appositions respectively reach GnRH neurons (Clarkson & Herbison, 2006). Evidence for potential inputs at GnRH terminals in the median eminence has also been reported (Lehman et al., 2010a). Anyway, the presence of GPR54 at GnRH nerve terminal level has yet to be demonstrated.

Most work has emphasized the ability of kisspeptin neurons to mediate gonadal steroid feedback on GnRH release. Indistinctly from the specific kisspeptin population observed, all neurons co-localize estradiol (ER), progesterone (PR) and androgen (AR) receptors; particular attention concerns ER, since the greatest percentage of both ARC and AVPV kisspeptin neurons expresses the isoform ER α , in comparison with GnRH neurons that express the isoform β (Lehman et al., 2010b and references therein). In female, throughout most of the cycle, GnRH/LH secretion is under a negative feedback from ovarian steroids, with estradiol that controls pulse amplitude and progesterone pulse frequency. The feedback becomes positive at the end of the follicular phase to sustain pre-ovulatory GnRH/LH surge and the ovulation (Karsch, 1987). This sexually dimorphic event - only females of most species exhibit it - specifically depends on ER α and occurs *via* a classical

mechanism involving transcriptional regulation of gene expression. Of important note, AVPV kisspeptin neurons mediate this positive feedback triggering ovulation; in this respect, kisspeptin signalling might have potential therapeutic roles in the control of ovulation (Clarkson & Herbison, 2009). On the other hand, ARC KNDy neurons convey to GnRH neurons the estrogen negative feedback, at least in rodents, sheep and primates (Roa et al., 2008). Differently, in ewes, KNDy neurons alone govern both inhibitory and stimulatory estrogen effects on GnRH secretion, leaving open the question of how the same sub-population of neurons is able to discern these inputs; a possible explanation could lie in DYN that has the well known ability to mediate the inhibitory feedback of progesterone on GnRH secretion (Goodman et al., 2004). Although some aspects of ARC KNDy neurons have been elucidated, their physiology and their ability to integrate inputs in the control of reproduction are still partially unknown. The use of experimental approach such as cell ablation, that consists in the complete elimination of a whole cell type, can result in quite different phenotypes from complementary gene KO approach. This is the case of kisspeptin/GPR54 neuron ablation (Mayer & Boehm, 2011). The major upsetting observations coming from this study concern the loss of effect of kisspeptin/GPR54 ablation on the timing of puberty onset in female mice, suggesting a regulatory circuit upstream GnRH pulse, independent from kisspeptin signalling. Despite the formation of smaller ovaries, these animals are fertile and, even if GPR54 neuron ablation notably reduces the number of GnRH neurons to almost 10%, this limited percentage seems to be sufficient for a normal reproductive development. Only in the case of acute kisspeptin neuron ablation, animals show acyclicity and infertility, suggesting an essential role for kisspeptin neurons themselves, but not for kisspeptin signalling in ovulatory cyclicity (Mayer & Boehm, 2011).

Reproductive success strongly depends on energy balance, environmental and stressor cues (Bouret et al., 2004). In this perspective, ARC KNDy neurons have been suggested to convey metabolic and stressor information on HPG by means of the expression of glucocorticoid and leptin receptor - where leptin is a hormone secreted by white adipose tissue in proportion to the amount of body energy stores (Bouret et al., 2004, Kinsey-Jones et al., 2009). Conditions of hypoleptinemia have been linked to decreased hypothalamic expression of *kiss1* mRNA and to the inhibition of reproductive functions (Smith et al., 2006). Since GnRH neurons do not directly respond to nutritional conditions (Louis et al., 2011), current hypothesis is that kisspeptin neurons represent intermediate obligate pathways to transmit leptin actions to GnRH neurons that do not express leptin receptor (Castellano et al., 2009, Smith et al., 2006, Tena-Sempere, 2010).

Thus, not only kisspeptins represent a new well consolidated class of modulators of GnRH neuron activity, but also there is an intricate network of regulators that operate up-stream kisspeptin neurons making the study of their physiology much more complicate. Therefore, it is interesting to include in this scenario also the GnIH; it is remarkable that GnIH - through its ability to induce an increase of *kiss1* expression in the the ARC - could act up-stream and not always opposite to kisspeptin neurons, thus to completely upset the idea of the balance kisspeptin/GnIH conceived to date (Ancel et al., 2012).

<i>Evidence</i>	<i>Reference</i>
<i>- Hypothalamus</i>	
GnRH antagonists completely abrogate kisspeptin releasing effects on LH and FSH	Matsui et al., 2004; Navarro et al., 2005a; Navarro et al., 2005b
GnRH neurons in the rat forebrain express <i>GPR54</i> gene	Irwig et al., 2004
Kisspeptin induces <i>c-fos</i> expression in rodent GnRH neurons	Irwig et al., 2004
Kisspeptin induces long-lasting depolarization responses of GnRH neurons	Han et al., 2005
Kisspeptin loses its stimulatory effect upon LH secretion in GnRH-deficient <i>hpg</i> mice	Roa et al., 2009
<i>- Hypothalamus cell lines</i>	
GT1-7 cell lines - parentally related to GnRH neurons - express <i>GPR54</i> and respond to kisspeptin stimulation	Quaynor et al., 2007
<i>- Pituitary</i>	
<i>Kiss1</i> and <i>GPR54</i> expression is differentially regulated by steroids	Richard et al., 2008
<i>- Pituitary cell lines</i>	
<i>GPR54</i> is expressed in fractions of ovine pituitary cells enriched for gonadotropes, somatotropes and lactotropes	Smith et al., 2008b
Gonadotrope and somatotrope functions are regulated by kisspeptins in nonhuman primate pituitary	Luque et al., 2011
Kisspeptin inhibits LH expression in eel	Pasquier et al., 2011

This table provides a list of references in which the involvement of kisspeptins in the control of GnRH neuron activity as well as of gonadotropin discharge has been demonstrated.

Table 1. Experimental evidences supporting the mechanisms of action of kisspeptins.

As puberty, physiological conditions as pregnancy, lactation and aging suppose a strong contribution of kisspeptins in the regulation of gonadotropin secretion.

During pregnancy, in particular, hypothalamic *kiss1* expression increases and global state of hyper-responsiveness to kisspeptin, probably due to elevated levels of estradiol and progesterone, emerges. Moreover, circulating levels of kisspeptin - mainly derived from the placenta - dramatically increase, suggesting a possible involvement of this signalling in the regulation of trophoblast invasion during the first trimester (Horikoshi et al., 2003). Instead, the suckling stimulus during lactation might be responsible for the suppression of *kiss1* expression in the ARC and so for LH reduction (Yamada et al., 2007). Lastly, aging causes a high degree of reproductive system disruption, most obviously in females; this defect mainly concerns AVPV nucleus. Of note, during menopause, kisspeptin neurons are subjected to a high degree of hypertrophy (Downs & Wise, 2009).

Besides mammals, some important evidences concerning kisspeptin system have also been obtained in non-mammalian species. Recent advances have led to the identification of multiple isoforms of both *kiss* and *GPR54* in non-mammalian vertebrates, contrarily to mammals where only one gene coding for both the ligand and the receptor is present. This complication has been suggested as a consequence of genome duplication events (Tena-Sempere et al., 2012; Um et al., 2010). In this light, fish have two forms of kisspeptin (*kiss1* and *kiss2*) and *GPR54* (*GPR54-1* and *GPR54-2*), whereas in amphibians a second round of gene duplication may have contributed to the generation of subtypes for both ligand and receptor (*kiss1a*, -1*b*, -2; *GPR54-1a*, -1*b*, -2) (Tena-Sempere et al., 2012; Um et al., 2010). As previously suggested for GnRH, the presence of two kisspeptin systems might indicate a diversity of roles. Accordingly, zebrafish possess two independent kisspeptin systems, with *kiss2* mostly involved in the control of reproductive functions, through interactions with GnRH neurons, whereas *kiss1* is supposed to be implicated in the perception of environmental and metabolic signals (Servili et al., 2011). A possible relationship between food intake and kisspeptin signalling has been proposed in Senegalese sole, as well; this function has been attributed to *kiss2*, being the only isoform found in this specie, in contrast to the evident situation in most fish (Mechaly et al., 2011).

Lower vertebrates, as in particular fish models, strongly respond to environmental conditions, mostly thermal and light cues. In tilapia brain, *GPR54* co-localizes with neurons expressing GnRH1, GnRH2 and GnRH3, suggesting an involvement of kisspeptin system in the regulation of GnRH system expression as in mammals (Parhar et al., 2004). Higher levels of *GPR54* expression in the brain have been discovered in mature than in immature animals, thus to hypothesize a link between gonadal development and encephalic *GPR54* expression (Parhar et al., 2004). Similarly to mammals, *kiss1* gene has been identified in two distinct neuronal populations that exhibited a differential response to steroid milieu (Kanda et al., 2008). Furthermore, in fish, light has been shown to have a strong impact on kisspeptin system; in particular, long day - a typical permissive light condition - induces an increase in kisspeptin neuron number, supporting a role for this system in the mediation of light response and reproduction. Such a mediation has been evaluated in mammals, as well. In particular, *kiss1* mRNA and protein expression in ARC of Syrian hamsters decreases in short day condition, in correlation with decreased reproductive activity (Revel et al., 2006). Taken together these findings assign to kisspeptins an important role in the photoperiodic control of reproduction, processes under control of melatonin signals, whose actions on GnRH neurons are not direct. Currently, another wedge has been added to the picture, just in Syrian hamster with the demonstration that RFRP-3 is able to reactivate the reproductive axis blocked under photoinhibitory short-day conditions, thus suggesting that RFRP-3 could be the missing link between melatonin and kisspeptins (Ancel et al., 2012). In sheep, reproductive function is activated by short-day and inhibited by long-day photoperiods; during the breeding season, ARC shows an increase of *Kiss1* and a decrease of *RFRP* expression. Furthermore, the number of kisspeptin fibres onto GnRH neurons increases in comparison to that of RFRP, thus suggesting that these two peptides act in concert, with opposing effects, to regulate GnRH neuron activity (Smith et al., 2008a).

In goldfish, *in vivo* administration of kisspeptin induces LH release, indicating a conserved role from fish to mammals (Li et al., 2009). Anyway, the physiology of kisspeptin system seems to be rather controversial even among piscine species. This is the case of the eel, *Anguilla anguilla*, where kisspeptin - through a direct activity on the pituitary has an inhibitory - and not a stimulatory - effect on *LH β* expression; this effect is also specific, since no action on other pituitary glycoprotein hormone subunits has been shown (Pasquier et al., 2011).

In amphibian model, the presence of three isoforms for both ligand and receptor makes the evaluation of possible physiological roles for each component likely complex. As in fish, in amphibian all isoforms of kisspeptin and GPR54 are highly expressed in the brain, notably in the hypothalamus, allowing to hypothesize a conserved neuroendocrine and neuromodulatory role in the control of puberty onset and reproduction. Moreover, the expression of *GPR54-1a* and -2 in the pituitary would support a direct neuroendocrine action at pituitary level (Lee et al., 2009). In *Rana catesbeiana*, in particular, GPR54 is primarily expressed in the hypothalamus and pituitary, and weakly expressed in the testis. Of note, neither a faint signal has been detected in other peripheral tissues, such as heart, spleen, liver, supporting an exclusive role for kisspeptin system in the control of central functions (Moon et al., 2009).

Peripheral administration of kisspeptin also stimulates LH as well as GH and PRL secretion (Kadokawa et al., 2008; Yang et al., 2010). Therefore, an additional target of kisspeptin might be outside the blood-brain barrier (Matsui et al., 2004) and some evidences suggest autocrine/paracrine actions of kisspeptins at the pituitary level. Hence, indicative results as the presence of a functional kisspeptin receptor in the pituitary, combined with the finding that kisspeptin is released in hypophyseal portal blood, reinforce the idea that kisspeptin could be able of a dual action at both the hypothalamus and pituitary level. In support of this hypothesis, both *kiss* and *GPR54* are expressed in the pituitary of several species investigated to date (Richard et al., 2009); GPR54 has also been detected in ovine cellular fractions enriched of gonadotropes, somatotropes and lactotropes (Smith et al., 2008b). Once again, this duality suggested in many species is not the rule, since in a sheep model of hypothalamo-pituitary disconnection, kisspeptin loses its stimulatory action on LH secretion, thus assuming an exclusive effect upstream of the pituitary (Smith et al. 2008b). Anyway, this debate is still open and surely warrants further investigations.

3. Gonadic action of GnRH molecular forms

3.1. GnRH molecular forms and GnRHRs expression and activity in male and female gonads

An intriguing question is the synthesis of GnRH at gonadal level. Despite in hypophysectomised animals GnRH agonist administration induces steroidogenesis and GnRH like molecules have been detected in the main circulation of elasmobranches, in tetrapods the peptide has never been detected in plasma (King et al., 1992). Extra brain synthesis and function of GnRH - especially GnRH2 - and GnRHRs (both mRNA and

protein) have been detected in vertebrate gonads, humans included, endometrium, placenta, and endometrial cancer cells (Pierantoni et al., 2002; Ramakrishnappa et al., 2005; Singh et al., 2007; Wu et al., 2009).

In males, GnRH involvement in paracrine Sertoli/Leydig cell communication has been postulated in both mammals and lower vertebrates (Pierantoni et al., 2002; Sharpe 1986). In rodent and human testis, the main source of testicular GnRH are Sertoli cells, as well as spermatogenetic cells whereas GnRHR has been mainly located in interstitial Leydig cells (Bahk et al., 1995). Thus, GnRH likely acts as paracrine mediator for steroidogenesis and spermatogenesis progression. GnRH involvement in sperm release has also been reported since GnRH agonist/antagonist, induces/suppresses the spermiation in amphibians and lampreys (Deragon & Sower 1994; Pierantoni et al., 1984a, 1984b). Despite the presence of a functional GnRHR in spermatozoa is questionable, GnRH involvement in sperm function at fertilization has also been proposed, since GnRH antagonists inhibit *in vivo* and *in vitro* fertilization in rodents (Morales et al., 2002a). Most information concerning the local activity of GnRH derived from studies carried out in lower vertebrates, where, as in humans, GnRH2 is the main form detected at peripheral level (Pierantoni et al., 2002; White et al., 1998). However, it is not excluded that the existence of multiple forms of GnRH might be linked to the development of specific functions. For instance, in the amphibian, *Rana esculenta*, we have just cloned and characterized two GnRH molecular forms (GnRH1 and GnRH2) and three GnRHRs, with a specific expression pattern in testis, a specific testicular localization and probably, a specific function during the spermatogenesis (Chianese et al., unpublished). In the frog *Rana esculenta*, GnRH cooperates with estradiol in order to gain spermatogonial proliferation in a mechanism involving the protooncogene *c-fos* (Cobellis et al., 2002). During the frog annual sexual cycle, FOS protein appeared inside the cytoplasm of spermatogonia before the proliferative period, whereas it appeared inside the nucleus as soon as spermatogonia proliferation resumes (Cobellis et al., 2002). Estradiol, produced by Leydig cells, induces the transcription of *c-fos* inside the spermatogonia and the protein is stored in cytoplasmic compartment (Cobellis et al., 2002); then, at the end of the winter stasis, GnRH, produced by Sertoli cells, induces FOS activity by means of FOS protein traslocation from cytoplasmic to nuclear compartment (Cobellis et al., 2003) with a consequent increase of spermatogonial mitotic index. The involvement of *c-fos* as well as of estradiol in spermatogonial proliferation has recently been confirmed also in cell lines (He et al., 2008; Sirianni et al., 2008). However, GnRH role in spermatogonial proliferation is an evolutionarily conserved mechanism since it has been demonstrated also in bivalve mollusc (Treen et al., 2012).

As for testis, also in the ovary of mammalian and non-mammalian vertebrates GnRH binding sites as well as at least one GnRH molecular form and one GnRHR have been detected (Pierantoni et al., 2002). Besides testis, in the ovary GnRH activity highly depends on the state of maturation (Guilgur et al., 2009; Pierantoni et al., 2002; Uzbekova et al., 2002; Wu et al., 2009). In lamprey, GnRH induces both steroidogenesis and ovulation, whereas in teleost fish different GnRH molecular forms differentially modulate both meiosis resumption and steroidogenesis (Nabissi et al. 2000; Pati & Habibi, 2000). In rodents,

GnRHR expression rate and localization change during the reproductive cycle, with high expression levels observed in granulosa cells of atretic follicles as well as in mural granulosa cells of Graffian and preovulatory follicles (Bauer-Dantoin & Jameson, 1995; Kogo et al., 1995). Thus, GnRH seems to have a direct role in follicular atresia, a well known phenomenon of cell death. In *in vitro* cultures, GnRH inhibits DNA synthesis (Saragueta et al., 1997) or induces apoptosis in rat granulosa cells (Billig et al., 1994). Studies have shown the evidence for GnRH-induced remodelling of the extra cellular matrix by inducing structural luteolysis in superovulated rats through stimulation of specific matrix metalloproteinase (Goto et al., 1999). In human ovary, GnRH acts as an autocrine modulator of granulosa cells and has the ability to inhibit progesterone biosynthesis (Peng et al., 1994). Lastly, in addition to endocrine regulation, GnRH is also known to act in an autocrine and paracrine manner in order to suppress cell proliferation and to activate apoptosis in the endometrium and endometrial cancer cells through several mechanisms (Wu et al., 2009). In human ovary cell lines, such a mechanism involves GnRH2 and not GnRH1, and opens question of a functional GnRHR2 in humans (Leung et al., 2003). Hypothesis of remnant *GnRHR2* genes in human, mouse and rat genomes is reported in literature (Pawson et al., 2003); at present, it remains unknown whether or not GnRHR2 is expressed as a full-length, properly processed and functional transcript in humans.

However, both GnRH1 and GnRH2 exhibit regulatory roles in tissue remodelling during embryo implantation and placentation, which suggests that these hormones may have important roles in embryo implantation and early pregnancy (Wu et al., 2009).

3.2. Are ECBs and kisspeptin putative local modulators of GnRH/gonadotropin activity?

As for GnRH, also ECBs and kisspeptin biosynthesis and activity have been reported at gonadal levels, thus opening new questions in their possible local crosstalk. Besides the well known suppression of LH in both marijuana smokers and animal model, events due to hypothalamic GnRH suppression, ECBs deeply affect male and female reproductive functions (Wang et al., 2006; Wenger et al., 2001) and they have been detected in reproductive fluids (Schuel et al., 2002; Wang et al., 2006). In males, ECS modulates the progression of spermatogenesis, spermatozoa functions and the activity of testicular somatic cells in mammalian, non-mammalian vertebrates as well as in invertebrates. (Battista et al., 2012; Cacciola et al 2008; Cobellis et al., 2006; Cottone et al., 2008; Grimaldi et al., 2009; Maccarrone et al., 2003, 2005; Pierantoni et al., 2009, Schuel et al., 1991; Wang et al., 2006). In females, ECBs represent fertility signals in folliculogenesis, follicle maturation, oocyte maturation and ovulation (El-Talatini et al., 2009). Then, ECBs and CBs drive embryo transport, survival, implantation, development and growth, placentation and labour (Battista et al., 2012). Consistently, marijuana smokers exhibit several reproductive dysfunctions such as decreased LH levels in both sexes, decreased testosterone level, decreased sperm quality (oligospermia), sperm abnormality and block of acrosome reaction in males, whereas menstrual cycle disorders, reduced birth rates, preterm birth, low foetal birth weight have been described in women (Bari et al., 2011; Wang et al., 2006).

As for kisspeptin concerns, at present, the physiological significance of kisspeptin signalling at gonadal level is under investigation. Besides hypothalamic kisspeptin signalling is critical for puberty onset (Mayer & Boehm, 2011), human and rodent gonads express both *GPR54* and *kiss1* genes (Funes et al., 2003; Terao et al., 2004). With respect to GnRH and ECS, in rat ovary, in particular, *kiss1* mRNA expression shows a fluctuation dependent on the phase of the cycle, with a strong increase before ovulation and a dramatic decrease when ovary is at an immature state (Roa et al., 2007). Also in zebrafish females, *GPR54* expression follows the ovarian development with a decline of expression going toward the reproductive maturity; conversely, an increase of *kiss1* expression coincides with the appearance of mature oocytes (Biran et al., 2008; Filby et al., 2008).

At 7 weeks of age, *GPR54* KO mice display a reduced size of the internal and external reproductive organs with hypoplasia of seminiferous tubules, interstitial Leydig cells, uterine horns and mammary glands; these results let to hypothesize an involvement of *GPR54* in cell proliferation and differentiation that are properly necessary for gonadal development (Funes et al., 2003). Interestingly, in zebrafish male, high levels of both ligand and receptor expression have been observed during the first stages of spermatogenesis, when testis is mainly populated by type A spermatogonia to decrease after puberty (Biran et al., 2008; Filby et al., 2008). Kisspeptin expression is modulated by estradiol (Clarkson & Herbison, 2009) and estradiol cooperates with GnRH in order to induce proliferation of spermatogonia (Cobellis et al., 2003), thus raising the possibility of kisspeptin involvement in such a process. The presence of CB2 protein in mouse differentiating spermatogonia (Grimaldi et al., 2009) makes such an item an interesting issue for future investigations.

In disagreement with the general idea that kisspeptin signalling might have a positive impact on reproductive functions, also at a local level, a degenerative effect of kisspeptin administration on maturing rat testes has been reported (Ramzan & Qureshi, 2011). In particular, LH and testosterone suppression and severe degeneration of spermatogenesis in prepubertal testes in a dose-dependent manner have been reported. Such a disruption consists in: Sertoli cells impairment, meiosis inhibition concomitantly to increased spermatogonial proliferation. Already at low doses of kisspeptins, seminiferous tubules show intraepithelial vacuolizations that could be the cause of their massive degeneration, germ cells undergo necrosis, round and elongated spermatids have abnormal acrosome and the interstitial compartment is enlarged. Anyway, testicular degeneration observed after kisspeptin treatment has been suggested to be centrally mediated, and specifically due to an acute hyper-stimulation of the HPG axis (Thompson et al., 2009).

Worth mentioning, the negative effect of kisspeptin on testosterone secretion just reported in rats is in total disagreement with many other evidences provided in both rodents and humans where kisspeptin administration increases testosterone levels (Dhillon et al., 2005; Patterson et al., 2006). Such a point might represent a key switch in autocrine-paracrine communications in Leydig-Sertoli cells - germ cells circuitry involving also GnRH and ECS. AEA administration suppresses testosterone levels (Wenger et al., 2001). Such an issue is surely a consequence of hypothalamic GnRH inhibition, but it is not excluded a direct action

at testicular level. Leydig cells are the main source of testicular steroids and express both *GnRHR* and *CB1* (Bahk et al., 1995; Wenger et al 2001); Sertoli cells are the suggested source of GnRH and, in a FSH dependent fashion, modulate ECB tone and aromatase activity (Bahk et al., 1995; Rossi et al., 2007); germ cells are a suggested source of ECBs and have the ability to respond to ECBs and GnRH (Grimaldi et al., 2009). In such a story, the localization of kisspeptin/GPR54 inside the testis, to our knowledge, is completely lacking.

Really interesting might be the sequential activation of ECS, kisspeptin and GnRH in the diachronic process of epididymal sperm motility acquisition, post-ejaculatory events (capacitation and hyperactivation), and the capacity to recognize and to bind to the oocyte investments and egg plasma membrane. Mammalian spermatozoa acquire the ability to swim during their transit from the testis to the oviduct under the control of several external and intracellular factors. In vertebrates, spermatozoa possess a complete ECS and, at least in humans, evidence of kisspeptin system activity has been provided (Pinto et al., 2012; Wang et al., 2006). ECBs, *via* CB1, operate into the epididymis to regulate sperm motility acquisition and to prevent premature acrosome reaction (Cobellis et al., 2010; Ricci et al., 2007; Wang et al., 2006). To date, in species with external fertilization ECBs control the number of motile spermatozoa keeping sperm motility quiescent until their release in aquatic environment (Cobellis et al., 2006). *In vitro*, kisspeptin stimulates an irregular flagellar beating that is typical of a hyperactivation state, a condition critical for fertilization (Pinto et al., 2012). Then, it is intriguing to note that CB1, but not kisspeptin, controls the zona pellucida induced acrosome reaction (Wang et al., 2006) and GnRH increases sperm-zona pellucida binding in humans (Morales et al., 2002b).

4. Closing remarks

Astonishing progress has been accomplished the understanding of how intricate is the scenario that sustains the functionality of the reproductive axis. However, numerous new regulators are still emerging thus suggesting that many other key aspects have to be unravelled. A deep involvement of ECS and kisspeptin system in the modulation of GnRH activity clearly emerged at hypothalamic level. Nevertheless, a functional crosstalk between kisspeptin system, ECS and GnRH has never been investigated so far, neither in the brain nor in male and female gonads. At central level, ECBs and kisspeptins have opposite effects upon GnRH secreting neurons and glutamatergic glial cell surrounding GnRH neurons. In this respect, the balance between AEA and kisspeptin tone might represent a cooperative switch on/off signal for the activity of HPG axis. This aspect becomes more intriguing whether related to the existence of a complicate and multifunctional hypothalamic/gonadal GnRH system in non- mammalian vertebrates and humans.

Most work has attempted to rapidly decipher molecular mechanisms that control kisspeptin, ECBs and GnRH activity at peripheral level, but at moment many aspects of this debate are far from being fully elucidated and warrant further investigation.

Thus, altogether, the putative crosstalk among ECS, kisspeptin system and GnRH might provide a deep insight into the complex field of reproductive biology, opening the avenue to novel therapeutic approaches able to cure and prevent human infertility.

Author details

Rosaria Meccariello*

Dipartimento di Studi delle Istituzioni e dei Sistemi Territoriali, Università di Napoli Parthenope, Napoli, Italy

Rosanna Chianese, Silvia Fasano and Riccardo Pierantoni

Dipartimento di Medicina Sperimentale sez "F. Bottazzi", Seconda Università di Napoli, Napoli, Italy

Acknowledgement

Financial support by Italian Ministry of the University (COFIN 2008/BXCKKX to R.P) and Sorveglianza Sanitaria ex Esposti Mesoteliomi.

Abbreviations

AEA, anandamide; 2-AG, 2-arachidonoylglycerol; AR, androgen receptor; ARC, arcuate nucleus; AVPV, anteroventral periventricular nucleus; CBs, ECB receptors; ER, estradiol receptor; DAGL, diacylglycerol lipase; Δ^9 -THC, Δ^9 -tetrahydrocannabinol; DYN, dynorphin; ECBs, endocannabinoids; ECS, endocannabinoid system; EMT, ECB transporters; FABPs, fatty acid binding proteins; FAAH, fatty acid amide hydrolase; FLAT, FAAH-like AEA transporter; FSH, follicular Stimulating Hormone; GH, growth hormone; GnIH; gonadotropin-inhibitory hormone; GnRH, gonadotropin releasing hormone; HPG, hypothalamus-pituitary-gonadal axis; KNDy, Kisspeptin, Neurokinin B, Dynorphin cells; LH, Luteinizing Hormone; MAGL, monoacylglycerol lipase; NAAA, N-acylethanolamine-hydrolyzing acid amidase; NAPE-PLD, N-acyl phosphatidylethanolamine phospholipase D; NKB, neurokinin B; POA, preoptic area; PPAR, peroxisome-proliferator-activated receptor; PR, progesterone receptor; PRL, prolactin; RFRP-3, RFamide-related peptide.

5. References

- Ancel, C., Bentsen, AH., Sebert, ME., Tena-Sempere, M., Mikkelsen, JD., & Simmoneaux, V. (2012). Stimulatory effect of RFRP-3 on the gonadotrophic axis in the male Syrian Hamster: the exception proves the rule. *Endocrinol*, 153, 1352-1363.
- Bahk, JY., Hyun, JS., Chung, SH., Lee, H., Kim, MO., Lee, BH., & Choi, WS. (1995). Stage specific identification of the expression of GnRH mRNA and localization of the GnRH receptor in mature rat and adult human testis. *J Urol*, 154, 1958-1961.
- Bari, M., Battista, N., Pirazzi, V., & Maccarrone, M. (2011). The manifold actions of endocannabinoids on female and male reproductive events. *Front Biosci*, 16, 498-516.
- Battista, N., Meccariello, R., Cobellis, G., Fasano, S., Di Tommaso, M., Pirazzi, V., Konje, JC., Pierantoni, R., & Maccarrone, M. (2012). The role of endocannabinoids in gonadal

* Corresponding Author

- function and fertility along the evolutionary axis. *Mol Cell Endocrinol* [Epub ahead of print].
- Bauer-Dantoin, AC., & Jameson, JL. (1995). Gonadotropin-releasing hormone receptor messenger ribonucleic acid expression in the ovary during the rat estrous cycle. *Endocrinol*, 136, 4432-4438.
- Bern, HA. (1967). Hormones and endocrine glands in fishes. Studies on fish endocrinology reveal major physiologic and evolutionary problems. *Science*, 158, 455-462.
- Billig, H., Furuta, I., & Hsueh, AJ. (1994). Gonadotropin-releasing hormone directly induces apoptotic cell death in the rat ovary: biochemical and in situ detection of deoxyribonucleic acid fragmentation in granulosa cells. *Endocrinol*, 134, 245-252.
- Biran, J., Ben-Dor, S., & Levavi-Sivan, B. (2008). Molecular identification and functional characterization of the kisspeptin/kisspeptin receptor system in lower vertebrates. *Biol Reprod*, 79, 776-786.
- Bisogno, T., Howell, F., Williams, G., Minassi, A., Cascio, MG., Ligresti, A., Matias, I., Schiano-Moriello, A., Paul, P., Williams, EJ., Gangadharan, U., Hobbs, C., Di Marzo, V., & Doherty, P. (2003). Cloning of the first sn1-DAG lipases points to the spatial and temporal regulation of endocannabinoid signaling in the brain. *J Cell Biol*, 63, 463-468.
- Bouret, SG., Draper, SJ., & Simerly, RB. (2004). Trophic action of leptin on hypothalamic neurons that regulate feeding. *Science*, 304, 108-110.
- Brown, SM., Wager-Miller, J., & Mackie, K. (2002). Cloning and molecular characterization of the rat CB2 cannabinoid receptor. *Biochim Biophys Acta*, 1576, 255-264.
- Buznikov, GA., Nikitina, LA., Bezuglov, VV., Francisco, ME., Boysen, G., Obispo-Peak, IN., Peterson, RE., Weiss, ER., Schuel, H., Temple, BR., Morrow, AL., & Lauder, JM. (2010). A putative 'pre-nervous' endocannabinoid system in early echinoderm development. *Dev Neurosci*, 32, 1-18.
- Cacciola, G., Chioccarelli, T., Mackie, K., Meccariello, R., Ledent, C., Fasano, S., Pierantoni, R., & Cobellis, G. (2008). Expression of type-1 cannabinoid receptor during rat postnatal testicular development: Possible Involvement in adult Leydig cell differentiation. *Biol Reprod*, 79, 758-765.
- Castaño, JP., Martínez-Fuentes, AJ., Gutiérrez-Pascual, E., Vaudry, H., Tena-Sempere, M., & Malagón, MM. (2009). Intracellular signalling pathways activated by kisspeptins through GPR54: do multiple signals underlie function diversity? *Peptides*, 30, 10-15.
- Castellano, JM., Roa, J., Luque, RM., Dieguez, C., Aguilar, E., Pinilla, L., & Tena-Sempere, M. (2009). KiSS-1/kisspeptins and the metabolic control of reproduction: physiologic roles and putative physiopathological implications. *Peptides*, 30, 139-145.
- Cesa, R., Guastalla, A., Cottone, E., Mackie, K., Beltramo, M., & Franzoni, MF. (2002). Relationships between CB1 cannabinoid receptors and pituitary endocrine cells in *Xenopus laevis*: an immunohistochemical study. *Gen Comp Endocrinol*, 25, 17-24.
- Cheng, G., Coolen, LM., Padmanabhan, V., Goodman, RL., & Lehman, MN. (2010). The Kisspeptin/Neurokinin B/Dynorphin (KNDy) cell population of the arcuate nucleus: sex differences and effects of prenatal testosterone in sheep. *Endocrinol*, 151, 301-311.

- Chianese, R., Cobellis, G., Pierantoni, R., Fasano, S., & Meccariello, R. (2008). Non mammalian vertebrate models and the endocannabinoid system: relationships with gonadotropin-releasing hormone. *Mol Cell Endocrinol*, 286, S46-51.
- Chianese, R., Chioccarelli, T., Cacciola, G., Ciaramella, V., Fasano, S., Pierantoni, R., Meccariello, R., & Cobellis G. (2011a). The contribution of lower vertebrate animal models in human reproduction research. *Gen Comp Endocrinol*, 171, 17-27.
- Chianese, R., Ciaramella, V., Fasano, S., Pierantoni, R., & Meccariello, R. (2011b). Anandamide modulates the expression of *GnRH-II* and *GnRHRs* in frog, *Rana esculenta*, diencephalon. *Gen Comp Endocrinol*, 173, 389-395.
- Clarkson, J., & Herbison, AE. (2006). Postnatal development of kisspeptin neurons in mouse hypothalamus; sexual dimorphism and projections to gonadotropin-releasing hormone neurons. *Endocrinol*, 147, 5817-5825.
- Clarkson, J., & Herbison, AE. (2009). Oestrogen, kisspeptin, GPR54 and the pre-ovulatory luteinising hormone surge. *J Neuroendocrinol*, 21, 305-311.
- Clements, MK., McDonald, TP., Wang, R., Xie, G., O'Dowd, BF., George, SR., Austin, CP., & Liu, Q. (2001). FMRamide-related neuropeptides are agonists of the orphan G-protein-coupled receptor GPR54. *Biochem Biophys Res Commun*, 284, 1189-1193.
- Cobellis, G., Meccariello, R., Finga, G., Pierantoni, R., & Fasano, S. (2002). Cytoplasmic and nuclear Fos protein forms regulate resumption of spermatogenesis in the frog, *Rana esculenta*. *Endocrinol*, 143, 163-170.
- Cobellis, G., Meccariello, R., Minucci, S., Palmiero, C., Pierantoni, R., & Fasano, S. (2003). Cytoplasmic versus nuclear localization of Fos-related proteins in the frog, *Rana esculenta*, testis: *in vivo* and direct *in vitro* effect of a gonadotropin-releasing hormone agonist. *Biol Reprod*, 68, 954-960.
- Cobellis, G., Cacciola, G., Scarpa, D., Meccariello, R., Chianese, R., Franzoni, MF., Mackie, K., Pierantoni, R., & Fasano, S. (2006). Endocannabinoid system in frog and rodent testis: type-1 cannabinoid receptor and fatty acid amide hydrolase activity in male germ cells. *Biol Reprod*, 75, 82-89.
- Cobellis, G., Ricci, G., Cacciola, G., Orlando, P., Petrosino, S., Cascio, MG., Bisogno, T., De Petrocellis, L., Chioccarelli, T., Altucci, L., Fasano, S., Meccariello, R., Pierantoni, R., Ledent, C., & Di Marzo, V. (2010). A gradient of 2-arachidonoylglycerol regulates mouse epididymal sperm cell start-up. *Biol Reprod*, 82, 451-458.
- Cottone, E., Salio, C., Conrath, M., & Franzoni, MF. (2003). *Xenopus laevis* CB1 cannabinoid receptor: molecular cloning and mRNA distribution in the central nervous system. *J Comp Neurol*, 464, 487-496.
- Cottone, E., Forno, S., Campantico, E., Guastalla, A., Viltono, L., Mackie, K., & Franzoni, MF. (2005). Expression and distribution of CB1 cannabinoid receptors in the central nervous system of the African cichlid fish *Pelvicachromis pulcher*. *J Comp Neurol*, 485, 293-303.
- Cottone, E., Guastalla, A., Mackie, K., & Franzoni, MF. (2008). Endocannabinoids affect the reproductive functions in teleosts and amphibians. *Mol Cell Endocrinol*, 286, S41-S45.
- Cravatt, BF., Giang, DK., Mayfield, SP., Boger, DL., Lerner, RA., & Gilula, NB. (1996). Molecular characterization of an enzyme that degrades neuromodulatory fatty-acid amides. *Nature*, 384, 83-87.

- Deragon, K.L., & Sower, S.A. (1994). Effects of lamprey gonadotropin-releasing hormone-III on steroidogenesis and spermiation in male sea lampreys. *Gen Comp Endocrinol*, 95, 363-367.
- de Roux, N., Genin, E., Carel, J.C., Matsuda, F., Chaussain, J.L., & Milgrom, E. (2003). Hypogonadotropic hypogonadism due to loss of function of the Kiss1-derived peptide receptor GPR54. *Proc Natl Acad Sci USA*, 100, 10972-10976.
- Devane, W.A., Hanus, L., Breuer, A., Pertwee, R.G., Stevenson, L.A., Griffin, G., Gibson, D., Mandelbaum, A., Etinger, A., & Mechoulam, R. (1992). Isolation and structure of a brain constituent that binds to the cannabinoid receptor. *Science*, 258, 1946-1949.
- Dhillon, W.S., Chaudhri, O.B., Patterson, M., Thompson, E.L., Murphy, K.G., Badman, M.K., McGowan, B.M., Amber, V., Patel, S., Ghatei, M.A., & Bloom S.R. (2005). Kisspeptin-54 stimulates the hypothalamic-pituitary gonadal axis in human males. *J Clin Endocrinol Metab*, 90, 6609-6615.
- Dinh, T.P., Freund, T.F., & Piomelli, D. (2002). A role for monoglyceride lipase in 2-arachidonoylglycerol inactivation. *Chem Phys Lipids*, 121, 149-158.
- Downs, J.L., & Wise, P.M. (2009). The role of the brain in female reproductive aging. *Mol Cell endocrinol*, 299, 32-38.
- Dumalska, I., Wu, M., Morozova, E., Liu, R., van den Pol, A., & Alreja, M. (2008). Excitatory effects of the puberty-initiating peptide kisspeptin and group I metabotropic glutamate receptor agonists differentiate two distinct subpopulations of gonadotropin-releasing hormone neurons. *J Neurosci*, 28, 8003-8013.
- Egertová, M., Giang, D.K., Cravatt, B.F., & Elphick, M.R. (1998). A new perspective on cannabinoid signalling: complementary localization of fatty acid amide hydrolase and the CB1 receptor in rat brain. *Proc Biol Sci*, 265, 2081-2085.
- El-Talatini, M.R., Taylor, A.H., Elson, J.C., Brown, L., Davidson, A.C., & Konje, J.C. (2009). Localisation and function of the endocannabinoid system in the human ovary. *PLoS One*, 4, e4579.
- Estrada, K.M., Clay, C.M., Pompolo, S., Smith, J.T., & Clarke, I.J. (2006). Elevated Kiss-1 expression in the arcuate nucleus prior to the cyclic preovulatory gonadotrophin-releasing hormone/luteinising hormone surge in the ewe suggests a stimulatory role for kisspeptin in oestrogen-positive feedback. *J Neuroendocrinol*, 18, 806-809.
- Farkas, I., Kalló, I., Deli, L., Vida, B., Hrabovszky, E., Fekete, C., Moenter, S.M., Watanabe, M., & Liposits, Z. (2010). Retrograde endocannabinoid signaling reduces GABAergic synaptic transmission to Gonadotropin-Releasing Hormone neurons. *Endocrinol*, 151, 5818-5829.
- Fasano, S., Meccariello, R., Cobellis, G., Chianese, R., Cacciola, G., Chioccarelli, T., & Pierantoni, R. (2009). The Endocannabinoid System: An Ancient Signaling Involved in the Control of Male Fertility. *Ann NY Acad Sci*, 1163, 112-124.
- Fernandez-Fernandez, R., Martini, A.C., Navarro, V.M., Castellano, J.M., Dieguez, C., Aguilar, E., Pinilla, L., & Tena-Sempere, M. (2006). Novel signals for the integration of energy balance and reproduction. *Mol Cell Endocrinol*, 254-255, 127-132.
- Filby, A.L., van Aerle, R., Duitman, J.W., & Tyler, C.R. (2008). The Kisspeptin/gonadotropin-releasing hormone pathway and molecular signaling of puberty in fish. *Biol Reprod*, 78, 278-289.

- Fu, J., Bottegoni, G., Sasso, O., Bertorelli, R., Rocchia, W., Masetti, M., Guijarro, A., Lodola, A., Armirotti, A., Garau, G., Bandiera, T., Reggiani, A., Mor, M., Cavalli, A., & Pomelli, D. (2011). A catalytically silent FAAH-1 variant drives anandamide transport in neurons. *Nat Neurosci*, 15, 64-69.
- Funes, S., Hedrick, J.A., Vassleva, G., Markowitz, L., Abbondanzo, S., Golovko, A., Yang, S., Monsma, F.J., & Gustafson, E.L. (2003). The KISS-1 receptor GPR54 is essential for the development of the murine reproductive system. *Biochem Biophys Res Commun*, 312, 1357-1363.
- Gammon, C.M., Freeman, G.M. Jr., Xie, W., Petersen, S.L., & Wetsel, W.C. (2005). Regulation of gonadotropin-releasing hormone secretion by cannabinoids. *Endocrinol*, 146, 4491-4499.
- Galiegue, S., Mary, S., Marchand, J., Dussossoy, D., Carriere, D., Carayon, P., Bouaboula, M., Shire, D., Le Fur, G., & Casellas, P. (1995). Expression of central and peripheral cannabinoid receptors in human immune tissues and leukocyte subpopulations. *Eur J Biochem*, 232, 54-61.
- Glanowska, K.M., & Moenter, S.M. (2011). Endocannabinoids and prostaglandins both contribute to GnRH neuron-GABAergic afferent local feedback circuits. *J Neurophysiol*, 106, 3073-3081.
- González, S., Manzanares, J., Berrendero, F., Wenger, T., Corchero, J., Bisogno, T., Romero, J., Fuentes, J.A., Di Marzo, V., Ramos, J.A., & Fernández-Ruiz, J. (1999). Identification of endocannabinoids and cannabinoid CB(1) receptor mRNA in the pituitary gland. *Neuroendocrinol*, 70, 137-145.
- González, S., Bisogno, T., Wenger, T., Manzanares, J., Milone, A., Berrendero, F., Di Marzo, V., Ramos, J.A., & Fernández-Ruiz, J.J. (2000). Sex steroid influence on cannabinoid CB(1) receptor mRNA and endocannabinoid levels in the anterior pituitary gland. *Biochem Biophys Res Commun*, 270, 260-266.
- Goodman, R.L., Coolen, L.M., Anderson, G.M., Hardy, S.L., Valent, M., Connors, J.M., Fitzgerald, M.E., & Lehman, M.N. (2004). Evidence that dynorphin plays a major role in mediating progesterone negative feedback on gonadotropin-releasing hormone neurons in sheep. *Endocrinol*, 145, 2959-2967.
- Goto, T., Endo, T., Henmi, H., Kitajima, Y., Kiya, T., Nishikawa, A., Manase, K., Sato, H., & Kudo, R. (1999). Gonadotropin-releasing hormone agonist has the ability to induce increased matrix metalloproteinase (MMP)-2 and membrane type 1-MMP expression in corpora lutea, and structural luteolysis in rats. *J Endocrinol*, 161, 393-402.
- Grimaldi, P., Orlando, P., Di Sena, S., Lolicato, F., Petrosino, S., Bisogno, T., Geremia, R., De Petrocellis, L., & Di Marzo, V. (2009). The endocannabinoid system and pivotal role of the CB2 receptor in mouse spermatogenesis. *Proc Natl Acad Sci USA*, 106, 11131-11136.
- Guilgur, L.G., Strüßmann, C.A., & Somoza, G.M. (2009). mRNA expression of GnRH variants and receptors in the brain, pituitary and ovaries of pejerrey (*Odontesthes bonariensis*) in relation to the reproductive status. *Fish Physiol Biochem*, 35, 157-166.
- Han, S.K., Gottsch, M.L., Lee, K.J., Popa, S.M., Smith, J.T., Jakawich, S.K. (2005). Activation of gonadotropin-releasing hormone neurons by kisspeptin as a neuroendocrine switch for the onset of puberty. *J Neurosci*, 25, 11349-11356.

- He, Z., Jiang, J., Kokkinaki, M., Golestaneh, N., Hofmann, MC., & Dym, M. (2008). Gdnf upregulates c-Fos transcription via the Ras/Erk1/2 pathway to promote mouse spermatogonial stem cell proliferation. *Stem Cells*, 26, 266-278.
- Herbison, AE., & Pape, JR. (2001). New evidence for estrogen receptors in gonadotropin-releasing hormone neurons. *Front Neuroendocrinol*, 22, 292-308.
- Herbison, AE., d'Anglemont de Tassigny, X., Doran, J., & Colledge, WH. (2010). Distribution and postnatal development of Gpr54 gene expression in mouse brain and gonadotropin-releasing hormone neurons. *Endocrinol*, 151, 312-321.
- Herbison, AE., & Moenter, SM. (2011). Depolarising and hyperpolarising actions of GABA(A) receptor activation on gonadotrophin-releasing hormone neurones: towards an emerging consensus. *J Neuroendocrinol*, 23, 557-569.
- Ho, SY., Delgado, L., & Storch, J. (2002). Monoacylglycerol metabolism in human intestinal Caco-2 cells: evidence for metabolic compartmentation and hydrolysis. *J Biol Chem*, 277, 1816-1823.
- Horikoshi, Y., Matsumoto, H., Takatsu, Y., Ohtaki, T., Kitada, C., Usuki, S., & Fujino, M. (2003). Dramatic elevation of plasma metastin concentrations in human pregnancy: metastin as a novel placenta-derived hormone in humans. *J Clin Endocrinol Metab*, 88, 914-919.
- Irwig, MS., Fraley, GS., Smith, JT., Acohido, BV., Popa, SM., Cunningham, MJ., Gottsch, ML., Clifton, DK., & Steiner, RA. (2004). Kisspeptin activation of gonadotropin releasing hormone neurons and regulation of KISS-1 mRNA in the male rat. *Neuroendocrinol*, 80, 264-272.
- Kaczocha, M., Vivieca, S., Sun, J., Glaser, ST., & Deutsch, DG. (2012). Fatty acid-binding proteins transport N-acyl ethanolamines to nuclear receptors and are targets of endocannabinoid transport inhibitors. *J Biol Chem*, 287, 3415-3424.
- Kadokawa, H., Suzuki, S., & Hashizume, T. (2008). Kisspeptin-10 stimulates the secretion of growth hormone and prolactin directly from cultured bovine anterior pituitary cells. *Anim Reprod Sci*, 105, 404-408.
- Kah, O., Lethimonier, C., Somoza, G., Guilgur, LG., Vaillant, C., Lareyre, JJ. (2007). GnRH and GnRH receptors in metazoa: a historical, comparative, and evolutive perspectives. *Gen Comp Endocrinol*, 153, 346-364.
- Kanda, S., Akazome, Y., Matsunaga, T., Yamamoto, N., Yamada, S., Tsukamura, H., Maeda, K., & Oka, Y. (2008). Identification of Kiss-1 product kisspeptin and steroid-sensitive sexually-dimorphic kisspeptin neurons in medaka (*Oryzias latipes*). *Endocrinol*, 149, 2467-2476.
- Karsch, FJ. (1987). Central actions of ovarian steroids in the feedback regulation of pulsatile secretion of luteinizing hormone. *Annu Rev Physiol*, 49, 365-382.
- Kauffman, AS., Gottsch, ML., Roa, J., Byquist, AC., Crown, A., Clifton, DK., Hoffman, GE., Steiner, RA., & Tena-Sempere, M. (2007). Sexual differentiation of Kiss1 gene expression in the brain of the rat. *Endocrinol*, 148, 1774-1783.
- Kavanaugh, SI., Nozaki, M., & Sower, SA. (2008). Origins of gonadotropin-releasing hormone (GnRH) in vertebrates: identification of a novel GnRH in a basal vertebrate, the sea lamprey. *Endocrinol*, 149, 3860-3869.

- King, JA., Steneveld, AA., Millar, RP., Fasano, S., Romano, G., Spagnuolo, A., Zanetti, L., & Pierantoni, R. (1992). Gonadotropin-releasing hormone in elasmobranch (electric ray, *Torpedo marmorata*) brain and plasma: chromatographic and immunological evidence for chicken GnRH-II and novel molecular forms. *Peptides*, 13, 27-35.
- Kinsey-Jones, JS., Li, XF., Knox, AM., Wilkinson, ES., Zhu, XL., Chaudhary, AA., Millian, SR., Lightman, SL., & O'Byrne, KT. (2009). Down-regulation of hypothalamic kisspeptin and its receptor, Kiss1r, mRNA expression is associated with stress-induced suppression of luteinising hormone secretion in the female rat. *J Neuroendocrinol* 209, 21, 20-29.
- Kogo, H., Kudo, A., Park, MK., Mori, T., & Kawashima, S. (1995). In situ detection of gonadotropin-releasing hormone (GnRH) receptor mRNA expression in the rat ovarian follicles. *J Exp Zool*, 272, 62-68.
- Kotani, M., Detheux, M., Vandenbogaerde, A., Communi, D., vanderwinden, JM., Le Poul, E., Brezillon, S., Tyldesley, R., Suarez-Huerta, N., Vandeput, F., Blanpain, C., Schiffmann, SN., Vassart, G., & Parmentier, M. (2001). The metastasis suppressor gene KISS-1 encodes kisspeptins, the natural ligands of the orphan G protein-coupled receptor GPR54. *J Biol Chem*, 276, 34631-34636.
- Lam, CS., Rastegar, S., & Strähle, U. (2006). Distribution of cannabinoid receptor 1 in the CNS of zebrafish. *Neurosci*, 138, 83-95.
- Lapatto, R., Pallais, JC., Zhang, D., Chan, YM., Mahan, A., Cerrato, F., Le, WW., Hoffman, GE., & Seminara, SB. (2007). Kiss1^{-/-} mice exhibit more variable hypogonadism than Gpr54^{-/-} mice. *Endocrinol*, 148, 4927-4936.
- Lauckner, JE., Jensen, JB., Chen, HY., Lu, HC., Hille, B., & Mackie, K. (2008). GPR55 is a cannabinoid receptor that increases intracellular calcium and inhibits M current. *Proc Natl Acad Sci USA*, 105, 2699-2704.
- Lee, JH., Miele, ME., Hicks, DJ., Phillips, KK., Trent, JM., Weissman, BE., & Welch, DR. (1996). Kiss-1, a novel human malignant melanoma metastasis-suppressor gene. *J Natl Cancer Inst*, 88, 1731-1737.
- Lee, YR., Tsunekawa, K., Moon, MJ., Um, HN., Hwang, JI., Osugi, T., Otaki, N., Sunakawa, Y., Kim, K., Vaudry, H., Kwon, HB., Seong, JY., & Tsutsui, K. (2009). Molecular evolution of multiple forms of kisspeptins and GPR54 receptors in vertebrates. *Endocrinol*, 150, 2837-2846.
- Lehman, MN., Coolen, LM., & Goodman, RL. (2010a). Minireview: Kisspeptin/Neurokinin B/Dynorphin (KNDy) cells of the arcuate nucleus: a central node in the control of gonadotropin-releasing hormone secretion. *Endocrinol*, 151, 3479-3489.
- Lehman, MN., Merkley, CM., Coolen, LM., & Goodman, RL. (2010b). Anatomy of the kisspeptin neural network in mammals. *Brain Res*, 1364, 90-102.
- Leung, PC., Cheng, CK., & Zhu, XM. (2003). Multi-factorial role of GnRH-I and GnRH-II in the human ovary. *Mol Cell Endocrinol*, 202, 145-153.
- Li, S., Zhang, Y., Liu, Y., Huang, X., Huang, W., Lu, D., Zhu, P., Shi, Y., Cheng, CH., Liu, X., & Lin, H. (2009). Structural and functional multiplicity of the kisspeptin/GPR54 system in goldfish (*Carassius auratus*). *J Endocrinol*, 201, 407-418.

- Louis, GW., Greenwald-Yarnell, M., Phillips, R., Coolen, LM., Lehman, MN., & Myers, MG Jr. (2011). Molecular mapping of the neural pathways linking leptin to the neuroendocrine reproductive axis. *Endocrinol*, 152, 2302-2310.
- Luque, RM., Córdoba-Chacón, J., Gahete, MD., Navarro, VM., Tena-Sempere, M., Kineman, RD., & Castaño JP. (2011). Kisspeptin regulates gonadotroph and somatotroph function in nonhuman primate pituitary via common and distinct signaling mechanisms. *Endocrinol*, 152, 957-966.
- Lynn, AB., & Herkenham, M. (1994). Localization of cannabinoid receptors and nonsaturable high-density cannabinoid binding sites in peripheral tissues of the rat: implications for receptor-mediated immune modulation by cannabinoids. *J Pharmacol Exp Ther*, 268, 1612-1623.
- Maccarrone, M., Cecconi, S., Rossi, G., Battista, N., Pauselli, R., & Finazzi-Agrò, A. (2003). Anandamide activity and degradation are regulated by early postnatal aging and follicle-stimulating hormone in mouse Sertoli cells. *Endocrinol*, 144, 20-28.
- Maccarrone, M., Barboni, B., Paradisi, A., Bernabò, N., Gasperi, V., Pistilli, MG., Fezza, F., Lucidi, P., & Mattioli, M. (2005). Characterization of the endocannabinoid system in boar spermatozoa and implications for sperm capacitation and acrosome reaction. *J Cell Sci*, 118, 4393-4404.
- Matsuda, LA., Lolait, SJ., Brownstein, MJ., Young, AC., & Bonner, TI. (1990). Structure of a cannabinoid receptor and functional expression of the cloned cDNA. *Nature*, 346, 561-564.
- Matsui, H., Takatsu, Y., Kumano, S., Matsumoto, H., & Ohtaki, T. (2004). Peripheral administration of metastatin induces marked gonadotropin release and ovulation in the rat. *Biochem Biophys Res Commun*, 320, 383-388.
- Mayer, C., & Boehm, U. (2011). Female reproductive maturation in the absence of kisspeptin/GPR54 signaling. *Nat Neurosci*, 14, 704-710.
- McPartland, JM., Matias, I., Di Marzo, V., & Glass, M. (2006). Evolutionary origins of the endocannabinoid system. *Gene*, 370, 64-74.
- Meccariello, R., Franzoni, MF., Chianese, R., Cottone, E., Scarpa, D., Donna, D., Cobellis, G., Guastalla, A., Pierantoni, R., & Fasano, S. (2008). Interplay between the endocannabinoid system and GnRH-I in the forebrain of the anuran amphibian *Rana esculenta*. *Endocrinol*, 149, 2149-2158.
- Mechaly, AS., Vinas, J., & Piferrer, F. (2011). Gene structure analysis of kisspeptin-2 (*kiss2*) in the Senegalese sole (*Solea senegalensis*): characterization of two splice variants of *kiss2*, and novel evidence for metabolic regulation of kisspeptin signaling in non-mammalian species. *Mol Cell Endocrinol*, 339, 14-24.
- Migliarini, B., Marucci, G., Ghelfi, F., & Carnevali, O. (2006). Endocannabinoid system in *Xenopus laevis* development: CB1 receptor dynamics. *FEBS Lett*, 580, 1941-1945.
- Mikkelsen, JD., & Simonneaux, V. (2009). The neuroanatomy of the kisspeptin system in the mammalian brain. *Peptides*, 30, 26-33.
- Moon, JS., Lee, YR., Oh, DY., Hwang, JI., Lee, JY., Kim, JI., Vaudry, H., Kwon, HB., & Seong, JY. (2009). Molecular cloning of the bullfrog kisspeptin receptor GPR54 with high sensitivity to *Xenopus* kisspeptin. *Peptides*, 30, 171-179.

- Morales, P., Pasten, C., & Pizzarro, E. (2002a). Inhibition of *in vivo* and *in vivo* fertilization in rodents by gonadotropin-releasing hormone antagonist. *Biol Reprod*, 67, 1360-1365.
- Morales, P., Pizarro, E., Kong, M., & Pasten, C. (2002b). Sperm binding to the human zona pellucida and calcium influx in response to GnRH. *Andrologia*, 34, 301-307.
- Muir, AI., Chamberlain, L., Elshourbagy, NA., Michalovich, D., Moore, DJ., Calamari, A., Szekeres, PG., Sarau, HM., Chambers, JK., Murdock, P., Steplewski, K., Shabon, U., Miller, JE., Middleton, SE., Darker, JG, Larminie, CG., Wilson, S., Bergsma, DJ., Emson, P., Faull, R., Philpott, KL., & Harrison, DC. (2001). AXOR12, a novel human G protein-coupled receptor, activated by the peptide Kiss-1. *J Biol Chem*, 276, 28969-28975.
- Munro, S., Thomas, KL., & Abu-Shaar, M. (1993). Molecular characterization of a peripheral receptor for cannabinoids. *Nature*, 365, 61-65.
- Murphy, LL., Muñoz, RM., Adrian, BA., & Villanúa, MA. (1998). Function of cannabinoid receptors in the neuroendocrine regulation of hormone secretion. *Neurobiol Dis*, 5, 432-446.
- Nabissi, M., Soverchia, L., Polzonetti-Magni, AM., & Habibi, HR. (2000). Differential splicing of three gonadotropin-releasing hormone transcript in the ovary of the sea bream *Sparus aurata*. *Biol Reprod*, 62, 1329-1334.
- Navarro, VM., Castellano, JM., Fernandez-Fernandez, R., Tovar, S., Roa, J., Majen, A., Barreiro, ML., Casanueva, FF., Aguilar, E., Dieguez, C., Pinilla, L., Tena-Sempere, M. (2005a). Effects of Kiss-1 peptide, the natural ligand of GPR54, on follicle-stimulating hormone secretion in the rat. *Endocrinol*. 146, 1689-1697.
- Navarro, VM., Castellano, JM., Fernández-Fernández, R., Tovar, S., Roa, J., Mayen, A., Nogueiras, R., Vazquez, MJ., Barreiro, ML., Magni, P., Aguilar, E., Dieguez, C., Pinilla, L., Tena-Sempere, M. (2005b). Characterization of the potent luteinizing hormone-releasing activity of Kiss-1 peptide, the natural ligand of GPR54. *Endocrinol*, 146, 156-163.
- Oakley, AE., Clifton, DK., Steiner, RA. (2009). Kisspeptin signalin in the brain. *Endocr Rev*, 30, 713-743.
- Ohtaki, T., Shintani, Y., Honda, S., Matsumoto, H., Hori, A., Kanehashi, K., Terao, Y., Kumano, S., Takatsu, Y., Masuda, Y., Ishibashi, Y., Watanabe, T., Asada, M., Yamada, T., Suenaga, M., Kitada, C., Usuki, S., Kurokawa, T., Onda, H., Nishimura, O., & Fujino, M. (2001). Metastasis suppressor gene KISS-1 encodes peptide ligand of a G protein-coupled receptor. *Nature*, 411, 613-617.
- Okamoto, Y., Morishita, J., Tsuboi, K., Tonai, T., & Ueda, N. (2004). Molecular characterization of a phospholipase D generating anandamide and its congeners. *J Biol Chem*, 279, 5298-5305.
- Oláh, M., Mилоh, H., & Wenger, T. (2008). The role of endocannabinoids in the regulation of luteinizing hormone and prolactin release. Differences between the effects of AEA and 2AG. *Mol Cell Endocrinol*, 286, S36-40.
- O'Sullivan, S.E. (2007). Cannabinoids go nuclear: evidence for activation of peroxisome proliferator-activated receptors. *Br J Pharmacol*, 152, 576-582.
- Pagotto, U., Marsicano, G., Cota, D., Lutz, B., & Pasquali, R. (2006). The emerging role of the endocannabinoid system in endocrine regulation and energy balance. *Endocrine Rev*, 27, 73-100.

- Parhar, IS., Ogawa, S., & Sakuma, Y. (2004). Laser-captured single digoxigenin-labeled neurons of gonadotropin-releasing hormone types reveal a novel G protein-coupled receptor (GPR54) during maturation in cichlid fish. *Endocrinol*, 145, 3613-3618.
- Pasquier, J., Lafont, A-G., Leprince, J., Vaudry, H., Rousseau, K., & Dufour, S. (2011). First evidence for a direct inhibitory effect of kisspeptins on LH expression in the eel, *Anguilla anguilla*. *Gen Comp Endocrinol*, 173, 216-225.
- Pati, D., & Habibi, HR. (2000). Direct action of GnRH variants on goldfish oocyte meiosis and follicular steroidogenesis. *Mol Cell Endocrinol*, 160, 75-88.
- Patterson, M., Murphy, K.G., Thompson, E.L., Patel, S., Ghatei, M.A., & Bloom, S.R. (2006). Administration of kisspeptin-54 into discrete regions of the hypothalamus potently increases plasma luteinising hormone and testosterone in male adult rats. *J Neuroendocrinol* 2006, 18, 349-354.
- Pawson, AJ., Morgan, K., Maudsley, SR., & Millar, RP. (2003). Type II gonadotrophin-releasing hormone (GnRH-II) in reproductive biology. *Reprod*, 126, 271-278.
- Peng, C., Fan, NC., Ligier, M., Väänänen, J., & Leung, PC. (1994). Expression and regulation of gonadotropin-releasing hormone (GnRH) and GnRH receptor messenger ribonucleic acids in human granulosa-luteal cells. *Endocrinol*, 135, 1740-1746.
- Pierantoni, R., Fasano, S., Di Matteo, L., Minucci, S., Varriale, B., & Chieffi, G. (1984a). Stimulatory effect of a GnRH agonist (buserelin) in *in vitro* and *in vivo* testosterone production by the frog (*Rana esculenta*) testis. *Mol Cell Endocrinol*, 38, 215-219.
- Pierantoni, R., Iela, L., d'Istria, M., Fasano, S., Rastogi, RK., & Delrio, G. (1984b). Seasonal testosterone profile and testicular responsiveness to pituitary factors and gonadotrophin releasing hormone during two different phases of the sexual cycle of the frog (*Rana esculenta*). *J Endocrinol*, 102, 387-392.
- Pierantoni, R., Cobellis, G., Meccariello, R., & Fasano, S. (2002). Evolutionary aspects of cellular communication in the vertebrate hypothalamo-hypophysio-gonadal axis. *Int Rev Cytol*, 218, 69-141.
- Pierantoni, R., Cobellis, G., Meccariello, R., Cacciola, G., Chianese, R., Chioccarelli, T., & Fasano, S. (2009). CB1 Activity in Male Reproduction: Mammalian and Nonmammalian Animal Models. *Vitam Horm*, 81, 367-387.
- Pinto, FM., Cejudo-Román, A., Ravina, CG., Fernández-Sánchez, M., Martín-Lozano, D., Illanes, M., Tena-Sempere, M., & Candenas, ML. (2012). Characterization of the kisspeptin system in human spermatozoa. *Int J Androl*, 35, 63-73.
- Quaynor, S., Hu, L., Leung, PK., Feng, H., Mores, N., Krsmanovic, LZ., & Catt, KJ. (2007). Expression of a functional g protein-coupled receptor 54-kisspeptin autoregulatory system in hypothalamic gonadotropin-releasing hormone neurons. *Mol Endocrinol*, 21, 3062-3070.
- Ramakrishnappa, N., Rajamahendran, R., Lin, YM., & Leung, PC. (2005). GnRH in non-hypothalamic reproductive tissues. *Anim Reprod Sci*, 88, 95-113.
- Ramzan, F., & Qureshi, IZ. (2011). Intraperitoneal kisspeptin-10 administration induces dose-dependent degenerative changes in maturing rat testes. *Life Sci*, 88, 246-256.

- Revel, FG., Saboureau, M., Masson-Pévet, M., Pévet, P., Mikkelsen, JD., & Simonneaux V. (2006). Kisspeptin mediates the photoperiodic control of reproduction in hamsters. *Curr Biol*, 16, 1730-1735.
- Ricci, G., Cacciola, G., Altucci, L., Meccariello, R., Pierantoni, R., Fasano, S., & Cobellis, G. (2007). Endocannabinoid control of sperm motility: the role of epididymus. *Gen Comp Endocrinol*, 153, 320-322.
- Richard, N., Galmiche, G., Corvaisier, S., Caraty, A., & Kottler, ML. (2008). Kiss-1 and GPR54 genes are co-expressed in rat gonadotrophs and differentially regulated in vivo by oestradiol and gonadotrophin-releasing hormone. *J Neuroendocrinol*, 20, 381-393.
- Richard, N., Corvaisier, S., Camacho, E., & Kottler, ML. (2009). Kiss-1 and GPR54 at the pituitary level: overview and recent insights. *Peptides*, 30, 123-129.
- Roa, J., & Tena-Sempere M. (2007). Kiss-1 system and reproduction: comparative aspects and roles in the control of female gonadotropic axis in mammals. *Gene Comp Endocrinol*, 153, 132-140.
- Roa, J., Aguilar, E., Dieguez, C., Pinilla, L., & Tena-Sempere, M. (2008). New frontiers in kisspeptins/GPR54 physiology as fundamental gatekeepers of reproductive function. *Front Neuroendocrinol*, 29, 48-69.
- Roa, J., Castellano, JM., Navarro, VM., Handelsman, DJ., Pinilla, L., & Tena-Sempere, M. (2009). Kisspeptins and the control of gonadotropin secretion in male and female rodents. *Peptides*, 30, 57-66.
- Rossi, G., Gasperi, V., Paro, R., Barsacchi, D., Cecconi, S., & Maccarrone, M. (2007). Follicle-stimulating hormone activates fatty acid amide hydrolase by protein kinase A and aromatase-dependent pathways in mouse primary Sertoli cells. *Endocrinol*, 148, 1431-1439.
- Saragüeta, PE., Lanuza, GM., & Barañao, JL. (1997). Inhibitory effect of gonadotrophin-releasing hormone (GnRH) on rat granulosa cell deoxyribonucleic acid synthesis. *Mol Reprod Dev*, 47, 170-174.
- Schuel, H., Chang, MC., Berkery, D., Schuel, R., Zimmerman, AM., & Zimmerman S. (1991). Cannabinoids inhibit fertilization in sea urchins by reducing the fertilizing capacity of sperm. *Pharmacol Biochem Behav*, 40, 609-15.
- Schuel, H., Burkman, LJ., Lippes, J., Crickard, K., Forester, E., Piomelli, D., & Giuffrida, A. (2002). N-Acylethanolamines in human reproductive fluids. *Chem Phys Lipids*, 121, 211-227.
- Scorticati, C., Fernandez-Solari, J., De Laurentiis, A., Mohn, C., Prestifilippo, JP., Lasaga, M., Seilicovich, A., Billi, S., Franchi, A., McCann, S., & Rettori, V. (2004). The inhibitory effect of anandamide on luteinizing hormone-releasing hormone secretion is reversed by oestrogen. *Proc Natl Acad Sci USA*, 32, 11891-11896.
- Seminara, SB., Messenger, S., Chatzidaki, EE., Thresher, RR., Acierno, JS. Jr, Shagoury, JK., Bo-Abbas, Y., Kuohung, W., Schwinof, KM., Hendrick, AG., Zahn, D., Dixon, J., Kaiser, UB., Slaugenhaupt, SA., Gusella, JF., O'Rahilly, S., Carlton, MB., Crowley, WF. Jr, Aparicio, SA., & Colledge, WH. (2003). The GPR54 gene as a regulator of puberty. *N Engl J Med*, 349, 1614-1627.
- Servili, A., Le Page, Y., Leprince, J., Caraty, A., Escobar, S., Parhar, IS., Seong, JY., Vaudry, H., & Kah, O. (2011). Organization of two independent kisspeptin systems derived from evolutionary-ancient kiss genes in the brain of zebrafish. *Endocrinol*, 152, 1527-1540.

- Sharpe, RM. (1986). Paracrine control of the testis. *Clin Endocrinol Metab*, 15, 185-207.
- Shire, D., Carillon, C., Kaghad, M., Calandra, B., Rinaldi-Carmona, M., Le Fur, G., Caput, D., & Ferrara, P. (1995). An amino-terminal variant of the central cannabinoid receptor resulting from alternative splicing. *J Biol Chem*, 270, 3726-3731.
- Singh, P., Krishna, A., Sridaran, R. (2007). Localization of gonadotrophin-releasing hormone I, bradykinin and their receptors in the ovaries of non-mammalian vertebrates. *Reprod*, 133, 969-981.
- Sirianni, R., Chimento, A., Ruggiero, C., De Luca, A., Lappano, R., Andò, S., Maggiolini, M., Pezzi, V. (2008). The novel estrogen receptor, G protein-coupled receptor 30, mediates the proliferative effects induced by 17beta-estradiol on mouse spermatogonial GC-1 cell line. *Endocrinol*, 149, 5043-5051.
- Smith, JT., Acohido, BV., Clifton, DK., & Steiner, RA. (2006). Kiss-1 neurons are direct target for leptin in the ob/ob mouse. *J Neuroendocrinol*, 18, 298-303.
- Smith, JT., Coolen, LM., Kriegsfeld, LJ., Sari, IP., Jaafarzadehshirazi, MR., Maltby, M., Bateman, K., Goodman, RL., Tilbrook, AJ., Ubuka, T., Bentley, GE., Clarke, IJ., & Lehman, MN. (2008a). Variation in kisspeptin and RFamide-related peptide (RFRP) expression and terminal connections to gonadotropin-releasing hormone neurons in the brain: a novel medium for seasonal breeding in the sheep. *Endocrinol*, 149, 5770-5782.
- Smith, JT., Rao, A., Pereira, A., Caraty, A., Millar RP., & Clarke, IJ. (2008b). Kisspeptin is present in ovine hypophysial portal blood does not increase during the preovulatory luteinizing hormone surge: evidence that gonadotropes are not direct targets of kisspeptin in vivo. *Endocrinol*, 149, 1951-1959.
- Sugiura, T., Kondo, S., Sukagawa, A., Nakane, S., Shinoda, A., Itoh, K., Yamashita, A., & Waku, K. (1995). 2-Arachidonoylglycerol: a possible endogenous cannabinoid receptor ligand in brain. *Biochem Res Commun*, 215, 89-97.
- Sun, Y., & Bennett, A. (2007). Cannabinoids: A New Group of Agonists of PPARs. *PPAR Res*. Vol 2007;ARTICLE ID:23513.
- Tena-Sempere M. (2010). Kisspeptin signalling in the brain: recent developments and future challenges. *Mol Cell Endocrinol*, 314, 164-169.
- Tena-Sempere, M., Felip, A., Gomez, A., Zanuy, S., & Carillo, M. (2012). Comparative insights of the kisspeptin/kisspeptin receptor system: lessons from non-mammalian vertebrates. *Gen Comp Endocrinol*, 175, 234-243.
- Terao, Y., Kumano, S., Takatsu, Y., Hattori, M., Nishimura, A., Ohtaki, T., & Shintani Y. (2004). Expression of KISS-1, a metastasis suppressor gene, in trophoblast giant cells of the rat placenta. *Biochim Biophys Acta*, 1678, 102-110.
- Thompson, EL., Amber, V., Stamp, GW., Patterson, M., Curtis, AE., Cooke, JH., Appleby, GF., Dhillon, WS., Ghatei, MA., Bloom, SR., & Murphy, KG. (2009). Kisspeptin-54 at high doses acutely induces testicular degeneration in adult male rats via central mechanisms. *Br J Pharmacol*, 156, 609-625.
- Todman, MG., Han, SK., & Herbison, AE. (2005). Profiling neurotransmitter receptor expression in mouse gonadotropin-releasing hormone neurons using green fluorescent protein-promoter transgenics and microarrays. *Neurosci*, 132, 703-712.

- Topaloglu, AK., Reimann, F., Guclu, M., Yalin, AS., Kotan, LD., Porter, KM., Serin, A., Mungan, NO., Cook, JR., Ozbek, MN., Imamoglu, S., Akalin, NS., Yuksel, B., O'Rahilly, S., & Semple, RK. (2009). TAC3 and TACR3 mutations in familial hypogonadotropic hypogonadism reveal a key role for Neurokinin B in the central control of reproduction. *Nat Gen*, 41, 354-358.
- Treen, N., Itoh, N., Miura, H., Kikuchi, I., Ueda, T., Takahashi, KG., Ubuka, T., Yamamoto, K., Sharp, PJ., Tsutsui, K., & Osada, M. (2012). Mollusc gonadotropin-releasing hormone directly regulates gonadal functions: A primitive endocrine system controlling reproduction. *Gen Comp Endocrinol* [Epub ahead of print].
- Tsuboi, K., Sun, YX., Okamoto, Y., Araki, N., Tonai, T., & Ueda, N. (2005). Molecular characterization of N-acylethanolamine-hydrolyzing acid amidase, a novel member of the choloylglycine hydrolase family with structural and functional similarity to acid ceramidase. *J Biol Chem*, 280, 11082-11092.
- Ueda, N., Tsuboi, K., & Uyama, T. (2010). Enzymological studies on the biosynthesis of N-acylethanolamines. *Biochim Biophys Acta*, 1801, 1274-1285.
- Um, HN., Han, JM., Hwang, JI., Hong, SI., Vaudry, H., & Seong, JY. (2010). Molecular coevolution of kisspeptins and their receptors from fish to mammals. *Ann N Y Acad Sci*, 1200, 67-74.
- Uzbekova, S., Lareyre, JJ., Madigou, T., Davail, B., Jalabert, B., & Breton, B. (2002). Expression of prepro-GnRH and GnRH receptor messengers in rainbow trout ovary depends on the stage of ovarian follicular development. *Mol Reprod Dev*, 62, 47-56.
- Valenti, M., Cottone, E., Martinez, R., De Pedro, N., Rubio, M., Viveros, MP., Franzoni, MF., Delgado, MJ., & Di Marzo, V. (2005). The endocannabinoid system in the brain of *Carassius auratus* and its possible role in the control of food intake. *J Neurochem*, 95, 662-672.
- van der Stelt, M., & Di Marzo, V. (2004). Endovanilloids. Putative endogenous ligands of transient receptor potential vanilloid 1 channels. *Eur J Biochem*, 271, 1827-1834.
- van der Stelt, M., & Di Marzo, V. (2005). Anandamide as an intracellular messenger regulating ion channel activity. *Prostaglandins Other Lipid Mediat*, 77, 111-122.
- Vida, B., Hrabovsky, E., Caraty, A., Ciofi, P., Coen, CW., Liposits, Z., & Kallo, Dr I. (2009). Gender differences in the co-localisation of neuropeptides with kisspeptin in the hypothalamic neurons of the mouse brain. In Annual Meeting of the Society of Neuroscience. Vol Ed, Chicago, IL. pp. Poster 865.9.
- Waleh, NS., Cravatt, BF., Apte-Deshpande, A., Terao, A., & Kilduff, TS. (2002). Transcriptional regulation of the mouse fatty acid amide hydrolase gene. *Gene*, 291, 203-210.
- Wang, H., Dey, SK., & Maccarrone, M. (2006). Jekyll and Hyde: two faces of cannabinoid signalling in male and female fertility. *Endocr Rev*, 27, 427-448.
- Wei, BQ., Mikkelsen, TS., McKinney, MK., Lander, ES., & Cravatt, BF. (2006). A second fatty acid amide hydrolase with variable distribution among placental mammals. *J Biol Chem*, 281, 36569-36578.
- Wenger, T., Fernández-Ruiz, JJ., & Ramos, JA. (1999). Immunocytochemical demonstration of CB1 cannabinoid receptors in the anterior lobe of the pituitary gland. *J Neuroendocrinol*, 11, 873-878.

- Wenger, T., Jamali, KA., Juaneda, C., Bascsy, E., & Tramu, G. (2000). The endogenous cannabinoid, anandamide regulates anterior pituitary secretion in vitro. *Addiction Biol*, 5, 59-64.
- Wenger, T., Ledent, C., Csernus, V., & Gerendai, I. (2001). The central cannabinoid receptor inactivation suppresses endocrine reproductive functions. *Biochem Biophys Res Commun*, 284, 363-368.
- White, RB., Eisen, JA., Kasten, TL., & Fernald, RD. (1998). Second gene for gonadotropin-releasing hormone in humans. *Proc Natl Acad Sci USA*, 95, 305-309.
- Wu, HM., Wang, HS., Huang, HY., Soong, YK., MacCalman, CD., & Leung, PC. (2009). GnRH signaling in intrauterine tissues. *Reprod*, 137, 769-777.
- Yamada, S., Uenoyama, Y., Kinoshita, M., Iwata, K., Takase, K., Matsui, H., Adachi, S., Inoue, K., Maeda, KI., & Tsukamura, H. (2007). Inhibition of metastin (kisspeptin-54)-GPR54 signaling in the arcuate nucleus-median eminence region during lactation in rats. *Endocrinol*, 148, 2226-2232.
- Yang, B., Jiang, Q., Chan, T., Ko, WK., & Wong, AO. (2010). Goldfish kisspeptin: molecular cloning, tissue distribution of transcript expression, and stimulatory effects on prolactin, growth hormone and luteinizing hormone secretion and gene expression via direct actions at the pituitary level. *Gen Comp Endocrinol*, 165, 60-71.
- Yasuo, S., Unfried, C., Kettner, M., Geisslinger, G., & Korf, HW. (2010a). Localization of an endocannabinoid system in the hypophysial pars tuberalis and pars distalis of man. *Cell Tissue Res*, 342, 273-281.
- Yasuo, S., Koch, M., Schmidt, H., Ziebell, S., Bojunga, J., Geisslinger, G., & Korf, HW. (2010b). An endocannabinoid system is localized to the hypophysial pars tuberalis of Syrian hamsters and responds to photoperiodic changes. *Cell Tissue Res*, 40, 127-136.