We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Role of Adipose Secreted Factors and Kisspeptin in the Metabolic Control of Gonadotropin Secretion and Puberty¹

Clay A. Lents, C. Richard Barb and Gary J. Hausman

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/48802>

1. Introduction

1.1. Adipose tissue as an endocrine organ

Recent investigations from many species continue to reinforce and validate adipose tissue as an endocrine organ that impacts physiological mechanisms and whole-body homeostasis. Factors secreted by adipose tissue or “adipokines” continue to be discovered and are linked to important physiological roles (Ahima, 2006) including the innate immune response (Schäffler & Schölmerich, 2010). In a number of recent experiments transcriptional profiling demonstrated that 5,000 to 8,000 adipose tissue genes were differentially expressed during central stimulation of the melanocortin 4 receptor (Barb et al., 2010a) and several conditions such as fasting (Lkhagvadorj et al., 2009) and feed restriction (Lkhagvadorj et al., 2010). In contrast, 300 to 1,800 genes were differentially expressed in livers in these three studies (Barb et al., 2010a; Lkhagvadorj et al., 2009, 2010). This degree of differential gene expression in adipose depots reflects the potential influence of adipose tissue as a secretory organ on multiple systems in the body. Furthermore, advances in the study of adipose tissue gene expression include high throughput technologies in transcriptome profiling and deep sequencing of the adipose tissue microRNA transcriptome (review, Basu et al., 2012).

Recent proteomic studies of human and rat adipocytes have revealed the true scope of the adipose tissue secretome (Chen et al., 2005; Kheterpal et al., 2011; Lehr et al., 2012; Lim et al., 2008; Zhong et al., 2010). With refined and advanced proteomics techniques, these studies have revealed that many of the adipose tissue secreted factors identified at the gene level do indeed encode secreted proteins (Chen et al., 2005; Kheterpal et al., 2011; Lehr et al., 2012;

¹ Mention of trade names or commercial products in this publication is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the U.S. Department of Agriculture.

Lim et al., 2008; Zhong et al., 2010). The presence of an N-terminal secretion signal peptide validates secreted proteins in conditioned media (Renes et al., 2009). In many of these studies, the presence or absence of a signal peptide was used to validate or identify truly secreted adipocyte proteins (Chen et al., 2005; Lehr et al., 2012; Lim et al., 2008; Zhong et al., 2010). In these studies, the percentage of total apparent secreted proteins that were considered secreted (+ signal peptide) ranged from 39 to 75% and the total number of secreted proteins ranged from 164 to 263 (Chen et al., 2005; Lehr et al., 2012; Lim et al., 2008; Zhong et al., 2010). However, the signal peptide approach could underestimate the adipocyte derived proteins present in the extracellular space (review, Renes et al., 2009). For instance, a blocking strategy has been used to distinguish between true secreted proteins and proteins that simply “leak” from the cell (review, Renes et al., 2009). Continued development and refinement of proteomic approaches in the study of the adipose tissue secretome will ultimately confirm the endocrine status of adipose tissue.

1.2. Adipose tissue as a modulator of gonadotropin secretion

Adipose tissue plays a role in whole-body homeostasis by acting as an endocrine organ, which was clearly demonstrated with the discovery of leptin. Evidence indicates a strong link between neural influences and adipocyte expression and secretion of leptin and other adipokines such as other cytokines (interleukins), neurotrophic factors (ciliary neurotrophic factor, CNTF; brain-derived neurotrophic factor, BDNF), insulin-like growth factor (IGF-I, and -II), binding protein (IGFBP-5), and neuropeptides such as neuropeptide Y (NPY) and nesfatin-1 (Table 1). Developmental changes in these relationships are considered important for onset of puberty. Leptin augments secretion of gonadotropins which are essential for initiation and maintenance of normal reproductive function, by acting centrally at the hypothalamus to regulate the gonadotropin-releasing hormone (GnRH) and neuronal activity. The effects of leptin on GnRH are mediated through interneuronal pathways involving NPY, proopiomelanocortin (POMC) and kisspeptin. Increased infertility associated with diet induced obesity or central leptin resistance are likely mediated through the kisspeptin-GnRH pathway. Furthermore, leptin regulates reproductive function by altering the sensitivity of the pituitary gland to GnRH. Other putative metabolic signals are circulating long chain fatty acid which can signal nutrient availability to the central nervous system (CNS) and alter feed intake and glucose availability.

2. Free Fatty Acids (FFA)

2.1. Long-chain fatty acids act in the CNS

The control of appetite and metabolism in response to changes in nutrient availability occurs in part at the level of the hypothalamus (Barb et al., 1999, 2001a; Woods et al., 1998). Thus, macronutrients, such as carbohydrates and lipids, play a role in regulating peripheral concentrations of leptin and insulin (Ahima et al., 1996), which in turn has a direct effect on appetite and energy expenditure primarily through the hypothalamus (Barb et al., 2006; Woods et al., 1998). Levin et al. (1999) reported that hypothalamic neurons may directly detect nutrients. To that extent, treatment with a fatty acid synthase inhibitor reduced food

Regulatory – secreted factors			Receptors
adiponectin	IFNG	IL-5	ADIPOR2
adipsin	IGFBP-1	IL-6	BMPR2
agouti	IGFBP-3	IL-8	EDNRB
ANG	IGFBP-4	leptin	ESR1
ANGPTL2, ANGPTL4	IGFBP-5	PAI-1	FGFR1, FGFR4
RBP1, RBP4	IGFBP-7	RANTES	GNRHR2
APO-A1	IL-10	RTN	IFNGR1
APO-CIII	IL-12	TGF- β , TGF- β 3	IGF-IR, IGF-IIR
APO-E	IL-15	THBS1	IL-4R, IL-10R
APO-R1	IL-18	TNF α	PGRMC1
BDNF	IL-1A	VEGFC	INSR
bFGF	IL-1B	visfatin	NGFR
CNTF	IL-1RN	CTGF	OB-rb
IGF-I, IGF-II	IL-4	NPY	THRA, THRA2, TSHR
Chemokine ligands 2, 3, 4, 12	Compliment component 1, 2, 4A, 6X, C7	TGF- α	EGFR
BMP-4, BMP-15	CTRP4, CIQTN4	MCP-1	LDLR
RLN	PDGFD	NUCB2, nesfatin-1	LHCGR
LPL			TLR 4
			AGTR1

Abbreviations: ADIPOR2 = adiponectin receptor 2, AGTR1 = angiotensin II receptor, ANG = angiotensin, ANGPTL = angiopoietin-like protein, APO = apolipoprotein, BDNF = brain-derived neurotrophic factor, bFGF = basic fibroblast growth factor, BMP = bone morphogenic protein, BMPR2 = bone morphogenic protein receptor 2, CIQTN4 = complement-c1q tumor necrosis factor-related protein 4, CNTF = ciliary neurotrophic factor, CTGF = connective tissue growth factor, CTRP4 = complement-c1q tumor necrosis factor-related protein 4, EDNRB = endothelin receptor type B, EGFR = epidermal growth factor receptor, ESR1 = estrogen receptor 1, GNRHR2 = gonadotropin-releasing hormone receptor 2, IFNG = interferon gamma, IGF = insulin-like growth factor, IGF-IR = IGF-I receptor, IGFBP = insulin-like growth factor binding protein, IL = interleukin, INSR = insulin receptor, LDLR = low density lipoprotein receptor, LHCGR = luteinizing hormone-choriogonadotropin receptor, LPL = lipoprotein lipase, MCP-1 = monocyte chemoattractant protein-1, NGFR = nerve growth factor receptor, NPY = neuropeptide Y, OB-rb = long form leptin receptor, NUCB2 = nucleobindin 2, PAI-1 = plasminogen activator inhibitor-1, PDGFD = platelet derived growth factor D, PGRMC1 = progesterone receptor membrane component 1, RANTES = chemokine (c-c motif) ligand 5, RBP = retinol binding protein, RLN = relaxin, TGF = transforming growth factor, RTN = reticulon, THR = thyroid hormone receptor, TLR = toll-like receptor, TNF = tumor necrosis factor, TSHR = thyroid-stimulating hormone receptor, VEGFC = vascular endothelial growth factor C.

References: Barb et al., 2010a; Basu et al., 2012; Chen et al., 2005; Hausman & Hausman, 2004; Hausman et al., 2009; Lehr et al., 2012; Lim et al., 2008; Lkhagvadorj et al., 2009, 2010; Renes et al., 2009; Zhong et al., 2010

Table 1. List of representative genes and proteins reported to be expressed by adipose tissue of humans, large animals, and rats.

intake and body weight in mice by reducing expression of NPY in the hypothalamus via a malonyl-Coenzyme A mechanism, which supports the idea that lipid metabolism in the CNS plays a role in the control of appetite (Loftus et al., 2000). Furthermore, long-chain fatty acyl CoAs (LC-CoAs), such as oleyl-CoA, can activate ATP-sensitive K^+ channels in non-neuronal cells (Larsson et al., 1996). Circulating fatty acids gain rapid access to the brain, where they equilibrate with neuronal LC-CoAs (J.C. Miller et al., 1987; Rapaport, 1996). They are then further metabolized via mitochondria β -oxidation or incorporated into phospholipids (J.C. Miller et al., 1987; Rapaport, 1996). Obici et al. (2002) hypothesized that fatty acids may signal nutritional status to selective neurons in the CNS and activate a feedback loop designed to curtail further influx of nutrients into the circulation. To that extent, Obici and coworkers (2002) reported that intracerebroventricular (i.c.v.) administration of the long-chain fatty acid, oleic acid, suppressed glucose production and feed intake. In addition, this was accompanied by a reduction in hypothalamic expression of NPY. This neuronal circuit plays a role in maintaining energy homeostasis by switching fuel sources from carbohydrates to lipids and by limiting circulating endogenous and exogenous nutrients. Disruption of this circuit may play a role in obesity, type 2 diabetes and other endocrine abnormalities (for a review, see Obici, 2009), which are often accompanied by gonadotropin insufficiency.

2.2. Regulation of gonadotropin secretion by long-chain fatty acids

In the pig, feed deprivation results in a rapid mobilization of FFA from peripheral fat depots, but maintenance of euglycemia suggests increased hydrolysis of triglycerides and FFA oxidation resulting in a glucose sparing effect (Barb et al., 1997). We previously reported that metabolic response to acute feed deprivation occurred more rapidly in prepubertal gilts compared to mature gilts, likely because prepubertal gilts have a higher metabolic rate, smaller energy reserves and thus a greater nutrient intake requirement for growth (Barb et al., 1997). In mature animals, chronic feed restriction resulted in cessation of estrous cycles and lower concentrations of plasma insulin, increased levels of FFA and reduced LH pulse frequency compared to controls (Armstrong & Britt, 1987). This brings into question, therefore, if alterations in serum concentrations of FFA influence hypothalamic-pituitary function. To address this matter, prepubertal gilts received intravenous (i.v.) injection of a lipid emulsion which consisted of the following fatty acids: linoleic (65.87%), oleic (17.7%), palmitic (8.8%), linolenic (4.2%) and stearic (3.43%) acid. The fatty acid content of the lipid emulsion was comparable to that present in the circulation of the pig (Cera et al., 1989). Lipid emulsion injection enhanced the LH response to GnRH (Barb et al., 1991), whereas infusion of lipid emulsion at 1 hour intervals increased serum LH pulse amplitude without effecting LH pulse frequency (Barb et al., 1991). Dispersed cells of the anterior pituitary gland of the pig were cultured to determine whether the effects of FFA *in vivo* occur at the pituitary without the benefit of input from the CNS. The long-chain fatty acids, oleic and linoleic acids increased basal LH release. In contrast oleic acid suppressed the GnRH-induced release of LH (Figure 1). The response for linoleic acid was equivocal (Barb et al., 1995). These events seem to be mediated at the plasma membrane because oleic and linoleic acids did not block the forskolin-induced release of LH (Barb et al., 1995). These results may explain the altered

neuroendocrine activity observed during periods of feed restriction and fast. To that extent, administration of oleic acid into the third ventricle suppressed food intake and hypothalamic expression of NPY in the rat (Obici et al., 2002).

Figure 1. Anterior pituitary cells from prepubertal gilts ($n = 11$) were cultured in the presence of media alone (C, control wells; basal secretion in absence of any treatment) or gonadotropin releasing hormone (GnRH) at 10^{-8} M. Oleic or linoleic acid were included at 10^{-6} M, 10^{-5} M or 10^{-4} M in wells containing GnRH. Pituitary cells were exposed to oleic or linoleic acid for 30 min before the addition of GnRH. Media was collected 4 h after GnRH treatment. ^aDifferent from C ($P < 0.03$). ^bDifferent from GnRH alone ($P < 0.03$). Data from Barb et al. (1995).

An acute 28 h fast increased serum FFA concentrations, and decreased leptin pulse frequency but not mean concentrations of leptin in serum nor LH secretion in the ovariectomized prepubertal gilt (Barb et al., 2001b), while treatment with a competitive inhibitor of glycolysis suppressed LH secretion without affecting serum concentrations of leptin (Barb et al., 2001b). In contrast, short term feed restriction for 8 days decreased leptin secretion and LH pulse frequency in the mature ovariectomized gilt (Whisnant & Harrel, 2002). The ability of the pig to maintain euglycemia during acute fast may account for the failure of acute food deprivation to effect LH secretion (Barb et al., 1997). Although, leptin may serve as a metabolic signal which communicates metabolic status to the brain, the neuroendocrine response to acute energy deprivation may depend on age or mass of adipose tissue.

3. Nesfatin-1

3.1. Nesfatin-1 as an adipokine

While searching for new satiety factors, Oh-I et al. (2006) discovered a troglitazone- (PPAR γ ligand) stimulated transcript expressed in SQ-5 (lung squamous carcinoma cell line) cells

that was homologous to the nucleobindin 2 (NUCB2) gene, which codes for a DNA binding/EF hand/acidic protein (NEFA). The NUCB2 gene product is a 396 amino acid protein with several cleavage sites for prohormone convertase. Post-translational processing of the NUCB2 preprotein produces three cleavage products corresponding to amino acid residues 1-82, 85-163, and 166-396. Upon the observation that i.c.v. injection of the first 82 amino acid cleavage product suppressed feed intake resulting in reduced body and fat depot weights in mice, Oh-I et al. (2006) termed the protein nesfatin-1 for NEFA/nucleobindin2-encoded satiety- and fat-influencing protein-1.

Immediately upstream of the nesfatin-1 protein is a 26 amino acid signal sequence indicating that nesfatin-1 is likely a secreted factor that may have endocrine or paracrine action. Expression of NUCB2 mRNA is observed in predifferentiated 3T3-L1 cells (Oh-I et al., 2006; Ramanjaneya et al., 2010) and induction of differentiation resulted in a marked increase in expression of NUCB2 mRNA and secretion of nesfatin-1 into culture media (Ramanjaneya et al., 2010). Nesfatin-1 also is expressed and secreted from human and mouse adipose tissue explants (Ramanjaneya et al., 2010), with subcutaneous adipose tissue having greater expression of NUCB2/nesfatin-1 than omental adipose tissue (Ramanjaneya et al., 2010). Moreover, NUCB2 expression was greater in the adipocyte fraction of adipose tissue than in the stromal vascular fraction (Ramanjaneya et al., 2010) adding further support to the concept of nesfatin-1 as an adipose derived factor. Further studies are needed to define the precise roles of nesfatin-1, or the other NUCB2 gene products, in adipose tissue, but current evidence suggests involvement in chronic inflammatory response of adipose tissue associated with metabolic disease. Treating adipose tissue explants with energy partitioning hormones (insulin, dexamethasone) and cytokines, interleukin-6 (IL-6) and tumor necrosis factor α (TNF α), altered NUCB2 expression and nesfatin-1 secretion (Ramanjaneya et al., 2010). Furthermore, NUCB2 is involved in IL-1 β stimulated release of soluble tumor necrosis factor receptor 1 to the extracellular space (Islam et al., 2006).

It is important to note that NUCB2 mRNA and nesfatin-1 protein have been found to be expressed in several endocrine cells and glands throughout the body including gastric glands of digestive tract (Stengel et al., 2009a; Zhang et al., 2010), islet cells of the pancreas (Gonzalez et al., 2009), and Leydig cells of the testes (Garcia-Galiano et al., 2012). This is indicative of the role nesfatin-1 plays in gastric emptying and nutrient absorption (Stengel et al., 2009b), glucose utilization (Gonzalez et al., 2011; Nakata et al., 2011; Su et al., 2010), and testosterone production (Garcia-Galiano et al., 2012). At present, it is unclear how these tissues may contribute to circulating concentrations of nesfatin-1; however, given that adipose tissue is the largest endocrine organ of the body, the contribution that fat depots would have to plasma concentrations of nesfatin-1 seems obvious. Concentrations of nesfatin-1 in the blood are, for the most part, positively correlated with body mass index (BMI) in healthy human subjects (Aydin et al., 2009; Li et al., 2010; Ogiso et al., 2011; Ramanjaneya et al., 2010) as are several single nucleotide polymorphisms within the NUCB2 gene (Zegers et al., 2011). Expression of nesfatin-1 in subcutaneous adipose tissue of mice is suppressed with fasting and increased when mice were fed a high fat diet (Ramanjaneya et al., 2010) indicating that nesfatin-1 concentrations in serum could be regulated by nutritional

status. In point of fact, circulating concentrations of nesfatin-1 were less in patients with anorexia nervosa (Ogiso et al., 2011) and type 2 diabetes (Li et al., 2010). Together with the fact that nesfatin-1 crosses the blood-brain barrier via a nonsaturatable mechanism (Pan et al., 2007; Price et al., 2007), these data collectively indicate that nesfatin-1 is secreted from adipose tissue into the circulation and can enter the brain to regulate appetite.

3.2. Nesfatin-1 as a central regulator of food intake

The anorexogenic effects of nesfatin-1 are observed when nesfatin-1 is given either centrally (Shimizu et al., 2009) or peripherally (Stengel et al., 2009b). It is not clear, however, if suppression of appetite is entirely due to peripherally derived nesfatin-1 or the paracrine action of the protein produced within the hypothalamus. Expression of NUCB2/nesfatin-1 mRNA and protein has been demonstrated in several areas of the CNS. Within the hypothalamus, NUCB2/nesfatin-1 is expressed in nuclei that have important roles for control of appetite including the arcuate (ARC), paraventricular (PVN), lateral hypothalamic area and supraoptic nucleus (Brailoiu et al., 2007; Foo et al., 2008; Kohno et al., 2008; Oh-I et al., 2006). Areas of the brain stem that play pivotal roles in regulating energy homeostasis including the area postrema and the nucleus tractus solitaries (NTS) as well as the nucleus dorsalis of the vagus nerve all express NUCB2/nesfatin-1. Functional evidence that hypothalamic NUCB2/nesfatin-1 is involved in control of energy balance is derived from the observations that NUCB2/nesfatin-1 expression in the PVN is suppressed after fasting in adult and juvenile rats (Garcia-Galiano et al., 2010; Oh-I et al., 2006), and that refeeding activates nesfatin-1 neurons (as assessed by c-Fos) in the PVN (Kohno et al., 2008). Anorexogenic effects of nesfatin-1 require melanocortin receptors (Oh-I et al., 2006) and NPY neurons in hypothalamic slices of the ARC from mice were inhibited by nesfatin-1 *in vitro* (Price et al., 2008); although expression of NPY mRNA in the ARC of the rat *in vivo* was unchanged with nesfatin-1 treatment (Oh-I et al., 2006). Furthermore, alpha melanocyte-stimulating hormone treatment increased NUCB2 expression in the PVN (Oh-I et al., 2006) and nesfatin-1 has potent anorectic action in animals that are resistant to the effects of leptin (Oh-I et al., 2006; Su et al., 2010). This led to the initial thought that nesfatin-1 might be a down-stream effector of the action of leptin; however, i.c.v. injection of nesfatin-1 antibodies did not block the anorectic effect of leptin in the rat (Oh-I et al., 2006). Instead, the anorexogenic actions of nesfatin-1 appear to be relayed through a mechanism independent from leptin. For instance, nesfatin-1 stimulates oxytocin cells in the PVN which in turn activate POMC neurons in the NTS of the brain stem (Maejima et al., 2009). Moreover, cholecystokinin (CCK) activates NUCB2/nesfatin-1 cell bodies in the PVN and NTS. The inhibition of food intake by CCK is mediated, at least partially, through NUCB2/nesfatin-1 neurons via a corticotrophin-releasing hormone (CRH) 2-receptor. Blocking the action of the CRH2 receptor with an antagonist ameliorated the suppressive effects of nesfatin-1 on food intake (Stengel et al., 2009b).

3.3. Nesfatin-1 as a neuroendocrine regulator of gonadotropin secretion

The neuroanatomical distribution of nesfatin-1 cell bodies in areas of the hypothalamus involved in integration of energy balance and reproduction (i.e., the ARC) and the fact that

peripheral concentrations of nesfatin-1 reflect BMI suggest a role for nesfatin-1 in metabolic regulation of gonadotropin secretion. Hypothalamic expression of NUCB2/nesfatin-1 increases during the pubertal transition in the activity of the gonadotropic axis of rats (Garcia-Galiano et al., 2010). When young pubertal female rats were given i.c.v. injection of nesfatin-1, LH secretion increased two- to threefold; however, the effects of centrally administered nesfatin-1 on LH were much greater (9-fold increase) when rats were fasted for 48 h (Garcia-Galiano et al., 2010). The later observation is likely related to the fact that fasting or less severe but long-term nutrient restriction reduced NUCB2/nesfatin-1 expression in the brain and may explain a possible mechanism whereby fluctuations in energy balance impact gonadotropin secretion in a leptin independent manner. The stimulatory effects of i.c.v. nesfatin-1 on LH were not evident in adult female rats (Garcia-Galiano et al., 2010) suggesting nesfatin-1 plays an important role in regulating gonadotropin secretion during the pubertal transition; a period when increasing adiposity and sensitivity to adipokines is generally thought to be important for activation of the reproductive axis. Consistent with this is the fact that central infusion of nesfatin-1 antisense-morpholino oligonucleotides suppressed LH secretion and delayed puberty (as determined by absence of vaginal opening) in approximately 60% of peripubertal female rats but failed to alter ovulatory surges of LH in adult females (Garcia-Galiano et al., 2010). The effects of nesfatin-1 on LH and follicle-stimulating hormone (FSH) secretion may be sexually dimorphic as i.c.v. treatment with nesfatin-1 stimulated LH and FSH secretion in male rats that were fasted (Tadross et al., 2010). Moreover, nesfatin-1 stimulated release of GnRH from hypothalamic explants taken from male rats (Tadross et al., 2010).

Collectively these data indicated that nesfatin-1 is a protein hormone that participates in metabolic regulation of appetite and energy homeostasis. Reproductive function is sensitive to nutritional status and nesfatin-1 appears to have a role in conferring metabolic state to the gonadotropic axis, particularly during pubertal development. The mechanisms whereby this occurs have not been revealed yet, but likely involve action at the GnRH neuron. Whether this is a direct paracrine action of hypothalamic nesfatin-1 or an alteration in plasma concentrations of nesfatin-1 entering the brain is not known at present. Expression of nesfatin-1 in the testis and its role in regulating testosterone release (Garcia-Galiano et al., 2012) adds further complexity, and raises the possibility that nesfatin-1 can have indirect action on gonadotropin secretion through changes in gonadal steroid feed-back to the hypothalamus or anterior pituitary gland.

4. Leptin

4.1. Effects of leptin on the hypothalamic-pituitary axis

In the pig, presence of biologically-active leptin receptor (OB-rb) in the hypothalamus and pituitary (Lin et al., 2000) and the fact that leptin increased LH secretion from pig pituitary cells (Barb et al., 2004) and GnRH release from hypothalamic tissue (Figure 2; Barb et al., 2004) *in vitro* suggests that leptin acts through the hypothalamic-pituitary axis to modulate

Figure 2. Hypothalamic explants (hypothalamic-preoptic area) were collected from ovariectomized prepubertal gilts and were placed in perfusion culture. Tissue was treated as shown with recombinant human leptin (Lep) at 10⁻¹² M (n = 4), 10⁻¹⁰ M (n = 4), 10⁻⁸ M (n = 4), 10⁻⁶ M (n = 5) or control (n = 5). All fragments were exposed to K⁺ (60 mM) to verify tissue viability. Effluent was continuously collected as 5-min fractions (500 μ l). *Increased above baseline ($P < 0.05$). Data from Barb et al. (2004).

LH secretion. There is strong evidence from co-localization of leptin receptor mRNA with NPY gene expression that hypothalamic NPY is a potential target for leptin in the pig (Czaja et al., 2002). Moreover, central administration of NPY suppressed LH secretion and stimulated feed intake by reversing the inhibitory action of leptin (Barb et al., 2006). These results support the idea that leptin may serve as a metabolic signal in the activation of the reproductive axis.

Leptin treatment stimulated basal LH secretion directly from pig anterior pituitary cells in culture and GnRH release from hypothalamic-preoptic tissue explants from intact and ovariectomized prepubertal gilts on maintenance rations (Barb et al., 2004). Interestingly, i.c.v. administration of leptin failed to stimulate LH secretion in the well-fed intact prepubertal gilt (Barb et al., 2004). Obviously, hypothalamic explants are deprived of neuro-anatomical connections with other extra-hypothalamic tissues that may convey the heightened negative feedback action of estradiol on the GnRH pulse generator that occurs during pubertal development (Barb et al., 2010a), which may in part explain the failure of a LH response to i.c.v. administration of leptin in the pig.

Intracerebroventricular injection of leptin stimulated LH secretion in steroid-implanted castrated male sheep (D.W. Miller et al., 2002), and chronic i.c.v. administration of leptin stimulated LH secretion in the feed-restricted ovariectomized cow (Amstalden et al., 2002) and ewe (Henry et al., 2001). In contrast, chronic i.c.v. administration of leptin failed to stimulate LH secretion in well nourished ovariectomized ewes with no steroid replacement (Henry et al., 1999), and in intact ewe lambs (Morrison et al., 2001). *In vitro* studies demonstrated that leptin treatment stimulated basal and GnRH-mediated LH secretion from pituitary explants from fasted, but not control-fed cows, while having no effect on GnRH release from hypothalamic explants from either group of cows (Amstalden et al., 2003). Thus, metabolic state appears to be a primary determinant of the hypothalamic-pituitary response to leptin in ruminants.

4.2. The role of leptin in onset of puberty

Onset of puberty may be linked to attainment of a critical body weight or a minimum percentage of body fat (Frisch, 1984). Alternatively, metabolic mass and food intake or its correlated metabolic rate may be the triggering mechanism (Frisch, 1984). Initiation of puberty also may be influenced by metabolic factors of peripheral origin. In this regard, it has been postulated that metabolic signals are important in the initiation of puberty (Barb et al., 1997; Cameron et al., 1985). The discovery of leptin has improved our understanding of the relationship between adipose tissue and energy homeostasis (Campfield et al., 1995). Leptin treatment advanced sexual maturation in restricted and *ad lib* fed animals (Ahima et al., 1997; Barash et al., 1996). In addition, chronic leptin treatment not only reduced food intake and body weight in *ob/ob* (leptin deficient) mice, but also restored fertility (Barash et al., 1996). Serum leptin concentrations increased during puberty in the mouse (Chehab et al., 1997), heifer (Garcia et al., 2002) and pig (Qian et al., 1999) and, in the human female, age at first menarche was inversely related to serum leptin concentrations (Matkovic et al., 1997).

There exists, however, controversy as to the precise role of leptin in the onset of puberty. Several reports demonstrated that blood leptin concentrations remain relatively unchanged during pubertal development in the female mouse and rat (Ahima et al., 1998; Bronson, 2001; Cheung et al., 2001), while leptin administration failed to advance puberty onset in well nourished female mice (Cheung et al., 2001). Although, serum leptin concentrations increased during puberty in the gilt, other factors in addition to leptin may regulate onset of

puberty. As indicated above, it is hypothesized that estradiol modulates the hypothalamic-pituitary axis response to leptin (Barb et al., 2004). Moreover, estradiol may regulate the pubertal related changes in Ob-rb gene expression (Figure 3). In the ovariectomized prepubertal gilt, estrogen-induced increase in leptin mRNA expression in adipose tissue occurred at the time of expected puberty but not in younger animals (Qian et al., 1999). This was associated with an increase in LH pulse frequency (Barb et al., 2010b) and an age dependent increase in hypothalamic OB-rb expression (Lin et al., 2001).

Figure 3. The frequency of luteinizing hormone (LH) pulses (A) and expression of leptin mRNA in subcutaneous (s.q.) adipose tissue (B) of ovariectomized (OVX) gilts. Gilts were OVX at 90, 150, or 210 d of age. Osmotic pumps were implanted s.q. and delivered control (vehicle; polypropylene glycol) or 0.19 mg of estradiol benzoate per kg of body weight daily for 7 d. Messenger RNA for leptin was quantified with RNA protection assays. Means without a common superscript are different; for (A) $a,bP < 0.04$ and for (B) $a,bP < 0.01$. Data from Qian et al. (1999) and Barb et al. (2010b). Redrawn from Barb et al. (1999).

Several human studies, both cross-sectional and longitudinal, have demonstrated a sharp rise in serum leptin concentrations in young girls starting as early as age 7 and continuing to rise as they progressed through puberty at least age 15 (Ahmed et al., 1999; Blum et al., 1997; Garcia-Mayor et al., 1997). In contrast, in boys, leptin concentrations seem to increase transiently and then decline after Tanner stage 2 to prepubertal concentrations that are approximately one third of those observed in the late-pubertal girl. These changes in concentrations of leptin were paralleled by increasing body fat during female puberty and decreasing body fat during male puberty. Garcia-Mayor et al. (1997) reported in one cross-

sectional study, that the rise in serum concentrations of leptin were well established 2 years prior to marked increases in serum LH and estradiol concentrations were detected. The authors (Garcia-Mayor et al., 1997) suggest this is consistent with the hypothesis that leptin concentrations reach a putative threshold which allows puberty to progress; as opposed to a critical factor that triggers puberty.

Matkovic et al. (1997) examined the idea that if the relationship between body fat and early menarche in humans is mediated by leptin, then leptin concentrations would be related to age at menarche. This study consisted of 343 healthy girls (Tanner stage 2 of puberty) between 8.3 and 13.1 years of age. Menstrual history, height and weight, body composition by dual-energy X-ray absorptiometry, and leptin were measured every 6 to 12 months during a 4-year period. Leptin concentration was highly correlated with body fat mass ($r = 0.81$). Greater leptin concentrations up to 12 ng/mL were associated with a decline in the age of menarche by approximately 1 month per 1 ng/mL increase in leptin. Furthermore, a group of girls who remained premenarcheal for the entire 4 years of the study had significantly lower leptin concentrations compared to the groups of girls who reached menarche during the study. Matkovic and coworkers (1997) concluded that a threshold blood concentration of leptin may be needed for establishment of normal menses. Furthermore, in a recent review, Kaplowitz (2008) reports that current data supports the idea that leptin plays a permissive role as opposed to a metabolic signal that initiates puberty.

In the prepubertal ruminant, short term feed restriction reduced adipose leptin gene expression and leptin secretion, but increased hypothalamic OB-rb expression (Amstalden et al., 2000; Dyer et al., 1997). This was associated with decreased serum insulin concentration, IGF-I concentration and LH pulse frequency (Amstalden et al., 2000; Morrison et al., 2001). In addition, serum leptin concentrations increased as did leptin gene expression in heifers during pubertal development, which coincided with increases in serum IGF-I concentrations and body weight (Garcia et al., 2002). In contrast to the prepubertal heifer (Amstalden et al., 2000), short-term fasting failed to reduce pulsatile LH secretion in the mature cow (Amstalden et al., 2002). This suggests that there is a heightened sensitivity of the hypothalamic-pituitary axis to variations in energy availability in the heifer. Previous reports demonstrated that inhibition of LH secretion by nutrient restriction in the ovariectomized ewe (Henry et al., 2001) or the ewe lamb (Morrison et al., 2001) was reversed by leptin treatment demonstrating a positive association between LH secretion and leptin. Although leptin treatment reversed the fasting mediated reduction in LH pulse frequency in prepubertal heifers as cited above, chronic administration of ovine leptin by subcutaneous injections twice daily to 12- to 13-month old heifers for 40 days (Maciel et al., 2004) or 3 i.v. leptin injections per hour for 5 hours at 5-week intervals during pubertal development (Zieba et al., 2004) were unable to accelerate LH pulse frequency or onset of puberty. In contrast to data obtained from the cow, it is proposed that the effect of leptin on LH secretion in the pig during pubertal development is associated with stage of sexual maturation and subsequent change in the negative feedback action of estradiol on LH secretion (see Figure 3 and Barb et al., 2004, 2010a).

5. Kisspeptin

5.1. Kisspeptin regulates gonadotropin secretion and pubertal development

Kisspeptin is a hypothalamic neuropeptide and a potent stimulator of gonadotropin secretion (Caraty et al., 2007; Lents et al., 2008; Navarro et al., 2004a, 2005) due to its action directly on GnRH neurons (Constantin et al., 2009; Herbison et al., 2010; Irwig et al., 2004) to stimulate release of GnRH into the hypophyseal portal vessels (Messenger et al., 2005; Smith et al., 2011). A substantial body of evidence has accumulated that demonstrates kisspeptin plays a pivotal role in the timing of the onset of puberty. Hypothalamic expression of kisspeptin-1 (KiSS-1) and the kisspeptin receptor (GPR54) are developmentally regulated with expression increasing near the expected time of puberty (Castellano et al., 2005; Navarro et al., 2004a; Shahab et al., 2005). Furthermore, expression of KiSS-1 in the ARC and the rostral preoptic area are differentially regulated by gonadal steroids (Estrada et al., 2006; Smith et al., 2005, 2007; Tomikawa et al., 2010). It has recently been shown that increased LH pulsatility during sexual maturation in the ewe is associated with a reduction in the suppressive effects of estradiol on KiSS-1 expression (Redmond et al., 2011). The fundamental importance of kisspeptin in the onset of puberty raises the question as to whether kisspeptin has a central role in the timing of pubertal events associated with metabolic state or body energy reserves.

5.2. Kisspeptin is sensitive to energy balance

Restricted feeding and fasting reduces hypothalamic expression of KiSS-1 in rodents (Castellano et al., 2005; Luque et al., 2007), sheep (Backholer et al., 2010a), and nonhuman primates (Wahab et al., 2011). Expression of KiSS-1 also is suppressed during negative energy balance associated with lactation (True et al., 2011; Yamada et al., 2007). These data demonstrate that kisspeptin neurons in the hypothalamus are an important component to how the reproductive axis senses nutritional state. Castellano et al. (2005) used long-term caloric restriction to inhibit the occurrence of puberty (as defined by absence of vaginal opening and suppressed gonadotropin and estradiol concentrations) in female rats. Treating these rats with kisspeptin rescued gonadotropin secretion and induced puberty (vaginal opening) in approximately 60% of the animals, indicating that kisspeptin may have a role in integrating the effects of energy balance with the pubertal transition in gonadotropin secretion.

Recent data from growth restricted castrate male lambs indicates that the nutritional control of gonadotropin release may also involve alterations in sensitivity to kisspeptin. At 4 weeks after weaning, castrate male lambs were randomly assigned to different diets so that they either continued to grow or maintained body weight. After 12 weeks of treatment, animals in each group were then assigned to receive either i.v. infusion of 0.77 μ moles of kisspeptin or saline control. Area under the LH curve (AUC) for the saline treated animals was similar for growth and restricted lambs; however, the kisspeptin-induced release of LH was greater and lasted longer, as indicated by AUC, in the growth restricted lambs than in the full

growth lambs (Figure 4). Our findings in the growth restricted male lamb corroborate those of Castellano et al. (2005) in rats. These authors used prepubertal male and female rats that were fed either *ad libitum* or were fasted for 72 h. In fed animals, both prepubertal female and male rats demonstrated a 9 to 10 fold increase in LH concentrations in serum 15 minutes after i.c.v. injection of kisspeptin. In contrast, fasted rats demonstrated a much greater 50 to 60 fold increase in LH release. Moreover, the kisspeptin-stimulated release of GnRH from

Figure 4. Four weeks after weaning, castrate male lambs ($n = 16$) were divided and assigned to either continue normal growth (growth) or to maintain body weight (restricted). After 12 weeks, animals in both groups received $0.77 \mu\text{moles}$ of kisspeptin or saline as a single intravenous injection (time 0). A) Serum concentrations of luteinizing hormone (LH) during the 120 minutes before (period 1) and after (period 2) injection. B) Area under the curve (AUC) for each period (* $P < 0.05$). C) AUC in period 2 expressed as the change from period 1. ^{a,b,c}Means without a common superscript are different ($P < 0.05$). Data from C. A. Lents (unpublished).

hypothalamic explants collected from rats that were fasted for 72 h was greater than that from hypothalamic explants collected from *ad libitum* fed rats (Castellano et al., 2005). The increased responsiveness of the hypothalamus to kisspeptin and the subsequently greater release of LH in underfed animals are likely related to changes in expression of kisspeptin receptors. Expression of GPR54 mRNA in the hypothalamus was greater in fasted rats than in *ad libitum* fed controls (Castellano et al., 2005). Thus it appears that the pubertal transition in gonadotropin secretion involves not only increased expression and release of kisspeptin itself (Bentsen et al., 2010), but also a heightened sensitivity of the hypothalamus to the action of kisspeptin (Shahab et al., 2005). Both of these aspects can be modulated by metabolic state and are important for the overall tone of the kisspeptin system (Castellano et al., 2011; Roa et al., 2010).

5.3. Kisspeptin mediates the action of leptin on sexual development and gonadotropin secretion

The effect of energy balance on the kisspeptin system appears to be a consequence of the action of leptin. Expression of KiSS-1 in the ARC of the hypothalamus of *ob/ob* mice, which lack functional leptin, is significantly less when compared to expression in wild-type mice (Quennell et al., 2011; Smith et al., 2006); however, KiSS-1 expression in the anteroventral periventricular nucleus (AVPV) of *ob/ob* mice was similar to wild-type animals. This indicates that leptin acts on a specific population of kisspeptin cells within the ARC to modulate gonadotropin release. Leptin stimulated firing of kisspeptin neurons in hypothalamic slices of the ARC from guinea pigs (Qiu et al., 2011) and treating either *ob/ob* mice or *KiSS1-Cre* mice with leptin stimulated increased hypothalamic expression of kisspeptin mRNA in the ARC (Quennell et al., 2011; Smith et al., 2006) but not the AVPV (Cravo et al., 2011; Quennell et al., 2011; Smith et al., 2006). Leptin probably affects kisspeptin neurons in the ARC directly because kisspeptin cells localized within this hypothalamic area of guinea pigs (Qiu et al., 2011) and mice (Quennell et al., 2011; Smith et al., 2006) express leptin receptor mRNA. Moreover, second messengers that are important in signaling of leptin receptor (i.e., STAT-3) were expressed in kisspeptin cells in the ARC, but not within kisspeptin cells of the AVPV (Quennell et al., 2011). This indicates that increasing concentrations of leptin associated with greater body energy reserves may impact activity of the GnRH pulse generator through increasing the tone of the kisspeptin system via its action on kisspeptin neurons within the ARC.

The consequences of negative energy balance on KiSS-1 expression aren't fully offset by the positive effect of leptin. For example, leptin treatment did not fully reverse the lactation-induced reduction in KiSS-1 expression in rats (Xu et al., 2009). In a similar fashion, continuous i.c.v. infusion of leptin during a 72 h fast of ovariectomized ewes that were thin (made so with chronic nutritional restriction) rescued LH pulses (Backholer et al., 2010b) but KiSS-1 expression was only partially restored when compared with ewes that had greater body fat (Backholer et al., 2010a). Consequently, the suppressive effect of negative energy balance or nutrient deprivation on the gonadotropic axis via the KiSS-1 system

likely involves more than simply altered leptin signaling alone. Other metabolic factors, such as insulin for example, are reflective of metabolic state or availability of food and likely have an important role in regulating the kisspeptin system to augment gonadotropin release.

The possibility that adipocyte derived factors may also inhibit gonadotropin release in undernourished subjects should not be dismissed. Adiponectin is secreted by adipose tissue in response to nutrient restriction and body weight loss. It activates adenosine monophosphate-activated protein kinase (AMPK) to stimulate glucose uptake and β -oxidation of free fatty acids (Gil-Campos et al., 2004). Receptors for adiponectin are expressed not only in the hypothalamus (Kos et al., 2007) but also the anterior pituitary gland (Rodriguez-Pacheco et al., 2007). Furthermore, mice that overexpress adiponectin have an infertile phenotype (Combs et al., 2004). This is indicative of a role for adiponectin in modulating gonadotropin secretion during periods when nutrient intake is insufficient to meet energy demands. Treating anterior pituitary cells *in vitro* (Rodriguez-Pacheco et al., 2007) or L β T2 cells (immortalized embryonic gonadotrope cell line) with adiponectin suppressed both basal and GnRH-stimulated LH release (Lu et al., 2008). When adiponectin was administered i.c.v. to male rats, mean concentrations of LH were decreased owing to a suppression of LH pulse amplitude (Cheng et al., 2011). The later observation would indicate that adiponectin could be functioning to suppress activity of the GnRH neuronal network in subjects experiencing reductions in energy balance. In line with this is the fact that adiponectin inhibits the release of GnRH from GT1-7 cells (immortalized hypothalamic cell line) via an AMPK pathway (Cheng et al., 2011; Wen et al., 2008). It may well be that increased secretion and activity of adiponectin in animals during food deprivation or nutrient restriction off-set, to some degree, the stimulatory action of exogenous leptin on KiSS-1 expression in the hypothalamus. It is yet to be determined, however, if the suppressive effects of adiponectin on GnRH/LH release involve changes in the hypothalamic kisspeptin system.

Expression of KiSS-1 in immortalized hypothalamic N6 cells was increased after treatment with NPY (Luque et al., 2007). This would suggest that neuronal pathways downstream of leptin can impact the kisspeptin system. In the ewe, 13 to 30% of kisspeptin neurons in the ARC are in close apposition to NPY fibers (Backholer et al., 2010a). Moreover, 30 to 40% of kisspeptin cells in the ARC were contacted by POMC fibers (Backholer et al., 2010a). Since both NPY and POMC expressing cells are direct targets for leptin's action, the effects of leptin on gonadotropin secretion may be mediated through kisspeptin neurons indirectly via NPY and POMC pathways. It is also noted that kisspeptin neuronal fibers are located in close apposition to approximately 7% of NPY cell bodies and 20% of POMC cell bodies in the ovine hypothalamus (Backholer et al., 2010a). This anatomical evidence implies that the reproductive axis can influence neuronal pathways to modulate appetite. In fact, i.c.v. injection of kisspeptin increased NPY mRNA and reduced POMC mRNA in the ARC of the hypothalamus of sheep (Backholer et al., 2010a). Thus, other factors that may drive NPY or POMC expression during conditions of underfeeding may further limit the ability of leptin to stimulate increased KiSS-1 expression in the hypothalamus.

5.4. Kisspeptin is involved in the reproductive pathobiology of diabetes and obesity

Metabolic disorders such as diabetes and obesity are accompanied by alterations in adipose tissue biology and impaired fertility. Given the impacts of leptin on the kisspeptin system in the hypothalamus, one could easily speculate that metabolic diseases that impinge upon circulating concentrations of leptin could have negative consequences for reproductive function via alterations in the hypothalamic expression of kisspeptin. Using the streptozotocin-induced diabetic male rat model, Castellano et al. (2006) observed that LH release was rescued when rats were treated with exogenous kisspeptin. Moreover, expression of KiSS-1 was reduced in the hypothalamus of these diabetic male rats. When the authors treated the diabetic rats with leptin, they found that KiSS-1 expression was restored along with increased concentrations of LH and testosterone in serum.

Obesity is an ever growing epidemic and patients that are obese present a number of pathologies. One of these is a reduction in the sensitivity to the action of leptin. Iwasa et al. (2010) observed that female rats which underwent intrauterine growth retardation during their growth as fetuses developed leptin resistance after birth. These leptin resistant female rats demonstrated delayed onset of puberty associated with reduced expression of KiSS-1 in the hypothalamus. Thus, infertility associated with obesity and central leptin resistance may be related to tone of the kisspeptin system within the hypothalamus. Navarro et al. (2004b) found kisspeptin treatment restored LH secretion in *fa/fa* Zucker rats; a genetic model for leptin resistance. Furthermore, diet induced leptin resistance in mice, resulting from prolonged feeding of a high fat diet, was associated with reduced KiSS-1 expression and LH concentrations in serum (Quennell et al., 2011). Therefore, infertility resulting from hypogonadotropism that arises in diabetic or obese patients is likely due to alterations in the expression and secretion of kisspeptin in the hypothalamus.

6. Conclusion

Adipose tissue expresses and secretes a wide array of regulatory factors that have diverse biological roles. These factors contribute to the regulation of energy homeostasis by acting on neural circuits within the hypothalamus. Gonadotropin-releasing hormone is secreted from hypothalamic neurons and acts on gonadotrope cells within the anterior pituitary gland to stimulate the synthesis and release of LH. Activity of this gonadotropic-axis is sensitive to metabolic state. Free fatty acids are released from adipose tissue to have a glucose sparing effect and can be directly sensed by neurons in the hypothalamus. Cyclic changes in availability of FFA associated with meal frequency act to sustain continued release of LH pulses over short periods of time, but chronically elevated FFA likely impairs reproductive function by decreasing the sensitivity of the pituitary gland to GnRH. Conversely, leptin can enhance pituitary GnRH sensitivity and increase LH secretion. Within the hypothalamus, leptin stimulates release of GnRH by acting through interneuronal pathways involving NPY, POMC, and kisspeptin. Other adipose derived factors such as adiponectin and nesfatin-1 can have negative or positive effects on LH

release, respectively (figure 5). Metabolic control of puberty onset likely involves developmental changes in these relationships.

Figure 5. A proposed model for how metabolic signals, including adipokines such as leptin, affect gonadotropin secretion. Insulin fluctuates with consumption of meals at regular intervals to promote adipose accretion. Increased mass of adipose tissue is reflected in concentrations of adipokines such as leptin and nesfatin-1 that circulate in the blood to act as specific neural circuits within the hypothalamus. Leptin suppresses feed intake by modifying activity of POMC and NPY neurons in the arcuate (ARC) and paraventricular (PVN) nuclei, and stimulates release of gonadotropin hormones (LH and FSH). Many neurons in these areas of the hypothalamus express leptin receptor and directly innervate adipose tissue, thus constituting a hypothalamic-adipose neuroendocrine axis involving the sympathetic nervous system (SNS). Leptin directly activates kisspeptin (KiSS) cell bodies to stimulate GnRH release and to cause an upregulation of LH pulses. Nesfatin-1, which also stimulates LH release, suppresses food intake by acting through second order neurons to modulate activity of POMC systems in the nucleus tractus solitaries (NTS) of the hind brain as well as the nucleus dorsalis of the vagus nerve; thus altering liver function, which results in shifting availability of oxidizable fuels. Elevated free fatty acids (FFA) have a glucose sparing affect and can be directly sensed by neurons in the hypothalamus to sustain continued release of LH pulses over a short period.

Author details

Clay A. Lents*

United States Department of Agriculture² (USDA), Agricultural Research Service (ARS),
U.S. Meat Animal Research Center (USMARC), Clay Center, Nebraska, USA

* Corresponding Author

Richard C. Barb and Gary J. Hausman

*United States Department of Agriculture² (USDA), Agricultural Research Service (ARS),
Richard B. Russell Research Center (RRC), Athens, Georgia, USA*

Acknowledgement

The authors thank Linda Parnell for assistance in manuscript preparation. This project was supported by Agriculture and Food Research Initiative Competitive Grant no. 2005-35203-16852 (C.A. Lents) from the USDA National Institute of Food and Agriculture.

Abbreviations

ADIPOR2	adiponectin receptor 2	KISS-1	kisspeptin-1
AGTR1	angiotensin II receptor	LC-CoAs	long-chain fatty acyl CoAs
AMPK	adenosine monophosphate-activated protein kinase	LDLR	low density lipoprotein receptor
ANG	angiotensin	LH	luteinizing hormone
ANGPTL	angiopoietin-like protein	LHA	lateral hypothalamic area
APO	apolipoprotein	LHCGR	luteinizing hormone-choriogonadotropin receptor
ARC	arcuate	LPL	lipoprotein lipase
ARS	Agricultural Research Service	LβT2	immortalized embryonic gonadotrope cell line
AUC	Area under curve	MCP-1	monocyte chemoattractant protein-1
AVPV	anteroventral periventricular nucleus	N6	neuronal 6 cells
BDNF	brain-derived neurotrophic factor	NEFA	DNA binding/EF hand/acidic amino acid rich protein
bFGF	basic fibroblast growth factor	NGFR	nerve growth factor receptor
BMI	body mass index	NPY	neuropeptide Y
BMP	bone morphogenic protein	NTS	nucleus tractus solitaries
BMPR2	bone morphogenic protein receptor 2	NUCB2	nucleobindin 2
CCK	cholecystokinin	OB-rb	biologically-active long form leptin receptor
CIQTN4	complement-c1q tumor necrosis factor-related protein 4	PAI-1	plasminogen activator inhibitor-1
CNS	central nervous system	PDGFD	platelet derived growth factor D
CNTF	ciliary neurotrophic factor	PGRMC1	progesterone receptor membrane component 1

² The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410, or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

CRH	corticotrophin-releasing hormone	POMC	proopiomelanocortin
CTGF	connective tissue growth factor	PPAR γ	proxysome proliferator activated receptor γ
CTRP4	complement-c1q tumor necrosis factor-related protein 4	PVN	paraventricular
EDNRB	endothelin receptor type B	RANTES	chemokine (c-c motif) ligand 5
EGFR	epidermal growth factor receptor	RBP	retinol binding protein
ESR1	estrogen receptor 1	RLN	relaxin
FFA	free fatty acids	RRC	Richard B. Russell Research Center
FSH	follicle-stimulating hormone	RTN	reticulon
GnRH	gonadotropin-releasing hormone	SQ-5	lung squamous carcinoma cell line 5
GNRHR2	gonadotropin-releasing hormone receptor 2	STAT-3	signal transducer and activator of transcription 3
GPR54	g protein coupled receptor 54	TGF	transforming growth factor
GT1-7	GT1-7 cells (immortalized hypothalamic cell line)	THR	thyroid hormone receptor
i.c.v.	intracerebroventricular	TLR	toll-like receptor
i.v.	intravenous	TNF	tumor necrosis factor
IFNG	interferon gamma	TNF α	tumor necrosis factor α
IGF	insulin-like growth factor	TSHR	thyroid-stimulating hormone receptor
IGF-IR	IGF-I receptor	USDA	United States Department of Agriculture
IGFBP	insulin-like growth factor binding protein	USMARC	U.S. Meat Animal Research Center
IL	interleukin	VEGFC	vascular endothelial growth factor C
INSR	insulin receptor		

7. References

- Ahima, R. S. (2006). Adipose Tissue As an Endocrine Organ. *Obesity*, Vol.14, No.S8, (August 2006), pp. 242S-249S
- Ahima, R. S.; Prabakaran, D.; Mantzoros, C.; Qu, D.; Lowell, B.; Maratos-Flier, E. & Flier, J. S. (1996). Role of Leptin in the Neuroendocrine Response to Fasting. *Nature*, Vol.382, No.6588, (July 1996), pp. 250-252
- Ahima, R. S.; Dushay, J.; Flier, S. N.; Prabakaran, D. & Flier, J. S. (1997). Leptin Accelerates the Onset of Puberty in Normal Female Mice. *The Journal of Clinical Investigation*, Vol.99, No.3, (February 1997), pp. 391-395
- Ahima, R. S.; Prabakaran, D. & Flier, J. S. (1998). Postnatal Leptin Surge and Regulation of Circadian Rhythm of Leptin by Feeding. Implications for Energy Homeostasis and Neuroendocrine Function. *The Journal of Clinical Investigation*, Vol.101, No.5, (March 1998), pp. 1020-1027

- Ahmed, M. L.; Ong, K. K. L.; Morrell, D. J.; Cox, L.; Drayer, N.; Perry, L.; Preece, M. A. & Dunger, D. B. (1999). Longitudinal Study of Leptin Concentrations During Puberty: Sex Differences and Relationship to Changes in Body Composition. *The Journal of Clinical Endocrinology & Metabolism*, Vol.84, No.3, (March 1999), pp. 899-905
- Amstalden, M.; Garcia, M. R.; Williams, S. W.; Stanko, R. L.; Nizielski, S. E.; Morrison, C. D.; Keisler, D. H. & Williams, G. L. (2000). Leptin Gene Expression, Circulating Leptin, and Luteinizing Hormone Pulsatility Are Acutely Responsive to Short-Term Fasting in Prepubertal Heifers: Relationships to Circulating Insulin and Insulin-Like Growth Factor I. *Biology of Reproduction*, Vol.63, No.1, (July 2000), pp. 127-133
- Amstalden, M.; Garcia, M. R.; Stanko, R. L.; Nizielski, S. E.; Morrison, C. D.; Keisler, D. H. & Williams, G. L. (2002). Central Infusion of Recombinant Ovine Leptin Normalizes Plasma Insulin and Stimulates a Novel Hypersecretion of Luteinizing Hormone After Short-Term Fasting in Mature Beef Cows. *Biology of Reproduction*, Vol.66, No.5, (May 2002), pp. 1555-1561
- Amstalden, M.; Zieba, D. A.; Edwards, J. F.; Harms, P. G.; Welsh, T. H., Jr.; Stanko, R. L. & Williams, G. L. (2003). Leptin Acts at the Bovine Adenohypophysis to Enhance Basal and Gonadotropin-Releasing Hormone-Mediated Release of Luteinizing Hormone: Differential Effects Are Dependent Upon Nutritional History. *Biology of Reproduction*, Vol.69, No.5, (November 2003), pp. 1539-1544
- Armstrong, J. D. & Britt, J. H. (1987). Nutritionally-Induced Anestrus in Gilts: Metabolic and Endocrine Changes Associated With Cessation and Resumption of Estrous Cycles. *Journal of Animal Science*, Vol.65, No.2, (August 1987), pp. 508-523
- Aydin, S.; Dag, E.; Ozkan, Y.; Erman, F.; Dagli, A. F.; Kilic, N.; Sahin, İ.; Karatas, F.; Yoldas, T.; Barim, A. O. & Kendir, Y. (2009). Nesfatin-1 and Ghrelin Levels in Serum and Saliva of Epileptic Patients: Hormonal Changes Can Have a Major Effect on Seizure Disorders. *Molecular and Cellular Biochemistry*, Vol.328, No.1, (August 2009), pp. 49-56
- Backholer, K.; Smith, J. T.; Rao, A.; Pereira, A.; Iqbal, J.; Ogawa, S.; Li, Q. & Clarke, I. J. (2010a). Kisspeptin Cells in the Ewe Brain Respond to Leptin and Communicate With Neuropeptide Y and Proopiomelanocortin Cells. *Endocrinology*, Vol.151, No.5, (May 2010), pp. 2233-2243
- Backholer, K.; Bowden, M.; Gamber, K.; Bjørnbæk, C.; Iqbal, J. & Clarke, I. J. (2010b). Melanocortins Mimic the Effects of Leptin to Restore Reproductive Function in Lean Hypogonadotropic Ewes. *Neuroendocrinology*, Vol.91, No.1, (January 2010), pp. 27-40
- Barash, I. A.; Cheung, C. C.; Weigle, D. S.; Ren, H.; Kabisigting, E. B.; Kuijper, J. L.; Clifton, D. K. & Steiner, R. A. (1996). Leptin Is a Metabolic Signal to the Reproductive System. *Endocrinology*, Vol.137, No.7, (July 1996), pp. 3144-3147
- Barb, C. R.; Kraeling, R. R.; Barrett, J. B.; Rampacek, G. B.; Campbell, R. M. & Mowles, T. F. (1991). Serum Glucose and Free Fatty Acids Modulate Growth Hormone and Luteinizing Hormone Secretion in the Pig. *Proceedings of the Society for Experimental Biology and Medicine*, Vol.198, No.1, (October 1991), pp. 636-642

- Barb, C. R.; Kraeling, R. R. & Rampacek, G. B. (1995). Glucose and Free Fatty Acid Modulation of Growth Hormone and Luteinizing Hormone Secretion by Cultured Porcine Pituitary Cells. *Journal of Animal Science*, Vol.73, No.5, (May 1995), pp. 1416-1423
- Barb, C. R.; Kraeling, R. R.; Rampacek, G. B. & Dove, C. R. (1997). Metabolic Changes During the Transition From the Fed to the Acute Feed-Deprived State in Prepuberal and Mature Gilts. *Journal of Animal Science*, Vol.75, No.3, (March 1997), pp. 781-789
- Barb, C. R.; Barrett, J. B.; Kraeling, R. R. & Rampacek, G. B. (1999). Role of Leptin in Modulating Neuroendocrine Function: A Metabolic Link Between the Brain-Pituitary and Adipose Tissue. *Reproduction in Domestic Animals*, Vol.34, No.3-4, (August 1999), pp. 111-125
- Barb, C. R.; Kraeling, R. R. & Rampacek, G. B. (2001a). Nutritional Regulators of the Hypothalamic-Pituitary Axis in Pigs. *Reproduction Supplement*, Vol.58, (2001), pp. 1-15
- Barb, C. R.; Barrett, J. B.; Kraeling, R. R. & Rampacek, G. B. (2001b). Serum Leptin Concentrations, Luteinizing Hormone and Growth Hormone Secretion During Feed and Metabolic Fuel Restriction in the Prepuberal Gilt. *Domestic Animal Endocrinology*, Vol.20, No.1, (January 2001), pp. 47-63
- Barb, C. R.; Barrett, J. B. & Kraeling, R. R. (2004). Role of Leptin in Modulating the Hypothalamic-Pituitary Axis and Luteinizing Hormone Secretion in the Prepuberal Gilt. *Domestic Animal Endocrinology*, Vol.26, No.3, (April 2004), pp. 201-214
- Barb, C. R.; Kraeling, R. R.; Rampacek, G. B. & Hausman, G. J. (2006). The Role of Neuropeptide Y and Interaction With Leptin in Regulating Feed Intake and Luteinizing Hormone and Growth Hormone Secretion in the Pig. *Reproduction*, Vol.131, No.6, (June 2006), pp. 1127-1135
- Barb, C. R.; Hausman, G. J.; Rekaya, R.; Lents, C. A.; Lkhagvadorj, S.; Qu, L.; Cai, W.; Couture, O. P.; Anderson, L. L.; Dekkers, J. C. M. & Tuggle, C. K. (2010a). Gene Expression in Hypothalamus, Liver, and Adipose Tissues and Food Intake Response to Melanocortin-4 Receptor Agonist in Pigs Expressing Melanocortin-4 Receptor Mutations. *Physiological Genomics*, Vol.41, No.3, (May 2010), pp. 254-268
- Barb, C. R.; Hausman, G. J. & Kraeling, R. R. (2010b). Luteinizing Hormone Secretion As Influenced by Age and Estradiol in the Prepubertal Gilt. *Animal Reproduction Science*, Vol.122, No.3-4, (December 2010), pp. 324-327
- Basu, U.; Romao, J. M. & Guan, L. L. (2012). Adipogenic Transcriptome Profiling Using High Throughput Technologies. *Journal of Genomics*, Vol.1, (2012), pp. 22-28
- Bentsen, A. H.; Ansel, L.; Simonneaux, V.; Tena-Sempere, M.; Juul, A. & Mikkelsen, J. D. (2010). Maturation of Kisspeptinergic Neurons Coincides With Puberty Onset in Male Rats. *Peptides*, Vol.31, No.2, (February 2010), pp. 275-283
- Blum, W. F.; Englaro, P.; Hanitsch, S.; Juul, A.; Hertel, N. T.; Müller, J.; Skakkebaek, N. E.; Heiman, M. L.; Birkett, M.; Attanasio, A. M.; Kiess, W. & Rascher, W. (1997). Plasma Leptin Levels in Healthy Children and Adolescents: Dependence on Body Mass Index, Body Fat Mass, Gender, Pubertal Stage, and Testosterone. *The Journal of Clinical Endocrinology & Metabolism*, Vol.82, No.9, (September 1997), pp. 2904-2910

- Brailoiu, G. C.; Dun, S. L.; Brailoiu, E.; Inan, S.; Yang, J.; Chang, J. K. & Dun, N. J. (2007). Nesfatin-1: Distribution and Interaction With a G Protein-Coupled Receptor in the Rat Brain. *Endocrinology*, Vol.148, No.10, (October 2007), pp. 5088-5094
- Bronson, F. H. (2001). Puberty in Female Mice Is Not Associated With Increases in Either Body Fat or Leptin. *Endocrinology*, Vol.142, No.11, (November 2001), pp. 4758-4761
- Cameron, J. L.; Koerker, D. J. & Steiner, R. A. (1985). Metabolic Changes During Maturation of Male Monkeys: Possible Signals for Onset of Puberty. *American Journal of Physiology - Endocrinology and Metabolism*, Vol.249, No.4, (October 1985), p. E385-E391
- Campfield, L. A.; Smith, F. J.; Guisez, Y.; Devos, R. & Burn, P. (1995). Recombinant Mouse OB Protein: Evidence for a Peripheral Signal Linking Adiposity and Central Neural Networks. *Science*, Vol.269, No.5223, (July 1995), pp. 546-549
- Caraty, A.; Smith, J. T.; Lomet, D.; Ben Saïd, S.; Morrissey, A.; Cognie, J.; Doughton, B.; Baril, G.; Briant, C. & Clarke, I. J. (2007). Kisspeptin Synchronizes Preovulatory Surges in Cyclical Ewes and Causes Ovulation in Seasonally Acyclic Ewes. *Endocrinology*, Vol.148, No.11, (November 2007), pp. 5258-5267
- Castellano, J. M.; Navarro, V. M.; Fernández-Fernández, R.; Nogueiras, R.; Tovar, S.; Roa, J.; Vazquez, M. J.; Vigo, E.; Casanueva, F. F.; Aguilar, E.; Pinilla, L.; Dieguez, C. & Tena-Sempere, M. (2005). Changes in Hypothalamic KiSS-1 System and Restoration of Pubertal Activation of the Reproductive Axis by Kisspeptin in Undernutrition. *Endocrinology*, Vol.146, No.9, (September 2005), pp. 3917-3925
- Castellano, J. M.; Navarro, V. M.; Fernández-Fernández, R.; Roa, J.; Vigo, E.; Pineda, R.; Dieguez, C.; Aguilar, E.; Pinilla, L. & Tena-Sempere, M. (2006). Expression of Hypothalamic KiSS-1 System and Rescue of Defective Gonadotropic Responses by Kisspeptin in Streptozotocin-Induced Diabetic Male Rats. *Diabetes*, Vol.55, No.9, (September 2006), pp. 2602-2610
- Castellano, J. M.; Bentsen, A. H.; Sánchez-Garrido, M. A.; Ruiz-Pino, F.; Romero, M.; Garcia-Galiano, D.; Aguilar, E.; Pinilla, L.; Diéguez, C.; Mikkelsen, J. D. & Tena-Sempere, M. (2011). Early Metabolic Programming of Puberty Onset: Impact of Changes in Postnatal Feeding and Rearing Conditions on the Timing of Puberty and Development of the Hypothalamic Kisspeptin System. *Endocrinology*, Vol.152, No.9, (September 2011), pp. 3396-3408
- Cera, K. R.; Mahan, D. C. & Reinhart, G. A. (1989). Postweaning Swine Performance and Serum Profile Responses to Supplemental Medium-Chain Free Fatty Acids and Tallow. *Journal of Animal Science*, Vol.67, No.8, (August 1989), pp. 2048-2055
- Chehab, F. F.; Mounzih, K.; Lu, R. & Lim, M. E. (1997). Early Onset of Reproductive Function in Normal Female Mice Treated With Leptin. *Science*, Vol.275, No.5296, (January 1997), pp. 88-90
- Chen, X.; Cushman, S. W.; Pannell, L. K. & Hess, S. (2005). Quantitative Proteomic Analysis of the Secretory Proteins From Rat Adipose Cells Using a 2D Liquid Chromatography-MS/MS Approach. *Journal of Proteome Research*, Vol.4, No.2, (March 2005), pp. 570-577

- Cheng, X.-B.; Wen, J.-P.; Yang, J.; Yang, Y.; Ning, G. & Li, X.-Y. (2011). GnRH Secretion Is Inhibited by Adiponectin Through Activation of AMP-Activated Protein Kinase and Extracellular Signal-Regulated Kinase. *Endocrine*, Vol.39, No.1, (February 2011), pp. 6-12
- Cheung, C. C.; Thornton, J. E.; Nurani, S. D.; Clifton, D. K. & Steiner, R. A. (2001). A Reassessment of Leptin's Role in Triggering the Onset of Puberty in the Rat and Mouse. *Neuroendocrinology*, Vol.74, No.1, (July 2001), pp. 12-21
- Combs, T. P.; Pajvani, U. B.; Berg, A. H.; Lin, Y.; Jelicks, L. A.; Laplante, M.; Nawrocki, A. R.; Rajala, M. W.; Parlow, A. F.; Cheeseboro, L.; Ding, Y.-Y.; Russell, R. G.; Lindemann, D.; Hartley, A.; Baker, G. R. C.; Obici, S.; Deshaies, Y.; Ludgate, M.; Rossetti, L. & Scherer, P. E. (2004). A Transgenic Mouse With a Deletion in the Collagenous Domain of Adiponectin Displays Elevated Circulating Adiponectin and Improved Insulin Sensitivity. *Endocrinology*, Vol.145, No.1, (January 2004), pp. 367-383
- Constantin, S.; Caligioni, C. S.; Stojilkovic, S. & Wray, S. (2009). Kisspeptin-10 Facilitates a Plasma Membrane-Driven Calcium Oscillator in Gonadotropin-Releasing Hormone-1 Neurons. *Endocrinology*, Vol.150, No.3, (March 2009), pp. 1400-1412
- Cravo, R. M.; Margatho, L. O.; Osborne-Lawrence, S.; Donato Jr, J.; Atkin, S.; Bookout, A. L.; Rovinsky, S.; Frazão, R.; Lee, C. E.; Gautron, L.; Zigman, J. M. & Elias, C. F. (2011). Characterization of Kiss1 Neurons Using Transgenic Mouse Models. *Neuroscience*, Vol.173, (January 2011), pp. 37-56
- Czaja, K.; Lakomy, M.; Sienkiewicz, W.; Kaleczyc, J.; Pidsudko, Z.; Barb, C. R.; Rampacek, G. B. & Kraeling, R. R. (2002). Distribution of Neurons Containing Leptin Receptors in the Hypothalamus of the Pig. *Biochemical and Biophysical Research Communications*, Vol.298, No.3, (November 2002), pp. 333-337
- Dyer, C. J.; Simmons, J. M.; Matteri, R. L. & Keisler, D. H. (1997). Leptin Receptor mRNA Is Expressed in Ewe Anterior Pituitary and Adipose Tissues and Is Differentially Expressed in Hypothalamic Regions of Well-Fed and Feed-Restricted Ewes. *Domestic Animal Endocrinology*, Vol.14, No.2, (March 1997), pp. 119-128
- Estrada, K. M.; Clay, C. M.; Pompolo, S.; Smith, J. T. & Clarke, I. J. (2006). Elevated KiSS-1 Expression in the Arcuate Nucleus Prior to the Cyclic Preovulatory Gonadotrophin-Releasing Hormone/Lutenising Hormone Surge in the Ewe Suggests a Stimulatory Role for Kisspeptin in Oestrogen-Positive Feedback. *Journal of Neuroendocrinology*, Vol.18, No.10, (October 2006), pp. 806-809
- Foo, K. S.; Brismar, H. & Broberger, C. (2008). Distribution and Neuropeptide Coexistence of Nucleobindin-2 mRNA/Nesfatin-Like Immunoreactivity in the Rat CNS. *Neuroscience*, Vol.156, No.3, (October 2008), pp. 563-579
- Frisch, R. E. (1984). Body Fat, Puberty and Fertility. *Biol. Rev. Camb. Philos. Soc.*, Vol.59, No.2, (May 1984), pp. 161-188
- Garcia, M. R.; Amstalden, M.; Williams, S. W.; Stanko, R. L.; Morrison, C. D.; Keisler, D. H.; Nizielski, S. E. & Williams, G. L. (2002). Serum Leptin and Its Adipose Gene Expression During Pubertal Development, the Estrous Cycle, and Different Seasons in Cattle. *Journal of Animal Science*, Vol.80, No.8, (August 2002), pp. 2158-2167

- García-Galiano, D.; Navarro, V. M.; Roa, J.; Ruiz-Pino, F.; Sánchez-Garrido, M. A.; Pineda, R.; Castellano, J. M.; Romero, M.; Aguilar, E.; Gaytán, F.; Diéguez, C.; Pinilla, L. & Tena-Sempere, M. (2010). The Anorexigenic Neuropeptide, Nesfatin-1, Is Indispensable for Normal Puberty Onset in the Female Rat. *The Journal of Neuroscience*, Vol.30, No.23, (June 2010), pp. 7783-7792
- García-Galiano, D.; Pineda, R.; Ilhan, T.; Castellano, J. M.; Ruiz-Pino, F.; Sánchez-Garrido, M. A.; Vazquez, M. J.; Sangiao-Alvarellos, S.; Romero-Ruiz, A.; Pinilla, L.; Diéguez, C.; Gaytán, F. & Tena-Sempere, M. (2012). Cellular Distribution, Regulated Expression, and Functional Role of the Anorexigenic Peptide, NUCB2/Nesfatin-1, in the Testis. *Endocrinology*, Vol.153, Published online before print February 14, 2012, doi: 10.1210/en.2011-2032 (Available from <http://endo.endojournals.org/content/early/2012/02/08/en.2011-2032.abstract>)
- Garcia-Mayor, R. V.; Andrade, M. A.; Rios, M.; Lage, M.; Dieguez, C. & Casanueva, F. F. (1997). Serum Leptin Levels in Normal Children: Relationship to Age, Gender, Body Mass Index, Pituitary-Gonadal Hormones, and Pubertal Stage. *The Journal of Clinical Endocrinology & Metabolism*, Vol.82, No.9, (September 1997), pp. 2849-2855
- Gil-Campos, M.; Cañete, R. & Gil, A. (2004). Adiponectin, the Missing Link in Insulin Resistance and Obesity. *Clinical Nutrition*, Vol.23, No.5, (October 2004), pp. 963-974
- Gonzalez, R.; Tiwari, A. & Unniappan, S. (2009). Pancreatic Beta Cells Colocalize Insulin and Pronesfatin Immunoreactivity in Rodents. *Biochemical and Biophysical Research Communications*, Vol.381, No.4, (April 2009), pp. 643-648
- Gonzalez, R.; Perry, R. L. S.; Gao, X.; Gaidhu, M. P.; Tsushima, R. G.; Ceddia, R. B. & Unniappan, S. (2011). Nutrient Responsive Nesfatin-1 Regulates Energy Balance and Induces Glucose-Stimulated Insulin Secretion in Rats. *Endocrinology*, Vol.152, No.10, (October 2011), pp. 3628-3637
- Hausman, G. J. & Hausman, D. B. (2004). Ontogeny: Adipose Tissue, In: *Encyclopedia of Animal Science*, W.G. Pond & A.W. Bell, (Eds.), pp. 684-687, Marcel Dekker, Inc., ISBN 0-415-80286-5 [ei 1-4389-0932-1], New York, NY
- Hausman, G. J.; Dodson, M. V.; Ajuwon, K.; Azain, M.; Barnes, K. M.; Guan, L. L.; Jiang, Z.; Poulos, S. P.; Sainz, R. D.; Smith, S.; Spurlock, M.; Novakofski, J.; Fernyhough, M. E. & Bergen, W. G. (2009). Board-Invited Review: The Biology and Regulation of Preadipocytes and Adipocytes in Meat Animals. *Journal of Animal Science*, Vol.87, No.4, (April 2009), pp. 1218-1246
- Henry, B. A.; Goding, J. W.; Alexander, W. S.; Tilbrook, A. J.; Canny, B. J.; Dunshea, F.; Rao, A.; Mansell, A. & Clarke, I. J. (1999). Central Administration of Leptin to Ovariectomized Ewes Inhibits Food Intake Without Affecting the Secretion of Hormones From the Pituitary Gland: Evidence for a Dissociation of Effects on Appetite and Neuroendocrine Function. *Endocrinology*, Vol.140, No.3, (March 1999), pp. 1175-1182
- Henry, B. A.; Goding, J. W.; Tilbrook, A. J.; Dunshea, F. R. & Clarke, I. J. (2001). Intracerebroventricular Infusion of Leptin Elevates the Secretion of Luteinising Hormone Without Affecting Food Intake in Long-Term Food-Restricted Sheep, but

- Increases Growth Hormone Irrespective of Bodyweight. *The Journal of Endocrinology*, Vol.168, No.1, (January 2001), pp. 67-77
- Herbison, A. E.; d'Anglemont de Tassigny, X.; Doran, J. & Colledge, W. H. (2010). Distribution and Postnatal Development of *Gpr54* Gene Expression in Mouse Brain and Gonadotropin-Releasing Hormone Neurons. *Endocrinology*, Vol.151, No.1, (January 2010), pp. 312-321
- Irwig, M. S.; Fraley, G. S.; Smith, J. T.; Acohido, B. V.; Popa, S. M.; Cunningham, M. J.; Gottsch, M. L.; Clifton, D. K. & Steiner, R. A. (2004). Kisspeptin Activation of Gonadotropin Releasing Hormone Neurons and Regulation of KiSS-1 MRNA in the Male Rat. *Neuroendocrinology*, Vol.80, No.4, (2004), pp. 264-272
- Islam, A.; Adamik, B.; Hawari, F. I.; Ma, G.; Rouhani, F. N.; Zhang, J. & Levine, S. J. (2006). Extracellular TNFR1 Release Requires the Calcium-Dependent Formation of a Nucleobindin 2-ARTS-1 Complex. *Journal of Biological Chemistry*, Vol.281, No.10, (March 2006), pp. 6860-6873
- Iwasa, T.; Matsuzaki, T.; Murakami, M.; Fujisawa, S.; Kinouchi, R.; Gereltsetseg, G.; Kuwahara, A.; Yasui, T. & Irahara, M. (2010). Effects of Intrauterine Undernutrition on Hypothalamic *Kiss1* Expression and the Timing of Puberty in Female Rats. *The Journal of Physiology*, Vol.588, No.5, (March 2010), pp. 821-829
- Kaplowitz, P. B. (2008). Link Between Body Fat and the Timing of Puberty. *Pediatrics*, Vol.121, No.Supplement 3, (February 2008), p. S208-S217
- Kheterpal, I.; Ku, G.; Coleman, L.; Yu, G.; Ptitsyn, A. A.; Floyd, Z. E. & Gimble, J. M. (2011). Proteome of Human Subcutaneous Adipose Tissue Stromal Vascular Fraction Cells Versus Mature Adipocytes Based on DIGE. *Journal of Proteome Research*, Vol.10, No.4, (January 2011), pp. 1519-1527
- Kohno, D.; Nakata, M.; Maejima, Y.; Shimizu, H.; Sedbazar, U.; Yoshida, N.; Dezaki, K.; Onaka, T.; Mori, M. & Yada, T. (2008). Nesfatin-1 Neurons in Paraventricular and Supraoptic Nuclei of the Rat Hypothalamus Coexpress Oxytocin and Vasopressin and Are Activated by Refeeding. *Endocrinology*, Vol.149, No.3, (March 2008), pp. 1295-1301
- Kos, K.; Harte, A. L.; da Silva, N. F.; Tonchev, A.; Chaldakov, G.; James, S.; Snead, D. R.; Hoggart, B.; O'Hare, J. P.; McTernan, P. G. & Kumar, S. (2007). Adiponectin and Resistin in Human Cerebrospinal Fluid and Expression of Adiponectin Receptors in the Human Hypothalamus. *The Journal of Clinical Endocrinology & Metabolism*, Vol.92, No.3, (March 2007), pp. 1129-1136
- Larsson, O.; Deeney, J. T.; Bränström, R.; Berggren, P.-O. & Corkey, B. E. (1996). Activation of the ATP-Sensitive K⁺ Channel by Long Chain Acyl-CoA. *The Journal of Biological Chemistry*, Vol.271, No.18, (May 1996), pp. 10623-10626
- Lehr, S.; Hartwig, S.; Lamers, D.; Famulla, S.; Müller, S.; Hanisch, F.-G.; Cuvelier, C.; Ruige, J.; Eckardt, K.; Ouwers, D. M.; Sell, H. & Eckel, J. (2012). Identification and Validation of Novel Adipokines Released From Primary Human Adipocytes. *Molecular & Cellular Proteomics*, Vol.11, No.1, (January 2012), p. M111.011734

- Lents, C. A.; Heidorn, N. L.; Barb, C. R. & Ford, J. J. (2008). Central and Peripheral Administration of Kisspeptin Activates Gonadotropin but Not Somatotropin Secretion in Prepubertal Gilts. *Reproduction*, Vol.135, No.6, (March 2008), pp. 879-887
- Levin, B. E.; Dunn-Meynell, A. A. & Routh, V. H. (1999). Brain Glucose Sensing and Body Energy Homeostasis: Role in Obesity and Diabetes. *AJP - Regulatory, Integrative and Comparative Physiology*, Vol.276, No.5, (May 1999), p. R1223-R1231
- Li, Q.-C.; Wang, H.-Y.; Chen, X.; Guan, H.-Z. & Jiang, Z.-Y. (2010). Fasting Plasma Levels of Nesfatin-1 in Patients With Type 1 and Type 2 Diabetes Mellitus and the Nutrient-Related Fluctuation of Nesfatin-1 Level in Normal Humans. *Regulatory Peptides*, Vol.159, No.1-3, (January 2010), pp. 72-77
- Lim, J.-M.; Sherling, D.; Teo, C. F.; Hausman, D. B.; Lin, D. & Wells, L. (2008). Defining the Regulated Secreted Proteome of Rodent Adipocytes Upon the Induction of Insulin Resistance. *Journal of Proteome Research*, Vol.7, No.3, (March 2008), pp. 1251-1263
- Lin, J.; Barb, C. R.; Matteri, R. L.; Kraeling, R. R.; Chen, X.; Meinersmann, R. J. & Rampacek, G. B. (2000). Long Form Leptin Receptor mRNA Expression in the Brain, Pituitary, and Other Tissues in the Pig. *Domestic Animal Endocrinology*, Vol.19, No.1, (July 2000), pp. 53-61
- Lin, J.; Barb, C. R.; Kraeling, R. R. & Rampacek, G. B. (2001). Developmental Changes in the Long Form Leptin Receptor and Related Neuropeptide Gene Expression in the Pig Brain. *Biology of Reproduction*, Vol.64, No.6, (June 2001), pp. 1614-1618
- Lkhagvadorj, S.; Qu, L.; Cai, W.; Couture, O. P.; Barb, C. R.; Hausman, G. J.; Nettleton, D.; Anderson, L. L.; Dekkers, J. C. M. & Tuggle, C. K. (2009). Microarray Gene Expression Profiles of Fasting Induced Changes in Liver and Adipose Tissues of Pigs Expressing the Melanocortin-4 Receptor D298N Variant. *Physiological Genomics*, Vol.38, No.1, (June 2009), pp. 98-111
- Lkhagvadorj, S.; Qu, L.; Cai, W.; Couture, O. P.; Barb, C. R.; Hausman, G. J.; Nettleton, D.; Anderson, L. L.; Dekkers, J. C. M. & Tuggle, C. K. (2010). Gene Expression Profiling of the Short-Term Adaptive Response to Acute Caloric Restriction in Liver and Adipose Tissues of Pigs Differing in Feed Efficiency. *AJP - Regulatory, Integrative and Comparative Physiology*, Vol.298, No.2, (February 2010), p. R494-R507
- Loftus, T. M.; Jaworsky, D. E.; Frehywot, G. L.; Townsend, C. A.; Ronnett, G. V.; Lane, M. D. & Kuhajda, F. P. (2000). Reduced Food Intake and Body Weight in Mice Treated With Fatty Acid Synthase Inhibitors. *Science*, Vol.288, No.5475, (June 2000), pp. 2379-2381
- Lu, M.; Tang, Q.; Olefsky, J. M.; Mellon, P. L. & Webster, N. J. G. (2008). Adiponectin Activates Adenosine Monophosphate-Activated Protein Kinase and Decreases Luteinizing Hormone Secretion in L β T2 Gonadotropes. *Molecular Endocrinology*, Vol.22, No.3, (March 2008), pp. 760-771
- Luque, R. M.; Kineman, R. D. & Tena-Sempere, M. (2007). Regulation of Kiss-1 and GPR54 Genes by Metabolic Factors: Analyses Using Mouse Models and a Cell Line. *Endocrinology*, Vol.148, No.10, (October 2007), pp. 4601-4611
- Maciel, M. N.; Zieba, D. A.; Amstalden, M.; Keisler, D. H.; Neves, J. P. & Williams, G. L. (2004). Chronic Administration of Recombinant Ovine Leptin in Growing Beef Heifers:

- Effects on Secretion of LH, Metabolic Hormones, and Timing of Puberty. *Journal of Animal Science*, Vol.82, No.10, (October 2004), pp. 2930-2936
- Maejima, Y.; Sedbazar, U.; Suyama, S.; Kohno, D.; Onaka, T.; Takano, E.; Yoshida, N.; Koike, M.; Uchiyama, Y.; Fujiwara, K.; Yashiro, T.; Horvath, T. L.; Dietrich, M. O.; Tanaka, S.; Dezaki, K.; Oh-I, S.; Hashimoto, K.; Shimizu, H.; Nakata, M.; Mori, M. & Yada, T. (2009). Nesfatin-1-Regulated Oxytocinergic Signaling in the Paraventricular Nucleus Causes Anorexia Through a Leptin-Independent Melanocortin Pathway. *Cell Metabolism*, Vol.10, No.5, (November 2009), pp. 355-365
- Matkovic, V.; Ilich, J. Z.; Skugor, M.; Badenhop, N. E.; Goel, P.; Clairmont, A.; Klisovic, D.; Nahhas, R. W. & Landoll, J. D. (1997). Leptin Is Inversely Related to Age at Menarche in Human Females. *The Journal of Clinical Endocrinology & Metabolism*, Vol.82, No.10, (October 1997), pp. 3239-3245
- Messenger, S.; Chatzidaki, E. E.; Ma, D.; Hendrick, A. G.; Zahn, D.; Dixon, J.; Thresher, R. R.; Malinge, I.; Lomet, D.; Carlton, M. B. L.; Colledge, W. H.; Caraty, A. & Aparicio, S. A. J. R. (2005). Kisspeptin Directly Stimulates Gonadotropin-Releasing Hormone Release Via G Protein-Coupled Receptor 54. *Proceedings of the National Academy of Sciences of the United States of America*, Vol.102, No.5, (February 2005), pp. 1761-1766
- Miller, D. W.; Findlay, P. A.; Morrison, M. A.; Raver, N. & Adam, C. L. (2002). Seasonal and Dose-Dependent Effects of Intracerebroventricular Leptin on LH Secretion and Appetite in Sheep. *The Journal of Endocrinology*, Vol.175, No.2, (November 2002), pp. 395-404
- Miller, J. C.; Gnaedinger, J. M. & Rapoport, S. I. (1987). Utilization of Plasma Fatty Acid in Rat Brain: Distribution of [¹⁴C]Palmitate Between Oxidative and Synthetic Pathways. *Journal of Neurochemistry*, Vol.49, No.5, (November 1987), pp. 1507-1514
- Morrison, C. D.; Daniel, J. A.; Holmberg, B. J.; Djiane, J.; Raver, N.; Gertler, A. & Keisler, D. H. (2001). Central Infusion of Leptin into Well-Fed and Undernourished Ewe Lambs: Effects on Feed Intake and Serum Concentrations of Growth Hormone and Luteinizing Hormone. *The Journal of Endocrinology*, Vol.168, No.2, (February 2001), pp. 317-324
- Nakata, M.; Manaka, K.; Yamamoto, S.; Mori, M. & Yada, T. (2011). Nesfatin-1 Enhances Glucose-Induced Insulin Secretion by Promoting Ca²⁺ Influx Through L-Type Channels in Mouse Islet β -Cells. *Endocrine Journal*, Vol.58, No.4, (April 2011), pp. 305-313
- Navarro, V. M.; Castellano, J. M.; Fernández-Fernández, R.; Barreiro, M. L.; Roa, J.; Sanchez-Criado, J. E.; Aguilar, E.; Dieguez, C.; Pinilla, L. & Tena-Sempere, M. (2004a). Developmental and Hormonally Regulated Messenger Ribonucleic Acid Expression of KiSS-1 and Its Putative Receptor, GPR54, in Rat Hypothalamus and Potent Luteinizing Hormone-Releasing Activity of KiSS-1 Peptide. *Endocrinology*, Vol.145, No.10, (October 2004), pp. 4565-4574
- Navarro, V. M.; Fernández-Fernández, R.; Castellano, J. M.; Roa, J.; Mayen, A.; Barreiro, M. L.; Gaytan, F.; Aguilar, E.; Pinilla, L.; Dieguez, C. & Tena-Sempere, M. (2004b). Advanced Vaginal Opening and Precocious Activation of the Reproductive Axis by KiSS-1 Peptide, the Endogenous Ligand of GPR54. *The Journal of Physiology*, Vol.561, No.2, (December 2004), pp. 379-386

- Navarro, V. M.; Castellano, J. M.; Fernández-Fernández, R.; Tovar, S.; Roa, J.; Mayen, A.; Barreiro, M. L.; Casanueva, F. F.; Aguilar, E.; Dieguez, C.; Pinilla, L. & Tena-Sempere, M. (2005). Effects of KiSS-1 Peptide, the Natural Ligand of GPR54, on Follicle-Stimulating Hormone Secretion in the Rat. *Endocrinology*, Vol.146, No.4, (April 2005), pp. 1689-1697
- Obici, S.; Feng, Z.; Morgan, K.; Stein, D.; Karkanias, G. & Rossetti, L. (2002). Central Administration of Oleic Acid Inhibits Glucose Production and Food Intake. *Diabetes*, Vol.51, No.2, (February 2002), pp. 271-275
- Obici, S. (2009). Molecular Targets for Obesity Therapy in the Brain. *Endocrinology*, Vol.150, No.6, (June 2009), pp. 2512-2517
- Ogiso, K.; Asakawa, A.; Amitani, H.; Nakahara, T.; Ushikai, M.; Haruta, I.; Koyama, K.-I.; Amitani, M.; Harada, T.; Yasuhara, D. & Inui, A. (2011). Plasma Nesfatin-1 Concentrations in Restricting-Type Anorexia Nervosa. *Peptides*, Vol.32, No.1, (January 2011), pp. 150-153
- Oh-I, S.; Shimizu, H.; Satoh, T.; Okada, S.; Adachi, S.; Inoue, K.; Eguchi, H.; Yamamoto, M.; Imaki, T.; Hashimoto, K.; Tsuchiya, T.; Monden, T.; Horiguchi, K.; Yamada, M. & Mori, M. (2006). Identification of Nesfatin-1 As a Satiety Molecule in the Hypothalamus. *Nature*, Vol.443, No.7112, (October 2006), pp. 709-712
- Pan, W.; Hsueh, H. & Kastin, A. J. (2007). Nesfatin-1 Crosses the Blood-Brain Barrier Without Saturation. *Peptides*, Vol.28, No.11, (November 2007), pp. 2223-2228
- Price, T. O.; Samson, W. K.; Niehoff, M. L. & Banks, W. A. (2007). Permeability of the Blood-Brain Barrier to a Novel Satiety Molecule Nesfatin-1. *Peptides*, Vol.28, No.12, (December 2007), pp. 2372-2381
- Price, C. J.; Samson, W. K. & Ferguson, A. V. (2008). Nesfatin-1 Inhibits NPY Neurons in the Arcuate Nucleus. *Brain Research*, Vol.1230, (September 2008), pp. 99-106
- Qian, H.; Barb, C. R.; Compton, M. M.; Hausman, G. J.; Azain, M. J.; Kraeling, R. R. & Baile, C. A. (1999). Leptin mRNA Expression and Serum Leptin Concentrations As Influenced by Age, Weight, and Estradiol in Pigs. *Domestic Animal Endocrinology*, Vol.16, No.2, (February 1999), pp. 135-143
- Qiu, J.; Fang, Y.; Bosch, M. A.; Rønnekleiv, O. K. & Kelly, M. J. (2011). Guinea Pig Kisspeptin Neurons Are Depolarized by Leptin Via Activation of TRPC Channels. *Endocrinology*, Vol.152, No.4, (April 2011), pp. 1503-1514
- Quennell, J. H.; Howell, C. S.; Roa, J.; Augustine, R. A.; Grattan, D. R. & Anderson, G. M. (2011). Leptin Deficiency and Diet-Induced Obesity Reduce Hypothalamic Kisspeptin Expression in Mice. *Endocrinology*, Vol.152, No.4, (April 2011), pp. 1541-1550
- Ramanjaneya, M.; Chen, J.; Brown, J. E.; Tripathi, G.; Hallschmid, M.; Patel, S.; Kern, W.; Hillhouse, E. W.; Lehnert, H.; Tan, B. K. & Rande, H. S. (2010). Identification of Nesfatin-1 in Human and Murine Adipose Tissue: A Novel Depot-Specific Adipokine With Increased Levels in Obesity. *Endocrinology*, Vol.151, No.7, (July 2010), pp. 3169-3180
- Rapoport, S. (1996). *In Vivo* Labeling of Brain Phospholipids by Long-Chain Fatty Acids: Relation to Turnover and Function. *Lipids*, Vol.31, No.1, (March 1996), p. S97-S101

- Redmond, J. S.; Baez-Sandoval, G. M.; Spell, K. M.; Spencer, T. E.; Lents, C. A.; Williams, G. L. & Amstalden, M. (2011). Developmental Changes in Hypothalamic *Kiss1* Expression During Activation of the Pulsatile Release of Luteinising Hormone in Maturing Ewe Lambs. *Journal of Neuroendocrinology*, Vol.23, No.9, (September 2011), pp. 815-822
- Renes, J.; Rosenow, A. & Mariman, E. (2009). Novel Adipocyte Features Discovered by Adipoproteomics. *Adipobiology*, Vol.1, (2009), pp. 7-18
- Roa, J.; García-Galiano, D.; Castellano, J. M.; Gaytan, F.; Pinilla, L. & Tena-Sempere, M. (2010). Metabolic Control of Puberty Onset: New Players, New Mechanisms. *Molecular and Cellular Endocrinology*, Vol.324, No.1-2, (August 2010), pp. 87-94
- Rodriguez-Pacheco, F.; Martinez-Fuentes, A. J.; Tovar, S.; Pinilla, L.; Tena-Sempere, M.; Dieguez, C.; Castaño, J. P. & Malagon, M. M. (2007). Regulation of Pituitary Cell Function by Adiponectin. *Endocrinology*, Vol.148, No.1, (January 2007), pp. 401-410
- Schäffler, A. & Schölmerich, J. (2010). Innate Immunity and Adipose Tissue Biology. *Trends in Immunology*, Vol.31, No.6, (June 2010), pp. 228-235
- Shahab, M.; Mastronardi, C.; Seminara, S. B.; Crowley, W. F.; Ojeda, S. R. & Plant, T. M. (2005). Increased Hypothalamic GPR54 Signaling: A Potential Mechanism for Initiation of Puberty in Primates. *Proceedings of the National Academy of Sciences of the United States of America*, Vol.102, No.6, (February 2005), pp. 2129-2134
- Shimizu, H.; Ohsaki, A.; Oh-I, S.; Okada, S. & Mori, M. (2009). A New Anorexigenic Protein, Nesfatin-1. *Peptides*, Vol.30, No.5, (May 2009), pp. 995-998
- Smith, J. T.; Cunningham, M. J.; Rissman, E. F.; Clifton, D. K. & Steiner, R. A. (2005). Regulation of *Kiss1* Gene Expression in the Brain of the Female Mouse. *Endocrinology*, Vol.146, No.9, (September 2005), pp. 3686-3692
- Smith, J. T.; Acohido, B. V.; Clifton, D. K. & Steiner, R. A. (2006). KiSS-1 Neurones Are Direct Targets for Leptin in the *Ob/Ob* Mouse. *Journal of Neuroendocrinology*, Vol.18, No.4, (April 2006), pp. 298-303
- Smith, J. T.; Clay, C. M.; Caraty, A. & Clarke, I. J. (2007). KiSS-1 Messenger Ribonucleic Acid Expression in the Hypothalamus of the Ewe Is Regulated by Sex Steroids and Season. *Endocrinology*, Vol.148, No.3, (March 2007), pp. 1150-1157
- Smith, J. T.; Li, Q.; Yap, K. S.; Shahab, M.; Roseweir, A. K.; Millar, R. P. & Clarke, I. J. (2011). Kisspeptin Is Essential for the Full Preovulatory LH Surge and Stimulates GnRH Release From the Isolated Ovine Median Eminence. *Endocrinology*, Vol.152, No.3, (March 2011), pp. 1001-1012
- Stengel, A.; Goebel, M.; Yakubov, I.; Wang, L.; Witcher, D.; Coskun, T.; Taché, Y.; Sachs, G. & Lambrecht, N. W. G. (2009a). Identification and Characterization of Nesfatin-1 Immunoreactivity in Endocrine Cell Types of the Rat Gastric Oxyntic Mucosa. *Endocrinology*, Vol.150, No.1, (January 2009), pp. 232-238
- Stengel, A.; Goebel, M.; Wang, L.; Rivier, J.; Kobelt, P.; Mönnikes, H.; Lambrecht, N. W. G. & Taché, Y. (2009b). Central Nesfatin-1 Reduces Dark-Phase Food Intake and Gastric Emptying in Rats: Differential Role of Corticotropin-Releasing Factor₂ Receptor. *Endocrinology*, Vol.150, No.11, (November 2009), pp. 4911-4919

- Su, Y.; Zhang, J.; Tang, Y.; Bi, F. & Liu, J.-N. (2010). The Novel Function of Nesfatin-1: Anti-Hyperglycemia. *Biochemical and Biophysical Research Communications*, Vol.391, No.1, (January 2010), pp. 1039-1042
- Tadross, J. A.; Patterson, M.; Wynne, K. J.; Patel, S.; Suzuki, K.; Ghatei, M. A. & Bloom, S. R. (2010). Nesfatin Suppresses Feeding and Stimulates the Hypothalamo-Pituitary-Gonadal Axis. *Endocrine Journal*, Vol.57 (Supplement 2), (2010), p. S442 (P3-1-3)
- Tomikawa, J.; Homma, T.; Tajima, S.; Shibata, T.; Inamoto, Y.; Takase, K.; Inoue, N.; Ohkura, S.; Uenoyama, Y.; Maeda, K. & Tsukamura, H. (2010). Molecular Characterization and Estrogen Regulation of Hypothalamic *KISS1* Gene in the Pig. *Biology of Reproduction*, Vol.82, No.2, (February 2010), pp. 313-319
- True, C.; Kirigiti, M.; Ciofi, P.; Grove, K. L. & Smith, M. S. (2011). Characterisation of Arcuate Nucleus Kisspeptin/Neurokinin B Neuronal Projections and Regulation During Lactation in the Rat. *Journal of Neuroendocrinology*, Vol.23, No.1, (January 2011), pp. 52-64
- Wahab, F.; Ullah, F.; Chan, Y.-M.; Seminara, S. B. & Shahab, M. (2011). Decrease in Hypothalamic *Kiss1* and *Kiss1r* Expression: A Potential Mechanism for Fasting-Induced Suppression of the HPG Axis in the Adult Male Rhesus Monkey (*Macaca Mulatta*). *Hormone and Metabolic Research*, Vol.43, No.2, (February 2011), pp. 81-85
- Wen, J.-P.; Lv, W.-S.; Yang, J.; Nie, A.-F.; Cheng, X.-B.; Yang, Y.; Ge, Y.; Li, X.-Y. & Ning, G. (2008). Globular Adiponectin Inhibits GnRH Secretion From GT1-7 Hypothalamic GnRH Neurons by Induction of Hyperpolarization of Membrane Potential. *Biochemical and Biophysical Research Communications*, Vol.371, No.4, (July 2008), pp. 756-761
- Whisnant, C. S. & Harrell, R. J. (2002). Effect of Short-Term Feed Restriction and Refeeding on Serum Concentrations of Leptin, Luteinizing Hormone and Insulin in Ovariectomized Gilts. *Domestic Animal Endocrinology*, Vol.22, No.2, (April 2002), pp. 73-80
- Woods, S. C.; Seeley, R. J.; Porte, D., Jr. & Schwartz, M. W. (1998). Signals That Regulate Food Intake and Energy Homeostasis. *Science*, Vol.280, No.5368, (May 1998), pp. 1378-1383
- Xu, J.; Kirigiti, M. A.; Grove, K. L. & Smith, M. S. (2009). Regulation of Food Intake and Gonadotropin-Releasing Hormone/Luteinizing Hormone During Lactation: Role of Insulin and Leptin. *Endocrinology*, Vol.150, No.9, (September 2009), pp. 4231-4240
- Yamada, S.; Uenoyama, Y.; Kinoshita, M.; Iwata, K.; Takase, K.; Matsui, H.; Adachi, S.; Inoue, K.; Maeda, K.-I. & Tsukamura, H. (2007). Inhibition of Metastin (Kisspeptin-54)-GPR54 Signaling in the Arcuate Nucleus-Median Eminence Region During Lactation in Rats. *Endocrinology*, Vol.148, No.5, (May 2007), pp. 2226-2232
- Zegers, D.; Beckers, S.; Mertens, I. L.; Van Gaal, L. F. & Van Hul, W. (2011). Association Between Polymorphisms of the *Nesfatin* Gene, *NUCB2*, and Obesity in Men. *Molecular Genetics and Metabolism*, Vol.103, No.3, (July 2011), pp. 282-286
- Zhang, A.-Q.; Li, X.-L.; Jiang, C.-Y.; Lin, L.; Shi, R.-H.; Chen, J.-D. & Oomura, Y. (2010). Expression of Nesfatin-1/NUCB2 in Rodent Digestive System. *World Journal of Gastroenterology*, Vol.16, No.14, (April 2010), pp. 1735-1741

- Zhong, J.; Krawczyk, S. A.; Chaerkady, R.; Huang, H.; Goel, R.; Bader, J. S.; Wong, G. W.; Corkey, B. E. & Pandey, A. (2010). Temporal Profiling of the Secretome During Adipogenesis in Humans. *Journal of Proteome Research*, Vol.9, No.10, (August 2010), pp. 5228-5238
- Zieba, D. A.; Amstalden, M.; Morton, S.; Maciel, M. N.; Keisler, D. H. & Williams, G. L. (2004). Regulatory Roles of Leptin at the Hypothalamic-Hypophyseal Axis Before and After Sexual Maturation in Cattle. *Biology of Reproduction*, Vol.71, No.3, (September 2004), pp. 804-812