We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Potential Therapeutic Strategies to Prevent the Progression of Alzheimer to Disease States

Ester Aso and Isidre Ferrer

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/54783>

1. Introduction

Alzheimer is an age-dependent neurodegenerative process distinct from normal aging and characterized morphologically by the presence of senile plaques and neurofibrillary tangles, which progress from the brain stem and inner parts of the temporal lobes to most the telencephalon.

Senile plaques are mainly composed of different species of fibrillar β -amyloid ($A\beta$), a product of the cleavage of the β -amyloid precursor protein (APP), and they are surrounded by dystrophic neurites, reactive astrocytes and microglia. $A\beta$ fibrillar deposits also occur in diffuse plaques, subpial deposits and in the wall of the cerebral and meningeal blood vessels in the form of amyloid angiopathy. A substantial part of β -amyloid is not fibrillar but soluble and forms oligomers of differing complexity which are toxic to nerve cells.

Neurofibrillary tangles are mainly composed of various isoforms of tau protein, which is hyper-phosphorylated and nitrated. It has an altered conformation and is truncated at different sites through the action of a combination of several proteolytic enzymes giving rise to species of low molecular weight which are toxic to nerve cells. Abnormal tau deposition also occurs in the dystrophic neurites of senile plaques and within the small neuronal processes, resulting in the formation of neuropil threads.

The mechanisms of disease progression are not completely understood but $A\beta$ initiates the pathological process in the small percentage of familial cases due to mutations in genes encoding APP, presenilin 1 and presenilin 2, the latter involved in the cleavage of APP, and potentiates tau phosphorylation in sporadic cases that represent the majority of affected individuals (β -amyloid cascade hypothesis). Moreover, $A\beta$ act as a seed of new β -amyloid production and deposition under appropriate settings, and abnormal tau promotes the

production and deposition of hyper-phosphorylated tau. Therefore, A β and hyper-phosphorylated tau promote the progression of the process and this may occur in an exponential way once these abnormal proteins are accumulated in the brain.

In addition to these pathological hallmarks, multiple alterations play roles in the degenerative process. Several genetic factors, such as apolipoprotein ϵ 4 (APOE4), and external factors, such as vascular and circulatory alterations and repeated cerebral traumatism, among others, facilitate disease progression in sporadic forms. Furthermore, metabolic components mainly, but not merely, associated with aging have a cardinal influence, including mitochondrial defects and energy production deficiencies, production of free radicals (oxidative and nitrosative reactive species: ROS and NOS) and oxidative and nitrosative damage, increased reticulum stress damage, altered composition of membranes, inflammatory responses and impaired function of degradation pathways such as autophagy and ubiquitin-proteasome system.

It has been proven that the degenerative process, at least the presence of neurofibrillary tangles, starts in middle age in selected nuclei of the brain stem and entorhinal cortex, and then progresses to other parts of the brain. Instrumental stages of Braak cover stages I and II with involvement of the entorhinal and transentorhinal cortices; stages II and IV also affect the hippocampus and limbic system together with the basal nucleus of Meynert; and stages V and VI involve the whole brain although neurofibrillary tangles are not found in selected regions such as the cerebellar cortex and the dentate gyrus. The distribution of senile plaques is a bit different as they first appear in the orbitofrontal cortex and temporal cortex and then progress to the whole convexity.

A concomitant decline in neuronal organization occurs most often in parallel with senile plaques and neurofibrillary tangles manifested as synaptic dysfunction and synaptic loss, and neuronal death and progressive isolation of remaining neurons.

An important observation is that about 80% of individuals aged 65 years have Alzheimer-related changes, at least at stages I-III, whereas only 5% have cognitive impairment and dementia. About 25% of individuals aged 85 years suffer from cognitive impairment and dementia of Alzheimer type. Stages I-IV are often silent with no clinical symptoms. Cognitive impairment and dementia usually occur at stages V and VI when the neurodegenerative process is very advanced. Importantly, the progression from stage I to stage IV may last decades, whereas the progression to stages V and VI is much more rapid. Therefore, Alzheimer is a well-tolerated degenerative process during a relatively long period of time, but it may have devastating effects once thresholds are crossed. Moreover, clinical symptoms may be complicated by concomitant vascular pathology.

Several attempts have been made to predict the evolution to disease states. Neuroimaging, including high resolution and functional magnetic resonance imaging, positron emission tomography and the use of relative selective markers of β -amyloid and tau deposition in the brain, together with reduced levels of A β and increased index of phospho-tau/total tau in the cerebrospinal fluid, are common complementary probes (biomarkers) in addition to the data

provided by the neuropsychological examination. Unfortunately, these tests, at present, detect relatively advanced stages of the process in pathological terms.

It is very illustrating to visualize under the microscope how a brain at middle stages of the degenerative process has been working without apparent neurological deficits during life. The adaptive capacities of the brain in coping with current functions in spite of the decrepitude of composition and organization resulting from the chronic progression of the degenerative process are impressive.

Taking into consideration this scenario, it is compulsory to increase understanding of the first stages of the degenerative process and to act on selective targets before the appearance of clinical symptoms.

The present review is not a mere list of putative treatments of Alzheimer's disease (AD) but rather an approach to learning about observations made on experimental models and early stages of disease aimed at curbing or retarding disease progression on the basis of definite rationales. It is also our aim to encourage the consideration of Alzheimer as a degenerative process not necessarily leading to dementia [1]. This concept has important clinical implications as it supports early preventive measures in the population at risk (i.e. persons over 50 years) even in the absence of clinical symptoms.

2. Experimental therapeutic strategies to prevent Alzheimer progression to Alzheimer Disease (AD) states

Several reviews have focused on various aspects related to habits and dietary elements which may act as protective factors against AD, including physical and mental exercise, low caloric intake, various diets with low fat content, and vitamin complements [2, 3]. It is worth noting that neuropathological studies in old-aged individuals usually present combined pathologies, and combination of Alzheimer changes and vascular lesions are very common [4]. It is well documented that vascular pathology potentiates primary neurodegenerative pathology and that vascular factors may be causative of cognitive impairment and dementia [5]. Therefore, therapies geared to reduce vascular risk factors are also protective factors against AD clinical manifestations.

2.1. Targeting A β

Most of the current drug development for the prevention or treatment of AD is based on the β -amyloid cascade hypothesis and aims at reducing the levels of A β in the brain. Overproduction, aggregation and deposition of the A β peptide begin before the onset of symptoms and they are considered an essential early event in AD pathogenesis. Thus, targeting these early A β alterations is assumed to reduce the progression to disease states. The different strategies developed to achieve this objective include decreasing A β production through modulating secretase activity, interfering with A β aggregation, and promoting A β clearance.

2.1.1. Secretase-targeting therapies

APP is processed in the brain exclusively by three membrane-bound proteases, α -, β - and γ -secretase. Therefore, specifically modifying such enzyme activity should result in a reduction of A β production [6].

- *α -secretase activators*: α -secretase initiates the non-amyloidogenic pathway by cleaving APP within the A β sequence, thereby preventing the production of A β and producing a non-toxic form of APP derivative which is neuroprotective and growth-promoting [7]. Therefore, compounds that stimulate α -secretase activity could become an attractive strategy to reduce A β production. In fact, some indirect methods of promoting α -secretase activity, such as the stimulation of the protein kinase C (PKC) or Mitogen-activated protein kinases (MAPK) pathways, the use of α -7-nicotinic acetylcholine (ACh) receptor and 5-hydroxytryptamine (5-HT) receptor 4 agonists, and γ -aminobutyric acid A receptor modulators, result in α -secretase-mediated cleavage of APP and reduced A β levels *in vivo* [8]. However, the development of a direct activator of α -secretase as a drug treatment for AD seems premature because of the lack of knowledge about the consequences of chronic up-regulation of α -secretase-mediated cleavage on other substrates [6].
- *β -secretase inhibitors*: the β -secretase enzyme initiates the amyloidogenic pathway, cleaving APP at the amino terminus of the A β peptide. Further cleavage of the resulting carboxy-terminal fragment by γ -secretase results in the release of A β . β -secretase activity is specifically mediated by the β -site APP cleaving enzyme 1 (BACE1), which is also involved in the processing of numerous substrates in addition to APP. The research of drugs inhibiting BACE1 activity was encouraged by studies revealing that the expression of mutated BACE1 reduces amyloidogenesis and cognitive impairment in APP transgenic mice [9, 10]. The first generation of BACE1 inhibitors was peptide-based mimetics of the APP β -cleavage site. Unfortunately, these compounds exhibited some difficulties because of the large substrate binding site of BACE1 and because of the difficulty in crossing the blood-brain barrier (BBB) and penetrating the plasma and endosomal membranes to gain access to the intracellular compartments where endogenous BACE1 plays its function. Recently, non-peptide small-molecule BACE1 inhibitors have been reported to improve bioavailability and to lower cerebral A β levels in animal models of AD [11, 12]. However, the involvement of BACE1 in other important physiological processes raises concerns about minimizing the potential adverse effects derived from generalized BACE1 inhibition.
- *γ -secretase inhibitors (GSIs)*: γ -secretase is a complex composed of presenilin 1 and presenilin 2 (PS1 and PS2) forming the catalytic core and three accessory proteins, anterior pharynx-defective 1 (APH-1), nicastrin and presenilin enhancer protein 2 (PEN2). The γ -secretase complex displays a high degree of subunit heterogeneity and little is known about the physiological roles of the diverse complexes and how they process different trans-membrane substrates in addition to APP. This heterogeneity suggests that selective targeting of one particular subunit might be a more effective treatment strategy than non-selective γ -secretase inhibition [13]. Thus, removal of APH-1B and APH-1C isoforms in a mouse model of AD decreased A β plaque formation and improved behavioral deficits [14]. A number of orally bioavailable and brain-penetrating GSIs have been shown to decrease A β production

and deposition in APP mouse models and in humans [15-17]. However, target-based toxicity of GSIs has been a major obstacle to the clinical development of these compounds. In fact, two large Phase III clinical trials of *Semagacestat*, the only GSI extensively studied in AD, were prematurely interrupted because of the observation of detrimental cognitive and functional effects of the drug [18]. Several dozen γ -secretase substrates have been identified, including Notch1 trans-membrane receptor, which plays an important role in a variety of developmental and physiological processes by controlling cell fate decisions. To overcome these toxicity issues, pharmaceutical companies have been trying to develop a second generation of 'Notch-sparing' GSIs, which revealed beneficial effects in *in vitro* and in animal models of AD [19-21]. They are currently under clinical studies. Such 'Notch-sparing' GSIs have higher pharmacological selectivity than the first GSIs probably due to the distinct binding to the substrate docking site on γ -secretase of Notch and APP. Identification of several γ -secretase inhibitors has been reviewed elsewhere [22].

2.1.2. $A\beta$ degrading enzymes

Almost 20 enzymes are currently known to contribute to $A\beta$ degradation in the brain, although the most studied are two zinc metalloproteases, neprilysin (NEP) and insulin-degrading enzyme (IDE). NEP is one of the major $A\beta$ -degrading enzymes in the brain [23] and NEP levels are decreased in the brain of AD and animal models [24, 25]. Lentiviral delivery of the NEP gene to the brain of AD transgenic mice reduced $A\beta$ pathology [26]. A number of subsequent studies with NEP and other related peptidases such as endothelin-converting enzymes 1 and 2 (ECE-1 and ECE-2) further supported this observation [27]. Similarly, over-expression of IDE in neurons significantly reduces brain $A\beta$ levels, prevents $A\beta$ plaque formation and its associated cytopathology, and rescues the premature lethality present in these particular APP transgenic mice [28]. A growing body of evidence has been accumulated supporting the potential therapeutic properties of IDE in AD [29].

Other specific $A\beta$ -cleaving proteases such as angiotensin-converting enzyme (ACE), matrix metalloproteinase-9 (MMP-9) and the serine protease plasmin, which have distinct sub-cellular localizations and differential responses to aging, oxidative stress and pharmacological agents, are also potential candidates to become novel therapeutic strategies for AD prevention and treatment [27].

Targeting the delivery of these compounds to the brain remains a major challenge. The most promising current approaches include peripheral administration of agents that enhance the activity of $A\beta$ -degrading enzymes and direct intra-cerebral release of enzymes by convection-enhanced delivery. Genetic procedures geared at increasing cerebral expression of $A\beta$ -degrading enzymes may offer additional advantages [30].

2.1.3. Decreasing $A\beta$ aggregation

Compounds that suppress the aggregation or reduce the stability of $A\beta$ oligomers may bind monomers in order to attenuate formation of both the oligomeric and senile plaque fibrillar $A\beta$ constituents. One of the amyloid-binding drugs more extensively studied in animal models

and AD patients is tramiprosate (3-amino-1-propanesulfonic acid; Alzhemed). Tramiprosate was effective in reducing A β polymerisation *in vitro*, inhibiting the formation of neurotoxic aggregates, and decreasing A β plaque formation in animal models [31]. However, recent phase III clinical trials did not produce any significant improvement in cognition in AD patients chronically treated with tramiprosate in spite of the significant reduction in hippocampus volume loss [32]. Similarly, some other compounds known to inhibit A β aggregation and fibril formation showed positive effects in animal and *in vitro* models of AD but failed to produce conclusive results in human clinical trials. This is the case with scyllo-inositol and PBT2. Scyllo-inositol inhibited cognitive deficits in TgCRND8 mice and significantly ameliorated disease pathology, even in animals at advanced stages of AD-like pathology, without interfering with endogenous phosphatidylinositol lipid production [33, 34]. Yet a phase II clinical trial failed in supporting or refuting a benefit of scyllo-inositol in mild to moderate AD patients [35]. PBT2 is a copper/zinc ionophore which targets metal-induced aggregation of A β . When given orally to two models of A β -bearing transgenic mice, PBT2 was able to markedly decrease soluble brain A β levels within hours and to improve cognitive performance within days [36]. These results correlated with a rapid cognitive improvement in AD patients in a recent phase IIa clinical trial [37], an observation that argues for large-scale testing of PBT2 for AD.

Another promising recent experimental approach is the use of dendrimers as agents interfering with A β fibrilization. Dendrimers are globular branched polymers, typically symmetric around the core with a spherical three-dimensional morphology. Their chemical structure allows dendrimers to couple to active amyloid species through hundreds of possible sites. Dendrimers have been shown to be able to modulate A β peptide aggregation by interfering in different ways with the polymerization process, including fibril breaking, inhibition of fibril formation and acceleration of fibril formation [38, 39]. However, some dendrimers assayed in amyloidogenic systems are toxic to cells. The development of non-toxic glycodendrimers, which reduce toxicity by clumping fibrils together [40], opens the possibility of using dendrimers with low intrinsic toxicity in AD. Additional difficulties in dendrimer administration involve the crossing of the BBB so as to reach their targets in the brain.

2.1.4. Facilitating A β clearance: Immunotherapy against A β

Active and passive immunotherapy against A β peptide has been explored as a therapeutic approach to stimulate the clearance of A β in the brain at the preclinical and clinical stages of the disease in animal models. Pioneering studies proved that vaccination of young APP transgenic mice using a synthetic aggregated form of A β_{42} (AN-1792) effectively prevented A β plaque formation, neuritic dystrophy and astrogliosis in adult brains [41]. Subsequent studies further demonstrated improvement of memory loss in those APP transgenic mice vaccinated against A β [42, 43]. Different models, methods and ways of administration showed the beneficial effects of active and passive immunization in animal models of AD. Nevertheless, the phase II trial in humans was discontinued because of the occurrence of aseptic meningoencephalitis in a number of cases [44-46]. The cause of the meningoencephalitis was a concomitant T-cell-mediated autoimmune response [45, 46]. Moreover, several studies in APP transgenic mice have reported an increased risk of microhemorrhages at sites of cerebro-

vascular A β deposits [47]. Yet important conclusions were drawn from the studies in humans: immunization reduced the number of A β plaques and the number of dystrophic neurites, including tau phosphorylation around plaques, but not A β burden in blood vessels; however, immunization increased intracerebral levels of soluble A β [48-50].

New vaccines containing immunodominant B-cell epitopes of A β [51] and recognizing other A β residues [52, 53], and the use of passive immunization with deglycosylated antibodies [54] have demonstrated positive effects in the clearance of A β without causing inflammatory response or hemorrhages in animal models of AD [55]. These findings have prompted new clinical trials which are currently evaluating the toxicity and effectiveness of at least ten vaccines in mild-to-moderate AD patients worldwide [56]. While vaccines hold great hope as AD therapies, it is important to stress that immunization at pre-symptomatic stages is essential in order to avoid the irreversible brain damage occurring even at the early symptomatic stages [57].

2.2. Targeting tau

The interest in tau-related therapies is still emerging and very few clinical studies are underway, in part because of the difficulties encountered with anti-A β strategies that captured most efforts in the two last decades, but also because of the challenging identification of tractable therapeutic targets related to tau. Current research in the prevention of tau pathology developed in animal models of AD has resulted in some promising results [58]. Main rationales in tau pathology are based on: 1: inhibition of tau aggregation, 2: reduction of tau phosphorylation by inhibition of tau kinases or activation of phosphatases (including PP2a activity), 3: reduction of tau levels by increasing tau degradation or by using active immunization, and 4: stabilization of microtubule [59].

2.2.1. Inhibition of tau aggregation

Some compounds that are known to inhibit tau-tau interactions have been tested as agents aimed at slowing Alzheimer progression to disease states. Among them, phenothiazine methylene blue inhibits tau-tau interactions, is neuroprotective and is able to facilitate soluble tau clearance in a mouse model of human tauopathy [60, 61]. Moreover, phenothiazine methylene blue has shown beneficial effects in a phase II clinical trial conducted for one year [62]. Another promising inhibitor of tau aggregation is the immunosuppressant FK506, which exerts its beneficial effects in transgenic mice by directly binding tau to the FK506 binding protein 52 and by modulating microglial activation [63, 64].

However, some concerns arise from the use of tau aggregation inhibitors in that at least some tau aggregation inhibitors enhance the formation of potentially toxic tau oligomers [65].

2.2.2. Reduction of tau hyperphosphorylation

Kinases which participate in the phosphorylation of tau and phosphatases which dephosphorylate tau are clear putative therapeutic targets for AD [66]. The most widely studied tau kinases in AD pathogenesis are Glycogen synthase kinase 3 beta (GSK-3 β) and Cyclin-

dependent kinase (CDK5) [67, 68]. Several GSK-3 β inhibitors, including lithium, aloisines, flavopiridol, hymenialdisine, paullones, and staurosporine, are under active investigation and development [69]. Lithium revealed some promising results when administered in transgenic mice expressing the P301L human 4R0N tau at pre-symptomatic stages; it improved behavior and reduced the levels of phosphorylation, aggregation and insoluble tau in transgenic mice [70]. However, several concerns have arisen in relation of the use of GSK-3 β in the treatment of AD; these are based on the fact that lithium lacks specificity over GSK-3 β activity and it has a narrow safety margin [71]. Moreover, GSK-3 β acts on multiple metabolic pathways that are also impaired with unknown consequences after chronic treatment.

CDK5 inhibitors prevent A β -induced tau hyper-phosphorylation and cell death *in vitro* [72, 73]. A recent *in vivo* study further demonstrates that inhibition of CDK5 activates GSK-3 β , which plays a more dominant role in overall tau phosphorylation than does CDK5 [74]. Thus, considering that CDK5 inhibitors might be unable to reverse abnormal hyper-phosphorylation of tau and treat neurofibrillary degeneration because of the interplay between CDK5 and GSK-3 β , as well as the essential role played by CDK5 in multiple cell signaling pathways [75], the interest of such compounds as a tau-targeting therapy for AD is limited.

Another approach to reverse tau hyper-phosphorylation is up-regulation of tau phosphatases [66]. The major tau phosphatase, PP2A, is down-regulated in AD brain. In consequence, correcting PP2A levels is the primary target to be considered. Among the compounds known to reverse PP2A inhibition, memantine is the most outstanding because of the demonstrated clinical benefit in AD. In an animal model, memantine was able to reverse okadaic acid-induced PP2A inhibition and to prevent tau hyper-phosphorylation, restoring MAP2 expression [76]. Similarly, melatonin has also been shown to restore PP2A activity and reverse tau hyper-phosphorylation, both *in vitro* and in experimental animals [77]. One important concern in considering PP2A as a potential therapeutic target is that all protein phosphatases have much broader substrate specificities than protein kinases. Thus, more undesirable effects might be expected than when using kinase inhibitors [66]. A further intriguing point is that PP2A function and activity depend on multiple subunits and cofactors which are dysregulated in AD [78]. It is not clear how all these elements can be resolved to result in maintained balanced activity.

2.2.3. Reduction of tau levels

A potential alternative to modulate tau phosphorylation is reducing overall tau levels [58]. Experiments carried out in genetically-modified mice expressing reduced tau levels revealed diminished cognitive impairment and A β -induced neuronal damage [79-81]. An alternative method to reduce tau levels could be by targeting molecules that regulate the expression or clearance of tau. Tau can be degraded via the ubiquitin-proteasome system and the lysosomal pathways. Reduction of the levels of the tau ubiquitin-ligase CHIP increases the accumulation of tau aggregates in JNPL3 mice, suggesting that increasing the expression of CHIP could result in reduced tau levels [82]. Acetylation of tau inhibits its degradation [83], alters its microtubule binding, and enhances aggregation [84]. Thus, the combination of tau acetylation inhibition and ubiquitination-proteasome enhancement might produce a synergy that lowers the levels of pathogenic tau species.

Tau degradation can also be enhanced by immunization. Active immunization targeting phosphorylated tau reduces filamentous tau inclusions and neuronal dysfunction in JNPL3 transgenic mice [85, 86]. Moreover, recent studies have raised the possibility of modulating tau pathology by passive immunization revealing reduced behavioral impairment and tau pathology in two transgenic models of tauopathies [87].

2.2.4. Microtubule stabilizers

Since microtubule disruption occurs in several models of AD and is associated with tau dysfunction, microtubule stabilizers have been assayed in preclinical and clinical trials for AD [88]. The anti-mitotic drug paclitaxel prevents A β -induced toxicity in cell culture [89], as well as axonal transport deficits and behavioral impairments in tau transgenic mice [90]. Unfortunately, paclitaxel is a P-glycoprotein substrate and it has very low capacity to cross the BBB, making it unsuitable for the treatment of human tauopathies. Epothilone D, which has better BBB permeability, improves microtubule density and cognition in tau transgenic mice [91]. Finally, the peptide NAP stabilizes microtubules and reduces tau hyper-phosphorylation [92]. NAP can be administered intra-nasally and has shown promising results in a phase II clinical trial [93].

2.3. Oxidative stress

Several pieces of evidence demonstrate that oxidative stress precedes other hallmarks of the neurodegenerative process in human brains and animal models of AD, including A β deposition, NFT formation, and metabolic dysfunction and cognitive decline. It plays a functional role in the pathogenesis of the disease [94-100]. These findings sustain the possibility of using anti-oxidants in the prevention and treatment of Alzheimer [101, 102]. Several studies in AD transgenic mouse models support the potential beneficial effect of antioxidant compounds as preventive drugs.

2.3.1. Naturally-occurring anti-oxidants

Several nutritional antioxidants such as resveratrol, curcumin, epigallocatechin gallate, L-acetyl-carnitine, RRR- α -tocopherol (vitamin E) and ascorbic acid (vitamin C) have been tested to counteract oxidative stress-induced brain damage in AD.

- *Resveratrol* is a polyphenolic compound found in grapes, berries and peanuts with well known anti-oxidant, anti-cancer, anti-inflammatory and estrogenic activities. *In vitro* and animal experiments reveal that resveratrol protects against A β toxicity by promoting the non-amyloidogenic cleavage of APP, thus enhancing the clearance of A β peptides by promoting their degradation through the ubiquitin-proteasome system, as well as reducing neuronal damage by decreasing the expression of inducible nitric oxide synthase (iNOS) and cyclooxygenase 2 (COX-2), and the pro-apoptotic factors Bax and c-Jun N-terminal kinase (JNK). Moreover, the capacity of resveratrol to induce the over-expression of sirtuins, proteins having a role in cell survival, probably contributes to its neuroprotective effect [103, 104].

- *Curcumin* is a polyphenolic compound present in the rhizome of *Curcuma longa*, commonly used as a spice to color and flavor food, which has anti-inflammatory, anti-carcinogenic and anti-infectious properties. The first evidence of a protective role of curcumin in AD was derived from epidemiological studies based on populations subjected to a curcumin-enriched diet. Additionally, *in vitro* studies have shown that curcumin protects neurons from A β toxicity whereas the use of AD transgenic mouse models show that curcumin suppresses inflammation and oxidative damage as well as accelerating the A β rate of clearance and inhibiting A β aggregation. Curcumin is considered a bi-functional antioxidant because it is a direct scavenger of oxidants as well as a long-lasting protector promoting the expression of cytoprotective proteins through the induction of Nrf2-dependent genes [105, 106]. Regrettably, no significant improvement in cognitive function between placebo and curcumin-treated groups has been observed in the only two clinical trials carried out until now [107].
- *Epigallocatechin gallate (EGCG)* is a polyphenolic flavonoid encountered in green tea. Human epidemiological and animal data suggest that tea may decrease the incidence of dementia and AD. EGCG has been demonstrated to exert its neuroprotective activity by reducing A β production and inflammation, and increasing mitochondrial stabilization, iron chelation and ROS scavenging [108]. However, to date no clinical trials have been performed to verify whether EGCG neuroprotective/neurorestorative actions can be successfully translated into human beings.
- *Acetyl-L-Carnitine (ALC)* is a natural compound found in red meat whose biological role is to facilitate the transport of fatty acids to the mitochondria. Thus, the main mechanism of action of ALC is the improvement of mitochondrial respiration, which allows the neurons to produce the necessary ATP to maintain normal membrane potential. Yet ALC is neuroprotective through a variety of additional effects, including an increase in protein kinase C activity and modulation of synaptic plasticity by counteracting the loss of NMDA receptors in the neuronal membrane and by increasing the production of neurotrophins [105]. Moreover, ALC reduces A β toxicity in primary cortical neuronal cultures by increasing both heme-oxygenase 1 (HO-1) and heat-shock protein 70 (Hsp70) expression, probably through transcription factor Nrf2. In two clinical studies, ALC administered for one year significantly reduced cognitive decline in early-onset AD patients [109, 110] thus sustaining the potential use of ALC in AD prevention and treatment at early stages.
- *RRR- α -tocopherol (Vitamin E)* is probably the most important lipid-soluble natural antioxidant in mammalian cells. Most vegetable oils, nuts and some fruits are important dietary sources of vitamin E. The interest in evaluating its potential beneficial properties in AD is also sustained by its known ability to cross the BBB and to accumulate in the central nervous system. Deficiency in the α -tocopherol transfer protein mediating vitamin E activity induces an increase in brain lipid peroxidation, earlier and more severe cognitive dysfunction, and increased A β deposits in the brain of Tg2576 mice; this phenotype was ameliorated with vitamin E supplementation [111]. However, although epidemiological studies have demonstrated that increasing the intake of fruit and vegetables rich in vitamins prevents or retards the onset of AD, clinical trials for vitamin E treatment have revealed paradoxical

results: whereas vitamin E supplementation partially prevents the memory loss associated with the progression of the disease in some cases, the same treatment was detrimental in others [112].

- *Ascorbic acid (Vitamin C)* is an essential nutrient since it acts as a cofactor in elemental enzymatic reactions, but in contrast to most of organisms, humans are not able to synthesize ascorbic acid. The main dietary source of vitamin C is fresh fruit and vegetables. The main interest in vitamin C for the treatment of neurodegenerative processes is related to its potent anti-oxidant properties. Some studies have revealed that vitamin C supplementation reduces oxidative stress, and mitigates A β oligomer formation and behavioral decline, but it did not decrease plaque deposition in AD mouse models [113, 114]. Despite epidemiological studies reporting reduced prevalence and incidence of AD in consumers of vitamin supplements [115], meta-analyses revealed the risks of chronic consumption of high doses of vitamin C thus discouraging its routine use in AD. [116]
- *Egb76* is a standardized *Ginkgo biloba* extract already approved in some countries as symptomatic treatment for dementia although the evidence for its effectiveness remains inconclusive [117]. However, Egb761 has anti-oxidant properties, inhibits A β oligomerization *in vitro*, reduces impaired memory and learning capacities and enhances hippocampal neurogenesis in AD transgenic mice [118]. For these reasons, *Ginkgo biloba* extract is currently under evaluation as a preventive drug in AD.

In spite of the experimental evidence of beneficial effects of natural anti-oxidants in cultured cells and transgenic models, clinical studies have demonstrated only minimal effect in humans probably due to the bioavailability and pharmacokinetics of these substances [102, 105]. What's more, a slight acceleration in cognitive decline has been observed in patients treated for 16 weeks with a cocktail of natural antioxidants [119].

2.3.2. Mitochondrial antioxidants

In contrast to other antioxidants, those designed to target the free radical damage to mitochondria provide greater therapeutic potential.

- *Lipoic acid (LA)* is a naturally-occurring precursor of an essential cofactor of many mitochondrial enzymes, including pyruvate dehydrogenase and alpha-ketoglutarate dehydrogenase, which is found in almost all foods. LA has been shown to present a variety of properties that can interfere with pathogenic processes of AD. LA increases ACh production, stimulates glucose uptake, protects against A β toxicity, chelates redox-active transition metals, scavenges reactive oxygen species (ROS) and induces anti-oxidant protective enzymes probably through the activation of the transcription factor Nrf2. Via the same mechanisms, down-regulation of redox-sensitive inflammatory processes is also achieved [120]. Data from cell culture and animal models suggest that LA can be combined with other dietary anti-oxidants to synergistically decrease oxidative stress, inflammation, A β levels, and thus provide a combined benefit in the treatment of AD. However, clinical benefits after LA administration were quite small in patients with mild or moderate dementia [121].

- *N-acetyl-cysteine* (NAC) is a precursor of glutathione (GSH), the most abundant endogenous anti-oxidant. NAC acts itself as an anti-oxidant by directly interacting with free radicals, as well as by increasing GSH levels. NAC protects against A β -induced cognitive deficits by decreasing the associated oxidative stress and related neuroinflammation, but also by activating anti-apoptotic signaling pathways in neuronal cultures [122]. Late-stage AD patients supplemented with NAC over a period of six months showed significantly improved performance in some cognitive tasks, although levels of oxidative stress in peripheral blood did not differ significantly from untreated patients [123].
- *Coenzyme Q₁₀* (CoQ₁₀) is a small electron-carrier of the respiratory chain with anti-oxidant properties due to its role in carrying high-energy electrons from complex I to complex II during oxidative phosphorylation. CoQ₁₀ and its analogues, idebenone and mitoquinone (or MitoQ), have been widely used for the treatment of mitochondrial disorders, as well as for the treatment of Friedreich's ataxia, and they are also being tested in other neurodegenerative disorders such as amyotrophic lateral sclerosis, and Huntington's, Parkinson's and Alzheimer's diseases [124]. CoQ₁₀ reduces oxidative stress damage and A β plaque burden, and ameliorates behavioral performance in mouse models of AD [125, 126]. However, CoQ₁₀ presents two major weaknesses. First, the function of the enzyme is entirely dependent on the electron transport chain (ETC) which is usually damaged in AD mitochondria. Second, CoQ₁₀ does not efficiently cross the BBB when administered systemically, being unable to directly protect neurons from damage. Consequently, CoQ₁₀ derivatives such as MitoQ, which is a more soluble compound able to penetrate the BBB and that does not depend on ETC, are seen to offer more promising results [127].

2.4. Inflammation

There is a general consensus that neuroinflammation is a prominent feature in AD with activated microglia being one of the main manifestations. Neuroinflammation is a complex process that has both beneficial effects, in terms of maintaining brain homeostasis after various kinds of insults, and detrimental effects when sustained chronically [128]. This latter situation is what occurs in AD, in which neuroinflammation is driven by different mechanisms including A β production and plaque formation, tau pathology, oxidative stress, and autocrine and paracrine release of cytokines and other inflammatory molecules which contribute to a feed-forward spiral favoring the self-propagation of neuroinflammation.

Early epidemiological studies suggesting that long-term use of antiinflammatories might reduce the risk for developing AD [129] prompted several studies designed to evaluate the preventive properties of non-steroid anti-inflammatory drugs (NSAIDs). The main NSAID mechanism of action is to inhibit the activity of cyclooxygenase-1 and -2 (COX-1 and COX-2) which are the enzymes responsible of the production of prostaglandins and other inflammatory agents [130]. The administration of the NSAID ibuprofen at early stages of the pathological process resulted in the reduction of the A β burden, dystrophic neurites and activated microglia in at least three different AD transgenic models [131-134]. Another study indicated that ibuprofen was effective even in older mice once lesions are well established [135]. Other NSAIDs such as indomethacin and nimuselide exhibit milder effects compared to ibuprofen

in the Tg2576 mice [136, 137]. In contrast, the selective COX-2 inhibitor celecoxib failed to reduce the inflammatory burden and, even worse, increased the $A\beta_{42}$ levels when administered to young Tg2576 mice [138].

In spite of the promising results in animal models and the data from retrospective human epidemiological studies identifying long-term use of NSAIDs as being protective against AD, prospective clinical trials have not confirmed the efficiency of this group of drugs in the amelioration of symptoms and in the progression of AD [139].

Other anti-inflammatory agents such as trifusal have been shown to be beneficial in certain AD transgenic mice models [140].

2.5. Energetic failure: Metabolic deficiency and mitochondrial impairment

Several findings indicate that brain glucose hypometabolism, deficient bioenergetics and mitochondrial dysfunction precede clinical symptoms in AD [1, 141-143]. The energetic failure observed even in the prodromal phase of the Alzheimer process is thought to be produced by the combination of mitochondria dysfunction, alteration of energy metabolism at pore-mitochondrial level, and increase in energetic demands of altered nerve cells. Thus, strategies to improve brain energy supply and to preserve mitochondrial functions becomes relevant in the prevention of progression to disease states [1, 144-146].

2.5.1. Metabolic deficiency

The primary fuel for the brain under normal conditions is glucose, whereas the energetic contribution made by fatty acids is minor. Therefore, facilitation of energy metabolism and energy availability has been assayed in animal models and AD by facilitating glucose metabolism and shifting towards the use of alternative fuels.

- *Targeting reduced glucose metabolism:* Reduction in the utilization of glucose in AD [147] can be due to several causes including deficient insulin signaling, impairment in glucose transport mechanisms and dysfunction in glycolysis. Preclinical studies in animal models of AD have revealed some beneficial effects of anti-diabetic treatments. Thus, the use of the insulin sensitizer rosiglitazone, an activator of peroxisome-proliferator-activated receptor gamma (PPAR γ) receptor, resulted in the rescue of behavioral deficits and insulin responsiveness in Tg2576 mice [148, 149]. Similarly, exendin-4, an antidiabetic agent that stimulates the insulin signaling pathway through activation of glucagon-like peptide -1 (GLP1) receptors, shows beneficial effects in AD, and reduces brain soluble $A\beta$ levels, amyloid plaque burden, and cognitive impairment in treated APP/PS1 transgenic mice [150, 151]. Therefore, it seems that the positive effects of targeting insulin signaling in AD are related to the role played by insulin receptor in memory formation, inflammation and $A\beta$ neuro-protective effects rather than to the facilitation of glucose transport into the brain [149, 150]. This hypothesis seems also to be supported by a recent study revealing that insulin did not ameliorate the disruption of energetic homeostasis induced by $A\beta$ oligomers in cultured neurons [152]. In the end, clinical trials designed to test whether PPAR γ agonists could be beneficial in AD patients provided negative results [153].

- *Shift to alternative energy source:* Under metabolically challenging conditions neurons can utilize acetyl-CoA generated from ketone body metabolism, produced distally in the liver or locally in the brain by glial cells. In this way, ketone bodies can bypass defects in glucose metabolism and enter the tricarboxylic acid cycle in the mitochondria of neurons as a source of ATP. The use of ketogenic diets reduces A β 40 and A β 42 levels in young AD transgenic mice [154] and enhances mitochondrial bioenergetic capacity, reducing A β generation and increasing mechanisms of A β clearance in a mouse model of AD [155]. The ketogenic compound AC-1202 administered in patients with AD has shown a significant improvement in some cognitive parameters more notable in individuals APOE4(-) [156]. Another possible alternative source of ATP is creatine. Preliminary studies have shown that creatine has protective effects against A β *in vitro* [157] and against injury *in vivo* by maintaining ATP levels and mitochondrial function [158], suggesting a potential therapeutic effect of creatine supplementation in AD.

2.5.2. Mitochondrial dysfunction

In addition to the already discussed antioxidant compounds, other potential drugs targeting mitochondrial dysfunction in AD are available. Several findings point towards a role for A β toxicity in the mitochondrial dysfunction found in AD.

The progressive A β accumulation in mitochondria is associated with diminished enzymatic activity of respiratory chain complexes (III and IV) and reduction in the rate of oxygen consumption, contributing to cellular dysfunction in AD [159]. A β in mitochondria binds to A β -binding alcohol dehydrogenase (ABAD) to block ABAD activity, increasing the production of ROS, reducing the mitochondrial membrane potential and the activity of the respiratory chain complex IV, and ultimately leading to a decrease in ATP levels [160]. In fact, double transgenic mice over-expressing mutated APP and ABAD exhibit exaggerated oxidative stress and memory impairment [160]. Therefore, compounds designed to block A β -ABAD interactions are considered putative therapeutic agents in AD. In line with this hypothesis, a recent study has shown that AG18051, a novel small ABAD-specific compound inhibitor, partially blocked the A β -ABAD interaction, prevented the A β 42-induced down-regulation of ABAD activity and protected cultured neurons against A β 42 toxicity by reducing A β 42-induced impairment of mitochondrial function and oxidative stress [161]. Furthermore, the introduction of an ABAD-decoy peptide into transgenic APP mice reduces A β -ABAD interaction and protects against A β -mediated mitochondrial toxicity [162].

Another line of research suggests that drugs that activate ATP-sensitive potassium (K_{ATP}) channels present in the mitochondrial inner membrane exhibit therapeutic potential in the treatment of AD, as K_{ATP} channels are activated when cellular ATP levels fall below a critical value thereby reducing excitability so as to maintain ion homeostasis and preserve ATP levels [163]. Long-term administration of diazoxide improves neuronal bioenergetics, suppresses A β and tau pathologies, and ameliorates memory deficits in the 3xTgAD mouse model of AD [164].

Finally, another potential drug in the treatment of AD that acts on mitochondrial pathways is latrepirdine, also known as Dimebon™ [165]. Latrepirdine reduces A β -induced mitochondrial impairment and increases the threshold of inductors to mitochondrial pore transition, making mitochondria more resistant to lipid peroxidation and increasing neuronal survival *in vitro* [166-168]. The interest in developing latrepirdine as a drug against AD is also supported by its multiple potential mechanism of action apart from mitochondrial effects, including anti-excitotoxic agent, inhibitor of AChE, channel-regulation and neurotrophic stimulator [165]. A preliminary clinical trial revealed that latrepirdine was safe and well tolerated, and significantly improved the clinical course of the disease in patients with mild-to-moderate AD [169]. Current phase III clinical trials are already being conducted [165].

2.6. Neurotransmitter dysfunction

The alteration of several transmitter systems is assumed to trigger both cognitive and neuropsychiatric symptoms in AD. A number of *post-mortem* studies indicate that neurotransmitter systems are not uniformly affected in AD. Thus, while cholinergic, serotonergic and glutamatergic deficits are present at relatively early stages of AD, dopaminergic and GABAergic systems appear to be affected later [170].

2.6.1. Cholinergic system

A large body of evidence has shown that basal forebrain cholinergic neurons are vulnerable to AD leading to a progressive cholinergic denervation of the cerebral neocortex [171, 172]. Taking into account the involvement of this system in the cognitive processing of memory and attention, the current attempts in cholinergic therapy in AD are justified [172, 173]. The various cholinergic strategies include the use of ACh precursors, inhibitors of cholinesterases, muscarinic and nicotinic agonists, and ACh releasers, in addition to the rescue of cholinergic function by nerve growth factor (NGF) which is reviewed in section 2.8.

- *ACh precursor.* Animal studies report that choline and lecithin increased the production of brain ACh which argues for their use in the treatment of cholinergic deficits in AD. However, evidence from randomized trials did not sustain this hypothesis [174].
- *Cholinesterase inhibitors (ChEIs).* Physostigmine, tacrine and derivatives donepezil, galantamine and rivastigmine have been tested in AD patients during the last three decades. Their therapeutic properties have been profusely reviewed [172, 175-177] and for this reason a detailed revision of ChEIs is beyond the scope of this chapter. Nevertheless, it is worth briefly indicating additional mechanisms of action of these compounds beyond inhibition of cholinesterases, including increase of nicotinic ACh receptor expression, facilitation of APP processing and attenuation of A β -induced toxicity [173, 178]. In spite of the fact that their efficacy has been proved in several clinical trials, only approximately 50% of patients respond positively. This limited effect of ChEIs on cognitive decline, together with the occurrence of undesirable side-effects such as diarrhea, nausea, insomnia, fatigue and loss of appetite, reduces the therapeutic capacities of ChEIs.

- *Muscarinic receptor 1 agonist.* The cholinergic deficiency in AD appears to be mainly pre-synaptic. Thus, the pharmacological stimulation of the post-synaptic M1 muscarinic receptors, which are preserved until late stages of AD, may balance the degeneration of pre-synaptic cholinergic terminals unable to properly synthesize and release ACh [173]. In fact, the selective M1 agonist AF267B reduces memory impairment, A β 42 levels, and tau hyperphosphorylation in AD triple transgenic mice [179], corroborating some early studies *in vitro* [180, 181]. This selective agonist is currently under clinical evaluation for safety and tolerability and a number of other M1 agonists are being investigated [173].
- *Nicotinic agonists.* Preclinical studies in animal models and some pilot studies in AD have shown that the activation of pre-synaptic nicotinic ACh receptors may reduce cognitive impairment by increasing ACh release and may have beneficial effects on A β metabolism [182, 183]. Thus, chronic nicotine treatment results in a significant reduction in plaque burden and in cortical A β concentrations in Tg2576/PS1-A246E mice [184]. However, nicotine exacerbates tau pathology in 3xTg-AD mice [185]. These apparently contradictory results may be due to the presence of several subtypes of nicotinic receptors, the activation of which may have disparate effects in AD. Therefore, more specific nicotine agonists are needed to act exclusively on determinate subtypes of nicotinic receptor [186]. In this line, α 7 nAChR gene delivery into mouse hippocampal neurons leads to functional receptor expression and improves spatial memory-related performance and hyperphosphorylation of tau [187]. Regarding α 4 β 2 nicotinic receptor, the selective agonist cytisine inhibits A β cytotoxicity in cortical neurons [188].
- *ACh releasers.* Facilitation of ACh release can be achieved with depolarizing agents of the cholinergic neurons acting via potassium-channel blockade as happens with linopirdine and analogues [189] or by the blockade of the pre-synaptic inhibitory M2 muscarinic receptor via specific antagonists [190, 191]. However, clinical trials using linopirdine did not demonstrate effectiveness in improving cognitive function [192]. On the other hand, certain selective M2 antagonists, such as SCH-57790 and SC-72788, restore memory impairments in animal models that mimic to some extent the cholinergic failure in AD [193]. It must be kept in mind that the potential benefit of M2 antagonists is limited because of the progressive pre-synaptic cholinergic degeneration in AD and because of the possible side-effects derived from the blockade of peripheral M2 receptors including cardiac M2 receptors.

2.6.2. Glutamatergic system

Low concentrations of A β oligomers are able to activate certain glutamate receptors including NMDA receptors. The activation of NMDA receptors may increase glutamate activity, raise intracellular Ca²⁺ concentration and promote excitotoxicity and neuronal damage [194, 195]. Another process contributing to the excessive glutamate activity in AD is the impairment of glial cells to remove glutamate from the synaptic cleft possibly due to the action of free radicals on the glutamate transporter 1 (GLT-1) [196]. Glutamatergic activation, in turn, may disrupt synaptic plasticity promoting long term depression (LTD) and inhibiting long term potentiation (LTP) of 2-amino-3-(5-methyl-3-oxo-1,2-oxazol-4-yl)propanoic acid (AMPA) receptor-mediated synaptic transmission [197]. The associated persistent reduction in the number of

functional synaptic AMPA receptors reduces fast excitatory transmission and eventually triggers spine retraction and synaptic loss [198]. Moreover, glutamate receptors are not only involved in the process of A β -mediated synaptic dysfunction but also play important roles in A β production [199, 200].

Based on these observations, several studies have been designed in an attempt to correct glutamatergic dysfunction in AD, including the modulation of both AMPA and NMDA receptors [201]. First attempts were carried out with AMPAKines [202], which are drugs that prolong the action of glutamate on AMPA receptors by increasing their sensitivity. Interestingly, AMPAKines proved effective in restoring cognitive deficits in aging rats [203, 204]. These compounds were tested in AD patients [205]. The modulation of the NMDA receptor was assessed via the glycine co-agonist site in rats with disrupted glutamatergic temporal systems resulting in improved learning and memory [206]. Preliminary clinical studies suggested some promising effects in AD [207] but full-scale trials have not yet been initiated.

The most relevant glutamatergic strategy against AD is the non-competitive NMDA antagonist memantine [201, 208], which has succeeded in clinical trials in moderate and severe AD as reviewed in detail elsewhere [209, 210]. Several studies performed in animal models of AD corroborate the beneficial properties of memantine as a symptomatic and neuroprotective treatment in AD [211-215]. Nevertheless, memantine has no benefits in cases with mild AD [216] suggesting that this drug is not a good choice for preventing the progression to disease states.

2.6.3. Serotonergic system

Loss of serotonergic nerve terminals in AD was described several years ago [217, 218]. Although the suggested serotonergic dysfunction was initially related almost exclusively with the neuropsychiatric symptoms of AD, including anxiety, irritability, fear and depression, recent studies have demonstrated that serotonin signaling also plays an important role in cognition and in the development of A β and tau pathologies [219].

Antidepressant compounds, acting through serotonin signaling, result in cognitive improvements and reduce the levels of A β and tau pathology in animal models of AD [220, 221]. Similar compounds reduce amyloid burden in humans [221]. Additional serotonergic compounds that are currently being investigated in AD are 5-hydroxytryptamine (5-HT or serotonin) receptors: 5-HT₁ and 5-HT₆ antagonists, and 5-HT₄ agonists. The 5-HT_{1A} antagonist lecozotan (SRA-333) enhances cognition in primates and is now being tested in AD [222-224]. The pro-cognitive effects of 5-HT_{1A} antagonists are probably due to the facilitation of glutamatergic and cholinergic transmission after reduction of the inhibitory effects of serotonin. Similarly, 5-HT₆ antagonists improve cognitive performance in animal models and human beings by modulating multiple neurotransmitter systems [225]. These properties mark 5-HT₆ antagonists as potential symptomatic drugs in AD. In addition, 5-HT₄ receptor agonists are neuroprotective, modulating the production of A β , and have the property of ameliorating cognitive deficits [226, 227].

2.7. Synaptic dysfunction

Synaptic dysfunction and failure are processes that occur early in the Alzheimer process and progress during the course of the disease from an initially reversible functionally-responsive stage of down-regulated synaptic function to stages irreversibly associated with degeneration.

These alterations are manifested early as impaired metabotropic glutamate receptor/phospholipase C signaling pathway [230] and up-regulation of adenosine receptors in the frontal cortex in AD [231].

The initial reversible stages are important targets for protective treatments to slow progression and preserve cognitive and functional abilities [232, 233]. *In vivo* and *in vitro* studies have demonstrated that high levels of A β impair structural and functional plasticity of synapses by affecting the balance between excitation and inhibition and contributing to the destabilization of neuronal networks, eventually causing synaptic loss [234]. Two main designs have been proposed to antagonize synaptic plasticity-disrupting actions of A β oligomers in preclinical AD: maintenance of the structure and fluidity of the lipid membranes forming the synaptic buttons, and stimulation of synaptic plasticity by neurotrophic factors.

Minor changes in the fluidity of phospholipidic membranes might have an important impact on the function of synapses by influencing neurotransmitter receptor activity. In fact, AD brains exhibit altered lipid composition of lipid rafts, key membrane microdomains that facilitate the transfer of substrates and protein-protein and lipid-protein interactions, as a result of the abnormally low levels of n-3 long-chain polyunsaturated fatty acids, mainly docosahexaenoic acid (DHA), increasing viscosity and energy consumption and contributing to synaptic dysfunction [142, 235]. Abnormal lipid raft composition may also modify the activity of key enzymes that modulate the cleavage of APP to form toxic A β . Thus, the preservation of adequate membrane composition has become an alternative way to prevent the deleterious effect of A β at the synapses. DHA is a major lipid constituent of synaptic end-sites and its delivery is a prerequisite for the conversion of nerve growth cones to mature synapses [236]. Numerous epidemiological studies have highlighted the beneficial influence of DHA on the preservation of synaptic function and memory capacity in aged individuals or after A β exposure, whereas DHA deficiency is presented as a risk factor for AD [237]. Moreover, a number of studies have reported the beneficial effects of dietary DHA supplementation on cognition and synaptic integrity in various AD models [238]. According to this evidence, DHA, which can be synthesized or obtained directly from fish oil, appear to be one of the most valuable diet ingredients whose neuroprotective properties contribute to preventing AD.

Cytidine 5'-diphosphocholine, CDP-choline, or citicoline is an essential intermediate in the biosynthetic pathway of structural phospholipids in cell membranes, particularly phosphatidylcholine. Chronic administration has been beneficial in patients with mild cognitive impairment [239].

Another emerging potential line to preserve synaptic function is the targeting of scaffolding proteins that modulate neurotransmitter receptor activity at the synapses. Scaffolding proteins stabilize post-synaptic receptors at the spines in close proximity to their intracellular signaling

proteins, phosphatases and kinases, thereby facilitating signal-transduction cascades. Evidence from *in vitro* cell and animal models of AD indicates that reductions in the post-synaptic density membrane-associated guanylate kinase (PSD-MAGUK) proteins are linked to synaptic dysfunction that might trigger plastic changes at early stages of the Alzheimer process [240]. However, specific molecules that affect interactions between scaffolding proteins and neurotransmitter receptors are still in development and further research is necessary to evaluate their potential benefit in AD.

2.8. Neurotrophic factors

Neurotrophins represent a family of proteins that play a pivotal role in the mechanisms underlying neuronal survival, differentiation, modulation of dendritic branching and dendritic spine morphology as well as synaptic plasticity and apoptosis [241]. All the members of the neurotrophin family, including NGF, brain-derived neurotrophic factor (BDNF) and neurotrophins 3 to 7, transduce their biological effects by interacting with two types of cell surface receptors, the tyrosine kinase receptor (Trk) and the p75 pan-neurotrophin receptor (p75^{NTR}) [241]. Other growth factor families also related to synaptic plasticity include the cytokine family of growth factors, the transforming growth factor- β (TGF β) family, the fibroblast growth factor family and the insulin-like growth factor family. Evidence accumulated during recent years suggests that targeting neurotrophic factor signaling can retard nerve cell degeneration and to some extent preserve synaptic function. The most studied neurotrophic factors in AD are NGF, BDNF and TGF β 1.

- **NGF:** Mature basal forebrain cholinergic neurons are highly dependent on the availability of NGF for the maintenance of their biochemical and morphological phenotype, and for survival after lesions or variegated insults [242, 243]. For this reason, exploitation of NGF activity on cholinergic neurons may provide an attractive therapeutic option for preventing cholinergic cell degeneration in AD. Levels of proNGF, the precursor form of NGF, are highly elevated in AD brains and animal models, a feature that may be associated with a reduced conversion to NGF and augmented degradation of mature NGF. These combined effects have been interpreted as causative of cholinergic atrophy in AD [244]. A role for A β peptide in the induction of such NGF altered metabolism has been described [245]. Minocycline, a second-generation tetracycline antibiotic known to potentiate NGF activity, is able to normalize proNGF levels and to reverse the increased activity of the NGF-degrading enzyme matrix metalloproteinase 9, as well as to increase the expression of iNOS and microglial activation, leading to improved cognitive behavior in a transgenic mouse model of AD [245]. Yet a disturbing finding is the demonstration of AD proNGF when compared to proNGF of control individuals [246-248]. Whether this abnormal form of AD-related proNGF has any impact on the pathogenesis of AD needs further investigation. Another putative therapy is the use NGF, but NGF does not readily cross the BBB and requires intra-cerebroventricular infusion to reach targeted brain areas. Pilot clinical trials were discontinued because of the side-effects of NGF infusions [249]. Therefore, the development of NGF therapy is constrained by the need to achieve adequate concentrations in the relevant brain areas with susceptible target neurons while preventing unwanted

adverse effects in non-target regions or cells. Alternative strategies that are currently under development include gene therapy and nasal delivery of recombinant forms of NGF, the use of small molecules with NGF agonist activity, NGF synthesis inducers, NGF processing modulators, and proNGF antagonists [250].

- *BDNF*: This neurotrophin is normally produced in the cerebral cortex with high levels in the entorhinal cortex and hippocampus in adulthood [241]. BDNF levels are reduced in the cerebral cortex and hippocampus in AD [251-254]. Several studies have shown beneficial effects of BDNF in animal models of AD [255]. For instance, sustained BDNF gene delivery using viral vectors after disease onset resulted in elevated BDNF levels in the entorhinal cortex and hippocampus which were associated with improvement in learning and memory, and with restoration of most genes altered as a result of mutant APP expression in that specific transgenic mice model [256]. Similar results were obtained in a different mouse model of AD, and in aged rats and primates by using distinct BDNF delivery systems [256, 257]. It is worth pointing out that BDNF did not change β -amyloid plaque density in any case suggesting that the therapeutic effects of BDNF occur independently of direct action on APP processing. However, the multiple variegated effects of BDNF on neuronal function also raise the hypothetical possibility that unintended adverse effects of BDNF may limit its clinical efficacy in AD [256]. An additional point must be considered; BDNF signaling pathway is also altered in AD as TrkB expression is reduced and truncated TrkB is highly expressed in astrocytes at least in advanced stages of the disease [251]. Therefore, regarding BDNF function in AD, there is not only an alteration in the expression of BDNF but also an impaired downstream pathway that may corrupt the signal of the trophic factor acting on inappropriate receptors. Preliminary clinical trials are currently in progress to evaluate the safety and efficacy of BDNF.
- *TGF β 1*: Astrocytes and microglia are the major sources of TGF- β 1 in the injured brain [258, 259]. Impaired TGF- β 1 signaling has been demonstrated in AD brain, particularly at the early phase of the disease; this is associated with A β pathology and neurofibrillary tangle formation in animal models [260]. Reduced TGF- β 1 seems to induce microglial activation [259] and ectopic cell-cycle re-activation in neurons [261]. Several drugs may induce TGF- β 1 release by glial cells, including estrogens [262], mGlu2/3 agonists [263], lithium [264], the antidepressant venlafaxine [265] and glatiramer, which is a synthetic amino acid co-polymer currently approved for the treatment of multiple sclerosis [266]. All of them have neuroprotective effects in different *in vitro* and *in vivo* models of AD pathology [260]. Additionally, small molecules with specific TGF- β 1-like activity are being developed as neuroprotectors [267].

A final point must be considered. A generalized sprouting is produced around β -amyloid deposits in senile plaques in both humans and in animal models [268-270]. The reasons for such sprouting are not well defined but amyloid species may play a trigger role. In any case, trophic factors might increase aberrant sprouting at the senile plaques through receptors expressed at these localizations.

2.9. Autophagy

Autophagy is a catabolic process occurring in all cell types in which the machinery of the lysosome degrades cellular components such as long-lived or damaged proteins and organelles. Thus, a failure of autophagy in neurons results in the accumulation of aggregate-prone proteins that might exacerbate neurodegenerative process [271, 272]. Autophagy is also implicated in the accumulation of altered mitochondria and polymorphous inclusions in the dystrophic neurites around amyloid plaques [273-278].

Indeed, autophagic dysfunction is implicated in the progression of Alzheimer from the earliest stage, when a defective lysosomal clearance of autophagic substrates and impaired autophagy initiation occurs and leads to massive buildup of incompletely digested substrates within dystrophic axons and dendrites [279]. The pharmacological induction of 'preserved' autophagy might enhance the clearance of intracytoplasmic aggregate-prone proteins and therefore ameliorate pathology [272]. Attempts to restore more normal lysosomal proteolysis and autophagy efficiency in mouse models of AD pathology have revealed promising therapeutic effects on neuronal function and cognitive performance, demonstrating the relevance of the failure of autophagy in the pathogenesis of AD, and the potential of autophagy modulation as a therapeutic strategy. Autophagy induction with the mTOR-inhibiting drug rapamycin in young mice resulted in a reduction in A β plaques, NFT and cognitive deficits in the adulthood in two different models of AD [280-283]. Interestingly, rapamycin did not alter any of those parameters when administered in old animals once the pathology was established, highlighting the importance of early treatment in the disease progression [282]. However, the kinase mTOR plays an important role in multiple signaling pathways apart from negatively regulating autophagy [284]. Therefore, rapamycin treatment is also a putative inducer of undesirable side-effects. Other drugs including lithium, sodium valproate and carbamazepine acting have been proved to induce autophagy through the inhibition of inositol monophosphatase in an mTOR-independent pathway [285]. These compounds reveal positive effects by reducing the accumulation and toxic effects of aggregation-prone proteins in cell models as well as by protecting against neurodegeneration in *in vivo* models of Huntington's disease [286]. Further research is needed to learn whether they can also be useful tools in the treatment of AD.

2.10. Multi-target treatments

Considering the multifactorial etiology of AD, and the numerous and complex pathological mechanisms involved in the progression of the disease, it is quite reasonable that treatments targeting a single causal or modifying factor may have limited benefits. Therefore, growing interest is focused on therapeutic agents with pleiotropic activity, which will be able to target, in parallel, several processes affected in AD [287, 288]. Several compounds already mentioned in the previous sections fulfill these properties, such as DHA which presents anti-inflammatory, anti-oxidant, neuroprotective and anti-tau phosphorylation properties apart from the modulation of synaptic membrane composition [289], and curcumin, which in addition to anti-oxidant properties also exhibits anti-inflammatory and A β - and tau-binding properties [106]. Similarly, rosiglitazone and dimebon are known to produce beneficial effects through insulin receptor signaling mod-

ulation and mitochondrial protection [153, 165]. Other multi-target potential treatments currently under development for AD are based on the use of the following compounds:

- *Caffeine*: This is one of the most consumed psychoactive drugs which mainly acts blocking adenosine receptors 1 and 2 [290, 291]. In addition, caffeine reduces amyloid burden in animal models of AD [292, 293]. Epidemiological studies in humans have also shown protection against cognitive decline [294-296].
- *Estrogen*: This steroid hormone is known to play an important role in neuronal survival, mitochondrial function, neuroinflammation and cognition, with important neuroprotective effects [297-299]. Some of the neuroprotective actions mediated by estrogens are related to the insulin-like growth factor-1 (IGF-1) signaling pathway [300]. Several studies in animal models of AD have revealed therapeutic properties of estrogen against the progression of the disease. For instance, the treatment of ovariectomized 3xTg-AD mice with estrogen resulted in prevention of the increased A β accumulation and worsening memory performance induced by the depletion of sex steroid hormones [301]. Clinical and epidemiological studies in AD support the beneficial effects of estrogens [302]. However, a critical factor for success in estrogen therapy for AD is the age at the initiation of the treatment; the efficacy of estrogens is greatest in younger women and in women who initiated the estrogen therapy at the time of menopause [303].
- *Cannabinoids*: The natural compounds derived from *Cannabis sativa* or synthetic compounds acting on endogenous cannabinoid system have emerged as potential agents against several neurodegenerative processes [305]. Cannabinoids offer a multi-faceted approach for the treatment of AD as the stimulation of the widely brain-expressed cannabinoid receptors provides neuroprotection against A β [305, 306] and reduces neuroinflammation [306-308] and tau phosphorylation [306, 309] in AD-like transgenic mice. In addition, cannabinoids support brain repair mechanisms by augmenting neurotrophin expression and enhancing neurogenesis [310]. Moreover, cannabinoids are able to reduce A β -dependent oxidative stress [311] and A β -mediated lysosomal destabilization related to apoptosis [312]. In addition, some cannabinoids are able to inhibit acetylcholinesterase activity [313]. It is worth stressing that molecular achievements of cannabinoids are accompanied by cognitive improvement and reduction of several degenerative markers in two different animal models of AD [306, 308]. Examination of the potential beneficial effects of chronic administration of low doses of cannabinoids with little psychotropic effect at early stages of the degenerative process in humans seems very promising.
- *Erythropoietin (EPO) and derivatives*: EPO is effective in neuroprotection against ischemia and traumatic brain injury [314]. In addition, animal studies reveal that EPO both reduces tau phosphorylation through modulation of PI3K/Akt-GSK-3 β pathway [315] and protects against A β -induced cell death through anti-oxidant mechanisms [316]. An additional characteristic of EPO that confers potential utility in AD is the specific effect on cognition: EPO enhances hippocampal LTP and memory by modulating plasticity, synaptic connectivity and activity of memory-related neuronal networks [317]. In spite of these benefits, chronic administration of EPO is problematic because of the concomitant excessive erythropoiesis. In this sense, some new derivatives of EPO that do not bind to the classical EPO

receptor (carbamylated EPO) or that have such a brief half-life in the circulation that they do not stimulate erythropoiesis (asialo EPO and neuro EPO) have demonstrated neuroprotective activities without the potential adverse effects on circulation associated with EPO [318]. Therefore, these new compounds are considered as potential treatments in AD.

- *Statins*: Evidence has accumulated that a high cholesterol level may increase the risk of developing AD and that the use of statins to treat hyper-cholesterolemia is useful in treating and preventing AD [319]. Statins reduce the production of cholesterol and isoprenoid intermediates. These isoprenoids modulate the turnover of small GTPase molecules that are essential in numerous cell-signaling pathways, including vesicular trafficking and inflammation [320]. Thus, statins reduce the production of A β by disrupting secretase enzyme function and by curbing neuroinflammation in experimental models of AD [321, 322].
- *Ladostigil* is a dual acetylcholine-butyrylcholinesterase and brain selective monoamine oxidase (MAO)-A and -B inhibitor *in vivo*. Interest in this compound in AD treatment research is sustained by the potential increase in brain cholinergic activity properties but also by the capacity of ladostigil to prevent gliosis and oxidative-nitrosative stress damage. Moreover, ladostigil has been demonstrated to possess potent anti-apoptotic and neuroprotective properties *in vitro* and in various neurodegenerative animal models including AD transgenic mice [323]. These neuroprotective activities involve regulation of APP processing, activation of protein kinase C and mitogen-activated protein kinase signaling pathways, inhibition of neuronal death markers, prevention of the fall in mitochondrial membrane potential, up-regulation of neurotrophic factors, and anti-oxidative activity.
- *Huperzine A* is an extract of the Chinese plant *Huperzia serrata*. Huperzine A is a selective potent inhibitor of AChE [324]. In addition, some studies have shown that huperzine A may shift APP metabolism towards the non-amyloidogenic α -secretase pathway [325]. In addition, huperzine A reduces glutamate-induced cytotoxicity by antagonizing cerebral NMDA receptors [326]. Finally, huperzine A reverses or attenuates cognitive deficits in some animal models of AD [325]. Large-scale, randomized, placebo-controlled trials are necessary to establish the role of huperzine A in the treatment of AD [327].
- *Phytochemicals* as curcumin, catechins and resveratrol beyond their antioxidant activity are also involved in anti-amyloidogenic, anti-inflammatory mechanisms and inhibitors of NF κ B [328-330].
- *Celastrrol* is another compound which appears to have multiple functions as anti-inflammatory, anti-oxidant and reductor of amyloid via BACE 1 [331, 332].

3. Concluding remarks

Main targets of therapeutic intervention at early stages of Alzheimer are summarized in Figure 1. Based on the presently available data several conclusions can be drawn. Combination therapies with drugs targeting different pathological factors or the use of multi-target compounds appear to be the most effective strategy in the treatment of the neurodegenerative

process in Alzheimer. Most potential experimental therapies exhibit the highest efficiency when applied during the pre-symptomatic phase of the disease. Therefore, it is essential to develop diagnostic tools to detect Alzheimer at early stages. Moreover, considering that Alzheimer, as a degenerative process not necessarily leading to dementia, affects a large percentage of individuals in the sixth decade of life, it would be wise to introduce habits and low-cost, safe treatments to prevent the progression of Alzheimer early in life, as occurs in arteriosclerosis, to transform AD into a chronic, incomplete and non-devastating disease thereby allowing for normal life in the elderly.

Figure 1. Schematic representation of the main cellular targets that are currently under development to prevent or retard the progression of Alzheimer to disease states. Most of the experimental approaches are designed to block or mitigate (red lines) pathological events occurring at the earliest stages, including abnormal A β and tau aggregation, chronic inflammatory responses, and oxidative stress damage. Other strategies (blue lines) aim at stimulating the metabolism to reduce Alzheimer's energetic failure as well as to promote intrinsic mechanisms that protect or repair cellular damage, including synaptic plasticity, preservation of the lipid membrane composition, and the promotion of damaged protein and organelle turnover. Therapeutic approaches based on the modulation of neurotransmission (green dashed lines) are designed to bypass deficient cholinergic neurotransmission whereas other compounds aim to block glutamatergic excitotoxicity. Considering the complex scenario of the Alzheimer neurodegenerative process, multi-target therapies applied at early stages of the disease appear to be the most effective strategy.

In addition to these general conclusions, several points deserve a particular comment. Recognition of the genotypic background, clinical and neuropathological subtypes and different pace of clinical manifestations is important to refine personalized treatments [333-335]. This includes modifications of the treatment as Alzheimer is not a mere accumulation of defects but rather a combination of deficiencies and plastic changes that imply shifts in molecular pathways with disease progression. Drugs and treatments

beneficial at first stages of the degenerative process may be harmful at advanced stages. Special effort must be put into practice to learn about the combination of drugs at which determinate time for every particular individual.

Acknowledgements

Parts of the work used in this review were supported by the project BESAD-P (Instituto Carlos III), Mutua Madrileña and Agrupación Mútua. We wish to thank T. Yohannan for editorial assistance.

Author details

Ester Aso and Isidre Ferrer*

*Address all correspondence to: 8082ifa@gmail.com

Institut de Neuropatologia, Hospital Universitari de Bellvitge, Universitat de Barcelona, CIBERNED, Spain

References

- [1] Ferrer I. Defining Alzheimer as a common age-related neurodegenerative process not inevitably leading to dementia. *Prog Neurobiol.* 2012; 97: 38-51.
- [2] Petot GJ, Friedland RP. Lipids, diet and Alzheimer disease: an extended summary. *J Neurol Sci.* 2004; 226: 31-3.
- [3] Luchsinger JAQ, Tang Noble JM, Scarmeas N. Diet and Alzheimer's disease. *Curr Neurol Neurosci Rep.* 2007; 7: 366-72.
- [4] Kovacs GG, Alafuzoff I, Al-Sarraj S, Arzberger T, Bogdanovic N, Capellari S, Ferrer I, Gelpi E, Kövari V, Kretschmar H, Nagy Z, Parchi P, Seilhean D, Soininen H, Troakes C, Budka H. Mixed brain pathologies in dementia: the brainNet Europe consortium experience. *Dementia.* 2008; 26: 343-50.
- [5] Ferrer I. Cognitive impairment of vascular origin: Neuropathology of cognitive impairment of vascular origin. *J Neurol Sci.* 2010; 299-339-49.
- [6] De Strooper B, Vassar R, Golde T. The secretases: enzymes with therapeutic potential in Alzheimer disease. *Nat Rev Neurol.* 2010; 6: 99-107
- [7] Ring S, Weyer SW, Kilian SB, Waldron E, Pietrzik CU, Filippov MA, Herms J, Buchholz C, Eckman CB, Korte M, Wolfer DP, Müller UC. The secreted beta-amyloid precursor

- protein ectodomain APPs alpha is sufficient to rescue the anatomical, behavioral, and electrophysiological abnormalities of APP-deficient mice. *J Neurosci.* 2007; 27: 7817-26.
- [8] Bandyopadhyay S, Goldstein LE, Lahiri DK, Rogers JT. Role of the APP non-amyloidogenic signaling pathway and targeting alpha-secretase as an alternative drug target for treatment of Alzheimer's disease. *Curr Med Chem.* 2007; 14: 2848-64.
- [9] Laird FM, Cai H, Savonenko AV, Farah MH, He K, Melnikova T, Wen H, Chiang HC, Xu G, Koliatsos VE, Borchelt DR, Price DL, Lee HK, Wong PC. BACE1, a major determinant of selective vulnerability of the brain to amyloid-beta amyloidogenesis, is essential for cognitive, emotional, and synaptic functions. *J Neurosci.* 2005; 25:11693-709.
- [10] McConlogue L, Buttini M, Anderson JP, Brigham EF, Chen KS, Freedman SB, Games D, Johnson-Wood K, Lee M, Zeller M, Liu W, Motter R, Sinha S. Partial reduction of BACE1 has dramatic effects on Alzheimer plaque and synaptic pathology in APP Transgenic Mice. *J Biol Chem.* 2007; 282: 26326-34.
- [11] Hills ID, Vacca JP. Progress toward a practical BACE-1 inhibitor. *Curr Opin Drug Discov Devel.* 2007; 10: 383-91.
- [12] Rajendran L, Schneider A, Schlechtingen G, Weidlich S, Ries J, Braxmeier T, Schwille P, Schulz JB, Schroeder C, Simons M, Jennings G, Knölker HJ, Simons K. Efficient inhibition of the Alzheimer's disease beta-secretase by membrane targeting. *Science.* 2008; 320: 520-3.
- [13] Serneels L, Van Biervliet J, Craessaerts K, Dejaegere T, Horr K, Van Houtvin T, Esselmann H, Paul S, Schfer MK, Berezovska O, Hyman BT, Sprangers B, Sciot R, Moons L, Jucker M, Yang Z, May PC, Karran E, Wiltfang J, D'Hooge R, De Strooper B. gamma-Secretase heterogeneity in the Aph1 subunit: relevance for Alzheimer's disease. *Science.* 2009; 324: 639-42.
- [14] Serneels L, Dejaegere T, Craessaerts K, Horr K, Jorissen E, Tousseyn T, Hbert S, Coolen M, Martens G, Zwijsen A, Annaert W, Hartmann D, De Strooper B. Differential contribution of the three Aph1 genes to gamma-secretase activity in vivo. *Proc Natl Acad Sci U S A.* 2005; 102:1719-24.
- [15] Dovey HF, John V, Anderson JP, Chen LZ, de Saint Andrieu P, Fang LY, Freedman SB, Folmer B, Goldbach E, Holsztynska EJ, Hu KL, Johnson-Wood KL, Kennedy SL, Kholodenko D, Knops JE, Latimer LH, Lee M, Liao Z, Lieberburg IM, Motter RN, Mutter LC, Nietz J, Quinn KP, Sacchi KL, Seubert PA, Shopp GM, Thorsett ED, Tung JS, Wu J, Yang S, Yin CT, Schenk DB, May PC, Altstiel LD, Bender MH, Boggs LN, Britton TC, Clemens JC, Czilli DL, Dieckman-McGinty DK, Droste JJ, Fuson KS, Gitter BD, Hyslop PA, Johnstone EM, Li WY, Little SP, Mabry TE, Miller FD, Audia JE. Functional gamma-secretase inhibitors reduce beta-amyloid peptide levels in brain. *J Neurochem.* 2001; 76: 173-81.
- [16] Abramowski D, Wiederhold KH, Furrer U, Jatton AL, Neuenschwander A, Runser MJ, Danner S, Reichwald J, Ammaturo D, Staab D, Stoeckli M, Rueeger H, Neumann U,

- Staufenbiel M. Dynamics of Abeta turnover and deposition in different beta-amyloid precursor protein transgenic mouse models following gamma-secretase inhibition. *J Pharmacol Exp Ther.* 2008; 327: 411-24.
- [17] Bateman RJ, Siemers ER, Mawuenyega KG, Wen G, Browning KR, Sigurdson WC, Yarasheski KE, Friedrich SW, Demattos RB, May PC, Paul SM, Holtzman DM. A gamma-secretase inhibitor decreases amyloid-beta production in the central nervous system. *Ann Neurol.* 2009; 66: 48-54.
- [18] Schor NF. What the halted phase III γ -secretase inhibitor trial may (or may not) be telling us. *Ann Neurol.* 2011; 69: 237-9.
- [19] Netzer WJ, Dou F, Cai D, Veach D, Jean S, Li Y, Bornmann WG, Clarkson B, Xu H, Greengard P. Gleevec inhibits beta-amyloid production but not Notch cleavage. *Proc Natl Acad Sci USA.* 2003; 100:12444-9.
- [20] Mayer SC, Kreft AF, Harrison B, Abou-Gharbia M, Antane M, Aschmies S, Atchison K, Chlenov M, Cole DC, Comery T, Diamantidis G, Ellingboe J, Fan K, Galante R, Gonzales C, Ho DM, Hoke ME, Hu Y, Huryn D, Jain U, Jin M, Kremer K, Kubrak D, Lin M, Lu P, Magolda R, Martone R, Moore W, Oganessian A, Pangalos MN, Porte A, Reinhart P, Resnick L, Riddell DR, Sonnenberg-Reines J, Stock JR, Sun SC, Wagner E, Wang T, Woller K, Xu Z, Zaleska MM, Zeldis J, Zhang M, Zhou H, Jacobsen JS. Discovery of begacestat, a Notch-1-sparing gamma-secretase inhibitor for the treatment of Alzheimer's disease. *J Med Chem.* 2008; 51: 7348-51.
- [21] Borgegard T, Jurus A, Olsson F, Rosqvist S, Sabirsh A, Rotticci D, Paulsen K, Klintenberg R, Yan H, Waldman M, Stromberg K, Nord J, Johansson J, Regner A, Parpal S, Malinowsky D, Radesater AC, Li T, Singh R, Eriksson H, Lundkvist J. First and second generation γ -secretase modulators (GSMs) modulate amyloid- β (A β) peptide production through different mechanisms. *J Biol Chem.* 2012; 287:11810-9.
- [22] D'Onofrio G, Panza F, Frisardi V, Solfrizzi V, Imbimbo BP, Paroni G, Cascavilla L, Seripa D, Pilotto A. Advances in the identification of γ -secretase inhibitors for the treatment of Alzheimer's disease. *Expert Opin Drug Discov.* 2012; 7: 19-37.
- [23] Shirotani K, Tsubuki S, Iwata N, Takaki Y, Harigaya W, Maruyama K, Kiryu-Seo S, Iwata H, Tomita T, Iwatsubo T, Saido TC. Neprilysin degrades both amyloid beta peptides 1-40 and 142 most rapidly and efficiently among thiorphan- and phosphoramidon-sensitive endopeptidases. *J Biol Chem.* 2001; 276: 21895-901.
- [24] Iwata N, Takaki Y, Fukami S, Tsubuki S, Saido TC. Region-specific reduction of A beta-degrading endopeptidase, neprilysin, in mouse hippocampus upon aging. *J Neurosci Res.* 2002; 70: 493-500.
- [25] Wang DS, Iwata N, Hama E, Saido TC, Dickson DW. Oxidized neprilysin in aging and Alzheimer's disease brains. *Biochem Biophys Res Commun.* 2003; 310: 236-41.

- [26] Marr RA, Rockenstein E, Mukherjee A, Kindy MS, Hersh LB, Gage FH, Verma IM, Masliah E. Neprilysin gene transfer reduces human amyloid pathology in transgenic mice. *J Neurosci.* 2003; 23: 1992-6.
- [27] Nalivaeva NN, Beckett C, Belyaev ND, Turner AJ. Are amyloid-degrading enzymes viable therapeutic targets in Alzheimer's disease? *J Neurochem.* 2012; 120 Suppl 1:167-85.
- [28] Leissring MA, Farris W, Chang AY, Walsh DM, Wu X, Sun X, Frosch MP, Selkoe DJ. Enhanced proteolysis of beta-amyloid in APP transgenic mice prevents plaque formation, secondary pathology, and premature death. *Neuron.* 2003; 40: 1087-93.
- [29] Qiu WQ, Folstein MF. Insulin, insulin-degrading enzyme and amyloid-beta peptide in Alzheimer's disease: review and hypothesis. *Neurobiol Aging.* 2006; 27: 190-8.
- [30] Miners JS, Barua N, Kehoe PG, Gill S, Love S. A β -degrading enzymes: potential for treatment of Alzheimer disease. *J Neuropathol Exp Neurol.* 2011; 70: 944-59.
- [31] Aisen PS, Gauthier S, Vellas B, Briand R, Saumier D, Laurin J, Garceau D. Alzhemed: a potential treatment for Alzheimer's disease. *Curr Alzheimer Res.* 2007; 4: 473-8.
- [32] Aisen PS, Gauthier S, Ferris SH, Saumier D, Haine D, Garceau D, Duong A, Suhy J, Oh J, Lau WC, Sampalis J. Tramiprosate in mild-to-moderate Alzheimer's disease _ a randomized, double-blind, placebo-controlled, multi-centre study (The Alphase study). *Arch Med Sci.* 2011; 7: 102-11.
- [33] Fenili D, Brown M, Rappaport R, McLaurin J. Properties of scyllo-inositol as a therapeutic treatment of AD-like pathology. *J Mol Med (Berl).* 2007; 85: 603-11.
- [34] Hawkes CA, Deng LH, Shaw JE, Nitz M, McLaurin J. Small molecule beta-amyloid inhibitors that stabilize protofibrillar structures in vitro improve cognition and pathology in a mouse model of Alzheimer's disease. *Eur J Neurosci.* 2010; 31: 203-13.
- [35] Salloway S, Sperling R, Keren R, Porsteinsson AP, van Dyck CH, Tariot PN, Gilman S, Arnold D, Abushakra S, Hernandez C, Crans G, Liang E, Quinn G, Bairu M, Pastrak A, Cedarbaum JM; ELND005-AD201 Investigators. A phase 2 randomized trial of ELND005, scyllo-inositol, in mild to moderate Alzheimer disease. *Neurology.* 2011; 77: 1253-62.
- [36] Adlard PA, Cherny RA, Finkelstein DI, Gautier E, Robb E, Cortes M, Volitakis I, Liu X, Smith JP, Perez K, Loughton K, Li QX, Charman SA, Nicolazzo JA, Wilkins S, Deleva K, Lynch T, Kok G, Ritchie CW, Tanzi RE, Cappai R, Masters CL, Barnham KJ, Bush AI. Rapid restoration of cognition in Alzheimer's transgenic mice with 8-hydroxy quinoline analogs is associated with decreased interstitial A β . *Neuron.* 2008 59: 43-55.
- [37] Faux NG, Ritchie CW, Gunn A, Rembach A, Tsatsanis A, Bedo J, Harrison J, Lannfelt L, Blennow K, Zetterberg H, Ingelsson M, Masters CL, Tanzi RE, Cummings JL, Herd

CM, Bush AI. PBT2 rapidly improves cognition in Alzheimer's Disease: additional phase II analyses. *J Alzheimers Dis.* 2010; 20: 509-16.

- [38] Klajnert B, Cladera J, Bryszewska M. Molecular interactions of dendrimers with amyloid peptides: pH dependence. *Biomacromolecules.* 2006; 7: 2186-91.
- [39] Klajnert B, Cortijo-Arellano M, Cladera J, Bryszewska M. Influence of dendrimer's structure on its activity against amyloid fibril formation. *Biochem Biophys Res Commun.* 2006; 345: 21-8.
- [40] Klementieva O, Benseny-Cases N, Gella A, Appelhans D, Voit B, Cladera J. Dense shell glycodendrimers as potential nontoxic anti-amyloidogenic agents in Alzheimer's disease. Amyloid-dendrimer aggregates morphology and cell toxicity. *Biomacromolecules.* 2011; 12: 3903-9.
- [41] Schenk D, Barbour R, Dunn W, Gordon G, Grajeda H, Guido T, Hu K, Huang J, Johnson-Wood K, Khan K, Kholodenko D, Lee M, Liao Z, Lieberburg I, Motter R, Mutter L, Soriano F, Shopp G, Vasquez N, Vandeventer C, Walker S, Wogulis M, Yednock T, Games D, Seubert P. Immunization with amyloid-beta attenuates Alzheimer-disease-like pathology in the PDAPP mouse. *Nature.* 1999; 400:173-7.
- [42] Janus C, Pearson J, McLaurin J, Mathews PM, Jiang Y, Schmidt SD, Chishti MA, Horne P, Heslin D, French J, Mount HT, Nixon RA, Mercken M, Bergeron C, Fraser PE, St George-Hyslop P, Westaway D. A beta peptide immunization reduces behavioural impairment and plaques in a model of Alzheimer's disease. *Nature.* 2000; 408: 979-82.
- [43] Morgan D, Diamond DM, Gottschall PE, Ugen KE, Dickey C, Hardy J, Duff K, Jantzen P, DiCarlo G, Wilcock D, Connor K, Hatcher J, Hope C, Gordon M, Arendash GW. A beta peptide vaccination prevents memory loss in an animal model of Alzheimer's disease. *Nature.* 2000; 408: 982-5.
- [44] Orgogozo JM, Gilman S, Dartigues JF, Laurent B, Puel M, Kirby LC, Jouanny P, Dubois B, Eisner L, Flitman S, Michel BF, Boada M, Frank A, Hock C. Subacute meningoencephalitis in a subset of patients with AD after Abeta42 immunization. *Neurology.* 2003; 61: 46-54.
- [45] Nicoll JA, Wilkinson D, Holmes C, Steart P, Markham H, Weller RO. Neuropathology of human Alzheimer disease after immunization with amyloid-beta peptide: a case report. *Nat Med.* 2003; 9: 448-52.
- [46] Ferrer I, Boada Rovira M, Sánchez Guerra ML, Rey MJ, Costa-Jussá F. Neuropathology and pathogenesis of encephalitis following amyloid-beta immunization in Alzheimer's disease. *Brain Pathol.* 2004; 14: 11-20.
- [47] Meyer-Luehmann M, Mora JR, Mielke M, Spires-Jones TL, de Calignon A, von Andrian UH, Hyman BT. T cell mediated cerebral hemorrhages and microhemorrhages during passive Abeta immunization in APPPS1 transgenic mice. *Mol Neurodegener.* 2011; 6: 22.

- [48] Patton RL, Kalback WM, Esh CL, Kokjohn TA, Van Vickie GD, Luehrs DC, Kuo YM, Lopez J, Brune D, Ferrer I, Masliah E, Newel AJ, Beach TG, Castaño EM, Roher AE. Amyloid-beta peptide remnants in AN-1792-immunized Alzheimer's disease patients: a biochemical analysis. *Am J Pathol.* 2006; 169: 1048-63.
- [49] Nicoll JA, Barton E, Boche D, Neal JW, Ferrer I, Thompson P, Vlachouli C, Wilkinson D, Bayer A, Games D, Seubert P, Schenk D, Holmes C. Abeta species removal after abeta42 immunization. *J Neuropathol Exp Neurol.* 2006 ; 65: 1040-8.
- [50] Serrano-Pozo A, William CM, Ferrer I, Uro-Coste E, Delisle MB, Maurage CA, Hock C, Nitsch RM, Masliah E, Growdon JH, Frosch MP, Hyman BT. Beneficial effect of human anti-amyloid-beta active immunization on neurite morphology and tau pathology. *Brain.* 2010; 133: 1312-27.
- [51] Agadjanyan MG, Ghochikyan A, Petrushina I, Vasilevko V, Movsesyan N, Mkrtichyan M, Saing T, Cribbs DH. Prototype Alzheimer's disease vaccine using the immunodominant B cell epitope from beta-amyloid and promiscuous T cell epitope pan HLA DR-binding peptide. *J Immunol.* 2005; 174:1580-6.
- [52] McLaurin J, Cecal R, Kierstead ME, Tian X, Phinney AL, Manea M, French JE, Lambermon MH, Darabie AA, Brown ME, Janus C, Chishti MA, Horne P, Westaway D, Fraser PE, Mount HT, Przybylski M, St George-Hyslop P. Therapeutically effective antibodies against amyloid-beta peptide target amyloid-beta residues 4-10 and inhibit cytotoxicity and fibrillogenesis. *Nat Med.* 2002; 8: 1263-9.
- [53] Lemere CA, Maier M, Jiang L, Peng Y, Seabrook TJ. Amyloid-beta immunotherapy for the prevention and treatment of Alzheimer disease: lessons from mice, monkeys, and humans. *Rejuvenation Res.* 2006; 9: 77-84.
- [54] Wilcock DM, Alamed J, Gottschall PE, Grimm J, Rosenthal A, Pons J, Ronan V, Symmonds K, Gordon MN, Morgan D. Deglycosylated anti-amyloid-beta antibodies eliminate cognitive deficits and reduce parenchymal amyloid with minimal vascular consequences in aged amyloid precursor protein transgenic mice. *J Neurosci.* 2006; 26: 5340-6.
- [55] Solomon B, Frenkel D. Immunotherapy for Alzheimer's disease. *Neuropharmacology.* 2010; 59: 303-9.
- [56] Delrieu J, Ousset PJ, Caillaud C, Vellas B. 'Clinical trials in Alzheimer's disease': immunotherapy approaches. *J Neurochem.* 2012; 120 Suppl 1:186-93.
- [57] Golde TE, Schneider LS, Koo EH. Anti-a β therapeutics in Alzheimer's disease: the need for a paradigm shift. *Neuron.* 2011; 69: 203-13.
- [58] Morris M, Maeda S, Vossel K, Mucke L. The many faces of tau. *Neuron.* 2011; 70: 410-26.
- [59] Huang Y, Mucke L. Alzheimer mechanisms and therapeutic strategies. *Cell* 2012; 148: 1204-22.

- [60] Wischik CM, Edwards PC, Lai RY, Roth M, Harrington CR. Selective inhibition of Alzheimer disease-like tau aggregation by phenothiazines. *Proc Natl Acad Sci USA*. 1996; 93:11213-8.
- [61] O'Leary JC 3rd, Li Q, Marinec P, Blair LJ, Congdon EE, Johnson AG, Jinwal UK, Koren J 3rd, Jones JR, Kraft C, Peters M, Abisambra JF, Duff KE, Weeber EJ, Gestwicki JE, Dickey CA. Phenothiazine-mediated rescue of cognition in tau transgenic mice requires neuroprotection and reduced soluble tau burden. *Mol Neurodegener*. 2010; 5: 45.
- [62] Gura T. Hope in Alzheimer's fight emerges from unexpected places. *Nat Med*. 2008; 14: 894.
- [63] Yoshiyama Y, Higuchi M, Zhang B, Huang SM, Iwata N, Saido TC, Maeda J, Suhara T, Trojanowski JQ, Lee VM. Synapse loss and microglial activation precede tangles in a P301S tauopathy mouse model. *Neuron*. 2007; 53: 337-51.
- [64] Chambraud B, Sardin E, Giustiniani J, Dounane O, Schumacher M, Goedert M, Baulieu EE. A role for FKBP52 in Tau protein function. *Proc Natl Acad Sci U S A*. 2010; 107: 2658-63.
- [65] Taniguchi S, Suzuki N, Masuda M, Hisanaga S, Iwatsubo T, Goedert M, Hasegawa M. Inhibition of heparin-induced tau filament formation by phenothiazines, polyphenols, and porphyrins. *J Biol Chem*. 2005; 280: 7614-23.
- [66] Gong CX, Iqbal K. Hyperphosphorylation of microtubule-associated protein tau: a promising therapeutic target for Alzheimer disease. *Curr Med Chem*. 2008; 15: 2321-8.
- [67] Mi K, Johnson GV. The role of tau phosphorylation in the pathogenesis of Alzheimer's disease. *Curr Alzheimer Res*. 2006; 3: 449-63.
- [68] Hernández F, de Barreda EG, Fuster-Matanzo A, Goñi-Oliver P, Lucas JJ, Avila J. The role of GSK3 in Alzheimer disease. *Brain Res Bull*. 2009; 80: 248-50.
- [69] Mazanetz MP, Fischer PM. Untangling tau hyperphosphorylation in drug design for neurodegenerative diseases. *Nat Rev Drug Discov*. 2007; 6: 464-79.
- [70] Noble W, Planel E, Zehr C, Olm V, Meyerson J, Suleman F, Gaynor K, Wang L, LaFrancois J, Feinstein B, Burns M, Krishnamurthy P, Wen Y, Bhat R, Lewis J, Dickson D, Duff K. Inhibition of glycogen synthase kinase-3 by lithium correlates with reduced tauopathy and degeneration in vivo. *Proc Natl Acad Sci U S A*. 2005; 102: 6990-5.
- [71] Grandjean EM, Aubry JM. Lithium: updated human knowledge using an evidence-based approach. Part II: Clinical pharmacology and therapeutic monitoring. *CNS Drugs*. 2009; 23: 331-49.
- [72] Alvarez A, Toro R, Cáceres A, Maccioni RB. Inhibition of tau phosphorylating protein kinase cdk5 prevents beta-amyloid-induced neuronal death. *FEBS Lett*. 1999; 459: 421-6

- [73] Zheng YL, Kesavapany S, Gravel M, Hamilton RS, Schubert M, Amin N, Albers W, Grant P, Pant HC. A Cdk5 inhibitory peptide reduces tau hyperphosphorylation and apoptosis in neurons. *EMBO J*. 2005; 24: 209-20.
- [74] Wen Y, Planel E, Herman M, Figueroa HY, Wang L, Liu L, Lau LF, Yu WH, Duff KE. Interplay between cyclin-dependent kinase 5 and glycogen synthase kinase 3 beta mediated by neuregulin signaling leads to differential effects on tau phosphorylation and amyloid precursor protein processing. *J Neurosci*. 2008; 28: 2624-32.
- [75] Cheung ZH, Ip NY. Cdk5: a multifaceted kinase in neurodegenerative diseases. *Trends Cell Biol*. 2012; 22: 169-75.
- [76] Li L, Sengupta A, Haque N, Grundke-Iqbal I, Iqbal K. Memantine inhibits and reverses the Alzheimer type abnormal hyperphosphorylation of tau and associated neurodegeneration. *FEBS Lett*. 2004; 566: 261-9.
- [77] Cheng Y, Feng Z, Zhang QZ, Zhang JT. Beneficial effects of melatonin in experimental models of Alzheimer disease. *Acta Pharmacol Sin*. 2006; 27: 129-39.
- [78] Torrent L, Ferrer I. PP2A and Alzheimer disease. *Curr Alzheimer Res*. 2012; 9: 248-56.
- [79] Ittner LM, Ke YD, Delerue F, Bi M, Gladbach A, van Eersel J, Wölfling H, Chieng BC, Christie MJ, Napier IA, Eckert A, Staufenbiel M, Hardeman E, Götz J. Dendritic function of tau mediates amyloid-beta toxicity in Alzheimer's disease mouse models. *Cell*. 2010; 142: 387-97.
- [80] Roberson ED, Searce-Levie K, Palop JJ, Yan F, Cheng IH, Wu T, Gerstein H, Yu GQ, Mucke L. Reducing endogenous tau ameliorates amyloid beta-induced deficits in an Alzheimer's disease mouse model. *Science*. 2007; 316: 750-4.
- [81] Roberson ED, Halabisky B, Yoo JW, Yao J, Chin J, Yan F, Wu T, Hamto P, Devidze N, Yu GQ, Palop JJ, Noebels JL, Mucke L. Amyloid- β /Fyn-induced synaptic, network, and cognitive impairments depend on tau levels in multiple mouse models of Alzheimer's disease. *J Neurosci*. 2011; 31: 700-11.
- [82] Sahara N, Murayama M, Mizoroki T, Urushitani M, Imai Y, Takahashi R, Murata S, Tanaka K, Takashima A. In vivo evidence of CHIP up-regulation attenuating tau aggregation. *J Neurochem*. 2005; 94:1254-63.
- [83] Min SW, Cho SH, Zhou Y, Schroeder S, Haroutunian V, Seeley WW, Huang EJ, Shen Y, Masliah E, Mukherjee C, Meyers D, Cole PA, Ott M, Gan L. Acetylation of tau inhibits its degradation and contributes to tauopathy. *Neuron*. 2010; 67: 953-66.
- [84] Cohen TJ, Guo JL, Hurtado DE, Kwong LK, Mills IP, Trojanowski JQ, Lee VM. The acetylation of tau inhibits its function and promotes pathological tau aggregation. *Nat Commun*. 2011; 2: 252.

- [85] Asuni AA, Boutajangout A, Quartermain D, Sigurdsson EM. Immunotherapy targeting pathological tau conformers in atangle mouse model reduces brain pathology with associated functional improvements. *J Neurosci.* 2007; 27: 9115-29.
- [86] Boimel M, Grigoriades N, Lourbopoulos A, Haber E, Abramsky O, Rosennmann H. Efficacy and safety of immunization with phosphorylated tau against neurofibrillary tangles in mice. *Exp Neurol* 2010; 224: 472-85.
- [87] Chai X, Wu S, Murray TK, Kinley R, Cella CV, Sims H, Buckner N, Hanmer J, Davies P, O'Neill MJ, Hutton ML, Citron M. Passive immunization with anti-Tau antibodies in two transgenic models: reduction of Tau pathology and delay of disease progression. *J Biol Chem.* 2011; 286: 34457-67.
- [88] Brunden KR, Yao Y, Potuzak JS, Ferrer NI, Ballatore C, James MJ, Hogan AM, Trojanowski JQ, Smith AB 3rd, Lee VM. The characterization of microtubule-stabilizing drugs as possible therapeutic agents for Alzheimer's disease and related tauopathies. *Pharmacol Res.* 2011; 63: 341-51.
- [89] Zempel H, Thies E, Mandelkow E, Mandelkow EM. Abeta oligomers cause localized Ca(2+) elevation, missorting of endogenous Tau into dendrites, Tau phosphorylation, and destruction of microtubules and spines. *J Neurosci.* 2010; 30: 11938-50.
- [90] Zhang B, Maiti A, Shively S, Lakhani F, McDonald-Jones G, Bruce J, Lee EB, Xie SX, Joyce S, Li C, Toleikis PM, Lee VM, Trojanowski JQ. Microtubule-binding drugs offset tau sequestration by stabilizing microtubules and reversing fast axonal transport deficits in a tauopathy model. *Proc Natl Acad Sci USA.* 2005; 102:: 227-31.
- [91] Brunden KR, Zhang B, Carroll J, Yao Y, Potuzak JS, Hogan AM, Iba M, James MJ, Xie SX, Ballatore C, Smith AB 3rd, Lee VM, Trojanowski JQ. Epothilone D improves microtubule density, axonal integrity, and cognition in a transgenic mouse model of tauopathy. *J Neurosci.* 2010; 30:13861-6.
- [92] Vulih-Shultzman I, Pinhasov A, Mandel S, Grigoriadis N, Touloumi O, Pittel Z, Gozes I. Activity-dependent neuroprotective protein snippet NAP reduces tau hyperphosphorylation and enhances learning in a novel transgenic mouse model. *J Pharmacol Exp Ther.* 2007; 323: 438-49.
- [93] Gozes I, Stewart A, Morimoto B, Fox A, Sutherland K, Schmeche D. Addressing Alzheimer's disease tangles: from NAP to AL-108. *Curr Alzheimer Res.* 2009; 6: 455-60.
- [94] Nunomura A, Perry G, Aliev G, Hirai K, Takeda A, Balraj EK, Jones PK, Ghanbari H, Wataya T, Shimohama S, Chiba S, Atwood CS, Petersen RB, Smith MA. Oxidative damage is the earliest event in Alzheimer disease. *J Neuropathol Exp Neurol.* 2001; 60:759-67.
- [95] Nunomura A, Perry G, Pappolla MA, Friedland RP, Hirai K, Chiba S, Smith MA. Neuronal oxidative stress precedes amyloid-beta deposition in Down syndrome. *J Neuropathol Exp Neurol.* 2000; 59: 1011-7

- [96] Perry G, Smith MA. Is oxidative damage central to the pathogenesis of Alzheimer disease? *Acta Neurol Belg.* 1998; 98:175-9.
- [97] Praticò D, Uryu K, Leight S, Trojanowski JQ, Lee VM. Increased lipid peroxidation precedes amyloid plaque formation in an animal model of Alzheimer amyloidosis. *J Neurosci.* 2001; 21: 4183-7.
- [98] Praticò D, Clark CM, Liun F, Rokach J, Lee VY, Trojanowski JQ. Increase of brain oxidative stress in mild cognitive impairment: a possible predictor of Alzheimer disease. *Arch Neurol.* 2002 59(6):972-6. Erratum in: *Arch Neurol* 2002; 59:1475.
- [99] Martínez A, Portero-Otin M, Pamplona R, Ferrer I. Protein targets of oxidative damage in human neurodegenerative diseases with abnormal protein aggregates. *Brain Pathol.* 2010; 20: 281-97.
- [100] Terni B, Boada J, Portero-Otin M, Pamplona R, Ferrer I. Mitochondrial ATP-synthase in the entorhinal cortex is a target of oxidative stress at stages I/II of Alzheimer's disease pathology. *Brain Pathol.* 2010; 20: 222-33.
- [101] Praticò D. Evidence of oxidative stress in Alzheimer's disease brain and antioxidant therapy: lights and shadows. *Ann N Y Acad Sci.* 2008; 1147: 70-8.
- [102] Bonda DJ, Wang X, Perry G, Nunomura A, Tabaton M, Zhu X, Smith MA. Oxidative stress in Alzheimer disease: a possibility for prevention. *Neuropharmacology.* 2010; 59: 290-4
- [103] Richard T, Pawlus AD, Iglésias ML, Pedrot E, Waffo-Teguo P, Mérillon JM, Monti JP. Neuroprotective properties of resveratrol and derivatives. *Ann N Y Acad Sci.* 2011; 1215: 103-8.
- [104] Li F, Gong Q, Dong H, Shi J. Resveratrol, a neuroprotective supplement for Alzheimer's disease. *Curr Pharm Des.* 2012; 18: 27-33.
- [105] Mancuso C, Bates TE, Butterfield DA, Calafato S, Cornelius C, De Lorenzo A, Dinkova Kostova AT, Calabrese V. Natural antioxidants in Alzheimer's disease. *Expert Opin Investig Drugs.* 2007; 16:1921-31.
- [106] Belkacemi A, Doggui S, Dao L, Ramassamy C. Challenges associated with curcumin therapy in Alzheimer disease. *Expert Rev Mol Med.* 2011; 13:e34.
- [107] Hamaguchi T, Ono K, Yamada M. REVIEW: Curcumin and Alzheimer's disease. *CNS Neurosci Ther.* 2010; 16: 285-97.
- [108] Mandel SA, Amit T, Weinreb O, Reznichenko L, Youdim MB. Simultaneous manipulation of multiple brain targets by green tea catechins: a potential neuroprotective strategy for Alzheimer and Parkinson diseases. *CNS Neurosci Ther.* 2008; 14: 352-65.
- [109] Spagnoli A, Lucca U, Menasce G, Bandera L, Cizza G, Forloni G, Tettamanti M, Frattura L, Tiraboschi P, Comelli M, et al. Long-term acetyl-L-carnitine treatment in Alzheimer's disease. *Neurology.* 1991; 41: 1726-32.

- [110] Montgomery SA, Thal LJ, Amrein R. Meta-analysis of double blind randomized controlled clinical trials of acetyl-L-carnitine versus placebo in the treatment of mild cognitive impairment and mild Alzheimer's disease. *Int Clin Psychopharmacol*. 2003; 18: 61-71.
- [111] Nishida Y, Yokota T, Takahashi T, Uchihara T, Jishage K, Mizusawa H. Deletion of vitamin E enhances phenotype of Alzheimer disease model mouse. *Biochem Biophys Res Commun*. 2006; 350: 530-6.
- [112] Viña J, Lloret A, Giraldo E, Badia MC, Alonso MD. Antioxidant pathways in Alzheimer's disease: possibilities of intervention. *Curr Pharm Des* 2011; 17: 3861-4.
- [113] Harrison FE, Hosseini AH, McDonald MP, May JM. Vitamin C reduces spatial learning deficits in middle-aged and very old APP/PSEN1 transgenic and wild-type mice. *Pharmacol Biochem Behav*. 2009; 93: 443-50.
- [114] Murakami K, Murata N, Ozawa Y, Kinoshita N, Irie K, Shirasawa T, Shimizu T. Vitamin C restores behavioral deficits and amyloid- oligomerization without affecting plaque formation in a mouse model of Alzheimer's disease. *J Alzheimers Dis*. 2011; 26: 7-18.
- [115] Zandi PP, Anthony JC, Khachaturian AS, Stone SV, Gustafson D, Tschanz JT, Norton MC, Welsh-Bohmer KA, Breitner JC; Cache County Study Group. Reduced risk of Alzheimer disease in users of antioxidant vitamin supplements: the Cache County Study. *Arch Neurol*. 2004; 61: 82-8
- [116] Boothby LA, Doering PL. Vitamin C and vitamin E for Alzheimer's disease. *Ann Pharmacother*. 2005; 39: 2073-80.
- [117] Birks J, Grimley Evans J. Ginkgo biloba for cognitive impairment and dementia. *Cochrane Database Syst Rev*. 2009; (1):CD003120.
- [118] Tchantchou F, Xu Y, Wu Y, Christen Y, Luo Y. EGb 761 enhances adult hippocampal neurogenesis and phosphorylation of CREB in transgenic mouse model of Alzheimer's disease. *FASEB J*. 2007; 21: 2400-8.
- [119] Galasko DR, Peskind E, Clark CM, Quinn JF, Ringman JM, Jicha GA, Cotman C, Cottrell B, Montine TJ, Thomas RG, Aisen P; for the Alzheimer's Disease Cooperative Study. Antioxidants for Alzheimer disease: a randomized clinical trial with cerebrospinal fluid biomarker measures. *Arch Neurol*. 2012 Mar 19.
- [120] Holmquist L, Stuchbury G, Berbaum K, Muscat S, Young S, Hager K, Engel J, Münch G. Lipoic acid as a novel treatment for Alzheimer's disease and related dementias. *Pharmacol Ther*. 2007; 113:154-64.
- [121] Maczurek A, Hager K, Kenklies M, Sharman M, Martins R, Engel J, Carlson DA, Münch G. Lipoic acid as an anti-inflammatory and neuroprotective treatment for Alzheimer's disease. *Adv Drug Deliv Rev*. 2008; 60: 1463-70.

- [122] Pocernich CB, Butterfield DA. Elevation of glutathione as a therapeutic strategy in Alzheimer disease. *Biochim Biophys Acta*. 2012; 1822: 625-30.
- [123] Adair JC, Knoefel JE, Morgan N. Controlled trial of N-acetylcysteine for patients with probable Alzheimer's disease. *Neurology*. 2001; 578: 1515-7.
- [124] Orsucci D, Mancuso M, Ienco EC, LoGerfo A, Siciliano G. Targeting mitochondrial dysfunction and neurodegeneration by means of coenzyme Q10 and its analogues. *Curr Med Chem*. 2011; 18: 4053-64.
- [125] Beal MF. Mitochondrial dysfunction and oxidative damage in Alzheimer's and Parkinson's diseases and coenzyme Q10 as a potential treatment. *J Bioenerg Biomembr*. 2004; 36: 381-6.
- [126] Dumont M, Kipiani K, Yu F, Wille E, Katz M, Calingasan NY, Gouras GK, Lin MT, Beal MF. Coenzyme Q10 decreases amyloid pathology and improves behavior in a transgenic mouse model of Alzheimer's disease. *J Alzheimers Dis*. 2011; 27: 211-23.
- [127] McManus MJ, Murphy MP, Franklin JL. The mitochondria-targeted antioxidant MitoQ prevents loss of spatial memory retention and early neuropathology in a transgenic mouse model of Alzheimer's disease. *J Neurosci*. 2011; 31:15703-15.
- [128] Hensley K. Neuroinflammation in Alzheimer's disease: mechanisms, pathologic consequences, and potential for therapeutic manipulation. *J Alzheimers Dis*. 2010; 21:1-14.
- [129] McGeer PL, McGeer E, Rogers J, Sibley J. Anti-inflammatory drugs and Alzheimer disease. *Lancet*. 1990; 335:1037.
- [130] Kaufmann WE, Andreasson KI, Isakson PC, Worley PF. Cyclooxygenases and the central nervous system. *Prostaglandins*. 1997; 54: 601-24.
- [131] Lim GP, Yang F, Chu T, Chen P, Beech W, Teter B, Tran T, Ubeda O, Ashe KH, Frautschy SA, Cole GM. Ibuprofen suppresses plaque pathology and inflammation in a mouse model for Alzheimer's disease. *J Neurosci*. 2000; 20: 5709-14.
- [132] Jantzen PT, Connor KE, DiCarlo G, Wenk GL, Wallace JL, Rojiani AM, Coppola D, Morgan D, Gordon MN. Microglial activation and beta -amyloid deposit reduction caused by a nitric oxide-releasing nonsteroidal anti-inflammatory drug in amyloid precursor protein plus presenilin-1 transgenic mice. *J Neurosci*. 2002; 22: 2246-54.
- [133] Yan Q, Zhang J, Liu H, Babu-Khan S, Vassar R, Biere AL, Citron M, Landreth G. Anti-inflammatory drug therapy alters beta-amyloid processing and deposition in an animal model of Alzheimer's disease. *J Neurosci*. 2003; 23: 7504-9.
- [134] Heneka MT, Sastre M, Dumitrescu-Ozimek L, Hanke A, Dewachter I, Kuiperi C, O'Banion K, Klockgether T, Van Leuven F, Landreth GE. Acute treatment with the PPARgamma agonist pioglitazone and ibuprofen reduces glial inflammation and Abeta1-42 levels in APPV717I transgenic mice. *Brain*. 2005; 128: 1442-53.

- [135] Lim GP, Yang F, Chu T, Gahtan E, Ubeda O, Beech W, Overmier JB, Hsiao-Ashec K, Frautschy SA, Cole GM. Ibuprofen effects on Alzheimer pathology and open field activity in APPsw transgenic mice. *Neurobiol Aging*. 2001; 22: 983-91.
- [136] Quinn J, Montine T, Morrow J, Woodward WR, Kulhanek D, Eckenstein F. Inflammation and cerebral amyloidosis are disconnected in an animal model of Alzheimer's disease. *J Neuroimmunol*. 2003; 137: 32-41.
- [137] Sung S, Yang H, Uryu K, Lee EB, Zhao L, Shineman D, Trojanowski JQ, Lee VM, Praticò D. Modulation of nuclear factor-kappa B activity by indomethacin influences A beta levels but not A beta precursor protein metabolism in a model of Alzheimer's disease. *Am J Pathol*. 2004; 165: 2197-206.
- [138] Kukar T, Murphy MP, Eriksen JL, Sagi SA, Weggen S, Smith TE, Ladd T, Khan MA, Kache R, Beard J, Dodson M, Merit S, Ozols VV, Anastasiadis PZ, Das P, Fauq A, Koo EH, Golde TE. Diverse compounds mimic Alzheimer disease-causing mutations by augmenting Abeta42 production. *Nat Med*. 2005; 11: 545-50.
- [139] Jaturapatporn D, Isaac MG, McCleery J, Tabet N. Aspirin, steroidal and non-steroidal anti-inflammatory drugs for the treatment of Alzheimer's disease. *Cochrane Database Syst Rev*. 2012; CD006378.
- [140] Coma M, Serenó L, Da Rocha-Souto B, Scotton TC, España J, Sánchez MB, Rodríguez M, Agulló J, Guardia-Laguarta C, Garcia-Alloza M, Borrelli LA, Clarimón J, Lleó A, Bacskaí BJ, Saura CA, Hyman BT, Gómez-Isla T. Triflusal reduces dense-core plaque load, associated axonal alterations and inflammatory changes, and rescues cognition in a transgenic mouse model of Alzheimer's disease. *Neurobiol Dis*. 2010; 38: 482-91.
- [141] Mosconi L, Pupi A, De Leon MJ. Brain glucose hypometabolism and oxidative stress in preclinical Alzheimer's disease. *Ann NY Acad Sci* 2008; 147: 180-95.
- [142] Ferrer I. Altered mitochondria, energy metabolism, voltage-dependent anion channel, and lipid rafts converge to exhaust neurons in Alzheimer's disease. *J Bioenerg Bio-membr*. 2009; 41: 425-31.
- [143] Ferreira IL, Resende R, Ferreira E, Rego AC, Pereira CF. Multiple defects in energy metabolism in Alzheimer's disease. *Curr Drug Targets*. 2010; 11: 1193-206.
- [144] Ankarcrona M, Mangialasche F, Winblad B. Rethinking Alzheimer's disease therapy: are mitochondria the key? *J Alzheimers Dis*. 2010; 20 Suppl 2: S579-90
- [145] Cunnane S, Nugent S, Roy M, Courchesne-Loyer A, Croteau E, Tremblay S, Castellano A, Pifferi F, Bocti C, Paquet N, Begdouri H, Bentourkia M, Turcotte E, Allard M, Barberger-Gateau P, Fulop T, Rapoport SI. Brain fuel metabolism, aging, and Alzheimer's disease. *Nutrition*. 2011; 27: 3-20.
- [146] Kapogiannis D, Mattson MP. Disrupted energy metabolism and neuronal circuit dysfunction in cognitive impairment and Alzheimer's disease. *Lancet Neurol*. 2011; 10: 187-98.

- [147] Jagust WJ, Seab JP, Huesman RH, Valk PE, Mathis CA, Reed BR, Coxson PG, Budinger TF. Diminished glucose transport in Alzheimer's disease: dynamic PET studies. *J Cereb Blood Flow Metab.* 1991; 11: 323-30.
- [148] Pedersen WA, Flynn ER. Insulin resistance contributes to aberrant stress responses in the Tg2576 mouse model of Alzheimer's disease. *Neurobiol Dis* 2004; 17: 500-6.
- [149] Landreth G, Jiang Q, Mandrekar S, Heneka M. PPARgamma agonists as therapeutics for the treatment of Alzheimer's disease. *Neurotherapeutics.* 2008; 5: 481-9.
- [150] Bak AM, Egefjord L, Gejl M, Steffensen C, Stecher CW, Smidt K, Brock B, Rungby J. Targeting amyloid-beta by glucagon-like peptide -1 GLP-1) in Alzheimer's disease and diabetes. *Expert Opin Ther Targets.* 2011; 15: 1153-62.
- [151] Bomfim TR, Forny-Germano L, Sathler LB, Brito-Moreira J, Houzel JC, Decker H, Silverman MA, Kazi H, Melo HM, McClean PL, Holscher C, Arnold SE, Talbot K, Klein WL, Munoz DP, Ferreira ST, De Felice FG. An anti-diabetes agent protects the mouse brain from defective insulin signaling caused by Alzheimer's disease-associated A β oligomers. *J Clin Invest.* 2012; 122: 1339-53.
- [152] Miichi Y, Sakurai T, Akisaki T, Yokono K. Effects of insulin and amyloid β (1-42) oligomers on glucose incorporation and mitochondrial function in cultured rat hippocampal neurons. *Geriatr Gerontol Int.* 2011; 11: 517-24.
- [153] Miller BW, Willett KC, Desilets AR. Rosiglitazone and pioglitazone for the treatment of Alzheimer's disease. *Ann Pharmacother.* 2011; 45: 1416-24.
- [154] Van der Auwera I, Wera S, Van Leuven F, Henderson ST. A ketogenic diet reduces amyloid beta 40 and 42 in a mouse model of Alzheimer's disease. *Nutr Metab.* 2005; 2: 28.
- [155] Yao J, Chen S, Mao Z, Cadenas E, Brinton RD. 2-Deoxy-D-glucose treatment induces ketogenesis, sustains mitochondrial function, and reduces pathology in female mouse model of Alzheimer's disease. *PLoS One.* 2011; 6:e21788.
- [156] Henderson ST, Vogel JL, Barr LJ, Garvin F, Jones JJ, Costantini LC. Study of the ketogenic agent AC-1202 in mild to moderate Alzheimer's disease: a randomized, double-blind, placebo-controlled, multicenter trial. *Nutr Metab.* 2009; 6:31.
- [157] Brewer GJ, Wallimann TW. Protective effect of the energy precursor creatine against toxicity of glutamate and beta-amyloid in rat hippocampal neurons. *J Neurochem.* 2000; 74: 1968-78.
- [158] Sullivan PG, Geiger JD, Mattson MP, Scheff SW. Dietary supplement creatine protects against traumatic brain injury. *Ann Neurol.* 2000; 48: 723-9.
- [159] Caspersen C, Wang N, Yao J, Sosunov A, Chen X, Lustbader JW, Xu HW, Stern D, McKhann G, Yan SD. Mitochondrial A β : a potential focal point for neuronal metabolic dysfunction in Alzheimer's disease. *FASEB J.* 2005; 19: 2040-1.

- [160] Lustbader JW, Cirilli M, Lin C, Xu HW, Takuma K, Wang N, Caspersen C, Chen X, Pollak S, Chaney M, Trinchese F, Liu S, Gunn-Moore F, Lue LF, Walker DG, Kuppasamy P, Zewier ZL, Arancio O, Stern D, Yan SS, Wu H. ABAD directly links Abeta to mitochondrial toxicity in Alzheimer's disease. *Science*. 2004; 304: 448-52.
- [161] Lim YA, Grimm A, Giese M, Mensah-Nyagan AG, Villafranca JE, Ittner LM, Eckert A, Götz J. Inhibition of the mitochondrial enzyme ABAD restores the amyloid- β -mediated deregulation of estradiol. *PLoS One*. 2011; 6: e28887
- [162] Yao J, Du H, Yan S, Fang F, Wang C, Lue LF, Guo L, Chen D, Stern DM, Gunn Moore FJ, Xi Chen J, Arancio O, Yan SS. Inhibition of amyloid-beta (Abeta)peptide-binding alcohol dehydrogenase-Abeta interaction reduces Abeta accumulation and improves mitochondrial function in a mouse model of Alzheimer's disease. *J Neurosci*. 2011; 31: 2313-20.
- [163] Yamada K, Inagaki N. Neuroprotection by KATP channels. *J Mol Cell Cardiol*. 2005 ; 38: 945-9.
- [164] Liu D, Pitta M, Lee JH, Ray B, Lahiri DK, Furukawa K, Mughal M, Jiang H, Villarreal J, Cutler RG, Greig NH, Mattson MP. The KATP channel activator diazoxide ameliorates amyloid- β and tau pathologies and improves memory in the 3xTgAD mouse model of Alzheimer's disease. *J Alzheimers Dis*. 2010; 22: 443-57.
- [165] Sabbagh MN, Shill HA. Latrepirdine, a potential novel treatment for Alzheimer's disease and Huntington's chorea. *Curr Opin Investig Drugs*. 2010; 11: 80-91.
- [166] Bachurin S, Lermontova N, Shevtzova E, Serkova T, Kireeva E. Comparative study of Tacrine and Dimebon action on mitochondrial permeability transition and β -amyloid-induced neurotoxicity. *J Neurochem*. 1999; 73(Suppl S): S185.
- [167] Lermontova NN, Redkozubov AE, Shevtsova EF, Serkova TP, Kireeva EG, Bachurin SO. Dimebon and tacrine inhibit neurotoxic action of β -amyloid in culture and block L-type Ca^{2+} channels. *Bull Exp Biol Med*. 2001; 132:1079-83.
- [168] Shevtsova EP, Grigoriev VV, Kireeva EG, Koroleva IV, Bachurin SO. Dimebon as mitoprotective and antiaging drug. *Biochim Biophys Acta*. 2006 (Suppl S) Abs P2.3.4.
- [169] Doody RS, Gavrilova SI, Sano M, Thomas RG, Aisen PS, Bachurin SO, Seely L, Hung D; dimebon investigators. Effect of dimebon on cognition, activities of daily living, behaviour, and global function in patients with mild-to-moderate Alzheimer's disease: a randomised, double-blind, placebo-controlled study. *Lancet*. 2008; 372: 207-15.
- [170] Francis PT, Ramírez MJ, Lai MK. Neurochemical basis for symptomatic treatment of Alzheimer's disease. *Neuropharmacology*. 2010; 59: 221-9.
- [171] Geula C, Nagykerly N, Nicholas A, Wu CK. Cholinergic neuronal and axonal abnormalities are present early in aging and in Alzheimer disease. *J Neuropathol Exp Neurol*. 2008; 67: 309-18.

- [172] Mufson EJ, Counts SE, Perez SE, Ginsberg SD. Cholinergic system during the progression of Alzheimer's disease: therapeutic implications. *Expert Rev Neurother*. 2008; 8: 1703-18.
- [173] Fisher A. Cholinergic treatments with emphasis on m1 muscarinic agonists as potential disease-modifying agents for Alzheimer's disease. *Neurotherapeutics*. 2008; 5: 433-42.
- [174] Higgins JP, Flicker L. Lecithin for dementia and cognitive impairment. *Cochrane Database Syst Rev*. 2003; CD001015.
- [175] Birks J. Cholinesterase inhibitors for Alzheimer's disease. *Cochrane Database Syst Rev*. 2006; (1):CD005593.
- [176] Wilcock GK, Dawbarn D. Current pharmacological approaches to treating Alzheimer's disease. In: *Neurobiology of Alzheimer's Disease*. Third edition. Edited by David Dawbarn and Shelley J. Allen. Oxford University Press. 2007. Pp 359-390.
- [177] Herrmann N, Chau SA, Kircanski I, Lanctôt KL. Current and emerging drug treatment options for Alzheimer's disease: a systematic review. *Drugs*. 2011; 71: 2031-65.
- [178] Fischer W, Wictorin K, Björklund A, Williams LR, Varon S, Gage FH. Amelioration of cholinergic neuron atrophy and spatial memory impairment in aged rats by nerve growth factor. *Nature*. 1987; 329: 65-8.
- [179] Caccamo A, Oddo S, Billings LM, Green KN, Martinez-Coria H, Fisher A, LaFerla FM. M1 receptors play a central role in modulating AD-like pathology in transgenic mice. *Neuron*. 2006; 49: 671-82.
- [180] Nitsch RM, Slack BE, Wurtman RJ, Growdon JH. Release of Alzheimer amyloid precursor derivatives stimulated by activation of muscarinic acetylcholine receptors. *Science*. 1992; 258: 304-7.
- [181] Wolf BA, Wertkin AM, Jolly YC, Yasuda RP, Wolfe BB, Konrad RJ, Manning D, Ravi S, Williamson JR, Lee VM. Muscarinic regulation of Alzheimer's disease amyloid precursor protein secretion and amyloid beta-protein production in human neuronal NT2N cells. *J Biol Chem*. 1995; 270: 4916-22.
- [182] Woodruff-Pak DS, Gould TJ. Neuronal nicotinic acetylcholine receptors: involvement in Alzheimer's disease and schizophrenia. *Behav Cogn Neurosci Rev*. 2002; 1: 5-20.
- [183] Parri RH, Dineley TK. Nicotinic acetylcholine receptor interaction with beta-amyloid: molecular, cellular, and physiological consequences. *Curr Alzheimer Res*. 2010; 7: 27-39.
- [184] Nordberg A, Hellström-Lindahl E, Lee M, Johnson M, Mousavi M, Hall R, Perry E, Bednar I, Court J. Chronic nicotine treatment reduces beta-amyloidosis in the brain of a mouse model of Alzheimer's disease (APPsw). *J Neurochem*. 2002; 81:655-8.

- [185] Oddo S, Caccamo A, Green KN, Liang K, Tran L, Chen Y, Leslie FM, LaFerla FM. Chronic nicotine administration exacerbates tau pathology in a transgenic model of Alzheimer's disease. *Proc Natl Acad Sci U S A*. 2005; 102: 3046-51.
- [186] Haydar SN, Dunlop J. Neuronal nicotinic acetylcholine receptors - targets for the development of drugs to treat cognitive impairment associated with schizophrenia and Alzheimer's disease. *Curr Top Med Chem*. 2010; 10:144-52.
- [187] Ren K, Thinschmidt J, Liu J, Ai L, Papke RL, King MA, Hughes JA, Meyer EM. alpha7 Nicotinic receptor gene delivery into mouse hippocampal neurons leads to functional receptor expression, improved spatial memory-related performance, and tau hyperphosphorylation. *Neuroscience*. 2007; 145: 314-22
- [188] Kihara T, Shimohama S, Urushitani M, Sawada H, Kimura J, Kume T, Maeda T, Akaike A. Stimulation of alpha4beta2 nicotinic acetylcholine receptors inhibits beta-amyloid toxicity. *Brain Res*. 1998; 792: 331-4.
- [189] Tam SW, Zaczek R. Linopirdine. A depolarization-activated releaser of transmitters for treatment of dementia. *Adv Exp Med Biol*. 1995; 363:47-56.
- [190] Lachowicz JE, Duffy RA, Ruperto V, Kozlowski J, Zhou G, Clader J, Billard W, Binch H 3rd, Crosby G, Cohen-Williams M, Strader CD, Coffin V. Facilitation of acetylcholine release and improvement in cognition by a selective M2 muscarinic antagonist, SCH 72788. *Life Sci*. 2001; 68: 2585-92.
- [191] Clader JW, Wang Y. Muscarinic receptor agonists and antagonists in the treatment of Alzheimer's disease. *Curr Pharm Des*. 2005; 11: 3353-61.
- [192] Rockwood K, Beattie BL, Eastwood MR, Feldman H, Mohr E, Pryse-Phillips W, Gauthier S. A randomized, controlled trial of linopirdine in the treatment of Alzheimer's disease. *Can J Neurol Sci*. 1997; 24:140-5.
- [193] Boyle CD, Lachowicz JE. Orally active and selective benzyldene ketal M2 muscarinic receptor antagonists for the treatment of Alzheimer's disease. *Drug Dev Res* 2002; 56: 310 -20.
- [194] Palop JJ, Mucke L. Amyloid-beta-induced neuronal dysfunction in Alzheimer's disease: from synapses toward neural networks. *Nat Neurosci*. 2010; 13: 812-8.
- [195] Hu NW, Ondrejcek T, Rowan MJ. Glutamate receptors in preclinical research on Alzheimer's disease: update on recent advances. *Pharmacol Biochem Behav* 2012; 100: 855-62.
- [196] Keller JN, Mark RJ, Bruce AJ, Blanc E, Rothstein JD, Uchida K, Waeg G, Mattson MP. 4-Hydroxynonenal, an aldehydic product of membrane lipid peroxidation, impairs glutamate transport and mitochondrial function in synaptosomes. *Neuroscience*. 1997; 80: 685-96.

- [197] Li S, Hong S, Shepardson NE, Walsh DM, Shankar GM, Selkoe D. Soluble oligomers of amyloid Beta protein facilitate hippocampal long-term depression by disrupting neuronal glutamate uptake. *Neuron*. 2009; 62: 788-801.
- [198] Cissé M, Halabisky B, Harris J, Devidze N, Dubal DB, Sun B, Orr A, Lotz G, Kim DH, Hamto P, Ho K, Yu GQ, Mucke L. Reversing EphB2 depletion rescues cognitive functions in Alzheimer model. *Nature*. 2011; 469: 47-52.
- [199] Hoey SE, Williams RJ, Perkinton MS. Synaptic NMDA receptor activation stimulates alpha-secretase amyloid precursor protein processing and inhibits amyloid-beta production. *J Neurosci*. 2009; 29: 4442-60.
- [200] Bordji K, Becerril-Ortega J, Nicole O, Buisson A. Activation of extrasynaptic, but not synaptic, NMDA receptors modifies amyloid precursor protein expression pattern and increases amyloid- β production. *J Neurosci*. 2010; 30:15927-42.
- [201] Francis PT. Glutamatergic approaches to the treatment of cognitive and behavioural symptoms of Alzheimer's disease. *Neurodegener Dis*. 2008; 5: 241-3.
- [202] Lynch G, Gall CM. Glutamate-based therapeutic approaches: ampakines. *Curr Opin Pharmacol*. 2006; 6: 82-8.
- [203] Granger R, Deadwyler S, Davis M, Moskovitz B, Kessler M, Rogers G, Lynch G. Facilitation of glutamate receptors reverses an age-associated memory impairment in rats. *Synapse*. 1996; 22: 332-7.
- [204] Bartolini L, Casamenti F, Pepeu G. Aniracetam restores object recognition impaired by age, scopolamine, and nucleus basalis lesions. *Pharmacol Biochem Behav*. 1996; 53: 277-83.
- [205] Johnson SA, Simmon VF. Randomized, double-blind, placebo-controlled international clinical trial of the Ampakine CX516 in elderly participants with mild cognitive impairment: a progress report. *J Mol Neurosci*. 2002; 19: 197-200.
- [206] Myhrer T, Paulsen RE. Infusion of D-cycloserine into temporal-hippocampal areas and restoration of mnemonic function in rats with disrupted glutamatergic temporal systems. *Eur J Pharmacol*. 1997; 328: 1-7.
- [207] Schwartz BL, Hashtroudi S, Herting RL, Schwartz P, Deutsch SI. d-Cycloserine enhances implicit memory in Alzheimer patients. *Neurology*. 1996; 46: 420-4.
- [208] Parsons CG, Stöffler A, Danysz W. Memantine: a NMDA receptor antagonist that improves memory by restoration of homeostasis in the glutamatergic system — too little activation is bad, too much is even worse. *Neuropharmacology*. 2007; 53: 699-723.
- [209] Reisberg B, Doody R, Stöffler A, Schmitt F, Ferris S, Möbius HJ; Memantine Study Group. Memantine in moderate-to-severe Alzheimer's disease. *N Engl J Med*. 2003; 348:1333-41.

- [210] McKeage K. Memantine: a review of its use in moderate to severe Alzheimer's disease. *CNS Drugs*. 2009; 23: 881-97.
- [211] Danysz W, Parsons CG. The NMDA receptor antagonist memantine as asymptomatic and neuroprotective treatment for Alzheimer's disease: preclinical evidence. *Int J Geriatr Psychiatry*. 2003; 18(Suppl 1): S23-32.
- [212] Dong H, Yuede CM, Coughlan C, Lewis B, Csernansky JG. Effects of memantine on neuronal structure and conditioned fear in the Tg2576 mouse model of Alzheimer's disease. *Neuropsychopharmacology*. 2008; 33: 3226-36.
- [213] Scholtzova H, Wadghiri YZ, Douadi M, Sigurdsson EM, Li YS, Quartermain D, Banerjee P, Wisniewski T. Memantine leads to behavioral improvement and amyloid reduction in Alzheimer's-disease-model transgenic mice shown as by micromagnetic resonance imaging. *J Neurosci Res*. 2008; 86: 2784-91.
- [214] Martinez-Coria H, Green KN, Billings LM, Kitazawa M, Albrecht M, Rammes G, Parsons CG, Gupta S, Banerjee P, LaFerla FM. Memantine improves cognition and reduces Alzheimer's-like neuropathology in transgenic mice. *Am J Pathol*. 2010; 176: 870-80.
- [215] Filali M, Lalonde R, Rivest S. Subchronic memantine administration on spatial learning, exploratory activity, and nest-building in an APP/PS1 mouse model of Alzheimer's disease. *Neuropharmacology*. 2011; 60: 930-6.
- [216] Schneider LS, Dagerman KS, Higgins JP, McShane R. Lack of evidence for the efficacy of memantine in mild Alzheimer disease. *Arch Neurol*. 2011; 68: 991-8.
- [217] Bowen DM, Allen SJ, Benton JS, Goodhardt MJ, Haan EA, Palmer AM, Sims NR, Smith CC, Spillane JA, Esiri MM, Neary D, Snowden JS, Wilcock GK, Davison AN. Biochemical assessment of serotonergic and cholinergic dysfunction and cerebral atrophy in Alzheimer's disease. *J Neurochem*. 1983; 41: 266-72.
- [218] Arai H, Ichimiya Y, Kosaka K, Moroji T, Iizuka R. Neurotransmitter changes in early- and late-onset Alzheimer-type dementia. *Prog Neuropsychopharmacol Biol Psychiatry*. 1992; 16: 883-90.
- [219] Geldenhuys WJ, Van der Schyf CJ. Role of serotonin in Alzheimer's disease: a new therapeutic target? *CNS Drugs*. 2011; 25: 765-81.
- [220] Nelson RL, Guo Z, Halagappa VM, Pearson M, Gray AJ, Matsuoka Y, Brown M, Martin B, Iyun T, Maudsley S, Clark RF, Mattson MP. Prophylactic treatment with paroxetine ameliorates behavioral deficits and retards the development of amyloid and tau pathologies in 3xTgAD mice. *Exp Neurol*. 2007; 205:166-76.
- [221] Cirrito JR, Disabato BM, Restivo JL, Verges DK, Goebel WD, Sathyan A, Hayreh D, D'Angelo G, Benzinger T, Yoon H, Kim J, Morris JC, Mintun MA, Sheline YI. Serotonin signaling is associated with lower amyloid- β levels and plaques in transgenic mice and humans. *Proc Natl Acad Sci U S A*. 2011; 108: 14968-73.

- [222] Schechter LE, Dawson LA, Harder JA. The potential utility of 5-HT_{1A} receptor antagonists in the treatment of cognitive dysfunction associated with Alzheimer's disease. *Curr Pharm Des.* 2002; 8:139-45.
- [223] Schechter LE, Smith DL, Rosenzweig-Lipson S, Sukoff SJ, Dawson LA, Marquis K, Jones D, Piesla M, Andree T, Nawoschik S, Harder JA, Womack MD, Buccafusco J, Terry AV, Hoebel B, Rada P, Kelly M, Abou-Gharbia M, Barrett JE, Childers W. Lecozotan (SRA-333): a selective serotonin 1A receptor antagonist that enhances the stimulated release of glutamate and acetylcholine in the hippocampus and possesses cognitive-enhancing properties. *J Pharmacol Exp Ther.* 2005; 314:1274-89.
- [224] Patat A, Parks V, Raje S, Plotka A, Chassard D, Le Coz F. Safety, tolerability, pharmacokinetics and pharmacodynamics of ascending single and multiple doses of lecozotan in healthy young and elderly subjects. *Br J Clin Pharmacol.* 2009; 67: 299-308.
- [225] Upton N, Chuang TT, Hunter AJ, Virley DJ. 5-HT₆ receptor antagonists as novel cognitive enhancing agents for Alzheimer's disease. *Neurotherapeutics* 2008; 5: 458-69.
- [226] Cho S, Hu Y. Activation of 5-HT₄ receptors inhibits secretion of beta-amyloid peptides and increases neuronal survival. *Exp Neurol.* 2007; 203: 274-8.
- [227] Russo O, Cachard-Chastel M, Rivière C, Giner M, Soulier JL, Berthouze M, Richard T, Monti JP, Sicsic S, Lezoualc'h F, Berque-Bestel I. Design, synthesis, and biological evaluation of new 5-HT₄ receptor agonists: application as amyloid cascade modulators and potential therapeutic utility in Alzheimer's disease. *J Med Chem.* 2009; 52: 2214-25.
- [228] Knafo S, Alonso-Nanclares L, Gonzalez-Soriano J, Merino-Serrais P, Fernaud-Espinosa I, Ferrer I, DeFelipe J. Widespread changes in dendritic spines in a model of Alzheimer's disease. *Cereb Cortex.* 2009; 19: 586-92.
- [229] Smith DL, Pozueta J, Gong B, Arancio O, Shelanski M. Reversal of long-term dendritic spine alterations in Alzheimer disease models. *PNAS* 2009; 106: 16877-82.
- [230] Albasanz JL, Dalfó E, Ferrer I, Martín M. Impaired metabotropic glutamate receptor/phospholipase C signaling pathway in the cerebral cortex in Alzheimer's disease and dementia with Lewy bodies correlates with stage of Alzheimer's-disease-related changes. *Neurobiol Dis.* 2005; 20: 685-93.
- [231] Albasanz JL, Perez S, Barrachina M, Ferrer I, Martín M. Up-regulation of adenosine receptors in the frontal cortex in Alzheimer's disease. *Brain Pathol.* 2008; 18: 211-9.
- [232] Coleman P, Federoff H, Kurlan R. A focus on the synapse for neuroprotection in Alzheimer disease and other dementias. *Neurology.* 2004; 63:1155-62.
- [233] Arendt T. Alzheimer's disease as a disorder of mechanisms underlying structural brain self-organization. *Neuroscience.* 2001; 102: 723-65.
- [234] Palop JJ, Mucke L. Synaptic depression and aberrant excitatory network activity in Alzheimer's disease: two faces of the same coin? *Neuromolecular Med.* 2010; 12: 48-55.

- [235] Martin V, Fabelo N, Santpere G, Puig B, Marín R, Ferrer I, Diaz M. Lipid alterations in lipid rafts from Alzheimer's disease human brain cortex. *J Alzheimers Dis* 2010; 19: 489-502.
- [236] Darios F, Davletov B. Omega-3 and omega-6 fatty acids stimulate cell membrane expansion by acting on syntaxin 3. *Nature*. 2006; 440: 813-7.
- [237] Florent-Bécharde S, Desbène C, Garcia P, Allouche A, Youssef I, Escanyé MC, Koziel V, Hanse M, Malaplate-Armand C, Stenger C, Kriem B, Yen-Potin FT, Olivier JL, Pillot T, Oster T. The essential role of lipids in Alzheimer's disease. *Biochimie*. 2009; 91: 804-9.
- [238] Oster T, Pillot T. Docosahexaenoic acid and synaptic protection in Alzheimer's disease mice. *Biochim Biophys Acta*. 2010; 1801: 791-8.
- [239] Secades JJ, Lorenzo JL. Citicoline: pharmacological and clinical review, 2006 update. *Methods Find Exp Clin Pharmacol*. 2006; Suppl B:1-56.
- [240] Proctor DT, Coulson EJ, Dodd PR. Post-synaptic scaffolding protein interactions with glutamate receptors in synaptic dysfunction and Alzheimer's disease. *Prog Neurobiol*. 2011; 93: 509-21.
- [241] Kaplan DR, Miller FD. Neurotrophin signal transduction in the nervous system. *Curr Opin Neurobiol*. 2000; 10: 381-91.
- [242] Montero CN, Hefti F. Rescue of lesioned septal cholinergic neurons by nerve growth factor: specificity and requirement for chronic treatment. *J Neurosci*. 1988; 8: 2986-99.
- [243] Allard S, Leon WC, Pakavathkumar P, Bruno MA, Ribeiro-da-Silva A, Cuello AC. Impact of the NGF maturation and degradation pathway on the cortical cholinergic system phenotype. *J Neurosci*. 2012; 32: 2002-12.
- [244] Cuello AC, Bruno MA, Allard S, Leon W, Iulita MF. Cholinergic involvement in Alzheimer's disease. A link with NGF maturation and degradation. *J Mol Neurosci*. 2010; 40: 230-5.
- [245] Bruno MA, Leon WC, Frago G, Mushynski WE, Almazan G, Cuello AC. Amyloid beta-induced nerve growth factor dysmetabolism in Alzheimer disease. *J Neuropathol Exp Neurol*. 2009; 68: 857-69.
- [246] Pedraza CE, Podlesniy P, Vidal N, Arévalo JC, Lee R, Hempstead B, Ferrer I, Iglesias M, Espinet C. Pro-NGF isolated from the human brain affected by Alzheimer's disease induces neuronal apoptosis mediated by p75NTR. *Am J Pathol*. 2005; 166: 533-43.
- [247] Podlesniy P, Kichev A, Pedraza C, Saurat J, Encinas M, Perez B, Ferrer I, Espinet C. Pro-NGF from Alzheimer's disease and normal human brain displays distinctive abilities to induce processing and nuclear translocation of intracellular domain of p75NTR and apoptosis. *Am J Pathol*. 2006; 169:119-31.
- [248] Kichev A, Ilieva EV, Piñol-Ripoll G, Podlesniy P, Ferrer I, Portero-Otín M, Pamplona R, Espinet C. Cell death and learning impairment in mice caused by in vitro modified

pro-NGF can be related to its increased oxidative modifications in Alzheimer disease. *Am J Pathol.* 2009; 175: 2574-85.

- [249] Eriksdotter Jönhagen M, Nordberg A, Amberla K, Bäckman L, Ebendal T, Meyerson B, Olson L, Seiger, Shigeta M, Theodorsson E, Viitanen M, Winblad B, Wahlund LO. Intracerebroventricular infusion of nerve growth factor in three patients with Alzheimer's disease. *Dement Geriatr Cogn Disord.* 1998; 9: 246-57.
- [250] Cattaneo A, Capsoni S, Paoletti F. Towards noninvasive nerve growth factor therapies for Alzheimer's disease. *J Alzheimers Dis.* 2008; 15:255-83.
- [251] Ferrer I, Marín C, Rey MJ, Ribalta T, Goutan E, Blanco R, Tolosa E, Martí E. BDNF and full-length and truncated TrkB expression in Alzheimer disease. Implications in therapeutic strategies. *J Neuropathol Exp Neurol.* 1999; 58: 729-39.
- [252] Connor B, Young D, Yan Q, Faull RL, Synek B, Dragunow M. Brain-derived neurotrophic factor is reduced in Alzheimer's disease. *Brain Res Mol Brain Res.* 1997; 49: 71-81.
- [253] Holsinger RM, Schnarr J, Henry P, Castelo VT, Fahnstock M. Quantitation of BDNF mRNA in human parietal cortex by competitive reverse transcription-polymerase chain reaction: decreased levels in Alzheimer's disease. *Brain Res Mol Brain Res.* 2000; 76: 347-54.
- [254] Hock C, Heese K, Hulette C, Rosenberg C, Otten U. Region-specific neurotrophin imbalances in Alzheimer disease: decreased levels of brain-derived neurotrophic factor and increased levels of nerve growth factor in hippocampus and cortical areas. *Arch. Neurol* 2000; 57: 846–51.
- [255] Nagahara AH, Tuszynski MH. Potential therapeutic uses of BDNF in neurological and psychiatric disorders. *Nat Rev Drug Discov.* 2011; 10: 209-19.
- [256] Nagahara AH, Merrill DA, Coppola G, Tsukada S, Schroeder BE, Shaked GM, Wang L, Blesch A, Kim A, Conner JM, Rockenstein E, Chao MV, Koo EH, Geschwind D, Masliah E, Chiba AA, Tuszynski MH. Neuroprotective effects of brain-derived neurotrophic factor in rodent and primate models of Alzheimer's disease. *Nat Med.* 2009; 15: 331-7.
- [257] Blurton-Jones M, Kitazawa M, Martinez-Coria H, Castello NA, Müller FJ, Loring JF, Yamasaki TR, Poon WW, Green KN, LaFerla FM. Neural stem cells improve cognition via BDNF in a transgenic model of Alzheimer disease. *Proc Natl Acad Sci U S A.* 2009; 106: 13594-9.
- [258] Finch CE, Laping NJ, Morgan TE, Nichols NR, Pasinetti GM. TGF- β 1 is an organizer of response to neurodegeneration. *J Cell Biochem.* 1993; 53: 314–22.
- [259] Brionne TC, Tesseur I, Masliah E, Wyss Coray T. Loss of TGF- β 1 leads to increased neuronal cell death and microgliosis in mouse brain. *Neuron.* 2003; 40:1133–45.

- [260] Caraci F, Spampinato S, Sortino MA, Bosco P, Battaglia G, Bruno V, Drago F, Nicoletti F, Copani A. Dysfunction of TGF- β 1 signaling in Alzheimer's disease: perspectives for neuroprotection. *Cell Tissue Res.* 2012 347;; 291-301.
- [261] Caraci F, Battaglia G, Busceti C, Biagioni F, Mastroiacovo F, Bosco P, Drago F, Nicoletti F, Sortino MA, Copani A. TGF-beta 1 protects against Abeta-neurotoxicity via the phosphatidylinositol-3-kinase pathway. *Neurobiol Dis.* 2008; 30: 234-42.
- [262] Sortino MA, Chisari M, Merlo S, Vancheri C, Caruso M, Nicoletti F, Canonico PL, Copani A. Glia mediates the neuroprotective action of estradiol on beta-amyloid-induced neuronal death. *Endocrinology.* 2004; 145: 5080-6.
- [263] Bruno V, Battaglia G, Casabona G, Copani A, Caciagli F, Nicoletti F. Neuroprotection by glial metabotropic glutamate receptors is mediated by transforming growth factor-beta. *J Neurosci.* 1998; 18: 9594-600.
- [264] Caraci F, Battaglia G, Bruno V, Bosco P, Carbonaro V, Giuffrida ML, Drago F, Sortino MA, Nicoletti F, Copani A. TGF-beta1 pathway as a new target for neuroprotection in Alzheimer's disease. *CNS Neurosci Ther.* 2009; 17: 237-249.
- [265] Vollmar P, Haghighia A, Dermietzel R, Faustmann PM. Venlafaxine exhibits an anti-inflammatory effect in an inflammatory co-culture model. *Int J Neuropsychopharmacol.* 2008; 11:111-7.
- [266] Arnon R, Aharoni R. Mechanism of action of glatiramer acetate in multiple sclerosis and its potential for the development of new applications. *Proc Natl Acad Sci USA.* 2004; 101 (Suppl 2): 14593-8.
- [267] Zhang H, Zou K, Tesseur I, Wyss-Coray T. Small molecule tgf-beta mimetics as potential neuroprotective factors. *Curr Alzheimer Res.* 2005; 2: 183-6.
- [268] Scheibel AB, Tomiyasu U. Dendritic sprouting in Alzheimer's presenile dementia. *Exp Neurol.* 1978; 60:1-8.
- [269] Ferrer I, Aymami A, Rovira A, Grau Veciana JM. Growth of abnormal neurites in atypical Alzheimer's disease. A study with the Golgi method. *Acta Neuropathol.* 1983; 59:167-70.
- [270] Arendt T. Synaptic degeneration in Alzheimer's disease. *Acta Neuropathol.* 2009; 118: 167-79.
- [271] Barnett A, Brewer GJ. Autophagy in aging and Alzheimer's disease: pathologic or protective? *J Alzheimers Dis.* 2011; 25: 385-94.
- [272] Harris H, Rubinsztein DC. Control of autophagy as a therapy for neurodegenerative disease. *Nat Rev Neurol.* 2011; 8: 108-17.
- [273] Nixon RA, Wegiel J, Kumar A, Yu WH, Peterhoff C, Cataldo A, Cuervo AM. Extensive involvement of autophagy in Alzheimer disease: an immuno-electron microscopy study. *J Neuropathol Exp Neurol.* 2005; 64: 113-22.

- [274] Barrachina M, Maes T, Buesa C, Ferrer I. Lysosome-associated membrane protein 1 (LAMP-1) in Alzheimer's disease. *Neuropathol Appl Neurobiol.* 2006; 32: 505-16.
- [275] Li L, Zhang X, Le W. Autophagy dysfunction in Alzheimer's disease. *Neurodegener Dis.* 2010; 7: 265-71.
- [276] Moreira PI, Santos RX, Zhu X, Lee HG, Smith MA, Casadesus G, Perry G. Autophagy in Alzheimer's disease. *Expert Rev Neurother.* 2010; 10: 1209-18.
- [277] Lee S, Sato Y, Nixon RA. Primary lysosomal dysfunction causes cargo-specific deficits of axonal transport leading to Alzheimer-like neuritic dystrophy. *Autophagy.* 2011; 7: 1562-3.
- [278] Sanchez-Varo R, Trujillo-Estrada L, Sanchez-Mejias E, Torres M, Baglietto-Vargas D, Moreno-Gonzalez I, De Castro V, Jimenez S, Ruano D, Vizuite M, Davila JC, Garcia-Verdugo JM, Jimenez AJ, Vitorica J, Gutierrez A. Abnormal accumulation of autophagic vesicles correlates with axonal and synaptic pathology in young Alzheimer's mice hippocampus. *Acta Neuropathol.* 2012; 123: 53-70.
- [279] Nixon RA, Yang DS. Autophagy failure in Alzheimer's disease--locating the primary defect. *Neurobiol Dis.* 2011; 43: 38-45.
- [280] Caccamo A, Majumder S, Richardson A, Strong R, Oddo S. Molecular interplay between mammalian target of rapamycin (mTOR), amyloid-beta, and Tau: effects on cognitive impairments. *J Biol Chem.* 2010; 285: 13107-20.
- [281] Spilman P, Podlutskaya N, Hart MJ, Debnath J, Gorostiza O, Bredesen D, Richardson A, Strong R, Galvan V. Inhibition of mTOR by rapamycin abolishes cognitive deficits and reduces amyloid-beta levels in a mouse model of Alzheimer's disease. *PLoS One.* 2010; 5: e9979.
- [282] Majumder S, Richardson A, Strong R, Oddo S. Inducing autophagy by rapamycin before, but not after, the formation of plaques and tangles ameliorates cognitive deficits. *PLoS One.* 2011; 6: e25416.
- [283] Yang DS, Stavrides P, Mohan PS, Kaushik S, Kumar A, Ohno M, Schmidt SD, Wesson D, Bandyopadhyay U, Jiang Y, Pawlik M, Peterhoff CM, Yang AJ, Wilson DA, St George-Hyslop P, Westaway D, Mathews PM, Levy E, Cuervo AM, Nixon RA. Reversal of autophagy dysfunction in the TgCRND mouse model of Alzheimer's disease ameliorates amyloid pathologies and memory deficits. *Brain.* 2011; 134: 258-77.
- [284] Laplante M, Sabatini DM. mTOR signaling in growth control and disease. *Cell.* 2012; 149: 274-93.
- [285] Sarkar S, Floto RA, Berger Z, Imarisio S, Cordenier A, Pasco M, Cook LJ, Rubinsztein DC. Lithium induces autophagy by inhibiting inositol monophosphatase. *J Cell Biol.* 2005; 170:1101-11.
- [286] Williams A, Sarkar S, Cuddon P, Ttofi EK, Saiki S, Siddiqi FH, Jahreiss L, Fleming A, Pask D, Goldsmith P, O'Kane CJ, Floto RA, Rubinsztein DC. Novel targets for Hun-

tington's disease in an mTOR-independent autophagy pathway. *Nat Chem Biol.* 2008; 4: 295-305.

- [287] Frautschy SA, Cole GM. Why pleiotropic interventions are needed for Alzheimer's disease. *Mol Neurobiol.* 2010; 41: 392-409.
- [288] Bajda M, Guzior N, Ignasik M, Malawska B. Multi-target-directed ligands in Alzheimer's disease treatment. *Curr Med Chem.* 2011; 18: 4949-75.
- [289] Hashimoto M, Hossain S. Neuroprotective and ameliorative actions of polyunsaturated fatty acids against neuronal diseases: beneficial effect of docosahexaenoic acid on cognitive decline in Alzheimer's disease. *J Pharmacol Sci.* 2011; 116: 150-62.
- [290] Fredholm BB, Battig K, Holmen J, Nehlig A, Zvartau EE. Actions of caffeine in the brain with special reference to factors that contribute to its widespread use. *Pharmacol Rev.* 1999; 51: 83-133.
- [291] Marques S, Batalha VL, Vaqueiro Lops L, Fleming Outeiro T. Modulating Alzheimer's disease through caffeine: a putative link to epigenetics. *J Alzheimers Dis* 2011; 24: 161-71.
- [292] Arendash GW, Scleif W, Rezai-Zadeh K, Jackson EK, Zacharia LC, Cracchiolo JR, Shippy D, Tan J. Caffeine protects Alzheimer's mice against cognitive impairment and reduces brain beta-amyloid production. *Neuroscience* (2006) 142: 941-52.
- [293] Dall'Igna OP, Fett P, Gomes MW, Souza DO, Cunha RA, Lara DR. Caffeine and adenosine A (2a) receptor antagonists prevent beta-amyloid (25-30)-induced cognitive deficits in mice. *Exp Neurol.* 2007; 203: 241-5.
- [294] Ritchie K, Carriere I, de Mendonca A, Portet F, Dartigues JF, Rouaud O, Barberger-Gateau P, Ancelin ML. The neuroprotective effects of caffeine: a prospective population study. *Neurology.* 2007; 69: 536-45.
- [295] Santos C, Lunet N, Azevedo A, de Mendonca A, Richtie K, Barros H. Caffeine intake and dementia: systematic review and meta-analysis. *J Alzheimers Dis.* 2010; suppl 1: S187-204.
- [296] Eskelinen MH, Kivipelto M. Caffeine as a protective factor in dementia and Alzheimer's disease. *J Alzheimers Dis* 2010; 20 suppl 1: S167-74.
- [297] Singh M, Dykens JA, Simpkins JW. Novel mechanisms for estrogen-induced neuroprotection. *Exp Biol Med.* 2006; 231: 514-21.
- [298] Pike CJ, Carroll JC, Rosario ER, Barron AM. Protective actions of sex steroid hormones in Alzheimer's disease. *Front Neuroendocrinol.* 2009; 30: 239-58.
- [299] Correia SC, Santos RX, Cardoso S, Carvalho C, Santos MS, Oliveira CR, Moreira PI. Effects of estrogen in the brain: is it a neuroprotective agent in Alzheimer's disease? *Curr Aging Sci.* 2010; 3:113-26.

- [300] Alonso A, Gonzalez C. Neuroprotective role of estrogens: relationship with insulin/IGF-1 signaling. *Front Biosci.* 2012; 4: 607-19.
- [301] Carroll JC, Rosario ER, Chang L, Stanczyk FZ, Oddo S, LaFerla FM, Pike CJ. Progesterone and estrogen regulate Alzheimer-like neuropathology in female 3xTg-AD mice. *J Neurosci.* 2007; 27:13357-65.
- [302] Fillit H, Weinreb H, Cholst I, Luine V, McEwen B, Amador R, Zabriskie J. Observations in a preliminary open trial of estradiol therapy for senile dementia-Alzheimer's type. *Psychoneuroendocrinology.* 1986; 11: 337-45.
- [303] Craig MC, Murphy DG. Estrogen therapy and Alzheimer's dementia. *Ann N Y Acad Sci.* 2010; 1205: 245-53.
- [304] Campbell VA, Gowran A. Alzheimer's disease: taking the edge off with cannabinoids? *Br J Pharmacol* 2007; 152: 655-62.
- [305] Van Der Stelt M, Mazzola C, Esposito G, Mathias I, Petrosino S, De Filippis D et al. Endocannabinoids and β -amyloid-induced neurotoxicity in vivo: effect of pharmacological elevation of endocannabinoid levels. *Cell Mol Life Sci.* 2006; 63: 1410-24.
- [306] Aso E, Palomer E, Juvés S, Maldonado R, Muñoz FJ, Ferrer I. CB1 agonist ACEA protects neurons and reduces the cognitive impairment of A β PP/PS1 mice. *J Alzheimers Dis.* 2012; 30: 439-59.
- [307] Ramírez BG, Blázquez C, Gómez del Pulgar T, Guzmán M, de Ceballos ML. Prevention of Alzheimer's disease pathology by cannabinoids: neuroprotection mediated by blockade of microglial activation. *J Neurosci.* 2005; 25:1904-13.
- [308] Martín-Moreno AM, Brera B, Spuch C, Carro E, García-García L, Delgado M, Pozo MA, Innamorato NG, Cuadrado A, Ceballos ML. Prolonged oral cannabinoid administration prevents neuroinflammation, lowers β -amyloid levels and improves cognitive performance in Tg APP 2576 mice. *J Neuroinflammation.* 2012; 9:8.
- [309] Esposito G, De Filippis D, Carnuccio R, Izzo AA, Iuvone T. The marijuana component cannabidiol inhibits β -amyloid-induced tau protein hyperphosphorylation through Wnt/ β -catenin pathway rescue in PC12 cells. *J Mol Med* 2006; 84: 253-8.
- [310] Galve-Roperh I, Aguada T, Palazuelos J, Guzman M. The endocannabinoid system and neurogenesis in health and disease. *Neuroscientist* 2007; 13: 109-14.
- [311] Harvey BS, Ohlsson KS, Mååg JL, Musgrave IF, Smid SD. Contrasting protective effects of cannabinoids against oxidative stress and amyloid- β evoked neurotoxicity in vitro. *Neurotoxicology.* 2012; 33: 138-46.
- [312] Noonan J, Tanveer R, Klompas A, Gowran A, McKiernan J, Campbell VA. Endocannabinoids prevent β -amyloid-mediated lysosomal destabilization in cultured neurons. *J Biol Chem.* 2010; 285: 38543-54.

- [313] Eubanks LM, Rogers CJ, Beuscher AE 4th, Koob GF, Olson AJ, Dickerson TJ, Janda KD. A molecular link between the active component of marijuana and Alzheimer's disease pathology. *Mol Pharm.* 2006; 3: 773-7.
- [314] Sirén AL, Fasshauer T, Bartels C, Ehrenreich H. Therapeutic potential of erythropoietin and its structural or functional variants in the nervous system. *Neurotherapeutics.* 2009; 6: 108-27.
- [315] Sun ZK, Yang HQ, Pan J, Zhen H, Wang ZQ, Chen SD, Ding JQ. Protective effects of erythropoietin on tau phosphorylation induced by beta-amyloid. *J Neurosci Res.* 2008; 86: 3018-27.
- [316] Li G, Ma R, Huang C, Tang Q, Fu Q, Liu H, Hu B, Xiang J. Protective effect of erythropoietin on beta-amyloid-induced PC12 cell death through antioxidant mechanisms. *Neurosci Lett.* 2008; 442:143-7.
- [317] Adamcio B, Sargin D, Stradomska A, Medrihan L, Gertler C, Theis F, Zhang M, Müller M, Hassouna I, Hannke K, Sperling S, Radyushkin K, El-Kordi A, Schulze L, Ronnenberg A, Wolf F, Brose N, Rhee JS, Zhang W, Ehrenreich H. Erythropoietin enhances hippocampal long-term potentiation and memory. *BMC Biol.* 2008; 6:37.
- [318] Ponce LL, Navarro JC, Ahmed O, Robertson CS. Erythropoietin neuroprotection with traumatic brain injury. *Pathophysiology.* 2012 Mar 13.
- [319] Shepardson NE, Shankar GM, Selkoe DJ. Cholesterol level and statin use in Alzheimer disease: I. Review of epidemiological and preclinical studies. *Arch Neurol.* 2011; 68: 1239-44.
- [320] Pac-Soo C, Lloyd DG, Vizcaychipi MP, Ma D. Statins: the role in the treatment and prevention of Alzheimer's neurodegeneration. *J Alzheimers Dis.* 2011; 27: 1-10.
- [321] Chauhan NB, Siegel GJ, Feinstein DL. Effects of lovastatin and pravastatin on amyloid processing and inflammatory response in TgCRND8 brain. *Neurochem Res.* 2004; 29: 1897-911.
- [322] Li L, Cao D, Kim H, Lester R, Fukuchi K. Simvastatin enhances learning and memory independent of amyloid load in mice. *Ann Neurol.* 2006; 60: 729-39.
- [323] Weinreb O, Amit T, Bar-Am O, Youdim MB. Ladostigil: a novel multimodal neuroprotective drug with cholinesterase and brain-selective monoamine oxidase inhibitory activities for Alzheimer's disease treatment. *Curr Drug Targets.* 2012; 13: 483-94.
- [324] Wang R, Yan H, Tang XC. Progress in studies of huperzine A, a natural cholinesterase inhibitor from Chinese herbal medicine. *Acta Pharmacol Sin.* 2006; 27:1-26.
- [325] Peng Y, Lee DY, Jiang L, Ma Z, Schachter SC, Lemere CA. Huperzine A regulates amyloid precursor protein processing via protein kinase C and mitogen-activated protein kinase pathways in neuroblastoma SK-N-SH cells over-expressing wild type human amyloid precursor protein 695. *Neuroscience.* 2007; 150: 386-95.

- [326] Ved HS, Koenig ML, Dave JR, Doctor BP. Huperzine A, a potential therapeutic agent for dementia, reduces neuronal cell death caused by glutamate. *Neuroreport*. 1997; 8: 963-8.
- [327] Li J, Wu HM, Zhou RL, Liu GJ, Dong BR. Huperzine A for Alzheimer's disease. *Cochrane Database Syst Rev*. 2008; 16:CD005592.
- [328] Huang TC, Lu KT, Wo YY, Wu YJ, Yang YL. Resveratrol protects rats from A β -induced neurotoxicity by the reduction of iNOS expression and lipid peroxidation. *PLoS One*. 2011; 6: e29102.
- [329] Li F, Gong Q, Dong H, Shi J. Resveratrol, a neuroprotective supplement for Alzheimer's disease. *Curr Pharm Des*. 2012; 18: 27-33.
- [330] Davinelli S, Sapere N, Zella D, Bracale R, Intrieri M, Scapagnini G. Pleiotropic protective effects of phytochemicals in Alzheimer's disease. *Oxid Med Cell Longev*. 2012; 2012:386527.
- [331] Allison AC, Cacabelos R, Lombardi VR, Alvarez XA, Vigo C. Celastrol, a potent antioxidant and anti-inflammatory drug, as a possible treatment for Alzheimer's disease. *Neuropsychopharmacol Biol Psychiatry*. 2001; 25: 1341-1357.
- [332] Paris D, Ganey NJ, Laporte V, Patel NS, Beaulieu-Abdelahad D, Bachmeier C, March A, Ait-Ghezala G, Mullan MJ. Reduction of beta-amyloid pathology by celastrol in a transgenic mouse model of Alzheimer's disease. *J Neuroinflammation*. 2010;7: 17.
- [333] Murray ME, Graff-Radford NR, Ross OA, Petersen RC, Duara R, Dickson DW. Neuropathologically defined subtypes of Alzheimer's disease with distinct clinical characteristics: a retrospective study. *Lancet Neurol*. 2011; 10: 785-796.
- [334] Reiman EM, Langbaum JB, Fleisher AS, Caselli RJ, Chen K, Ayutyanont N, Quiroz YT, Kosik KS, Lopera F, Tariot PN. Alzheimer's Prevention Initiative: a plan to accelerate the evaluation of presymptomatic treatments. *J Alzheimers Dis*. 2011; 26 Suppl 3:321-329.
- [335] Henderson ST, Poirier J. Pharmacogenetic analysis of the effects of polymorphisms in APOE, IDE and IL1B on a ketone body based therapeutic on cognition in mild to moderate Alzheimer's disease; a randomized, double-blind, placebo-controlled study. *BMC Med Genet*. 2011; 12:137.