We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

siRNA and Gene Formulation for Efficient Gene Therapy

Ian S. Blagbrough and Abdelkader A. Metwally

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/55518>

1. Introduction

Whilst small interfering RNA (siRNA, also known as short interfering RNA) has a somewhat chequered history with regard to its discovery and initial usage, the “mammalian” research community singularly neither reading nor citing the output from the “plant” research community, it is now recognised in terms of \$bn being invested and spent that RNA interference (RNAi), sequence specific post-transcriptional gene silencing (PTGS) by siRNA, has many potential therapeutic applications [1] as well as being an important tool in the study of functional genomics. The site and mechanism of action of siRNA requires that these short double-stranded nucleic acids are delivered to the cytosol of target cells. Therefore, formulation is required in a strategy similar to that for gene therapy, although not requiring access to the nucleus. Efficient medicines design should come with an understanding of the problem at the molecular level. Our contributions are aimed at the use of non-viral gene therapy and this Chapter therefore has such a focus.

2. RNA interference

2.1. History and mechanism of RNA interference

siRNA is a double-stranded RNA (dsRNA) typically of 21-25 nucleotides per strand. siRNA operates as a part of the cellular mechanism called RNAi, which was first noticed in petunia flowers (*Petunia hybrida*) which showed reduced pigmentation on the introduction of exogenous genes that were meant to increase pigmentation [2, 3]. These experiments aimed at increasing the pigmentation of the petunia flowers by means of introducing additional gene constructs expressing either chalcone synthase [2, 3] or dihydroflavonol-4-reductase [2]. However, the resultant plants produced completely white flowers and/or flowers with white

or pale sectors on a pigmented background. The exact mechanism was not identified at the time and was simply termed co-suppression. The transcription level of the suppressed chalcone synthase genes in petunia flowers was found to be similar to that of the non-suppressed genes, and thus the co-suppression must have been at the post-transcriptional level [4]. Later in 1997, the suppression of chalcone synthase endogene in petunia flowers was suggested to be related to formation of RNA duplexes by intermolecular pairing of complementary sequences between the coding sequence and the 3'-UTR sequence of the transgene mRNA [5]. Indeed, the seminal contributions the plant RNAi community have made to this RNAi field are also reflected in the research of Hamilton and Sir David C. Baulcombe in the Sainsbury Laboratory, Norwich, UK, on PTGS as a nucleotide sequence-specific defence mechanism that can target both cellular and viral mRNAs with RNA molecules of a uniform length, ~25 nucleotides [6]. That RNA silencing involves the processing of dsRNA into 21-26 long siRNA to mediate gene suppression (correspondingly complementary to the dsRNA) was demonstrated in *Arabidopsis*, "RNA silencing pathways in plants that may also apply in animals" [7]. That *Arabidopsis* ARGONAUTE1 RNA-binding protein is an RNA slicer that selectively recruits microRNAs and siRNAs was shown to be by a key mechanism similar to but different from that found in animals [8]. In 1998, Fire, Mello and co-workers reported the reduction or inhibition (hence genetic "interference") of the expression of the *unc-22* gene in *Caenorhabditis elegans* by means of dsRNA that is homologous to 742 nucleotides in the targeted gene [9], a discovery that was awarded the Nobel Prize in medicine or physiology in 2006. The target gene expresses an abundant although nonessential myofilament protein. Decreasing *unc-22* activity resulted in an increasingly severe twitching phenotype, while complete inhibition resulted in impaired motility and muscle structural defects. The target gene inhibition was best achieved with dsRNA, while using the individual sense or anti-sense RNA strands resulted only in modest silencing. The authors also noticed that only few copies of the dsRNA are required per cell to initiate a potent and specific response, rejecting the hypothesis that the mechanism of interaction with target gene mRNA is stoichiometric in nature, and thus the role of the dsRNA in the interference machinery must be catalytic or amplifying.

Elbashir et al. reported in 2001 that sequence-specific gene silencing of endogenous and heterologous genes with 21 nucleotide siRNA occurs in mammalian cell cultures [10]. The reporter genes coding for sea pansy (*Renilla reniformis*) and firefly (*Photinus pyralis*) luciferases were silenced successfully in different cell lines including human embryonic kidney cells (293) and the cervix cancer cells (HeLa cell line, the first human cell line grown in vitro with success [11]), as well as the endogenous gene coding for the nuclear envelope proteins lamin A and lamin C in HeLa cells. The authors used dsRNA of length 21 or 22 nucleotides with 3'-symmetrical 2-nucleotide overhangs on each strand, as dsRNA with length >30 nucleotides initiates an immune response e.g. inducing interferon synthesis) that leads to non-specific mRNA degradation, which was evident from non-specific silencing of luciferase with 50 and 500 nucleotides dsRNA in HeLa S3 cells, COS-7 cells (kidney cells of the African green monkey), and NIH/3T3 cells (mouse fibroblasts) [10]. The RNAi mechanism of action continues to be investigated in detail and reviewed thoroughly [12-17]. The RNAi mechanism involves the incorporation of dsRNA segments (e.g. siRNA) that have a sequence complementary to the targeted mRNA in a protein com-

plex. This core complex which carries-out mRNA degradation is the RNA induced silencing complex (RISC) [18-20]. The degradation process requires the key argonaute family of proteins, which contain a domain with RNase H (endonuclease) type of activity that catalyse cleavage of the phosphodiester bonds of the targeted mRNA. RISC assembly and subsequently its function to mediate sequence specific mRNA degradation occur in the cytoplasm of the cell [16]. The source of the dsRNA segments incorporated in RISC can be endogenously processed microRNA (miRNA), short hairpin RNA (shRNA), or synthetic siRNA. miRNA is produced from endogenous DNA through the action of RNA polymerase II resulting in the formation of non-coding RNA called primary miRNA (pri-miRNA), which is processed in the nucleus by a protein complex containing an enzyme known as Drosha and a dsRNA binding protein cofactor called Pasha (DGCR8). Drosha cleaves pri-miRNA to produce (pre-miRNA), a dsRNA of 70-90 nucleotides and having a hairpin loop, which binds to Exportin 5 protein and is transferred from the nucleus into the cytoplasm. Pre-miRNA is processed by Dicer (RNase III enzyme) in the cytoplasm to give miRNA, typically of 22 nucleotides in length and having two nucleotide overhangs at the 3'-position [16, 21], shRNA is produced by transcription from an exogenous DNA that is delivered to the nucleus, and codes for a hairpin shaped RNA with segments of length 19-29 nucleotides and loop of 9 nucleotides [22, 23] which can then be processed by Dicer and incorporated in the RNAi machinery.

Once in the cytoplasm, the processed dsRNA (miRNA, processed shRNA, or siRNA) is then incorporated into a protein complex (RISC-loading complex, RLC). In *Drosophila* the RLC is composed of the dsRNA, heterodimer protein DCR2 (Dicer variant)/R2D2, possibly including the catalytic argonaute proteins as well in this complex. The active RISC is formed when one of the RNA strands in the complex is cleaved (the passenger strand) and the strand with the less thermodynamic stable 5'-end (guide/anti-sense strand) remains in the complex. The mRNA with complementary sequence to the guide strand binds to the active RISC and is cleaved by the endoribonuclease activity of the argonaute component of the complex (Figure 1).

2.2. RNA duplex structure

RNA is a polymer of ribonucleotides. Each RNA nucleotide is composed of one nucleobase, the monosaccharide pentose ribose, and one phosphate group. The nucleobases in RNA are adenine (purine base), guanine (purine base), uracil (pyrimidine base), and cytosine (pyrimidine base) (Figure 2). A nucleoside is formed when each base is connected via a glycosidic bond to the anomeric carbon 1' of ribose, thus when glycosylated, adenine, guanine, uracil, and cytosine nucleobases give adenosine, guanosine, uridine, and cytidine nucleosides. Each two nucleosides are connected via a phosphate diester bond between the 3' of one nucleoside and 5' of the next nucleoside to form the RNA polynucleotide strand. The main differences in the primary structure of RNA and DNA are that RNA pentose is ribose while DNA pentose is 2'-deoxyribose, and the RNA incorporates the nucleobase uracil instead of thymine.

Figure 1. RNAi mechanism in a eukaryotic cell. The source of the antisense strand incorporated in RISC can be miRNA, processed exogenous long dsRNA, or synthetic siRNA delivered to the cell.

Figure 2. Nucleobases and pentoses of RNA and DNA.

Figure 3. siRNA duplex 22-mer targeting the enhanced green fluorescent protein (EGFP) mRNA. The two deoxythymidine residues at the 3'-end of the sense strand are not shown. Sense strand: 5'-GCAAGCUGACCCUGAAGUUCAUTT-3' Anti-sense strand: 5'-AUGAACUUCAGGGUCAGCUUGCCG-3' Target DNA sequence: 5'-CGGCAAGCTGACCCTGAAGTTCAT-3'

In order to form an RNA duplex (Figure 3), the strands with complementary nucleotide sequence bind together by hydrogen bonds. Adenine is bound to uracil with two hydrogen bonds while guanine is bound to cytosine with three hydrogen bonds, thus forming what is known as Watson-Crick base pairs. RNA duplexes under normal physiological conditions are in the form of A-helix. This type of duplex is a right-handed helix [24-26].

The presence of the 2'-hydroxyl group of the ribose and the lack of the methyl group on the nucleotide uridine (in contrast to the methylated thymidine) results in structural differences between RNA and DNA, with the 2'-hydroxyl group of RNA being the major cause of the differences. The sugar phosphate backbone of RNA duplexes is stabilized by the 2'-hydroxyl in the C3'-endo position, while DNA adopts the C2'-endo position (Figure 4). Thus, the RNA duplex takes the A-helix form while the DNA helix takes the B-form. The A-helix form is suggested to have a greater hydration shell, giving RNA duplexes more thermodynamic stability and more rigidity compared to DNA duplexes [24-26]. RNA A-helix completes one complete rotation in 11-12 base pair (bp) compared to 10 bp for DNA, with a rise of 2.7 Å per bp of RNA [27]. The A-helix geometry has been suggested to be the major factor explaining why dsRNA and not dsDNA is involved in the RNAi machinery [28], where the A-helix geometry between the guide strand and the complementary target mRNA is essential for the catalytic activity of the argonaute 2 protein in the RISC.

As a result of the presence of a hydroxyl group in the 2'-position of the ribose in the RNA backbone, the RNA phosphodiester backbone is more susceptible to hydrolysis by nucleases compared to the DNA which lacks the 2'-hydroxyl in its 2'-deoxyribose [29]. Incubation of

Figure 4. 3'-endo ribose configuration of RNA (left) vs 2'-endo (right) of 2'-deoxyribose in DNA. Shown is cytidine (RNA) and deoxycytidine (DNA) with the 3'-hydroxyl phosphorylated. The hydrogen atoms at C2' and C3' are not displayed for clarity.

siRNA in fetal bovine or human serum at 37 °C resulted in the degradation and partial or complete loss of activity [30]. When incubated in human plasma at 37 °C, more than 50% of the unmodified siRNA was degraded within one minute, and practically all siRNA was completely degraded within 4 hours [31]. Although Ribonuclease A (RNase A, an endoribonuclease) cleaves single stranded RNA, siRNA degradation in serum was reported to be mainly due to RNase-like activity[32], which is suggested to occur during transient breaking of the hydrogen bonds joining the two siRNA strands. In addition to the RNase A family of enzymes, blood serum contains phosphatases and exoribonucleases which can also affect degradation of siRNA at nuclease-sensitive sites on both strands [33].

2.3. Therapeutic potential of RNAi based therapies

RNAi based therapies emerged in the period following its discovery in 1998 in plants, and are promising therapeutic candidates to treat various types of diseases, ranging from age related macular oedema to respiratory tract infections to various types of cancer [34-36]. In addition to siRNA based therapies, shRNA [37, 38] and miRNA [39] are potential therapeutic tools. siRNA based therapeutics are already in phase I and phase II clinical trials; representative examples of clinical trials involving siRNA are shown in Table 1. The basic concept is the reduction or inhibition of the expression of a protein that is involved in the pathophysiological pathway of the target disease (silencing/knocking-down the target gene). This concept is evident from using Cand5 siRNA targeting the mRNA translating the vascular endothelial growth factor (VEGF), thus reducing/inhibiting angiogenesis and preventing progression of wet age related macular oedema (Table 1) [40]. Atu027 siRNA targets the biosynthesis of protein kinase N3 which plays a role in cancer metastasis [41].

siRNA	Disease	Vector/ Route	Phase	Sponsor
Cand5/ Bevasiranib	Diabetic macular oedema	None/ <i>Intravitreal</i>	Phase II	Opko Health (Miami, USA)
Cand5/ Bevasiranib	Age-related macular degeneration	None/ <i>Intravitreal</i>	Phase II (Phase III halted)	Opko Health (Miami, USA)
ALN-RSV01	Respiratory syncytial virus infection	None/ <i>Intranasal</i>	Phase II	Alnylam Pharmaceuticals (Cambridge, USA)
CALAA-01	Solid tumour/melanoma	Cyclodextrin nanoparticles/ <i>Intravenous</i>	Phase I	Calando (Pasadena, CA, USA)
Atu027	Colorectal cancer metastasizing to the liver	AtuPlex- Liposome/ <i>Intravenous</i>	Phase I	Silence Therapeutics (London, UK)
Two siRNA against TGFBI and COX-2 STP705	Wound healing	Nanoparticles/ <i>Intravenous</i>	Phase I	Sirnaomics (Gaithersburg, MD, USA)
I5NP	Protection from acute kidney injury after cardiac bypass surgery	None/ <i>Intravenous</i>	Phase I	Quark Pharmaceuticals (Fremont, USA)
TKM-080301	Against PLK1 gene product in patients with hepatic cancer	Lipid nanoparticles/ <i>Hepatic intra- arterial administration</i>	Phase I	NCI (Maryland, USA)

Table 1. Representative clinical trials using siRNA (<http://clinicaltrials.gov/ct2/home>, accessed on 5/8/2012).

The therapeutic application of siRNA requires overcoming several barriers (Figure 5) for its intracellular delivery and the subsequent functional gene silencing activity [42-44]. Those barriers are mainly due to siRNA specific characteristics, most important are having a highly negative charge due to their phosphate backbone (on average 40-50 negative charges per siRNA), being susceptible to degradation by nucleases, and having relatively large molecular weight (13-15 kDa) compared to conventional small drug molecules. First, local delivery (such as intravitreal) is different from intravenous delivery, where the latter will subject the siRNA to the serum ribonucleases, which results in degrading non-modified siRNA within time periods that vary from minutes to hours [31]. siRNA injected intravenously in rats was reported to be cleared rapidly from circulation and accumulates in kidneys within minutes of injection [45], making it useful only if the target organ is the kidney.

In order to gain access into the cytoplasm where siRNA can exert its biological activity, the polyribonucleotide must pass first through the interstitial space then through the target cell membrane. This will be a difficult task, since both the extracellular matrix in many tissue types

and the cell membrane incorporate negatively charged glycosaminoglycans (e.g. heparan sulfate) [46]. In addition, cell membranes contain negatively charged phospholipids (e.g. phosphatidyl serine) therefore the membrane is negatively charged [46, 47]. The net result is an unfavourable repulsive interaction with naked siRNA. As a result, different strategies are being developed to overcome the barriers to reproducible and functional siRNA delivery, and these approaches fall into two general categories. One category is modifying the siRNA, the other is deploying a vector to protect the siRNA and increase its efficiency of delivery.

Figure 5. Summary of barriers to successful gene-silencing mediated by siRNA after intravenous injection, whether delivered naked or incorporated in nanoparticles.

3. Strategies to achieve efficient siRNA delivery and gene silencing

3.1. siRNA modifications

siRNA modifications include those carried out at the ribose residue, at the phosphate backbone, at the RNA nucleotides, the siRNA termini, and/or by conjugation of other molecules to the siRNA molecule. Modifications to the ribose at the 2'-position are common [48], and include 2'-O-alkylation (e.g. 2'-O-methyl and 2'-O-methylethoxy) modifications. 2'-Fluoro RNA is

another common modification. Locked nucleic acids (LNAs) have a methylene bridge connecting the 2'-O to the 4'-C of the ribose unit, locking the sugar in the 3'-endo conformation. These modifications led to increased ribonuclease resistance [48, 49]. Modifications at the phosphate backbone include phosphorthioate, boranophosphate, and methylphosphonate linkages [48, 49] and is reported to increase siRNA stability against various ribonucleases and phosphodiesterases [50]. siRNA nucleotides can be substituted with DNA nucleotides to increase stability and/or decrease unwanted siRNA off-target effects [51]. Modifications of the 3'-overhangs (usually two nucleotides in length) include incorporating deoxyribonucleotides to reduce costs and increase stability towards 3'-exoribonucleases. The 5'-terminus chemical phosphorylation of the antisense strand results in higher gene silencing efficiency, while blunt ended duplexes were reported to be more resistant to exonucleases. The advantages of each of the aforementioned techniques, other modification strategies, as well as the considerations related to the degree of modification and its effect on gene silencing efficiency and associated cytotoxic effects have been reviewed thoroughly [48, 52-54].

The conjugation of drug molecules, aptamers, lipids, polymers, and peptides/proteins to siRNA could enhance in vivo delivery [55]. The main aims of such conjugations are: to enhance siRNA stability, increase in vivo half-life, control biodistribution, increase efficiency of intracellular delivery, while maintaining the gene silencing activity.

One strategy is to increase the hydrophobicity of the siRNA. Cholesterol was conjugated to the 5'-terminus of siRNA, the cholesterol-siRNA conjugate (chol-siRNA) resulted in better intracellular delivery compared to unmodified siRNA and retained gene silencing activity in vitro in β -galactosidase expressing liver cells [56]. When cholesterol was conjugated to the 3'-terminus of the sense (passenger) strand of siRNA, the conjugate had improved in vivo pharmacokinetics as the intravenous administration of chol-siRNA in mice resulted in its distribution and detection in the fat tissues, heart, kidneys, liver, and lungs, even 24 h after intravenous injection [57]. No significant amounts of unmodified siRNA were detected in the tissues 24 h after the intravenous injection. Conjugation of siRNA to bile acids and long-chain fatty acids, in addition to cholesterol, mediates siRNA uptake into cells and gene silencing in vivo [58]. The medium chain fatty-acid conjugates, namely lauroyl (C12), myristoyl (C14) and palmitoyl (C16), did not silence the target apolipoprotein B mRNA levels in mouse livers after intravenous injection. However, siRNA fatty-acid conjugates having long saturated chains, stearoyl (C18) and docosanoyl (C22), significantly reduced apolipoprotein B mRNA levels.

Cell penetrating peptides (CPPs) are used to facilitate cellular membrane crossing of many molecules displaying various properties such as antisense oligonucleotides, peptides, and proteins and are already being tested in vivo [59]. siRNA was conjugated to penetratin and transportin, to silence luciferase and green fluorescent protein (GFP) in different types of mammalian cells [60]. However, in vivo lung delivery in mouse of siRNA conjugated to penetratin and TAT(48-60), targeting p38 MAP kinase mRNA showed that the reduction in gene expression was peptide induced and the penetratin conjugated siRNA resulted in innate immunity response [61].

siRNA functioning against the VEGF mRNA was conjugated to poly(ethylene glycol) (PEG, 25 kDa) via a disulfide bond at the 3'-terminus of the sense strand [62]. The siRNA-PEG

conjugate formed polyelectrolyte complex (PEC) micelles by electrostatic interaction with the cationic polymer polyethylenimine (PEI). The formed VEGF siRNA-PEG/PEI PEC micelles showed enhanced stability against nuclease degradation compared to the unmodified siRNA. These micelles efficiently silenced VEGF gene expression in prostate carcinoma cells (PC-3) and showed superior VEGF gene silencing compared to VEGF siRNA/PEI complexes in the presence of serum. PEG conjugation on its own enhanced the stability of the siRNA in serum containing medium. The prolonged stability of the PEC micelles was suggested to be due to the presence of PEG chains in the outer micellar shell layer, thus sterically hindering nuclease access into the siRNA in the micelle core [62]. Targeting molecules such as antibodies [63] and aptamers (peptides or single stranded DNA or RNA that have selective affinities toward target proteins) [64] have also been conjugated to siRNA, with the aim of increasing the efficiency of siRNA delivery to the target tissues.

Conjugating molecules to siRNA requires specific considerations. First, the site of conjugation (3'- and/or 5'-terminus, on sense and/or antisense strand) should be chosen such that it does not affect the activity of the siRNA and its ability to be incorporated in the RISC, or its ability to bind the target mRNA in the correct helix conformation. Second, the conjugated siRNA might have new properties that were not present in the unmodified parent siRNA. An example is the in vivo immune response resulting from the penetratin-siRNA conjugate [61]. Third, the conjugation process is multi-step, and the chemical reaction intermediates and products require efficient purification in order to meet the specifications of in vivo applications. These steps need to be repeated for each siRNA under investigation, which can be costly and time consuming. Thus, although there are clear advantages to synthesize siRNA conjugates, there are also disadvantages, and conjugation is therefore only one of two valuable approaches in the toolbox for preparing siRNA based therapies. The other valuable tool is complexation or incorporating the siRNA in a vector.

3.2. Viral vectors for shRNA delivery

Vectors for RNAi based therapies are either viral or non-viral vectors. Viral vectors (Table 2) are used to deliver genes encoding hairpin RNA structures such as shRNA and miRNA, which are then processed by the cellular RNAi machinery to the functional silencing dsRNA [65, 66].

Viral vectors offer two main advantages, the first is the very high efficiency compared to non-viral vectors [68], which can reach few orders of magnitude more than that achieved with non-viral vectors, and the second is the potential of long term expression of the delivered RNAi therapeutic, which is very useful in the treatment of chronic diseases such as HIV infection and viral hepatitis [69, 70]. Retroviruses are enveloped, single stranded RNA viruses and have a genome capacity of 7-10 kilobases (kb). They preferentially target dividing cells which limits their use to mitotic tissues (thus for example excluding brain and neurons). Retroviruses integrate their DNA in the host genome using an integrase enzyme, which provides the advantage of stable long term expression of the delivered transgene in the host cell and its descendants. However, integrating new DNA sequences into host genome carries the risk of

	Retrovirus/ Lentivirus	Adenovirus	Adeno- associated virus	Herpes virus
Viral vector properties	Genome	ssRNA	dsDNA	ssDNA
	Capsid	Icosahedral	Icosahedral	Icosahedral
	Envelope	Enveloped	None	Enveloped
	Viral Polymerase	Positive	Negative	Negative
	Diameter (nm)	80-130	70-90	18-26
	Genome size (kb)	7-10	38	5
Gene therapy related	Infection tropism	Dividing*	Dividing/ Non-dividing	Dividing/ Non-dividing
	Virus genome integration	Integrating	Non-integrating	Integrating
	Transgene expression	Lasting	Transient	Lasting
	Packaging capacity (kb)	7-8	8	4.5

* Lentiviral vectors can infect non-dividing cells as their pre-integration complex can traverse the nuclear membrane pores (NMP), in contrast to retrovirus pre-integration complex which does not traverse NMP, requiring the host-cell division to integrate the retroviral genome [67].

Table 2. Summary of properties of viral vectors that are commonly used in gene therapy (adapted from <http://www.genetherapy.net.com/viral-vectors.html>, accessed on 5/8/2012).

insertional mutagenesis [68, 71]. shRNA expression cassette delivered by a retroviral vector was used in rats to silence a RAS oncogene in order to suppress tumour growth [72]. Herpes virus was used successfully to deliver shRNA targeting exogenous β -galactosidase or endogenous trpv1 gene mRNA in the peripheral neurons in mice by injecting once directly into the sciatic nerve of the animals [73].

Unlike other retroviruses, lentiviruses can infect dividing as well as differentiated and non-dividing cells. The lentiviral genome can accommodate 7.5 kb [66], and their genome is integrated in the host cell genome, lentiviral vectors are generally preferred for long-term expression of transgenes, and efficient delivery in vivo to the brain, eye, and liver to induce long-term transgene expression as reported [74]. A lentiviral vector was used to deliver shRNA targeting *Smad3* gene mRNA, and enhanced myogenesis of old and injured muscles [75].

Adenoviruses are non-enveloped viruses, with linear double stranded DNA. They preferably infect the upper respiratory tract and the ocular tissue. Their genome can accommodate up to 8 kb which can be extended to ≥ 25 kb in modified viruses that have their viral genes deleted [68]. These viruses can infect post mitotic cells and thus are good candidates for neurological diseases. Unless delivering genes that can exist as episomes in host cells, adenoviruses result only in transient expression of their cargo. However, although the host cells with the episome can express the delivered genes for the cell life time, these cells will eventually be removed by the host immune system [68]. shRNA targeting VEGF that was delivered by an adenoviral vector resulted in potent inhibition of angiogenesis and tumour growth in mice [76].

Adeno-associated virus (AAV) is a single stranded DNA non-pathogenic virus that can accommodate a 4.7 kb genome. They can infect dividing or non-dividing cells. The replication of AAV requires co-infection with adenovirus. The viral genome integrates into the host cell genome at a specific location on chromosome 19 [68]. Direct intracerebellar injection in a mouse model of spinocerebellar ataxia of an AAV vector delivering a cargo expressing shRNA targeting polyglutamine induced neurodegeneration significantly restored cerebellar morphology and improved motor coordination in mice [77].

Although highly efficient in delivering their cargo, viral vectors have their disadvantages. Adenoviral vectors have the disadvantage of triggering a strong immune (adaptive and innate) response by repeated administration, in addition to target organ immunotoxicity, specially hepatotoxicity [78-80], which resulted in 1999 in the death of one 18-year-old male who received high dose of adenovirus that was delivered directly in the hepatic artery in a clinical gene therapy safety study [81]. Clonal T-cell acute lymphoblastic leukemia caused by insertional mutagenesis in a gene therapy completed clinical trial involving patients suffering X-linked severe combined immunodeficiency (SCID-X1) was reported in one out of the 10 patients using a retroviral vector [82]. Integration of the vector genome material in the antisense orientation 35 kb upstream of the protooncogene (LMO2) caused over expression of the gene in the leukemic cells. In a similar study, 4 out of 9 patients developed leukemia within 3-6 years post-treatment mainly due to vector-mediated upregulation of host cellular oncogenes [83, 84]. In addition, immune responses (whether adaptive or innate) of varying degrees depending on the type of vector, dose, and target organs were reported for lentiviral, adenoviral, adeno-associated viral vectors [80].

Current research on viral vectors for gene therapy is focussed on approaches such as vector engineering e.g. modifying the viral capsid or pseudotyping the envelope, different delivery strategies, and administration to immune-privileged sites that can tolerate the delivered viral vectors without responding with an inflammatory response [80, 85]. Other research focusses on the essential scaling-up process of vector production and increasing the packaging efficiency of the vectors [85], the processes without which, the wide spread and successful therapeutic use of the viral vectors will be very difficult to achieve.

3.3. Non-viral vectors

Non-viral vectors for gene and siRNA delivery are an alternative to viral vectors, as they do not suffer many of the disadvantages of the viral vectors, especially immunogenicity and tumourigenicity. The non-viral vectors can be classified generally as peptides, polymeric based vectors, carbohydrate based, and lipid based [86]. CPPs, also known as peptide transduction domains (PTDs), have shown the ability to cross the cellular membrane despite their relatively high molecular weight and size (Table 3).

PTDs generally are short amphipathic and/or cationic peptides that can transport many hydrophilic molecules across the cell membrane. A wide range of molecules including liposomes [87, 88], peptides, proteins [89], peptide nucleic acids [90] and polynucleotides [91] are delivered intracellularly using PTDs and they have also been applied in vivo [59, 92, 93].

CPP	Sequence of CPP	Type of association with siRNA	Target mRNA
CADY	GLWRALWRLRLSLWRLLWRA	Non-covalent	GAPDH, p53 [94]
EB1	LIRLWSHLIHIWFQNRRLKWKKK	Non-covalent	Luc [95]
MPG	GALFLGFLGAAGSTMGAWSQPKKKRKV	Non-covalent	Luc, GAPDH [96] Oct-3/4 [97]
Poly-arginine	RRRRRRRRR	Non-covalent	VEGF [98]
Penetratin	RQIKIWFQNRRMKWKK	Covalent	Luciferase (Luc), EGFP [60]
		Covalent	SOD1, caspase-3 [99]
		Covalent	Luc, p38 MAP kinase [61, 100]
Transportan	LIKKALAALAKLNILLYGASNLTWG	Covalent	Luc, EGFP [100]
TAT	GRKKRRQRRRPQ	Covalent	EGFP, CDK9 [101]

Table 3. Selected CPPs used for siRNA delivery [59].

It was reported by Frankel and Pabo in 1988 that the HIV-1 derived TAT protein could be taken up by cells growing in tissue culture [102], and that a small basic region of TAT (48-60) was essential for uptake by the cells [103]. PTDs include antennapedia homeodomain protein (Antp, penetratin), mitogen-activated protein (MAP), poly-arginine, transportan, VP22 [59, 92]. Two major pathways are involved in the uptake of PTDs and PTD-cargos: direct translocation at 4 °C and 37 °C and endocytosis-translocation at 37 °C. These mechanisms depend on many factors: cargo size, cell line, PTD concentration, and the type of PTD [59, 104, 105]. siRNA can be conjugated covalently to the CPP or can be complexed with the cationic groups of basic amino acids that are present in the backbone of the CPP. As a representative example of non-covalent complexation, CADY [94], which is basic due to its five arginine residues can complex with the negatively charged siRNA. Another example of non-covalent complexation is the poly-arginine CPP [98].

PEI (Figure 6) is an efficient, but toxic, plasmid DNA delivery vector. However, as a siRNA delivery vector PEI is reported to be much less efficient [106, 107]. This decreased efficiency is due to the dissociation of the siRNA/PEI complex upon interaction with the negatively charged cell membrane, which is suggested to be because of the short length of siRNA and the associated weak electrostatic interaction with PEI [108, 109]. Another drawback of PEI is its relatively high toxicity [110]. Thus, in addition to linear PEI, PEI polymers with a wide range of molecular weights were developed to increase PEI efficiency and/or decrease toxicity, although not all PEI are suitable for siRNA delivery [111]. The main advantage of PEI is the ability of its variety of amino groups to be protonated at lower pH (inside endosomes) leading to what is known as the “proton-sponge effect” [112], and efficient escape of the nucleic acid cargo from endosomes.

One approach to enhance siRNA delivery with PEI is increasing the hydrophobicity of PEI by covalently conjugating alkyl chains [113], where increasing the hydrophobic alkyl chain length generally improved the stability of the PEI/siRNA complex. In a similar strategy, cholesterol was conjugated to PEI with decreased toxicity of the conjugates [114]. Low molecular weight PEI (MW < 5 kDa) is less toxic than the higher molecular weight PEI (≈25 kDa), but less efficient

Figure 6. Representative examples of polyamines used in siRNA delivery either as in lipid conjugates of polyamine alkaloids e.g. spermine and spermidine, or as in polyethylenimine (PEI), a cationic polymer.

in polynucleotide delivery, thus, cross-linking of the low molecular weight PEI with disulfide bonds which are cleaved in the reducing environment of the cytoplasm increased the efficiency of siRNA delivery through the enhanced release of siRNA in the cytoplasm [115].

Chitosan is a biocompatible and biodegradable polysaccharide that is a copolymer of *N*-acetyl-D-glucosamine and D-glucosamine. Chitosan has weakly basic properties due to the presence of the D-glucosamine residue with a pK_a value 6.2-7.0. The molecular weight of chitosan affects the complex stability, size, zeta-potential and in vitro gene knock-down of siRNA/chitosan nanoparticles [116]. High molecular weight (64.8-170 kDa) chitosan formed stable complexes with siRNA and resulted in high gene knock-down efficiency in human lung carcinoma (H1299) cells, while low molecular weight (10 kDa) chitosan could not complex the siRNA into stable nanoparticles and showed almost no knock-down [117]. The method of association affects gene silencing efficiency, where chitosan-TPP/siRNA nanoparticles (siRNA entrapped inside the nanoparticles, and TPP is sodium tripolyphosphate and used as a polyanion to cross-link with the cationic chitosan groups by electrostatic interactions) showed high siRNA binding and better gene silencing in vitro compared to siRNA/chitosan particles prepared by

simple complexation and adsorption of siRNA onto chitosan [118]. Although chitosan has good potential as a non-viral gene delivery vector, widespread use is largely limited by its poor solubility (because of their pKa, chitosan amino groups are only partially protonated at the physiological pH 7.4), poor stability of its siRNA complexes at the physiological pH, and low transfection efficiency. Various strategies have been adopted to overcome these drawbacks, such as covalently conjugating PEG to chitosan and binding targeting ligands to enhance cell specificity [116].

Cyclodextrins (CD) are cyclic oligosaccharides composed of 6, 7, or 8 D-(+)-glucose units, known as α -CD, β -CD, γ -CD respectively, bound through α -1,4-linkages. Polymers conjugated to β -CD lack immunogenicity and hence are attractive vectors for polynucleotide delivery. β -CD have a hydrophilic outer surface and a hydrophobic inner cavity which enable them to form inclusion complexes. Efficient cellular transfection of siRNA labelled with a fluorescent tag into human embryonic lung fibroblasts (MRC-5 cells) was observed by siRNA complexes with the β -CD guanidine derivatized bis-(guanidinium)-tetrakis-(β -cyclodextrin) tetrapod (having four β -CD units) [119]. The ability of β -CD to form inclusion complexes was used to develop a siRNA delivery vector. β -CD was covalently bound to a polycationic segment (to electrostatically bind siRNA), while adamantane-PEG-transferrin (adamantane can fit in the β -CD cavity) formed an inclusion complex which can enhance the stability of siRNA nanoparticles in vivo [120]. This system was used to deliver siRNA silencing the *EWS-FLI1* gene thus inhibiting tumour growth in a murine model of metastatic Ewing's sarcoma. The first experimental siRNA therapeutic to provide targeted delivery in humans was reported by Davis and co-workers [121]. siRNA was formulated into a nanoparticle (CALAA-01), which consisted of a cyclodextrin-containing polymer that contains amidine and primary amine functional groups, a PEG for steric stabilization in the in vivo environment (via inclusion complexes of β -CD with adamantane-PEG conjugate), and human transferrin (Tf) as the targeting ligand to binds to the transferrin receptors that are over-expressed on cancer cells. The siRNA/nanoparticles components self-assembled in the pharmacy. CALAA-01 was administered intravenously to the first patient with a solid cancer in a phase I clinical trial (safety study) in May 2008 [121]. Tumour biopsies from patients' melanoma after treatment (phase I clinical trial) showed the presence of intracellular nanoparticles. Reductions in the levels of both the specific mRNA (M2 subunit of ribonucleotide reductase, RRM2) and the protein (RRM2) were found when compared to levels in pre-dosing tissues. These results demonstrated that siRNA nanoparticles administered systemically to a human patient can produce a specific gene knock-down via an RNAi mechanism of action [122]. A recent and novel approach to the synthesis of cationic or neutral PEGylated amphiphilic β -CD used copper-catalysed "click" chemistry to modify selectively the secondary 2-hydroxyl group of the β -CD. The 6-position of these β -CD conjugates was conjugated to a dodecane alkyl chain. Complexation of cationic β -CD alone with siRNA resulted in good silencing of the luciferase reporter gene in Caco2 cells in culture. Co-formulation of cationic β -CD with a PEGylated β -CD and siRNA resulted in lower surface charges and reduced aggregation. The transfection efficiency of the cationic β -CD vector was lowered by co-formulation with the PEGylated β -CD, although the siRNA binding was not affected and the surface charge of the complexes did not reach complete neutrality [123].

Dendrimers have a central core to which are connected several branched arms in a manner that can be symmetrical or asymmetrical. During the synthesis of dendrimers, arms (branches) are added to the core structure. Each addition is called a generation and increases the previous generation number by one. Due to their unique structure, dendrimers can have a planar, elliptical, or spherical shape depending on generation number. Among the most widely used dendrimers are polyamidoamine (PAMAM) and polypropylenimine (PPI) dendrimers [124]. Dendrimers which have positively charged cationic groups on their outer surface are commonly used for polynucleotide delivery. The transfection efficiency of dendrimers increases with increasing the charge density or generation number [125]. However, dendrimers with high generation number are generally more cytotoxic compared to dendrimers with low generation number [126]. Usually the inner space near the core is larger compared to outer space near the surface due to the lower density of molecules (less number of arms) near the core, which allow small molecules to be incorporated in the inner space. Owing to the relatively large molecular weight of polynucleotides, they are usually bound to the surface of cationic dendrimers and not in the inner space of the dendrimer. Generally, the toxicity of dendrimers is lower than that of PEI or poly-L-lysine (PLL) [127]. One advantage of dendrimers is that they have pH buffering capacity (proton-sponge effect), an important feature for endosomal escape and enhancing the release of polynucleotides [125, 128].

PPI dendrimers with high generation numbers (4 and 5) were more efficient in forming discrete nanoparticles with siRNA and in gene silencing in human lung cancer (A549) cells than lower generation dendrimers (2 and 3). Generation 5 PPI dendrimers were more toxic, probably due to the increased positive charge density per dendrimer, than generation 4 dendrimers [129]. Complex formation between PAMAM dendrimers with an ethylenediamine core and siRNA as a function of three variables has been reported [130]. The ionic strength of the medium (without or with 150 mM NaCl), the generation number (4, 5, 6 and 7) and the *N/P* ratio (ratio of positively charged amine groups per negative phosphate) were varied. The size of the complexes depended on the ionic strength of the media, with the strong electrostatic interactions in medium without NaCl making siRNA/dendrimer complexes smaller than those obtained in 150 mM NaCl. Both the intracellular delivery and the silencing of EGFP expression in cell culture was dependent on complex size, with smaller complexes efficiently delivered, and resulting in the highest silencing of EGFP expression. siRNA complexed with generation 7 dendrimers resulted in the highest silencing of EGFP expression both in human brain tumour cell line T98G-EGFP (35%) and mouse macrophage cell line J-774-EGFP (45%) cells, in spite of having lower protection of siRNA against degradation with RNase A, showing the importance of formulation procedures on the efficiency of transfection [130].

4. Cationic lipids as non-viral vectors for siRNA and DNA delivery

4.1. Gene delivery by cationic lipids

Gene delivery (DNA transfection) with cationic lipids (Figure 7) dates back to 1987 when it was reported by Felgner et al. [131], and the term “lipofection” was coined. Small unilamellar

liposomes containing the cationic lipid *N*-[1-(2,3-dioleoyloxy)propyl]-*N,N,N*-trimethyl ammonium chloride (DOTMA) was reported to spontaneously complex DNA completely entrapping the DNA, and enhanced fusion with the cell membrane in vitro in cell cultures, resulting in efficient delivery and expression of the delivered DNA. The lipofection was 5-100-fold more effective than the commonly used transfection techniques at the time by either calcium phosphate or DEAE-dextran (diethylaminoethyl-dextran), depending on the cell line used [131]. Cationic lipids have polar and non-polar domains and thus are amphiphilic in nature, with three general structural domains: (a) a cationic hydrophilic head-group (positively charged). The head-group can carry a permanent positive charge as in quaternary ammonium groups, or can be protonated at the physiological pH 7.4, such as primary and secondary amine groups. There can be one cationic group per lipid molecule (monovalent cationic lipids) or more than one cationic group per lipid molecule (multivalent cationic lipids); (b) a hydrophobic domain covalently attached by a linker to the cationic head-group. This domain can be in the form of either alkyl chains (commonly 2 chains) of various chain lengths (with various oxidation states) or can be a steroid such as cholesterol; (c) the linker between the head-group and the hydrophobic domain [132, 133]. This linker controls the biodegradation of the cationic lipid and its stability under different conditions according to the type of chemical bonds (e.g. ester, ether, or amide). Each domain can be controlled to change a specific character of the cationic lipid, e.g. using a disulfide functional group as a bioresponsive linker [134] which is reduced in the intracellular environment by glutathione/glutathione reductase and enhance biodegradation characters of the lipid and decrease its cytotoxicity.

4.2. The cationic head-group

The cationic head-group's main function is to bind electrostatically the negatively charged phosphates of the polynucleotides. The complexes of cationic lipids with polynucleotides such as DNA and siRNA are called lipoplexes. This requires the presence of a positive charge on the head-group at the physiological pH 7.4, i.e. the pKa of the head-group is ideally at least one unit higher than the physiological pH. The most commonly used head-groups contain nitrogen (e.g. amines or guanidines). However other head-groups, e.g. arsonium and phosphonium have been reported [135]. Arsonium is less toxic than arsenic (III), and in vitro cytotoxicity evaluation showed that arsonium and phosphonium are surprisingly less toxic than the ammonium group [135, 136].

One property that can be changed by controlling the type of the head-group is the head-group cross-sectional area. The greater the difference between the cross-sectional area of the polar head-group and that of the hydrophobic domain, when the former is designed to be smaller than the latter, the greater is the ability of the cationic lipid to fuse with the cell membrane and endosomal membrane and the greater is the release of polynucleotides from their complex with the cationic lipid due to the decreased structural stability of the lipid assembly [133, 137]. The hydration of the head-group affects its cross-sectional area, thus, the conjugation of groups which decrease the hydration state (such as hydroxyalkyl groups that form intermolecular H-bonds) decreases the head-group cross-sectional area.

Thus, gene delivery by DOTMA and DOTAP (1,2-dioleoyloxy-3-(trimethylammonio)-propane) was enhanced by incorporation of a hydroxyethyl group to yield the lipids DORIE

(1,2-dioleoyloxypropyl-3-dimethyl-hydroxyethyl ammonium bromide) and DORI (*N*-[1-(2,3-dioleoyloxy)propyl-*N*-[1-(2-hydroxy)ethyl]-*N,N*-dimethyl ammonium iodide) respectively [138, 139]. The head-group cross-sectional area can be also controlled by subtle changes to the head-group structure. The DC-Chol (3β (*N*-(*N',N'*-dimethylaminoethane)carbamoyl)cholesterol) with dimethylamino head-group resulted in more efficient transfection compared to DC-Chol with diethylamino or diisopropylamino head-groups, probably due to increased steric repulsion of the head-groups.

The in vitro gene transfer with six non-cholesterol-based cationic lipids (each having two alkyl chains) with a single guanidinium head-group in Chinese hamster ovary (CHO), COS-1, MCF-7, A549, and HepG2 cells has been reported [140]. These lipids were able to form lipoplexes with size-range 200-600 nm and ζ -potential +3.4 to -34 mV. The efficiencies of the lipids which had an extra quaternized cationic centre were 2-4-fold more than that of the commercially available reagent Lipofectamine in transfecting COS-1, CHO, A-549, and MCF-7 cells. MTT viability assay in CHO cells showed high (>75%) cell viabilities at the lipid/DNA charge ratios used. DNase I protection assays showed that the lipids having the extra quaternized centre protected DNA better against enzyme catalysed hydrolysis. These results shed light on the importance of choosing the type of head-group and number of cationic centres in designing cationic lipids [140].

A series of cationic cholesterol derivatives were synthesized by covalently attaching the heterocycles imidazole, piperazine, pyridine, and morpholine groups (the head-groups) to the parent cholesterol via a biodegradable carbamoyl linker [141]. These lipids were compared with the parent DC-Chol with the linear amine head-group, and they generally showed better or comparable transfection efficiency of pCMV-luciferase into human HepG2 cells (a human liver cancer cell line) in the presence or absence of FCS. The most efficient two of these lipids were with morpholine and piperazine head-groups, and they gave higher levels of gene expression in HepG2 and human melanoma cell line (KZ2) which are generally very hard to transfect with the commonly used reagents e.g. DC-Chol, Lipofectamine, and PEI. In vivo studies with lipids having morpholine and piperazine head-groups resulted in successful delivery of the reporter gene to the target cells through intrasplenic injection [141]. Cationic lipids which have more than one cationic head-group (multivalent cationic lipids) have more surface charge density than their monovalent (with one head-group) analogues, and they are generally expected to better bind and complex polynucleotides. Many multivalent cationic lipids contain a natural occurring polyamine such as spermidine and spermine, which are believed to interact with the minor groove of B-DNA [142].

The triamine spermidine and the tetramine spermine (Figure 6), and their diamine precursor putrescine, are organic polycations that are widely but unevenly distributed in both mammalian and non-mammalian cells and tissues. They have an essential role in controlling DNA, RNA and protein synthesis during normal and neoplastic growth, in cell differentiation, and tissue regeneration [143]. These polyamines exhibit many metabolic and neurophysiological effects in the nervous system, and are important for the developing and mature nervous system and affect modulation of ionic channels and calcium-dependent transmitter release [143-149].

Figure 7. Representative examples of cationic lipids used in DNA and siRNA delivery.

Spermine is incorporated in the cationic polymer polyspermine imidazole-4,5-imine (PSI) and in dioctadecylamidoglycyl-spermine (DOGS) [150] (Figure 7); spermidine is bound in cholesteryl-spermidine [151]. The free amine groups of spermine in cholesteryl-spermine conjugates have different pKa values and provide a buffering effect in the endosomes facilitating the escape of lipoplex from the endosomes [152]. The length of the linear polyamine attached to the hydrophobic domain and the charge distribution on it affects the transfection efficiency of the cationic lipid [153]. Addition of amine groups separated by methylene groups to the linear polyamine attached to a cholesterol residue did not automatically increase transfection efficiency regardless of the increased charge density. Molecular modelling simulations suggested that increasing chain length led to an increased number of folded conformations due to greater flexibility of the conjugates, which is unfavourable for interaction with DNA [132, 153].

The central tetramethylene motif of spermine was reported to be essential in conferring high transfection efficiency in a series of cholesterol-polyamine conjugates [152]. It was suggested that the tetramethylene segment of spermine can bridge between the DNA complementary strands, while the polyamine with a central trimethylene segment would only bind with the adjacent phosphates on the same DNA strand. These results point to the importance of the structure of the polyamine head-group and the relation between its amine groups, and also point to the fact that increasing efficiency of transfection is not only a matter of increasing the number of positive charges per head-group.

4.3. The hydrophobic domain

The length, saturation state and type of the hydrophobic chains conjugated to cationic lipids affect their transfection efficiency [154-156]. Although these factors were studied extensively for the effect on transfection, and although the majority of studies accepted that the type of alkyl chains influence the outcome of transfection, it is difficult to set a definitive set of rules to describe the best type of alkyl chains to be conjugated to the polar head-groups. The contribution of the alkyl chains (and the hydrophobic domain) to the cationic lipid properties as a whole is what determines the transfection efficiency of the lipid.

Results obtained with DMRIE (1,2-dimyristyloxypropyl-3-dimethylhydroxyethyl-ammonium bromide) [157], glycine betaine conjugates [138] with two alkyl chains, alkyl acyl carnitine esters having chains of length C12 to C18 [158], lactic acid conjugates of *N,N*-dialkyl amine group [159], lipids related to DOTAP with two alkyl chains (C12-C18) linked to the head-group through ether bonds [160], and cationic lipids with different hydroxyethyl or dihydroxypropyl ammonium backbones and esterified hydrocarbon chains and hydroxyl substituents [161] showed that a comparison of the cationic lipids based only on the lengths of the two saturated aliphatic chains led to the observation of the superior transfection efficiency of C14 chains over the longer C16 and C18 chains [132, 133]. It was proposed that a shorter chain length facilitates mixing with cellular membranes [138] which is important for endosomal escape [162].

In another set of experiments, we showed the longer chain C18 oleoyl (with one *cis*-double bond) to be more efficient than cationic lipids with shorter chain lengths. Varying the chain length in *N*⁴,*N*⁹-diacyl spermines from C10 to C18, for plasmid DNA delivery, resulted in us establishing that the conjugate with C18 oleoyl chains is both more efficient and less toxic than

the shorter chain conjugates [163]. A series of multivalent Gemini-surfactants with the hydrophobic chains systematically varied resulted in the conjugates with C18 oleoyl chains to be better in transfection than the C16 and C14 alkyl chains [164]. Chain saturation was also shown to affect the efficiency of transfection. The results of studies on a set of cationic triester phosphatidyl choline derivatives (each having two alkyl chains) show a strong dependence of their transfection efficiency on the lipid hydrocarbon chain characteristics, where transfection activity increases with increasing chain unsaturation from fully saturated to having two double bonds. Transfection efficiency decreased with increasing chain length (increasing the total number of carbons per lipid molecule ~30-50). Maximum transfection was with monounsaturated myristoleoyl 14:1 chains [156]. The data obtained from transfection experiments with 20 cationic phosphatidylcholine (PC) derivatives show that hydrocarbon chain variations results in transfection efficiencies that varies by more than 2 orders of magnitude. The most important variables were chain saturation state and total number of carbon atoms in the lipid chains. Transfection increased with decreasing chain length and increasing chain unsaturation. Best transfection efficiency was found for cationic lipids with monounsaturated (myristoleoyl) 14:1 chains [154]. Higher levels of transfection were also reported with lipids having oleoyl chains in comparison with stearoyl chains [157, 158]. Unsaturated chains promote lipid fusion between the lipoplexes and the various cellular membranes, which is essential for delivery and endosomal escape [133, 154, 165].

Cholesterol derivatives with various cationic head-groups were synthesized to investigate their efficiency as siRNA delivery vectors. The transfection efficiencies of siRNA lipoplexes prepared with the cationic cholesterol derivatives DC-Chol, cholesteryl-3 β -carboxyamido-ethylene-*N*-hydroxyethylamine (OH-Chol), and *N*-hydroxyethylaminopropane carbamoyl cholesterol (HAPC) was investigated in human prostate tumour cells that stably express the luciferase gene (PC-3-Luc). When lipoplexes were prepared in water, HAPC was more effective in knocking-down luciferase activity than OH-Chol and DC-Chol [166]. The presence of NaCl while preparing the lipoplexes increased the gene silencing efficiency of luciferase, while it did not affect efficiency of HAPC. The commercially available transfection reagent, Lipofectamine 2000 (a cationic lipid liposomal preparation) resulted in strong gene silencing by siRNA, but exhibited increased toxicity (~40% cell viability), in contrast to OH-Chol, DC-Chol, and HAPC lipoplexes (~80–100% cell viability). These results indicated that siRNA lipoplexes prepared with OH-Chol, and HAPC can efficiently suppress gene expression without increased cytotoxicity [166].

4.4. The linker

The linker is dependent upon the type (hence properties) of the functional group and its length (number of carbon atoms). The linker has two main functions: (a) to conjugate covalently the polar head-group to the hydrophobic domain; (b) to control the biodegradability of the cationic lipid and/or introduce a new property to the cationic lipid, e.g. responding to the intracellular reducing environment [133, 167]. The most commonly used linker functional groups are: amide, carbamate, ester, ether, ortho ester, and disulfide.

Both amides and ester bonds are biodegradable and hence are hypothesized to be less toxic than other non-biodegradable bonds (e.g. ethers) [168]. Lipids with a pyridinium head-group (with palmitoyl 16:0 hydrophobic domains and with ester and amide linkers) were used to prepare liposomes with either DOPE or cholesterol at the cationic lipid/helper-lipid molar ratio of 1:1. Following transfection of CHO cells with lipoplexes delivering plasmids expressing EGFP, the cationic lipids having amide linkers significantly increased transfection efficiency in all liposomal formulations compared to their counterparts having the ester linker [169]. The high transfection efficiency of lipids with amide linker was suggested to be due to their lower phase-transition temperature which makes the liposome's bilayer structure more stable in aqueous media during the transfection process as well as liposome storage. The phase-transition temperature of a lipid is the temperature at which there is a change in the lipid's physical state from the ordered gel phase (where the hydrocarbon chains are closely packed and fully extended) to the disordered liquid crystalline phase (where the hydrocarbon chains are fluid and randomly orientated) [169].

Depending on the structure of the cationic lipid, the linker influence on transfection efficiency can be more than on cytotoxicity. Cholesterol-based cationic lipids that have different nitrogenous heterocyclic head-groups (*N*-methylimidazole, *N*-methylmorpholine, and pyridine) and acid-labile linkers (carbamate, ester, and *N,O*-acetal ether) were used to transfect human embryonic kidney 293 (HEK 293) cells with EGFP plasmid [170]. Choosing those linkers was based on the concept that incorporation of acid-labile bonds in the cationic lipid structure enhances the release of polynucleotides from the endosomes, therefore increasing the transfection efficiency [171]. *N,O*-Acetals are known to undergo hydrolysis in acidic environment [170, 171]. The results showed that the structure of these lipids only slightly affected their cytotoxicity but largely changes the efficiency of intracellular accumulation of the polynucleotides. The lipids having the cationic head-groups pyridine and/or methylimidazole head-groups with either an ester or a carbamate linker resulted in better transfection efficiency as compared with the cationic lipids with either the *N*-methylmorpholine head-groups and/or an ether linker. The lipid that has a pyridine head-group and a carbamate linker to deliver EGFP plasmid resulted in comparable transfection efficiency with that achieved with commercially available Lipofectamine 2000.

Two cleavable cationic lipids having a linear or a cyclic ortho-ester linker between the cationic head-group and the unsaturated hydrophobic domain (two oleoyl chains) were previously reported [172]. It is hypothesized that the acidic pH in the endosomes catalyzes the hydrolysis of the linker group to result in fragmentation products that destabilize the endosomal membranes. At pH 7.4, the lipids (with DOPE) formed stable lipoplexes with plasmid DNA. Decreasing the pH enhanced the hydrolysis of the ortho ester linkers which removed the cationic head-groups and caused lipoplex aggregation. At pH 5.5, the cationic lipid *N*-[2-methyl-2-(1',2'-dioleylglyceroxy)dioxolan-4-yl]methyl-*N,N,N*-trimethylammonium iodide that have a cyclic ortho-ester linker showed increased pH-sensitivity and caused the permeation of its lipoplexes to model biomembranes within the time span of endosomal processing before the lysosomal degradation. This lipid markedly increased gene transfection (~3-50-fold) of the luciferase reporter protein in monkey kidney fibroblast (CV-1) and human breast cancer (HTB-129) cells in culture compared to the pH-insensitive control lipid DOTAP lipoplexes [172].

Transfection with DNA lipoplexes of three thiocholesterol-derived gemini cationic lipids possessing disulfide linkages incorporated between the cationic head-group and the thiocholesterol backbone in order to render the lipids biodegradable has been reported [173]. Comparing transfection in a prostate cancer line (PC3AR) and a human keratinocyte cell line (HaCat) with two commercially available reagents showed comparable or better expression of GFP in the transfected cells. Cytotoxicity studies showed the nontoxic property of these lipid-DNA complexes at different *N/P* ratios used for transfection studies. The rationale behind this design was to ensure the destabilization of the lipid-polynucleotide lipoplexes in the cytoplasm after reduction of the disulfide linker by the intracellular glutathione (GSH), which is the most abundant low molecular weight thiol present in cells and is involved in controlling cellular redox environment. GSH is found at very high intracellular concentrations and at comparatively low extracellular concentrations e.g. blood plasma concentrations (2 μ M) are 1000-fold less than concentration in erythrocytes (2 mM). This large difference between intra- and extracellular environments provides a potential mechanism for release of polynucleotides from lipoplexes of lipids that have a disulfide functional group linker and is now a well-trodden research path [115, 134, 173].

5. Conclusions and future avenues

In our research, symmetrical and asymmetrical acyl polyamine derivatives (fatty acid amides of spermine) [152] have been synthesized, characterized, and evaluated as non-viral vectors for siRNA [163, 174-177]. The intracellular delivery of siRNA and the subsequent sequence specific gene silencing has been quantified by flow cytometry techniques (FACS analysis) [163]. The ability of the spermine conjugates to bind siRNA and form nanoparticles has been investigated and the effect of the complexes of siRNA lipoplexes on the cell viability 48 h post-transfection has been quantified. Our SAR studies allow the identification of the most efficient fatty acids in terms of high gene-silencing efficiency and high cell viability [174-178].

Whilst we were completing this Chapter, four interesting papers, each one on a different aspect of this topic, were published. Langer, Anderson and co-workers at MIT reported on the delivery of immunostimulatory RNA (isRNA) to Toll-like receptor (TLR)-expressing cells to drive innate and adaptive immune responses. The specific activation of TLRs has potential for a variety of therapeutic indications including antiviral immunotherapy and as vaccine adjuvants. Effective lipidoid-isRNA nanoparticles, when tested in mice, stimulated strong IFN- α responses following subcutaneous injection, had robust antiviral activity that suppressed influenza virus replication, and enhanced antiovalbumin humoral and cell-mediated responses when used as a vaccine adjuvant. Their lipidoid formulations, designed specifically for the delivery of isRNA to TLRs, were superior to the commonly used *N*-[1-(2,3-dioleoyloxy)propyl]-*N,N,N*-trimethylammonium methylsulfate-RNA delivery system and may provide new tools for the manipulation of TLR responses in vitro and in vivo [179]. This paper follows after their other recent major contribution on delivering naked siRNA as part of a self-assembled (due to DNA complementarity) tetrahedral nanoparticle construct considering the presentation of folate as a cancer targeting ligand [180]. These monodisperse nanoparticles of

essentially naked DNA, carrying siRNA as the cargo, have a defined size of only a few nm. They show that at least three folate molecules per nanoparticle are required for optimal delivery of the siRNA into cells and that gene silencing only occurs when the ligands are appropriately orientated. In vivo, these naked DNA nanoparticles showed a longer blood circulation time than the parent siRNA [180]. In another exciting development, Geall and co-workers at Novartis have also advanced the field of nucleic acid vaccines by taking advantage of the recent innovations in non-viral systemic delivery of siRNA using lipid nanoparticles (LNPs) to develop a self-amplifying RNA vaccine. This technology elicited broad, potent, and protective immune responses, comparable with those achieved by a viral delivery system, but without the inherent limitations of viral vectors [181]. Even today, a biologically responsive cationic polymer system based on spermine has been reported for the intracellular delivery of siRNA [182]. This polyspermine imidazole-4,5-imine (PSI) (Figure 7) carrier is designed to be hydrolysed at the mildly acidic pH found in the endosome.

It is clear that both ssRNA to activate the immune system and RNAi brought about by siRNA delivery have high therapeutic potential. The major remaining barrier, that of efficient and potentially selective delivery to target cells is now being addressed. The non-viral delivery of siRNA is a major tool in modern functional genomics. Medicines design, the formulation of drugs, in this case siRNA and plasmid DNA, is an essential requirement for efficient gene therapy.

Acknowledgements

We thank the Egyptian Government for a fully-funded studentship to AAM.

Author details

Ian S. Blagbrough¹ and Abdelkader A. Metwally^{1,2}

*Address all correspondence to: prsisb@bath.ac.uk

1 Department of Pharmacy and Pharmacology, University of Bath, Bath BA2 7AY, UK

2 Department of Pharmaceutics and Industrial Pharmacy, Faculty of Pharmacy, Ain Shams University, Abbasya, Cairo, Egypt

References

- [1] Blagbrough IS, Zara C. Animal Models for Target Diseases in Gene Therapy - Using DNA and siRNA Delivery Strategies. *Pharmaceutical Research* 2009;26(1) 1-18.

- [2] van der Krol AR, Mur LA, Beld M, Mol J, Stuitje AR. Flavonoid Genes in Petunia: Addition of a Limited Number of Gene Copies May Lead to a Suppression of Gene Expression. *The Plant Cell Online* 1990;2(4) 291-299.
- [3] Napoli C, Lemieux C, Jorgensen R. Introduction of a Chimeric Chalcone Synthase Gene into Petunia Results in Reversible Co-Suppression of Homologous Genes *in Trans*. *The Plant Cell Online* 1990;2(4) 279-289.
- [4] Vanbolkland R, Vandergeest N, Mol JNM, Kooter JM. Transgene-Mediated Suppression of Chalcone Synthase Expression in Petunia Hybrida Results from an Increase in RNA Turnover. *Plant Journal* 1994;6(6) 861-877.
- [5] Metzlauff M, Odell M, Cluster PD, Flavell RB. RNA-Mediated RNA Degradation and Chalcone Synthase a Silencing in Petunia. *Cell* 1997;88(6) 845-854.
- [6] Hamilton AJ, Baulcombe DC. A Species of Small Antisense RNA in Posttranscriptional Gene Silencing in Plants. *Science* 1999;286(5441) 950-952. DOI: 10.1126/science.286.5441.950.
- [7] Hamilton A, Voinnet O, Chappell L, Baulcombe D. Two Classes of Short Interfering RNA in RNA Silencing. *EMBO Journal* 2002;21(17) 4671-4679. DOI: 10.1093/emboj/cdf464.
- [8] Baumberger N, Baulcombe DC. Arabidopsis Argonaute1 Is an RNA Slicer That Selectively Recruits microRNAs and Short Interfering RNAs. *Proceedings of the National Academy of Sciences of the United States of America* 2005;102(33) 11928-11933. DOI: 10.1073/pnas.0505461102.
- [9] Fire A, Xu SQ, Montgomery MK, Kostas SA, Driver SE, Mello CC. Potent and Specific Genetic Interference by Double-Stranded RNA in *Caenorhabditis Elegans*. *Nature* 1998;391(6669) 806-811.
- [10] Elbashir SM, Harborth J, Lendeckel W, Yalcin A, Weber K, Tuschl T. Duplexes of 21-Nucleotide RNAs Mediate RNA Interference in Cultured Mammalian Cells. *Nature* 2001;411(6836) 494-498.
- [11] Skloot R. *The Immortal Life of Henrietta Lacks*. London, UK: Macmillan; 2010.
- [12] Cogoni C, Macino G. Post-Transcriptional Gene Silencing across Kingdoms. *Current Opinion in Genetics and Development* 2000;10(6) 638-643.
- [13] Sontheimer EJ. Assembly and Function of RNA Silencing Complexes. *Nature Reviews Molecular Cell Biology* 2005;6(2) 127-138.
- [14] Collins RE, Cheng XD. Structural and Biochemical Advances in Mammalian RNAi. *Journal of Cellular Biochemistry* 2006;99(5) 1251-1266.
- [15] Hutvagner G, Simard MJ. Argonaute Proteins: Key Players in RNA Silencing. *Nature Reviews Molecular Cell Biology* 2008;9(1) 22-32.

- [16] Gaynor JW, Campbell BJ, Cosstick R. RNA Interference: A Chemist's Perspective. *Chemical Society Reviews* 2010;39(11) 4169-4184.
- [17] Liu QH, Paroo Z. Biochemical Principles of Small RNA Pathways. *Annual Review of Biochemistry* 2010;79 295-319.
- [18] Haley B, Zamore PD. Kinetic Analysis of the RNAi Enzyme Complex. *Nature Structural & Molecular Biology* 2004;11(7) 599-606.
- [19] Hutvagner G. Small RNA Asymmetry in RNAi: Function in Risc Assembly and Gene Regulation. *FEBS Letters* 2005;579(26) 5850-5857.
- [20] Wang HW, Noland C, Siridechadilok B, Taylor DW, Ma EB, Felderer K, Doudna JA, Nogales E. Structural Insights into RNA Processing by the Human Risc-Loading Complex. *Nature Structural & Molecular Biology* 2009;16(11) 1148-1153.
- [21] Wahid F, Shehzad A, Khan T, Kim YY. MicroRNAs: Synthesis, Mechanism, Function, and Recent Clinical Trials. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research* 2010;1803(11) 1231-1243.
- [22] Paddison PJ, Caudy AA, Bernstein E, Hannon GJ, Conklin DS. Short Hairpin RNAs (shRNAs) Induce Sequence-Specific Silencing in Mammalian Cells. *Genes & Development* 2002;16(8) 948-958.
- [23] Rao DD, Vorhies JS, Senzer N, Nemunaitis J. siRNA Vs. shRNA: Similarities and Differences. *Advanced Drug Delivery Reviews* 2009;61(9) 746-759.
- [24] Freier SM, Kierzek R, Jaeger JA, Sugimoto N, Caruthers MH, Neilson T, Turner DH. Improved Free-Energy Parameters for Predictions of RNA Duplex Stability. *Proceedings of the National Academy of Sciences of the United States of America* 1986;83(24) 9373-9377.
- [25] Freier SM, Altmann KH. The Ups and Downs of Nucleic Acid Duplex Stability: Structure-Stability Studies on Chemically-Modified DNA:RNA Duplexes. *Nucleic Acids Research* 1997;25(22) 4429-4443.
- [26] Beverly MB. Applications of Mass Spectrometry to the Study of siRNA. *Mass Spectrometry Reviews* 2010; 10.1002/mas.20260.
- [27] Shah SA, Brunger AT. The 1.8 Angstrom Crystal Structure of a Statically Disordered 17 Base-Pair RNA Duplex: Principles of RNA Crystal Packing and Its Effect on Nucleic Acid Structure. *Journal of Molecular Biology* 1999;285(4) 1577-1588.
- [28] Rana TM. Illuminating the Silence: Understanding the Structure and Function of Small RNAs. *Nature Reviews Molecular Cell Biology* 2007;8(1) 23-36.
- [29] Banan M, Puri N. The Ins and Outs of RNAi in Mammalian Cells. *Current Pharmaceutical Biotechnology* 2004;5(5) 441-450.

- [30] Hickerson RP, Vlassov AV, Wang Q, Leake D, Ilves H, Gonzalez-Gonzalez E, Contag CH, Johnston BH, Kaspar RL. Stability Study of Unmodified siRNA and Relevance to Clinical Use. *Oligonucleotides* 2008;18(4) 345-354.
- [31] Layzer JM, McCaffrey AP, Tanner AK, Huang Z, Kay MA, Sullenger BA. In Vivo Activity of Nuclease-Resistant siRNA. *RNA-a Publication of the Rna Society* 2004;10(5) 766-771.
- [32] Turner JJ, Jones SW, Moschos SA, Lindsay MA, Gait MJ. Maldi-Tof Mass Spectral Analysis of siRNA Degradation in Serum Confirms an RNase a-Like Activity. *Molecular Biosystems* 2007;3(1) 43-50.
- [33] Volkov AA, Kruglova NyS, Meschaninova MI, Venyaminova AG, Zenkova MA, Vlassov VV, Chernolovskaya EL. Selective Protection of Nuclease-Sensitive Sites in siRNA Prolongs Silencing Effect. *Oligonucleotides* 2009;19(2) 191-202.
- [34] Melnikova I. RNA-Based Therapies. *Nature Reviews Drug Discovery* 2007;6(11) 863-864.
- [35] Skoblov M. Prospects of Antisense Therapy Technologies. *Molecular Biology* 2009;43(6) 917-929.
- [36] Ghosal A, Kabir AH, Mandal A. RNA Interference and Its Therapeutic Potential. *Central European Journal of Medicine* 2011;6(2) 137-147.
- [37] Cheng H, Luo C, Wu X, Zhang Y, He Y, Wu Q, Xia Y, Zhang J. shRNA Targeting Plc ϵ Inhibits Bladder Cancer Growth in Vitro and in Vivo. *Urology* 2011;78(2).
- [38] Qin X-J, Dai D-J, Gao Z-G, Huan J-L, Zhu L. Effect of Lentivirus-Mediated shRNA Targeting Vegfr-3 on Proliferation, Apoptosis and Invasion of Gastric Cancer Cells. *International Journal of Molecular Medicine* 2011;28(5) 761-768.
- [39] Liu Q-S, Zhang J, Liu M, Dong W-G. Lentiviral-Mediated miRNA against Liver-Intestine Cadherin Suppresses Tumor Growth and Invasiveness of Human Gastric Cancer. *Cancer Science* 2010;101(8) 1807-1812.
- [40] Bumcrot D, Manoharan M, Koteliensky V, Sah DWY. RNAi Therapeutics: A Potential New Class of Pharmaceutical Drugs. *Nature Chemical Biology* 2006;2(12) 711-719.
- [41] Aleku M, Schulz P, Keil O, Santel A, Schaeper U, Dieckhoff B, Janke O, Endruschat J, Durieux B, Roeder N, Loeffler K, Lange C, Fechtner M, Moepert K, Fisch G, Dames S, Arnold W, Jochims K, Giese K, Wiedenmann B, Scholz A, Kaufmann J. Atu027, a Liposomal Small Interfering RNA Formulation Targeting Protein Kinase N3, Inhibits Cancer Progression. *Cancer Research* 2008;68(23) 9788-9798.
- [42] Behlke MA. Progress Towards in Vivo Use of siRNA. *Molecular Therapy* 2006;13(4) 644-670.
- [43] Whitehead KA, Langer R, Anderson DG. Knocking Down Barriers: Advances in siRNA Delivery. *Nature Reviews Drug Discovery* 2009;8(2) 129-138.

- [44] Reischl D, Zimmer A. Drug Delivery of siRNA Therapeutics: Potentials and Limits of Nanosystems. *Nanomedicine: Nanotechnology, Biology and Medicine* 2009;5(1) 8-20.
- [45] van de Water FM, Boerman OC, Wouterse AC, Peters JGP, Russel FGM, Masereeuw R. Intravenously Administered Short Interfering RNA Accumulates in the Kidney and Selectively Suppresses Gene Function in Renal Proximal Tubules. *Drug Metabolism and Disposition* 2006;34(8) 1393-1397.
- [46] Ruponen M, Ronkko S, Honkakoski P, Pelkonen J, Tammi M, Urtti A. Extracellular Glycosaminoglycans Modify Cellular Trafficking of Lipoplexes and Polyplexes. *Journal of Biological Chemistry* 2001;276(36) 33875-33880.
- [47] Singh AK, Kasinath BS, Lewis EJ. Interaction of Polycations with Cell-Surface Negative Charges of Epithelial-Cells. *Biochimica et Biophysica Acta* 1992;1120(3) 337-342.
- [48] Watts JK, Deleavey GF, Damha MJ. Chemically Modified siRNA: Tools and Applications. *Drug Discovery Today* 2008;13(19-20) 842-855.
- [49] Behlke MA. Chemical Modification of siRNAs for in Vivo Use. *Oligonucleotides* 2008;18(4) 305-319.
- [50] Hall AHS, Wan J, Shaughnessy EE, Shaw BR, Alexander KA. RNA Interference Using Boranophosphate siRNAs: Structure-Activity Relationships. *Nucleic Acids Research* 2004;32(20) 5991-6000.
- [51] Ui-Tei K, Naito Y, Zenno S, Nishi K, Yamato K, Takahashi F, Juni A, Saigo K. Functional Dissection of siRNA Sequence by Systematic DNA Substitution: Modified siRNA with a DNA Seed Arm Is a Powerful Tool for Mammalian Gene Silencing with Significantly Reduced Off-Target Effect. *Nucleic Acids Research* 2008;36(7) 2136-2151.
- [52] Chiu YL, Rana TM. siRNA Function in RNAi: A Chemical Modification Analysis. *RNA-a Publication of the Rna Society* 2003;9(9) 1034-1048.
- [53] Bell NM, Micklefield J. Chemical Modification of Oligonucleotides for Therapeutic, Bioanalytical and Other Applications. *ChemBioChem* 2009;10(17) 2691-2703.
- [54] Corey DR. Chemical Modification: The Key to Clinical Application of RNA Interference? *Journal of Clinical Investigation* 2007;117(12) 3615-3622.
- [55] Nishina K, Unno T, Uno Y, Kubodera T, Kanouchi T, Mizusawa H, Yokota T. Efficient in Vivo Delivery of siRNA to the Liver by Conjugation of Alpha-Tocopherol. *Molecular Therapy* 2008;16(4) 734-740.
- [56] Lorenz C, Hadwiger P, John M, Vornlocher HP, Unverzagt C. Steroid and Lipid Conjugates of siRNAs to Enhance Cellular Uptake and Gene Silencing in Liver Cells. *Bioorganic & Medicinal Chemistry Letters* 2004;14(19) 4975-4977.
- [57] Soutschek J, Akinc A, Bramlage B, Charisse K, Constien R, Donoghue M, Elbashir S, Geick A, Hadwiger P, Harborth J, John M, Kesavan V, Lavine G, Pandey RK, Racie T,

- Rajeev KG, Rohl I, Toudjarska I, Wang G, Wuschko S, Bumcrot D, Koteliensky V, Limmer S, Manoharan M, Vornlocher HP. Therapeutic Silencing of an Endogenous Gene by Systemic Administration of Modified siRNAs. *Nature* 2004;432(7014) 173-178.
- [58] Wolfrum C, Shi S, Jayaprakash KN, Jayaraman M, Wang G, Pandey RK, Rajeev KG, Nakayama T, Charrise K, Ndungo EM, Zimmermann T, Koteliensky V, Manoharan M, Stoffel M. Mechanisms and Optimization of in Vivo Delivery of Lipophilic siRNAs. *Nature Biotechnology* 2007;25(10) 1149-1157.
- [59] Eguchi A, Dowdy SF. siRNA Delivery Using Peptide Transduction Domains. *Trends in Pharmacological Sciences* 2009;30(7) 341-345.
- [60] Muratovska A, Eccles MR. Conjugate for Efficient Delivery of Short Interfering RNA (siRNA) into Mammalian Cells. *FEBS Letters* 2004;558(1-3) 63-68.
- [61] Moschos SA, Jones SW, Perry MM, Williams AE, Erjefalt JS, Turner JJ, Barnes PJ, Sproat BS, Gait MJ, Lindsay MA. Lung Delivery Studies Using siRNA Conjugated to Tat(48-60) and Penetratin Reveal Peptide Induced Reduction in Gene Expression and Induction of Innate Immunity. *Bioconjugate Chemistry* 2007;18(5) 1450-1459.
- [62] Kim SH, Jeong JH, Lee SH, Kim SW, Park TG. Peg Conjugated Vegf siRNA for Anti-Angiogenic Gene Therapy. *Journal of Controlled Release* 2006;116(2) 123-129.
- [63] Xia CF, Zhang Y, Boado RJ, Pardridge WM. Intravenous siRNA of Brain Cancer with Receptor Targeting and Avidin-Biotin Technology. *Pharmaceutical Research* 2007;24(12) 2309-2316.
- [64] Chu TC, Twu KY, Ellington AD, Levy M. Aptamer Mediated siRNA Delivery. *Nucleic Acids Research* 2006;34(10).
- [65] Li MJ, Bauer G, Michienzi A, Yee JK, Lee NS, Kim J, Li S, Castanotto D, Zaia J, Rossi JJ. Inhibition of HIV-1 Infection by Lentiviral Vectors Expressing Pol Iii-Promoted Anti-HIV RNAs. *Molecular Therapy* 2003;8(2) 196-206.
- [66] Sliva K, Schnierle BS. Selective Gene Silencing by Viral Delivery of Short Hairpin RNA. *Virology Journal* 2010;7 248.
- [67] Berkowitz R, Ilves H, Lin WY, Eckert K, Coward A, Tamaki S, Veres G, Plavec I. Construction and Molecular Analysis of Gene Transfer Systems Derived from Bovine Immunodeficiency Virus. *Journal of Virology* 2001;75(7) 3371-3382.
- [68] Atkinson H, Chalmers R. Delivering the Goods: Viral and Non-Viral Gene Therapy Systems and the Inherent Limits on Cargo DNA and Internal Sequences. *Genetica* 2010;138(5) 485-498.
- [69] Chen Y, Du D, Wu J, Chan CP, Tan Y, Kung HF, He ML. Inhibition of Hepatitis B Virus Replication by Stably Expressed shRNA. *Biochemical and Biophysical Research Communications* 2003;311(2) 398-404.

- [70] Kumar A, Panda SK, Durgapal H, Acharya SK, Rehman S, Kar UK. Inhibition of Hepatitis E Virus Replication Using Short Hairpin RNA (shRNA). *Antiviral Research* 2010;85(3) 541-550.
- [71] Bushman FD. Retroviral Integration and Human Gene Therapy. *Journal of Clinical Investigation* 2007;117(8) 2083-2086.
- [72] Brummelkamp TR, Bernards R, Agami R. Stable Suppression of Tumorigenicity by Virus-Mediated RNA Interference. *Cancer Cell* 2002;2(3) 243-247.
- [73] Anesti AM, Peeters PJ, Royaux I, Coffin RS. Efficient Delivery of RNA Interference to Peripheral Neurons in Vivo Using Herpes Simplex Virus. *Nucleic Acids Research* 2008;36(14).
- [74] Manjunath N, Wu H, Subramanya S, Shankar P. Lentiviral Delivery of Short Hairpin RNAs. *Advanced Drug Delivery Reviews* 2009;61(9) 732-745.
- [75] Carlson ME, Hsu M, Conboy IM. Imbalance between Psmad3 and Notch Induces Cdk Inhibitors in Old Muscle Stem Cells. *Nature* 2008;454(7203) 528-532.
- [76] Yoo JY, Kim JH, Kwon YG, Kim EC, Kim NK, Choi HJ, Yun CO. Vegf-Specific Short Hairpin RNA-Expressing Oncolytic Adenovirus Elicits Potent Inhibition of Angiogenesis and Tumor Growth. *Molecular Therapy* 2007;15(2) 295-302.
- [77] Xia H, Mao Q, Eliason SL, Harper SQ, Martins IH, Orr HT, Paulson HL, Yang L, Kottin RM, Davidson BL. RNAi Suppresses Polyglutamine-Induced Neurodegeneration in a Model of Spinocerebellar Ataxia. *Nature Medicine* 2004;10(8) 816-820.
- [78] Kuhlmann KF, Gouma DJ, Wesseling JG. Adenoviral Gene Therapy for Pancreatic Cancer: Where Do We Stand? *Digestive Surgery* 2008;25(4) 278-292.
- [79] Descamps D, Benihoud K. Two Key Challenges for Effective Adenovirus-Mediated Liver Gene Therapy: Innate Immune Responses and Hepatocyte-Specific Transduction. *Current Gene Therapy* 2009;9(2) 115-127.
- [80] Nayak S, Herzog RW. Progress and Prospects: Immune Responses to Viral Vectors. *Gene Therapy* 2010;17(3) 295-304.
- [81] Raper SE, Chirmule N, Lee FS, Wivel NA, Bagg A, Gao GP, Wilson JM, Batshaw ML. Fatal Systemic Inflammatory Response Syndrome in a Ornithine Transcarbamylase Deficient Patient Following Adenoviral Gene Transfer. *Molecular Genetics and Metabolism* 2003;80(1-2) 148-158.
- [82] Howe SJ, Mansour MR, Schwarzwaelder K, Bartholomae C, Hubank M, Kempinski H, Brugman MH, Pike-Overzet K, Chatters SJ, de Ridder D, Gilmour KC, Adams S, Thornhill SI, Parsley KL, Staal FJ, Gale RE, Linch DC, Bayford J, Brown L, Quaye M, Kinnon C, Ancliff P, Webb DK, Schmidt M, von Kalle C, Gaspar HB, Thrasher AJ. Insertional Mutagenesis Combined with Acquired Somatic Mutations Causes Leuke-

mogenesis Following Gene Therapy of Scid-X1 Patients. *Journal of Clinical Investigation* 2008;118(9) 3143-3150.

- [83] Hacein-Bey-Abina S, von Kalle C, Schmidt M, Le Deist F, Wulffraat N, McIntyre E, Radford I, Villeval JL, Fraser CC, Cavazzana-Calvo M, Fischer A. A Serious Adverse Event after Successful Gene Therapy for X-Linked Severe Combined Immunodeficiency. *New England Journal of Medicine* 2003;348(3) 255-256.
- [84] Qasim W, Gaspar HB, Thrasher AJ. Progress and Prospects: Gene Therapy for Inherited Immunodeficiencies. *Gene Therapy* 2009;16(11) 1285-1291.
- [85] Sheridan C. Gene Therapy Finds Its Niche. *Nature Biotechnology* 2011;29(2) 121-128.
- [86] Mintzer MA, Simanek EE. Nonviral Vectors for Gene Delivery. *Chemical Reviews* 2009;109(2) 259-302.
- [87] Torchilin VP, Rammohan R, Weissig V, Levchenko TS. Tat Peptide on the Surface of Liposomes Affords Their Efficient Intracellular Delivery Even at Low Temperature and in the Presence of Metabolic Inhibitors. *Proceedings of the National Academy of Sciences of the United States of America* 2001;98(15) 8786-8791.
- [88] Torchilin VP, Levchenko TS, Rammohan R, Volodina N, Papahadjopoulos-Sternberg B, D'Souza GG. Cell Transfection in Vitro and in Vivo with Nontoxic Tat Peptide-Liposome-DNA Complexes. *Proceedings of the National Academy of Sciences of the United States of America* 2003;100(4) 1972-1977.
- [89] Snyder EL, Dowdy SF. Cell Penetrating Peptides in Drug Delivery. *Pharmaceutical Research* 2004;21(3) 389-393.
- [90] Gait MJ. Peptide-Mediated Cellular Delivery of Antisense Oligonucleotides and Their Analogues. *Cellular and Molecular Life Sciences* 2003;60(5) 844-853.
- [91] Eguchi A, Akuta T, Okuyama H, Senda T, Yokoi H, Inokuchi H, Fujita S, Hayakawa T, Takeda K, Hasegawa M, Nakanishi M. Protein Transduction Domain of HIV-1 Tat Protein Promotes Efficient Delivery of DNA into Mammalian Cells. *Journal of Biological Chemistry* 2001;276(28) 26204-26210.
- [92] Foerg C, Merkle HP. On the Biomedical Promise of Cell Penetrating Peptides: Limits Versus Prospects. *Journal of Pharmaceutical Sciences* 2008;97(1) 144-162.
- [93] Mäe M, Andaloussi SE, Lehto T, Langel Ü. Chemically Modified Cell-Penetrating Peptides for the Delivery of Nucleic Acids. *Expert Opinion on Drug Delivery* 2009;6(11) 1195-1205.
- [94] Crombez L, Aldrian-Herrada G, Konate K, Nguyen QN, McMaster GK, Brasseur R, Heitz F, Divita G. A New Potent Secondary Amphipathic Cell-Penetrating Peptide for siRNA Delivery into Mammalian Cells. *Molecular Therapy* 2009;17(1) 95-103.

- [95] Lundberg P, El-Andaloussi S, Sutlu T, Johansson H, Langel U. Delivery of Short Interfering RNA Using Endosomolytic Cell-Penetrating Peptides. *FASEB Journal* 2007;21(11) 2664-2671.
- [96] Simeoni F, Morris MC, Heitz F, Divita G. Insight into the Mechanism of the Peptide-Based Gene Delivery System Mpg: Implications for Delivery of siRNA into Mammalian Cells. *Nucleic Acids Research* 2003;31(11) 2717-2724.
- [97] Zeineddine D, Papadimou E, Chebli K, Gineste M, Liu J, Grey C, Thurig S, Behfar A, Wallace VA, Skerjanc IS, Puceat M. Oct-3/4 Dose Dependently Regulates Specification of Embryonic Stem Cells toward a Cardiac Lineage and Early Heart Development. *Developmental Cell* 2006;11(4) 535-546.
- [98] Kim WJ, Christensen LV, Jo S, Yockman JW, Jeong JH, Kim YH, Kim SW. Cholesteryl Oligoarginine Delivering Vascular Endothelial Growth Factor siRNA Effectively Inhibits Tumor Growth in Colon Adenocarcinoma. *Molecular Therapy* 2006;14(3) 343-350.
- [99] Davidson TJ, Harel S, Arboleda VA, Prunell GF, Shelanski ML, Greene LA, Troy CM. Highly Efficient Small Interfering RNA Delivery to Primary Mammalian Neurons Induces microRNA-Like Effects before mRNA Degradation. *Journal of Neuroscience* 2004;24(45) 10040-10046.
- [100] Turner JJ, Jones S, Fabani MM, Ivanova G, Arzumanov AA, Gait MJ. RNA Targeting with Peptide Conjugates of Oligonucleotides, siRNA and Pna. *Blood Cells, Molecules, and Diseases* 2007;38(1) 1-7.
- [101] Chiu YL, Ali A, Chu CY, Cao H, Rana TM. Visualizing a Correlation between siRNA Localization, Cellular Uptake, and RNAi in Living Cells. *Chemistry and Biology* 2004;11(8) 1165-1175.
- [102] Frankel AD, Pabo CO. Cellular Uptake of the Tat Protein from Human Immunodeficiency Virus. *Cell* 1988;55(6) 1189-1193.
- [103] Vives E, Brodin P, Lebleu B. A Truncated HIV-1 Tat Protein Basic Domain Rapidly Translocates through the Plasma Membrane and Accumulates in the Cell Nucleus. *Journal of Biological Chemistry* 1997;272(25) 16010-16017.
- [104] Jones AT. Macropinocytosis: Searching for an Endocytic Identity and Role in the Uptake of Cell Penetrating Peptides. *Journal of Cellular and Molecular Medicine* 2007;11(4) 670-684.
- [105] Patel LN, Zaro JL, Shen WC. Cell Penetrating Peptides: Intracellular Pathways and Pharmaceutical Perspectives. *Pharmaceutical Research* 2007;24(11) 1977-1992.
- [106] Hassani Z, Lemkine GF, Erbacher P, Palmier K, Alfama G, Giovannangeli C, Behr JP, Demeneix BA. Lipid-Mediated siRNA Delivery Down-Regulates Exogenous Gene Expression in the Mouse Brain at Picomolar Levels. *The Journal of Gene Medicine* 2005;7(2) 198-207.

- [107] Grayson AC, Doody AM, Putnam D. Biophysical and Structural Characterization of Polyethylenimine-Mediated siRNA Delivery in Vitro. *Pharmaceutical Research* 2006;23(8) 1868-1876.
- [108] Spagnou S, Miller AD, Keller M. Lipidic Carriers of siRNA: Differences in the Formulation, Cellular Uptake, and Delivery with Plasmid DNA. *Biochemistry* 2004;43(42) 13348-13356.
- [109] Bolcato-Bellemin AL, Bonnet ME, Creusatt G, Erbacher P, Behr JP. Sticky Overhangs Enhance siRNA-Mediated Gene Silencing. *Proceedings of the National Academy of Sciences of the United States of America* 2007;104 16050-16055.
- [110] Moghimi SM, Symonds P, Murray JC, Hunter AC, Debska G, Szewczyk A. A Two-Stage Poly(Ethylenimine)-Mediated Cytotoxicity: Implications for Gene Transfer/Therapy. *Molecular Therapy* 2005;11(6) 990-995.
- [111] Werth S, Urban-Klein B, Dai L, Hobel S, Grzelinski M, Bakowsky U, Czubyko F, Aigner A. A Low Molecular Weight Fraction of Polyethylenimine (PEI) Displays Increased Transfection Efficiency of DNA and siRNA in Fresh or Lyophilized Complexes. *Journal of Controlled Release* 2006;112(2) 257-270. DOI: 10.1016/j.jconrel.2006.02.009.
- [112] Boussif O, Lezoualc'h F, Zanta MA, Mergny MD, Scherman D, Demeneix B, Behr JP. A Versatile Vector for Gene and Oligonucleotide Transfer into Cells in Culture and in Vivo: Polyethylenimine. *Proceedings of the National Academy of Sciences of the United States of America* 1995;92(16) 7297-7301.
- [113] Philipp A, Zhao X, Tarcha P, Wagner E, Zintchenko A. Hydrophobically Modified Oligoethylenimines as Highly Efficient Transfection Agents for siRNA Delivery. *Bioconjugate Chemistry* 2009;20(11) 2055-2061.
- [114] Bajaj A, Kondaiah P, Bhattacharya S. Synthesis and Gene Transfection Efficacies of PEI-Cholesterol-Based Lipopolymers. *Bioconjugate Chemistry* 2008;19(8) 1640-1651.
- [115] Breunig M, Hozsa C, Lungwitz U, Watanabe K, Umeda I, Kato H, Goepferich A. Mechanistic Investigation of Poly(Ethyleneimine)-Based siRNA Delivery: Disulfide Bonds Boost Intracellular Release of the Cargo. *Journal of Controlled Release* 2008;130(1) 57-63.
- [116] Mao S, Sun W, Kissel T. Chitosan-Based Formulations for Delivery of DNA and siRNA. *Advanced Drug Delivery Reviews* 2010;62(1) 12-27.
- [117] Liu X, Howard KA, Dong M, Andersen MO, Rahbek UL, Johnsen MG, Hansen OC, Besenbacher F, Kjems J. The Influence of Polymeric Properties on Chitosan/siRNA Nanoparticle Formulation and Gene Silencing. *Biomaterials* 2007;28(6) 1280-1288.
- [118] Katas H, Alpar HO. Development and Characterisation of Chitosan Nanoparticles for siRNA Delivery. *Journal of Controlled Release* 2006;115(2) 216-225.

- [119] Menuel S, Fontanay S, Clarot I, Duval RE, Diez L, Marsura A. Synthesis and Complexation Ability of a Novel Bis-(Guanidinium)-Tetrakis-(Beta-Cyclodextrin) Dendrimeric Tetrapod as a Potential Gene Delivery (DNA and siRNA) System. Study of Cellular siRNA Transfection. *Bioconjugate Chemistry* 2008;19(12) 2357-2362.
- [120] Hu-Lieskovan S, Heidel JD, Bartlett DW, Davis ME, Triche TJ. Sequence-Specific Knockdown of Ews-Fli1 by Targeted, Nonviral Delivery of Small Interfering RNA Inhibits Tumor Growth in a Murine Model of Metastatic Ewing's Sarcoma. *Cancer Research* 2005;65(19) 8984-8992.
- [121] Davis ME. The First Targeted Delivery of siRNA in Humans Via a Self-Assembling, Cyclodextrin Polymer-Based Nanoparticle: From Concept to Clinic. *Molecular Pharmaceutics* 2009;6(3) 659-668.
- [122] Davis ME, Zuckerman JE, Choi CH, Seligson D, Tolcher A, Alabi CA, Yen Y, Heidel JD, Ribas A. Evidence of RNAi in Humans from Systemically Administered siRNA Via Targeted Nanoparticles. *Nature* 2010;464(7291) 1067-1070.
- [123] O'Mahony AM, Ogier J, Desgranges S, Cryan JF, Darcy R, O'Driscoll CM. A Click Chemistry Route to 2-Functionalised Pegylated and Cationic Beta-Cyclodextrins: Co-Formulation Opportunities for siRNA Delivery. *Organic and Biomolecular Chemistry* 2012;10(25) 4954-4960.
- [124] Zhu L, Mahato RI. Lipid and Polymeric Carrier-Mediated Nucleic Acid Delivery. *Expert Opinion on Drug Delivery* 2010;7(10) 1209-1226.
- [125] Dufès C, Uchegbu IF, Schätzlein AG. Dendrimers in Gene Delivery. *Advanced Drug Delivery Reviews* 2005;57(15) 2177-2202.
- [126] Zinselmeyer BH, Mackay SP, Schätzlein AG, Uchegbu IF. The Lower-Generation Polypropyleneimine Dendrimers Are Effective Gene-Transfer Agents. *Pharmaceutical Research* 2002;19(7) 960-967.
- [127] Fischer D, Li Y, Ahlemeyer B, Kriegelstein J, Kissel T. In Vitro Cytotoxicity Testing of Polycations: Influence of Polymer Structure on Cell Viability and Hemolysis. *Biomaterials* 2003;24(7) 1121-1131.
- [128] Sonawane ND, Szoka FC, Verkman AS. Chloride Accumulation and Swelling in Endosomes Enhances DNA Transfer by Polyamine-DNA Polyplexes. *Journal of Biological Chemistry* 2003;278(45) 44826-44831.
- [129] Taratula O, Savla R, He HX, Minko T. Poly(Propyleneimine) Dendrimers as Potential siRNA Delivery Nanocarrier: From Structure to Function. *International Journal of Nanotechnology* 2011;8(1-2) 36-52.
- [130] Perez AP, Romero EL, Morilla MJ. Ethylenediamine Core PAMAM Dendrimers/siRNA Complexes as in Vitro Silencing Agents. *International Journal of Pharmaceutics* 2009;380(1-2) 189-200.

- [131] Felgner PL, Gadek TR, Holm M, Roman R, Chan HW, Wenz M, Northrop JP, Ringold GM, Danielsen M. Lipofection: A Highly Efficient, Lipid-Mediated DNA-Transfection Procedure. *Proceedings of the National Academy of Sciences of the United States of America* 1987;84(21) 7413-7417.
- [132] Martin B, Sainlos M, Aissaoui A, Oudrhiri N, Hauchecorne M, Vigneron JP, Lehn JM, Lehn P. The Design of Cationic Lipids for Gene Delivery. *Current Pharmaceutical Design* 2005;11(3) 375-394.
- [133] Bhattacharya S, Bajaj A. Advances in Gene Delivery through Molecular Design of Cationic Lipids. *Chemical Communications* 2009;(31) 4632-4656.
- [134] Shirazi RS, Ewert KK, Leal C, Majzoub RN, Boussein NF, Safinya CR. Synthesis and Characterization of Degradable Multivalent Cationic Lipids with Disulfide-Bond Spacers for Gene Delivery. *Biochimica et Biophysica Acta-Biomembranes* 2011;1808(9) 2156-2166.
- [135] Guenin E, Herve AC, Floch V, Loisel S, Yaouanc JJ, Clement JC, Ferec C, des Abbayes H. Cationic Phosphonolipids Containing Quaternary Phosphonium and Arsonium Groups for DNA Transfection with Good Efficiency and Low Cellular Toxicity. *Angewandte Chemie International Edition* 2000;39(3) 629-631.
- [136] Floch V, Loisel S, Guenin E, Herve AC, Clement JC, Yaouanc JJ, des Abbayes H, Ferec C. Cation Substitution in Cationic Phosphonolipids: A New Concept to Improve Transfection Activity and Decrease Cellular Toxicity. *Journal of Medicinal Chemistry* 2000;43(24) 4617-4628.
- [137] Hasegawa S, Hirashima N, Nakanishi M. Comparative Study of Transfection Efficiency of Cationic Cholesterols Mediated by Liposomes-Based Gene Delivery. *Bioorganic and Medicinal Chemistry Letters* 2002;12(9) 1299-1302.
- [138] Felgner JH, Kumar R, Sridhar CN, Wheeler CJ, Tsai YJ, Border R, Ramsey P, Martin M, Felgner PL. Enhanced Gene Delivery and Mechanism Studies with a Novel Series of Cationic Lipid Formulations. *Journal of Biological Chemistry* 1994;269(4) 2550-2561.
- [139] Bennett MJ, Aberle AM, Balasubramaniam RP, Malone JG, Malone RW, Nantz MH. Cationic Lipid-Mediated Gene Delivery to Murine Lung: Correlation of Lipid Hydration with in Vivo Transfection Activity. *Journal of Medicinal Chemistry* 1997;40(25) 4069-4078.
- [140] Sen J, Chaudhuri A. Design, Syntheses, and Transfection Biology of Novel Non-Cholesterol-Based Guanidinylated Cationic Lipids. *Journal of Medicinal Chemistry* 2005;48(3) 812-820.
- [141] Gao H, Hui KM. Synthesis of a Novel Series of Cationic Lipids That Can Act as Efficient Gene Delivery Vehicles through Systematic Heterocyclic Substitution of Cholesterol Derivatives. *Gene Therapy* 2001;8(11) 855-863.

- [142] Schmid N, Behr JP. Location of Spermine and Other Polyamines on DNA as Revealed by Photoaffinity Cleavage with Polyaminobenzenediazonium Salts. *Biochemistry* 1991;30(17) 4357-4361.
- [143] Bernstein H-G, Müller M. The Cellular Localization of the L-Ornithine Decarboxylase/Polyamine System in Normal and Diseased Central Nervous Systems. *Progress in Neurobiology* 1999;57(5) 485-505.
- [144] Janne J, Poso H, Raina A. Polyamines in Rapid Growth and Cancer. *Biochimica et Biophysica Acta* 1978;473(3-4) 241-293.
- [145] Fozard JR, Part ML, Prakash NJ, Grove J, Schechter PJ, Sjoerdsma A, Koch-Weser J. L-Ornithine Decarboxylase: An Essential Role in Early Mammalian Embryogenesis. *Science* 1980;208(4443) 505-508.
- [146] Pegg AE, Seely JE, Poso H, della Ragione F, Zagon IA. Polyamine Biosynthesis and Interconversion in Rodent Tissues. *Federation Proceedings* 1982;41(14) 3065-3072.
- [147] Pegg AE, McCann PP. Polyamine Metabolism and Function. *American Journal of Physiology* 1982;243(5) C212-221.
- [148] Janne J, Alhonen L, Leinonen P. Polyamines - from Molecular-Biology to Clinical-Applications. *Annals of Medicine* 1991;23(3) 241-259.
- [149] Hougaard DM. Polyamine Cytochemistry: Localization and Possible Functions of Polyamines. *International Review of Cytology* 1992;138 51-88.
- [150] Behr JP, Demeneix B, Loeffler JP, Perez-Mutul J. Efficient Gene Transfer into Mammalian Primary Endocrine Cells with Lipopolyamine-Coated DNA. *Proceedings of the National Academy of Sciences of the United States of America* 1989;86(18) 6982-6986.
- [151] Moradpour D, Schauer JI, Zurawski VR, Wands JR, Boutin RH. Efficient Gene Transfer into Mammalian Cells with Cholesteryl-Spermidine. *Biochemical and Biophysical Research Communications* 1996;221(1) 82-88.
- [152] Geall AJ, Eaton MAW, Baker T, Catterall C, Blagbrough IS. The Regiochemical Distribution of Positive Charges Along Cholesterol Polyamine Carbamates Plays Significant Roles in Modulating DNA Binding Affinity and Lipofection. *FEBS Letters* 1999;459(3) 337-342.
- [153] Fujiwara T, Hasegawa S, Hirashima N, Nakanishi M, Ohwada T. Gene Transfection Activities of Amphiphilic Steroid-Polyamine Conjugates. *Biochimica et Biophysica Acta-Biomembranes* 2000;1468(1-2) 396-402.
- [154] Koynova R, Tenchov B, Wang L, MacDonald RC. Hydrophobic Moiety of Cationic Lipids Strongly Modulates Their Transfection Activity. *Molecular Pharmaceutics* 2009;6(3) 951-958.

- [155] Incani V, Lavasanifar A, Uludag H. Lipid and Hydrophobic Modification of Cationic Carriers on Route to Superior Gene Vectors. *Soft Matter* 2010;6(10) 2124-2138.
- [156] Koynova R, Tenchov B. Cationic Phospholipids: Structure-Transfection Activity Relationships. *Soft Matter* 2009;5(17) 3187-3200.
- [157] Floch V, Legros N, Loisel S, Guillaume C, Guilbot J, Benvegny T, Ferrieres V, Plusquellec D, Ferec C. New Biocompatible Cationic Amphiphiles Derivative from Glycine Betaine: A Novel Family of Efficient Nonviral Gene Transfer Agents. *Biochemical and Biophysical Research Communications* 1998;251(1) 360-365.
- [158] Wang J, Guo X, Xu Y, Barron L, Szoka FC, Jr. Synthesis and Characterization of Long Chain Alkyl Acyl Carnitine Esters. Potentially Biodegradable Cationic Lipids for Use in Gene Delivery. *Journal of Medicinal Chemistry* 1998;41(13) 2207-2215.
- [159] Laxmi AA, Vijayalakshmi P, Kaimal TN, Chaudhuri A, Ramadas Y, Rao NM. Novel Non-Glycerol-Based Cytfectins with Lactic Acid-Derived Head Groups. *Biochemical and Biophysical Research Communications* 2001;289(5) 1057-1062.
- [160] Heyes JA, Niculescu-Duvaz D, Cooper RG, Springer CJ. Synthesis of Novel Cationic Lipids: Effect of Structural Modification on the Efficiency of Gene Transfer. *Journal of Medicinal Chemistry* 2002;45(1) 99-114.
- [161] Lindner LH, Brock R, Arndt-Jovin D, Eibl H. Structural Variation of Cationic Lipids: Minimum Requirement for Improved Oligonucleotide Delivery into Cells. *Journal of Controlled Release* 2006;110(2) 444-456.
- [162] Xu YH, Szoka FC. Mechanism of DNA Release from Cationic Liposome/DNA Complexes Used in Cell Transfection. *Biochemistry* 1996;35(18) 5616-5623.
- [163] Ghonaim HM, Ahmed OAA, Pourzand C, Blagbrough IS. Varying the Chain Length in N^4, N^9 -Diacyl Spermines: Non-Viral Lipopolyamine Vectors for Efficient Plasmid DNA Formulation. *Molecular Pharmaceutics* 2008;5(6) 1111-1121.
- [164] McGregor C, Perrin C, Monck M, Camilleri P, Kirby AJ. Rational Approaches to the Design of Cationic Gemini Surfactants for Gene Delivery. *Journal of the American Chemical Society* 2001;123(26) 6215-6220.
- [165] Heyes J, Palmer L, Bremner K, MacLachlan I. Cationic Lipid Saturation Influences Intracellular Delivery of Encapsulated Nucleic Acids. *Journal of Controlled Release* 2005;107(2) 276-287.
- [166] Hattori Y, Hagiwara A, Ding W, Maitani Y. NaCl Improves siRNA Delivery Mediated by Nanoparticles of Hydroxyethylated Cationic Cholesterol with Amido-Linker. *Bioorganic and Medicinal Chemistry Letters* 2008;18(19) 5228-5232.
- [167] Rao NM, Gopal V. Cationic Lipids for Gene Delivery in Vitro and in Vivo. *Expert Opinion on Therapeutic Patents* 2006;16(6) 825-844.
- [168] Ilies MA, Seitz WA, Johnson BH, Ezell EL, Miller AL, Thompson EB, Balaban AT. Lipophilic Pyrylium Salts in the Synthesis of Efficient Pyridinium-Based Cationic

- Lipids, Gemini Surfactants, and Lipophilic Oligomers for Gene Delivery. *Journal of Medicinal Chemistry* 2006;49(13) 3872-3887.
- [169] Zhu L, Lu Y, Miller DD, Mahato RI. Structural and Formulation Factors Influencing Pyridinium Lipid-Based Gene Transfer. *Bioconjugate Chemistry* 2008;19(12) 2499-2512.
- [170] Medvedeva DA, Maslov MA, Serikov RN, Morozova NG, Serebrenikova GA, Sherglov DV, Latyshev AV, Vlassov VV, Zenkova MA. Novel Cholesterol-Based Cationic Lipids for Gene Delivery. *Journal of Medicinal Chemistry* 2009;52(21) 6558-6568.
- [171] Guo X, Szoka FC. Chemical Approaches to Triggerable Lipid Vesicles for Drug and Gene Delivery. *Accounts of Chemical Research* 2003;36(5) 335-341.
- [172] Chen HG, Zhang HZ, McCallum CM, Szoka FC, Guo X. Unsaturated Cationic Ortho Esters for Endosome Permeation in Gene Delivery. *Journal of Medicinal Chemistry* 2007;50(18) 4269-4278.
- [173] Bajaj A, Kondaiah P, Bhattacharya S. Effect of the Nature of the Spacer on Gene Transfer Efficacies of Novel Thiocholesterol Derived Gemini Lipids in Different Cell Lines: A Structure-Activity Investigation. *Journal of Medicinal Chemistry* 2008;51(8) 2533-2540.
- [174] Metwally AA, Pourzand C, Blagbrough IS. Efficient Gene Silencing by Self-assembled Complexes of siRNA and Symmetrical Fatty Acid Amides of Spermine, *Pharmaceutics* 2011;3 125-140. DOI: 10.3390/pharmaceutics3020125.
- [175] Metwally AA, Blagbrough IS. Self-assembled Lipoplexes of Short Interfering RNA (siRNA) Using Spermine-based Fatty Acid Amide Guanidines: Effect on Gene Silencing Efficiency, *Pharmaceutics* 2011;3 406-424. DOI: 10.3390/pharmaceutics3030406.
- [176] Blagbrough IS, Metwally AA, Ghonaim HM, Asymmetrical N4,N9-Diacyl Spermines: SAR studies of Nonviral Lipopolyamine Vectors for Efficient siRNA Delivery with Silencing of EGFP Reporter Gene, *Molecular Pharmaceutics* 2012;9 1853-1861.
- [177] Metwally AA, Reelfs O, Pourzand C, Blagbrough IS, Efficient Silencing of EGFP Reporter Gene with siRNA Delivered by Asymmetrical N4,N9-Diacyl Spermines, *Molecular Pharmaceutics* 2012;9 1862-1876.
- [178] Metwally AA, Blagbrough IS, Mantell JM, Quantitative Silencing of EGFP Reporter Gene by Self-assembled siRNA Lipoplexes of LinOS and Cholesterol, *Molecular Pharmaceutics*, 2012;9 3384-3395. DOI: 10.1021/mp300435x.
- [179] Nguyen DN, Mahon KP, Chikh G, Kim P, Chung H, Vicari AP, Love KT, Goldberg M, Chen S, Krieg AM, Chen JZ, Langer R, Anderson DG. Lipid-derived Nanoparticles for Immunostimulatory RNA Adjuvant Delivery. *Proceedings of the National Academy of Sciences of the United States of America* 2012;109(14) E797-E803. DOI: 10.1073/pnas.1121423109.

- [180] Lee H, Lytton-Jean AKR, Chen Y, Love KT, Park AI, Karagiannis ED, Sehgal A, Querbes W, Zurenko CS, Jayaraman M, Peng CG, Charisse K, Borodovsky A, Manoharan M, Donahoe JS, Truelove J, Nahrendorf M, Langer R, Anderson DG. Molecularly Self-assembled Nucleic Acid Nanoparticles for Targeted in Vivo siRNA Delivery. *Nature Nanotechnology* 2012;7(6) 389-393. DOI: 10.1038/NNANO.2012.73.
- [181] Geall AJ, Verma A, Otten GR, Shaw CA, Hekele A, Banerjee K, Cu Y, Beard CW, Brito LA, Krucker T, O'Hagan DT, Singh M, Mason PW, Valiante NM, Dormitzer PR, Barnett SW, Rappuoli R, Ulmer JB, Mandl CW. Nonviral Delivery of Self-amplifying RNA Vaccines. *Proceedings of the National Academy of Sciences of the United States of America* 2012;109(36) 14604-14609. DOI: 10.1073/pnas.1209367109.
- [182] Duan S, Yuan W, Wu F, Jin T. Polyspermine Imidazole-4,5-imine, a Chemically Dynamic and Biologically Responsive Carrier System for Intracellular Delivery of siRNA. *Angewandte Chemie International Edition* 2012;51(32), 7938-7941. DOI: 10.1002/anie.201201793.