We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Avian Mycotoxicosis in Developing Countries

Adeniran Lateef Ariyo, Ajagbonna Olatunde Peter, Sani Nuhu Abdulazeez and Olabode Hamza Olatunde

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/56050>

1. Introduction

Avian mycotoxicosis refers to all the diseases caused by the effect of mycotoxin in birds. These diseases may not be pathognomonic and sometimes subclinical and difficult to diagnose. The problem is worldwide but effort will be made to localise the effect of these diseases in developing countries. Developing countries have more than 50% of total meat and egg production in global poultry market: [1]. The global poultry meat market is 86.8 million tonnes, consisting of chicken: 85.6%, turkey: 6.8%; duck: 4.6%; goose and guinea fowl: 2.6%.

Years	world	Developed countries	Developing countries	Share (%) of developing countries
1970	15	11	4	26
1975	19	13	5	31
1980	26	18	8	31
1985	31	21	10	33
1990	41	26	15	37
1995	55	28	26	48
2000	69	33	36	53
2005	81	37	44	55
(Increase %)	437	227	1,043	-

Table 1. Development of poultry meat production in developed and developing countries (million tonnes).

Years	world	Developed countries	Developing countries	Share (%) of developing countries
1970	20	15	5	24
1975	22	16	6	27
1980	26	18	8	32
1985	31	19	12	39
1990	35	19	16	46
1995	43	17	25	59
2000	51	18	33	64
2005	59	19	40	68
(Increase %)	195	29	758	-

Source: [1]

Table 2. Development of poultry hen egg production in developed and developing countries (million tonnes)

The world market poultry meat and egg market is been influenced by the production and managemental style from the developing countries. Avian mycotoxicosis is a great constraint in poultry industry, because the disease is characterized by immunosuppresion, hepatotoxicity, nephrotoxicity, loss of egg production, mutagenicity and tetratogenicity.

Mycotoxin are antinutritive factor present in feed ingredients and in concentrated feed, they are a group of secondary fungal metabolites of low molecular weight, diverse and ambiguous in nature, which are specifically implicated in causing toxic effect in animals and man [2,3]. Mycotoxicosis has been a major but unrecognized food safety issue for several centuries. They are naturally occurring contaminants that causes health related problems when it gets into the body through natural route of ingestion, inhalation or may be absorbed through the skin [4]. They are endogenously generated in foods as a result of secondary metabolism [5]. These metabolites are synthesized in or on food surfaces and transported through the food chain[6]. Mycotoxin production takes place in the mycelium after active fungal growth, but may accumulate in specialized structures such as sclerotia, conidia or in surrounding area [7]. Animal studies have shown that, besides acute effects, mycotoxins can cause carcinogenic, mutagenic and teratogenic effects. Mycotoxins-contaminated poultry feed can lead to the transfer of toxins through meat and egg to human beings.

The Food and Agriculture Organization [8] estimating that as much as 25% of the world's Agricultural commodities are contaminated with mycotoxins, leading to significant economic losses. Mycotoxigenic fungi genera include; *Aspergillus*, *Penicillium* and *Fusarium*. The important mycotoxin in the developing countries include aflatoxins, ochratoxins, citrinin, T-2 toxin, deoxynivalenol (DON), fumonisins and zearalenone.

Fungi genera	Associated mycotoxin
<i>Aspergillus</i>	Aflatoxin, Ochratoxins, cyclopiazonic acid, patulin, sterigmatocystin, gliotoxin, citrinin.
<i>Penicillium</i>	Ochratoxin, citrinin, patulin, penicillic acid, cyclopiazonic acid, penitrem A, griseofulvin.
<i>Fusarium</i>	Fumonisin, moniliformin, zearalenone, zearalenol, deoxynivalenol, nivalenol, 15-acetyldeoxynivalenol, 3-acetyldeoxynivalenol, T-2 toxin, iso T-2 toxin, acetyl T-2 toxin, t-2 triol, T-2 tetraol, fusarenon- X, diacetoxyscripentriol, neosalaniol, fusaric acid.
<i>Claviceps</i>	Ergot alkaloids.

[9]

Table 3. Showing fungi genera and the associated mycotoxin.

2. Avian aflatoxicosis

2.1. Aetiology

Avian aflatoxicosis is a disease of poultry caused by aflatoxin. Aflatoxins (AF) are widely distributed toxins produced by *Aspergillus*. Of the over 180 species of *Aspergillus*, only a few are aflatoxigenic. After the discovery of AF in the 1960s, *A. flavus* and *A. parasiticus* of the section *Flavi* were the only known AF producers producing the B and B/G types of AF, respectively [10]. Other aflatoxigenic species that subsequently emerged are *A. nomius* (B and G types), *A. bombycis* (B and G AF), *A. ochraceoroseus*, and *A. pseudotamarii* (B type), but they occur less frequently [11,12]. *A. tamarii*, *A. parvisclerotigenus* (B types), *A. rambellii* and certain members of *Aspergillus* subgenus *Nidulantes* namely: *Emericella venezualensis* [13] and *E. astellata* [14] have now been included in the growing list of aflatoxigenic species. *A. arachidicola* sp. Nov. and *A. minisclerotigenes* sp. Nov that produce both forms of the toxin, are the latest emerging aflatoxigenic species. The unexpected new comer is *A. niger*, an ochratoxin (OT) producer which was discovered over four decades ago but was never associated with AF synthesis. However, in a search for aflatoxigenic fungi in Romanian medicinal herbs, [15] showed the capacity of some strains of *A. niger* to produce AFB1 [16].

AFBI is a member of aflatoxin group is one of the most carcinogenic natural product formed in nature [17]. AF has been detected in most countries of the world. Four toxins soon identified: Aflatoxin B1, B2, G1, G2-blue or green florescence under UV-light. •AflatoxinB1most important -highly carcinogenic and widespread occurrence in foods •(B1> M1> G1> B2> M2~ G2). Aflatoxin M1: hydroxylated product of B1appears in milk, urine, and feces as metabolic product

2.2. Factor enhancing AF prevalence in feed ingredients and poultry feed

There are several factors enhancing the prevalence of aflatoxigenic fungi and aflatoxin production in developing countries. The factor include the following:

The food materials must be infected by aflatoxigenic fungi which deposit the toxins on feed ingredients and concentrated feeds.

The substrate which may be feed ingredients and concentrated feeds possesses a source of energy in the form of carbohydrates and organic and inorganic source of nitrogen, trace elements and moisture for growth of mould and toxin production [8].

Among cereal, the size and integrity of the seed coat also affects the susceptibility of fungal infection and mycotoxin formation [18]

The environmental condition favouring mould growth and AF production are hot and humid conditions, the optimal temperature varies between 24°C and 28°C [19] and seed moisture content of at least 17.5% [20]. These conditions that favours mould growth are present in most developing countries.

Soil type also affect the level of AF contamination of crops for example, light sandy soil support rapid growth of fungi [21]

Presence of other microorganism either bacteria for example presence of *Streptococcus lactis* and *Lactobacillus casei* causes reduction in AF production by *A. parasiticus* [20]. Meanwhile fungal metabolites like rubratoxin and cerulenin enhances AF production [16].

Agricultural practices also affect AF contamination of feed ingredient. Off season harvesting and harvesting system that enhances seed breakage would also increase the degree of AF production [22].

A well aerated storage condition used in most developing country to store feed ingredients increases metabolism and subsequent AF production [23].

2.3. Occurrence of AF

AF has been found as contaminants in animal feed ingredient worldwide. The occurrence in developing countries is more because there is no strict food and feed quality control programmes to reduce the burden of AF. Also their environmental condition presented as hot and humid climate makes most developing country vulnerable to AF in poultry feed. Among the four AF that are of significance in poultry include; AFB1, B2, G1 and G2. AFB1 was detected mostly from animal and feed ingredients from developing countries. Among the feeds ingredients, sorghum, wheat, maize were the most investigated, data on groundnut cake, cotton seed meal and fish meal showed high level of AF contamination. The highest level of AFB1 contamination of feed ingredient were reported in corn from Pakistan 25 µg/kg [24], Nigeria wheat was found to be contaminated with 17.10-20.53 µg/kg [25].

Higher level of contamination of AF were found in the animal feed than the feed ingredient possibly because of the storage condition which allowed growth and proliferation of mycotoxigenic fungi. In Nigeria AF was detected in poultry feed at 0.0-67.9 µg/kg [26], wheat 17.10-20.53 µg/kg [25], millet 1370-3475 µg/kg [23].

AF contamination of feed and feed ingredients has been a major concern in many developing country like Pakistan where concentration ranging from 24-37.62 µg/kg were found in poultry feed and poultry feed ingredient [24].

Commodity	Country	Type of AF	Incidence ($\mu\text{g/kg}$)	Range $\pm\text{SD}$ ($\mu\text{g/kg}$)	Mean level	References
Poultry/livestock	Nigeria	AFB1	6/13	0.0-67.9	15.5	[26]
Animal feed	Kenya	AFB1	703/830	0.9-595	8.9-46.0	[27]
Animal feed	South Africa	AF	17/23	0.8-156	39.8	[28]
Poultry feed	Morocco	AFB1	14/21	0.3-58	8.4	[29]
Sorghum	Malawi	AFB1	2/15	1.7-3.0	2.35 $\pm$ 0.65	[30]
Wheat	Kenya	AFB1	23/50	0-7	1.93	[31]
Wheat	Tunisia	AFs	15/51	4.0-12.9	6.7 $\pm$ 2.4	[32]
Wheat	Nigeria	AFB1	-	17.10-20.53	19.00 $\pm$ 1.67	[25]
Wheat	Algeria	AFB1	30/53	0.13-37.42	>5	[33]
Wheat	South Africa	AFB1	13/238	0.5-2.0	>2	[34]
Maize	Ghana	AF	30/30	6.20-29.50	13-596	[35]
Maize	Uganda	AF	22/49	1.00-1000	-	[22]
Millet	Nigeria	AFB1	12/49	1370.28-3495	2587.47 $\pm$ 78.23	[23]
Mouldy Sorghum	Nigeria	AFB1	93/168		199.51 $\pm$ 26	[36]
Poultry feed	Pakistan	AFB1	60%		37.62	[24]
Corn	Pakistan	AFB1	8/13		25	[24]
Rice broken	Pakistan	AFB1	3/5		21	[24]
Wheat	Pakistan	AFB1	3/5		19	[24]
Cotton seed meal	Pakistan	AFB1	5/5		22	[24]
Fish meal	Pakistan	AFB1	3/5		24	[24]
Broiler starter (Crumbs)	Pakistan	AFB1	6/11		20	[24]
Broiler starter (Mash)	Pakistan	AFB1	8/14		21	[24]
Layer Starter (Crumbs)	Pakistan	AFB1	16/30		26	[24]
Layer Grower (Crumbs)	Pakistan	AFB1	3/7		23	[24]
Layer Grower (Mash)	Pakistan	AFB1	3/5		32.5	[24]
Layers (Crumbs)	Pakistan	AFB1	23/60		21	[24]
Layers (Mash)	Pakistan	AFB1	25/58		23.5	[24]
Poultry Feed	China	AFB1	38.2		29.7	[37]
Poultry Feed	South Africa	Aflatoxin	22/62		0.7 $\pm$ 0.7	[38]
Maize	Morocco	AFB1	16/20		1.57 $\pm$ 0.78	[29]

[16]

Table 4. Showing occurrence of AF in animal feed and feed ingredients in developing countries

An unacceptably high level of concentration of AF that ranges from 1-1000 µg/kg was found in maize from Uganda[22] About 70% of wheat samples investigated in Algeria were contamination at a range of 0.13-3742mg/kg [33]

Animal species	Clinical signs Performance effect	Gross pathology	Histopathology	References
Duckling	inappetance, reduced growth. abnormal vacuolation, feather. picking, discoloration of leg/feet. lameness, ataxia, convulsion.	Liver enlarged, pale and shrunken. Kidney enlarged and pale Hydropericardium, ascitis.		[40]
Turkey	inappetance, unsteady gait, Recumbency, anaemia.	body in good condition, generalized and edema. Liver and kidney Congested enlarged and firm. Gallbladder was full.		[41, 42]
Chicken	Same in turkey and duckling Increase mortality Loss of production Increased susceptibility to infectious disease and vaccination failure.	Same in turkey and duckling	Fatty vacuolation of the liver. Karyomegaly and prominent nucleoli in hepatocytes, Bile duct proliferation.	[40-44]
Chicken (broiler)	impaired performance lower testis weight and Semen volume, Spermatocrit and Testosterone value.		Abnormal spermatozoa, cessation of spermatogenesis in seminiferous tubules.	[45-46]

Table 5. Main clinical signs, performance and pathological features in food producing animals exposed to AF in selected studies.

2.4. Pharmacological interaction

Aflatoxicosis has effect on plasma half-life, thus it affects drug effect in the body. [47] observed that chlortetracycline plasma concentrations were lowered due to decreased drug binding to plasma protein [48]. Though opinion differed considerably on sparing or aggravating effect caused by the addition of chlortetracycline to feed contaminated with aflatoxin [49,50].

2.5. Metabolism and residues

In broilers, metabolites of aflatoxins BI and B2 concentrated in kidney and liver but cleared within 4 days. Then metabolism of Aflatoxin B1 into conjugated aflatoxins B2a

and M1 occurred in the liver, which will be metabolized to aflatoxicol [51-54]. Aflatoxin B1 was excreted in the bile, urine, and feces as 6 major metabolites [55]. The half-life of aflatoxin B1 in laying hens is about 67 hours [56], though feed: egg transmission is about 5000:1 [57]. Most aflatoxin excreted through the bile and intestine, but aflatoxin B1 and aflatoxicol were detected in ova and eggs for 7 days or longer [58-59]. Aflatoxin B1 accumulated in reproductive organs and its subsequent transmission to eggs and hatched progeny (yolk sac and liver) in poultry [60]. It is well established that AFB₁ is both carcinogenic and cytotoxic. For example, synthesis of both RNA and DNA was inhibited when AF (5mg/kg of feed) was given to rats over a 6-week-period. The activated AFB₁ metabolite (i.e. AFB₁-8, 9-epoxide) forms a covalent bond with the N₇ of guanine [61] and forms AFB₁-N₇-guanine adducts in the target cells. The results are G_→T transversions, DNA repair, lesions, mutations and subsequently tumor formation [57]. The reactive epoxide can also be hydrolyzed to AFB₁-8, 9-dihydrodiol which ionizes to form a Schiff's base with primary amine groups in the proteins [58]. The short-lived epoxide AFB₁ has also been associated with coagulopathy due to reduced synthesis of vitamin K and other clotting factors as a result of sub-lethal intoxication of animals [62]. With regard to the cytotoxic effects, AFB₁ has been shown to induce lipid peroxidation in rat livers leading to oxidative damage to hepatocytes [63]. A more recent study [64], has demonstrated that AFB₁ can inhibit cyclic nucleotide phosphodiesterase activity in the brain, liver, heart, and kidney tissues.

3. Avian ochratoxicosis

3.1. Aetiology

This is a disease of bird caused by Ochratoxins. Ochratoxins are among the most toxic mycotoxins to poultry. They are nephrotoxins found in grains and feeds worldwide [65- 66]. Ochratoxins are isocoumarin compounds linked to L-b-phenylalanine and are designated A, B, C, and D, because of their methyl and ethyl esters. Ochratoxin A (OTA) is the most common and most toxic, and is relatively stable. OTA production is dependent on different factors such as temperature, water activity (*aw*) and medium composition, which affect the physiology of fungal producers. In cool and temperate regions, OTA is mainly produced by *Penicillium verrucosum* [67, 68] or *P. Nordicum* [69, 70]. *P. verrucosum* mainly contaminates plants such as cereal crops, whereas *P. nordicum* has been mainly detected in meat products and cheese [69]. In tropical and semitropical regions, OTA is mainly produced by *Aspergillus ochraceus* [71-73]. *A. ochraceus* is also referred to as *A. allutaceus* var *allutaceus* [71]. *A. ochraceus* have been reported in a large variety of matters like nuts, dried peanuts, beans, spices green coffee beans and dried fruits, but also in processed meat and smoked and salted fish [71]. Two other species of *Aspergillus* section *nigri*, *A. niger* var *niger* [74-75] and *A. carbonarius* [76,77] have been reported as OTA producers. The OTA contamination of substrate such as cereals, oilseeds and mixed feeds in warm zones is thought to be due to *A.niger* var *niger* in addition to *A. ochraceus* species [78], whereas *A. carbonarius* seems to be more common in grapes, raisins and coffee [79-80].

Recently, [81] isolated two new OTA producing *Aspergillus* species from coffee beans. These species, *A. lacticoffeatus* and *A. sclerotiumniger*, need further investigations and are provisionally accepted in section *Nigri*. In addition, another *Aspergillus* species, *A. alliaceus* also named *Petromyces alliaceus* isolated from onions [82] has been previously reported as OTA producer under laboratory conditions [83]. This species has been suspected to be responsible for the occasional OTA contamination in Californian figs [84, 85] an Argentinean medicinal herbs [67]. The biosynthetic pathway for OTA has not yet been completely established. However, labeling experiments using both ¹⁴C- and ¹³C-labelled precursors showed that the phenylalanine moiety originates from the shikimate pathway and the dihydroisocoumarin moiety from the pentaketide pathway. The first step in the synthesis of the isocoumarin polyketide consists in the condensation of one acetate unit (acetyl-CoA) to four malonate units. Recent data showed that this step requires the activity of a polyketide synthase [86]. Moreover, the gene encoding polyketide synthase appears to be very different between *Penicillium* and *Aspergillus* species [86]. In *A. ochraceus*, the gene of polyketide synthase is expressed only under OTA permissive conditions and only during the early stages of the mycotoxin synthesis [86]. No such data are presently available on *penicillium*. In *Penicillium* species, ' [86]' observed that *P.nordicum* and *P. verrucosum* use two different polyketide synthases for OTA synthesis. This difference is probably related to the *P. verrucosum* ability to produce citrinin, also a polyketide-based mycotoxin, in addition to OTA.

Destined Specie	Country	Type of samples	Type of Grain composition	LOD (LOQ)	Incidence (%)	Average Range		References
Poultry	Kuwait	Raw ingredients	Wheat bran	5	10/14(71.4%)	4.6	n.d. -12.1	[87]
			soybean meal		18/21(85.8%)	7.9	n.d - 40	
			Yellow maize		31/32 (96.8%)	6.38	n.d. -14.5	
		Feed	Layer mash	-	20/20 (100%)	9.6	5-16.7	
			broiler starter		13/14 (93%)	8.0	nd-9.1	
			Broiler finisher	-	17/19 (90%)	6.1	nd-14.3	
Poultry	Venezuela	Concentrated feed	-	-	47 /50 (94%)	-	2.56 -31.98	[88]
Poultry	Argentina	Feed	Corn (60%)	10	38%	27		[89]
Poultry	China	Feed			61.5%	7.0		[37]
Poultry	India	Raw ingredients	Wheat		4/12 (33.3%)	60.75	n.d. -98	[90]
			Maize		5/12 (41.7%)	104.4	n.d. -140	
			Rice		3/12 (25%)	20	n.d. -25	
			Sorghum		2/12 (16.7%)	34	n.d. -38	
			Barley		3/12 (25%)	26	n.d. -38	
			Grams		2/12 (16.7%)	12.5	n.d - 15	
			Ground nut		2/12 (16.7%)	39	n.d.48	
			Millet		3/12 (25%)	2.5	n.d. -3	
			Cotton seed meal		2/10 (20%)	31	n.d. -42	
			Soybean meal		2/10 (20%)	31	n.d. -42	
			Rapeseed meal		2/10 (20%)	19	n.d. -28	
			Sunflower meal		3/10 (30%)	49	n.d. -68	
			Guar meal		2/10 (20%)	21	n.d. -22	
			Corn gluten meal		2/10 (20%)	21	n.d. -22	

Source: [91]

Table 6. Showing the occurrence of Ochratoxin in developing countries in a selected surveys.

3.2. Occurrence of ochratoxin

OTA was reported by [90] in raw ingredient for making poultry feed were found to be contaminated at range of 00-140 µg/kg in maize 00-98mg/kg in wheat followed by sunflower meal at 00-68 µg/kg. The incidence in India ranges from 16-41% [90]. Also high level of contamination was reported in soya meal from Kuwait n-d-40 µg/kg [87]. [88] reported a contamination level of 2.56-31.98 µg/kg in Venezuela concentrated poultry feed investigated.

Animal specie	Production Phase (age)	Experimental (dosage)	Clinical signs and (duration)performance effects	Gross pathology	References
Laying hens) (<i>Isa brown</i>)	Laying (28 week-old)	Artificially contaminated feed (100-2000) ug/kg (30 days)	Decrease of egg mean weight Decrease of number of eggs placed Shell decalcification/thinning Egg altered conformation	Kidney Congestion and hemorrhages Increased size Liver - Yellowish	[92]
Broiler chickens	Fattening (6 day-old)	Artificially contaminated feed (100-2000) µg/kg (28 days)	Decreased growth rate Reduced feed consumption Decreased feed conversion Reduced serum protein and albumin.	Not evaluated	[93]
Laying hen (<i>Isa brown</i>)	Laying (11 month old)	Naturally contaminated diet (160-332) ug/kg	Debility Reduced egg production	Kidneys Congestion Hemorrhage Renomegalia Liver yellowish color Diffuse Hepatomegalia Scattered necrotic foci Congestion Hemorrhage	[94]
Broiler chickens	Fattening (1 day-old)	Artificially contaminated Feed (0-800) ug/kg (5 weeks)	Significantly decreased of -Body weight -Feed consumption - Feed conversion ratio - Increased mortality	Not assessed	[95]

Source: [91]

Table 7. Showing the main clinical signs, performance and pathological features in poultry exposed to OTA in selected studies

4. Funmonism

4.1. Aetiology

The fumonism, are food borne carcinogenic mycotoxin that affect animals and man. They are evaluated as group 2B component carcinogen [96]. It has various analog. Twenty eight

fumonisin analogs separated into four main groups, as fumonisin A, B, C, and P series has been identified. The fumonisin B (FB) analogs, comprising toxicologically important FB₁, FB₂, and FB₃, are the most abundant naturally occurring fumonisins, with FB₁ predominating and usually being found at the highest levels [96] in feed ingredients and poultry feed in world wide. FB₁ accounts for about 80% of the total fumonisins produced in this substrates, while FB₂ usually makes around 20% and FB₃ usually makes up from about 5% when cultured on corn or rice or in liquid medium [97-100].

Different *Fusarium* species have been reported to produce fumonisins

4.2. Occurrence of fumonisin

FB₁ a potent carcinogen was found in maize investigated from developing countries like Argentina, Benin, Egypt, Nepal, Honduras, Malawi, Zambia Botswana, and Tanzania at a range between 35- 65,000 µg/kg. [101-109]. Poultry feed investigated in china was contaminated with 1854.3mg/kg(37).

No doubt the climatic condition to the agricultural practices in these country allow the growth of fungi and subsequent elaboration of toxin in their substrate.

Country	Nature of samples	Number of samples Positive/total	Highest level (µg/kg)	References
Argentina	maize	16/17	2000	[101]
Benin	maize	20/21	2310	[102]
Egypt	maize	2/2	2380	[103]
Nepal	maize	12/24	4600	[104]
Honduras	Maize	24/24	6555	[105]
Zambia	maize	20/20	1710	[102]
China	Poultry Feed	82%	1854.3	[37]
Botswana	Maize		35-255	[106]
Mozambique	Maize		240-296	[106]
Malawi	Maize		ND-115	[106]
Tanzania	Maize		ND-160	[106]
Honduras	Maize		68-6555	[107]
Uruguay	Maize		ND-3688	[108]
Coastarica	Maize		1700-4780	[109]

Table 8. Occurrence of FB₁ In Feed Ingredient In Developing Countries

Animal Species	Type of toxin	Clinical signs	Gross pathology	Histopathology	References
Laying hen	FB1	Black sticky diarrhea Reduction in food intake, egg production and body weight. Lameness, increased mortality and impaired immunity.	Kidney, pancreas and liver enlargement. Enlargement of proventriculus. Atrophy of lymphoid Organ. Rickets.	Liver has multifocal necrosis of hepatocytes hyperplasia of hepatocytes and bile ductules. Intestine had villous atrophy Globlet cell hyperplasia lymphoid cell depletion from thymus.	[110] [111]
	FB1+ FB2 + moliniformin.			Decrease in blood cell counts Haemoglobin, PCV and white Blood cell count. Abnormal erythrocyte Formation	[112]
Chicken Embryo	FB1	100% mortality	pathological changes in liver, heart, lung, Musculoskeletal system, Intestine, testis and brain		[113]
Duckling	FBs		slightly swollen and Reddened liver, low body fat And loss of weight		[113, 114]
Turkey And poult	FB1	loss of weight	cerebral encephalomalasia	increased sphingamine to Sphingosine ratio in the Serum by disrupting Biosynthesis of ceramide and sphingolipid metabolism Hepatocellular and biliary hyperplasia	[115, 116]

Table 9. Showing the main clinical signs, performance and pathological features in poultry exposed to fumonism in selected studies

5. Fusariotoxin poisoning

Synonyms Fusariomycotoxicosis, trichothecene mycotoxicosis, T-2 toxicosis, vomitotoxicosis, zearalenone toxicosis. They are responsible for various diseases of birds in developing countries.

5.1. Aetiology

The trichothecenes include deoxynivalenol (DON), 3, monoacetyldeoxynivalenol (3-AcDON), 15, mono-acetyldeoxynivalenol (15-AcDON), nivalenol (NIV), HT-2 toxin (HT-2), neosolaniol (NEO), T-2 toxin (T-2), T-2 tetraol and T-2 triol, diacetoxyscirpenol (DAS), MAS-monoacetoxyscirpenol (MAS) and fusarenone-X.

Different fungi species of the general *Fusarium* are responsible for the production of this group of mycotoxins. Major producers of trichothecenes are *F. graminearum*, *F. culmorum*, *F. cerealis*, *F. poae*.

5.2. Occurrence of trichothecenes

Occurrence of trichothecenes in feed ingredients and poultry feed in developing countries is as a result of ubiquitous nature of the fungi which are generally found when certain cereal crops like maize, wheat, corn, millet where grown under stressful condition such as drought. These mould occur in soil, hay and especially grains undergoing microbial and possibly enzymatic degradation [20]. Direct contamination or indirect contamination of feed may occur. Direct contamination occur when the poultry feed were infected with mycotoxigenic fungi.

Indirect contamination may be as a result of fungi elaborating its toxin into the substrate, the incriminating fungi may be removed but the toxin remained in the poultry feed made from such feed ingredients.

Since moulded feed are part of the diet of animal in developing counties, thus all poultry feed are suspect and may contain different level of Trichothecus toxin.

DON has been reported in maize from Nigeria, South Africa, Argentina, Brazil Pakistan at different concentration that ranges from 0.05-2650 mg/kg [116-120].

6. Diagnosis

Diagnosis is made through observing the appropriate field signs, finding gross as well as microscopic tissue lesions, and detecting the suspected toxin in grains, forages, or the ingesta of affected animals. However, the tests required to detect these toxins are complex and few diagnostic laboratories offer tests for multiple trichothecenes in developing countries. The samples of choice include both refrigerated and frozen carcasses for necropsy examination and a representative sample of the suspected contaminated grain source. Because the toxin is produced under cold conditions, the grain sample should be frozen rather than refrigerated for shipment to the diagnostic laboratory.

Country	Feed samples	Type of toxin	Range (µg/kg)	References
Nigeria	Maize	DON	9.5-745.1	[116]
	Maize	3-acDON	0.7-72.4	
	Maize	DAS	1.0-51.0	
Poland	Wheat	DON	2-40	[121]
	Wheat	NIV	0.01	
	Wheat	ZEN	0.01-2	
Bulgaria	Wheat	DON	up to 1.8	[122]
	Wheat	ZEN	up to 0.12	
Finland	Feed and grains	DON	0.007-0.3	[123]
	Feed and grain	ZEN	0.022-0.095	
Norway	Wheat	DON	0.45-4.3	[124]
Netherland	Wheat	DON	0.020-0.231	[125]
	Wheat	NIV	0.007-0.203	
	Wheat	ZEN	0.002-0.174	
South Africa	Maize	DON	up to 1.83	[126]
	Maize	NIV	up to 0.37	
South Africa	Cereals and Animal feed	ZEN	0.05-8.0	[117]
Philippine	Maize	NIV	0.018-0.102	[127]
	Maize	ZEN	0.059-0.505	
Thailand	Maize	ZEN	0.923	[127]
Korea	Maize	DON	Mean-0.145	[128]
	Maize	NIV	Mean-168	
Vietnam	Maize powder	DON	1.53-6.51	[129]
	Maize powder	NIV	0.78-1.95	
China	Maize	DON	0.49-3.10	[129]
	Maize	NIV	0.6	
Japan	Wheat	DON	0.029-11.7	[130]
	Wheat	NIV	0.01-4.4	
	Wheat	ZEN	0.053-0.51	
New Zealand	Maize	DON	Max 3.4-8.5	[131]
	Maize	NIV	Max 4.4-7.0	
	Maize	ZEN	Max 2.7-10.5	
U.S.A	Wheat	DON	up to 9.3	[132]
Canada	Wheat and barley	DON	up to 0.5	[133]
	Wheat and barley	ZEN	up to 0.3	
Argentina	Maize	DON	0.02-4.09	[134]
	Wheat	DON	0.10-9.25	
Brazil	Wheat	DON	0.47-0.59	[119]
	Wheat	NIV	0.16-0.40	
	Wheat	ZEN	0.04-0.21	
Pakistan	Maize	Nivalenol	500-2650	[120]
	Maize	DON	136-2656	
	Maize	3-ac DON	100-850	
	Maize	ZON	1250	
	Maize	15ac DON	100	
	Maize	DAS	364-750	
	Maize	HT-2	100-500	
	Maize	T-2	143-1125	

Table 10. Occurrence of Trichothecenes in Feed Ingredients In Developing Countries

Domestic Animal	Type of toxin	clinical signs	Gross lesions	Histopathology	References
Domestic poultry		Gastrointestinal bleeding Neurological abnormality Flaccid paralysis, weakness Of the neck and wing muscles Characteristic drooping head And wings. Immune suppression which May predispose bird to Secondary bacteria infection	Skin and mucosal surface inflammation which affected subcutaneous fluid over head and neck. Multiple haemorrhage and pale area in skeletal muscle.		[136]
Chicken	DAS T-2 toxin	necrotic lesion on the tongue reduces growth rate and efficiency	Lesion in palatine, sublingual, internal angle of the tongue.		[137]
	DAS	feed refusal Haemorrhagic disorder Prolong clothing time Mild diarrhea, fatigue, ataxia Poor feathering, poor feather Quality and soft bones, Subcutaneous edema, Degeneration of neck muscle.	Enlargement of liver, and gallbladder. perivascular edema of brain, depletion of lymphocyte in thymus and bursa of fabriciuos	it affects Coagulation factors VII, prothrombin fibrinogen	[138]
	T-2 And Aflatoxin	Decrease egg production increase egg shell breakage			[139]
Duck	DON	100% mortality and high Concentration	dehydration and heamorrhage along the intestinal epithelium		[140]
	T-2 toxin	reduced body weight		altered serum And plasma Level	[141]
	DON	reduced egg production			[142]
Goose And turkey	T-2 toxin	decrease egg production	degenerative changes in ovaries Interruption of maturation of Follicle. Peritonitis, bleeding, Wrapped oviducts, necrosis and Spleen amilodiosis. Catarrhal Enteritis.		

Table 11. Showing the main clinical signs, gross and histopathological changes in some poultry species.

7. Management of avian mycotoxicosis

This involves various practises that reduces fungal contamination of feedstuff and possible use of mycotoxin binders in feed. The clinical signs seen in avian mycotoxicosis are not pathognomonic so the presence of toxin in feed with the associated clinical signs may help clinician to make effective diagnosis and withdrawal of contaminated feed can help to ameliorated the disease condition. Preventive and control of mould in feed is key to achieving control of avian mycotoxicosis.

Following good agricultural practices during both pre-harvest and post-harvest conditions would, minimize the problem of contamination by mycotoxins such as aflatoxins, ochratoxin and trichothecene mycotoxins. These include appropriate drying techniques, maintaining proper storage facilities and taking care not to expose grains.

7.1. Prevention and control of mycotoxins in stored grains and seeds

7.1.1. *Dry the feed ingredients*

Fungi cannot grow or mycotoxins be produced in properly dried foods, so efficient drying of commodities and maintenance of the dry state is an effective control measure against fungal growth and mycotoxin production.

To reduce or prevent production of most mycotoxins, drying should take place as soon after harvest and as rapidly as feasible. The critical water content for safe storage corresponds to a water activity (a_w) of about 0.7. Maintenance of feeds below 0.7 a_w is an effective technique used throughout the world for controlling fungal spoilage and mycotoxin production in foods.

Problems in maintaining an adequately low a_w often occur in the tropics, where high ambient humidity make control of commodity moisture difficult. Where grain is held in bags, systems that employ careful drying and subsequent storage in moisture-proof plastic sheeting may overcome this problem.

While it is possible to control fungal growth in stored commodities by controlled atmospheres or use of preservatives or natural inhibitors, such techniques are almost always more expensive than effective drying, and are thus rarely feasible in developing countries.

7.1.2. *Avoid grain damage*

Damaged grain is more prone to fungal invasion and therefore mycotoxin contamination. It is thus important to avoid damage before and during drying, and in storage. Drying of maize on the cob, before shelling, is a very good practice.

Insects are a major cause of damage. Field insect pests and some storage species damage grain on the head and promote fungal growth in the moist environment of the ripening grain. In storage, many insect species attack grain, and the moisture that can accumulate from their activities provides ideal conditions for the fungi. To avoid moisture and mould problems, it is essential that numbers of insects in stored grain be kept to a minimum. Such

problems are compounded if the grain lacks adequate ventilation, particularly if metal containers are used.

7.1.3. Ensure proper storage conditions

While keeping commodities dry during storage in tropical areas can be difficult, the importance of dry storage cannot be overemphasized. On a small scale, polyethylene bags are effective; on a large scale, safe storage requires well-designed structures with floors and walls impermeable to moisture. Maintenance of the water activity of the stored commodity below 0.7 is crucial.

In tropical areas, outdoor humidities usually fall well below 70% on sunny days. Appropriately timed ventilation, fan-forced if necessary, will greatly assist the maintenance of the commodity at below 0.7 aw. Ideally, all large-scale storage areas should be equipped with instruments for measuring humidity, so that air appropriate for ventilation can be selected.

Sealed storage under modified atmospheres for insect control is also very effective for controlling fungal growth, provided the grain is adequately dried before storage, and provided diurnal temperature fluctuations within the storage are minimised.

If commodities must be stored before adequate drying this should be for only short periods of no more than, say, three days. Use of sealed storage or modified atmospheres will prolong this safe period, but such procedures are relatively expensive and gaslight conditions are essential.

A proven system of storage management is needed, with mycotoxin considerations an integral part of it. A range of decision-support systems is becoming available covering the varying levels of sophistication and scale involved.

7.2. Control

Control of mycotoxin in poultry feed is important and it should be hinged on eliminating mycotoxin from the food chain. Mycotoxygenic fungi are naturally found in soil and air, which makes it difficult to prevent their contamination of agriculture commodities. Nevertheless attempts should be made to control factors that affects the growth of mycotoxigenic fungi and the subsequent toxin production. Factors which include warm temperature between (20°C to 30°C), high moisture content (20-25%), water activity (a_w) of about 0.7 a_w and relative humidity of 70% and above. These factors enhance fungi growth and mycotoxin production (143,145).

Before harvesting of crops damage to grains as a result of field insect pest and some storage species damage grains and promote fungal growth in the environment of ripening grain. Strategies used in various preventive measure in poultry feed involves good agronomic practices, detoxification of mycotoxin in grains use of mould inhibitors, genetic approach through improved breeds of plants.

7.2.1. Physical decontamination

Decontamination of mycotoxin from cereal crops used in the production of poultry feed can be classified as physical decontamination, biological decontamination and chemical decontamination [146]. [145] suggest the following method of elimination of mycotoxin in grains which include, Density segregation and floatation, cleaning and washing, sieving, dehulling, hand picking, irradiation, milling, thermal degeneration.

[147] observed that washing using distill water resulted in 65%-95% reduction of DON (16-24mg/kg) and 2% to 61% of ZEN (0.9-1.6mg/kg) in contaminated barley and corn.

Density segregation in certain liquid or fractionation by specific gravity help to segregate fungi infected and mycotoxin contamination grains used in the production of poultry feed.[148]. It was also observed that fumonisins present in broken corn kernels is about 10 fold higher than that in intact corn therefore the separation based on the size has been suggested. [147, 146]

Irradiation is also a useful tool in inactivation of some mycotoxin reported ultrasonication been used in contaminated corn without affecting the grain composition. Several workers reported the use of Gamma irradiation to reduce Zearelenone DON and T-2 toxin in corn, wheat. [148-150]

7.2.2. Biological decontamination

This involved systemic degradation of toxins leading to a less toxin product. [151] reports a fungus yeast *Expoliata spinnifera* was able to grow on fumonisin B1 as a sole of carbon source. The hydrolysis of fumonisin B1 yields free Tricarboxylic acid and aminopental, the intermediate aminopentol undergo oxidative deamination. *Sacchromyces cerevisiae* ferment zearelenone converting it into beta-zearelenol, which has less activity compared to the parent compound.[152]

Feed additive like mycotoxin inactivates trichothecenes by enzymatic decontamination of the 12-13 epoxy ring, and zearelenone by the enzymatic opening of the lactone ring [148-152]

7.2.3. Detoxification of mycotoxin feed

Moist ozone and dry ozone were able to reduce DON concentration in contaminated corn up to 90% and 70% respectively [153]. [154] reports 79% reduction in fumonism level in corn.

Author details

Adeniran Lateef Ariyo and Ajagbonna Olatunde Peter

Department of Physiology and Biochemistry, Faculty of Veterinary Medicine, University of Abuja, Nigeria

Sani Nuhu Abdulazeez

Department of Veterinary Pathology, Faculty of Veterinary Medicine, University of Abuja, Nigeria

Olabode Hamza Olatunde

Department of Veterinary Microbiology, Faculty of Veterinary Medicine, University of Abuja, Nigeria

8. References

- [1] Simons P. Global production, Consumption and International market of Poultry meat and eggs. World's Poultry Science Association (WPSA) Poultry Seminar, Lonovala, In September 12 2009
- [2] Moss M O. Mycotoxic fungi. In: Microbial Food Poisoning. Elley, A. R. (Ed) Chapman and Hall. London, Glassgow, New York, Tokyo, Melbourne, Madras. 1992 ; Pp 73-106.
- [3] D'Mello JPF. Mycotoxins in cereal grains, nuts and other plant products. In: Food Safety Contaminants and Toxins. D'Mello, J P. F.(Ed).Cromwell Press, Trowbridge, UK; 2003. 65-90.
- [4] Pitt J I. What are mycotoxins? Mycotoxin News 7:1.1996
- [5] Scimeca J A. Naturally occurring orally active dietary carcinogens. In: Handbook of humantoxicology. Massaro, E. J. (Ed), Boca Raton, New York, U.S.A.; 1997. Pp 409-466.
- [6] van Egmond HP, Speijers GJA. Survey of data on the incidence and levels of ochratoxin A in food and animal feed worldwide. Natural Toxins 1994; 3: 125-144.
- [7] Bhatnagar D, Yu J. Ehrlich K. C. Toxins of filamentous fungi. Chemical Immunology 2002; 81: 167-206.
- [8] FA O 1983. Post harvest losses in quality of food grains. Food and Agriculture Organisation (Food and Nutrition Paper No 29; 1983. p. 103.
- [9] DeVries JW., Trucksess MW., Jackson LS. Mycotoxins and Food Safety. Kluwer Academic/Plenum Publishers, New York, NY, USA. 2002.
- [10] Blount WP. Turkey X disease Turkeys. Journal of the British Turkey Association 1961; 9, 55-58.
- [11] Peterson SW, Ito Y, Horn BW, Goto T. *Aspergillus bombycis*, a new aflatoxigenic species and genetic variation in its sibling species, *A. nomius*. Mycologia 2001; 93, 689–703.
- [12] Ito Y, Peterson SW, Wicklow DT, Goto T. *Aspergillus pseudotamarii*, a new Aflatoxin producing species in *Aspergillus* section *Flavi*. Mycological Research 2001; 105, 233–239.
- [13] Frisvad JC, Skouboe P, Samson RA. Taxonomic comparison of three different groups of aflatoxin producers and a new efficient producer of aflatoxin B1, sterigmatocystin and 3-O-methylsterigmatocystin, *Aspergillus rambellii* sp. nov Systematic Applied Microbiology 2005; 28, 442–453.
- [14] Frivad JC, Samson RA, Smedsgaard J. *Emericella astellata*, a new producer of aflatoxin B1, B2 and sterigmatocystin. Letters in Applied Microbiology 2004; 38, 440–445.
- [15] Mircea C, Poiata A, Tuchilus C, Agoroae L, Butnaru E, Stanescu U. Aflatoxigenic fungi isolated from medicinal herbs Toxicology Letters 2008; 180: 32-246.
- [16] Makun HA, Dutton MF, Njobeh PB, Gbodi TA, Ogbadu GH. Aflatoxin Contamination in Foods and Feeds: A Special Focus on Africa .Intech; 2012.

- [17] D'Mello JPF. Mycotoxins in cereal grains, nuts and other plant products. In: Food Safety Contaminants and Toxins. D'Mello, J P. F.(Ed).Cromwell Press, Trowbridge, UK; 2003. 65-90.
- [18] Stössel P. Aflatoxin contamination in soybeans: role of proteinase inhibitors, zinc availability, and seed coat integrity. *Applied and Environmental Microbiology* 1986; 52, 68-72.
- [19] Schindler AF. Temperature limits for production of aflatoxin by twenty-five isolates of *Aspergillus flavus* and *Aspergillus parasiticus*. *Journal of Food Protection*. 1977; 40:39-40.
- [20] Ominski KH., Marquardi RR., Sinha RN., Abramson D . Ecological aspects of growth and mycotoxin production by storage fungi. In: Miller, J.D and Trenholm, HL. *Mycotoxins in grains: Compounds other than aflatoxins*. Eagan Press, St. Paul Minnesota, USA.1994; 287-314.
- [21] Codex Alimentarius Commission. Code of Practice for the Prevention and Reduction of Aflatoxin Contamination in Peanuts. 2004. Retrieved August, 2012 from http://www.codexalimentarius.net/download/standards/10084/CXC_055_2004e.pdf.
- [22] Kaaya NA., Warren HL. A review of past and present research on aflatoxin in Uganda *African Journal of Food Agriculture Nutrition and Development* 2005, 18 pages (On Line Free Access)
- [23] Agboola S.D 1992. Post harvest technologies to reduce mycotoxin contamination of food crops. In Z.S.C Okoye (ed) *Book of proceedings of the first National Workshop on Mycotoxins held at University Jos, on the 29th November, 1990*, 73-88.
- [24] Anjum MA., Khan SH., Sahota AW, Sardar R. Assessment of Aflatoxin B1 in commercial poultry feed and feed ingredients. *The journal of animal and plant sciences* 2012; 22, 268-272.
- [25] Odoemelam SA., Osu CI. Aflatoxin B1 contamination of some edible grains marketed in Nigeria. *E-Journal of Chemistry* 2009; 6 (2):308-314.
- [26] Adebayo-Tayo BC, Ettah AE. Microbiological quality and aflatoxin B1 level in poultry and livestock feeds. *Nigerian Journal of Microbiology*, 2010; 24(1): 2145 – 2152.
- [27] Kang'ethe EK., Lang,a KA. Aflatoxin B1 and M1 contamination of animal feeds and milk from urban centers in Kenya. *African Health Sciences* 2009; 9 (4): 218-226
- [28] Mngadi PT, Govinden R, Odhav B. Co-occurring mycotoxins in animal feeds. *African Journal of Biotechnology* 2008; 7 (13): 2239-2243.
- [29] Zinedine, A Juan, C Soriano, JM Moltó, JC Idrissi, L and Mañes. J. Limited survey for the occurrence of aflatoxins in cereals and poultry feeds from Rabat, Morocco *International Journal of Food Microbiology* 2007; 115, 124-127.
- [30] Matumba L, Monjerezi M, Khonga EB, Lakudzala DD. Aflatoxins in sorghum, sorghum malt and traditional opaque beer in southern Malawi. *Food Control* 2011; 22, 266-268
- [31] Muthomi JW, Ndung'u JK, Gathumbi JK, Mutitu EW, Wagacha JM. The occurrence of *Fusarium species* and mycotoxins in Kenyan wheat. *Crop Protection* 2008; 27, 1215- 1219.
- [32] Ghali R, Khelifa KH, Ghorbel H, Maaroufi K, Hedilli A. Incidence of aflatoxins, ochratoxin A and zearalenone in Tunisian foods. *Food Control* 2008; 19, 921-924.

- [33] Riba A., Bouras N., Mokrane S., Mathieu F., Lebrihi A., Sabaou N. *Aspergillus section Flavi* and aflatoxins in Algerian wheat and derived products. Food and Chemical Toxicology 2010; 48, 2772–2777
- [34] Mashinini K, Dutton MF. The incidence of fungi and mycotoxins in South Africa wheat and wheat-based products Journal of Environmental Science and Health Part B, 2006; 41:285-296
- [35] Akrobortu DE. Aflatoxin contamination of maize from different storage locations in Ghana. An M.Sc. Thesis submitted to the Department of Agricultural Engineering, Kwame Nkrumah University of Science and Technology, Ghana. 2008; 27-32.
- [36] Makun HA, Gbodi TA, Akanya HO, Sakalo AE, Ogbadu HG. Fungi and some mycotoxins contaminating rice (*Oryza sativa*) in Niger state, Nigeria. African Journal of Biotechnology 2007; 6(2):99–108
- [37] Njobeh PB., Dutton MF., Aberg AT., Haggblom P. Estimation of multi-mycotoxin contamination in South African compound feeds. Toxins 2012; 4, 836-848.
- [38] Binder EM. managing the risk of mycotoxins in modern feed production Animal Science and Technology. 2007; 133:149-166.
- [39] Anjum AD. Outbreak of infectious bursar disease in vaccinated chickens due to aflatoxicosis. Indian Veterinary Journal 1994; 71:322-324.
- [40] Asplin FD, Carnaghan RBA. The toxicity of certain groundnut meals for poultry with special reference to their effect on ducklings and chickens. Veterinary Record 1961; 73:1215-1219.
- [41] Siller WG, Ostler DC. The histopathology of an entero- hepatic syndrome of turkey poults. Veterinary Record 1961; 73:134-138.
- [42] Wannop CC. The histopathology of turkey "x" disease in Great Britain. Avian Disease 1961; 5:371-381.
- [43] Archibald RM, Smith HJ, Smith JD. Brazilian groundnut toxicosis in Canadian broiler chickens. Canadian Veterinary Journal 1962; 3:322-325.
- [44] Batra PA, Pruthi K, Sandana IR. Effect of aflatoxin B1 on the efficacy of turkey herpesvirus vaccine against Marek's disease. Research in Veterinary Science 1991 ;51:115-119.
- [45] Sharlin IS, Howarth B, Jr. Wyatt RD. Effect of dietary aflatoxin on reproductive performance of mature White Leghorn males. Poultry Science 1980; 59:1311-1315.
- [46] Ortalatli M, Ciftci MK, Tuzcu M, Kaya A. The effects of aflatoxin on the reproductive system of roosters. Research in Veterinary Science 2002; 72:29-36.
- [47] Larsen C, Acha M, Ehrich M. Research note: Chlortetracycline and aflatoxin interaction in two lines of chicks. Poultry Science 1988; 67: 1229-1232.
- [48] Smith IW, Hill CH, Hamilton PB. The effect of dietary modifications on aflatoxicosis in the broiler chicken. Poultry Science 1971; 50:768-774.
- [49] Basic Food Safety for Health Workers.WHO. 1999
- [50] Chipley IR, Mabee MS, Applegate KL, Dreyt"us.i MS. Further characterization of tissue distribution and metabolism of (I4C) aflatoxin BI in chickens. Applied Microbiology 1974; 28:1027-1029.

- [51] Patterson OSP, Roberts BA. The in vitro reduction of aflatoxins Bland 82 by soluble avian liver enzymes. Food Additive and Contaminant 1971; 9:821-831.
- [52] Dutton ME. Fumonisin, Mycotoxins of Increasing Importance: Their Nature and Their Effects. Pharmacology and Therapeutics. 1996; 70(2) 137-161
- [53] Harland EC, Cardeilhac PT. Excretion of carbon14-labeled aflatoxin B₁ via bile, urine, and intestinal contents of the chicken. American Journal of Veterinary Research 1975; 36:909-912.
- [54] Sawhney DS, Vadehra DV, Baker RC. The metabolism of [14C] aflatoxins in laying hens. Poultry Science 1973; 52:1302-1309.
- [55] Oliveira CA, Kobashigawa E, Reis TA, Mestieri L, Albuquerque R, Correa B. Aflatoxin B₁ residues in eggs of laying hens fed a diet containing different levels of the mycotoxin. Food Additive and Contaminant 2000; 17:459-462.
- [56] Jacobson WC, Wiseman HG. The transmission of aflatoxin B₁ into eggs. Poultry Science 1974; 53:1743-1745.
- [57] Trucksess MW, Stoloff L, Young K, Wyatt RD, Miller BL. Aflatoxinol and aflatoxins B₁ and M₁ in eggs and tissues of laying hens consuming aflatoxin-contaminated feed. Poultry Science 1983; 62:2176--2182.
- [58] Sova Z, Fukal L, Trefuy D, Prosek I, Slamova A. B₁ aflatoxin (AFB₁) transfer from reproductive organs of farm birds into their eggs and hatched young. Conference of European Agriculture 1986;7:602-603.
- [59] Lillehoj EB, Aalund O, Hald B. Bioproduction of 14C Ochratoxin A in submerged culture, Applied Environmental Microbiology. 1978; 36 : 720-723..
- [60] Foster PL, Eisenstadt E, Miller JH. Base substitution mutations induced by metabolically activated aflatoxin B₁. Proceeding of National Academic of Science 1983; 80 : 2695-2698.
- [61] Raney KD, Meyer DJ, Ketterer B, Harris TM, Guengerich FP. Glutathione conjugation of aflatoxin B-1 exo- and endo-epoxides by rat and human glutathione-S-transferases. Chemical Research Toxicology 1992; 5 :470-478.
- [62] Bababunmi EA, Thabrew I, Bassir O. Aflatoxin induced coagulopathy in different nutritionally classified animal species. World Review of Nutrition and Diet 1997; 34 : 161-181.
- [63] Shen M, Ong CN, Shi CY. Involvement of reactive oxygen species in aflatoxin B₁-induced cell injury in cultured rat hepatocytes. Toxicology 1997; 99 :115-123
- [64] Bonsi GP, Agusti-Tocco M, Palmery M, Giorgi M. Aflatoxin B₁ is an inhibitor of cyclic nucleotide phosphodiesterase activity. General Pharmacology 1991; 32: 615-619.
- [65] Dwivedi P, Burns RB. The natural occurrence of ochratoxin A and its effects in poultry. A review. Part I. Epidemiology and toxicity. World Poultry Science Journal 1986; 42:32-47.
- [66] Rosa C A R, Ribeiro J M M, Fraga M J, Gatti M, Cavaglieri L R, Magnoli C E, Dalcero A M, Lopez C WG. Mycoflora of poultry feeds and ochratoxin-producing ability of isolated *Aspergillus* and *Penicillium* species. Veterinary Microbiology 2006; 113:89-96.

- [67] Rizzo I, Vedoya G, Maurotto S, Haidukowski M, Varkavsky E. Assessment of oxinogenic fungi on Argentinean medicinal herbs, *Microbiology Research* 2004; 159:113–120.
- [68] Sawhney DS, Vadehra DV, Baker RC. The metabolism of [14C] aflatoxins in laying hens. *Poultry Science* 1973; 52:1302-1309.
- [69] Larsen TO, Svendsen A, Smedsgaard J. Biochemical characterization of ochratoxin A-producing strains of the genus *Penicillium*, *Applied Environmental Microbiology* 2001; 67: 3630–3635.
- [70] Castella G, Larsen TO, Cabanes J, Schmidt H, Alboresi A, Niessen L, Geisen R. Molecular characterization of ochratoxin A producing strains of the genus *Penicillium*, *Systemic and Applied Microbiology* 2002; 25 : 74–83
- [71] Kozakiewicz Z. *Aspergillus* species on stored products, Mycological paper no. 161, CAB International Mycological Institute, Kew, UK, 1989, p. 1.
- [72] Pardo E, Marín S, Ramos AJ, Sanchis V. Effect of water activity and temperature on mycelial growth and ochratoxin A production by isolates of *Aspergillus ochraceus* on irradiated green coffee beans, *Journal of Food Protection* 2005; 68 : 133–138.
- [73] Basic Food Safety for Health Workers.WHO. 1999
- [74] Abarca M, Bragault M R, Castella G, Cabanes F J. Ochratoxin A production by strains of *Aspergillus niger* var *niger*, *Applied Environmental Microbiology* 1994; 60: 2650–2652.
- [75] Belli N, Marin S, Sanchis V, Ramos AJ. Influence of water activity and temperature on growth of isolates of *Aspergillus section nigri* obtained from grapes, *International Journal Food Microbiology* 2004; 96 : 19–27.
- [76] Teren J, Varga J, Hamari Z. Immunochemical detection of Ochratoxin A in black *Aspergillus* strain, *Mycopathologia* 1996;134 :171–176.
- [77] Mitchell D, Parra R, Aldred D, Magan N. Water and temperature relations of growth and ochratoxin A production by *Aspergillus carbonarius* strains from grapes in Europe and Israel, *Journal Applied Microbiology* 2004; 97: 439–445.
- [78] Rosa C A R, Ribeiro J M M, Fraga M J, Gatti M, Cavaglieri L R, Magnoli C E, Dalcero A M, Lopez C WG. Mycoflora Of poultry feeds and ochratoxin-producing ability of isolated *Aspergillus* and *Penicillium* species. *Veterinary Microbiology* 2006; 113:89-96.
- [79] Accensi F, Abarca M.L, Cabanes FJ. Occurrence of *Aspergillus* species in Mixed feeds and component raw materials and their ability to produce ochratoxin A, *Food Microbiology* 2004; 21 : 623–627.
- [80] Beg MU, Al-Mutairi M, Beg KR, Al-Mazeedi HM, Ali LN, Saeed T, 2006. Mycotoxins in poultry feed in Kuwait. *Archive of Environmental Contaminant. Toxicology* 2006; 50, 594–602.
- [81] Cabanes FJ, Accensi F, Bragault MR, Abarca ML, Minguez G, Pons S. What is the source of ochratoxin A in wine? *International Journal Food Microbiology* 2002;79 : 213–215.
- [82] Samson RA, Houbroken JAMP, Kujipers AFA, Frank JM, Frisvard, JC. New ochratoxin A or sclerotium producing species in *Aspergillus section Nigri*, *Study.Mycology* 2004;50 : 45–61.

- [83] Moss MO. Mode of formation of ochratoxin A, Food Additive and Contaminant. 1996;13 : 5–9.
- [84] ayman P, Baker JL, Doster MA, Michailidis TJ, Mahoney NE. (2002).Ochratoxin production by the *Aspergillus ochraceus* group and *Aspergillus alliaceus*, Applied Environmental Microbiology 2002; 68 : 2326–2329.
- [85] Varga J, Rigo K, Toth B, Teren J, Kozakiewicz Z. Evolutionary relationships among *Aspergillus* species producing economically important mycotoxins, Food Technology and Biotechnology 2003; 41 : 29–36.
- [86] Callaghan JO, Caddick MX, Dobson AD. A polyketide synthase gene required for ochratoxin A biosynthesis in *Aspergillus ochraceus*, Microbiology 2003;149 : 3485–3491.
- [87] Beg MU., Al-Mutairi M., Beg KR., Al-Mazeedi HM., Ali LN., Saeed T. Mycotoxins in poultry feed in Kuwait. Archive of Environmental Contaminant & Toxicology. 2006;50, 594–602
- [88] Figueroa S, Centeno, S, Calvo MA, Rengel A, Adelantado C. Mycobiodata and concentration of ochratoxin A in concentrated poultry feed from Venezuela. Pakistan Journal of Biological Science 2009; 12: 589–594
- [89] Dalcero A, Magnoli C, Hallak C, Chiacchiera SM, Palacio G, Rosa CAR. Detection of ochratoxin A in animal feeds and capacity to produce this mycotoxin for *Aspergillus section Nigri* in Argentina. Food Additives and Contaminants. 2002; 19 (11), 1065–1072.
- [90] Zafar, F., Yasmin, N., Hassan, R., Naim, T., Qureshi, A.A. A study on the analysis of ochratoxin A in different poultry feed ingredients. Pakistan Journal of Pharmaceutical Sciences 2001;14, 5–7.
- [91] Duarte SC, Lino CL, Pena A. Ochratoxin A in feed of food-producing animals: An undesirable mycotoxin with health and performance effects. Veterinary Microbiology 2011; 154, 1–13.
- [92] Bozzo G., Bonerba E, Ceci E, Colao V, Tantillo G. Determination of ochratoxin A in eggs and target tissues of experimentally drugged hens using HPLC–FLD. Food Chemistry 2011; 126, 1278–1282.
- [93] Sakthivelan SM, Rao GVS. Effect of ochratoxin A on body weight, feed intake and feed conversion in broiler chicken. Veterinary Medical Int. 2010; 1–4.
- [94] Bozzo G, Ceci E, Bonerba E, Desantis S, Tantillo G. Ochratoxin A in laying hens: high-performance liquid chromatography detection and cytological and histological analysis of target tissues. Journal of Applied Poultry Research 2008; 17, 151–156.
- [95] Elaroussi MA, Mohamed FR, Barkouky EME, Atta AM, Abdou AM, Hatab MH. Experimental ochratoxicosis in broiler chickens. Avian Pathology 2006; 35, 263–269.
- [96] International Agency for Research on Cancer. 1993. Toxins derived from *Fusarium moniliforme*: fumonisins B₁ and B₂ and *Fusarium C*, p. 445–466. In IARC Monographs on the evaluation of the carcinogenic risks to humans: some naturally occurring substances: food items and constituents, heterocyclic aromatic amines and mycotoxins, vol. 56. International Agency for Research on Cancer, Lyon, France.

- [97] Marasas WFO. Fumonisin: history, worldwide occurrence and impact, p. 1-17. In L. S. Jackson, J. W. DeVries, and L. B. Bullerman (ed.), *Fumonisin in food*. Plenum Press, New York, N.Y; 1996.
- [98] Branham, B. E., and R. D. Plattner. 1993. Alanine is a precursor in the biosynthesis of fumonisin B₁ by *Fusarium moniliforme*. *Mycopathologia* 124:99-1
- [99] Marín S, Sanchis V, Magan N. Water activity, temperature, and pH effects on growth of *Fusarium moniliforme* and *Fusarium proliferatum* isolates from maize. *Canadian Journal of Microbiology* 1995a; 41:1063-1070.
- [100] Marín S, Sanchis V, Vinas I, Canela R, Magan N. Effect of water activity and temperature on growth and fumonisin B₁ and B₂ production by *Fusarium proliferatum* and *F. moniliforme* in grain. *Letter of Applied Microbiology* 1995b; 21:298-301.
- [101] Sydenham EW, Shephard G S, Thiel P G, Marasas, W E, Rheeder J I, Peralta C E, Gonzalez H L, Resnik S L. Fumonisin in Argentinian field trial corn. *Journal of Agriculture and Food Chemistry* 1993; 41: 891-895.
- [102] Doko M B, Rapior S, Visconti A, Schjoth J E. Incidence and levels of fumonisin contamination in maize endotypes grown in Europe and Africa. *Journal of Agriculture and Food Chemistry* 1995. 43: 429-434.
- [103] Sydenham E W, Shephard G S, Thiel PG, Marasas W F 0, Stockenstrom S. Fumonisin contamination of commercial corn-based human foodstuffs. *Journal of Agriculture and Food Chemistry* 1991; 39: 2014-2018.
- [104] Ueno Y, Aoyama S, Suglaru Y, Wang D S, Lee U S, Hirooka EY, Ham S, Karki T, Chen G, Yu S Z. A limited survey of fumonisins in corn and corn-based products in Asia countries. *Mycotoxin. Research* 1993; 9: 27- 34
- [105] Julian A M, Wareing IW, Phillips SI, Medlock V F, MacDonald M V, de1 Rio L E. Fungal contamination and selected mycotoxins in pre and postharvest maize in Honduras. *Mycopathologia* 1995; 129: 5-16.
- [106] Doko MB., Canet C., Brown N., Sydenham EW., Mpuchane S., Siame B A. Natural co-occurrence of fumonisins and zearalenone in cereals and cereal-based foods from eastern and southern Africa. *Journal of Agriculture and Food Chemistry*.1996; 44, 3240-3243.
- [107] Julian A M, Wareing IW, Phillips SI, Medlock V F, MacDonald M V, de1 Rio L E. Fungal contamination and selected mycotoxins in pre and postharvest maize in Honduras. *Mycopathologia* 1995; 129: 5-16.
- [108] Pineiro MS., Silva GE., Scott PM., Lawrence GA., Stack ME. Fumonisin level in Uruguayan corn products. *Journal of AOAC International* 1997; 80, 825-828.
- [109] Viquez OM., Castell-Pere M.E., Shelby RA. Occurrence of fumonisin B₁ in maize grown in Costa Rica. *Journal of Agriculture and Food Chemistry*. 1996; 44, 2789-2791.
- [110] Prathapkumar S H, RaoVS, Paramkishanand UR, Bha RV. Disease outbreak in laying hens arising from the consumption of fumonisin-contaminated food. *Br Poultry Science* 1997; 38:475-479.

- [111] Chatterjee D S, Mukhejee K, Dey A. Nuclear disintegration in chicken peritoneal macrophages exposed to fumonisin B1 from Indian maize. *Applied Microbiology* 1995; 20:184-185.
- [112] Dombrink-Kurtzman MA, Javed T, Bennett G A, Richard J L, Cow L, Buck W B. Lymphocyte cytotoxicity and erythrocytic abnormalities induced in broiler chicks by fumonisins B, and B1 and moniliformin from *Fusarium proliferatum*. *Mycopathologia* 1993; 124: 47-54.
- [113] Javed I, Bennett G A, Richard J L, Dombrink Kurtzman MA, Cote L M, Buck W B. Mortality in broiler chicks on feed amended with *Fusarium proliferatum* culture materials or with purified fumonisin B and moniliformin. *Mycopathologia*; 1993; 123: 171-184.
- [114] Marasas, VI E. Mycotoxicological investigations on corn produced in esophageal cancer areas in Transkei. In: *Cancer of the Esophagus*, 1982 Vol. 1, pp. 29-40, Pfeiffer, C J (ed.) CRC Press Inc, Boca Raton.
- [115] Vesonder R, Halihurton J, Golinski P. Toxicity of field samples and *Fusarium moniliforme* from feed associated with equine- leucoencephalomalacia *Archive of Environmental Contaminant* 1989; 18: 439-442.
- [116] Adejumo TO, Hettwer U, Karlovsky P. Occurrence of *Fusarium* species and trichothecenes in Nigerian maize. *International Journal of Food Microbiology*. 2007; 116: 350-357
- [117] Dutton MF., Kinsey A. A note on the occurrence of mycotoxins in cereals and animal feedstuffs in Kwazulu Natal, South Africa 1984-1993. *South Africa Journal of Animal Science*. 1996; 26, 53-57.
- [118] Pacin AM., Resnik SL., Neira MS., Molto G., Martinez E. Natural occurrence of deoxynivalenol in wheat, wheat flour and bakery products in Argentina. *Food Additive and Contaminant*. 1997; 14, 327-331.
- [119] Furlong EB., Soares LMV., Lasca CC., Kohara EY., 1995. Mycotoxins and fungi in wheat harvested during in test plots in the state of Sao Paulo Brazil. *Mycopathologia* 1990; 131, 185-190.
- [120] Salma K, Nafeesa Q H, Tahira I, Sultana N, Sultana K, and Ayub N. Natural occurrence of aflatoxins, zearalenone and Trichothecenes in maize grown in Pakistan, *Pakistan Journal of Botanical* 2012; 44(1): 231-236
- [121] Perkowski J., Plattner RD., Golinski P., Vesonder RF. Natural occurrence of deoxynivalenol, 3-acetyldeoxynivalenol, 15-acetyl-deoxynivalenol, nivalenol, 4,7-dideoxynivalenol and zearalenone in Polish wheat. *Mycotoxin Research*. 1990; 6, 7-12.
- [122] Vrabcheva T., Gebler R., Usleber E., Martlbauer E. First survey on the natural occurrence of *Fusarium* mycotoxins in Bulgarian wheat. *Mycopathologia*. 1996; 136, 47-52.
- [123] Hietaniemi V., Kumpulainen J. Contents of *Fusarium* toxins in Finnish and imported grains and feeds. *Food Additive and Contaminants*. 1991; 8, 171-182.
- [124] Langseth W., Elen O. Differences between barley, oats and wheat in the occurrence of deoxynivalenol and other trichothecenes in Norwegian grain. *J. Phytopathology*. 1996; 144, 113-118.

- [125] Tanaka T., Yamamoto S., Hasegawa A., Aoki N., Besling JR., Sugiura, Y., Ueno Y. A survey of the natural occurrence of *Fusarium* mycotoxins, deoxynivalenol, nivalenol and zearalenone, in cereals harvested in The Netherlands. *Mycopathologia*. 1990; 110, 19-22.
- [126] Rheeder JP., Sydenham EW., Marasas WFO., Thiel PG., Shephard GS., Schlechter M., Stockenstrom S. .. Viljoen JH. Fungal infestation and mycotoxin contamination of South African commercial maize harvested in 1989 and 1990. *South Africa Journal of Science*. 1995; 91, 127-131.
- [127] Yamashita A., Yoshizawa T., Aiura Y., Sanchez PC., Dizon EI., Arim, RH., Sardjono, , 1995. *Fusarium* mycotoxins (fumonisins, nivalenol and zearalenone) and aflatoxins in corn from southeast Asia. *Biological science. Biotechnology. Biochemistry*. 1995; 59, 1804-1807.
- [128] Ryu JC., Yang JS., Song YS., Kwon OS., Park J., Chang IM. 1996. Survey of natural occurrence of trichothecene mycotoxins and zearalenone in Korean cereals harvested in 1992 using gas chromatography/ mass spectrometry. *Food Additive and Contamination*. 1996; 13, 333-341.
- [129] Wang DS., Liang YX., Iijima K., Sugiura Y., Tanaka T., Chen G., Yu SZ., Ueno Y. Cocontamination of mycotoxins in corn harvested in Haimen, a high risk area of primary liver cancer in China. *Mycotoxins* 1995b; 41, 67-70.
- [130] Yoshizawa T. Geographic difference in trichothecene occurrence in Japanese wheat and barley. *Bull. Institute of. Comprehensive Agricultural Science. Kinki University* 1997; 5, 23-30.
- [131] Lauren DR., Jensen DJ., Smith WA., Dow BW., Sayer ST. Mycotoxins in New Zealand maize: a study of some factors influencing contamination levels in grain New Zealand. *Journal of Crop and Horticultural Science*. 1996; 24, 13-20.
- [132] Fernandez C., Stack ME., Musser SM. Determination of deoxynivalenol in 1991 U.S. winter and spring wheat by high-performance thin-layer chromatography. *Journal of AOAC Institute*. 1994; 77, 628-630.
- [133] Stratton GW., Robinson AR., Smith HC., Kittilsen L., Barbour M. Levels of five mycotoxins in grains harvested in Atlantic Canada as measured by high performance liquid chromatography. *Architecture of Environmental Contamination and Toxicology*. 1993; 24, 399-409.
- [134] Scott PM. Multi-year monitoring of Canadian grains and grain-based foods for trichothecenes and zearalenone. *Food Additive and Contaminant*. 14, 333-339.
- [135]
- [136] Diaz G.J., Squires E. J., Julian R. J. And Boermans H.J. Individual and combined effects of T-2 toxin and DAS in laying hens and Broiler *Poultry Science* 1994; 35; 393 - 405
- [137] Brake J, Hamilton PB, Kittrell R S. Effects of the Trichothecene Mycotoxin Diacetoxyscirpenol on Feed Consumption, Body Weight, and Oral Lesions of Broiler Breeders. *Poultry Science*, 2000; 79:856–863
- [138] Ademoyero A. A, Hamilton PB. Influence of degree of acetylation of scirpenol mycotoxins on feed refusal by chickens. *Poultry Science* 1989; 68:854–856.

- [139] Jakić-Dimić D, Nešić K, Šefer D. Mycotoxicoses of poultry caused by Trichothecenes. *Biotechnology in Animal Husbandry* 2011; 27 (3), p 713-719. DOI: 10.2298/BAH1103713J
- [140] Rafai P, Pettersson H, Bata A, Papa Z, Glavits S, Tubolys S, Vanyi A, Sooss P. Effect of dietary T-2 fusariotoxin concentrations on the health and production of white Pekin duck broilers. *Poultry Science* 2000; 79, 1548-1556.
- [141] Scott ML., Dean WF. In: *Nutrition and Management of Ducks*. M.& Scott, Ithaca, NY; 1991. 150-166.
- [142] Vanyi A, Bata A, Kovacs F. Effects of T-2 toxin treatment on the egg yield and hatchability in geese. *Acta Veterinaria Hungarica* 1994; 42, 1, 79-85.
- [143] Langseth W., Hqie R. Gullord M. The influence of cultivars, location and climate on deoxynivalenol contamination in Norwegian oats (1985-1990). *Acta Agriculture Scandinavica. Section B: Soil and Plant Science*, 1995 45: 63-67.
- [144] Ma H., Zhon M., Liu Z. Liu W. Progress on genetic improvement for resistance to wheat scab in KLA. *Journal of Applied Genetics*, 2002; 43: 259-266.
- [145] Negedu A., Atawodi SE. Ameh JB., Umoh VJ., Tanko HY. Economic and health perspectives of mycotoxins: a review. *Continental Journal of Biomedical Sciences* 2011; 5 (1): 5 – 26.
- [146] Gbodi TA., Makun HA., Kabiru YA., Ogbadoyi E., Tijani SA., Lawal SA., Bayero RA. Effect of local processing methods on aflatoxin contents of some common Nigerian foods prepared from artificially contaminated maize, rice and sorghum. *Journal of Agricultural Science. Mansoura University*. 2001; 26: 3759 -3769.
- [147] Desjardins AE., Manandhar GG., Plattner RD., Maragos CM., Shrestha K.. McCormick SP. (2000). Occurrence of *Fusarium* species and mycotoxins in Nepalese maize and wheat and the effect of traditional processing methods on mycotoxin levels. *Journal of Agricultural food Chemistry*. 2000; 48:1377 – 1383.
- [148] Scott PM. Mycotoxins transmitted into beer from contaminated grains during brewing. *Journal of the Association of Official Analytical Chemists International*. 1996; 79:675.
- [149] Diaz DE., Hopkins BA., Leonard LM. Hagler WM. and Whitlow LW. Effects of fumonisin on lactating dairy cattle. *Journal Dairy Science*. 2000. 83 (Abstract): 1171.
- [150] Fanelli C., Taddei F., Jestoi M. and Visconti A. Use of resveratrol and BHA to control fungal growth and mycotoxin production in wheat and maize seeds. *Aspect of Applied Biology*. 2003. 68: 63-71.
- [151] Duvick J., Rood TA., Maddox JR., and Gilliam J. Advances in ochratoxin A biosynthesis. Book of Abstracts, International Conference on “Advances on genomics, biodiversity and rapid systems for detection of toxigenic fungi and mycotoxins” September 26-29, 2006. Monopoli (Bari), Italy Pp.39 Retrieved from <http://www.Ispa.onr.n/mycoglobe-2006>. 1998.
- [152] Leggot NL and Shephard G. Patulin in South African commercial apple products. *Food Control*. 2001. 12 (2) 73-76
- [153] Diaz DE., and Smith TK. Mycotoxin sequestering agents: Practical tools for the neutralization of mycotoxins in the mycotoxin Blue Book. D. Diaz, ed. Nottingham Univ. Press, Nottingham, U. K. 2005.Pp. 313-339

- [154] Park JW., Scott PM., Lau BPY., and Levis D A. Analysis of heat processed corn foods for fumonisins and bound fumonisns. *Food Additives and Contaminants*. 2004; 21:1168-1178.

IntechOpen

IntechOpen