We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

The Role of Furin in the Development of Skin Cancer

Rethika Ravi and Terrence J Piva

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/55569>

---

1. Introduction

Skin cancer represents a major, and growing, public health problem, and is the most common type of cancer observed in Caucasians [1-3]. The three most common forms of skin cancer are basal cell carcinoma (BCC), squamous cell carcinoma (SCC) and melanoma. BCC and SCC are together known as non-melanoma skin cancers (NMSC), and are both derived from keratinocytes whereas melanomas are derived from melanocytes [3-6]. SCCs can undergo metastasis; BCCs rarely do, while melanomas can be highly metastatic [5, 6].

The ultraviolet (UV) radiation component of sunlight is acknowledged to be the main carcinogen implicated in the formation of skin cancer. UV radiation can be divided into three components: UVC (100-280 nm), UVB (280-320 nm) and UVA (320-400 nm). Ozone depletion, seasonal and weather variations affect the amount of UV radiation reaching the Earth's surface [7]. UVC and most of the UVB radiation emitted from the sun is blocked from reaching the Earth's surface by the ozone layer. The component of UV light that reaches the Earth's surface consists of 90-95% UVA and 5-10% UVB [3, 8]. The penetration of shorter-wavelength UVB radiation is predominantly confined to the epidermis while UVA penetrates into the dermis because of its longer wavelength [9].

UVB can cause sunburn, inflammation, DNA mutations and membrane damage as well as skin cancer [8, 10, 11]. It is known that UVB directly damages DNA and can induce Reactive Oxygen Species (ROS) by interactions with chromophores in the skin [12]. The DNA damage caused by UVB irradiation typically results in the formation of cyclobutane pyrimidine dimers (CPD) and pyrimidine (6-4) photoproducts. The mutations are frequently found in the p53, p16, PTCH and INK4 α /CDKN2A genes of skin cancer patients [13, 14]. Inflammation plays a significant role in creating an environment where cells possessing mutated DNA can become carcinogenic. UVA can cause premature skin ageing, wrinkle formation, blotching and induces sunburn cell formation in the epidermis, as well as skin cancer [8, 10, 15]. It affects keratinocytes

at a transcriptional level by altering the expression of genes involved in apoptosis, cell cycle, DNA repair, signal transduction, RNA processing and translation, and metabolism [9]. UVA can cause DNA damage by generating ROS [12] resulting in genomic damage e.g. single-stranded breaks, protein-DNA crosslinks, and oxidative base damage (i.e. 8-oxo-7,8-dihydroxyguanine) [16]. It can also initiate signal transduction pathways [13, 17] as well as inducing the expression of cytokines such as Interleukin (IL)-6, heme oxygenase-1, and cyclo-oxygenase [18] as well as inflammatory mediators such as tumor necrosis factor- α (TNF α) [15, 19].

While UVB has been thought to be the main contributor toward skin cancers, based largely on the DNA action spectrum of UV radiation, UVA has more recently been acknowledged as playing an important role in this process [9, 11, 20]. While UVA does not produce an inflammatory response like that of UVB, it produces ROS and as such activates many of the same signalling pathways [13]. It is clear that doses of UVB, UVA and solar stimulated UV that are too low to cause inflammation can induce mutations in epidermal cells. However, this does not exclude a role for ROS from inflammatory cells contributing to skin carcinogenesis, but it may be important for tumour progression [20]. UV-induced inflammation seen in the skin involves the action of many molecules. Of these inflammatory molecules TNF α plays a major role in UV-irradiated inflammation in the skin [15, 19]. TNF α is cleaved from its membrane-bound precursor by the action of the metalloprotease, Tumour Necrosis Factor- α Converting Enzyme (TACE) [21, 22]. While UVB radiation increases the release of TNF α from skin cells, it is not known whether this is due to increased TACE activity and/or expression. However, before TACE is activated it is cleaved from its proform by the action of furin, a proprotein convertase [23, 24]. Furin can cleave other proteases such as matrix metalloproteases (MMPs) [23, 24]. Exposure to UVB radiation also increases MMP activity in skin cells [25]. While furin is expressed in skin cells, the effect UV radiation has on its expression and/or activity and that of the proteases it activates is not fully known. As a result of elevated furin levels in a mutated cell, enhanced TACE activity would see an increase in the secretion of TNF α thereby sustaining a localised inflammatory environment allowing for the development of carcinogenic cells. As furin activates MMP activity, these carcinogenic cells have the potential to become metastatic. This review investigates the role that furin plays in the activation of TACE and MMPs and the effect that this has on a skin cells exposed to UV radiation, as well as that its role in cancer cells which undergoes metastasis, and how an understanding of the role played by this proprotein convertase, may assist in the design of new inhibitors which have therapeutic potential.

2. UV-induced inflammation

High doses of UV can induce inflammation in the skin that results in the appearance of macrophages and other leucocytes [15, 26]. Along with the activation of these cells, many mediators of inflammation are also seen including; prostaglandins [18], nitric oxide (NO) [27] and ROS [12, 17], and cytokines such as interleukin (IL)-1, interferon (IFN)- γ , and TNF α [14, 19, 28, 29]. ROS can cause DNA strand breaks as well as lipid peroxidation, membrane and protein damage [12, 30]. The effects of UV radiation on the levels of these proinflammatory molecules in skin cells are seen in Table 1 [8, 15, 31-33].

Inflammatory mediators such as IL-1, TNF α and IL-6 have been postulated to play a major role in both melanoma [34] and NMSC formation [26,35,36]. Male mice are known to be more sensitive to UVB-induced skin carcinogenesis than female mice [4], which is consistent with human studies showing men having a higher incidence of skin cancer than women [20]. Damian *et al.* [20] found that while women developed a larger inflammatory response to UVB radiation, men had lower antioxidant levels in the skin resulting in a higher level of oxidative damage to DNA, and were more sensitive to UV immunosuppression. This suggests that UV-induced immunosuppression and DNA damage plays a greater role in the formation of skin cancers in men compared to women [20]. IL-1 α and IL-1 β are both induced in keratinocytes exposed to UVB radiation [31, 37]. IL-1 α has been shown to enhance the expression and release of TNF α from UVB-irradiated keratinocytes [38,39], while IL-1 β enhances the expression of matrix metalloprotease (MMP)-9 in these irradiated cells [40]. Apart from IL-1 β , UVB can stimulate MMP-9 expression in human skin via the induction of Activator protein-1 (AP-1) and NF κ B activities [41].

Mediator	Produced By	Function	References
TNF α	Keratinocytes Mast cells Dermal fibroblasts Langerhans cells	Langerhans cell migration, sunburn cell information, stimulates prostaglandin (PG) synthesis, changes in adhesion molecule expression	[15, 31, 33]
IL-1 α	Keratinocytes Langerhans cells	Simulates PG synthesis, increases TNF α and IL-6, inhibited by IL-1 receptor antagonist	[15, 33]
IL-1 β	Keratinocytes Langerhans cells	Langerhans cell migration	[15, 31, 33]
IL-6	Keratinocytes, Langerhans cells	Fever Severe sunburn	[8, 15, 33]
IL-10	Macrophages Melanocytes	Blocks cytokine production by T cells, macrophages and NK cells Decreases antigen presentation, Increases IL-1 receptor antagonist	[33] [8, 15, 32, 33]
IL-12p40 (not bioactive)	Keratinocytes, Dendritic cell, Langerhans cells	Decreases Th1 response Decreases antigen presentation	[33]
IFN γ	T cells	Triggers apoptosis T-cell mediated tumour cell destruction	[33]
PGE ₂	Keratinocytes, Mast cells	Erythema Decreases antigen presentation Increases IL-4, decreases IL-12	[15, 33]
Histamine	Mast Cells	Increases release of PG Inhibit lymphocyte functions like IL-2 and IFN γ	[8, 15, 33]

Table 1. Effect of UV radiation on the expression of bioactive molecules in human skin cells

UVB radiation can increase cyclo-oxygenase (COX)-2 expression and activity in keratinocytes [20, 28, 42]. High levels of COX-2 activity have been observed in human epithelial skin cancers [43]. Nonsteroidal anti-inflammatory drugs can inhibit COX-2 activity and subsequent PGE formation in the skin, and have been used in the treatment of actinic keratosis (AK) [44], BCC, SCC and melanoma [45]. This suggests a role for COX-2 in the formation of skin cancers, and high levels of activity have been observed in many of these tumours [46].

Upregulation of TNF α is a key early response observed in keratinocytes exposed to UVB radiation [8, 38, 47] and represents an important component of the inflammatory cascade in skin. The expression of TNF α mRNA was enhanced a few hours post-UVB irradiation in both keratinocytes and dermal fibroblasts [38, 47]. IL-1 α was shown to stimulate TNF α expression in UVB-irradiated keratinocytes [47] and melanocytes [48]. While Bashir *et al.* [38] observed that TNF α expression in keratinocytes was only induced by UVB irradiation, others have shown that UVA can also induce expression in these cells [49, 50]. This increase in TNF α released by the cells is due to elevated gene transcription [38, 49]. The IL-1 α formed in the skin, can in turn, induce mast cells to express inflammatory cytokines (e.g. TNF α and IL-1 α), as well as prostaglandins which can enhance the inflammation caused by direct UV exposure on the epidermis [15, 20, 28, 51]. Histamine released from the mast cells can induce vasodilation of the surrounding blood vessels, which assists leucocytes in undergoing diapedesis and entering this region [20, 51]. UVB radiation can induce the synthesis and release of IL-6 and IL-8 from irradiated keratinocytes and fibroblasts [33, 36, 37, 51]. IL-8 assists in the homing of leucocytes, primarily neutrophils, from surrounding blood vessels into the inflamed region, while IL-6 can trigger the activation of monocytes and other infiltrating leucocytes to secrete cytokines and chemokines [51]. Figure 1 shows the complex interaction that occurs between different bioactive molecules in the skin following exposure to UV radiation.

TNF α can induce the expression of adhesion molecules and chemokines in surrounding epithelial cells, resulting in the recruitment of inflammatory leucocytes from surrounding blood vessels via diapedesis [15, 20, 51, 52]. These inflammatory cells in turn can express additional cytokines that form a positive feedback loop that further upregulates TNF α as well as downstream TNF α -induced chemokines, cytokines, and other pro-inflammatory mediators in irradiated skin [8, 38, 53]. The effects elicited by these infiltrating inflammatory cells occur some hours following exposure to UV irradiation, thereby prolonging the inflammatory response. UVB radiation also induces inducible nitric oxide synthase (iNOS) activity in dermal endothelial cells, through a TNF α -dependent pathway [38, 54].

TNF α plays a pro-inflammatory role in the skin due to; (a) the direct effects of UV radiation and (b) the indirect effects of inflammatory cells that chemotax to the skin. UV- and inflammatory cell-derived cytokines further enhance TNF α gene transcription in human skin cells [38], which can again increase its production by epidermal cells. In contrast, clustering and internalization of the TNF receptors may lessen the cell's response to TNF α , which may account for why the upregulation of TNF α mRNA is not sustained over time in culture [20]. For further information on the complex interplay of cytokines, chemokines and other mediators in UV-induced inflammation please refer to the following reviews [15, 20, 41, 42].

Figure 1. The inflammatory response seen in the skin following exposure to UV radiation. Inflammation can be induced as a direct result of UV exposure on epidermal cells, or due to the release of secreted molecules, which in turn induce the release of inflammatory mediators from the dermis, as well as attracting inflammatory cells from circulation into this region of the skin. The infiltrating monocytes and macrophages, which enter the irradiated skin tissue in turn, secrete mediators that prolong the inflammatory response. See text for details and references.

3. Tumor necrosis factor α

TNF α , is a member of the TNF ligand superfamily, and is a type II transmembrane glycoprotein of 234 amino acids possessing an extracellular carboxy-terminus and a cytoplasmic amino group [53, 55, 56]. It can exist in one of two forms; a 26 kDa membrane-bound form (mTNF α) and a 17 kDa soluble form (sTNF α). sTNF α is cleaved from its membrane bound precursor between Ala⁷⁶ - Val⁷⁷ by the action of the metalloprotease TACE [22, 55].

Numerous cells produce TNF α , including macrophages, leucocytes, dendritic cells, keratinocytes, melanocytes and fibroblasts [8, 47, 57, 58]. It plays a role in apoptosis, cellular proliferation, differentiation, inflammation, tumorigenesis, viral replication, immune response to extracellular stimuli, as well as in local and systemic inflammation [21, 53, 55-57, 59]. Most of the cellular actions described for TNF α correspond to its secreted, mature soluble form. There is increasing evidence that mTNF α is also biologically active [58]. Both forms of TNF α can specifically bind to one of two receptors: TNF-R1 (CD120a receptor), a 55 kDa protein; TNF-R2 (CD120b receptor), a 75 kDa protein [57]. The receptors are both transmembrane glycoproteins, and display a high degree of structural homology and are expressed on most cell types [60].

TNF-R1 is expressed on a wide range of cell types and its signalling mediates cytotoxicity, cell proliferation, antiviral activity and many of the proinflammatory actions of TNF α [58, 61]. TNF-R2 is expressed on a limited range of cells, including leucocytes, endothelial cells, Langerhans cells (LC) and epithelial cells but its actions are less clear [58, 61]. Membrane-bound TNF-R1 and TNF-R2 can be cleaved by TACE to release the soluble forms of these receptors and this process is activated by IL-10 [58]. The soluble forms of TNF-R may act as (a) an antagonist to the surface receptors by competing for sTNF α or (b) an agonist by stabilizing the TNF trimer; therefore maintaining saturating concentrations in extracellular fluids [58, 62].

When TNF α is bound to the TNF-R1 receptor it plays a role in UVB-induced apoptosis in keratinocytes [54, 63]. Transgenic mice deficient for either TNF-R1 and/or TNF-R2 have been shown to be less susceptible to UVB-induced skin tumours than were wild type controls [64]. Through the use of TNF-R1 [65, 66] and TNF-R2 [65] gene-targeted mutant mice, it has been shown that TNF-R1 plays a decisive role in the host's defence against microorganisms, while TNF-R2 plays a role in the induction of tissue necrosis. Through the use of agonist and antagonist antibodies, TNF-R1 was shown to be the main mediator of TNF α action in the cell [67].

Dermal injection of TNF α resulted in the accumulation of dendritic cells in draining lymph nodes as well as in impairment of contact hypersensitivity (CHS) in the skin [60, 68]. This suggests that TNF α induces the migration of LC from the skin to the surrounding regional lymph nodes. Streilein and colleagues [69, 70] observed that UVB indirectly induced TNF α , which then caused morphologic and functional changes on LC resulting in the impairment of CHS, suggesting that TNF α plays a role in this process.

Studies using TNF-R1(-) mutant mice have shown that TNF α was not involved in UVB-induced immunosuppression [71]. UVB-induced immunosuppression is implicated in the pathogenesis of skin cancers, and is mediated in part by *cis*-urocanic acid (*cis*-UCA) [72, 73]. *trans*-Urocanic acid, a deamination product of histidine, is a major chromophore present at high concentrations in the stratum corneum [73]. Upon exposure to UV radiation, *trans*-UCA undergoes a photoisomerization to its *cis*-isomer until equilibrium is reached. In humans, this occurs after one minimal erythral dose of UV radiation, which is the lowest dose that can induce a visibly perceptible erythema [72, 73]. *cis*-UCA does not exert its immunosuppressive effects via TNF α , but through other factors such as prostaglandin E₂ [72]. Amerio *et al.* [71] showed that in TNF-R1 and TNF-R2 double knockout mice, TNF α played a minimal role in UVB-induced immunosuppression and therefore cannot be considered as a major mediator of *cis*-UCA-induced immunosuppression. While TNF α does not play a major role in UV-induced immunosuppression [60, 71] it does play a significant role in UV-induced inflammation [20] as well as in other inflammatory diseases such as rheumatoid arthritis, psoriasis, systemic lupus erythematosus and cancer [21, 38, 46, 74].

4. TACE

TNF α is cleaved from its proform by the action of the metalloprotease TACE [75]. This enzyme is a member of the disintegrin and metalloprotease (ADAM) family of proteases, and is

also known as ADAM 17 [22, 75-77]. ADAM proteases belong to the adamalysin/reprolysin subfamily of the metzincin superfamily, and contain a Zn^{2+} -dependent catalytic domain [75, 77].

TACE was first purified, characterized and cloned in 1997 and is a multi-domain type I transmembrane protein of 824 amino acids in length [22, 76, 78]. While its amino acid sequence shows relatively low homology to other ADAM family members, its structure contains all the domain regions, which are characteristic for this family of metalloproteases [22, 76, 79]. Structurally TACE consists of a signal peptide followed by a pro, catalytic, disintegrin, cysteine-rich, transmembrane and cytoplasmic domain [55, 80]. The catalytic domain contains the zinc-binding consensus motif HEXGHXXGXXHD involved in coordinating Zn^{2+} with His residues and creating the active site of the enzyme [79, 81]. The cysteine-rich domain may play a role in enzyme maturation or substrate recognition [75, 76].

TACE is synthesized as an inactive zymogen, which is subsequently proteolytically processed to the catalytically active form. In order for TACE to be activated its prodomain is removed at the furin cleavage site RVKR (Arg-Val-Lys-Arg) localized between the pro- and the catalytic domain, and is due to the action of a furin-type proprotein convertase [24, 77, 82-84]. In mammalian cells, proTACE is located in the endoplasmic reticulum and the proximal Golgi body whereas the mature form is located both intracellularly and on the cell membrane [83, 85]. TACE maturation is closely linked to the transport of proTACE through the medial Golgi, where upon exit, prodomain removal occurs before the enzyme reaches the cell's surface [77].

Apart from $TNF\alpha$, TACE cleaves a wide range of molecules including transforming growth factor α ($TGF\alpha$), amphiregulin, neuregulin, growth hormone receptor, $TNF-R1$, $TNF-R2$, L-selectin, amyloid precursor protein and IL-6R [77, 86-89]. TACE-knockout mice are far less efficient at processing $TNF\alpha$ on the cell membrane compared to wild type controls [75, 86]. This suggests that TACE is the main protease responsible for the processing of $TNF\alpha$ in the cell. Although some matrix metalloproteases (MMP) can cleave $TNF\alpha$, the cleaved products are inactive due to hydrolysis occurring at different sites within the molecule [75, 81, 89].

Some metalloproteases are activated in epidermal cells following UV radiation [90-93]. Piva and co-workers found that there were a number of proteases whose activity was upregulated in UVC- or UVB-irradiated HeLa cells [91-93]. These enzymes included aminopeptidases and a "TGF α ase" [91, 92]. On re-evaluation of their data, the TGF α ase in questions is most likely TACE, because (a) the later enzyme is known to cleave $TGF\alpha$ among other growth factors [81, 88] and (b) the substrate used in these studies was a nonapeptide based on the N-terminal cleavage site of $TGF\alpha$ [90-93]. In cells undergoing UV-induced apoptosis, the level of cell surface protease activity (aminopeptidase and "TGF α ase") was shown to be higher than that seen in viable or necrotic cells [91, 93]. The results of these studies were the first to show that TACE activity was elevated in cells exposed to UV radiation. Recently Skiba *et al.* [29] reported that UVA and UVB irradiation increased TACE mRNA levels in HaCaT cells, with higher induction induced by UVA. The expression patterns for both UVA- and UVB-irradiated cells in general appeared to be constant, although mRNA levels were significantly higher than controls throughout the 48 h post-exposure period [29].

In UV-irradiated HaCaT cells, TACE was responsible for the increased cleavage of EGF family members [28, 94]. Inhibition of TACE by metalloprotease inhibitors reduced the release of these growth factors, resulting in an increase in apoptotic cell death [28, 94]. It appears that TACE mediates a EGF receptor/AKT signalling pathway in these cells that is activated as a result of its cleavage of EGF family members. In HaCaT cells exposed to UVA-radiation TACE mediated EGF receptor activation and cell cycle progression, which suggests that UVA, at non-lethal doses, has the potential to be a skin cancer promoter [28, 94]. TACE has also shown to be overexpressed in some tumours [21, 46, 56], as well in a large number of skin cancer cells lines compared to their non tumorous counterparts [28, 94]. It is also known that members of the EGF family are overexpressed in skin cancers [95], and this could be a mechanism by which skin cancer growth is stimulated by autogenic growth factors. The results of these recent studies suggest that inhibition of TACE following UV radiation may prevent the stimulation of surviving irradiated cells. This has the potential in reducing the incidence of skin cancer that may arise from prolonged sun exposure. It is not clear if the increase in TACE activity seen in UV-irradiated skin cells is due to increased numbers or a higher level of activity. Furin is known to activate TACE [83, 85, 96] as well as matrix metalloproteases (MMP) [97, 98] and may indirectly play a role in this process.

5. Furin

Furin, also known as PACE, is a 94 kDa, type I transmembrane, Ca^{2+} -dependant serine protease. It is a member of the proprotein convertase (PC) family which is related to the bacterial subtilisin enzyme [23, 97-99]. The PC family consists of seven distinct members (furin and PC1-PC7) that vary in regards to their tissue and subcellular distribution as well as enzymatic and biochemical properties [23, 24, 97, 100]. Furin, PACE 4, PC5/6 and PC7/8 are widely expressed in the epidermis whereas PC2 and PC1/3 are limited to neuroendocrine tissues and PC4 is restricted to the testis [23, 24, 98]. The PC enzymes recognize basic motifs, cleaving after paired basic residues (PC2 and PC1/3); or after a canonical Rx (R/K) R (Arg-x-(Arg/Lys)-Arg) motif (furin and PACE4) [24, 97, 98, 100-102]. Both PC7 and furin share cleave similar substrates and the selectivity of which depends on their cellular localization. As their cytosolic domains regulate intracellular trafficking it is likely that the cellular localization of PC7 differs to that for furin [85].

Structurally furin and other PCs consist of a signal peptide followed by pro, catalytic, middle, and cytoplasmic domains, respectively [24]. The signal peptide directs the translocation of the peptide chain to the endoplasmic reticulum and the secretory pathway [82, 97, 103]. The pro-region is cleaved in the endoplasmic reticulum, where it then associates with the catalytic domain and helps to guide the protein through this region to the Golgi apparatus where it becomes catalytically active [97, 103]. The trans-membrane region anchors the enzyme in the membrane of the trans Golgi network (TGN) or on the cell membrane. The cytosolic tail contains the information necessary for furin's sorting to various intracellular compartments [82, 97, 103]. In the epidermis, furin can exist either as: (a) a mature 97 kDa membrane bound enzyme or (b) a smaller 75 kDa form that lacks the transmembrane domain [97]. This suggests that post-translational cleavage at the C-terminus occurs within in the cell [97, 98, 103]. Furin

and other PC family members process inactive precursor proteins to their functional or mature form, and these include growth factor receptors, growth factors, hormones, plasma proteins, and MMPs [23, 24, 97, 98, 103] as seen in Table 2. PC family members play crucial roles in a variety of physiological processes and are involved in the pathology of diseases such as cancer and viral infection [23, 101, 103-106].

Functional group	Substrate	References
Serum proteins	Von Willebrand Factor	[107]
	Coagulation factor IX	[108]
Signalling peptides	Endothelin-1	[103, 109]
Growth factors	TGF β	[103, 110]
	Vascular endothelial growth factor (VEGF)	[111]
	β -Nerve growth factor	[112]
Membrane proteins	MT1-MMP	[86, 113, 114]
	TACE	[77, 99, 115]
Transmembrane receptors	Notch1 Receptor	[98, 116]
	Insulin growth factor 1 receptor	[117]
Extracellular matrix proteins	N-Cadherin	[113, 118]
	Integrin α -chain subunits	[119]
Viral proteins	Ebola virus glycoprotein	[103, 120]
	Papillomavirus minor capsid protein L2	[121]
Bacterial toxins	Anthrax toxin	[122]
	<i>Clostridium septicum</i> alpha-toxin	[103, 123]

Table 2. Some biological molecules cleaved by furin

As a result of the role furin plays in many disease states, considerable effort has been directed at designing specific inhibitors that may have therapeutic applications. The first furin inhibitors that were synthesised were peptidyl chloromethyl ketones [124]. The next major furin inhibitor that was developed, decanoyl-Arg-Val-Lys-Arg-chloromethylketone (dec-RVKR-cmk, or CMK) was less cytotoxic and is cell permeable and has been used in many experimental studies [86, 106, 125]. It was recently shown to reduce the incidence of skin cancer in transgenic mice by inhibiting PACE4 as well as other PCs [126]. However a limitation of CMK's use is that it is not furin specific, and is also known to inhibit other proprotein convertases [86, 102, 127]. Zhu *et al.* [127] has recently developed an antibody-based single domain nanobody which is a furin specific inhibitor. Through the use of this and other furin-specific inhibitors, it will be possible to delineate the role furin plays in the processing of specific substrates within in the cell. This will help in development of specific inhibitors, which will have therapeutic potential in the treatment of a variety of diseases.

Furin and other PCs have been shown to be involved in the maturation of both TACE and MMP within skin cells. ProTACE is processed by both furin and PC7 to its mature form thereby increasing its proteolytic activity [83, 85]. The maturation of TACE occurs as it transits through

the Golgi compartment where the prodomain was removed by a furin-type proprotein convertase [77, 84, 85]. As increased amounts of mature TACE are detected in furin over-expressing cells, it appears that proTACE is a better substrate for furin than it is for PC7 [85]. A similar observation has been seen in cells overexpressing TACE [58, 83, 99] where furin was shown to be responsible for its cleavage [83, 99]. This finding was confirmed using cell permeable furin inhibitors CMK and PDX in Cos7 cells [83] and keratinocytes [98] where reduced levels of mature TACE were formed.

Furin mRNA, protein and enzyme activity has been observed in human epidermal keratinocytes [29, 98, 111, 128, 129]. Skiba *et al.* [29] found that UVA and UVB radiation immediately increased furin mRNA levels in HaCaT cells. UVB irradiation induced higher levels of furin mRNA expression [29]. The time course for furin mRNA levels in cells irradiated with low dose of UVA or high dose of UVB was similar to that for TNF α , whereas maximal mRNA induction of both genes were detected 8 h post-irradiation [29]. Although UV irradiation does appear to have an effect on furin gene expression, no direct relationship was apparent between TACE and furin mRNA induction. A recent study has shown that following exposure to UVA and UVB, furin levels in HaCaT cells fell with respect to time [49, 129]. However, it was unknown whether this was due to the loss of the pro or mature form of the enzyme. Through its effect on stimulating MMPs, as well as activating TACE and the resultant effect this has on TNF α released by the cell, furin activity has an influence on the inflammation seen in the skin following exposure to UV radiation as seen in Figure 2.

Furin/PC processing of substrates has been shown to also contribute to tumour progression, aggressiveness, metastasis, and angiogenesis [23, 24, 104-106]. Tumour invasion and metastasis represent a multistep process that depends on the activity of many proteins [46, 101, 104, 130]. Proteolytic degradation of the ECM components is a central event of this process. Several classes of proteases, including MMPs, serine proteases and cysteine proteases have been implicated in the tumour cell invasive process [104, 130, 131]. Of these, MMPs appear to be primarily responsible for much of the ECM degradation observed during invasive processes [111, 130, 132-134]. They can contribute to tumour growth not only by degradation of the ECM but by the release of sequestered growth factors or the generation of bioactive fragments VEGF, bFGF or TGF β , the suppression of tumour cell apoptosis and the destruction of immune-modulating chemokine gradients [131, 132, 135]. Furin also cleaves a number of MMPs from their proform, and activating them as a result [23, 86, 102, 105].

6. MMPs

MMPs belong to the family of zinc-dependent endopeptidases collectively referred to as metzincins. The metzincins can be subdivided into four families: seralysins, astacins, ADAMs/adamalysins, and MMPs [130, 136]. So far to date, 28 members of the MMP family have been identified [130, 135, 136] which are primarily responsible for most of the ECM degradation observed during the invasive processes. MMPs are produced by skin cells (fibroblasts, keratinocytes, melanocytes) as well as macrophages, endothelial cells and mast cells [10, 25, 81, 137].

Figure 2. The role furin plays in the maturation of TACE and MMPs in skin cells. Furin cleaves and activates TACE, which in turn can process TNFα from its proform. Keratinocytes secrete TNFα following exposure to UVB radiation, and this is enhanced if IL-1α is present. Furin also cleaves MMPs from their respective proforms, and the expression and activity of these proteases are elevated when the cells have been exposed to UVB radiation, and they are enhanced if either IL-1α (MMP-9) or TNFα (MMP-2) is present. The effect of UVB radiation on the expression of the enzymes and pTNFα in the cell is represented by dashed lines, if it is enhanced it is represented by (+), and if it is unknown (?)

MMPs are also implicated in cell migration, proliferation, and tissue remodelling and thereby may also play a role in growth and development, angiogenesis, and atherosclerosis [138, 139].

Structurally MMPs consist of a signal peptide followed by pro, catalytic, hemopexin and cytoplasmic domains, respectively [130]. MMPs cleave peptides and proteins, which have a myriad of functions that are independent of their proteolytic activity [140]. They have distinct but often overlapping substrate specificities, hence leading to the absence of distinct phenotypes in most genetically-engineered mice with knockdown of specific MMPs [140].

MMPs are generally expressed in very low amounts and their transcription is tightly regulated either positively or negatively by cytokines and growth factors such as IL-1, IL-4, IL-6, TGFβ, or TNFα [130, 135, 141, 142]. Some of these regulatory molecules can be proteolytically activated or inactivated by MMPs (via a feedback loop). MMPs are synthesized as latent proenzymes, which are converted into mature, catalytically active forms in the TGN by PCs [111, 139]. Activation of MMPs following secretion from the cell depends on disruption of the prodomain interaction with the catalytic site, which may occur either by conformational changes or proteolytic removal of the prodomain. With the exception of MMP-2, the mechanism for *in vivo* activation of secreted MMPs is not well understood [135].

In normal skin, MMPs are not constitutively expressed but can be induced temporarily in response to exogenous signals such as UVR [10, 25, 143]. Elevated levels of MMP activity in human skin, as a result of prolonged periods of sun exposure, confirm that it plays a major role in photoageing [10, 25, 28]. Onoue *et al.* [144] suggested that MMP-9 secreted from keratinocytes after UVB irradiation might result from apoptotic events. UV radiation is known to elevate the expression of MMP-1, MMP-3 (stromelysin-1) and MMP-9 in human skin *in vivo* [25]. All three MMPs (1, 3 and 9) can degrade most of the proteins found in the extracellular matrix [25]. MMP-1, which is produced by both dermal fibroblasts and epidermal keratinocytes, cleaves type 1 collagen into specific fragments. These fragments can then be further hydrolysed by MMP-2 and MMP-9 [137, 145]. Steinbrenner *et al.* [137] found that UVA irradiation dose-dependently decreased the steady-state mRNA levels of MMP-2 and MMP-9 and lowered the gelatinolytic activity of both enzymes in cell culture supernatants. Of interest is that *in vivo*, following exposure to UV radiation only keratinocytes express large levels of MMP on the cell membrane [146], but when fibroblasts grown in culture are irradiated they express higher levels of MMP on their plasma membrane [147]. The reasons for the discrepancies between the responses of human skin cells under *in vivo* and *in vitro* conditions are not known.

TNF α has been shown to induce proMMP-2 in human dermal fibroblasts [8], while IL-1 α induced proMMP-9 levels in fibroblasts and keratinocytes [148]. In mesenchymal cells TNF α was shown to stimulate MMP-2 activity by activating a proteolytic cascade involving furin and MT1-MMP [139]. It is not known if TNF α activates MMP activity in epidermal cells via a similar mechanism. MMP-9 [144], has been shown to play an important role in the pathophysiology of many skin conditions such as wound healing [145], and angiogenesis [87]. Activation of pro-MMP2 takes place at the cell surface and involves interactions with active MT1-MMP, which is itself activated through rapid trafficking to the cell surface and proteolytic processing [139]. Maquoi *et al.* [114] demonstrated that furin-inhibitor reduces the level of mature MT1-MMP, which is paralleled by a decrease in pro-MMP-2 activation as well as in cell invasiveness, confirming the furin plays a role in this process. Direct cleavage of proMMP-2 by furin in the TGN has shown to inactivate this matrix metalloprotease [149]. Therefore changes to the level of MMP-2 activity on the surface of the cell can be directly or indirectly regulated by furin either through cleavage in the TGN to reduce activity or indirectly via MT1-MMP, which increases activity. The mechanism by which this is regulated is not clear.

7. Role of furin in skin cancer

Low doses of UV radiation while they are not inflammatory, can cause mutations in skin tissue [9, 19, 30] or in cultured fibroblasts [150]. Constant exposure to UV radiation will not only result in mutations in p53 [150, 151] but in other genes including p16, PTCH, B-Raf, which will result in these cells becoming carcinogenic [6, 152, 153]. Apart from causing these mutations, UV radiation enhances the release of inflammatory molecules (e.g. TNF α , IL-1 α , ROS, chemokines) from the skin (see section 2 for details). The result of which is the creation of an environment that allows for these mutated skin cells to become cancerous over time [15, 19, 21, 46, 56]. As a result of increased levels of MMP activity on the cell membrane, either as a

result of UV exposure or other factors, these skin cancer cells may become metastatic as a result of epithelial to mesenchymal transition (EMT) [154].

In the progression of a mutated cell to that of a tumorigenic or metastatic cell, PCs have been shown to cleave a range of precursors of growth factors, their receptors, adhesion molecules, proteases and MMPs. Some of these molecules include cadherins, TGF β , platelet derived growth factor as well as insulin-like growth factor 1 and its receptor [23, 101, 106, 117, 155]. Tumour progression and metastasis may enhanced by a number of factors such as (a) hypoxia-induced upregulation of furin activity within the solid tumour mass [105], (b) changes in cell adhesion through PC cleavage and activation of integrins and related adhesion molecules [156], (c) furin processing of vascular endothelial growth factor resulting in increased angiogenic activity [157], and (d) furin enhanced expression of MMPs [84, 158].

Furin/PC expression and processing can increase the incidence and severity of the cancer phenotype [104, 111]. Aberrant furin expression has been observed in a number of tumours including those from the breast [159], ovary [160], liver [125], brain [161], skin [111, 132] and from other tissues [23, 128, 162, 163]. Bassi *et al.* [101] observed that PACE4 transgenic mice were more susceptible to epidermal carcinogenesis and tumour progression compared to controls. These transgenic keratinocytes had higher rates of processing of MT1-MMP and MT2-MMP resulting in increased collagen degradation.

MMP-2 (gelatinase A) and MMP-9 (gelatinase B) have been frequently associated with the invasive and metastatic potential of tumour cells [10, 104, 111, 130, 132, 133, 137]. The expression of MMP-2, is regulated independently of MMP-9 [144]. The close correlation observed between MMP-2 activation and metastatic progression in various tumours suggests that it may act as a “master switch” triggering tumour spread [114]. The expression of MMP is low in keratinocytes but elevated in BCC and SCC [49, 164, 165]. In SCC, MMP proteins (1, 2, 3 and MT1-MMP) are expressed both in tumorous and stromal cells [165], while MMP proteins (1, 2, 3 and 9) are observed in BCCs and melanomas [165, 166]. This expression correlates with the progression and the metastatic potentials of these tumours [106, 135, 165, 167]. UVR can participate to the development of skin cancer by the activation of MMPs. Two molecular mechanisms contribute to the UV-induced MMPs expression. First, the activation of cell-surface receptors with subsequent activation of mitogen activated protein kinase (MAPK) cascade that in turn contributes to the transcriptional up-regulation of MMPs [168]. Second, through the expression of pro-inflammatory cytokines, which induce the expression of MMPs [46, 133, 142, 143]. The role of UV in the induction of MMPs is supported by two experimental findings. First, UV-irradiation of SCC cell lines results in an increased secretion of MMPs [49, 132]. Second, UV-induced phosphorylation of extracellular signal-regulated kinase (ERK) and stress kinases precedes the rapid stimulation of MMPs in SCC cells [10]. If there are cells in the skin which become cancerous as a result of DNA damage, some may go onto become metastatic due to increased MMP activity [111]. This increase in levels of activated MMPs on the surface of the cell could be due to either increased expression of proMMP protein and/or increased furin activity. The role furin plays in the development of skin cancer suggests that it could be significant, and as such the development of specific inhibitors may offer a new therapy to treat such tumours.

Recently, Fu *et al.* [106] showed that in transgenic mice overexpressing furin, exposed to chemical carcinogens formed more and larger tumours than did control mice. This suggests that furin enhances skin cancer growth. Of interest was that these tumours were induced by chemical carcinogens and not by UV radiation. While enhanced furin levels were shown to enhance skin tumour development, in studies using cultured melanoma and glioma cells, furin inhibition was shown to reduce the processing of the pro-N-cadherin adhesion molecule, enhancing their migratory and aggressive nature [118]. The result of this study suggests that high furin levels may not enhance the metastasis of some tumours. Huang *et al.* [125] when investigating the expression of furin in hepatocellular carcinoma and surrounding tissue in humans, observed that if this ratio was >3.5 this resulted in higher patient survival rates compared to those who had ratios <3.5 . This suggests that furin may play a dual role in cancer development, though the exact nature of which is not clear at this stage. Further study on this is warranted, and may result in a better understanding of its role in cancer development, and from which it may be possible to develop specific furin inhibitors which could be used clinically to treat these tumours.

8. Conclusion

While it has been shown that UVA and UVB radiation cause different effects on the immune response this could be related to the activity of cell surface metalloproteases found on skin cells. Although the effect of TNF α in UV-induced inflammation has been well documented, little is known if the changes in TACE activity are due to increased protein levels or changes in enzyme activity. The inflammatory environment, seen in the skin, following exposure to UV radiation is known to stimulate the development of mutated cells which possess DNA damage caused directly by UV radiation or indirectly through the generation of ROS. Apart from processing TNF α , TACE also cleaves EGF family members, which would stimulate the growth of these mutated cells, which over time may become cancerous. As TNF α is a powerful inflammatory cytokine, considerable research has been under taken to develop specific TACE inhibitors [55, 56, 80]. Such inhibitors may play an important role in preventing the development of UV-induced skin cancers. An increased knowledge of the roles played by metalloproteases in tumour progression, combined with the use of more selective inhibitors, could lead to effective use of these compounds in cancer therapies [55].

Similar to that of TACE, MMPs are also activated by UV radiation and also play a crucial role in skin tumour cell development and metastasis. Furin and other PCs have been shown to play an important role in activating both TACE and many MMPs in skin cells. Whereas the overexpression and activity of furin exacerbates the cancer phenotype, inhibition of its activity decreases or nullifies its effects, and thus, the development and use of specific inhibitors may also be a viable route to cancer therapy [23, 86, 102, 127].

It is known that UV induces furin mRNA in skin cells, though protein levels do not appear to change with respect to time, which suggests a rapid turnover of the enzyme. Further study is needed on how UV radiation activates furin, TACE and MMPs in skin cells. These proteases

play an important role in the changes observed in those epidermal cells exposed to UV radiation. The development of specific furin inhibitors has the potential to reduce the carcinogenic effects of sunlight by preventing the activation of TACE and MMPs and their subsequent downstream effects. Such compounds may have the potential to offer new therapies both in the prevention and treatment of skin cancer.

Acknowledgements

We would like to thank the School of Medical Sciences, RMIT University, Bundoora, Australia for supporting this study, and for the awarding of a School of Medical Science's Postgraduate Research Scholarship to RR.

Author details

Rethika Ravi¹ and Terrence J Piva^{2*}

*Address all correspondence to: terry.piva@rmit.edu.au

1 Endeavour College of Natural Health, Melbourne, Australia

2 School of Medical Sciences, RMIT University, Melbourne, Australia

References

- [1] Guy GP, Ekwueme DU. Years of potential life lost and indirect costs of melanoma and non-melanoma skin cancer: a systematic review of the literature. *Pharmacoeconomics*. 2011;29(10) 863-874.
- [2] Qureshi AA, Laden F, Colditz GA, Hunter DJ. Geographic variation and risk of skin cancer in US women. Differences between melanoma, squamous cell carcinoma, and basal cell carcinoma. *Archives of Internal Medicine*. 2008;168(5) 501-507.
- [3] Narayanan DL, Saladi RN, Fox JL. Ultraviolet radiation and skin cancer. *International Journal of Dermatology*. 2010;49(9) 978-986.
- [4] Thomas-Ahner JM, Wulff BC, Tober KL, Kusewitt DF, Riggenbach JA, Oberyszyn TM. Gender differences in UVB-induced skin carcinogenesis, inflammation, and DNA damage. *Cancer Research*. 2007;67(7) 3468-3474.

- [5] Erb P, Ji J, Kump E, Mielgo A, Wernli M. Apoptosis and pathogenesis of melanoma and nonmelanoma skin cancer. *Advances in Experimental Medicine and Biology*. 2008;624 283-295.
- [6] Rass K, Reichrath J. UV damage and DNA repair in malignant melanoma and non-melanoma skin cancer. *Advances in Experimental Medicine and Biology*. 2008;624 162-178.
- [7] Maverakis E, Miyamura Y, Bowen MP, Correa G, Ono Y, Goodarzi H. Light, including ultraviolet. *Journal of Autoimmunity*. 2010;34(3) J247-257.
- [8] Clydesdale GJ, Dandie GW, Muller HK. Ultraviolet light induced injury: immunological and inflammatory effects. *Immunology and Cell Biology*. 2001;79(6) 547-568.
- [9] Agar NS, Halliday GM, Barnetson RS, Ananthaswamy HN, Wheeler M, Jones AM. The basal layer in human squamous tumors harbors more UVA than UVB fingerprint mutations: a role for UVA in human skin carcinogenesis. *Proceedings of the National Academy of Sciences of the United States of America*. 2004;101(14) 4954-4959.
- [10] Dong KK, Damaghi N, Picart SD, Markova NG, Obayashi K, Okano Y, Masaki H, Grether-Beck S, Krutmann J, Smiles KA, Yarosh DB. UV-induced DNA damage initiates release of MMP-1 in human skin. *Experimental Dermatology*. 2008;17(12) 1037-1044.
- [11] Pfeifer GP, Besaratinia A. UV wavelength-dependent DNA damage and human non-melanoma and melanoma skin cancer. *Photochemical and Photobiological Sciences*. 2012;11(1) 90-97.
- [12] Sander CS, Chang H, Hamm F, Elsner P, Thiele JJ. Role of oxidative stress and the antioxidant network in cutaneous carcinogenesis. *International Journal of Dermatology*. 2004;43(5) 326-335.
- [13] Muthusamy V, Piva TJ. The UV response of the skin: a review of the MAPK, NF κ B and TNF α signal transduction pathways. *Archives for Dermatological Research (Archiv fur Dermatologische Forschung)*. 2010;302(1) 5-17.
- [14] Rangwala S, Tsai KY. Roles of the immune system in skin cancer. *British Journal of Dermatology*. 2011;165(5) 953-965.
- [15] Ullrich SE, Byrne SN. The immunologic revolution: photoimmunology. *Journal of Investigative Dermatology*. 2012;132(3 Pt 2) 896-905.
- [16] Shorrocks J, Paul ND, McMillan TJ. The dose rate of UVA treatment influences the cellular response of HaCaT keratinocytes. *Journal of Investigative Dermatology*. 2008;128(3) 685-693.

- [17] Valencia A, Kochevar IE. Nox1-based NADPH oxidase is the major source of UVA-induced reactive oxygen species in human keratinocytes. *Journal of Investigative Dermatology*. 2008;128(1) 214-222.
- [18] Mahns A, Wolber R, Stab F, Klotz LO, Sies H. Contribution of UVB and UVA to UV-dependent stimulation of cyclooxygenase-2 expression in artificial epidermis. *Photochemical and Photobiological Sciences*. 2004;3(3) 257-262.
- [19] Halliday GM, Lyons JG. Inflammatory doses of UV may not be necessary for skin carcinogenesis. *Photochemistry and Photobiology*. 2008;84(2) 272-283.
- [20] Damian DL, Patterson CR, Stapelberg M, Park J, Barnetson RS, Halliday GM. UV radiation-induced immunosuppression is greater in men and prevented by topical nicotinamide. *Journal of Investigative Dermatology*. 2008;128(2) 447-454.
- [21] Balkwill F. Tumour necrosis factor and cancer. *Nature Reviews Cancer*. 2009;9(5) 361-371.
- [22] Black RA, Rauch CT, Kozlosky CJ, Peschon JJ, Slack JL, Wolfson MF, Castner BJ, Stocking KL, Reddy P, Srinivasan S, Nelson N, Boiani N, Schooley KA, Gerhart M, Davis R, Fitzner JN, Johnson RS, Paxton RJ, March CJ, Cerretti DP. A metalloproteinase disintegrin that releases tumour-necrosis factor- α from cells. *Nature*. 1997;385(6618) 729-733.
- [23] Artenstein AW, Opal SM. Proprotein convertases in health and disease. *New England Journal of Medicine*. 2011;365(26) 2507-2518.
- [24] Seidah NG. The proprotein convertases, 20 years later. *Methods in Molecular Biology*. 2011;768 23-57.
- [25] Quan T, Qin Z, Xia W, Shao Y, Voorhees JJ, Fisher GJ. Matrix-degrading metalloproteinases in photoaging. *Journal of Investigative Dermatology Symposium Proceedings*. 2009;14(1) 20-24.
- [26] Sluyter R, Halliday GM. Infiltration by inflammatory cells required for solar-simulated ultraviolet radiation enhancement of skin tumor growth. *Cancer Immunology, Immunotherapy*. 2001;50(3) 151-156.
- [27] Weller R. Nitric oxide: a key mediator in cutaneous physiology. *Clinical and Experimental Dermatology*. 2003;28(5) 511-514.
- [28] Bennett MF, Robinson MK, Baron ED, Cooper KD. Skin immune systems and inflammation: protector of the skin or promoter of aging? *Journal of Investigative Dermatology Symposium Proceedings*. 2008;13(1) 15-19.
- [29] Skiba B, Neill B, Piva TJ. Gene expression profiles of TNF- α , TACE, furin, IL-1 β and matrilysin in UVA- and UVB-irradiated HaCat cells. *Photodermatology, Photoimmunology and Photomedicine*. 2005;21(4) 173-182.

- [30] Halliday GM. Inflammation, gene mutation and photoimmunosuppression in response to UVR-induced oxidative damage contributes to photocarcinogenesis. *Mutation Research*. 2005;571(1-2) 107-120.
- [31] Barr RM, Walker SL, Tsang W, Harrison GI, Ettehadi P, Greaves MW, Young AR. Suppressed alloantigen presentation, increased TNF- α , IL-1, IL-1Ra, IL-10, and modulation of TNF-R in UV-irradiated human skin. *Journal of Investigative Dermatology*. 1999;112(5) 692-698.
- [32] Byrne SN, Limon-Flores AY, Ullrich SE. Mast cell migration from the skin to the draining lymph nodes upon ultraviolet irradiation represents a key step in the induction of immune suppression. *Journal of Immunology*. 2008;180(7) 4648-4655.
- [33] Norval M. Effects of solar radiation on the human immune system. *Journal of Photochemistry and Photobiology B, Biology*. 2001;63(1-3) 28-40.
- [34] Morelli JG, Norris DA. Influence of inflammatory mediators and cytokines on human melanocyte function. *Journal of Investigative Dermatology*. 1993;100(2 Suppl) 191S-195S.
- [35] Berhane T, Halliday GM, Cooke B, Barnetson RS. Inflammation is associated with progression of actinic keratoses to squamous cell carcinomas in humans. *British Journal of Dermatology*. 2002;146(5) 810-815.
- [36] Meeran SM, Punathil T, Katiyar SK. IL-12 deficiency exacerbates inflammatory responses in UV-irradiated skin and skin tumors. *Journal of Investigative Dermatology*. 2008;128(11) 2716-2727.
- [37] Yoshizumi M, Nakamura T, Kato M, Ishioka T, Kozawa K, Wakamatsu K, Kimura H. Release of cytokines/chemokines and cell death in UVB-irradiated human keratinocytes, HaCaT. *Cell Biology International*. 2008;32(11) 1405-1411.
- [38] Bashir MM, Sharma MR, Werth VP. UVB and proinflammatory cytokines synergistically activate TNF- α production in keratinocytes through enhanced gene transcription. *Journal of Investigative Dermatology*. 2009;129(4) 994-1001.
- [39] Bashir MM, Sharma MR, Werth VP. TNF- α production in the skin. *Archives for Dermatological Research (Archiv fur Dermatologische Forschung)*. 2009;301(1) 87-91.
- [40] Lyons JG, Birkedal-Hansen B, Pierson MC, Whitelock JM, Birkedal-Hansen H. Interleukin-1 β and transforming growth factor- α /epidermal growth factor induce expression of M(r) 95,000 type IV collagenase/gelatinase and interstitial fibroblast-type collagenase by rat mucosal keratinocytes. *Journal of Biological Chemistry*. 1993;268(25) 19143-19151.
- [41] Fisher GJ, Datta SC, Talwar HS, Wang ZQ, Varani J, Kang S, Voorhees JJ. Molecular basis of sun-induced premature skin ageing and retinoid antagonism. *Nature*. 1996;379(6563) 335-339.

- [42] Rundhaug JE, Fischer SM. Cyclo-oxygenase-2 plays a critical role in UV-induced skin carcinogenesis. *Photochemistry and Photobiology*. 2008;84(2) 322-329.
- [43] An KP, Athar M, Tang X, Katiyar SK, Russo J, Beech J, Aszterbaum M, Kopelovich L, Epstein EH, Jr., Mukhtar H, Bickers DR. Cyclooxygenase-2 expression in murine and human nonmelanoma skin cancers: implications for therapeutic approaches. *Photochemistry and Photobiology*. 2002;76(1) 73-80.
- [44] Patel MJ, Stockfleth E. Does progression from actinic keratosis and Bowen's disease end with treatment: diclofenac 3% gel, an old drug in a new environment? *British Journal of Dermatology*. 2007;156 Suppl 3 53-56.
- [45] Butler GJ, Neale R, Green AC, Pandeya N, Whiteman DC. Nonsteroidal anti-inflammatory drugs and the risk of actinic keratoses and squamous cell cancers of the skin. *Journal of the American Academy of Dermatology*. 2005;53(6) 966-972.
- [46] Sethi G, Shanmugam MK, Ramachandran L, Kumar AP, Tergaonkar V. Multifaceted link between cancer and inflammation. *Bioscience Reports*. 2012;32(1) 1-15.
- [47] Yarosh D, Both D, Kibitel J, Anderson C, Elmetts C, Brash D, Brown D. Regulation of TNF α production and release in human and mouse keratinocytes and mouse skin after UV-B irradiation. *Photodermatology, Photoimmunology and Photomedicine*. 2000;16(6) 263-270.
- [48] Muthusamy V, Hodges LD, Macrides TA, Boyle GM, Piva TJ. Effect of novel marine nutraceuticals on IL-1 α -mediated TNF- α release from UVB-irradiated human melanocyte-derived cells. *Oxidative Medicine and Cellular Longevity*. 2011;2011 728645.
- [49] Ravi R. The effect of UV radiation on furin activity in human keratinocyte cell lines [PhD Thesis]. RMIT University; Melbourne: 2010.
- [50] Muthusamy V. The role cell signalling pathways play in TNF α release from UV-irradiated human skin cells [PhD Thesis]. RMIT University; Melbourne: 2010.
- [51] Fuller BB, Smith DR. Topical anti-inflammatories. In: Draelos ZD, Thaman LA, editors. *Cosmetic formulation of skin care products*. New York: Informa Healthcare; 2002. p. 351-376.
- [52] Ogilvie AL, Luftl M, Antoni C, Schuler G, Kalden JR, Lorenz HM. Leukocyte infiltration and mRNA expression of IL-20, IL-8 and TNF-R P60 in psoriatic skin is driven by TNF-alpha. *Int J Immunopathol Pharmacol*. 2006;19(2) 271-278.
- [53] Serwin AB, Sokolowska M, Chodyncka B. Tumour necrosis factor α (TNF- α)-converting enzyme (TACE) and soluble TNF- α receptor type 1 in psoriasis patients treated with narrowband ultraviolet B. *Photodermatology, Photoimmunology and Photomedicine*. 2007;23(4) 130-134.
- [54] Schwarz A, Bhardwaj R, Aragane Y, Mahnke K, Riemann H, Metze D, Luger TA, Schwarz T. Ultraviolet-B-induced apoptosis of keratinocytes: evidence for partial in-

- volvement of tumor necrosis factor- α in the formation of sunburn cells. *Journal of Investigative Dermatology*. 1995;104(6) 922-927.
- [55] Saftig P, Reiss K. The "A Disintegrin And Metalloproteases" ADAM10 and ADAM17: novel drug targets with therapeutic potential? *European Journal of Cell Biology*. 2011;90(6-7) 527-535.
- [56] Scheller J, Chalaris A, Garbers C, Rose-John S. ADAM17: a molecular switch to control inflammation and tissue regeneration. *Trends in Immunology*. 2011;32(8) 380-387.
- [57] DasGupta S, Murumkar PR, Giridhar R, Yadav MR. Current perspective of TACE inhibitors: a review. *Bioorganic and Medicinal Chemistry*. 2009;17(2) 444-459.
- [58] Goetz FW, Planas JV, MacKenzie S. Tumor necrosis factors. *Developmental and Comparative Immunology*. 2004;28(5) 487-497.
- [59] Bouwmeester T, Bauch A, Ruffner H, Angrand PO, Bergamini G, Croughton K, Cruciat C, Eberhard D, Gagneur J, Ghidelli S, Hopf C, Huhse B, Mangano R, Michon AM, Schirle M, Schlegl J, Schwab M, Stein MA, Bauer A, Casari G, Drewes G, Gavin AC, Jackson DB, Joberty G, Neubauer G, Rick J, Kuster B, Superti-Furga G. A physical and functional map of the human TNF- α /NF- κ B signal transduction pathway. *Nature Cell Biology*. 2004;6(2) 97-105.
- [60] Lewis M, Tartaglia LA, Lee A, Bennett GL, Rice GC, Wong GH, Chen EY, Goeddel DV. Cloning and expression of cDNAs for two distinct murine tumor necrosis factor receptors demonstrate one receptor is species specific. *Proceedings of the National Academy of Sciences of the United States of America*. 1991;88(7) 2830-2834.
- [61] Choo-Kang BS, Hutchison S, Nickdel MB, Bundick RV, Leishman AJ, Brewer JM, McInnes IB, Garside P. TNF-blocking therapies: an alternative mode of action? *Trends in Immunology*. 2005;26(10) 518-522.
- [62] Cumberbatch M, Kimber I. Tumour necrosis factor- α is required for accumulation of dendritic cells in draining lymph nodes and for optimal contact sensitization. *Immunology*. 1995;84(1) 31-35.
- [63] Zhuang L, Wang B, Shinder GA, Shivji GM, Mak TW, Sauder DN. TNF receptor p55 plays a pivotal role in murine keratinocyte apoptosis induced by ultraviolet B irradiation. *Journal of Immunology*. 1999;162(3) 1440-1447.
- [64] Starcher B. Role for tumour necrosis factor- α receptors in ultraviolet-induced skin tumours. *British Journal of Dermatology*. 2000;142(6) 1140-1147.
- [65] Erickson SL, de Sauvage FJ, Kikly K, Carver-Moore K, Pitts-Meek S, Gillett N, Sheehan KC, Schreiber RD, Goeddel DV, Moore MW. Decreased sensitivity to tumour-necrosis factor but normal T-cell development in TNF receptor-2-deficient mice. *Nature*. 1994;372(6506) 560-563.

- [66] Rothe J, Lesslauer W, Lotscher H, Lang Y, Koebel P, Kontgen F, Althage A, Zinkernagel R, Steinmetz M, Bluethmann H. Mice lacking the tumour necrosis factor receptor 1 are resistant to TNF-mediated toxicity but highly susceptible to infection by *Listeria monocytogenes*. *Nature*. 1993;364(6440) 798-802.
- [67] Tartaglia LA, Pennica D, Goeddel DV. Ligand passing: the 75-kDa tumor necrosis factor (TNF) receptor recruits TNF for signaling by the 55-kDa TNF receptor. *Journal of Biological Chemistry*. 1993;268(25) 18542-18548.
- [68] Cumberbatch M, Kimber I. Dermal tumour necrosis factor- α induces dendritic cell migration to draining lymph nodes, and possibly provides one stimulus for Langerhans' cell migration. *Immunology*. 1992;75(2) 257-263.
- [69] Kurimoto I, Streilein JW. cis-urocanic acid suppression of contact hypersensitivity induction is mediated via tumor necrosis factor- α . *Journal of Immunology*. 1992;148(10) 3072-3078.
- [70] Streilein JW. Sunlight and skin-associated lymphoid tissues (SALT): if UVB is the trigger and TNF α is its mediator, what is the message? *Journal of Investigative Dermatology*. 1993;100(1) 47S-52S.
- [71] Amerio P, Toto P, Feliciani C, Suzuki H, Shivji G, Wang B, Sauder DN. Rethinking the role of tumour necrosis factor- α in ultraviolet (UV) B-induced immunosuppression: altered immune response in UV-irradiated TNFR1R2 gene-targeted mutant mice. *British Journal of Dermatology*. 2001;144(5) 952-957.
- [72] Kaneko K, Smetana-Just U, Matsui M, Young AR, John S, Norval M, Walker SL. cis-Urocanic acid initiates gene transcription in primary human keratinocytes. *Journal of Immunology*. 2008;181(1) 217-224.
- [73] Moodycliffe AM, Bucana CD, Kripke ML, Norval M, Ullrich SE. Differential effects of a monoclonal antibody to cis-urocanic acid on the suppression of delayed and contact hypersensitivity following ultraviolet irradiation. *Journal of Immunology*. 1996;157(7) 2891-2899.
- [74] Dinarello CA. The paradox of pro-inflammatory cytokines in cancer. *Cancer and Metastasis Reviews*. 2006;25(3) 307-313.
- [75] Black RA. Tumor necrosis factor- α converting enzyme. *International Journal of Biochemistry and Cell Biology*. 2002;34(1) 1-5.
- [76] Moss ML, Jin SL, Milla ME, Bickett DM, Burkhart W, Carter HL, Chen WJ, Clay WC, Didsbury JR, Hassler D, Hoffman CR, Kost TA, Lambert MH, Leesnitzer MA, McCauley P, McGeehan G, Mitchell J, Moyer M, Pahel G, Rocque W, Overton LK, Schoenen F, Seaton T, Su JL, Becherer JD, et al. Cloning of a disintegrin metalloproteinase that processes precursor tumour-necrosis factor- α . *Nature*. 1997;385(6618) 733-736.

- [77] Schlondorff J, Becherer JD, Blobel CP. Intracellular maturation and localization of the tumour necrosis factor α convertase (TACE). *Biochemical Journal*. 2000;347 Pt 1 131-138.
- [78] Schlondorff J, Blobel CP. Metalloprotease-disintegrins: modular proteins capable of promoting cell-cell interactions and triggering signals by protein-ectodomain shedding. *Journal of Cell Science*. 1999;112 (Pt 21) 3603-3617.
- [79] Mezyk R, Bzowska M, Bereta J. Structure and functions of tumor necrosis factor- α converting enzyme. *Acta Biochimica Polonica*. 2003;50(3) 625-645.
- [80] Rabinowitz MH, Andrews RC, Becherer JD, Bickett DM, Bubacz DG, Conway JG, Cowan DJ, Gaul M, Glennon K, Lambert MH, Leesnitzer MA, McDougald DL, Moss ML, Musso DL, Rizzolio MC. Design of selective and soluble inhibitors of tumor necrosis factor- α converting enzyme (TACE). *Journal of Medicinal Chemistry*. 2001;44(24) 4252-4267.
- [81] Mohan R, Chintala SK, Jung JC, Villar WV, McCabe F, Russo LA, Lee Y, McCarthy BE, Wollenberg KR, Jester JV, Wang M, Welgus HG, Shipley JM, Senior RM, Fini ME. Matrix metalloproteinase gelatinase B (MMP-9) coordinates and effects epithelial regeneration. *Journal of Biological Chemistry*. 2002;277(3) 2065-2072.
- [82] Anderson ED, VanSlyke JK, Thulin CD, Jean F, Thomas G. Activation of the furin endoprotease is a multiple-step process: requirements for acidification and internal propeptide cleavage. *EMBO Journal*. 1997;16(7) 1508-1518.
- [83] Peiretti F, Canault M, Deprez-Beauclair P, Berthet V, Bonardo B, Juhan-Vague I, Nalbone G. Intracellular maturation and transport of tumor necrosis factor alpha converting enzyme. *Experimental Cell Research*. 2003;285(2) 278-285.
- [84] Komiyama T, Coppola JM, Larsen MJ, van Dort ME, Ross BD, Day R, Rehemtulla A, Fuller RS. Inhibition of furin/proprotein convertase-catalyzed surface and intracellular processing by small molecules. *Journal of Biological Chemistry*. 2009;284(23) 15729-15738.
- [85] Endres K, Anders A, Kojro E, Gilbert S, Fahrenholz F, Postina R. Tumor necrosis factor- α converting enzyme is processed by proprotein-convertases to its mature form which is degraded upon phorbol ester stimulation. *European Journal of Biochemistry*. 2003;270(11) 2386-2393.
- [86] de Cicco RL, Bassi DE, Benavides F, Conti CJ, Klein-Szanto AJ. Inhibition of proprotein convertases: approaches to block squamous carcinoma development and progression. *Molecular Carcinogenesis*. 2007;46(8) 654-659.
- [87] Mohan MJ, Seaton T, Mitchell J, Howe A, Blackburn K, Burkhart W, Moyer M, Patel I, Waitt GM, Becherer JD, Moss ML, Milla ME. The tumor necrosis factor- α converting enzyme (TACE): a unique metalloproteinase with highly defined substrate selectivity. *Biochemistry*. 2002;41(30) 9462-9469.

- [88] Peschon JJ, Slack JL, Reddy P, Stocking KL, Sunnarborg SW, Lee DC, Russell WE, Castner BJ, Johnson RS, Fitzner JN, Boyce RW, Nelson N, Kozlosky CJ, Wolfson MF, Rauch CT, Cerretti DP, Paxton RJ, March CJ, Black RA. An essential role for ectodomain shedding in mammalian development. *Science*. 1998;282(5392) 1281-1284.
- [89] Sunnarborg SW, Hinkle CL, Stevenson M, Russell WE, Raska CS, Peschon JJ, Castner BJ, Gerhart MJ, Paxton RJ, Black RA, Lee DC. Tumor necrosis factor- α converting enzyme (TACE) regulates epidermal growth factor receptor ligand availability. *Journal of Biological Chemistry*. 2002;277(15) 12838-12845.
- [90] Brown SB, Krause D, Ellem KA. Low fluences of ultraviolet irradiation stimulate HeLa cell surface aminopeptidase and candidate "TGF α ase" activity. *Journal of Cellular Biochemistry*. 1993;51(1) 102-115.
- [91] Piva TJ, Davern CM, Francis KG, Chojnowski GM, Hall PM, Ellem KA. Increased ecto-metallopeptidase activity in cells undergoing apoptosis. *Journal of Cellular Biochemistry*. 2000;76(4) 625-638.
- [92] Piva TJ, Krause DR, Ellem KO. UVC activation of the HeLa cell membrane "TGF α ase," a metalloenzyme. *Journal of Cellular Biochemistry*. 1997;64(3) 353-368.
- [93] Piva TJ, Davern CM, Hall PM, Winterford CM, Ellem KA. Increased Activity of Cell Surface Peptidases in HeLa Cells Undergoing UV-Induced Apoptosis Is Not Mediated by Caspase 3. *International Journal of Molecular Sciences*. 2012;13(3) 2650-2675.
- [94] Singh B, Schneider M, Knyazev P, Ullrich A. UV-induced EGFR signal transactivation is dependent on proligand shedding by activated metalloproteases in skin cancer cell lines. *International Journal of Cancer*. 2009;124(3) 531-539.
- [95] Krahm G, Leiter U, Kaskel P, Udart M, Utikal J, Bezold G, Peter RU. Coexpression patterns of EGFR, HER2, HER3 and HER4 in non-melanoma skin cancer. *European Journal of Cancer*. 2001;37(2) 251-259.
- [96] Adrain C, Zettl M, Christova Y, Taylor N, Freeman M. Tumor necrosis factor signaling requires iRhom2 to promote trafficking and activation of TACE. *Science*. 2012;335(6065) 225-228.
- [97] Denault J, Bissonnette L, Longpre J, Charest G, Lavigne P, Leduc R. Ectodomain shedding of furin: kinetics and role of the cysteine-rich region. *FEBS Letters*. 2002;527(1-3) 309-314.
- [98] Pearton DJ, Nirunsuksiri W, Rehemtulla A, Lewis SP, Presland RB, Dale BA. Proprotein convertase expression and localization in epidermis: evidence for multiple roles and substrates. *Experimental Dermatology*. 2001;10(3) 193-203.
- [99] Srour N, Lebel A, McMahon S, Fournier I, Fugere M, Day R, Dubois CM. TACE/ADAM-17 maturation and activation of sheddase activity require proprotein convertase activity. *FEBS Letters*. 2003;554(3) 275-283.

- [100] Tian S, Huang Q, Fang Y, Wu J. FurinDB: A Database of 20-Residue Furin Cleavage Site Motifs, Substrates and Their Associated Drugs. *International Journal of Molecular Sciences*. 2011;12(2) 1060-1065.
- [101] Bassi DE, Fu J, Lopez de Cicco R, Klein-Szanto AJ. Proprotein convertases: "master switches" in the regulation of tumor growth and progression. *Molecular Carcinogenesis*. 2005;44(3) 151-161.
- [102] Becker GL, Lu Y, Hards K, Strehlow B, Levesque C, Lindberg I, Sandvig K, Bakowsky U, Day R, Garten W, Steinmetzer T. Highly potent inhibitors of proprotein convertase furin as potential drugs for treatment of infectious diseases. *Journal of Biological Chemistry*. 2012;287(26) 21992-22003.
- [103] Thomas G. Furin at the cutting edge: from protein traffic to embryogenesis and disease. *Nature Reviews Molecular Cell Biology*. 2002;3(10) 753-766.
- [104] Coppola JM, Bhojani MS, Ross BD, Rehemtulla A. A small-molecule furin inhibitor inhibits cancer cell motility and invasiveness. *Neoplasia*. 2008;10(4) 363-370.
- [105] Arsenault D, Lucien F, Dubois CM. Hypoxia enhances cancer cell invasion through relocalization of the proprotein convertase furin from the trans-Golgi network to the cell surface. *Journal of Cellular Physiology*. 2012;227(2) 789-800.
- [106] Fu J, Bassi DE, Zhang J, Li T, Nicolas E, Klein-Szanto AJ. Transgenic overexpression of the proprotein convertase furin enhances skin tumor growth. *Neoplasia*. 2012;14(4) 271-282.
- [107] Wise RJ, Barr PJ, Wong PA, Kiefer MC, Brake AJ, Kaufman RJ. Expression of a human proprotein processing enzyme: correct cleavage of the von Willebrand factor precursor at a paired basic amino acid site. *Proceedings of the National Academy of Sciences of the United States of America*. 1990;87(23) 9378-9382.
- [108] Wasley LC, Rehemtulla A, Bristol JA, Kaufman RJ. PACE/furin can process the vitamin K-dependent pro-factor IX precursor within the secretory pathway. *Journal of Biological Chemistry*. 1993;268(12) 8458-8465.
- [109] Blais V, Fugere M, Denault JB, Klarskov K, Day R, Leduc R. Processing of proendothelin-1 by members of the subtilisin-like pro-protein convertase family. *FEBS Letters*. 2002;524(1-3) 43-48.
- [110] Dubois CM, Blanchette F, Laprise MH, Leduc R, Grondin F, Seidah NG. Evidence that furin is an authentic transforming growth factor- β 1-converting enzyme. *American Journal of Pathology*. 2001;158(1) 305-316.
- [111] Lopez de Cicco R, Watson JC, Bassi DE, Litwin S, Klein-Szanto AJ. Simultaneous expression of furin and vascular endothelial growth factor in human oral tongue squamous cell carcinoma progression. *Clinical Cancer Research*. 2004;10(13) 4480-4488.
- [112] Seidah NG, Benjannet S, Pareek S, Savaria D, Hamelin J, Goulet B, Laliberte J, Lazure C, Chretien M, Murphy RA. Cellular processing of the nerve growth factor precursor

by the mammalian pro-protein convertases. *Biochemical Journal*. 1996;314 (Pt 3) 951-960.

- [113] Bassi DE, Mahloogi H, Klein-Szanto AJ. The proprotein convertases furin and PACE4 play a significant role in tumor progression. *Molecular Carcinogenesis*. 2000;28(2) 63-69.
- [114] Maquoi E, Noel A, Franken F, Angliker H, Murphy G, Foidart JM. Inhibition of matrix metalloproteinase 2 maturation and HT1080 invasiveness by a synthetic furin inhibitor. *FEBS Letters*. 1998;424(3) 262-266.
- [115] Li X, Perez L, Pan Z, Fan H. The transmembrane domain of TACE regulates protein ectodomain shedding. *Cell Research*. 2007;17(12) 985-998.
- [116] Logeat F, Bessia C, Brou C, LeBail O, Jarriault S, Seidah NG, Israel A. The Notch1 receptor is cleaved constitutively by a furin-like convertase. *Proceedings of the National Academy of Sciences of the United States of America*. 1998;95(14) 8108-8112.
- [117] Khatib AM, Siegfried G, Prat A, Luis J, Chretien M, Metrakos P, Seidah NG. Inhibition of proprotein convertases is associated with loss of growth and tumorigenicity of HT-29 human colon carcinoma cells: importance of insulin-like growth factor-1 (IGF-1) receptor processing in IGF-1-mediated functions. *Journal of Biological Chemistry*. 2001;276(33) 30686-30693.
- [118] Maret D, Gruzglin E, Sadr MS, Siu V, Shan W, Koch AW, Seidah NG, Del Maestro RF, Colman DR. Surface expression of precursor N-cadherin promotes tumor cell invasion. *Neoplasia*. 2010;12(12) 1066-1080.
- [119] Lehmann M, Rigot V, Seidah NG, Marvaldi J, Lissitzky JC. Lack of integrin alpha-chain endoproteolytic cleavage in furin-deficient human colon adenocarcinoma cells LoVo. *Biochemical Journal*. 1996;317 (Pt 3) 803-809.
- [120] Volchkov VE, Feldmann H, Volchkova VA, Klenk HD. Processing of the Ebola virus glycoprotein by the proprotein convertase furin. *Proceedings of the National Academy of Sciences of the United States of America*. 1998;95(10) 5762-5767.
- [121] Richards RM, Lowy DR, Schiller JT, Day PM. Cleavage of the papillomavirus minor capsid protein, L2, at a furin consensus site is necessary for infection. *Proceedings of the National Academy of Sciences of the United States of America*. 2006;103(5) 1522-1527.
- [122] Klimpel KR, Molloy SS, Thomas G, Leppla SH. Anthrax toxin protective antigen is activated by a cell surface protease with the sequence specificity and catalytic properties of furin. *Proceedings of the National Academy of Sciences of the United States of America*. 1992;89(21) 10277-10281.
- [123] Gordon VM, Benz R, Fujii K, Leppla SH, Tweten RK. Clostridium septicum alpha-toxin is proteolytically activated by furin. *Infection and Immunity*. 1997;65(10) 4130-4134.

- [124] Angliker H, Wikstrom P, Shaw E, Brenner C, Fuller RS. The synthesis of inhibitors for processing proteinases and their action on the Kex2 proteinase of yeast. *Biochemical Journal*. 1993;293 (Pt 1) 75-81.
- [125] Huang YH, Lin KH, Liao CH, Lai MW, Tseng YH, Yeh CT. Furin overexpression suppresses tumor growth and predicts a better postoperative disease-free survival in hepatocellular carcinoma. *PLoS ONE*. 2012;7(7) e40738.
- [126] Bassi DE, Zhang J, Cenna J, Litwin S, Cukierman E, Klein-Szanto AJ. Proprotein convertase inhibition results in decreased skin cell proliferation, tumorigenesis, and metastasis. *Neoplasia*. 2010;12(7) 516-526.
- [127] Zhu J, Declercq J, Roucourt B, Ghassabeh GH, Meulemans S, Kinne J, David G, Vermorken AJ, Van de Ven WJ, Lindberg I, Muyldermans S, Creemers JW. Generation and characterization of non-competitive furin-inhibiting nanobodies. *Biochemical Journal*. 2012;448(1) 73-82.
- [128] Bassi DE, Lopez De Cicco R, Mahloogi H, Zucker S, Thomas G, Klein-Szanto AJ. Furin inhibition results in absent or decreased invasiveness and tumorigenicity of human cancer cells. *Proceedings of the National Academy of Sciences of the United States of America*. 2001;98(18) 10326-10331.
- [129] Huynh TT, Chan KS, Piva TJ. Effect of ultraviolet radiation on the expression of pp38MAPK and furin in human keratinocyte-derived cell lines. *Photodermatology, Photoimmunology and Photomedicine*. 2009;25(1) 20-29.
- [130] Hua H, Li M, Luo T, Yin Y, Jiang Y. Matrix metalloproteinases in tumorigenesis: an evolving paradigm. *Cellular and Molecular Life Sciences*. 2011;68(23) 3853-3868.
- [131] Noel A, Jost M, Maquoi E. Matrix metalloproteinases at cancer tumor-host interface. *Seminars in Cell and Developmental Biology*. 2008;19(1) 52-60.
- [132] Vosseler S, Lederle W, Airola K, Obermueller E, Fusenig NE, Mueller MM. Distinct progression-associated expression of tumor and stromal MMPs in HaCaT skin SCCs correlates with onset of invasion. *International Journal of Cancer*. 2009;125(10) 2296-2306.
- [133] Rucci N, Sanita P, Angelucci A. Roles of metalloproteases in metastatic niche. *Current Molecular Medicine*. 2011;11(8) 609-622.
- [134] Pytliak M, Vargova V, Mechirova V. Matrix metalloproteinases and their role in oncogenesis: a review. *Onkologie*. 2012;35(1-2) 49-53.
- [135] Murphy G, Nagase H. Progress in matrix metalloproteinase research. *Molecular Aspects of Medicine*. 2008;29(5) 290-308.
- [136] Stocker W, Bode W. Structural features of a superfamily of zinc-endopeptidases: the metzincins. *Current Opinion in Structural Biology*. 1995;5(3) 383-390.

- [137] Steinbrenner H, Ramos MC, Stuhlmann D, Sies H, Brenneisen P. UVA-mediated downregulation of MMP-2 and MMP-9 in human epidermal keratinocytes. *Biochemical and Biophysical Research Communications*. 2003;308(3) 486-491.
- [138] Newby AC. Dual role of matrix metalloproteinases (matrixins) in intimal thickening and atherosclerotic plaque rupture. *Physiological Reviews*. 2005;85(1) 1-31.
- [139] Tellier E, Negre-Salvayre A, Bocquet B, Itohara S, Hannun YA, Salvayre R, Auge N. Role for furin in tumor necrosis factor alpha-induced activation of the matrix metalloproteinase/sphingolipid mitogenic pathway. *Molecular and Cellular Biology*. 2007;27(8) 2997-3007.
- [140] Sternlicht MD, Werb Z. How matrix metalloproteinases regulate cell behavior. *Annual Review of Cell and Developmental Biology*. 2001;17 463-516.
- [141] Han YP, Tuan TL, Wu H, Hughes M, Garner WL. TNF- α stimulates activation of pro-MMP2 in human skin through NF- κ B mediated induction of MT1-MMP. *Journal of Cell Science*. 2001;114(Pt 1) 131-139.
- [142] Wan Y, Belt A, Wang Z, Voorhees J, Fisher G. Transmodulation of epidermal growth factor receptor mediates IL-1 β -induced MMP-1 expression in cultured human keratinocytes. *International Journal of Molecular Medicine*. 2001;7(3) 329-334.
- [143] Ramos MC, Steinbrenner H, Stuhlmann D, Sies H, Brenneisen P. Induction of MMP-10 and MMP-1 in a squamous cell carcinoma cell line by ultraviolet radiation. *Biological Chemistry*. 2004;385(1) 75-86.
- [144] Onoue S, Kobayashi T, Takemoto Y, Sasaki I, Shinkai H. Induction of matrix metalloproteinase-9 secretion from human keratinocytes in culture by ultraviolet B irradiation. *Journal of Dermatological Science*. 2003;33(2) 105-111.
- [145] McCawley LJ, Li S, Benavidez M, Halbleib J, Wattenberg EV, Hudson LG. Elevation of intracellular cAMP inhibits growth factor-mediated matrix metalloproteinase-9 induction and keratinocyte migration. *Molecular Pharmacology*. 2000;58(1) 145-151.
- [146] Fisher GJ, Voorhees JJ. Molecular mechanisms of photoaging and its prevention by retinoic acid: ultraviolet irradiation induces MAP kinase signal transduction cascades that induce Ap-1-regulated matrix metalloproteinases that degrade human skin in vivo. *Journal of Investigative Dermatology Symposium Proceedings*. 1998;3(1) 61-68.
- [147] Fagot D, Asselineau D, Bernerd F. Matrix metalloproteinase-1 production observed after solar-simulated radiation exposure is assumed by dermal fibroblasts but involves a paracrine activation through epidermal keratinocytes. *Photochemistry and Photobiology*. 2004;79(6) 499-505.
- [148] Han YP, Downey S, Garner WL. Interleukin-1 α -induced proteolytic activation of metalloproteinase-9 by human skin. *Surgery*. 2005;138(5) 932-939.

- [149] Cao J, Rehemtulla A, Pavlaki M, Kozarekar P, Chiarelli C. Furin directly cleaves proMMP-2 in the trans-Golgi network resulting in a nonfunctioning proteinase. *Journal of Biological Chemistry*. 2005;280(12) 10974-10980.
- [150] Kappes UP, Luo D, Potter M, Schulmeister K, Runger TM. Short- and long-wave UV light (UVB and UVA) induce similar mutations in human skin cells. *Journal of Investigative Dermatology*. 2006;126(3) 667-675.
- [151] Ratushny V, Gober MD, Hick R, Ridky TW, Seykora JT. From keratinocyte to cancer: the pathogenesis and modeling of cutaneous squamous cell carcinoma. *Journal of Clinical Investigation*. 2012;122(2) 464-472.
- [152] Nickoloff BJ, Qin JZ, Chaturvedi V, Bacon P, Panella J, Denning MF. Life and death signaling pathways contributing to skin cancer. *Journal of Investigative Dermatology Symposium Proceedings*. 2002;7(1) 27-35.
- [153] Besaratinia A, Pfeifer GP. Sunlight ultraviolet irradiation and BRAF V600 mutagenesis in human melanoma. *Human Mutation*. 2008;29(8) 983-991.
- [154] Thiery JP. Epithelial-mesenchymal transitions in tumour progression. *Nature Reviews Cancer*. 2002;2(6) 442-454.
- [155] Qi J, Chen N, Wang J, Siu CH. Transendothelial migration of melanoma cells involves N-cadherin-mediated adhesion and activation of the β -catenin signaling pathway. *Molecular Biology of the Cell*. 2005;16(9) 4386-4397.
- [156] Berthet V, Rigot V, Champion S, Secchi J, Fouchier F, Marvaldi J, Luis J. Role of endoproteolytic processing in the adhesive and signaling functions of α v β 5 integrin. *Journal of Biological Chemistry*. 2000;275(43) 33308-33313.
- [157] Siegfried G, Basak A, Cromlish JA, Benjannet S, Marcinkiewicz J, Chretien M, Seidah NG, Khatib AM. The secretory proprotein convertases furin, PC5, and PC7 activate VEGF-C to induce tumorigenesis. *Journal of Clinical Investigation*. 2003;111(11) 1723-1732.
- [158] McMahon S, Laprise MH, Dubois CM. Alternative pathway for the role of furin in tumor cell invasion process. Enhanced MMP-2 levels through bioactive TGF β . *Experimental Cell Research*. 2003;291(2) 326-339.
- [159] Cheng M, Watson PH, Paterson JA, Seidah N, Chretien M, Shiu RP. Pro-protein convertase gene expression in human breast cancer. *International Journal of Cancer*. 1997;71(6) 966-971.
- [160] Page RE, Klein-Szanto AJ, Litwin S, Nicolas E, Al-Jumaily R, Alexander P, Godwin AK, Ross EA, Schilder RJ, Bassi DE. Increased expression of the pro-protein convertase furin predicts decreased survival in ovarian cancer. *Cellular Oncology*. 2007;29(4) 289-299.

- [161] Mercapide J, Lopez De Cicco R, Bassi DE, Castresana JS, Thomas G, Klein-Szanto AJ. Inhibition of furin-mediated processing results in suppression of astrocytoma cell growth and invasiveness. *Clinical Cancer Research*. 2002;8(6) 1740-1746.
- [162] Bassi DE, Mahloogi H, Al-Saleem L, Lopez De Cicco R, Ridge JA, Klein-Szanto AJ. Elevated furin expression in aggressive human head and neck tumors and tumor cell lines. *Molecular Carcinogenesis*. 2001;31(4) 224-232.
- [163] Bassi DE, Mahloogi H, Lopez De Cicco R, Klein-Szanto A. Increased furin activity enhances the malignant phenotype of human head and neck cancer cells. *American Journal of Pathology*. 2003;162(2) 439-447.
- [164] Dumas V, Kanitakis J, Charvat S, Euvrard S, Faure M, Claudy A. Expression of basement membrane antigens and matrix metalloproteinases 2 and 9 in cutaneous basal and squamous cell carcinomas. *Anticancer Research*. 1999;19(4B) 2929-2938.
- [165] Varani J, Hattori Y, Chi Y, Schmidt T, Perone P, Zeigler ME, Fader DJ, Johnson TM. Collagenolytic and gelatinolytic matrix metalloproteinases and their inhibitors in basal cell carcinoma of skin: comparison with normal skin. *British Journal of Cancer*. 2000;82(3) 657-665.
- [166] Hofmann UB, Eggert AA, Blass K, Brocker EB, Becker JC. Expression of matrix metalloproteinases in the microenvironment of spontaneous and experimental melanoma metastases reflects the requirements for tumor formation. *Cancer Research*. 2003;63(23) 8221-8225.
- [167] Hernandez-Perez M, Mahalingam M. Matrix metalloproteinases in health and disease: insights from dermatopathology. *American Journal of Dermatopathology*. 2012;34(6) 565-579.
- [168] Wan YS, Wang ZQ, Shao Y, Voorhees JJ, Fisher GJ. Ultraviolet irradiation activates PI 3-kinase/AKT survival pathway via EGF receptors in human skin in vivo. *International Journal of Oncology*. 2001;18(3) 461-466.

IntechOpen