

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



---

# Quinolines, Isoquinolines, Angustureine, and Congeneric Alkaloids — Occurrence, Chemistry, and Biological Activity

---

Gaspar Diaz, Izabel Luzia Miranda and  
Marisa Alves Nogueira Diaz

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/59819>

---

## 1. Introduction

The alkaloids are a family of compounds widely found in nature. Therefore, they are nitrogenous secondary metabolites heterocyclic derivatives of amino acids or by the transamination process, which confers basic character [1].

The term alkaloid, linguistically derived from the Arabic *al-quali* (ash plants), is used to designate pharmacologically active nitrogen compounds found predominantly in Angiosperms [2]. Their distribution is not uniform in that division. It is estimated that approximately 40% of all plant families have at least one plant species that contains alkaloids. They are easily found in Fabaceae and Solanaceae but are rare in Gymnosperms and Pteridophytes such as ferns and monocots [3]. Although most occur in plants, alkaloids have also been isolated from algae, insects, marine and land animals, microorganisms, and fungi [4].

Since the dawn of civilization, there are reports of the use of plant extracts containing alkaloids for various purposes. The death of Socrates in 399 BC, for example, was attributed to the consumption of the extract of *Conium maculatum* containing the alkaloid coniine. There are also reports that during the last century BC, Cleopatra employed extracts of *Hyoscyamus muticus* containing the alkaloid atropine to dilate the pupils to look more attractive. It is known that medieval European women employed extract from *Atropa belladonna* with the same goal. Moreover, 25% of contemporary medicines are plant-derived. For example, morphine is extracted from the poppy. It is a narcotic drug used as analgesic and marketed since 1827 by Merck [2].

Several alkaloids (papaverine, morphine, and cocaine) can cause different stimuli in the nervous system of an animal; a common phenomenon in ecological relationships may constitute, for example, a defense mechanism by plants against herbivores [1].

The importance of alkaloids in the development of medicine is unquestionable since several advances made in the battle against diseases like malaria, leukemia, cancer, and neurodegenerative diseases would not have occurred without the use of these substances [1].

This chapter focuses on the occurrence occurring, chemistry, and biological activity of the quinoline and isoquinoline alkaloids, including the angustureine and congeneric alkaloids.

## 2. Quinoline and isoquinoline alkaloids


Among the various classes of alkaloid compounds, this section highlights the quinoline and isoquinoline alkaloids. They were originally obtained from natural sources, whose remarkable biological activities and relatively simple structures have attracted great interest in the scientific community, especially researchers involved in the chemistry of natural products. However, these compounds have also attracted the interest of synthetic organic chemists due to the need to obtain increased amounts aimed at additional biological research, as well as in developing efficient synthetic routes for these alkaloids and their derivatives, whose chemical and biological properties could become greatly enhanced by the design of new structures from these modifications.

### 2.1. Methods of extraction of quinoline and isoquinoline alkaloids

The general method for extracting and isolating alkaloids from plants consists of an acid-base extraction. The dried and pulverized plant is extracted with organic solvents or with acidified water. When the extraction is carried out with organic solvents immiscible with water, the plant sample is alkalized prior to being extracted. For this, the sample is wrapped in filter paper, as a cartridge, humidified with dilute basic solution, e.g., ammonium hydroxide, and extracted under heating using Soxhlet apparatus with organic solvent such as ethyl ether, chloroform, or dichloromethane. After extraction, the solvent volume is reduced by half in rotary evaporator under reduced pressure. The remaining volume of the solvent containing the alkaloid residue is transferred to a separatory funnel and extracted with aqueous phosphoric acid solution (pH 1 to 2) or hydrochloric acid. The wetting of the plant sample with basic solution allows the alkaloids are released from organic acids, thus facilitating its extraction by organic solvents. Meanwhile, when using an acidified solution, alkaloids tend to form salts with these strong acids, facilitating their removal from the two-phase system, water-solvent for the separation of these into the aqueous phase [5].


Another extraction technique uses polar solvent such as ethanol, for example. However, if a plant contains a high amount of lipids, a preextraction with a nonpolar solvent such as hexanes or petroleum ether is necessary for removing these lipids. Next, the organic solvent is concentrated under reduced pressure and the residue solubilized in water [6]. As most alkaloids are

basic or are found in salt form, aqueous solutions of phosphoric acid (pH 1 to 2) or hydrochloric acid are used for their removal. Neutral amide alkaloids, such as colchicine and piperine (Figure 1), remain in the organic phase when treated with an organic solvent (usually ethyl acetate) during the extraction process, while most of the other alkaloids may be extracted using an organic solvent only after neutralization of the aqueous phase with a base [4].



**Figure 1.** Structure of the neutral amide alkaloids colchicine and piperine.

Distillation methods are rarely employed for isolating alkaloids. This is possible only in cases of low molecular weight alkaloids like coniine and sparteine (Figure 2) [4].



**Figure 2.** Structure of the alkaloids coniine and sparteine.

Generally, the crude extract is purified by column chromatography (CC), employing silica gel or alumina as the stationary phase and a mixture of appropriate solvents as the mobile phase, following an appropriate choice made via thin layer chromatography (TLC). Often, alkaloids still need additional purification by recrystallization, employing solvent systems such as ethanol/water, methanol/chloroform, or acetone/methanol [4].

While these techniques are successful to extract and to isolate most alkaloids, for those highly water soluble and, therefore, partially or totally insoluble in organic solvents, these methods are not appropriate [6].

Although generally higher solubility in aqueous solutions involves greater bioactivities [6], the employment of more suitable methods for the extraction of highly water-soluble alkaloids is very useful.

In the case of quaternary alkaloids, the residual aqueous solution should be tested with a reagent suitable for the detection of alkaloids (Dragendorff reagent, for example) after completion of conventional partition methods. If it is positive, one should employ methods such as direct crystallization, precipitation as insoluble salts, extraction with polar

solvent water-immiscible, or formation of pseudobases to extract the alkaloids from the aqueous phase [6].

Alkaloids *N*-oxides can be isolated by indirect methods or chemical derivatization, through a reaction of reduction to the corresponding tertiary basis and extracted using the conventional method, and subsequently regenerated by an oxidation reaction, using oxidizing agents such as peroxide hydrogen or *m*-chloroperbenzoic acid. In some cases, the *N*-oxides containing higher aliphatic chains can be almost completely extracted by increasing the basicity of the aqueous phase, but for those who have high levels of hydroxylation, it is infeasible [6].





The methods employed to extract most common polyhydroxylated alkaloids are the direct extraction and ion exchange chromatography [6].












After extraction of alkaloids, a quantitative analysis that may be performed by traditional methods such as simple weighing or base titration after dilution under acidic conditions is recommended. Other widely used techniques for the quantification of alkaloids are by employing the HPLC apparatus with UV detection and by testing the refractive index depending on the structure of the isolated compound(s) [5].












## 2.2. Occurrence in nature of quinoline and isoquinoline alkaloids











The quinoline and isoquinoline alkaloids were initially extracted from coal tar in 1834 and 1835, respectively. Quinoline, which has high boiling temperature, is commonly employed in organic synthesis as a solvent. Isoquinoline has a low melting temperature and both have moderate basicity ( $pK_a = 4.9$  and  $5.1$ , respectively) [7].








Although the vast majority of quinoline and isoquinoline alkaloids derived from flowered plants, they can also be isolated from animals and microorganisms, some representative examples are given in Table 1 [7].

Structure	Usual name	Source	Biological activity
	Argemonine	<i>Argemone</i> sp.	-
	Berberine	Family of <i>Berberidaceae</i>	Antiprotozoal Antimalarial Antibacterial Antidiarrheal
	Berlambine	<i>Thalictrum foliolosum</i>	-
	Camptothecin	<i>Camptotheca acuminata</i>	Topoisomerase inhibitor (anticancer agent)

Structure	Usual name	Source	Biological activity
	Canadine	Family of <i>Berberidaceae</i>	-
	Cinchonine	<i>Cinchona</i> (tropical species belonging to the family <i>Rubiaceae</i> )	Antimalarial
	Codeine	<i>Opium</i>	Analgesic Antitexpectorant
	Dictamnine	<i>Dictamnus albus</i>	hepatotoxicity (cytotoxicity to HepG2 cells)
	Emetine	Family of <i>Rubiaceae</i>	Emetic Antiamoebic
	Erysotramidine	<i>Erythrina genus</i>	Antifeedant
	Erythraline	<i>Erythrina genus</i>	Antifeedant
	Glaucine	<i>Glaucium genus</i>	Antitexpectorant
	Haemanthamine	<i>Narcissus confusus</i>	-
	Hydrastine	<i>Hydrastis canadensis</i>	Uterine hemostatic Antiseptic
	Hydroquinine	<i>Cinchona</i> (tropical species belonging to the family <i>Rubiaceae</i> )	Depigmenting

Structure	Usual name	Source	Biological activity
	Jamine	<i>Cocculus hirsutus</i>	Anti-hyperglycemic
	Lavendamycin	<i>Streptomyces leventulae</i>	Antitumor antibiotic
	Lephocereine	<i>Lephocereus schotti</i>	-
	Lycorine	<i>Lycoris radiata</i>	hypotensive
	Maritidine	-	-
	Morphine	<i>Opium</i>	Analgesic
	Nandinine	<i>Nandina domestica</i>	-
	Nantenine	<i>Nandina domestica</i>	$\alpha$ 1 Adrenergic and serotonin antagonist 5-HT <sub>2A</sub>
	Nitidine	<i>Zanthoxylum nitidum</i>	Topoisomerase inhibitor (anticancer agent)
	N-norlaudanosine	-	-
	Palmitine	<i>Jateorhiza palmate</i>	-

Structure	Usual name	Source	Biological activity
	Papaverine	<i>Opium</i>	Vasodilator
	Pellotine	<i>Lophophora williamsii</i>	-
	Protoemetine	<i>Alangium lamarckii</i>	-
	Protoemetinol	<i>Alangium lamarckii</i>	Emetic Antiamoebic Anticancer
	Psychotrine	<i>Uragoga ipecacuanha</i>	Inhibitor of HIV-1 transcriptase
	Quinidine	<i>Cinchona</i> (tropical species belonging to the family Rubiaceae)	Antiarrhythmic Antimalarial
	Quinine	<i>Cinchona</i> (tropical species belonging to the family Rubiaceae)	Antimalarial Muscle relaxant
	d-Quinotoxine	-	-
	Reticuline	<i>Anona reticulata</i> <i>Opium</i>	-
	Sandramycin	<i>Norcardioides</i> sp. (microorganism)	Antitumor antibiotic


Structure	Usual name	Source	Biological activity
	Skimmianine	<i>Skimmia japonica</i>	Sedative Anticonvulsant
	Sanguinarine	<i>Sanguinaria</i>	Antibacterial
	Stepholidine	<i>Stephania intermedia</i>	D <sub>1</sub> receptor agonist D <sub>2</sub> receptor antagonist
	Streptonigrin	<i>Streptomyces flocculus</i>	Antitumor antibiotic
	Tetrahydropa-Imitine	<i>Corydalis family</i>	Analgesic
	Thalictricavine	<i>Corydalis tuberosa</i>	-
	Tubocurarin	<i>Chododendron tomentosum</i>	Neuromuscular blocking agent

**Table 1.** Quinoline and isoquinoline alkaloids: natural sources and biological activities.


### 2.3. Biosynthetic origin of quinoline and isoquinoline alkaloids

The most One of the systems used for the classification of alkaloids is based on the kind of nitrogen heterocycle in the structure and its botanical origin (plant family which is extracted); thus, the name given to an alkaloid derived from the genus or species of organism produced, including the term "ine." Another commonly used system takes into account its precursor amino acid. Thus, alkaloid derivatives of tryptophan and tyrosine are biosynthetically classified as quinoline and isoquinoline, respectively (Figure 3) [3].

Thus, quinoline alkaloids are derived from 3-hydroxyanthranilic acid, a metabolite formed from tryptophan through a sequence of enzymatic reactions (Figure 4). The condensation between 3-hydroxyanthranilic acid and malonyl-SCoA, followed by cyclization, produces quinoline alkaloid (Figure 4) [8-10].



**Figure 3.** Structure of quinoline and isoquinoline alkaloids.




**Figure 4.** Biosynthesis of quinoline alkaloids.

In the case of isoquinoline systems, the biosynthesis proceeds from the cyclization of the Schiff base formed between the dopamine and an aliphatic aldehyde [8]. Dopamine is obtained from the hydroxylation and decarboxylation of tyrosine. A second intermediate, *p*-hydroxyphenyl acetaldehyde, can also be formed by transamination, decarboxylation, and hydroxylation of tyrosine. The condensation of these intermediates followed by a sequence of steps (cyclization, hydroxylation, and methylation) produces the (*S*)-reticuline, a biosynthetic intermediate of all isoquinoline alkaloids (Figure 5) [11].


## 2.4. Biological activities of quinoline and isoquinoline alkaloids

In addition to the biological activities shown in Table 1, many quinoline alkaloids and their analogues represent the most important drugs currently used to combat malaria, with chloroquine, amodiaquine, piperazine, primaquine, quinine, and mefloquine (Figure 6)




**Figure 5.** Biosynthesis of isoquinoline alkaloids.

being the most representative drugs of this group [12]. Even today, many studies are still being conducted to increase the efficiency of quinoline derivatives such as AZT-chloro-quinoline, quinine-dihydroartemisinin, and MEFAS, a salt derived from mefloquine and artesunate [13] (Figure 7).




**Figure 6.** Structure of some quinoline alkaloids and their analogues.




**Figure 7.** Structure of other quinoline derivatives.

On the other hand, the quinoline nucleus has also demonstrated a critical role in the development of new anticancer drugs. Some of its derivatives showed excellent results on various types of cancer cells, through different mechanisms of action. Recently, three protein kinase inhibitors (Bosutinib, Lenvatinib, and Cabozantinib) and an inhibitor of farnesyltransferase (Tipifarnib) (Figure 8), considered as potential anticancer agents, entered into phase of clinical trials [14].



**Figure 8.** Structure of bosutinib, lenvatinib, cabozantinib, and tipifarnib quinoline nucleus.

Knowing that functionalized quinolines have also shown high anti-carcinogenic potential, in addition to acting as anti-angiogenic agents, and inhibitors of telomerase in various human tumor cells, Muñoz and coworkers [15] selected the DM8 and DM12 tetrahydroquinolines (Figure 9) for testing cytotoxic activity. They observed that individually these compounds significantly inhibited cell growth of breast cancer, and when tested concomitantly with other two anticancer drugs, they work synergistically to increase their cytotoxic activities. The results obtained for the DM12 quinoline made this compound a promising candidate as a new adjunctive anticancer agent.




**Figure 9.** Structure of the DM8 and DM12 tetrahydroquinoline alkaloids.

Furthermore, Villemagne and Okamura [16] obtained promising results with other quinoline derivatives used as selective ligands for Tau proteins, associated with brain damage, which represent a risk factor for diseases such as Alzheimer's, which allowed the investigation of the causes, diagnosis, and treatment of neurodegenerative diseases, encephalopathy, and traumatic brain injury.

Biological activities of hundreds of other substituted quinolines have been reported, many of which are promising in terms of their potential as pharmaceutical agents. 2-Methyl-5-hydroxy-1,2,3,4-tetrahydroquinoline exhibits analgesic activity with a potency one-eighth that of morphine [17]. 1,2,3,4-Tetrahydroquinoline-4-carboxylic acid is used in tissue-irrigating solutions [18]. A wide array of other biological activities have been reported, including *inter alia* inhibition of (H<sup>+</sup>/K<sup>+</sup>)-ATPase [19], blood serum monoamine oxidase [20], angiotensin I converting enzyme [21], lipoxygenase [22], lipid peroxidation [23], bone resorption [24], leukotriene synthesis [25, 26], and bacterial dihydrofolate reductase [27]. Relevance to many other indications has also been noted [28-54].


In addition to the numerous utilities in the pharmacological area, studies are being performed in order to investigate the potential activities of these alkaloids as agrochemical agents. Sanguinarine and chelerythrine (Figure 10), for example, have shown high activity against fungi and phytopathogenic bacteria *in vitro*, with sanguinarine being the most effective against *Rhizoctonia solani* fungus, which has a wide range of hosts, which causes disease in most plants grown in the world [55].



**Figure 10.** Structure of the sanguinarine and chelerythrine agrochemical agents.

### 3. Tetrahydroquinoline alkaloids angustureine, galipeine, galipinine, and cuspareine

Among the simplest natural alkaloids that contain the 1,2,3,4-tetrahydroquinoline core are angustureine and its congeners, galipeine, cuspareine, and galipinine (Figure 11). The structures are more narrowly described as 2-alkyl-1-methyl-1,2,3,4-tetrahydroquinolines. They are chiral by virtue of the stereogenic center at position 2. There is an excellent review of the chemistry and synthesis of 1,2,3,4-tetrahydroquinolines by Diaz and Dudley [56].



**Figure 11.** Structure of the 1,2,3,4-tetrahydroquinolin alkaloids angustureine, galipeine, cuspareine, and galipinine.

In 1999, Jacquemond-Collet and coworkers [57] isolated and characterized these four tetrahydroquinoline alkaloids (Figure 11) from the extract of the bark of *Galipea officinalis*, commonly known as “angostura” [58]. The genus *Galipea* comprises about 20 species that are found predominantly in the Northern of South America. *G. officinalis* is a shrub known by rural communities in the Venezuelan mountains for their pharmacological activities, and its extract is used in the control of dyspepsia, dysentery, and chronic diarrhea, in addition to presenting antimalarial, cytotoxic [59], molluscicidal activities, and antimicrobial properties, inhibiting the growth of *Mycobacterium tuberculosis*, the etiological agent of tuberculosis [60].


In later trials, galipinine and galipeine exhibited promising biological activities *in vitro* such as antimalarial ( $IC_{50}$ : 0.24-6.12 and 0.33-13.78 mM, respectively) for the protozoa species *Plasmodium falciparum*, one of the species causing malaria that is already resistant to chloroquine, one of the main drugs used to combat the disease [61].

Due to their structural simplicities associated with the promising pharmacological activities, these alkaloids, especially angustureine, have attracted the attention of the synthetic organic community. For this alkaloid alone, more than 24 different syntheses have already been described [62-85]. Some recent examples of the syntheses of these alkaloids are described below.

Diaz *et al.* [83] described the enantioselective synthesis of both alkaloids (*R*)-(-)- and (*S*)-(+)-angustureine with overall yields of 80% and 44%, respectively, and excellent enantiomeric excesses (95% and 96%, respectively), starting from the (*S*)- $\beta$ -amino ester and (*R*)-sodium carboxylate, prepared following enzymatic resolution of the  $\beta$ -amino ester racemate [86, 87].

Foubelo *et al.* [85] described a second-generation synthesis of (-)-angustureine by way of 2-allyl-tetrahydroquinoline. A further sequence of three steps from this intermediate provided angustureine in 36% overall yield. However, 2-allyl-tetrahydroquinolines can serve as common precursors to various 2-substituted tetrahydroquinoline alkaloids by making use of the cross-metathesis reaction with the Hoveyda-Grubbs reagent (second-generation ruthenium catalyst) to achieve the appropriate side-chain homologation events.

In turn, Lam *et al.* [88] synthesized analogues of tetrahydroquinoline alkaloids employing asymmetric hydrogenation catalyzed by iridium. These compounds were tested *in vitro* for antitumor activities against lung cancer, breast cancer, hepatocellular carcinoma, and sarcoma cancer and have achieved remarkable results, as well as *in vivo*, using mice as animal models receiving hepatocellular tumor grafts, being that 1,2,3,4-tetrahydroquin-8-ol (Figure 12) revealed the highest toxicity to human cancer cells.



1,2,3,4-tetrahydroquin-8-ol

**Figure 12.** Structure of 1,2,3,4-tetrahydroquin-8-ol anticancer agent.

## 4. Concluding remarks

This chapter is an illustration of the fascinating world of chemistry of quinoline and isoquinoline alkaloids. The range of pharmacological and agrochemical activities, among others, associated with their biosynthetic routes and various natural sources demonstrates the creative universe of the nature of these alkaloids. This universe of compounds arouses interest to organic chemists in general, stimulated by the simplicity and its varied forms of exploitation throughout constant search for the discovery of novel pharmacological agents targeting the treatment of several diseases that affect society.

## 5. Future directions

In this chapter, we can verify the importance of quinoline and isoquinoline alkaloids due to their potent biological activities demonstrated by *in vitro* and *in vivo* assays, as well as

agrochemical agents reported throughout this chapter. Based on these facts, this family of compounds is presented as an inexhaustible source for research groups working with natural and synthetic products for the pharmaceutical area. In this context, our group intends to continue the study of these alkaloids with the synthesis of novel quinoline and isoquinoline derivatives, aimed at the discovery of new pharmacological agents.

## Acknowledgements

The authors are grateful to CAPES and FAPEMIG for their financial support.

## Author details

Gaspar Diaz<sup>1\*</sup>, Izabel Luzia Miranda<sup>2</sup> and Marisa Alves Nogueira Diaz<sup>3</sup>

\*Address all correspondence to: [gaspardm@qui.ufmg.br](mailto:gaspardm@qui.ufmg.br)

1 Departament of Chemistry, Federal University of Minas Gerais, Belo Horizonte, Minas Gerais, Brazil

2 Departament of Chemistry, Federal University of Viçosa, Viçosa, Minas Gerais, Brazil

3 Departament of Biochemistry and Molecular Biology, Federal University of Viçosa, Viçosa, Minas Gerais, Brazil

## References

- [1] Aniszewski T. Alkaloids—Secrets of Life. Amsterdam: Elsevier Academic Press; 2007.
- [2] Kutchan TM. Alkaloid biosynthesis—the basis for metabolic engineering of medicinal plants. *Plant Cell* 1995;7(7) 1059-1070.
- [3] Ruiz PG. Productos Naturales. Pamplona: Universidad Pública de Navarra; 2002.
- [4] Mann J, Davidson RS, Hobbs JB, Banthorpe DV, Harborne JB. Natural Products: Their Chemistry and Biological Significance. London: Longman Scientific & Technical; 1996.
- [5] Simões CMO. Farmacognosia—Da planta ao medicamento. Editoras UFRGS e UFSC; 2010.

- [6] Colegate SM, Molyneus RJ. Bioactive Natural Products: Detection, Isolation and Structural Determination. Boca Raton, Florida: CRC Press; 1993.
- [7] Majumdar KC, Chattopadhyay SK. Heterocycles in Natural Product Synthesis. Singapore: Wiley-VCH; 2011.
- [8] Mann J. Secondary Metabolism. Oxford: Clarendon Press; 2005.
- [9] Souza MVN, Almeida MV, Hyaric M, Cardoso SH, Amarante WG. Métodos de preparação e atividade biológica do ácido quinolínico e derivados. Química Nova 2003;26(5) 694-698.
- [10] Sultana N. Medicinal properties and biosynthetic studies on indigenous medicinal plant *Skimmia laureola*. Critical Review in Pharmaceutical Sciences 2013;2(2) 13-42.
- [11] Civjan N. Natural Products in Chemical Biology. Hoboken, New Jersey: John Wiley & Sons; 2012.
- [12] Kaur K, Jain M, Reddy RP, Jain R. Quinolines and structurally related heterocycles as antimalarials. European Journal of Medicinal Chemistry 2010;45(8) 3245-3264.
- [13] Boechat N, Ferreira MLG, Pinheiro LCS, Jesus AML, Leite MMM, Junior CCS, Aguiar ACC, Andrade IM, Krettli AU. New Compounds Hybrids 1*H*-1,2,3-Triazole-Quinoline Against *Plasmodium falciparum*. Chemical Biology and Drug Design 2014;84(3) 325-332.
- [14] Afzal O, Kumar S, Haider MR, Ali MR, Kumar R, Jaggi M, Bawa S. A review on anti-cancer potential of bioactive heterocycle quinoline. European Journal of Medicinal Chemistry 2014. DOI: 10.1016/j.ejmech.2014.07.044. *In press*.
- [15] Muñoz A, Sojo F, Arenas DRM, Kouznetsov VV, Arvelo F. Cytotoxic effects of new trans-2,4-diaryl-r-3-methyl-1,2,3,4-tetrahydroquinolines and their interaction with antitumoral drugs gemcitabine and paclitaxel on cellular lines of human breast cancer. Chemico-Biological Interactions 2011;189(3) 215-221.
- [16] Villemagne VL, Okamura N. *In vivo* tau imaging: obstacles and progress. Alzheimer's and Dementia 2014;10(3) 254-264.
- [17] Ferranti A, Garuti L, Giovanninetti G, Gaggi R, Roncada P, Nardi P. Preparation and analgesic activity of tetrahydroquinolines and tetrahydroisoquinolines. Farmaco Edizione Scientifica 1987;42(4) 237-249.
- [18] LeClerc G, Ruhland B, Andermann G, De Burlet G, Dietz M. For controlling tissue stability during eye surgery: antisepting agent 1990; US Patent US 4,952,573, 1990.
- [19] Uchida M, Chihiro M, Morita S, Kanbe T, Yamashita H, Yamasaki K, Yabuuchi, Y, Nakagawa K. Studies on proton pump inhibitors. II. Synthesis and antiulcer activity of 8-[(2-benzimidazolyl)sulfinylmethyl]-1,2,3,4-tetrahydroquinolines and related compounds. Chemical and Pharmaceutical Bulletin 1989;37(8) 2109-2116.

- [20] Gracheva IN, Gridneva LI, Tochilkin AI, Gorkin VZ. Hydrazides and sulfonyl hydrazides from the series of quinoline are human serum amine oxidase inhibitors. *Khimiiko-Farmatsevticheskii Zhurnal* 1988;22(11) 1336-1339.
- [21] Miller KE, Huang CT, Portlock DE, Wright GC. Angiotensin I converting enzyme inhibitors containing unnatural alpha-amino acid analogues of phenylalanine. *Life Sciences* 1987;40(1) 63-70.
- [22] Stevens RW, Ikeda T, Wakahayashi H, Nakane M. Antiinflammatory hydroxamic acids and *N*-hydroxyureas. US Patent 1993; US 5,256,789.
- [23] Kihara N, Tomino I, Tan H, Ishihara T. Novel nitrogen-containing compound. European Patent 1988; EP 0289.36.5.
- [24] Kojima E, Saito K. Cyclic anthranilic acid derivatives as therapeutic agents for treatment of metabolic bone disease. European Patent 1990; EP 402859.
- [25] Youssefyeh RD, Magnien E, Lee TDY, Chan WK, Lin CJ, Galemme JrRA, Johnson JrWH, Tan J, Campbell HF, Huang FC, Nuss GW, Carnathan GW, Sutherland CA, Van Inwegen RG. Development of a novel series of (2-quinolinylmethoxy)phenyl-containing compounds as high-affinity leukotriene receptor antagonists. 1. Initial structure-activity relationships. *Journal of Medicinal Chemistry* 1990;33(4) 1186-1194.
- [26] Paris D, Cottin M, Demonchaux P, Augert G, Dupassieux P, Lenoir P, Peck MJ, Jaserand D. synthesis, structure-activity relationships, and pharmacological evaluation of pyrrolo[3,2,1-ij]quinoline derivatives: potent histamine and platelet activating factor antagonism and 5-lipoxygenase inhibitory properties. potential therapeutic application in asthma. *Journal of Medicinal Chemistry* 1995;38(4) 669-685.
- [27] Rauckman BS, Tidwell MY, Johnson JV, Roth B. 2,4-Diamino-5-benzylpyrimidines and analogs as antibacterial agents. 10. 2,4-Diamino-5-(6-quinolylmethyl)- and -[(tetrahydro-6-quinolyl)methyl]pyrimidine derivatives. Further specificity studies. *Journal of Medicinal Chemistry* 1989;32(8) 1927-1935.
- [28] Ogawa H, Miyamoto H, Kondo K, Yamashita H, Nakaya K, Komatsu H, Tanaka M, Takara S, Tominaga M, Yabuchi Y. Preparation of *N*-benzoyl benzo-fused heterocyclic compounds as vasopressin antagonists. *Kokai Tokkyo Koho* 1992; JP 04321669.
- [29] Bigg D, Mangane M. Tetrahydroquinoline derivatives, their preparation and their therapeutic application 1986; France Patent FR 2,576,308.
- [30] Nagasaka T, Kosugi Y, Kawahara T, Kakimoto M, Tamuru K, Hirata A. Preparation of 1,4-dihydro-4-phenyl-3,5-pyridinedicarboxylic acids as calcium antagonists 1993. Japanese Patent 05,339,263.
- [31] Moon MW, Morris JK, Heier RF, Chidester CG, Hoffmann WE, Piercey MF, Althaus JS, VonVoigtlander PF, Evans DL, Figur LM, Lahti RA. Dopaminergic and serotonergic activities of imidazoquinolinones and related compounds. *Journal of Medicinal Chemistry* 1992;35(6) 1076-1092.

- [32] Moon MW, His RSP. Synthesis of (R)-5-(Di[2,3-<sup>3</sup>H<sub>2</sub>]propylamino)5,6-dihydro-4*H*-imidazo[4,5,1-*ij*]quinolin-2(1*H*)-one ([<sup>3</sup>H]U-86170) and (R)-5-([2,3-<sup>3</sup>H<sub>2</sub>]Propylamino)-5,6-dihydro-4*H*-imidazo[4,5,1-*ij*]quinolin-2(1*H*)-one ([<sup>3</sup>H]U-91356). *Journal of Labelled Compounds and Radiopharmaceuticals* 1992;31(11), 933-943.
- [33] Buzas A, Ollivier R, El Ahmad Y, Laurent E. 1,4-Dialkylpiperazine derivatives, method for obtaining them, and pharmaceutical compositions containing them. PCT International Applications 1993; WO 9316057.
- [34] Agarwal SK, Saxena AK, Jain PC, Sur RN, Srimal RC, Dhawan BN, Anand N. Synthesis and structure activity relationship in 1-aryloxy-3-[*N'*-N4-arylpiperazinyl)] propanes. *Indian Journal of Chemistry, Section B* 1987; 26B, 642-646.
- [35] Uchida M, Chihiro M, Morita S, Yamashita H, Yamasaki K, Kunbe T, Yabuuchi Y, Nakagawa K. Synthesis and antiulcer activity of 4-substituted 8-[(2-benzimidazol)sulfinylmethyl]-1,2,3,4-tetra-hydroquinolines and related compounds. *Chemical and Pharmaceutical Bulletin* 1990;38(6) 1575-1586.
- [36] Uchida M, Morita S, Chihiro M. Preparation of 8-[(2-benzimidazolylthio)alkyl]hydroquinolines and their sulfoxides as antiulcer agents. *European Patent* 1987; EP 239129.
- [37] Atwal K. Indanyl- and quinolylureas and related compounds as cardiovascular agents. *European Patent* 1992; EP 488616.
- [38] Santangelo F, Casagrande C, Miragoli G, Vecchiotti V. Synthesis and positive inotropic effect of 1-alkyl- and 1-acyl-6,7-dimethoxy-3-dimethylamino-1,2,3,4-tetrahydroquinolines. *European Journal of Medicinal Chemistry* 1994;29(11), 877-882.
- [39] Galtier D, Lassalle G. Preparation of 1,2,3,4-tetrahydroquinolin-8-sulfonic acid and its chlorides as intermediates for antithrombotics. *European Patent* 1995; EP 643046.
- [40] Lassalle G, Galtier D, Galli F. Preparation of 1-[2-amino-5-[1-(triphenylmethyl)-1*H*-imidazol-4-yl]-1-oxopentyl]piperidines and their use as intermediates for antithrombotics. *European Patent* 1995; EP 643047.
- [41] Baumgarth M, Lues I, Minck KO, Beier N. Preparation of (benzocyclylethyl)arylperidines and -piperazines for treating arrhythmia and tachycardia. *German Patent* 1995; DE 4321366.
- [42] Biller SA, Misra RN. Inhibitors of leukotriene biosynthesis in macrophage cells. *US Patent* US 4,843,082, 1989.
- [43] Lukevics E, Lapina T, Segals I, Augustane I, Verovskii VN. synthesis and antitlastic activity of organosilicon derivatives of quinoline, isoquinoline, and *n*-methylpiperazine. *Khimiko-Farmatsevticheskii Zhurnal* 1988;22(8) 619-623.
- [44] Kohno Y, Kojima E. Preparation of 1,2,3,4-tetrahydroquinoline-4,8-dicarboxylates as drugs. *European Patent* 1990; EP 403980.

- [45] Kohno Y, Awano K, Ishizaki T, Kojima E, Kudoh S, Sekoe Y, Saito K. Preparation of tetrahydroquinoline acetic acid derivatives as immunosuppressants. PCT International Applications 1992; WO 9218482.
- [46] Kohno Y, Kojima E, Saito K, Kudoh S, Sekoe Y. Preparation of tetrahydroquinolines as immunosuppressants and inflammation inhibitors. Kokai Tokkyo Koho 1994; JP 06 56788.
- [47] Carling RW, Leeson PD, Moseley AM, Smith JD, Saywell K, Tricklebank MD, Kemp JA, Marshall GR, Foster AC, Grimwood S. Anticonvulsant activity of glycine-site NMDA antagonists. 2. *trans* 2-carboxy-4-substituted tetrahydroquinolines. Bioorganic and Medicinal Chemistry Letters 1993;3(1) 65-70.
- [48] Sangwan NK, Malik MS, Dhindsa KS. Synthesis of 3-benzyl-1,2,3,4-tetrahydro-7-methoxy-4-[4-(2-pyrrolidinoethoxy)phenyl]-1-tosylquinolines as potential antifertility agents. Chimica Acta Turcica 1985;13(1) 129-134.
- [49] Stevenson GL, Leeson PD, Rowley M, Sanderson I, Stansfield I. Synthesis of *cis* and *trans* 4-amido 2-carboxytetrahydroquinolines, high affinity ligands at the glycine site of the NMDA receptor. Bioorganic and Medicinal Chemistry Letters 1992;2(5) 371-374.
- [50] Regnier G, Guillonnet C, Lepagnol J, Lestage P. Morpholine derivatives, process for their preparation and pharmaceutical compositions containing them for treatment of cerebral conditions. European Patent 1991; EP 427605.
- [51] Boursier-Neyret C, Baune A, Klippert P, Castagne I, Sauveur C. Determination of S 12024 enantiomers in human plasma by liquid chromatography after chiral pre-column derivatization. Journal of Pharmaceutical and Biomedical Analysis 1993;11(11-12) 1161-1166.
- [52] Ventura C, Miller R, Wolf HP, Beier N, Jonas R, Klockow M, Lues I, Hano O, Spurgeon HA, Lakatta EG, Capogrossi MC. Novel diazinone derivatives separate myofilament  $\text{Ca}^{2+}$  sensitization and phosphodiesterase III inhibitory effects in guinea pig myocardium. Circulation Research 1992;70(6) 1081-1090.
- [53] Regnier G, Guillonnet C, Lepagnol J. Preparation of (morpholinylmethoxy) tetrahydroquinoline derivatives as drugs for treatment of cerebral ischemia. European Patent 1988; EP 286495.
- [54] Kohno Y, Saito K, Sekoe Y, Kojima E. Preparation of 6-aryl-cyclic anthranilic acids and their use as therapeutics for treatment of metabolic bone disorder. Japanese Patent 1992; JP 04316557.
- [55] Cantrell CL, Dayan FE, Duke SO. Natural products as sources for new pesticides. Journal of Natural Products 2012;75(6) 1231-1242.

- [56] Diaz MG, Dudley BG. Synthesis of 1,2,3,4-tetrahydroquinolines including angustureine and congeneric alkaloids. *Organic Preparations and Procedures International*, 2014. In press.
- [57] Jacquemond-Collet I, Hannedouche S, Fabre N, Fourasté I, Moulis C. Two tetrahydroquinoline alkaloids from *Galipea officinalis*. *Phytochemistry* 1999;51(8) 1167-1169.
- [58] Taylor LL, Goldberg FW, Hii KK. Asymmetric synthesis of 2-alkyl-substituted tetrahydroquinolines by an enantioselective aza-Michael reaction. *Organic Biomolecular Chemistry* 2012;10(22) 4424-4432.
- [59] Jacquemond-Collet I, Benoit-Vical F, Mustofa Valentin A, Stanislas E, Mallié M, Fourasté I. Antiplasmodial and cytotoxic activity of galipinine and other tetrahydroquinolines from *Galipea officinalis*. *Planta Medica* 2002;68(1) 68-69.
- [60] Houghton PJ, Watabe Y, Woldemariam TZ, Yates M. Activity of alkaloids from *Angostura* bark against *Mycobacterium tuberculosis*. *Journal of Pharmacy and Pharmacology*, 1998;50(S9) 230.
- [61] Osório EJ, Robledo SM, Bastida J. *The Alkaloids: Chemistry and Biology*. Amsterdam: Elsevier Academic Press; 2008.
- [62] Wang WB, Lu SM, Yang PY, Han XW, Zhou YG. Highly enantioselective iridium-catalyzed hydrogenation of heteroaromatic compounds, quinolines. *Journal of American Chemical Society* 2003;125(35) 10536-10537.
- [63] Avemaria F, Vanderheiden S, Braese S. The aza-xylylene Diels-Alder approach for the synthesis of naturally occurring 2-alkyl tetrahydroquinolines. *Tetrahedron* 2003;59(35) 6785-6796.
- [64] Lin XF, Li Y, Ma DW. Total synthesis of tetrahydroquinoline alkaloid (+)-angustureine. *Chinese Journal of Chemistry* 2004;22(9) 932-934.
- [65] Theeraladanon C, Arisawa M, Nakagawa M, Nishida A. Total synthesis of (+)-(S)-angustureine and the determination of the absolute configuration of the natural product angustureine. *Tetrahedron: Asymmetry* 2005;16(4) 827-831.
- [66] Lu SM, Wang YQ, Han XW, Zhou YG. Asymmetric hydrogenation of quinolines and isoquinolines activated by chloroformates. *Angewandte Chemie International Edition* 2006;45(14) 2260-2263.
- [67] Rueping M, Antonchick AP, Theissmann T. A highly enantioselective Bronsted acid catalyzed cascade reaction: organocatalytic transfer hydrogenation of quinolones and their application in the synthesis of alkaloids. *Angewandte Chemie International Edition* 2006;45(22) 3683-3686.
- [68] Ryu JS. Hydroarylation for the facile synthesis of 2-substituted tetrahydroquinoline: a concise synthesis of (+)-(S)-angustureine. *Bulletin of the Korean Chemical Society* 2006;27(5) 631-632.

- [69] Patil NT, Wu H, Yamamoto Y. A route to 2-substituted tetrahydroquinolines via palladium-catalyzed intramolecular hydroamination of aniline-alkynes. *Journal of Organic Chemistry* 2007;72(17) 6577-6579.
- [70] O'Byrne A, Evans P. Rapid synthesis of the tetrahydroquinoline alkaloids: angustureine, cuspareine and galipinine. *Tetrahedron* 2008;64(35) 8067-8072.
- [71] Shahane S, Louafi F, Moreau J, Hurvois JP, Renaud JL, Van de Weghe P, Roisnel, T. Synthesis of alkaloids of *Galipea officinalis* by alkylation of an  $\alpha$ -amino nitrile. *European Journal of Organic Chemistry* 2008;2008(27) 4622-4631.
- [72] Fustero S, Moscardó J, Jiménez D, Pérez-Carrión MD, Sánchez-Roselló M, del Pozo C. Organocatalytic approach to benzofused nitrogen-containing heterocycles: Enantioselective total synthesis of (+)-angustureine. *Chemistry a European Journal* 2008;14(32) 9868-9872.
- [73] Wang ZJ, Zhou HF, Wang TL, He YM, Fan QH. Highly enantioselective hydrogenation of quinolones under solvent-free or highly concentrated conditions. *Green Chemistry* 2009;11(6) 767-769.
- [74] Kothandaraman P, Foo SJ, Chan PW. H. Gold-catalyzed intramolecular allylic amination of 2-tosylaminophenylprop-1-en-3-ols. A concise synthesis of ( $\pm$ )-angustureine. *Journal of Organic Chemistry* 2009;74(16) 5947-5952.
- [75] Chen BL, Wang B, Lin GQ. Highly diastereoselective addition of alkynylmagnesium chlorides to N-tert-butanefulfinyl aldimines: a practical and general access to chiral  $\alpha$ -branched amines. *Journal of Organic Chemistry* 2010;75(3) 941-944.
- [76] Cruz-Lopez O, Nunez MC, Conejo-Garcia A, Kimatrai M, Campos JM. Syntheses of 2,3-dihydro-1,4-benzodioxins and bioisosteres as structural motifs for biologically active compounds. *Current Organic Chemistry* 2011;15(6) 869-887.
- [77] Bentley SA, Davies SG, Lee JA, Roberts PM, Thomson JE. Conjugate addition of lithium N-phenyl-N-( $\alpha$ -methylbenzyl)amide: application to the asymmetric synthesis of (R)-(-)-angustureine. *Organic Letters* 2011;13(10) 2544-2547.
- [78] Wang T, Zhuo LG, Li Z, Chen F, Ding Z, He Y, Fan QH, Xiang J, Yu ZX, Chan ASC. Highly enantioselective hydrogenation of quinolones using phosphine-free chiral cationic ruthenium catalysts: scope, mechanism, and origin of enantioselectivity. *Journal of the American Chemical Society* 2011;133(25) 9878-9891.
- [79] Satyanarayana G, Pflästerer D, Helmchen G. Enantioselective synthesis of tetrahydroquinolines based on iridium-catalyzed allylic substitutions: total synthesis of (+)-angustureine and (-)-cuspareine. *European Journal of Organic Chemistry* 2011;2011(34) 6877-6886.
- [80] Ye KY, He H, Liu WB, Dai LX, Helmchen G, You SL. Iridium-catalyzed allylic vinylation and asymmetric allylic amination reactions with o-aminostyrenes. *Journal of the American Chemical Society* 2011;133(46) 19006-19014.

- [81] Taylor LL, Goldberg FW, Hii KK. Asymmetric synthesis of 2-alkyl-substituted tetrahydroquinolines by an enantioselective aza-Michael reaction. *Organic and Biomolecular Chemistry* 2012;10(22) 4424-4432.
- [82] Tummatorn J, Diaz MG, Dudley GB. Synthesis of (-)-(R)-aungustureine by formal alkylation of a chiral  $\beta$ -amino ester. *Tetrahedron Letters* 2013;54(10) 1312-1314.
- [83] Diaz G, Diaz MAN, Reis MA. Enantioselective synthesis of both (-)-(R)- and (+)-(S)-aungustureine controlled by enzymatic resolution. *Journal of the Brazilian Chemical Society* 2013;24(9) 1497-1503.
- [84] Sirvent JA, Foubelo F, Yus M. Enantioselective synthesis of tetrahydroquinoline alkaloids (-)-aungustureine and (-)-cuspareine from chiral *tert*-butanesulfinyl imines. *Heterocycles* 2014;88(2) 1163-1174.
- [85] Sirvent JA, Foubelo F, Yus M. Stereoselective synthesis of indoline, tetrahydroquinoline, and tetrahydrobenzazepine derivatives from o-bromophenyl *N*-*tert*-butylsulfinyl aldiminas. *Journal of Organic Chemistry* 2014;79(3) 1356-1367.
- [86] Katayama S, Ae N, Nagata R. Enzymatic resolution of 2-substituted tetrahydroquinolines. Convenient approaches to tricyclic quinoxalinediones as potent NMDA-glycine antagonists. *Tetrahedron: Asymmetry* 1998;9(24) 4295-4299.
- [87] Nagata R, Tanno N, Kodo T, Ae N, Yamaguchi H, Nishimura T, Antoku F, Tatsuno T, Kato T, Yanaka Y, Nakamura M. Tricyclic quinoxalinediones: 5,6-dihydro-1H-pyrrolo[1,2,3-de]quinoxaline-2,3-diones and 6,7-Dihydro-1H,5H-pyrido[1,2,3-de]quinoxaline-2,3-diones as potent antagonists for the glycine binding site of the NMDA receptor. *Journal of Medicinal Chemistry* 1994;37(23) 3956-3968.
- [88] Lam K, Lee KK, Gambari R, Wong RS, Cheng GY, Tong S, Chan K, Lau F, Lai PB, Wong W, Chan AS, Kok SH, Tang JC, Chui C. Preparation of *Galipea officinalis* Hancock type tetrahydroquinoline alkaloid analogues as anti-tumor agents. *Phytomedicine* 2013;20(2) 166-171.

IntechOpen