We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Imaging of Brain Tumours

Mike Fay, Chris Bell, Nick Dowson,
Simon Puttick and Stephen Rose

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/59981>

---

1. Introduction

The last 20 years have seen a dramatic and rapid improvement in imaging technology. Developments in magnetic resonance (MR) imaging and more recently in positron emission tomography (PET) have allowed unique insights into tumour biology. Although careful evaluation this technology's clinical value is still being undertaken, improved imaging techniques have potential to assist many of the difficult problems currently faced in neuro-oncology. [1, 2]

One of the most significant clinical issues following treatment for high grade glioma is injury to the blood-brain barrier; so called pseudoprogression. [3] This results in oedema and apparent deterioration in the CT and MRI images, often associated with increased steroid dose. Brandes et al have highlighted the need for better imaging biomarkers for the problem of pseudo-progression. [1]

There are two potential outcomes of improved medical imaging for conventional imaging techniques currently adopted in the clinic. Firstly, treating clinicians may abandon treatment if imaging shows it to have an apparent negative impact. Secondly, if imaging shows that treatment regimes are not working, clinicians can decide whether or not to continue the current regime given the opportunity cost and risk of the treatment to the patient. As the number of treatment options slowly increase; the impact of this opportunity cost grows. In addition, data from a study by Stupp et al suggests the survival with glioblastoma treated with temozolomide is 14.6 months; [4] allowing clinicians little time to change treatment course.

Further insights into tumour biology could be gained from a better understanding of treatment efficacy. There is a need for better ways to choose patients for treatment, assess when these treatments are working and move onto something else when they are not. As an example, the

benefit of temozolomide has been obscured to some degree by overestimating progression when MRI alone is considered. [5]

Increasingly, radiotherapy (RT) technology, often combined with new drug treatments, is allowing clinicians to treat more aggressively than could have been the case in the past. Modern radiotherapy is a complex chain of procedures. [6] If the tumour is poorly delineated; the risks of geographic misses increase. If the volume of treatment is too large, the toxicity increases and limits the dose that can be achieved.

Clinical trials involving brain tumour imaging face a number of difficulties. Brain tumours are relatively rare (excluding skin tumours; glioblastoma accounts for about 1% of tumours diagnosed in Australia), often making patient recruitment difficult. There are also difficulties with patient consent, as patients have frequent cognitive problems; especially when it comes to salvage treatment regimes. Imaging trials are difficult to perform - requiring blinded comparison of current best practice methods with and without the new modality. Finally, patients may require multiple imaging sessions on separate machines, requiring a significant amount of organisation. With each significant advance in imaging the studies may have to be repeated. It is no surprise that progress has been slow.

The Cancer Genome Atlas has shown us a vast amount about the complex biology of brain tumours such as glioblastoma. We are gaining “a window into its mutational landscape”. [7] However, the more we know, the more complex the situation appears. U87, a commonly studied glioma cell line, has at least 512 homozygously mutated genes. [7] Targeted therapies are a last bearing fruit; as evidenced by CML treatment with imatinib or non-small cell lung cancer with EGFR inhibitors. We are moving from a single drug to the era of the rationally designed therapeutic regimen. Traditional response measures used in the chemotherapy era are much less successful when the response is cytostatic – as it is with many targeted agents. [8] The challenge is how to assess and integrate new agents within existing protocols.

2. Current imaging techniques

Computed Tomography (CT) provides good delineation of bony anatomy, can show haemorrhage or hydrocephalus and can be useful in planning resection of lesions. In combination with PET; CT provides both improved anatomical localisation and improved PET images by allowing attenuation correction. Although CT is often the initial imaging modality used, it has a relatively small role in the investigation of brain tumours and has not been shown to be useful in response assessment or to accurately delineate tumour outlines. [9]

In radiation oncology practice; a planning CT is routinely used to define the shape of the patient and to provide an electron density map to allow modelling of the dose deposited by the radiotherapy beam. The tumour itself, however, is usually delineated on a contrast enhanced MRI fused to a non-contrast planning CT. MRI provides much greater anatomic detail of the structures within the brain. Additionally, with increased magnet strength and better design of coils, images obtained have steadily improved. MRI is routinely used for delineation of the tumour both in neurosurgical and radiation oncology practice. [10]

A contrast agent containing gadolinium is usually added to assist with interpretation. While it has been established that MRI can identify oedema and necrosis along with tumour microstructure in great anatomical detail, as well as regions displaying dysfunction of the blood-brain barrier (BBB). However, it is often the case that these regions do not correlate with areas of infiltration or other neoplastic molecular events. [11] This is particularly concerning in low-grade gliomas, where the blood brain barrier may not be disrupted.

Recently the growth in the use of MRI has stimulated a body of research aimed at estimating electron density from the MR data and using this alone for radiotherapy planning. This avoids the necessity for a CT, reducing both treatment costs and patient radiation burden. [12]

A variety of MR sequences are usually obtained – T1/ T2/ DCE and spin echo. The detailed evaluation of a glioma patient usually involves about 45 minutes on the scanner for the patient and would usually be repeated about every 3 months.

Diffusion weighted imaging (DWI), or the more elaborate Diffusion Tensor Imaging (DTI), are developments of MRI based on the diffusion of water molecules. This diffusion occurs preferentially down the axons rather than across cell membranes. DTI post-processing can identify white matter fibre tracts within the brain, allowing ablative therapies (such as radiosurgery) to eloquent areas to be avoided. [10]

Spectroscopy can be useful in certain clinical scenarios, particularly when radionecrosis is suspected post radiotherapy. It is most commonly undertaken using proton nuclei on commercial scanners. [13] Ratios of choline to N-acetyl aspartate (NAA) have been found to correlate with tumour tissue. [13] NAA, associated with mature neuronal cell density, [14] is decreased in tumours, while choline, involved in cell membrane turnover, [14] is increased, leading to an increase in the ratio of choline/NAA. [13] However, Horska makes the point that spectroscopy looks very different depending on the zone of the tumour being examined (necrotic core, actively growing tumour or peritumoural oedema) [13]. In clinical practice, spectroscopy is complicated by the need for large voxels; limiting the resolution and application in small lesions. There have been attempts to incorporate wider field MRS in radiotherapy planning; even using whole brain acquisition. [14] With the limitation of difficulty in the anterior cranial fossa and prolonged scan time; it looks potentially useful. Spectroscopy's clinical role currently is to provide confirmatory information to other modalities.

The issue of response assessment is critical to development in neuro-oncology, from early phase testing of drugs through to community oncology clinics delivering care to patients. The traditional approach to tumour response has been to base assessment on the change in size of the lesion. Volumetric criteria perform poorly in slowly growing lesions such as low grade glioma. [15] Initially, response assessment was based on two dimensional imaging (MacDonald criteria) of high grade lesions but it was appreciated that these criteria did not account for non-enhancing progression and so Response Assessment for Neuro-Oncology (RANO) criteria were developed. [9, 15]

CT and MRI are not sensitive for early malignant transformation – a significant problem with low grade glioma. The usual clinical situation is – has the patient progressed? The MRI is very difficult to interpret for the 3 months post chemo-radiotherapy. Limitations of both CT and

MRI have led to the increased interest and development of functional imaging. [16-18] Combined PET-MR scanners are finding their way into the clinical setting and may become more than the sum of their parts. It is clear to us that the skill set required for development runs across chemistry, medical physics, pharmacology, computational informatics, radiology, nuclear medicine and clinical oncology. [19] This chapter will concentrate on the clinical applications across a number of entities and review where advanced imaging may help us improve treatment.

3. Metabolic imaging of the brain

PET imaging involves injection of a radioactive tracer, which undergoes decay, with positron emission. Positrons collide with electrons, resulting in positron annihilation and the emission two photons at 180 degrees, detected by the scanner’s detector ring. Following the acquisition, detected events are reconstructed into an image. PET scanning can be performed in static or dynamic mode. In static mode, a single PET image is acquired which is a sum of all events detected during the period of the scan. From this, a measure of tumour-to-background contrast can be used to provide both qualitative and quantitative results. In dynamic mode, the scan is divided up into temporal segments, allowing an insight into the pharmacokinetics and temporal path of the tracer, with kinetic modelling offering quantitative measures that can provide insights into tumour biology. Although dynamic imaging has been shown to provide more information, static acquisitions are still considered the norm in the clinical setting largely due to time and data storage constraints.

3.1. PET radiotracers

A large number of radiotracers have been developed for clinical imaging with many investigated for brain tumours. Selection of an appropriate radiotracer is critical for the investigation being performed, as many target a specific molecular process. A list of common radiotracers used for brain tumour imaging is presented in Table 1.

Radiotracer	Abbreviation	Molecular Target
[¹⁸ F]Fluorodeoxyglucose	¹⁸ F-FDG	Glycolysis
6-[¹⁸ F]fluoro-dihydroxy-l-phenylalanine	¹⁸ F-FDOPA	Dopamine / Amino acid transport
[¹⁸ F]fluoromisonidazole	¹⁸ F-FMISO	Hypoxia
[¹¹ C]-L-methionine	¹¹ C-MET	Amino Acid Transport
O-(2-[¹⁸ F]fluoroethyl)-l-tyrosine	¹⁸ F-FET	Amino Acid Transport
3-deoxy-3-[¹⁸ F]fluorothymidine	¹⁸ F-FLT	Thymidine Kinase Enzyme

Table 1. Common radiotracers used for imaging of brain tumours.

It should be remembered that the difficulty in synthesis and half-life of many of these tracers varies greatly. This can limit their availability, especially in areas with a widely dispersed population. The ultimate utility of an imaging molecule depends on a combination of cost, benefit over other modalities, availability and half-life of the tracer.

3.2. ^{18}F -Fluorodeoxyglucose (FDG)

Tumours such as glioma have a predisposition for increased proliferation and metabolism and there is a wealth of literature investigating the clinical applications of FDG,. These include comparison to MRI [20], detection of glioma [21-28], tumour grading [29-36], choice of biopsy site and radiotherapy planning [37-39], detecting tumour progression [40], recurrence and necrosis [41-53], predicting prognosis and survival [54-58], and patient follow up [59]. FDG PET has the advantage that it is available in most centres due to its wide, relatively simple synthesis and low cost of production. However, the literature highlights numerous limitations in application to brain tumours. The brain is an obligate metaboliser of glucose,. This results in a higher tracer uptake in normal brain tissue. While high-grade gliomas such as glioblastoma are still detectable over background uptake, low-grade and slow progressing tumours often show uptake equal to that of surrounding normal tissue. Other pathologies such as inflammatory diseases can cause tracer up regulation, [60] leading to false positive diagnoses. More sensitive scanner technology has improved this to some degree but a number of studies have been performed exploring alternative radiotracers that may be more useful in brain tumour imaging, such as ^{11}C -L-methionine (MET), 6- ^{18}F fluoro-dihydroxy-L-phenylalanine (FDO-PA), O-(2- ^{18}F fluoroethyl)-L-tyrosine (FET) and 3-deoxy-3- ^{18}F fluorothymidine (FLT).

3.3. ^{11}C -L-Methionine (MET)

MET is an amino acid. MET has advantages over FDG due to both its low cortical uptake and high uptake in neoplastic tissues. (i.e better signal to noise ratio) Methionine is involved in protein synthesis, energy production. It is clear that MET is a powerful diagnostic tool for the detection of primary [61-66] and recurrent [42, 52, 62, 66-70] glioma, as well as having some benefit in prediction of survival. [30, 55, 71-74] Numerous investigations have been performed comparing the diagnostic accuracy of FDG and MET, with reported findings clearly showing the superiority of MET in detection [21, 30, 75] and grading [30, 31] of primary glioma. Conflicting reports have been published, however, regarding the benefit of MET in detection of recurrent glioma. [20, 21, 41, 42, 47, 52]

The ability to provide complimentary information to that of MRI is a key strength of PET, assisting in diagnosis and treatment planning where differential diagnosis on MRI alone is difficult. However, while findings have been published showing the superiority of MET over standard MRI protocols for both primary [61] and recurrent [76] glioma, dynamic susceptibility contrast-enhanced magnetic resonance imaging (DSCE MRI) [77] and perfusion MRI [45, 78] have been shown to be superior to MET PET for tumour detection.

Conflicting findings have been reported for the use of MET PET for grading gliomas. Ceyssens et al [74] and Moulin-Romsee et al [79] reported that MET could not differentiate between

glioma grades. However, Singhal et al [30] found that MET could discriminate between high and low-grade gliomas and Miyake et al [31] and Santoni et al [62] found the tracer could discriminate between grades II and IV and grades II, III and IV, respectively.

While the literature does support the integration of MET PET into the clinical setting, there remain two significant caveats for the tracer. The half-life of ^{11}C isotopes such as MET is considerably short at 20.3 minutes, limiting this isotope and its tracers to centres with onsite production capabilities. Finally, similar to FDG, MET accumulates in inflammatory lesions, complicating differential diagnosis.

3.4. 6- [^{18}F]fluoro-dihydroxy-l-phenylalanine (FDOPA)

An alternative to MET is 6- [^{18}F]fluoro-dihydroxy-l-phenylalanine (FDOPA), a tracer that is commonly used for the assessment of Parkinson's Disease and other neurodegenerative diseases. However, a case study was published in 1996 by Heiss et al [80] revealed the tracer's potential application to brain tumour imaging. While investigating a patient with suspected Parkinson's disease, F-DOPA uptake was visualized in a small region of the brain later pathologically confirmed to be a low-grade glioma. This discovery generated significant interest in FDOPA, with the subsequent literature showing that it provides complimentary information to that of MRI, assisting in diagnosis, prognosis and guiding treatment in primary and recurrent glioma patients.

Studies by Ledezma et al [81] and Karunanithi et al [82] have shown the potential for FDOPA to provide information complementary to that of MRI. Ledezma et al showed that using FDOPA PET along with MRI provided a good mechanism for detecting lesions in primary and recurrent glioma. Karunanithi et al compared the two modalities, finding that FDOPA PET was able to detect tumour recurrence in more cases than MRI. Hence, while MRI is currently implemented in the diagnosis and treatment of glioma, these studies support the potential of FDOPA as modality for complimentary assessment to MRI. Specifically, ^{18}F -FDOPA could have utility where the MRI findings are negative in primary/recurrent tumours or inconclusive in recurrent tumours.

A number of studies have been performed comparing the efficacy of FDOPA PET to that of other radiotracers such as FDG, FLT, FET, Ammonia (NH_3) and MET. Investigations performed by Chen et al [50], Tripathi et al [28], Jora et al [27] and Jacob et al [83] all reported that FDOPA PET was more specific and sensitive to detecting primary and recurrent lesions than FDG. Tripathi et al and Jacob et al also showed that FDOPA PET was more sensitive than FLT and ammonia, respectively.

Given the aforementioned drawbacks of MET, Becherer et al [84] investigated FDOPA as a potential alternative, finding that both tracers had similar standard uptake values (SUVs) and almost identical patterns of spatial uptake.

Many factors are considered in determining prognosis of patients including Karnovsky performance status (KPS), age, tumour size, extent of surgery and tumour grade and histology. In addition, Karunanithi et al [85] have shown that numerous indices derived from FDOPA PET were independent predictors of survival, with Dowson et al further corroborating these

findings by showing that the uptake changes in the most treatment resistant region of tumour post-treatment are predictive of survival.

Another potential application of glioma PET imaging currently under investigation is tumour grading. Confirmation of a glioma diagnosis generally involves surgical biopsy or resection followed by a pathological assessment to determine tumour microvasculature. [86] There are some limitations to performing biopsies. Firstly, biopsy is an invasive procedure, sometimes not possible due to the anatomic location of the tumour, or the condition of the patient. Secondly, as biopsy tissue samples often represent only a small part of the entire tumour, there is a chance that the true tumour grade will be underestimated, as different tumour regions may be of a different grade. [74] Finally, surgery of certain regions may involve surrounding normal brain tissue, exacerbating the prognosis by causing additional neurological morbidity or dysfunction. [87] While FDG has been investigated for glioma grading, alternatives to biopsy methods are yet to be widely implemented in the clinical environment.

Numerous investigations have been performed in an attempt to discriminate high-grade gliomas from low-grade using FDOPA. However, reports are conflicting. Studies by Fueger et al [88], Pafundi et al [89] and Nioche et al [90] reported that FDOPA PET could discriminate between high and low-grade primary glioma. However, as part of their study, Chen et al found no significant difference in FDOPA uptake between the two classes. For recurrent glioma, only Nioche et al was able to discriminate between high and low-grade, while neither Fueger et al nor Kratochwil et al could differentiate the grades in their cohorts. While these studies do suggest that FDOPA has the potential to discriminate between tumour grades in newly diagnosed gliomas, further investigation is needed, especially for recurrent tumours.

While there are clear clinical benefits to using FDOPA for the assessment and management of patients with glioma, this tracer comes with its own set of caveats. The main drawback of FDOPA is synthesis. The most widely used method for FDOPA synthesis uses an electrophilic approach, based on a two-step process. [91, 92] A fully automated method for synthesis of FDOPA is now also available. [93, 94] However, there are limitations with this approach, such as the requirement of additional equipment, precautions for dealing with ^{18}F in gaseous form, and the low radiotracer production rate. [95, 96] Recent developments have seen the introduction of nucleophilic approaches. [97, 98] In contrast to electrophilic, nucleophilic pipelines produce ^{18}F in liquid form. In addition, nucleophilic methods for production of FDOPA are similar to that of FDG and thus require less additional equipment than electrophilic-based synthesis. Finally, the recent availability of single-use synthesis cassettes for FDOPA could make it more readily available..

3.5. O-(2- [^{18}F]fluoroethyl)-l-tyrosine (FET)

O-(2- [^{18}F]fluoroethyl)-l-tyrosine (FET) has also been widely investigated for imaging of brain tumours. FET is one of the first amino acid radiotracers that can be synthesised in large amounts for clinical purposes. Unlike MET and FDG, it has been shown in animal models that FET exhibits low uptake in inflammatory lesions.

A number of studies have been performed investigating FET in brain tumour imaging, with findings strongly supporting the use of the tracer for detecting both primary [99-10]⁴ and recurrent [52, 100, 105-108] glioma and showing superiority in detection, [25] treatment planning [24] and biopsy planning [24, 38] to that of FDG PET.

A key strength of FET PET is its apparent ability to differentiate between different grades of glioma. While some studies report that static imaging alone can provide the information necessary to grade tumours in vivo, [109-111] kinetic analysis from dynamic imaging appears to be superior for both primary [102, 110, 112, 113] and recurrent [114] cases. A combination of MRI, DWI and FET PET indices has also been investigated for grading tumours, providing promising results. [115]

In addition, like other radiotracers, FET PET has been investigated as a possible tool for predicting prognosis, [71, 100, 116-120] treatment planning, [121] and monitoring treatment response [122-124] with findings from all studies supporting its use.

Compared to FDG and MET, FET is a relatively new radiotracer, with the first human experiments performed in the late 1990s. It has been shown to be a cost effective tracer for adoption into the clinic, [123, 125] and future studies could see this tracer being brought closer towards clinical translation.

3.6. 3-deoxy-3- [¹⁸F]fluorothymidine (FLT)

3-deoxy-3- [¹⁸F]fluorothymidine (FLT) is another recently developed amino acid radiotracer that has been investigated for imaging of brain tumour proliferation. [126-131] FLT is an analogue of the thymidine nucleoside, altered to prevent it from following the full biochemical pathway of the nucleoside. Once the tracer is diffused into the cells by active transport, it is phosphorylated but not incorporated into the DNA, thus it becomes trapped within the cell.

FLT has been shown to effectively detect both primary [28, 132] and recurrent [22], [49, 127, 130, 133, 134] tumours, with a large body of work supporting it as a prognostic factor in both cases. [101, 135-141]

Further research has investigated FLT PET as a tool for assessing treatment response [142-145] as well as tumour grading. [129, 132, 134] A study by Jeong et al also found that FLT PET uptake in recurrent glioma correlated with the initial grade of the tumour.

However, despite the body of work supporting FLT as a surrogate biomarker for proliferation, [126-129] its use remains controversial [131, 146] and further studies are warranted to fully elucidate the role of FLT in future clinical practice.

3.7. Hypoxia and ¹⁸F-Fluoromisonidazole (FMISO)

Hypoxia has been shown to have a significant effect on treatment outcome for patients diagnosed with glioma. Low oxygen tension correlates with radioresistance, resulting in the development of local recurrence or metastasis. [147-149] It is well known that hypoxic tumour cells are significantly more resistant to radiation than normoxic cells, as the lack of oxygen inhibits permanent DNA damage. In fact, it has been shown that up to three times the radiation

dose is required to kill hypoxic cells. [150] While cell hypoxia is rare in normal tissue, it is a common occurrence in many solid tumours; [41, 42] classified as acute, chronic or anaemic.

Cells become acutely hypoxic when they are temporarily deprived of oxygen due to the closure or disruption of blood vessels, often caused by the hostile nature and abnormalities within the tumour environment. The abrupt changes in vasculature prevent the supply of oxygen and other nutrients to the cells. Upon return of oxygen to the cells, often due to angiogenesis or vasculogenesis, cells become normoxic again. Chronically hypoxic cells on the other hand, are cells that will remain hypoxic until they become necrotic, often due to the tumour outgrowing its blood supply or proliferation pressure forcing cells away from blood vessels. [151] In fact, necrosis is a requirement for the diagnosis of glioblastoma; as the highly proliferative nature of the tumour forces it to outgrow the blood supply in areas, resulting in a necrotic core usually visible from MRI. Anaemic hypoxia results from low oxygen concentrations within the blood supply and can be due to either pathology or the treatment itself.

There are numerous methods available for measuring hypoxia, both in vivo and in vitro, with the polarographic needle electrode regarded at the gold standard. While this method has been used extensively to assess hypoxia, [39, 44, 49-55] there are concerns with this approach. [74] Firstly, since the needle is invasive, not only is it uncomfortable for the patient, but there also exists a risk that it may cause trauma to the sample site, resulting in exacerbation of injury. Secondly, since only a small sample of the tumour is assessed, there is a risk of underestimating the level of hypoxia. Other, non-invasive methods exist for hypoxia assessment, including phosphorescent imaging, [152-154] which exploits the oxygen-depending quenching of phosphorescence, blood oxygen level dependent (BOLD) MRI, which is able to identify hypoxic vasculature using paramagnetic properties of deoxyhaemoglobin, ^{19}F MRI, [155] which utilizes the linear relationship between ^{19}F and tissue oxygen concentration, and near infrared spectroscopy (NIRS), [156-159] which exploits the low absorption of near infrared wavelengths by tissue chromophores.

In addition to the above methods, there is now significant interest in the use of PET radiotracers for assessing tumour hypoxia. Many radiotracers exist for investigating tumour hypoxia [160-170], including ^{18}F -HX4, ^{18}F -FAZA, ^{18}F -EF5, [124]I-IAZA, [123]/[125]I-IAZP, ^{64}Cu -ATSM, ^{18}F -FENI, ^{62}Cu -diacetyl-bis(N^4 -methylthiosemicarbazone), ^{18}F -FETA, ^{18}F -EF1 and [^{18}F]fluoroerythronitroimidazole. However, ^{18}F -Fluoromisonidazole (FMISO) PET is clearly the most widely used, as it provides the best compromise between sensitivity, specificity and invasiveness.

FMISO is a derivative of nitroimidazole, which is metabolized by intracellular nitroreductases at low oxygen concentrations. They form covalent bonds with macromolecules and become biochemically trapped within these hypoxic cells. [171] This has been shown to occur in oxygen concentrations below 10mmHg [70], [71]. Nitroimidazole metabolism relies on active enzymes and thus does not get absorbed by necrotic tissue.

The first to investigate FMISO as a potential hypoxia imaging agent was Valk et al, [172] who elegantly showed that high-grade gliomas were clearly delineable 40 minutes post-injection

of the tracer. Since this pioneering work, many studies have been performed on both humans and animal models, bringing this tracer closer to clinical translation.

Further research has compared FMISO PET to other imaging modalities such as MRI and FDG PET. MR imaging sequences such as FLAIR, T₂ and contrast enhanced T₁ are regularly used in the diagnosis, treatment planning and treatment evaluation of glioma. However, there are limitations with MRI as described above, especially for low-grade glioma, due to the invasive nature of glioma beyond areas of BBB disruption. Hypoxia has been shown to promote neovascularization through the angiogenic cascade and other molecular signalling events, leading to leakage of the blood-brain barrier. [173, 174] Hence, an understanding of the relationship between hypoxia, as assessed by FMISO PET and MR imaging may lead to an improved understanding of tumour physiology.

Studies by Spence et al, [175] Swanson et al [176], Szeto et al [177] and Kawai et al [178] have confirmed that a relationship does in fact exist between the two imaging modalities. Spence et al showed that hypoxic regions delineated using FMISO PET correlated with contrast enhanced T₁ defined regions. This result was verified by Swanson et al, who reported on two important findings in their study on pre- and post- operative glioma patients: regions of hypoxia correlated with MRI defined volumes; and regions of hypoxia were not restricted to inside MRI defined volumes. Further corroboration came from Kawai et al, who found that individual hypoxic volumes significantly correlated with volumes defined on contrast enhanced T₁. Finally, Szeto et al used a mathematical model adopted from Swanson et al to show that regions of hypoxia correlated with quantitative measures of tumour aggressiveness and sphericity.

It is well established that hypoxic cells can increase their glycolysis to maintain energy levels, suggesting that a relationship between hypoxia and glycolysis does exist. However, proliferation is considerably inhibited in chronic hypoxia, significantly reducing glycolysis. [179-181] This suggests that a complex relationship may exist, the understanding of which may lead to a more comprehensive understanding of the tumour microenvironment.

Investigations by Rejendran et al, [182] Hatano et al [183] and Cher et al [184] have confirmed that any relationship that may exist between the two microbiological processes is complex. While Rejendran et al and Hatano et al found little to no correlation between FMISO and FDG Uptake, Cher et al found that a correlation did exist within individual tumour grades. These results suggest that complementary information can be obtained from FDG and FMISO imaging. However, further studies are warranted to fully elucidate any relationship between the two tracers.

Accurate grading of glioma cases is imperative for both patient prognosis and treatment planning, as patients diagnosed with higher-grade glioma fair significantly worse compared those with lower grade. While necrosis, usually visualized with MRI, is a key requirement for diagnosis of glioblastoma, the most widely accepted method for grading is pathological assessment from biopsy. However as described previously, there are concerns with this approach: Biopsy is not always possible due to tumour location; where biopsy is possible, further neurological dysfunction is a risk with such an invasive procedure; and the sample

only represents a small section of the tumour volume, possibly resulting in an underestimation of the tumour grade.

However, studies into FMISO PET have found that tracer uptake correlates very well with tumour grade, offering up a non-invasive method for tumour grading. Cher et al, [184] Hirata et al, [185] Yamamoto et al [186] and Kawai et al [187] all reported a higher uptake of ^{18}F -FMISO in glioblastoma patients than all other grades. While Cher et al and Hirata et al reported no tracer uptake in lower-grade tumours compared to normal tissue, Yamamoto et al and Kawai et al reported higher uptake in grade III than surrounding cortex. These results suggest that, at the very least, FMISO PET could be used to confirm a diagnosis of high-grade (grade III-IV) glioma. However, further studies are needed to conclude distinction between grade III and IV cases.

Probably the most fascinating property of hypoxia is its apparent strong negative correlation with patient prognosis, which has been investigated since the development of the polarographic electrode needle. Cher et al [184] and Hirata et al [185] both reported FMISO PET defined hypoxia as a significant negative prognostic factor, with Cher et al able to predict the site of recurrence in two patients from ^{18}F -FMISO images. Spence et al was also able to correlate FMISO PET images with time-to-tumour-progression (TTP), survival and KPS. Finally, Swanson et al was able to negatively correlate FMISO PET images with patient survival in pre-operative cases of glioblastoma. These studies add to the growing body of evidence that suggests hypoxia is a key factor in treatment success and patient survival.

Finally, Valable et al [188] performed an investigation into hypoxia and the anti-angiogenic treatment sunitinib. Anti-angiogenic treatments were proposed as a possible cancer therapy, as they target VEGF pathways, leading to a decrease in tumour vasculature. However, treatment results were disappointing, with treatment benefits being temporary, leading to a short increase in time to progression at best. However, the concept of vascular normalisation rekindled the idea of anti-angiogenic treatments, whereby therapy improves tumours vasculature, reducing tumour hypoxia and improving the efficacy of radiotherapy. Valable et al showed in a mouse model that tumours treated with sunitinib had a significant decrease in FMISO PET after 7 days, suggesting that the use of anti-angiogenic treatments may be viable to improve tumour oxygenation and therefore radiotherapy treatments.

The body of work surrounding hypoxia strongly suggests that important information can be obtained from F-MISO PET imaging of glioma and, while more needs to be done to fully elucidate tumour biology, it is clear that hypoxia is an important aspect of future treatment for patients diagnosed with glioma.

3.8. Dual-tracer imaging (multiplexing)

The current literature strongly supports the idea that, given the complex and heterogeneous nature of glioma, knowledge of multiple biological characteristics is needed for effective patient treatment. For example, amino acid activity from FDOPA PET, proliferation from FLT PET and hypoxia from FMISO PET. Currently, however, the physical limitations of PET imaging prevents ascertaining information from two radiotracers simultaneously. Thus, to

acquire the necessary biological information, PET acquisitions must be performed on separate days, allowing the first radiotracer to decay.

Unfortunately, however, there are many disadvantages to this approach. Firstly, performing multiple PET acquisitions requires that the patient be prepared multiple times, significantly increases the scanner time and increases the associated costs. Secondly, performing acquisitions on separate days introduces spatial uncertainty in two ways: Tumour evolution between scans can be significant, especially given the aggressive nature of glioma. In addition, registration of the images introduces uncertainty due to change in patient positions and image artefacts. This, however, can be mitigated provided there is enough mutual anatomical information available between the two scans.

With the need to image multiple biological characteristics becoming evident, and the obvious inherent logistical limitations in performing multiple scans, the field of dual-tracer imaging is gaining momentum.

Dual-tracer imaging, or multiplexing, allows for imaging of multiple PET radiotracers within a single scanning session by employing novel post-processing techniques to separate the two tracer signals. This approach trades-off tracer signal fidelity for logistics, circumventing some of these issues encountered when performing multiple acquisitions: The patient need only be prepared for scanning once; the required scanning time is shorter than that of two individual acquisitions; tumour evolution between the two images is negligible; and, provided the patient does not move during the scan, registration is no longer required.

However, this technique has one major logistical limitation, requiring that multiple radiotracers be synthesized and available at the same time. This is a difficult task in most centres, as simultaneous synthesis of two ^{18}F radiotracers, for example, would be impossible without two cyclotrons. This limitation may be mitigated, however, by either using radiotracers with differing isotopes and/or synthesis methods, or using radiotracers with sufficiently long half-lives that allow the two tracers to be synthesized consecutively before application.

Since dual-tracer imaging was first proposed by Huang et al, [189] a number of methods have been proposed and can be loosely grouped by their method for signal separation. Methods that allow both tracers to be injected simultaneously reduce the scanning time but require that both tracers have different half-lives, with signal separation performed by exploiting the difference in tracer decay rates. Such methods include those by Huang et al, [189] who investigated ^{13}N , ^{18}F , ^{68}Ga and ^{64}Cu isotopes in phantoms, Figueiras et al, [190] who investigated FDG and ammonia in both phantoms and an ischemic rat model, and Kadrmas et al, [191] who exploited differing decay rates in conjunction with differing kinetics and distribution for FDG, ATSM and PTSM. Other methods, such as those by Koeppe et al [192] and Kadrmas et al, [193] employ a staggered injection that allows for the discrimination of tracer signals with the same half-life, increasing the selection of available radiotracers that can be used. Both methods rely on kinetic modeling to separate the two signals. In addition, Verhaeghe et al [194] and Ikoma et al [195] proposed methods that exploit differing tracer half-lives in addition to a staggered injection interval and kinetic analysis to separate tracer signals.

The above studies propose methods that require scanning to be initiated at the injection of the first tracer. To reduce the required scanning time further, a study by Bell et al [196] investigated separation of two ^{18}F radiotracers by staggering the injection interval and initiating the PET acquisition once the first tracer had reached steady state. Using this method, scanning time was only increased by 10 to 20 minutes over a standard PET acquisition.

Finally, two studies have been performed investigating performing signal separation at the reconstruction level. Gao et al [197] and Cheng et al [198] incorporate kinetic modelling into the reconstruction process to separate tracer information.

Dual-tracer imaging offers a viable alternative to performing multiple PET acquisitions, with the above methods illustrating its potential in future research practices. However, further studies are required to fully develop the concept of simultaneous multiplexing, with particular attention to circumventing the logistical issue of simultaneous radiotracer synthesis.

4. Clinical applications

Medical imaging is important at many junctures during the course of the treatment path for glioma patients, from the initial diagnosis and grading of the disease to planning and assessing response to therapy. MRI is the gold standard for obtaining morphological information on tumours, but has some limitations as mentioned previously. One role of imaging is in establishing a continually updated prognosis at all stages during treatment. So in addition to detecting and localising tumours, the differentiation of treatment induced pseudo-progression from recurrence is important. The latter is a particular challenge, as is establishing the grade of diffuse tumours and the late identification of recurrence. [199]

Non-structural MR modalities can assist in overcoming the difficulties of differentiating recurrence from post treatment radiation effect. A previous study [200] examined the use of relative cerebral blood volume, and measured a sensitivity of 92% with a specificity of 100%. Relative cerebral blood volume (rCBV) has shown superior abilities in distinguishing recurrence from radiation necrosis than either FDG or MET-PET. [45]

The sensitivity of molecular imaging techniques such as PET can be used to surmount some limitations of MRI, [16] for instance FDG, the most widely used PET tracer, can predict anaplastic transformation in low-grade gliomas. However PET should be used in conjunction with anatomical imaging as greater sensitivity and specificity is demonstrated when PET is combined with MRI. [16] In clinical practice, PET is considered useful for resolving ambiguities between recurrence and treatment induced changes. [201] It is broadly acknowledged that the augmentation of MRI and CT with other modalities such as PET, SPECT and diffusion imaging may assist in differentiating treatment effects from progression, but more research is needed, [202, 203] and efforts have been made to standardize assessments when considering metabolic imaging modalities, at least when using FDG. [204] As mentioned however, although FDG has demonstrated efficacy for many oncologic applications, it is of less utility in cortical regions where healthy tissue has high FDG-avidity preventing the unambiguous differentiation of

normal and neoplastic tissue. FDG retains some utility for glioma applications, for instance in a study of 19 patients changes in glucose metabolism measured using FDG-PET predicted response to temozolomide, but not when combined with radiotherapy. [205]

Amino acid tracers, such as MET and F-DOPA, are more sensitive for imaging recurrent tumours, especially LGG, [16] as amino acid transport is up-regulated in malignant lesions. [206] Some debate exists around the relative efficacy of different amino-acid tracers, with a number of tracers examined in the literature. [52] MET is popular choice due to its relative ease of synthesis [207] MET-PET is a better prognostic predictor and improves inter-observer consistency compared to FDG, [42] although it is noted combining FDG with MET further improves prognostic predictions. However, the aforementioned limitations with the half life of MET have seen FDOPA to be a viable alternative to MET, which is more accessible in the clinic due to its longer half-life. [84]

FDOPA precisely localizes tumours as verified on MRI [81] and can also identify tumour not visible on MRI. FDOPA has also been shown to better identify regions of higher density disease than MRI on a study examining 23 biopsy specimens from 10 patients. [89] When evaluating 28 of 30 patients on anti-angiogenic therapy, the absolute volume based on FDOPA two weeks after initiation of therapy, was the most significant predictor of overall survival. [208] This slightly out-performed MRI, possibly implying that the overall tumour burden is a factor in survival, and that metabolic images potentially provide better estimates of disease burden. Such early yet accurate predictions of treatment response offer the potential to select more appropriate therapeutic strategies.

FLT has also been examined as an alternative to FDG for the purposes of estimating tumour margins, and in a study of 25 patients, FLT demonstrated greater correlation with Ki-67 proliferation index. [127] FLT demonstrated superior performance to MET in the grading and assessment of proliferation in 41 newly diagnosed glioma of different grade, [132] with no complimentary information being observed. In comparison to FDOPA, the kinetic parameters of FLT were more predictive of overall survival in 126 glioma patients treated with Bevacizumab-Irinotecan, [145] although combining the modalities further improved predictive performance. However, for low grade glioma FDOPA offers better visualisation than either FLT or FDG for primary and recurrent low grade glioma. [28]

Considering low grade glioma in particular, FET has demonstrated efficacy. [209] In a retrospective study of 59 patients [210] found that the presence of FET uptake was not predictive of outcome, but in the subset of 30 patients with abnormal uptake and dynamic data, a decrease in FET was significantly predictive of outcome. FET PET has also accurately predicted antiangiogenic treatment failure in 11 patients, predicting treatment failure earlier than MRI alone. [211] However changes in FET was found to only be prognostic in astrocytic tumours and not for oligodendral tumours in a cohort of 83 patients. [212]

However, in addition to assessing glioma at the various stages of treatment, functional imaging has a role to play in the treatment of the disease by supplying metabolic information during the planning of surgery and of radiotherapy. The augmentation of MRI with PET to assist surgical planning, has demonstrated the potential to improve cost effectiveness for two

scenarios of varying disease severity. [213] Recently, a fluorescent biological markers that is tumour specific, 5-Aminolevulinic Acid (5-ALA), is becoming available following its proposal and the improvement in progression free survival in stage III trials. [214] 5-ALA is efficacious not only because tumour remains easily identifiable despite tissue movement, but because anaplastic foci are not always detectable on MRI. However 5-ALA is less sensitive than FET for low grade glioma, [215] although for high grade glioma regions of high FET uptake also fluoresce. Even so there is some motivation for the use of PET during surgical planning even when in-vivo biomarkers are available, especially since the greater tumour extent made visible by FDOPA imaging compared to MRI has been revealed by biopsies to be pathological in almost all cases. [89] The latter result also has implications for radiotherapy planning.

The availability of PET imaging influences the definition of GTV when it is used to augment MR images, which has the benefit of reducing inter-observer variability and better sparing normal tissues, at least for some oncology applications. [216] A further study [121] examined patterns of failure in a cohort of 10 patients, and found that treatment failures are generally in the central region, but that neither FET-PET nor MRI optimally predicted recurrence, and that the two modalities were complementary. However inadequate coverage of GTVs as defined by increased MET-PET activity in 5 of 34 patients were associated with an increased risk of treatment failure. [217] Such contradictions mean that although the potential of PET to assist radiotherapy planning is acknowledged, more research is required before consensus is reached on what strategy should be used to define treatment volumes. [218]

Although increased metabolic activity is useful to establishing target volumes, the margins added for infiltration and patient movement to respectively obtain clinical and planned target volumes are probably sufficient to prevent geographic misses, and focal radio-resistance plays a larger role in recurrence. A recent study [219] showed that the uptake in focal regions was a better predictor of recurrence, than uptake over the tumour as a whole, highlighting the importance of intra-tumour heterogeneity. The importance of heterogeneity was confirmed for FDOPA in a study examining high-grade gliomas, [220] where focal uptake was significantly correlated with survival time while tumour-global uptake was not. At the intra-tumour scale, patterns of FET kinetic parameters have been correlated with histopathology in suspected grade II gliomas, and focal regions of anaplastic change were identified. [103]

Given the large treatment volumes, the fact that treatment failure is typically focal, and that the radio-resistance associated with local hypoxia is a contender for the cause of treatment failures, the imaging of hypoxia in conjunction with MRI could have greater efficacy than imaging using amino-acid tracers. FMISO is a PET tracer that can be used to identify hypoxic regions.

Typically only static PET images are analysed, but PET is intrinsically a dynamic modality and kinetic analysis of the dynamic data contained by PET images can reveal supplementary biological factors that are of clinical relevance. Hence, the kinetic analysis of PET data is of increasing interest, for instance Pyka et al [219] demonstrated that recurrence can be best predicted using parameters derived from dynamic data in cohort of 34 patients, and static uptake had a lower significance for low grade gliomas. The kinetics of PET uptake have been investigated for some time. Kinetic parameters derived from FDOPA demonstrated the

potential to distinguishing between low and high grade gliomas for 37 patients. [221] Ellingson et al [222] have also proposed using parametric response maps to evaluate treatment response, especially for slow growing tumours. FET-PET has been shown to predict recurrence on a cohort of 14 patients with a sensitivity of 83% and a specificity 88% [219], when using the relative slope of FET uptake. Parameters derived from dynamic PET appear to offer advantages over the analysis static PET images for FET, when applied to grading gliomas. [114]

The role of medical imaging varies depending on type and grade of tumour in addition to the clinical application. When planning the surgery of low grade glioma, the intrinsic plasticity of the brain can be established using fMRI and diffusion imaging to better trade-off the resection of tumour tissue with the prevention of damage to eloquent areas. [223] PET has applicability in a wide range of clinical settings including biopsy guidance, estimating the true extent of tumours, establishing a prognosis, planning resections, detecting recurrence and monitoring therapy, [224] but benefits fewer patients for radiotherapy planning. [225] For low grade glioma the most promising PET tracers appear to be the amino acid markers FET and MET. [224] Examples of the complementary nature of PET to MRI have been noted by Berntsson et al, [226] where no correlation between tumour metabolism (MET uptake) and vascularity was found, and Rahm et al [227] where no agreement between FET uptake and anisotropic diffusion occurred. The pattern of MET uptake has also been found to differ from the pattern of neuronal cell loss and membrane proliferation shown by Cho/NAA ratios obtained from MRS and should potentially be taken into account when planning biopsies. [228]

FDOPA was found to be prognostic of low grade gliomas with non-enhancing T2 changes on MRI in a cohort of 93 patients by Rangan et al. [229] FET was also found to be prognostic in 59 patients by Jansen et al, [210] who found a rapid roll-off uptake dynamics to be predictive of shortened progression free survival although lack of uptake was not indicate of indolent disease. The pattern of uptake MRI uptake (circumscribed versus diffuse) augmented with FET PET has been found to significantly predict malignant transformation in 33 patients. [230] FET has also been used to establish whether brain lesions are neo-plastic. [111]

Static FET uptake when normalized to background is useful for predicting the malignant transformations, as is the dynamic parameter time-to-peak, [112] although the repeated PET scans suggested to detect malignant transformation early might not be possible in general clinical practice.

PET has potential utility in radiotherapy planning for recurrent and high-grade glioma. In a study of 39 patients with high-grade glioma MET defined gross tumour volumes exceeded those of MRI defined volumes in 74% of cases. [68] For the re-irradiation of recurrent high-grade glioma, in a cohort of 44 patients, patients planned using metabolic imaging (MET-PET and IMT-SPECT) had a significantly longer survival time than those planned using anatomical images alone. [231] However, Hutterer et al [232] note that the specificity of PET may be limited by passive tracer influx through the disrupted blood brain barrier and non-neoplastic lesions. The utility of PET for surgical planning for high-grade glioma has also been demonstrated, where 66 patients whose tumours were completely resected according to PET images used in planning had a prolonged survival. [233] In terms of response measurement, FET PET uptake has been demonstrated to be better than MRI at predicting the failure of Bevacizumab-

Irotocenin treatment. [234]Galldiks et al [122] also found tumour to background ratio (both mean and max) to be significant and independent predictors of progression free survival and overall survival.

For meningioma, the use of MET-PET was shown to improve inter-observer consistency for delineating margins of neoplasticity compared to MRI/CT alone in a study of 10 patients. [235] Another tracer, DOTATOC has also been used to delineate the extent of intracranial invasion [236] and has more recently been incorporated into radiotherapy planning. [237] Notably there were some masses that only DOTA could adequately identify. Cornelius et al [238] has highlighted the efficacy of PET for planning the surgery of meningioma as compliment to MRI and 5-ALA. [239]

Such observations motivate for the use of PET/MR during the planning of radiotherapy and surgery; initial experiences in the clinic have found few drawbacks for such systems in comparison to standard PET/CT and separate MRI systems. [240-242] While MRI will remain the standard baseline modality, PET has demonstrated applicability for several types of brain tumour. Applications include grading of tumours, detecting conversion to an aggressive phenotype, establishing a prognosis, planning surgery and planning radiotherapy. The enhanced sensitivity and specificity of metabolic imaging could assist in further delaying progression and improving survival time by reducing the risk of geographic misses and by identifying regions of infiltration that are occult to MRI, but only when used in concert with structural MRI. The increased availability and use of PET/MR scanners will be key to realising such ambitions.

5. Future developments in PET

5.1. Macromolecular imaging

Molecular imaging, in particular nuclear imaging, has been integral to the development and growth of the field of nanomedicine [243, 244]. The development of nanomedicines occurs over several stages, from materials development, characterisation and understanding at a molecular level to in vitro testing and eventually into in vivo models in the preclinical environment. Molecular imaging techniques have become the cornerstone of in vivo investigations in nanomedicine development and this, in turn, has led to the development of macromolecular imaging agents. The birth of the field of theranostics [245], therapeutic delivery and diagnostic reporter combined in the same molecule, has highlighted the potential of macromolecular imaging agents, which has translated into the field of neuro-oncology. The shift from a highly specific targeting molecule such as an antibody, fragment antibody, peptide or synthetic polymer to an imaging agent has become achievable with the development of facile and commercially available chemistries for the attachment of chelating ligands specific to radio-metals [246]. Essentially, the library of nuclear imaging agents has expanded to the almost infinite library of specific biomolecules that have been developed over the past 40 years in the biotechnology field. The power of this approach in the imaging of brain tumours is yet to be fully realised in the clinical environment. However, an increasing number of preclinical studies

are beginning to show the potential of the approach [243, 244, 247-257]. The attraction of a macromolecular imaging agent is very similar to the attractions of antibody-based therapeutics. Whereas with a targeted therapeutic the desired effect is increased anti-tumour efficacy with decreased toxicity, the advantage of an imaging agent is greater contrast between diseased and healthy tissue leading to a lower risk of misinterpretation. Additional benefits lie in the half-life of the radionuclides used to label macromolecules. Radiometals commonly used in labelling macromolecules are ^{68}Ga (68 minute half-life), ^{64}Cu (762.06 minute half-life) and ^{89}Zr (4704.6 minute half-life) each offering distinct advantages over the commonly used ^{11}C (half life = 20.3 minutes) and ^{18}F (half-life = 109.8 minutes). Due to the short half-life, a prerequisite for using ^{11}C and ^{18}F tracers is access to a cyclotron and radiochemistry facilities. The enormous expense of installation of these facilities often limits the use of PET based brain tumour imaging. The long half-life of ^{64}Cu and ^{89}Zr allows production of the isotope at one location, shipment overnight and conjugation to the macromolecular imaging agent at the imaging facility without significant loss of tracer activity. Although the short half-life of ^{68}Ga excludes this approach, ^{68}Ga is produced from a generator and not by cyclotron methods, which are comparatively inexpensive and have small footprints, leading to relative ease of installation in the majority of facilities [258].

5.2. Macromolecular theranostics

As the chelators used to conjugate imaging nuclides can also be used to conjugate therapeutics, beta emitting nuclides such as ^{90}Y and ^{177}Lu the use of highly specific labelled macromolecules leads to a unique solution for personalised medicine. Essentially, what you see is what you treat. Imaging with a radiolabelled macromolecule will give the clinician a quantitative map of macromolecule tracer within the body. This can be used to either recommend the patient for treatment with the diagnostic analogue of the tracer, or rule out the patient as a successful candidate and recommend another course of treatment. The potential of this approach can be evidenced with translation of materials and regimes to the clinic. The family of octreotate peptides specific to the somatostatin family of receptors are used with increasing success in the management of neuroendocrine tumours [259-261]. A ^{68}Ga labelled version of the peptide can be used for diagnosis and treatment planning and, if the patient is likely to display a response, a ^{177}Lu labelled version of the peptide is administered as the treatment.

5.3. Molecular imaging switches

Further potential for macromolecular imaging lies in switchable imaging agents. Hardware developments in MRI have led to the possibility of multinuclear MRI. ^{19}F (not to be confused with ^{18}F) MRI shows great promise as a molecular imaging modality and is being investigated for a number of applications [262-269]. The gyromagnetic ratio of ^{19}F is very close to that of ^1H ($^{19}\text{F} = 40.052$, $^1\text{H} = 42.576$), the natural abundance of ^{19}F is 100 % and the only endogenous ^{19}F in the body is in the bones and the teeth. Essentially this means that ^{19}F can be imaged on most clinical MRI scanners with the addition of a ^{19}F radiofrequency coil, providing a higher resolution molecular imaging modality. The use of ^{19}F MRI is most predominant in cell tracking and many cell types have been labelled with highly fluorinated polyether emulsions. Both

preclinical [269] and clinical [269, 270] trials have shown that imaging labelled cells in vivo is feasible at sub millimetre resolution, enough to visualise $10^3 - 10^5$ cells. [269] ^{19}F MRI however comes with an additional bonus: the signal can be switched on and off in response to a biological signal. There have been several reports of ^{19}F containing molecules that can be imaged in vivo and are sensitive to pH changes [271-275], temperature [276] and enzyme activity [277]. This is largely achieved through changes to the T_2 relaxation rate induced by a physical change to the environment of the polymer. The potential of this approach is high as, in theory, the same mechanism could be used to monitor the release of a drug from a nano-medicine. In this way, and with multimodal imaging platforms such as PET/MRI, key questions surrounding the efficacy of targeted therapeutics, 'how much reaches the target?', 'how fast is it cleared from the target?', 'is the drug delivered to the target?' and 'what is the response to the therapy?' can be answered by molecular imaging.

6. Conclusions

CT and MR imaging are currently considered the gold standard when treating patients diagnosed with a brain tumour but these techniques are much harder to interpret after treatment has commenced. Developing new therapies and improving care in the clinic will depend on overcoming these shortcomings. The introduction of PET offers the ability to gain complimentary information to that of MR and CT imaging, with many radiotracers developed for investigating tumour microbiology. While each radiotracer has its own set of caveats, there are continuing advances in tracer development and synthesis.

Hypoxia has been shown to significantly affect treatment outcomes, and although many radiotracers have been developed for imaging hypoxia, FMISO continues to be the most applicable. In addition, hypoxia has been shown to provide information on patient prognosis, tumour grading and response to therapy. Multiplexing PET imaging may provide further advances in treatment regimes, offering the ability to image with multiple radiotracers within a single acquisition, although further investigations will need to be performed to bring the technique towards clinical translation.

PET imaging has shown application far beyond the detection of tumours, including tumour grading, detecting transformation to a more aggressive phenotype, predicting outcome, planning surgery and radiotherapy. It's application to surgery and radiotherapy planning offers the potential to improve treatment by reducing the risk of geographic misses and by identifying regions of infiltration that are not seen on MRI.

Future developments in PET technology as well as radiopharmaceutical developments should see the wide adoption of theranostics, providing qualitative and quantitative measures of therapeutic efficacy. With the incorporation of molecular imaging switches in conjunction with newly developed PET/MR technology, the ability to answer vital questions about therapeutic delivery to the tumour can be answered, significantly advancing treatment development for the treatment of glioma.

Acronyms

PET - Positron emission tomography

MRI - Magnetic resonance imaging

FET - [18F]-fluoroethyl-L-tyrosine

FDG - [18F]-Fluorodeoxyglucose

FLT - [18F]fluorothymidine

LGG - Low grade glioma

FDOPA - [18F]fluoro-L-dihydroxyphenylalanine

MET - Methyl- [11C]-l-methionine

GTV - Gross tumour volume

IMT - 123I- α -methyl-tyrosine

5-ALA - 5-Aminolevulinic acid

DOTA - 68-Ga-DOTATOC-PET

SUV - Standard uptake value

KPS - Karnofsky performance status

NAA – N-acetyl aspartate

MRS - Magnetic resonance spectroscopy

BOLD MRI - Blood oxygen level dependant magnetic resonance imaging

NIRS - near infrared spectroscopy

FLAIR - Fluid attenuated inversion recovery

MVD – Microvessel density

VEGF - Vascular endothelial growth factor

HIF-1 α - Hypoxia inducible factor-1 α

TTP - Time to tumour progression

DSCE MRI - Dynamic susceptibility contrast-enhanced magnetic resonance imaging

BBB - Blood brain barrier

RANO – Response Assessment for Neuro-Oncology

CT - computed tomography

Author details

Mike Fay^{1,2,3}, Chris Bell⁴, Nick Dowson⁴, Simon Puttick⁵ and Stephen Rose^{4,6}

1 University of Queensland School of Medicine, Brisbane, Australia

2 Genesis Cancer Care NSW, Newcastle, Australia

3 Royal Brisbane and Women's Hospital, Queensland Health, Brisbane, Australia

4 The Australian e-Health Research Centre, CSIRO Digital Productivity & Services Flagship, Royal Brisbane and Women's Hospital, Brisbane, Australia

5 AIBN, University of Queensland, Brisbane, Australia

6 University of Queensland Centre for Clinical Research (UQCCR) Brisbane, Australia

References

- [1] Brandes AA, Tosoni A, Spagnoli F, et al. Disease progression or pseudoprogression after concomitant radiochemotherapy treatment: pitfalls in neurooncology. *Neuro Oncol* 2008;10:361-367.
- [2] Curran SD, Muellner AU, Schwartz LH. Imaging response assessment in oncology. *Cancer Imaging* 2006;6:S126-130.
- [3] Holdhoff M, Grossman SA. Controversies in the adjuvant therapy of high-grade gliomas. *Oncologist* 2011;16:351-358.
- [4] Stupp R, Mason WP, van den Bent MJ, et al. Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma. *N Engl J Med* 2005;352:987-996.
- [5] Brandes AA, Franceschi E, Tosoni A, et al. MGMT promoter methylation status can predict the incidence and outcome of pseudoprogression after concomitant radiochemotherapy in newly diagnosed glioblastoma patients. *J Clin Oncol* 2008;26:2192-2197.
- [6] Peters LJ, O'Sullivan B, Giralt J, et al. Critical impact of radiotherapy protocol compliance and quality in the treatment of advanced head and neck cancer: results from TROG 02.02. *J Clin Oncol* 2010;28:2996-3001.
- [7] Cloughesy TF, Mischel PS. New strategies in the molecular targeting of glioblastoma: how do you hit a moving target? *Clin Cancer Res* 2011;17:6-11.
- [8] Matthews PM, Rabiner EA, Passchier J, Gunn RN. Positron emission tomography molecular imaging for drug development. *Br J Clin Pharmacol* 2012;73:175-186.

- [9] Ahmed R, Oborski MJ, Hwang M, Lieberman FS, Mountz JM. Malignant gliomas: current perspectives in diagnosis, treatment, and early response assessment using advanced quantitative imaging methods. *Cancer Manag Res* 2014;6:149-170.
- [10] Sundgren PC, Dong Q, Gomez-Hassan D, Mukherji SK, Maly P, Welsh R. Diffusion tensor imaging of the brain: review of clinical applications. *Neuroradiology* 2004;46:339-350.
- [11] Chan JL, Lee SW, Fraass BA, et al. Survival and failure patterns of high-grade gliomas after three-dimensional conformal radiotherapy. *J Clin Oncol* 2002;20:1635-1642.
- [12] Prabhakar R, Julka PK, Ganesh T, Munshi A, Joshi RC, Rath GK. Feasibility of using MRI alone for 3D radiation treatment planning in brain tumors. *Jpn J Clin Oncol* 2007;37:405-411.
- [13] Horska A, Barker PB. Imaging of brain tumors: MR spectroscopy and metabolic imaging. *Neuroimaging Clin N Am* 2010;20:293-310.
- [14] Parra NA, Maudsley AA, Gupta RK, et al. Volumetric Spectroscopic Imaging of Glioblastoma Multiforme Radiation Treatment Volumes. *Int J Radiat Oncol Biol Phys* 2014.
- [15] van den Bent MJ, Wefel JS, Schiff D, et al. Response assessment in neuro-oncology (a report of the RANO group): assessment of outcome in trials of diffuse low-grade gliomas. *Lancet Oncol* 2011;12:583-593.
- [16] Chen W. Clinical applications of PET in brain tumors. *J Nucl Med* 2007;48:1468-1481.
- [17] Schiepers C, Dahlbom M, Chen W, et al. Kinetics of 3'-deoxy-3'-¹⁸F-fluorothymidine during treatment monitoring of recurrent high-grade glioma. *J Nucl Med* 2010;51:720-727.
- [18] Timmers HJ, Chen CC, Carrasquillo JA, et al. Comparison of ¹⁸F-fluoro-L-DOPA, ¹⁸F-fluoro-deoxyglucose, and ¹⁸F-fluorodopamine PET and ¹²³I-MIBG scintigraphy in the localization of pheochromocytoma and paraganglioma. *J Clin Endocrinol Metab* 2009;94:4757-4767.
- [19] Fay MF, Martin JH, Rose S. New imaging techniques for more effective treatment in glioblastoma. *Internal Medicine Journal* 2014;44:5-6.
- [20] Wang X, Hu X, Xie P, Li W, Li X, Ma L. Comparison of magnetic resonance spectroscopy and positron emission tomography in detection of tumor recurrence in post-treatment of glioma: A diagnostic meta-analysis. *Asia Pac J Clin Oncol* 2014.
- [21] Li DL, Xu YK, Wang QS, Wu HB, Li HS. (1)(1)¹C-methionine and (1)(8)¹⁸F-fluorodeoxyglucose positron emission tomography/CT in the evaluation of patients with suspected primary and residual/recurrent gliomas. *Chin Med J (Engl)* 2012;125:91-96.

- [22] Spence AM, Muzi M, Mankoff DA, et al. 18F-FDG PET of gliomas at delayed intervals: improved distinction between tumor and normal gray matter. *J Nucl Med* 2004;45:1653-1659.
- [23] Prieto E, Marti-Climent JM, Dominguez-Prado I, et al. Voxel-based analysis of dual-time-point 18F-FDG PET images for brain tumor identification and delineation. *J Nucl Med* 2011;52:865-872.
- [24] Pauleit D, Stoffels G, Bachofner A, et al. Comparison of (18)F-FET and (18)F-FDG PET in brain tumors. *Nucl Med Biol* 2009;36:779-787.
- [25] Lau EW, Drummond KJ, Ware RE, et al. Comparative PET study using F-18 FET and F-18 FDG for the evaluation of patients with suspected brain tumour. *J Clin Neurosci* 2010;17:43-49.
- [26] Das K, Mittal BR, Vasistha RK, Singh P, Mathuriya SN. Role of (18)F-fluorodeoxyglucose Positron Emission Tomography scan in differentiating enhancing brain tumors. *Indian J Nucl Med* 2011;26:171-176.
- [27] Jora C, Mattakarottu JJ, Aniruddha PG, et al. Comparative evaluation of 18F-FDOPA, 13N-AMMONIA, 18F-FDG PET/CT and MRI in primary brain tumors - A pilot study. *Indian J Nucl Med* 2011;26:78-81.
- [28] Tripathi M, Sharma R, D'Souza M, et al. Comparative evaluation of F-18 FDOPA, F-18 FDG, and F-18 FLT-PET/CT for metabolic imaging of low grade gliomas. *Clin Nucl Med* 2009;34:878-883.
- [29] Yoon JH, Kim JH, Kang WJ, et al. Grading of cerebral glioma with multiparametric MR imaging and 18F-FDG-PET: concordance and accuracy. *Eur Radiol* 2014;24:380-389.
- [30] Singhal T, Narayanan TK, Jacobs MP, Bal C, Mantil JC. 11C-methionine PET for grading and prognostication in gliomas: a comparison study with 18F-FDG PET and contrast enhancement on MRI. *J Nucl Med* 2012;53:1709-1715.
- [31] Miyake K, Shinomiya A, Okada M, Hatakeyama T, Kawai N, Tamiya T. Usefulness of FDG, MET and FLT-PET studies for the management of human gliomas. *J Biomed Biotechnol* 2012;2012:205818.
- [32] Tsuchida T, Takeuchi H, Okazawa H, Tsujikawa T, Fujibayashi Y. Grading of brain glioma with 1-11C-acetate PET: comparison with 18F-FDG PET. *Nucl Med Biol* 2008;35:171-176.
- [33] Derlon JM, Chapon F, Noel MH, et al. Non-invasive grading of oligodendrogliomas: correlation between in vivo metabolic pattern and histopathology. *Eur J Nucl Med* 2000;27:778-787.

- [34] Kim CK, Alavi JB, Alavi A, Reivich M. New grading system of cerebral gliomas using positron emission tomography with F-18 fluorodeoxyglucose. *J Neurooncol* 1991;10:85-91.
- [35] Mertens K, Acou M, Van Hauwe J, et al. Validation of 18F-FDG PET at conventional and delayed intervals for the discrimination of high-grade from low-grade gliomas: a stereotactic PET and MRI study. *Clin Nucl Med* 2013;38:495-500.
- [36] Shi X, Liu Y, Zhang X, et al. The comparison of 13N-ammonia and 18F-FDG in the evaluation of untreated gliomas. *Clin Nucl Med* 2013;38:522-526.
- [37] Lammering G, De Ruyscher D, van Baardwijk A, et al. The use of FDG-PET to target tumors by radiotherapy. *Strahlenther Onkol* 2010;186:471-481.
- [38] Plotkin M, Blechschmidt C, Auf G, et al. Comparison of F-18 FET-PET with F-18 FDG-PET for biopsy planning of non-contrast-enhancing gliomas. *Eur Radiol* 2010;20:2496-2502.
- [39] Pirotte B, Goldman S, Massager N, et al. Comparison of 18F-FDG and 11C-methionine for PET-guided stereotactic brain biopsy of gliomas. *J Nucl Med* 2004;45:1293-1298.
- [40] Imani F, Boada FE, Lieberman FS, Davis DK, Mountz JM. Molecular and metabolic pattern classification for detection of brain glioma progression. *Eur J Radiol* 2014;83:e100-105.
- [41] Takenaka S, Asano Y, Shinoda J, et al. Comparison of (11)C-methionine, (11)C-choline, and (18)F-fluorodeoxyglucose-PET for distinguishing glioma recurrence from radiation necrosis. *Neurol Med Chir (Tokyo)* 2014;54:280-289.
- [42] Van Laere K, Ceyssens S, Van Calenbergh F, et al. Direct comparison of 18F-FDG and 11C-methionine PET in suspected recurrence of glioma: sensitivity, inter-observer variability and prognostic value. *Eur J Nucl Med Mol Imaging* 2005;32:39-51.
- [43] Ozsunar Y, Mullins ME, Kwong K, et al. Glioma recurrence versus radiation necrosis? A pilot comparison of arterial spin-labeled, dynamic susceptibility contrast enhanced MRI, and FDG-PET imaging. *Acad Radiol* 2010;17:282-290.
- [44] Gomez-Rio M, Rodriguez-Fernandez A, Ramos-Font C, Lopez-Ramirez E, Llamas-Elvira JM. Diagnostic accuracy of 201Thallium-SPECT and 18F-FDG-PET in the clinical assessment of glioma recurrence. *Eur J Nucl Med Mol Imaging* 2008;35:966-975.
- [45] Kim YH, Oh SW, Lim YJ, et al. Differentiating radiation necrosis from tumor recurrence in high-grade gliomas: assessing the efficacy of 18F-FDG PET, 11C-methionine PET and perfusion MRI. *Clin Neurol Neurosurg* 2010;112:758-765.
- [46] Santra A, Kumar R, Sharma P, et al. F-18 FDG PET-CT in patients with recurrent glioma: comparison with contrast enhanced MRI. *Eur J Radiol* 2012;81:508-513.

- [47] Tripathi M, Sharma R, Varshney R, et al. Comparison of F-18 FDG and C-11 methionine PET/CT for the evaluation of recurrent primary brain tumors. *Clin Nucl Med* 2012;37:158-163.
- [48] Santra A, Kumar R, Sharma P, Bal C, Julka PK, Malhotra A. Detection of recurrence in glioma: a comparative prospective study between Tc-99m GHA SPECT and F-18 FDG PET/CT. *Clin Nucl Med* 2011;36:650-655.
- [49] Enslow MS, Zollinger LV, Morton KA, et al. Comparison of 18F-fluorodeoxyglucose and 18F-fluorothymidine PET in differentiating radiation necrosis from recurrent glioma. *Clin Nucl Med* 2012;37:854-861.
- [50] Chen W, Silverman DH, Delaloye S, et al. 18F-FDOPA PET imaging of brain tumors: comparison study with 18F-FDG PET and evaluation of diagnostic accuracy. *J Nucl Med* 2006;47:904-911.
- [51] Karunanithi S, Sharma P, Kumar A, et al. 18F-FDOPA PET/CT for detection of recurrence in patients with glioma: prospective comparison with 18F-FDG PET/CT. *Eur J Nucl Med Mol Imaging* 2013;40:1025-1035.
- [52] Nihashi T, Dahabreh IJ, Terasawa T. Diagnostic accuracy of PET for recurrent glioma diagnosis: a meta-analysis. *AJNR Am J Neuroradiol* 2013;34:944-950, S941-911.
- [53] Santra A, Kumar R, Sharma P, Bal C, Julka PK, Malhotra A. F-18 FDG PET-CT for predicting survival in patients with recurrent glioma: a prospective study. *Neuroradiology* 2011;53:1017-1024.
- [54] De Witte O, Lefranc F, Levivier M, Salmon I, Brothi J, Goldman S. FDG-PET as a prognostic factor in high-grade astrocytoma. *J Neurooncol* 2000;49:157-163.
- [55] Kim S, Chung JK, Im SH, et al. 11C-methionine PET as a prognostic marker in patients with glioma: comparison with 18F-FDG PET. *Eur J Nucl Med Mol Imaging* 2005;32:52-59.
- [56] Pardo FS, Aronen HJ, Fitzek M, et al. Correlation of FDG-PET interpretation with survival in a cohort of glioma patients. *Anticancer Res* 2004;24:2359-2365.
- [57] Colavolpe C, Chinot O, Metellus P, et al. FDG-PET predicts survival in recurrent high-grade gliomas treated with bevacizumab and irinotecan. *Neuro Oncol* 2012;14:649-657.
- [58] Padma MV, Said S, Jacobs M, et al. Prediction of pathology and survival by FDG PET in gliomas. *J Neurooncol* 2003;64:227-237.
- [59] Potzi C, Becherer A, Marosi C, et al. [11C] methionine and [18F] fluorodeoxyglucose PET in the follow-up of glioblastoma multiforme. *J Neurooncol* 2007;84:305-314.
- [60] van Waarde A, Cobben DC, Suurmeijer AJ, et al. Selectivity of 18F-FLT and 18F-FDG for differentiating tumor from inflammation in a rodent model. *J Nucl Med* 2004;45:695-700.

- [61] Galldiks N, Kracht LW, Dunkl V, et al. Imaging of non- or very subtle contrast-enhancing malignant gliomas with [(1)(1)C]-methionine positron emission tomography. *Mol Imaging* 2011;10:453-459.
- [62] Santoni M, Nanni C, Bittoni A, et al. [(11) C]-methionine positron emission tomography in the postoperative imaging and followup of patients with primary and recurrent gliomas. *ISRN Oncol* 2014;2014:463152.
- [63] Kracht LW, Miletic H, Busch S, et al. Delineation of brain tumor extent with [11C]L-methionine positron emission tomography: local comparison with stereotactic histopathology. *Clin Cancer Res* 2004;10:7163-7170.
- [64] Okada Y, Nihashi T, Fujii M, et al. Differentiation of newly diagnosed glioblastoma multiforme and intracranial diffuse large B-cell Lymphoma using (11)C-methionine and (18)F-FDG PET. *Clin Nucl Med* 2012;37:843-849.
- [65] Gumprecht H, Grosu AL, Souvatsoglou M, Dzewas B, Weber WA, Lumenta CB. 11C-Methionine positron emission tomography for preoperative evaluation of suggestive low-grade gliomas. *Zentralbl Neurochir* 2007;68:19-23.
- [66] Yamane T, Sakamoto S, Senda M. Clinical impact of (11)C-methionine PET on expected management of patients with brain neoplasm. *Eur J Nucl Med Mol Imaging* 2010;37:685-690.
- [67] Terakawa Y, Tsuyuguchi N, Iwai Y, et al. Diagnostic accuracy of 11C-methionine PET for differentiation of recurrent brain tumors from radiation necrosis after radiotherapy. *J Nucl Med* 2008;49:694-699.
- [68] Grosu AL, Weber WA, Riedel E, et al. L-(methyl-11C) methionine positron emission tomography for target delineation in resected high-grade gliomas before radiotherapy. *Int J Radiat Oncol Biol Phys* 2005;63:64-74.
- [69] Tsuyuguchi N, Takami T, Sunada I, et al. Methionine positron emission tomography for differentiation of recurrent brain tumor and radiation necrosis after stereotactic radiosurgery--in malignant glioma. *Ann Nucl Med* 2004;18:291-296.
- [70] Ullrich RT, Kracht L, Brunn A, et al. Methyl-L-11C-methionine PET as a diagnostic marker for malignant progression in patients with glioma. *J Nucl Med* 2009;50:1962-1968.
- [71] Galldiks N, Dunkl V, Kracht LW, et al. Volumetry of [(1)(1)C]-methionine positron emission tomographic uptake as a prognostic marker before treatment of patients with malignant glioma. *Mol Imaging* 2012;11:516-527.
- [72] Smits A, Westerberg E, Ribom D. Adding 11C-methionine PET to the EORTC prognostic factors in grade 2 gliomas. *Eur J Nucl Med Mol Imaging* 2008;35:65-71.

- [73] Nuutinen J, Sonninen P, Lehtikainen P, et al. Radiotherapy treatment planning and long-term follow-up with [(11)C]methionine PET in patients with low-grade astrocytoma. *Int J Radiat Oncol Biol Phys* 2000;48:43-52.
- [74] Ceyssens S, Van Laere K, de Groot T, Goffin J, Bormans G, Mortelmans L. [11C]methionine PET, histopathology, and survival in primary brain tumors and recurrence. *AJNR Am J Neuroradiol* 2006;27:1432-1437.
- [75] Yamamoto Y, Nishiyama Y, Kimura N, et al. 11C-acetate PET in the evaluation of brain glioma: comparison with 11C-methionine and 18F-FDG-PET. *Mol Imaging Biol* 2008;10:281-287.
- [76] Galldiks N, Ullrich R, Schroeter M, Fink GR, Jacobs AH, Kracht LW. Volumetry of [(11)C]-methionine PET uptake and MRI contrast enhancement in patients with recurrent glioblastoma multiforme. *Eur J Nucl Med Mol Imaging* 2010;37:84-92.
- [77] Deng SM, Zhang B, Wu YW, Zhang W, Chen YY. Detection of glioma recurrence by (1)(1)C-methionine positron emission tomography and dynamic susceptibility contrast-enhanced magnetic resonance imaging: a meta-analysis. *Nucl Med Commun* 2013;34:758-766.
- [78] Dandois V, Rommel D, Renard L, Jamart J, Cosnard G. Substitution of 11C-methionine PET by perfusion MRI during the follow-up of treated high-grade gliomas: preliminary results in clinical practice. *J Neuroradiol* 2010;37:89-97.
- [79] Moulin-Romsee G, D'Hondt E, de Groot T, et al. Non-invasive grading of brain tumours using dynamic amino acid PET imaging: does it work for 11C-methionine? *Eur J Nucl Med Mol Imaging* 2007;34:2082-2087.
- [80] Heiss WD, Wienhard K, Wagner R, et al. F-Dopa as an amino acid tracer to detect brain tumors. *J Nucl Med* 1996;37:1180-1182.
- [81] Ledezma CJ, Chen W, Sai V, et al. 18F-FDOPA PET/MRI fusion in patients with primary/recurrent gliomas: initial experience. *Eur J Radiol* 2009;71:242-248.
- [82] Karunanithi S, Sharma P, Kumar A, et al. Comparative diagnostic accuracy of contrast-enhanced MRI and (18)F-FDOPA PET-CT in recurrent glioma. *Eur Radiol* 2013;23:2628-2635.
- [83] Jacob MJ, Pandit AG, Jora C, Mudalsha R, Sharma A, Pathak HC. Comparative study of (18)F-DOPA, (13)N-Ammonia and F18-FDG PET/CT in primary brain tumors. *Indian J Nucl Med* 2011;26:139-143.
- [84] Becherer A, Karanikas G, Szabo M, et al. Brain tumour imaging with PET: a comparison between [18F]fluorodopa and [11C]methionine. *Eur J Nucl Med Mol Imaging* 2003;30:1561-1567.
- [85] Karunanithi S, Sharma P, Kumar A, et al. Can F-FDOPA PET/CT predict survival in patients with suspected recurrent glioma? A prospective study. *Eur J Radiol* 2013.

- [86] Behin A, Hoang-Xuan K, Carpentier AF, Delattre JY. Primary brain tumours in adults. *Lancet* 2003;361:323-331.
- [87] Spetzler RF, Martin NA. A proposed grading system for arteriovenous malformations. *J Neurosurg* 1986;65:476-483.
- [88] Fueger BJ, Czernin J, Cloughesy T, et al. Correlation of 6-18F-fluoro-L-dopa PET uptake with proliferation and tumor grade in newly diagnosed and recurrent gliomas. *J Nucl Med* 2010;51:1532-1538.
- [89] Pafundi DH, Laack NN, Youland RS, et al. Biopsy validation of 18F-DOPA PET and biodistribution in gliomas for neurosurgical planning and radiotherapy target delineation: results of a prospective pilot study. *Neuro Oncol* 2013;15:1058-1067.
- [90] Nioche C, Soret M, Gontier E, et al. Evaluation of quantitative criteria for glioma grading with static and dynamic 18F-FDopa PET/CT. *Clin Nucl Med* 2013;38:81-87.
- [91] Dolle F, Demphel S. 6- [18F]Fluoro-L-DOPA by radiofluorodestannylation: a short and simple synthesis of a new labelling precursor. *Journal of Labelled Compounds and Radiopharmaceuticals* 1998;40:105-112.
- [92] Namavari M, Bishop A, Satyamurthy N, Bida G, Barrio JR. Regioselective radiofluorodestannylation with [18F]F₂ and [18F]CH₃COOF: a high yield synthesis of 6- [18F]Fluoro-L-dopa. *Int J Rad Appl Instrum A* 1992;43:989-996.
- [93] Kao C-HK, Hsu W-L, Xie H-L, Lin M-C, Lan W-C, Chao H-Y. GMP production of [18F]FDOPA and issues concerning its quality analyses as in USP "Fluorodopa F 18 Injection". *Annals of nuclear medicine* 2011;25:309-316.
- [94] Vries EFJD, Luurtsema G, Bru M, Elsinga PH, Vaalburg W. Fully automated synthesis module for the high yield one- pot preparation of 6- [18 F] Fluoro- L -DOPA. *Journal of Labelled Compounds and Radiopharmaceuticals* 1999;41:1-10.
- [95] Behnam Azad B, Chirakal R, Schrobilgen GJ. Trifluoromethanesulfonic acid, an alternative solvent medium for the direct electrophilic fluorination of DOPA: new syntheses of 6- [18F]fluoro-L-DOPA and 6- [18F]fluoro-D-DOPA. *Journal of Labelled Compounds and Radiopharmaceuticals* 2007;50:1236-1242.
- [96] Forsback S, Eskola O, Haaparanta M, Bergmann Jr, Solin O. Electrophilic synthesis of 6- [18F]fluoro-L-DOPA using post-target produced [18F]F₂. *Radiochimica Acta* 2008;96:845-848.
- [97] Lemaire C, Gillet S, Guillouet Sp, Plenevaux A, Aerts JL, Luxen A. Highly Enantioselective Synthesis of No-Carrier-Added 6- [18F]Fluoro-L-dopa by Chiral Phase-Transfer Alkylation. *European Journal of Organic Chemistry* 2004;2004:2899-2904.
- [98] Wagner FM, Ermert J, Coenen HH. Three-step, "one-pot" radiosynthesis of 6-fluoro-3,4-dihydroxy-L-phenylalanine by isotopic exchange. *Journal of nuclear medicine : official publication, Society of Nuclear Medicine* 2009;50:1724-1729.

- [99] Benouaich-Amiel A, Lubrano V, Tafani M, et al. Evaluation of O-(2- [18F]-Fluoroethyl)-L-Tyrosine in the Diagnosis of Glioblastoma. *Arch Neurol* 2010;67:370-372.
- [100] Floeth FW, Pauleit D, Wittsack HJ, et al. Multimodal metabolic imaging of cerebral gliomas: positron emission tomography with [18F]fluoroethyl-L-tyrosine and magnetic resonance spectroscopy. *J Neurosurg* 2005;102:318-327.
- [101] Jeong SY, Lim SM. Comparison of 3'-deoxy-3'- [18F]fluorothymidine PET and O-(2- [18F]fluoroethyl)-L-tyrosine PET in patients with newly diagnosed glioma. *Nucl Med Biol* 2012;39:977-981.
- [102] Kratochwil C, Combs SE, Leotta K, et al. Intra-individual comparison of 18F-FET and 18F-DOPA in PET imaging of recurrent brain tumors. *Neuro-oncology* 2013;1-7.
- [103] Kunz M, Thon N, Eigenbrod S, et al. Hot spots in dynamic (18)FET-PET delineate malignant tumor parts within suspected WHO grade II gliomas. *Neuro Oncol* 2011;13:307-316.
- [104] Pauleit D, Floeth F, Hamacher K, et al. O-(2- [18F]fluoroethyl)-L-tyrosine PET combined with MRI improves the diagnostic assessment of cerebral gliomas. *Brain* 2005;128:678-687.
- [105] Grosu AL, Astner ST, Riedel E, et al. An interindividual comparison of O-(2- [18F]fluoroethyl)-L-tyrosine (FET)- and L- [methyl-11C]methionine (MET)-PET in patients with brain gliomas and metastases. *Int J Radiat Oncol Biol Phys* 2011;81:1049-1058.
- [106] Mehrkens JH, Popperl G, Rachinger W, et al. The positive predictive value of O-(2- [18F]fluoroethyl)-L-tyrosine (FET) PET in the diagnosis of a glioma recurrence after multimodal treatment. *J Neurooncol* 2008;88:27-35.
- [107] Popperl G, Gotz C, Rachinger W, Gildehaus FJ, Tonn JC, Tatsch K. Value of O-(2- [18F]fluoroethyl)- L-tyrosine PET for the diagnosis of recurrent glioma. *Eur J Nucl Med Mol Imaging* 2004;31:1464-1470.
- [108] Rachinger W, Goetz C, Popperl G, et al. Positron emission tomography with O-(2- [18F]fluoroethyl)-l-tyrosine versus magnetic resonance imaging in the diagnosis of recurrent gliomas. *Neurosurgery* 2005;57:505-511; discussion 505-511.
- [109] Calcagni ML, Galli G, Giordano A, et al. Dynamic O-(2- [18F]fluoroethyl)-L-tyrosine (F-18 FET) PET for glioma grading: assessment of individual probability of malignancy. *Clin Nucl Med* 2011;36:841-847.
- [110] Popperl G, Kreth FW, Mehrkens JH, et al. FET PET for the evaluation of untreated gliomas: correlation of FET uptake and uptake kinetics with tumour grading. *Eur J Nucl Med Mol Imaging* 2007;34:1933-1942.
- [111] Rapp M, Heinzel A, Galldiks N, et al. Diagnostic performance of 18F-FET PET in newly diagnosed cerebral lesions suggestive of glioma. *J Nucl Med* 2013;54:229-235.

- [112] Galldiks N, Stoffels G, Ruge MI, et al. Role of O-(2-18F-fluoroethyl)-L-tyrosine PET as a diagnostic tool for detection of malignant progression in patients with low-grade glioma. *J Nucl Med* 2013;54:2046-2054.
- [113] Jansen NL, Graute V, Armbruster L, et al. MRI-suspected low-grade glioma: is there a need to perform dynamic FET PET? *Eur J Nucl Med Mol Imaging* 2012;39:1021-1029.
- [114] Popperl G, Kreth FW, Herms J, et al. Analysis of 18F-FET PET for grading of recurrent gliomas: is evaluation of uptake kinetics superior to standard methods? *J Nucl Med* 2006;47:393-403.
- [115] Dunet V, Maeder P, Nicod-Lalonde M, et al. Combination of MRI and dynamic FET PET for initial glioma grading. *Nuklearmedizin* 2014;53.
- [116] Niyazi M, Jansen N, Ganswindt U, et al. Re-irradiation in recurrent malignant glioma: prognostic value of [18F]FET-PET. *J Neurooncol* 2012;110:389-395.
- [117] Piroth MD, Holy R, Pinkawa M, et al. Prognostic impact of postoperative, pre-irradiation (18F)-fluoroethyl-L-tyrosine uptake in glioblastoma patients treated with radiochemotherapy. *Radiother Oncol* 2011;99:218-224.
- [118] Piroth MD, Pinkawa M, Holy R, et al. Prognostic value of early [18F]fluoroethyltyrosine positron emission tomography after radiochemotherapy in glioblastoma multiforme. *Int J Radiat Oncol Biol Phys* 2011;80:176-184.
- [119] Sweeney R, Polat B, Samnick S, Reiners C, Flentje M, Verburg FA. O-(2- [(18F)]fluoroethyl)-L-tyrosine uptake is an independent prognostic determinant in patients with glioma referred for radiation therapy. *Ann Nucl Med* 2014;28:154-162.
- [120] Thiele F, Ehmer J, Piroth MD, et al. The quantification of dynamic FET PET imaging and correlation with the clinical outcome in patients with glioblastoma. *Phys Med Biol* 2009;54:5525-5539.
- [121] Weber DC, Casanova N, Zilli T, et al. Recurrence pattern after [(18F)]fluoroethyltyrosine-positron emission tomography-guided radiotherapy for high-grade glioma: a prospective study. *Radiother Oncol* 2009;93:586-592.
- [122] Galldiks N, Langen KJ, Holy R, et al. Assessment of treatment response in patients with glioblastoma using O-(2-18F-fluoroethyl)-L-tyrosine PET in comparison to MRI. *J Nucl Med* 2012;53:1048-1057.
- [123] Heinzl A, Muller D, Langen KJ, et al. The use of O-(2-18F-fluoroethyl)-L-tyrosine PET for treatment management of bevacizumab and irinotecan in patients with recurrent high-grade glioma: a cost-effectiveness analysis. *J Nucl Med* 2013;54:1217-1222.

- [124] Popperl G, Goldbrunner R, Gildehaus FJ, et al. O-(2- [18F]fluoroethyl)-L-tyrosine PET for monitoring the effects of convection-enhanced delivery of paclitaxel in patients with recurrent glioblastoma. *Eur J Nucl Med Mol Imaging* 2005;32:1018-1025.
- [125] Heinzl A, Stock S, Langen KJ, Muller D. Cost-effectiveness analysis of FET PET-guided target selection for the diagnosis of gliomas. *Eur J Nucl Med Mol Imaging* 2012;39:1089-1096.
- [126] Backes H, Ullrich R, Neumaier B, Kracht L, Wienhard K, Jacobs AH. Noninvasive quantification of 18F-FLT human brain PET for the assessment of tumour proliferation in patients with high-grade glioma. *Eur J Nucl Med Mol Imaging* 2009;36:1960-1967.
- [127] Chen W, Cloughesy T, Kamdar N, et al. Imaging proliferation in brain tumors with 18F-FLT PET: comparison with 18F-FDG. *J Nucl Med* 2005;46:945-952.
- [128] Price SJ, Fryer TD, Cleij MC, et al. Imaging regional variation of cellular proliferation in gliomas using 3'-deoxy-3'- [18F]fluorothymidine positron-emission tomography: an image-guided biopsy study. *Clin Radiol* 2009;64:52-63.
- [129] Saga T, Kawashima H, Araki N, et al. Evaluation of primary brain tumors with FLT-PET: usefulness and limitations. *Clin Nucl Med* 2006;31:774-780.
- [130] Shishido H, Kawai N, Miyake K, Yamamoto Y, Nishiyama Y, Tamiya T. Diagnostic Value of 11C-Methionine (MET) and 18F-Fluorothymidine (FLT) Positron Emission Tomography in Recurrent High-Grade Gliomas; Differentiation from Treatment-Induced Tissue Necrosis. *Cancers (Basel)* 2012;4:244-256.
- [131] Ullrich R, Backes H, Li H, et al. Glioma proliferation as assessed by 3'-fluoro-3'-deoxy-L-thymidine positron emission tomography in patients with newly diagnosed high-grade glioma. *Clin Cancer Res* 2008;14:2049-2055.
- [132] Hatakeyama T, Kawai N, Nishiyama Y, et al. 11C-methionine (MET) and 18F-fluorothymidine (FLT) PET in patients with newly diagnosed glioma. *Eur J Nucl Med Mol Imaging* 2008;35:2009-2017.
- [133] Spence AM, Muzi M, Link JM, et al. NCI-sponsored trial for the evaluation of safety and preliminary efficacy of 3'-deoxy-3'- [18F]fluorothymidine (FLT) as a marker of proliferation in patients with recurrent gliomas: preliminary efficacy studies. *Mol Imaging Biol* 2009;11:343-355.
- [134] Yamamoto Y, Wong TZ, Turkington TG, Hawk TC, Reardon DA, Coleman RE. 3'-Deoxy-3'- [F-18]fluorothymidine positron emission tomography in patients with recurrent glioblastoma multiforme: comparison with Gd-DTPA enhanced magnetic resonance imaging. *Mol Imaging Biol* 2006;8:340-347.
- [135] Belohlavek O, Fencel P, Majovsky M, Jaruskova M, Benes V. FLT-PET in previously untreated patients with low-grade glioma can predict their overall survival. *Nucl Med Rev Cent East Eur* 2014;17:7-12.

- [136] Chen W, Delaloye S, Silverman DH, et al. Predicting treatment response of malignant gliomas to bevacizumab and irinotecan by imaging proliferation with [18F] fluorothymidine positron emission tomography: a pilot study. *J Clin Oncol* 2007;25:4714-4721.
- [137] Harris RJ, Cloughesy TF, Pope WB, et al. 18F-FDOPA and 18F-FLT positron emission tomography parametric response maps predict response in recurrent malignant gliomas treated with bevacizumab. *Neuro Oncol* 2012;14:1079-1089.
- [138] Idema AJ, Hoffmann AL, Boogaarts HD, et al. 3'-Deoxy-3'-18F-fluorothymidine PET-derived proliferative volume predicts overall survival in high-grade glioma patients. *J Nucl Med* 2012;53:1904-1910.
- [139] Schwarzenberg J, Czernin J, Cloughesy TF, et al. 3'-deoxy-3'-18F-fluorothymidine PET and MRI for early survival predictions in patients with recurrent malignant glioma treated with bevacizumab. *J Nucl Med* 2012;53:29-36.
- [140] Wardak M, Schiepers C, Dahlbom M, et al. Discriminant analysis of (1)(8)F-fluorothymidine kinetic parameters to predict survival in patients with recurrent high-grade glioma. *Clin Cancer Res* 2011;17:6553-6562.
- [141] Zhao F, Cui Y, Li M, et al. Prognostic value of 3'-Deoxy-3'-18F-Fluorothymidine ([F] FLT PET) in patients with recurrent malignant gliomas. *Nucl Med Biol* 2014;41:710-715.
- [142] Laymon CM, Oborski MJ, Lee VK, et al. Combined imaging biomarkers for therapy evaluation in glioblastoma multiforme: correlating sodium MRI and F-18 FLT PET on a voxel-wise basis. *Magn Reson Imaging* 2012;30:1268-1278.
- [143] Sanghera B, Wong WL, Sonoda LI, et al. FLT PET-CT in evaluation of treatment response. *Indian J Nucl Med* 2014;29:65-73.
- [144] Viel T, Schelhaas S, Wagner S, et al. Early assessment of the efficacy of temozolomide chemotherapy in experimental glioblastoma using [18F]FLT-PET imaging. *PLoS One* 2013;8:e67911.
- [145] Wardak M, Schiepers C, Cloughesy TF, Dahlbom M, Phelps ME, Huang SC. (1)(8)F-FLT and (1)(8)F-FDOPA PET kinetics in recurrent brain tumors. *Eur J Nucl Med Mol Imaging* 2014;41:1199-1209.
- [146] Shinomiya A, Kawai N, Okada M, et al. Evaluation of 3'-deoxy-3'- [18F]-fluorothymidine (18F-FLT) kinetics correlated with thymidine kinase-1 expression and cell proliferation in newly diagnosed gliomas. *Eur J Nucl Med Mol Imaging* 2013;40:175-185.
- [147] Brizel DM, Dodge RK, Clough RW, Dewhirst MW. Oxygenation of head and neck cancer: changes during radiotherapy and impact on treatment outcome. *Radiother Oncol* 1999;53:113-117.

- [148] Gatenby RA, Kessler HB, Rosenblum JS, et al. Oxygen distribution in squamous cell carcinoma metastases and its relationship to outcome of radiation therapy. *Int J Radiat Oncol Biol Phys* 1988;14:831-838.
- [149] Kolstad P. Inter-capillary distance, oxygen tension and local recurrence in cervix cancer. *Scand J Clin Lab Invest Suppl* 1968;106:145-157.
- [150] Hall EJ. *Radiobiology for the Radiologist*, 4 ed. Philadelphia, PA: JB Lippincott Company, 1994.
- [151] Rampling R, Cruickshank G, Lewis AD, Fitzsimmons SA, Workman P. Direct measurement of pO₂ distribution and bioreductive enzymes in human malignant brain tumors. *Int J Radiat Oncol Biol Phys* 1994;29:427-431.
- [152] Rumsey WL, Vanderkooi JM, Wilson DF. Imaging of phosphorescence: a novel method for measuring oxygen distribution in perfused tissue. *Science* 1988;241:1649-1651.
- [153] Vinogradov SA, Grosul P, Rozhkov V, et al. Oxygen distributions in tissue measured by phosphorescence quenching. *Adv Exp Med Biol* 2003;510:181-185.
- [154] Zhang S, Hosaka M, Yoshihara T, et al. Phosphorescent light-emitting iridium complexes serve as a hypoxia-sensing probe for tumor imaging in living animals. *Cancer Res* 2010;70:4490-4498.
- [155] Foo SS, Abbott DF, Lawrentschuk N, Scott AM. Functional imaging of intratumoral hypoxia. *Mol Imaging Biol* 2004;6:291-305.
- [156] Kitai T, Tanaka A, Tokuka A, et al. Quantitative detection of hemoglobin saturation in the liver with near-infrared spectroscopy. *Hepatology* 1993;18:926-936.
- [157] Hull EL, Conover DL, Foster TH. Carbogen-induced changes in rat mammary tumour oxygenation reported by near infrared spectroscopy. *Br J Cancer* 1999;79:1709-1716.
- [158] Jobsis FF. Non-invasive, infra-red monitoring of cerebral O₂ sufficiency, blood volume, HbO₂-Hb shifts and bloodflow. *Acta Neurol Scand Suppl* 1977;64:452-453.
- [159] Pennekamp CW, Bots ML, Kappelle LJ, Moll FL, de Borst GJ. The value of near-infrared spectroscopy measured cerebral oximetry during carotid endarterectomy in peri-operative stroke prevention. A review. *Eur J Vasc Endovasc Surg* 2009;38:539-545.
- [160] Jette DC, Wiebe LI, Flanagan RJ, Lee J, Chapman JD. Iodoazomycin riboside (1-(5'-iodo-5'-deoxyribofuranosyl)-2-nitroimidazole), a hypoxic cell marker. I. Synthesis and in vitro characterization. *Radiat Res* 1986;105:169-179.
- [161] Mannan RH, Somayaji VV, Lee J, Mercer JR, Chapman JD, Wiebe LI. Radioiodinated 1-(5-iodo-5-deoxy-beta-D-arabinofuranosyl)-2-nitroimidazole (iodoazomycin arabinoside: IAZA): a novel marker of tissue hypoxia. *J Nucl Med* 1991;32:1764-1770.

- [162] Mannan RH, Mercer JR, Wiebe LI, Kumar P, Somayaji VV, Chapman JD. Radioiodinated azomycin pyranoside (IAZP): a novel non-invasive marker for the assessment of tumor hypoxia. *J Nucl Biol Med* 1992;36:60-67.
- [163] Hustinx R, Eck SL, Alavi A. Potential applications of PET imaging in developing novel cancer therapies. *J Nucl Med* 1999;40:995-1002.
- [164] Sorger D, Patt M, Kumar P, et al. [18F]Fluoroazomycinarabinofuranoside (18FAZA) and [18F]Fluoromisonidazole (18FMISO): a comparative study of their selective uptake in hypoxic cells and PET imaging in experimental rat tumors. *Nucl Med Biol* 2003;30:317-326.
- [165] Koch CJ, Scheuermann JS, Divgi C, et al. Biodistribution and dosimetry of (18)F-EF5 in cancer patients with preliminary comparison of (18)F-EF5 uptake versus EF5 binding in human glioblastoma. *Eur J Nucl Med Mol Imaging* 2010;37:2048-2059.
- [166] Rasey JS, Hofstrand PD, Chin LK, Tewson TJ. Characterization of [18F]fluoroetanidazole, a new radiopharmaceutical for detecting tumor hypoxia. *J Nucl Med* 1999;40:1072-1079.
- [167] Hwang DR, Dence CS, Bonasera TA, Welch MJ. No-carrier-added synthesis of 3-[18F]fluoro-1-(2-nitro-1-imidazolyl)-2-propanol. A potential PET agent for detecting hypoxic but viable tissues. *Int J Rad Appl Instrum A* 1989;40:117-126.
- [168] Kachur AV, Dolbier Jr WR, Evans SM, et al. Synthesis of new hypoxia markers EF1 and [18F]-EF1. *Appl Radiat Isot* 1999;51:643-650.
- [169] Wada H, Iwata R, Ido T, Takai Y. Synthesis of 1- [2- [18F]fluoro-1-(hydroxymethyl)-ethoxy]methyl-2-nitroimidazole ([18F]FENI), a potential agent for imaging hypoxic tissues by PET. *Journal of Labelled Compounds and Radiopharmaceuticals* 2000;43:785-793.
- [170] Yang DJ, Wallace S, Cherif A, et al. Development of F-18-labeled fluoroerythronitroimidazole as a PET agent for imaging tumor hypoxia. *Radiology* 1995;194:795-800.
- [171] Koh WJ, Rasey JS, Evans ML, et al. Imaging of hypoxia in human tumors with [F-18]fluoromisonidazole. *Int J Radiat Oncol Biol Phys* 1992;22:199-212.
- [172] Valk PE, Mathis CA, Prados MD, Gilbert JC, Budinger TF. Hypoxia in human gliomas: demonstration by PET with fluorine-18-fluoromisonidazole. *J Nucl Med* 1992;33:2133-2137.
- [173] Dunn IF, Heese O, Black PM. Growth factors in glioma angiogenesis: FGFs, PDGF, EGF, and TGFs. *J Neurooncol* 2000;50:121-137.
- [174] Fischer I, Gagner JP, Law M, Newcomb EW, Zagzag D. Angiogenesis in gliomas: biology and molecular pathophysiology. *Brain Pathol* 2005;15:297-310.
- [175] Spence AM, Muzi M, Swanson KR, et al. Regional hypoxia in glioblastoma multiforme quantified with [18F]fluoromisonidazole positron emission tomography be-

fore radiotherapy: correlation with time to progression and survival. *Clin Cancer Res* 2008;14:2623-2630.

- [176] Swanson KR, Chakraborty G, Wang CH, et al. Complementary but distinct roles for MRI and 18F-fluoromisonidazole PET in the assessment of human glioblastomas. *J Nucl Med* 2009;50:36-44.
- [177] Szeto MD, Chakraborty G, Hadley J, et al. Quantitative metrics of net proliferation and invasion link biological aggressiveness assessed by MRI with hypoxia assessed by FMISO-PET in newly diagnosed glioblastomas. *Cancer Res* 2009;69:4502-4509.
- [178] Kawai N, Maeda Y, Kudomi N, et al. Correlation of biological aggressiveness assessed by 11C-methionine PET and hypoxic burden assessed by 18F-fluoromisonidazole PET in newly diagnosed glioblastoma. *Eur J Nucl Med Mol Imaging* 2011;38:441-450.
- [179] Bos R, van Der Hoeven JJ, van Der Wall E, et al. Biologic correlates of (18)fluorodeoxyglucose uptake in human breast cancer measured by positron emission tomography. *J Clin Oncol* 2002;20:379-387.
- [180] Clavo AC, Wahl RL. Effects of hypoxia on the uptake of tritiated thymidine, L-leucine, L-methionine and FDG in cultured cancer cells. *J Nucl Med* 1996;37:502-506.
- [181] Evans SM, Hahn SM, Magarelli DP, Koch CJ. Hypoxic heterogeneity in human tumors: EF5 binding, vasculature, necrosis, and proliferation. *Am J Clin Oncol* 2001;24:467-472.
- [182] Rajendran JG, Mankoff DA, O'Sullivan F, et al. Hypoxia and glucose metabolism in malignant tumors: evaluation by [18F]fluoromisonidazole and [18F]fluorodeoxyglucose positron emission tomography imaging. *Clin Cancer Res* 2004;10:2245-2252.
- [183] Hatano T, Zhao S, Zhao Y, et al. Biological characteristics of intratumoral [F-18]fluoromisonidazole distribution in a rodent model of glioma. *Int J Oncol* 2013;42:823-830.
- [184] Cher LM, Murone C, Lawrentschuk N, et al. Correlation of hypoxic cell fraction and angiogenesis with glucose metabolic rate in gliomas using 18F-fluoromisonidazole, 18F-FDG PET, and immunohistochemical studies. *J Nucl Med* 2006;47:410-418.
- [185] Hirata K, Terasaka S, Shiga T, et al. (1)(8)F-Fluoromisonidazole positron emission tomography may differentiate glioblastoma multiforme from less malignant gliomas. *Eur J Nucl Med Mol Imaging* 2012;39:760-770.
- [186] Yamamoto Y, Maeda Y, Kawai N, et al. Hypoxia assessed by 18F-fluoromisonidazole positron emission tomography in newly diagnosed gliomas. *Nucl Med Commun* 2012;33:621-625.
- [187] Kawai N, Lin W, Cao WD, et al. Correlation between F-fluoromisonidazole PET and expression of HIF-1alpha and VEGF in newly diagnosed and recurrent malignant gliomas. *Eur J Nucl Med Mol Imaging* 2014.

- [188] Valable S, Petit E, Roussel S, et al. Complementary information from magnetic resonance imaging and (18)F-fluoromisonidazole positron emission tomography in the assessment of the response to an antiangiogenic treatment in a rat brain tumor model. *Nucl Med Biol* 2011;38:781-793.
- [189] Huang SC, Carson RE, Hoffman EJ, Kuhl DE, Phelps ME. An investigation of a double-tracer technique for positron computerized tomography. *J Nucl Med* 1982;23:816-822.
- [190] Figueiras FP, Jimnez X, Pareto D, et al. Simultaneous dual-tracer PET imaging of the rat brain and its application in the study of cerebral ischemia. *Molecular imaging and biology : MIB : the official publication of the Academy of Molecular Imaging* 2011;13:500-510.
- [191] Kadrmas DJ, Rust TC. Feasibility of rapid multitracer PET tumor imaging. *Nuclear Science, IEEE Transactions on* 2005;52:1341-1347.
- [192] Koeppe RA, Raffel DM, Snyder SE, Ficaro EP, Kilbourn MR, Kuhl DE. Dual-[11C]tracer single-acquisition positron emission tomography studies. *J Cereb Blood Flow Metab* 2001;21:1480-1492.
- [193] Kadrmas DJ, Rust TC, Hoffman JM. Single-scan dual-tracer FLT+FDG PET tumor characterization. *Phys Med Biol* 2013;58:429-449.
- [194] Verhaeghe J, Reader AJ. Simultaneous water activation and glucose metabolic rate imaging with PET. *Physics in medicine and biology* 2013;58:393-411.
- [195] Ikoma Y, Toyama H, Uemura K, Uchiyama A. Evaluation of the reliability in kinetic analysis for dual tracer injection of FDG and flumazenil PET study. In; 2001 2001, 2001: 2054-2057.
- [196] Bell C, Rose S, Puttick S, et al. Dual acquisition of (18)F-FMISO and (18)F-FDOPA. *Phys Med Biol* 2014;59:3925-3949.
- [197] Gao F, Liu H, Jian Y, Shi P. Dynamic dual-tracer PET reconstruction. *Inf Process Med Imaging* 2009;21:38-49.
- [198] Cheng X, Navab N, Ziegler S, Shi K. Direct Parametric Image Reconstruction of Rapid Multi-tracer PET. In: Mori K, Sakuma I, Sato Y, Barillot C, Navab N, eds.: Springer Berlin Heidelberg, 2013: 155-162.
- [199] Heiss WD, Raab P, Lanfermann H. Multimodality assessment of brain tumors and tumor recurrence. *J Nucl Med* 2011;52:1585-1600.
- [200] Hu LS, Baxter LC, Smith KA, et al. Relative cerebral blood volume values to differentiate high-grade glioma recurrence from posttreatment radiation effect: direct correlation between image-guided tissue histopathology and localized dynamic susceptibility-weighted contrast-enhanced perfusion MR imaging measurements. *AJNR Am J Neuroradiol* 2009;30:552-558.

- [201] Stupp R, Brada M, van den Bent MJ, Tonn JC, Pentheroudakis G. High-grade glioma: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2014.
- [202] Chaskis C, Neyns B, Michotte A, De Ridder M, Everaert H. Pseudoprogression after radiotherapy with concurrent temozolomide for high-grade glioma: clinical observations and working recommendations. *Surg Neurol* 2009;72:423-428.
- [203] Wen PY, Macdonald DR, Reardon DA, et al. Updated response assessment criteria for high-grade gliomas: response assessment in neuro-oncology working group. *J Clin Oncol* 2010;28:1963-1972.
- [204] Wahl RL, Jacene H, Kasamon Y, Lodge MA. From RECIST to PERCIST: Evolving Considerations for PET response criteria in solid tumors. *J Nucl Med* 2009;50 Suppl 1:122S-150S.
- [205] Charnley N, West CM, Barnett CM, et al. Early change in glucose metabolic rate measured using FDG-PET in patients with high-grade glioma predicts response to temozolomide but not temozolomide plus radiotherapy. *Int J Radiat Oncol Biol Phys* 2006;66:331-338.
- [206] Jager PL, Vaalburg W, Pruim J, de Vries EG, Langen KJ, Piers DA. Radiolabeled amino acids: basic aspects and clinical applications in oncology. *J Nucl Med* 2001;42:432-445.
- [207] Langstrom B, Antoni G, Gullberg P, et al. Synthesis of L- and D- [methyl-11C]methionine. *J Nucl Med* 1987;28:1037-1040.
- [208] Schwarzenberg J, Czernin J, Cloughesy TF, et al. Treatment Response Evaluation Using 18F-FDOPA PET in Patients with Recurrent Malignant Glioma on Bevacizumab Therapy. *Clin Cancer Res* 2014;20:3550-3559.
- [209] Rapp M, Floeth FW, Felsberg J, et al. Clinical value of O-(2- [(18)F]-fluoroethyl)-L-tyrosine positron emission tomography in patients with low-grade glioma. *Neurosurg Focus* 2013;34:E3.
- [210] Jansen NL, Suchorska B, Wenter V, et al. Dynamic 18F-FET PET in newly diagnosed astrocytic low-grade glioma identifies high-risk patients. *J Nucl Med* 2014;55:198-203.
- [211] Hutterer M, Nowosielski M, Putzer D, et al. O-(2-18F-fluoroethyl)-L-tyrosine PET predicts failure of antiangiogenic treatment in patients with recurrent high-grade glioma. *J Nucl Med* 2011;52:856-864.
- [212] Jansen N, Suchorska B, Wenter V, et al. Prognostic value of [F-18]FET-PET in newly diagnosed low grade glioma. *Society of Nuclear Medicine Annual Meeting Abstracts* 2013;54:355.

- [213] Heinzel A, Stock S, Langen KJ, Muller D. Cost-effectiveness analysis of amino acid PET-guided surgery for supratentorial high-grade gliomas. *J Nucl Med* 2012;53:552-558.
- [214] Stummer W, Pichlmeier U, Meinel T, Wiestler OD, Zanella F, Reulen HJ. Fluorescence-guided surgery with 5-aminolevulinic acid for resection of malignant glioma: a randomised controlled multicentre phase III trial. *Lancet Oncol* 2006;7:392-401.
- [215] Ewelt C, Floeth FW, Felsberg J, et al. Finding the anaplastic focus in diffuse gliomas: the value of Gd-DTPA enhanced MRI, FET-PET, and intraoperative, ALA-derived tissue fluorescence. *Clin Neurol Neurosurg* 2011;113:541-547.
- [216] van Baardwijk A, Baumert BG, Bosmans G, et al. The current status of FDG-PET in tumour volume definition in radiotherapy treatment planning. *Cancer Treat Rev* 2006;32:245-260.
- [217] Lee IH, Piert M, Gomez-Hassan D, et al. Association of ¹¹C-methionine PET uptake with site of failure after concurrent temozolomide and radiation for primary glioblastoma multiforme. *Int J Radiat Oncol Biol Phys* 2009;73:479-485.
- [218] Veas H, Senthamizhchelvan S, Miralbell R, Weber DC, Ratib O, Zaidi H. Assessment of various strategies for ¹⁸F-FET PET-guided delineation of target volumes in high-grade glioma patients. *Eur J Nucl Med Mol Imaging* 2009;36:182-193.
- [219] Pyka T, Gempt J, Ringel F, et al. Prediction of Glioma Recurrence Using Dynamic ¹⁸F-Fluoroethyltyrosine PET. *AJNR Am J Neuroradiol* 2014.
- [220] Dowson N, Thomas P, Fay M, et al. Early prediction of treatment response in advanced gliomas with (18)F-dopa positron-emission tomography. *Curr Oncol* 2014;21:e172-178.
- [221] Schiepers C, Chen W, Cloughesy T, Dahlbom M, Huang SC. ¹⁸F-FDOPA kinetics in brain tumors. *J Nucl Med* 2007;48:1651-1661.
- [222] Ellingson BM, Chen W, Harris RJ, et al. PET Parametric Response Mapping for Clinical Monitoring and Treatment Response Evaluation in Brain Tumors. *PET Clinics* 2013;8:201-217.
- [223] Duffau H. Lessons from brain mapping in surgery for low-grade glioma: insights into associations between tumour and brain plasticity. *Lancet Neurol* 2005;4:476-486.
- [224] Duffau H. *Diffuse Low-Grade Gliomas in Adults*. London: Springer London, 2013.
- [225] Grosu AL, Weber WA. PET for radiation treatment planning of brain tumours. *Radiother Oncol* 2010;96:325-327.
- [226] Berntsson SG, Falk A, Savitcheva I, et al. Perfusion and diffusion MRI combined with (1)(1)C-methionine PET in the preoperative evaluation of suspected adult low-grade gliomas. *J Neurooncol* 2013;114:241-249.

- [227] Rahm V, Boxheimer L, Bruehlmeier M, et al. Focal changes in diffusivity on apparent diffusion coefficient MR imaging and amino acid uptake on PET do not colocalize in nonenhancing low-grade gliomas. *J Nucl Med* 2014;55:546-550.
- [228] Bisdas S, Ritz R, Bender B, et al. Metabolic mapping of gliomas using hybrid MR-PET imaging: feasibility of the method and spatial distribution of metabolic changes. *Invest Radiol* 2013;48:295-301.
- [229] Rangan A, et al. Prognostic value of 18F-FDOPA PET in treated low-grade gliomas with non-enhancing T2 changes on MRI. *Society of Nuclear Medicine Annual Meeting Abstracts*, 2013;54(Supplement 2):359.
- [230] Floeth FW, Pauleit D, Sabel M, et al. Prognostic value of O-(2-18F-fluoroethyl)-L-tyrosine PET and MRI in low-grade glioma. *J Nucl Med* 2007;48:519-527.
- [231] Grosu AL, Weber WA, Franz M, et al. Reirradiation of recurrent high-grade gliomas using amino acid PET (SPECT)/CT/MRI image fusion to determine gross tumor volume for stereotactic fractionated radiotherapy. *Int J Radiat Oncol Biol Phys* 2005;63:511-519.
- [232] Hutterer M, Nowosielski M, Putzer D, et al. [18F]-fluoro-ethyl-L-tyrosine PET: a valuable diagnostic tool in neuro-oncology, but not all that glitters is glioma. *Neuro Oncol* 2013;15:341-351.
- [233] Pirotte BJ, Levivier M, Goldman S, et al. Positron emission tomography-guided volumetric resection of supratentorial high-grade gliomas: a survival analysis in 66 consecutive patients. *Neurosurgery* 2009;64:471-481; discussion 481.
- [234] Galldiks N, Rapp M, Stoffels G, et al. Response assessment of bevacizumab in patients with recurrent malignant glioma using [18F]Fluoroethyl-L-tyrosine PET in comparison to MRI. *Eur J Nucl Med Mol Imaging* 2013;40:22-33.
- [235] Grosu AL, Weber WA, Astner ST, et al. 11C-methionine PET improves the target volume delineation of meningiomas treated with stereotactic fractionated radiotherapy. *Int J Radiat Oncol Biol Phys* 2006;66:339-344.
- [236] Graf R, Plotkin M, Steffen IG, et al. Magnetic resonance imaging, computed tomography, and 68Ga-DOTATOC positron emission tomography for imaging skull base meningiomas with intracranial extension treated with stereotactic radiotherapy--a case series. *Head Face Med* 2012;8:1.
- [237] Thorwarth D, Muller AC, Pfannenberger C, Beyer T. Combined PET/MR imaging using (68)Ga-DOTATOC for radiotherapy treatment planning in meningioma patients. *Recent Results Cancer Res* 2013;194:425-439.
- [238] Cornelius JF, Langen KJ, Stoffels G, Hanggi D, Sabel M, Jakob Steiger H. Positron emission tomography imaging of meningioma in clinical practice: review of literature and future directions. *Neurosurgery* 2012;70:1033-1041; discussion 1042.

- [239] Cornelius JF, Slotty PJ, Stoffels G, Galldiks N, Langen KJ, Steiger HJ. 5-Aminolevulinic Acid and (18)F-FET-PET as Metabolic Imaging Tools for Surgery of a Recurrent Skull Base Meningioma. *J Neurol Surg B Skull Base* 2013;74:211-216.
- [240] Bisdas S, Nagele T, Schlemmer HP, et al. Switching on the lights for real-time multi-modality tumor neuroimaging: The integrated positron-emission tomography/MR imaging system. *AJNR Am J Neuroradiol* 2010;31:610-614.
- [241] Boss A, Bisdas S, Kolb A, et al. Hybrid PET/MRI of intracranial masses: initial experiences and comparison to PET/CT. *J Nucl Med* 2010;51:1198-1205.
- [242] Schwenzer NF, Stegger L, Bisdas S, et al. Simultaneous PET/MR imaging in a human brain PET/MR system in 50 patients--current state of image quality. *Eur J Radiol* 2012;81:3472-3478.
- [243] Jarzabek MA, Sweeney KJ, Evans RL, et al. Molecular imaging in the development of a novel treatment paradigm for glioblastoma (GBM): an integrated multidisciplinary commentary. *Drug Discov Today* 2013;18:1052-1066.
- [244] James ML, Gambhir SS. A Molecular Imaging Primer: Modalities, Imaging Agents, and Applications. *Physiol Rev* 2012;92:897-965.
- [245] Kelkar SS, Reineke TM. Theranostics: Combining Imaging and Therapy. *Bioconjugate Chem* 2011;22:1879-1903.
- [246] Cutler CS, Hennkens HM, Sisay N, Huclier-Markai S, Jurisson SS. Radiometals for Combined Imaging and Therapy. *Chem Rev* 2013;113:858-883.
- [247] Boase NRB, Blakey I, Thurecht KJ. Molecular imaging with polymers. *Polym Chem-Uk* 2012;3:1384-1389.
- [248] M. S, G. T. PET imaging using radiolabelled antibodies: future direction in tumor diagnosis and correlate applications. *Research and Reports in Nuclear Medicine* 2013;3:9-17.
- [249] Stockhofe K, Postema J, Schieferstein H, Ross T. Radiolabeling of Nanoparticles and Polymers for PET Imaging. *Pharmaceuticals* 2014;7:392-418.
- [250] Accardo A, Tesauro D, Morelli G. Peptide-based targeting strategies for simultaneous imaging and therapy with nanovectors. *Polym J* 2013;45:481-493.
- [251] Zhu Q, Qiu F, Zhu BS, Zhu XY. Hyperbranched polymers for bioimaging. *Rsc Adv* 2013;3:2071-2083.
- [252] Boase NRB, Blakey I, Rolfe BE, Mardon K, Thurecht KJ. Synthesis of a multimodal molecular imaging probe based on a hyperbranched polymer architecture. *Polymer Chemistry* 2014;5:4450-4458.
- [253] Wehrenberg-Klee E, Redjal N, Leece A, et al. A F(ab)'2-based EGFR targeted PET probe for glioma imaging. *J NUCL MED MEETING ABSTRACTS* 2014;55:1793-.

- [254] Petrulli JR, Sullivan JM, Zheng MQ, et al. Quantitative Analysis of [C-11]-Erlotinib PET Demonstrates Specific Binding for Activating Mutations of the EGFR Kinase Domain. *Neoplasia* 2013;15:1347-1353.
- [255] Olafsen T, Betting D, Kenanova VE, et al. Recombinant Anti-CD20 Antibody Fragments for Small-Animal PET Imaging of B-Cell Lymphomas. *J Nucl Med* 2009;50:1500-1508.
- [256] Natarajan A, Hackel BJ, Gambhir SS. A Novel Engineered Anti-CD20 Tracer Enables Early Time PET Imaging in a Humanized Transgenic Mouse Model of B-cell Non-Hodgkins Lymphoma. *Clin Cancer Res* 2013;19:6820-6829.
- [257] Cai WB, Ebrahimnejad A, Chen K, et al. Quantitative radioimmunoPET imaging of EphA2 in tumor-bearing mice. *Eur J Nucl Med Mol I* 2007;34:2024-2036.
- [258] Rosch F. Past, present and future of Ge-68/Ga-68 generators. *Appl Radiat Isotopes* 2013;76:24-30.
- [259] Hofman MS, Kong G, Neels OC, Eu P, Hong E, Hicks RJ. High management impact of Ga-68 DOTATATE (GaTate) PET/CT for imaging neuroendocrine and other somatostatin expressing tumours. *J Med Imag Radiat On* 2012;56:40-47.
- [260] Hofman MS, Hicks RJ. Changing Paradigms with Molecular Imaging of Neuroendocrine Tumors. *Discov Med* 2012;74:71-81.
- [261] Al-Nahhas A, Win Z, Szyszko T, et al. Gallium-68 PET: A new frontier in receptor cancer imaging. *Anticancer Res* 2007;27:4087-4094.
- [262] Thurecht KJ, Blakey I, Peng H, et al. Functional Hyperbranched Polymers: Toward Targeted in Vivo F-19 Magnetic Resonance Imaging Using Designed Macromolecules. *J Am Chem Soc* 2010;132:5336-+.
- [263] Rolfe BE, Blakey I, Squires O, et al. Multimodal Polymer Nanoparticles with Combined F-19 Magnetic Resonance and Optical Detection for Tunable, Targeted, Multimodal Imaging in Vivo. *J Am Chem Soc* 2014;136:2413-2419.
- [264] Srinivas M, Heerschap A, Ahrens ET, Figdor CG, de Vries IJM. F-19 MRI for quantitative in vivo cell tracking. *Trends Biotechnol* 2010;28:363-370.
- [265] Ruiz-Cabello J, Barnett BP, Bottomley PA, Bulte JWM. Fluorine (F-19) MRS and MRI in biomedicine. *Nmr Biomed* 2011;24:114-129.
- [266] Ahrens ET, Flores R, Xu HY, Morel PA. In vivo imaging platform for tracking immunotherapeutic cells. *Nat Biotechnol* 2005;23:983-987.
- [267] Du WJ, Nystrom AM, Zhang L, et al. Amphiphilic Hyperbranched Fluoropolymers as Nanoscopic (19)F Magnetic Resonance Imaging Agent Assemblies. *Biomacromolecules* 2008;9:2826-2833.

- [268] Peng H, Blakey I, Dargaville B, Rasoul F, Rose S, Whittaker AK. Synthesis and evaluation of partly fluorinated block copolymers as MRI imaging agents. *Biomacromolecules* 2009;10:374-381.
- [269] Ahrens ET, Bulte JWM. Tracking immune cells in vivo using magnetic resonance imaging. *Nat Rev Immunol* 2013;13:755-763.
- [270] Ahrens ET, Schwartzman D, inventors; Univ Carnegie Mellon (UYCM-C) Univ Pittsburgh Commonwealth System High (UPIT-C) Ahrens E T (AHRE-Individual) Schwartzman D (SCHW-Individual), assignee. Image-guided treatment for condition associated with inflammatory response in internal organ by administering fluorine-19 MRI fluorocarbon imaging reagent, identifying internal organ and administering medicament patent WO2014043677-A2; US2014093452-A1 WO2014043677-A2 20 Mar 2014 A61K-049/06 201425.
- [271] Oishi M, Sumitani S, Nagasaki Y. On-off regulation of F-19 magnetic resonance signals based on pH-Sensitive PEGylated nanogels for potential tumor-specific smart F-19 MRI probes. *Bioconjugate Chem* 2007;18:1379-1382.
- [272] Criscione JM, Le BL, Stern E, et al. Self-assembly of pH-responsive fluorinated dendrimer-based particulates for drug delivery and noninvasive imaging. *Biomaterials* 2009;30:3946-3955.
- [273] Nurmi L, Peng H, Seppala J, Haddleton DM, Blakey I, Whittaker AK. Synthesis and evaluation of partly fluorinated polyelectrolytes as components in F-19 MRI-detectable nanoparticles. *Polym Chem-Uk* 2010;1:1039-1047.
- [274] Wang KW, Peng H, Thurecht KJ, Puttick S, Whittaker AK. pH-responsive star polymer nanoparticles: potential F-19 MRI contrast agents for tumour-selective imaging. *Polym Chem-Uk* 2013;4:4480-4489.
- [275] Wang KW, Peng H, Thurecht KJ, Puttick S, Whittaker AK. Biodegradable core cross-linked star polymer nanoparticles as F-19 MRI contrast agents for selective imaging. *Polym Chem-Uk* 2014;5:1760-1771.
- [276] Langereis S, Keupp J, van Velthoven JLJ, et al. A Temperature-Sensitive Liposomal H-1 CEST and F-19 Contrast Agent for MR Image-Guided Drug Delivery. *J Am Chem Soc* 2009;131:1380-+.
- [277] Mizukami S, Takikawa R, Sugihara F, Shirakawa M, Kikuchi K. Dual-Function Probe to Detect Protease Activity for Fluorescence Measurement and F-19 MRI. *Angew Chem Int Edit* 2009;48:3641-3643.