We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Histone Methylation - A Cornerstone for Plant Responses to Environmental Stresses?

Roberto Bobadilla and Alexandre Berr

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/61733>

Abstract

Plants are sessile organisms and, as such, their survival relies on their ability to respond quickly all along their life cycle to any kind of environmental stimuli, including abiotic and biotic stresses. In this respect, plants have developed efficient mechanisms of protection and/or adaptation to minimize deleterious effects of stress on their growth and development. In a stress type-dependent manner, external signals are firstly sensed. This step is then followed by the activation of particular signalling pathways, resulting ultimately in the rapid and specific modulation of the plant transcriptome. Currently, transcriptional regulation is considered as a central process in the build-up of plant responses to both abiotic and biotic stresses. Among mechanisms involved in transcriptional regulation, the combined effect of different histone tail post-translational modifications (PTMs; e.g. acetylation and methylation) through the activity of particular histone-modifying enzymes can lead to changes in the local chromatin structure environment and hence the underlying DNA accessibility.

By focusing on histone lysine methylation, in this chapter we highlight our current understanding of the transcriptional roles played by chromatin-remodelling mechanisms in regulating plant response/adaptation to different biotic and abiotic stresses. Based on recent advances, we further discuss the stability and transmission of such methylation marks to subsequent generations, with the underlying idea of an epigenetically based transcriptional memory of stresses in plants.

Keywords: Histone methylation and demethylation, histone methyltransferases and demethylases, biotic and abiotic stresses

1. Introduction

Stress, as we currently think of it, is a highly subjective phenomenon defined as a state of threatened homeostasis. Depending on their nature, external stresses are usually divided into biotic (i.e. herbivorous insects and pathogens such as fungi, bacteria and viruses) or abiotic (i.e. including, among others, high or low temperature, submergence or drought and salinity). During their lifetime, all living organisms inevitably and constantly face all sorts of environmental stresses that often occur suddenly and/or simultaneously. Classically, different strategies can be applied to minimize deleterious effects of stresses, such as resistance, tolerance, avoidance or escape. Being sessile, plants cannot escape and are therefore more prone to the deleterious effect of unfavourable environmental growth conditions. Because responses are critical to ensure their survival, plants have developed specific and efficient strategies that allow them to precisely perceive different environmental stresses and respond and/or adapt to them [1, 2]. In addition to preformed defence traits, plants have evolved inducible defence strategies. Indeed, upon perception, each stress will raise a complex and more or less specific repertoire of cellular and molecular responses implemented by the plant to minimize or prevent damage. Particularly, the stimulation of a given stress-signalling pathway after pathogen detection will be integrated into the plant cell nucleus through a set of regulatory transcription factor cascades, which prioritizes defence over growth-related cellular functions, while conserving enough valuable resources for survival and reproduction [3, 4]. Supporting the idea that the capacity of a plant to rapidly reprogramme its gene expression at the transcriptional level is an essential and common component of all plant response strategies to stress and disturbance; more than 1,000 transcription factors were found to be involved in stress responses [5, 6]. Because eukaryotic genes function in the context of chromatin, modifications and remodelling of the chromatin configuration from permissive for transcription to restrictive, and vice versa, may be an integral part of mechanisms involved in this vital transcriptional reprogramming. In this chapter, we review and discuss the current knowledge about the functional impact of chromatin changes on the transcriptional regulation of genes under different stress conditions, with particular emphasis on histone methylation/demethylation.

2. Chromatin structure and histone methylation/demethylation

In eukaryotes, genomic DNA in the cell nucleus is packaged in a complex and evolutionarily conserved structure named chromatin, with nucleosome as the basic unit. The nucleosome complex contains about 160–241 base pairs (bp) of DNA, a nucleosome core particle and the H1 linker histone. The nucleosome core particle is composed of an octamer of core histones, consisting of two H3–H4 dimers associated with two H2A–H2B dimers. About 146 bp of DNA is wrapped in ~1.65 negatively supercoiled circles around the histone octamer, while the linker DNA associated with H1 varies in length from 8 to 114 bp [7]. At first sight, the chromatin as it is described appears as a barrier, restricting the access of all kinds of enzymes that process the DNA. However, nucleosomes are not merely static but highly dynamic entities. Indeed,

nucleosomes can be moved, stabilized/destabilized, disassembled/reassembled at particular genome locations in response to specific environmental signals or developmental cues [8]. This dynamic leads to a wide range of chromatin condensation states modulating the DNA accessibility, with euchromatin, being relaxed, and heterochromatin, being compacted. Therefore, in eukaryotic cells, an intimate connection exists between the structural organization of the genome and its functioning. For this reason, the level of chromatin condensation is directly related to all aspects of DNA metabolism, thus playing a major role in regulating transcription, DNA replication, DNA repair, recombination, transposition and chromosome segregation. In plants, changes in the chromatin structure were reported to affect various biological processes such as root growth, flowering, organogenesis, gametophyte or embryo formation [9–11].

In the nucleosome core particle, histones H2A, H2B, H3 and H4 possess two common regions, a histone-fold domain and a histone tail. The histone-fold domain is the most conserved region and the main element of histone dimerization [12]. The tail protrudes from the nucleosome core particle and is more variable and unstructured than the fold [13]. All four core histones have an N-terminal tail domain, but only histone H2A has an additional long C-terminal tail. Histone tails are extremely basic due to their particularly high content in basic amino acid residues, such as lysine and arginine [14]. Resulting positive charges allow them to closely associate with the negatively charged nucleosomal DNA through electrostatic interactions [15]. In addition, histone tails, especially N-terminal ones, may undergo diverse types of post-translational modifications such as acetylation or methylation. The great diversity of these modifications as well as the high number of amino acid residues that can be modified within histone tails, and the correlation between these modifications and various nuclear processes, lead to the hypothesis that the specific combination of histone modifications constitute a histone 'code' [16].

Technically, these histone marks can be localized by chromatin immunoprecipitation (ChIP) using specific antibodies against the modification [17]. Briefly, protein–DNA interactions are stabilized by cross-linking with formaldehyde; chromatin is sheared into small pieces to facilitate analysis and then immunoprecipitated using an antibody raised against a specific histone modification. Following enrichment, cross links are reversed to release DNA, which is then quantified by polymerase chain reaction (PCR) to measure the relative amount of the specific histone mark on selected plant genes. ChIP can also be combined with microarray hybridization (ChIP-chip) or high-throughput sequencing (ChIP-Seq), allowing the genome-wide discovery of DNA–histone modification interactions.

Methylation is the most abundant one compared with other histone PTMs. It can occur at both lysine (K) and arginine (R) residues of core histone tails. Further extending the indexing potential of this modification, mono-, di- and trimethylation of lysine and mono- and dimethylation (symmetric or asymmetric) of arginine are common at N-terminal tails of H2A, H2B, H3 and H4. Although histone acetylation is generally associated with active gene transcription, histone methylation can be associated with either active or silent gene expression, depending upon the histone, the methylated residue or the level (mono-, di- or tri-) of methylation. In *Arabidopsis*, genome-wide analyses revealed that trimethylations of H3K4 and H3K36

(H3K4me3 and H3K36me3) are generally enriched at actively transcribed genes, whereas H3K27me3 is associated with repressed genes and H3K9me2 and H4K20me1 are enriched at constitutive heterochromatin and silenced transposons [18]. For histone arginine methylation, a definitive role has not yet been clearly established. However, because the level of symmetric H3R2me2 and H4R3me2 was negatively correlated with the level of H3K4me3, a well-known mark reflecting active transcription, high levels of H3R2me2 and H4R3me2 are thought to cause transcriptional repression [19–21]. In contrast, asymmetric H4R3me2 was associated with gene activation [22, 23].

Histone methylation is relatively stable and can be established on lysine and arginine by two distinct families of enzymes, the histone lysine methyltransferases (HKMTs), all containing the evolutionary conserved catalytic SET domain in plants [24], and the protein arginine methyltransferases (PRMTs) [25], respectively. As a counterpart, methyl groups on histone can also be removed by at least two evolutionarily conserved classes of histone demethylases, the lysine-specific demethylase1 (LSD1) type and the Jumonji C (JmjC) domain-containing demethylases [26]. Histone methyltransferases and demethylases are well conserved in angiosperms and have been identified and classified on the basis of phylogenetic analyses and domain organization in several plants, including *Arabidopsis*, maize, tomato, rice, grapevine and *Brassica rapa*, [27–32]. However, cellular and molecular functions of many of these modifiers have not yet been addressed.

Although histone acetylation can directly modulate the chromatin structure, arginine and lysine methylation of histone tails can promote or prevent the docking of key transcriptional effector molecules, named readers, needed to ‘translate’ the code in order to determine the functional and structural outcome of the corresponding PTMs. Just as there are a large number of PTMs on histone tails, there are also numerous protein domains that recognize and bind to particular PTMs on these tails. For example, PTM-recognition domains such as plant homeo-domain (PHD) fingers, chromodomains and Tudor domains all recognize methylated lysine residues [33].

3. Histone methylation changes associated with biotic stress conditions

Biotic stress is the result of the damage done to plants by insects or pathogens, such as bacteria or fungi. Plant pathogens are generally divided into two distinct categories: biotrophs, which colonize living plant tissue and obtain nutrients from living host cells, and necrotrophs, which depend on dead host tissue for nutrients and reproduction. To fend off pathogens with different infection strategies, plants have evolved complex defence mechanisms. Classically, the pathogen-sensing machinery induces signalling cascades that promote the accumulation of hormones such as salicylic acid (SA) or jasmonic acid (JA)/ethylene (ET) [34]. These hormones then orchestrate the overall plant defence reaction locally and systemically by inducing the transcriptional activation of defence genes through an intricate signalling network. In this part, we highlight recent examples illustrating how histone methylations condition major steps leading to immunity, ranging from initial pathogen perception to hormonal homeostasis changes for antimicrobial effector expression.

3.1. Histone methylation/demethylation in the defence against biotrophic pathogens

The phytohormone SA plays an important role in plant defence, from the induction of pathogen resistance (*PR*) genes against biotrophic bacteria (e.g. *Pseudomonas syringae*) to the establishment of systemic acquired resistance (SAR) [35]. Several studies suggested that the SA signalling pathway is notably controlled by histone methylation. Under normal growth conditions, *Arabidopsis* mutants for *SNI1* (*Suppressor of NPR1, Inducible*), a negative regulator of SAR required to dampen the basal expression of *PR* genes, presented an increased H3K4me2 on *PR1* [36]. Rather than being a constitutive mark of transcription, H3K4me2 was proposed to be involved in the fine-tuning of tissue-specific expression [37]. Using the functional SA-analogue S-methyl benzo [1,2,3] thiadiazole-7-carbothioate (BTH), an increased level of H3K4me2 on *PR1* was observed in wild-type plants 48 h after treatment and was not detected in mutants. Interestingly, when expressed in yeast, SNI1 also repressed transcription, suggesting a highly conserved mechanism of transcriptional repression. These results together with the structural similarity of SNI1 with armadillo repeat (ARM) proteins (i.e. a motif known to mediate protein-protein interactions) imply that SNI1 may form a scaffold for interaction with proteins that modulates the chromatin structure of *PR* genes, thus repressing their transcription. In addition, the presence of H3K4me2 detected on *PR1* before induction suggested that this mark is readily in place, providing the appropriate chromatin configuration for the efficient induction of *PR1* upon need. Using a similar approach, Alvarez-Venegas et al. [38] reported no significant changes in levels of H3K4me2 and H3K4me3 on *PR1* 24 h after the SA treatment [38]. This discrepancy may reflect differences in experimental conditions. Indeed, the action of the so-called 'SA-analog' BTH on gene transcription is significantly broader than the action of SA itself [39]. Moreover, samplings were performed 48 h versus 24 h after treatment. Together, because the H3K4 methylation increase does not occur immediately after the induction of *PR1*, this mark may not be directly related to the transcriptional induction itself, but later, for the maintenance/reinforcement of *PR1* expression.

The ARABIDOPSIS HOMOLOG OF TRITHORAX (ATX1) is a H3K4 trimethyltransferase providing basal resistance against *Pseudomonas syringae* pv. *tomato* (*Pst*); [40]. Despite being not induced by either *Pst* infection or SA, ATX1 positively and directly regulates the expression of the transcription factor WRKY70 through H3K4 trimethylation at the WRKY70 promoter. In addition, *atx1* mutant shows induced expression of the JA-inducible *THI1.2* gene and the reduced *PR1* expression without detectable changes in their chromatin, resulting in impaired resistance to *Pst* infection. Since the transcriptional factor WRKY70 was positioned at the convergence node of the SA and JA signalling pathways, activating the SA-responsive *PR1* gene and repressing the JA inducible genes [41], ATX1 was proposed to indirectly regulate *PR1* and *THI1.2* through WRKY70. SET DOMAIN GROUP 8 (SDG8), another HKMT encoding the major *Arabidopsis* H3K36 di- and trimethyltransferase [42], was also involved in the plant-defence against *Pst*, but it was more upstream than ATX1 [43]. Indeed, SDG8 sustains the basal transcription of particular *R* genes (*RPM1* or *LAZ5*) by maintaining a basal level of H3K36me3, another histone mark tightly associated with active transcription. SDG8 is also required for the transcriptional induction of these *R* genes upon BTH treatment or *Pst* inoculation. However, this induction occurs without any detectable increase of H3K36me3. Therefore, in resting

plants, SDG8 may establish a 'permissive' chromatin structure at some *R* genes by methylating H3K36, thus ensuring their basal expression and their transcriptional inducibility upon need. Similarly as *atx1* and *sdg8* mutants, loss-of-function mutants for the putative HKMT SDG7 were also found to be more sensitive to *Pst* infection than wild-type plants [44]. The expression of other *R* genes seems to be under the control of histone methylation. Indeed, enhanced downy mildew 2 (EDM2) impacts disease resistance by controlling levels of H3K9me2 at an alternative polyadenylation site in the immune receptor gene *RPP7*, thus regulating the balance between full-length *RPP7* transcripts and prematurely polyadenylated transcripts, which do not encode the *RPP7* immune receptor [45, 46]. EDM2, as an epigenetic 'reader', contains two stretches of atypical PHD-finger motifs known to dock specifically several forms of methylated or unmethylated lysine residues on histones [47]. Besides this, EDM2 was also proposed to cooperate within a large protein complex with EMSY-like (AtEML) members, harbouring an Agenet domain related to the Tudor domain family of epigenetic 'readers' [48].

Apart from *Arabidopsis*, little is known about the regulatory role of histone methylation in the defence against pathogen attack in other plant species. In rice, the JmjC protein gene *JMJ705* encoding a histone lysine demethylase that specifically reverses H3K27me2/3 was found induced during infection with the bacterial pathogen *Xanthomonas oryzae*. *JMJ705* was further involved in the dynamic removal of the basal H3K27me3 over defence-related genes, thereby increasing their basal expression and/or potentiating their higher expression upon biotic stress. Interestingly, the *JMJ705* overexpression resulted in an enhanced resistance to the bacterial pathogen, while its mutation reduces the plant resistance [49].

3.2. Histone methylation/demethylation in the defence against necrotrophic pathogens

While to combat biotrophic pathogens the plant activates mainly the SA signalling pathway, the activation of the JA/ET signalling pathway is prominent to mediate defences against necrotrophic pathogens and herbivorous insect attacks [50]. The involvement of histone methylation in the defence against necrotrophic pathogens is far less documented as compared with the defence against biotrophic pathogens. Besides being more susceptible to *Pst* [43], *sdg8* mutants were also reported to be more sensitive to necrotrophic fungal pathogens such as *Alternaria brassicicola* (*Alt*) and *Botrytis cinerea* [51]. This increased susceptibility was the consequence of the inefficient transcriptional induction of different genes along the JA/ET signalling pathway that was correlated with a stably weak level of H3K36me3 at these genes. Inversely, in wild-type plants, H3K36me3 together with gene expression were increased upon *Alt* infection or stimulation with exogenous MeJA. Under resting conditions, a similarly weak level of H3K36me3 was correlated with a reduced basal expression in *sdg8*. On that account, H3K36 methylation was proposed to act as a 'permissive' mark correlated with gene activity and readily in place at a subset of JA/ET signalling-related genes to raise their rapid and efficient transcriptional induction when required [52]. Interestingly, a stable and very low level of H3K27me3 was detected in defence effector genes. Because H3K27me3 is often associated with epigenetic silencing [53], this low H3K27me3 level may provide these genes with a reduced probability for undesired silencing, thus participating in the reactivity of plants to pathogen infections.

4. Histone methylation changes associated with abiotic stress conditions

Abiotic stresses such as heat, cold, drought, salinity and nutrient deficiency are inherent to every ecosystem and essentially unavoidable. Abiotic stresses are considered the most harmful factors in terms of growth and productivity of crops worldwide [54, 55], especially when they occur in combinations [56]. Here, we summarize and discuss various studies in order to clarify the functional involvement of different histone methylation marks in setting up plant responses to adverse environmental growth conditions.

4.1. Histone methylation/demethylation and the plant stress hormone ABA

The phytohormone abscisic acid (ABA) is a crucial signalling molecule playing versatile functions in regulating many developmental processes, including seed dormancy and germination [57, 58]. ABA also plays a pivotal role in adaptive stress processes, integrating both biotic and abiotic environmental constraints in a complex network of interacting pathways with crosstalks at different levels [59–61]. Currently, ABA is considered as a global regulator of stress responses that can dominantly control the switch in priority between the responses to biotic or abiotic stress, allowing plants to respond to the most severe threat [62].

The transition from heterotrophic to autotrophic development at the post-germinative stage (i.e. embryonic state) is highly vulnerable to osmotic stress [63]. During a period of osmotic stress, ABA promotes the expression of transcription factors such as *ABI3* and *ABI5*, which in turn delay germination and lead to osmotolerance and survival [64]. In *Arabidopsis*, mutation in *PICKLE* (*PKL*), encoding a putative chromatin modifier, results in increased and abnormally sustained expression of *ABI3* and *ABI5* in response to exogenous ABA treatment. This sustained expression was correlated with reduced levels of H3K9me2 and H3K27me2, two methylation marks found in the chromatin of silent genes [65]. Based on these results, it was suggested that *PKL* might act on *ABI3* and *ABI5* to promote directly or indirectly the formation of a repressed chromatin state through a so-far-unknown mechanism.

In adult plants, the establishment of a response and tolerance to drought stress by ABA has been extensively studied and is well discussed in several outstanding reviews [58, 66]. Briefly, under drought conditions, water stress perception triggers ABA biosynthesis and increased tissue ABA accumulation, resulting in stomatal closure and reduced transpiration. Among major enzymes involved in the ABA biosynthesis pathway, NINE-CIS-EPOXYCAROTENOID DIOXYGENASE 3 (*NCED3*) is thought to be the rate-limiting enzyme [67]. In *Arabidopsis*, the increased transcription of *NCED3* upon dehydration was correlated with the binding of the HKMT ATX1 and the increased level of H3K4me3 at *NCED3* [68]. Therefore, the loss-of-function *atx1* mutant showed less tolerance to dehydration, notably because of the lower enrichment of RNA Polymerase II (RNAPII) and H3K4me3 at *NCED3* under stress. ATX1-modified H3K4me3 may thus have an important function in the transcriptional regulation of *NCED3*. However, it is still unclear whether this function is directly linked with the transcriptional induction or is more related to the reinforcement of the increased transcription upon stress perception.

4.2. Histone methylation/demethylation in response to water stresses

Water stresses including drought or submergence are major environmental factors limiting plant growth and crop productivity worldwide [69, 70]. Consequently, plants have evolved a variety of biochemical and physiological mechanisms to respond/adapt to these stresses [71, 72]. In the following section, we distinctly address the involvement of histone methylation in responses to drought and submergence.

4.2.1. Drought stress

Using *Arabidopsis*, the molecular response to water deficit was found to rely notably on the transcriptional regulation of stress-inducible genes with products thought to function in drought tolerance and response [73]. Using ChIP analyses in 15-day-old *Arabidopsis* seedlings, the level of the active mark H3K4me3 was found gradually enriched in response to dehydration stress, preferentially on the coding region of four drought-inducible genes (*RD29A*, *RD29B*, *RD20* and *RAP2.4*), and was correlated with their upregulation [74]. Consistent with this result, *ATX1* was reported to be required for the efficient transcriptional induction of *RD29A* and *RD29B* during a dehydration stress response in an ABA-independent manner [68]. Also, for unknown reasons, it is worth noting that upon exogenous ABA treatment the transcriptional induction of *RD29A* and *RD29B* was stronger in *atx1* mutants than in wild-type plants (their basal transcript levels were lower in *atx1* than in wild-type, while their transcript levels upon ABA treatment were identical [68]). Next, for the ABA-dependent *RD29A* and ABA-independent *RAP2.4* genes, a time lag was observed between their transcriptional induction and the increase in H3K4me3 [74]. Based on these findings and the rapid saturation in RNAPII enrichment compared with H3K4me3 (i.e. already saturated 1 h after stress exposure for RNAPII, while H3K4me3 was still increasing up to 5 h), Kim et al. [74] concluded that the H3K4me3 enrichment may be established, gradually, in response to drought stress after full transcriptional activation of *RD29A* and *RAP2.4*. Because the timing of H3K4me3 enrichment followed subsequent to the RNAPII enrichment, H3K4me3 might be dispensable for the initiation of transcription. Finally, the gradual increase of H3K4me3 further indicates that the longer the stress lasts, the more H3K4me3 will be enriched, suggesting that the epigenetic responsiveness must depend on the intensity of a stress [75].

In a similar approach but using ChIP-Seq, van Dijk et al. [76] established the whole-genome distribution patterns of H3K4me1, H3K4me2 and H3K4me3 in 4-week-old rosette *Arabidopsis* leaves under dehydration stress conditions. They also observed a strong correlation between H3K4me3 abundance and transcripts levels from responding genes. Indeed, among the most strongly downregulated genes, an increase in H3K4me1 and a decrease in H3K4me3 were detected, suggesting the involvement of a histone demethylase in modulating the expression of a subset of stress-responsive genes. Supporting this finding, a putative PKDM7 subfamily-like H3K4 demethylase homologue and two putative demethylase enzymes containing a JmjC domain were found to be drought-inducible in two barley cultivars and in young peanut plants, respectively [77, 78]. Surprisingly, in contrast to the classical genome-wide H3K4me3 enrichment around the transcriptional start site observed in all other eukaryotes [79], H3K4me3 displayed a broader distribution on dehydration and ABA-inducible

genes. Such an unusual profile may reflect a function not strictly related to transcription initiation, as will be discussed later in the stress memory section.

Through a genome-wide approach in rice seedling, Zong et al. [80] also uncovered a weak but positive correlation between H3K4me3 enrichment and the transcript level of some drought-responsive genes under drought stress. This correlation was extended to many genes involved in stress-related metabolite and hormone signalling pathways, further supporting the role played by H3K4me3 in the stress response [80]. However, because H3K4me3 is not the only histone mark for gene activation, this weak correlation may reflect that other active histone marks may also play important roles in regulating gene expression in response to stress in rice. Although these large data sets have provided much information on drought responses in rice, more detailed analyses will be required to elucidate whether the observed variations in H3K4 methylation are a cause or a consequence of the transcriptional changes triggered by water stress. Moreover, identifying key histone modification enzymes is indispensable to better understand the transcriptional regulatory network of the abiotic stress response.

4.2.2. Submergence

Submergence is a complex stress that encompasses many changes in environmental factors, including light intensity, pH and dissolved oxygen concentration. Alcoholic fermentation is important for the survival of plants especially under anaerobic environments [81]. In rice, *alcohol dehydrogenase 1* (*ADH1*) and *pyruvate decarboxylase 1* (*PDC1*) genes are involved in this anaerobic metabolism and their expression is reversibly induced (i.e. activated upon submergence and repressed upon re-aeration) [82]. Using these two genes as a model, Tsuji et al. [83] observed that the level of H3K4 methylation, specifically at both the 5'- and 3'-coding regions of *ADH1* and *PDC1*, was changed from a dimethylated state to a trimethylated state upon their transcriptional upregulation in response to submergence. This change was reverted back to its initial level following re-aeration, indicating that in this particular case, H3K4me3 does not serve as a memory mark of a prior transcriptional activity. Similarly to drought, these results highlight the dynamic and reversible change of histone H3K4 methylation at stress-related genes in response to the occurrence and disappearance of a stress.

4.3. Histone methylation/demethylation in response to salt stress

Salinity is also a serious factor affecting plants in several ways (i.e. water stress, ion toxicity, nutritional disorders, oxidative stress, alteration of metabolic processes, membrane disorganization, genotoxicity, reduction of cell division and expansion), thus limiting plant growth, development and survival [84]. In *Arabidopsis*, a NaCl or ABA treatment has been shown to induce the transcription of a range of abiotic stress-responsive genes (*ABI1*, *ABI2*, *KAT1*, *KAT2*, *DREB2A*, *RD29A* and *RD29B*). Using ChIP, this induction was found significantly correlated with an increase in the active mark H3K4me3 and a decrease in the repressive mark H3K9me2 [85]. Also, suggesting a link between different histone PTMs, mutations in the histone deacetylase HDA6 partially suppressed the H3K4me3 increase observed in response to stress, while the H3K9me2 decrease was not affected. Since histone acetylation and H3K4 trimethylation are often associated with gene activation [86], the repressive function of HDA6 may

suggest that acetylation on histone H3 is required for K4 methylation to occur [87]. Unfortunately, because ChIP experiments were not normalised against the total H3 density, it is difficult to attribute the decrease in H3K9me2 to either the active removal of methyl groups by a histone demethylases or the active removal of nucleosome in response to a stress. Supporting the second possibility, the nucleosome density of two other drought stress-inducible genes (*RD20* and *RAP2.4*) was found to gradually decrease in response to drought stress [74].

In soya bean and in response to a high NaCl concentration, these histone methylation marks were also found altered at some salinity-induced transcription factors (i.e. MYB, b-ZIP and AP2/DREB family members) that were primarily identified by microarray analysis [88]. For some genes, their transcriptional induction was correlated with an increased level of histone acetylation and H3K4me3, accompanied or not with a reduced level of DNA methylation and H3K9me2 in various parts of the promoter or coding regions. For other genes, DNA methylation had no influence on histone methylation. This work perfectly reflects the heterogeneity of the effect of salinity on histone methylation and DNA methylation, and supports the role(s) of histone methylation changes in the expression of some transcription factors important for salinity tolerance.

As mentioned above, H3K4me3 was found to be involved in the transcriptional induction of stress-responsive gene upon salt stress exposure. In plants, the JmjC-domain-containing histone demethylases JMJ14, JMJ15 and JMJ18 have been reported to display an H3K4me2/3 demethylase activity as well as to regulate diverse aspects of chromatin function and development [89–95]. Recently, the overexpression of *JMJ15* was reported to preferentially down-regulate many stress-related genes preferentially marked by H3K4me2/3 and to enhance salt stress tolerance [96]. In contrast, the loss-of-function mutant was more sensitive to salt. Despite the fact that an increased JMJ15 level may regulate stress-responsive gene transcription programmes in *Arabidopsis*, the role of H3K4me3 resetting in these processes is still very elusive.

Besides histone lysine methylation, arginine methylation was also involved in establishing the transcriptional response to salt stress. The protein arginine methyltransferase 5 (PRMT5), also named Shk1 kinase-binding protein1 (SKB1), is a type II methyltransferase that catalyses symmetric H4R3 dimethylation, a repressive mark known to promote flowering through the repression of the floral repressor *FLOWERING LOCUS C (FLC)* in *Arabidopsis* [20, 97]. Interestingly, gain-of-function *SKB1* mutants showed an enhanced salt stress tolerance and sensitivity to ABA [98]. As a consequence of PRMT5 disassociating from chromatin, the H4R3me2 level at stress-responsive genes was reduced during salt stress, resulting in their induced expression. Suggesting an additional function of PRMT5 on non-histone proteins, the methylation level of the U6 small nuclear ribonucleoprotein Sm-like4 (LSM4, a core protein of the spliceosome) was increased in response to salt stress and ABA. Since splicing defects were observed in the *prmt5/skb1* and *lsm4* mutants, with both of them being hypersensitive to salt stress, authors proposed that PRMT5 might mediate plant development and salt response by altering the methylation status of H4R3me2 and LSM4, linking transcriptional regulation to pre-mRNA splicing [98].

4.4. Histone methylation/demethylation in response to temperature

In plants, temperature stresses are classically classified into different types according to temperature exposure, which may be warm, high, chilling or freezing temperature. Due to global warming and because temperature stress greatly affects plant growth and development, immunity and circadian rhythm, and poses a serious threat to the global food supply, the genetic mechanisms of plant responses to heat have been well studied. Plants exposed to temperature stresses modulate the transcription of a large number of genes involved in distinct biochemical and physiological response pathways and networks of phytohormones or secondary metabolites, ultimately leading to increased tolerance to hazardous temperature stresses [99–102]. The role played by histone methylation during the plant response to a heat or a cold stress is discussed separately hereafter.

4.4.1. Heat stress

Heat stress during seed development decreases the seed size in many cereals, resulting in severe yield losses [103, 104]. In rice, a molecular mechanism involving the putative rice polycomb repressive complex 2 (PRC2) gene *Fertilization-Independent Endosperm1* (*OsFIE1*) was suggested as being a potential key component involved in regulating the thermal sensitivity of seed enlargement during endosperm development [105]. When developing seeds were exposed to a heat stress, both DNA methylation and H3K9 methylation were reduced on *OsFIE1* resulting in its derepression. Under heat stress, syncytial stage-specific MADS-box genes involved in seed size regulation were precociously repressed, due to the increased deposition of H3K27me3 silencing marks by the PRC2 complex [105]. In the unicellular green alga *Chlamydomonas reinhardtii*, histone modification was also affected by heat stress [106]. Indeed, after heat stress, the level of H3K4me1 was found decreased and the level of histone acetylation increased at promoter regions of active genes compared with inactive regions. As a hypothesis, authors proposed that upon heat stress, the heat shock transcription factor HSF1 might promote chromatin remodelling and RNAPII recruitment for transcription initiation/elongation [106]. Finally, while addressing molecular mechanisms of the response of cotton anthers to high temperature, two jmjC domain-containing genes, putatively involved in histone demethylation, were found significantly repressed during anther development under heat stress [107].

Because euchromatin is gene rich and usually transcriptionally active, investigation about the role of histone methylation in temperature stress acclimation was largely centred on euchromatin-associated coding regions. Focusing on the transcriptionally silent heterochromatin, mainly constituted of repetitive DNA sequences, some works demonstrated the transcriptional activation of normally silent transposable element embedded within heterochromatic regions under stress conditions [108]. Intriguingly, such activation under heat stress can occur without alteration of DNA methylation and with only minor changes in both H3K9me2 and H3K4me3 [109, 110]. In summary, these works suggest that temperature stress-mediated transcription of tandem-repeat elements might play a vital role in the adaptation of plants to temperature stimuli, offering an efficient mechanism by which heat or cold could promote the expression of some stress-responsive genes. Upon activation and when inserted into or very close to a

gene, such transposable elements could interfere with the expression of this gene, giving rise to deleterious mutations, genetic instability or positive contribution to gene regulation and adaptation [111].

4.4.2. Cold stress

The increased tolerance of plants to cold is referred as 'cold acclimation'. Cold acclimation differs from vernalization, as the last one requires a long-term exposure to cold temperatures, while cold acclimation can be achieved in a couple of days under non-freezing low temperatures [112]. Locally, histone methylation changes in cold-responsive genes were addressed in *Arabidopsis*. Upon cold stress, the repressive mark H3K27me3 decreased, in both a histone occupancy-dependent (i.e. arising from the lowering nucleosome density) and -independent (i.e. as the result of the activity of a not yet identified histone demethylase) manner, on the cold-responsive genes, cold-regulated 15A (*COR15A*) and galactinol synthase 3 (*ATGOLS3*; [113]). Interestingly, the decrease in H3K27me3 upon stimulation occurred more gradually than their rapid transcriptional induction, so that their activation may not be inhibited by H3K27me3 itself but rather lead to the removal of H3K27me3. Also, while the transcription of *COR15A* and *ATGOLS3* was completely repressed to the initial level upon returning plants to normal growth conditions, the H3K27me3 decrease was maintained. Given that this decrease does not affect the transcriptional induction of *COR15A* and *ATGOLS3* upon re-exposure to cold temperatures, such chromatin change can so far only be view as a 'reminiscence' of a recent transcriptional activity and not as a stress memory implicated in a gene priming process.

In maize during cold stress, changes in histone modifications, including the heterochromatic marks H3K9me2 and DNA methylation, were assessed through a genome-wide approach [114]. The more detailed analysis of the two knob-associated tandem-repetitive sequences, the 180-bp repeat and the 350-bp repeat termed TR-1, demonstrated that their selectively and transiently cold-activated transcription was correlated with a decreased H3K9me2 and DNA methylation, together with an increased H3K9 acetylation. Such cold-induced transcriptional activation of tandem repeats is selective and transient, and the silencing state is recovered as the treatment continues.

5. Histone methylation as a memory mark of stress

In animals, the formation of memory immune cells after primary antigen recognition confers long-lasting resistance, resulting in an accelerated and a more effective immune response in case of second exposure. Despite the absence of such memory immune cells, plants often acquire a systemic immunity to further infections after a primary localized infection [115]. This requires the accumulation of the plant hormone SA in systemic tissues and is called systemic acquired resistance (reviewed in [116]. The SAR is also associated with gene priming in systemic tissues, in which defence genes will be expressed more rapidly and robustly in case of a second attack [117]. At the transcriptional level, gene expression is primarily influenced by the chromatin structure, which in turn is controlled partly by processes, often referred to

as ‘epigenetic’ processes, which can be transmitted through mitosis and/or inherited through meiosis [118]. Therefore, chromatin remodelling through histone methylation offers a potential mechanism for short-/long-term stress memory within the lifespan of an individual, referred to as somatic memory, and/or across generations, referred to as transgenerational memory.

5.1. Somatic stress memory

In *Arabidopsis*, a priming event, either treatment with BTH or infection with *Pseudomonas syringae* *pv. maculicola*, systematically resulted in an increase in the level of H3K4me2/me3 at defence gene promoters (namely the WRKY transcription factors *WRKY6*, *WRKY29* and *WRKY52*) that are normally found on active genes, while the genes remain inactive [119]. More interestingly, this increase also occurs in leaves distal to localized foliar infection. Hence, even if the histone-modifying enzyme involved in this process remains unidentified, results from Jaskiewicz et al. [119] clearly suggest that histone methylation might create a ‘memory’ of the primary infection that is associated with an amplified reaction to a second stress stimulus. Further, Luna et al. [120] observed that promoters of SA-inducible *PR1*, *WRKY6* and *WRKY53* in the progeny of *Pst*-inoculated *Arabidopsis* plants were enriched with acetylated histone, while the promoter of the JA-inducible gene *PLANT DEFENSIN1.2* (*PDF1.2*) showed an increased level of H3K27me3. For decades, the signalling protein *NON EXPRESSOR OF PR1* (*NPR1*) has been implicated in mediating SAR induction [115] and also the crosstalk between SA- and JA/ET-dependent defence pathways, enabling plants to mount an appropriate defence reaction, depending on the nature of the attacker and the stage of infection [121, 122]. More recently, *NPR1* has been proposed to play a critical role in the expression of the transgenerational SAR as progeny from *npr1* failed to develop transgenerational defence phenotypes and failed to present enrichment for H3K27me3 at the *PDF1.2* promoter [120]. Together, these findings suggested that one or more systemic signals are stored as an immune memory on defence-related gene promoters in the form of histone modifications, thus providing the plant with a life-long protection, which can be transmitted to subsequent generations.

Besides being involved in defence priming related to biotic stress, histone methylation was also proposed as a priming strategy against drought. To further explore the functional impact of histone methylation on biotic stress responses in *Arabidopsis*, Kim et al. [123] followed chromatin dynamics of several drought genes (*RD20*, *RD29A* and *AtGOLS2*) and a rehydration-inducible gene (*ProDH*) during drought and rehydration. As previously discussed, a strong correlation was observed between H3K4me3 enrichment (i.e. especially in gene bodies) and transcription for drought genes upon drought [74, 123]. Such a correlation was also detected for the rehydration gene upon rehydration [123]. Suggesting a memory role for H3K4me3, RNAPII rapidly disappeared after rehydration at drought genes, while H3K4me3 was gradually decreased. Concomitantly, by training plants with up to four successive drought treatments, Ding et al. [124] uncovered the existence of two distinct subsets of genes within the dehydration stress–response gene fraction. The ‘non-trainable’ genes (e.g. the ABA-independent *RD29A* and *COR15A*) have repetitively similar transcription rates during each stress treatment, while ‘trainable’ ones (e.g. the ABA-dependent *RD29B* and *RAB18*) increased

the magnitude of their subsequent transcriptional response, relative to their initial stress response. Using ChIP, Ding et al. [124] observed that the H3K4me3 enrichment at 'trainable' genes, especially in gene bodies, was atypically retained from the preceding transcription after rehydration. Even more interestingly, the RNAPII phosphorylated at C-terminal domain (CTD) repeat serine 5 (Ser5P; associated with transcription initiation) was found stalled on these genes as a memory mark from a previously transcribed state. In contrast to 'trainable' genes, the stress-induced H3K4me3 and Ser5P enrichment at 'non-trainable' genes was decreased to its basal level during recovery [124]. Moreover, this transcriptional memory can persist in the absence of inducing signals at least for 5 days, but is lost after 7 days. Supporting a specific role for H3K4me3 in stress memory, other active chromatin marks such as acetylation of histones H3 and H4 were found rapidly increased at drought genes upon stress and decreased at comparable levels as before induction quickly after recovery [123, 124]. Consistent with an activating role of ATX1 at dehydration stress-response genes [68, 125, 126], dehydration-induced transcript levels were diminished in *atx1* plants [127]. However, ATX1 does not seem to have a critical impact on drought stress memory in *Arabidopsis*. Indeed, while being less increased than in wild-type plants, trainable genes still produced increased transcripts in trained, relative to untrained, *atx1* plants and retained high H3K4me3 levels during the watered recovery states. ATX1, ATX2, SDG25 and SET DOMAIN GROUP2/ATX-RELATED3 (SDG2/ATXR3) belong to the same class III of H3K4me3 methyltransferases and are thought to act, partially redundantly, as H3K4 methyltransferase [128–134]. More recently, SDG2 has been found to be essential for the full transcriptional activation of various hormone-responsive genes upon hormone treatment (i.e. including the ABA-dependant *RD29A*) via its H3K4 trimethyltransferase activity [135]. It is, therefore, likely that the other class III HKMT might also contribute to the drought stress memory. In summary, these results suggest that in addition to be a good marker of gene activation when found around promoter and 5' regions of genes, H3K4me3 might also play a role in establishing a transcriptional short-term somatic memory of drought stress when found in gene bodies.

Using a large-scale approach, the distribution of H3K4me2, H3K4me3, H3K9me2 and H3K27me3 was analysed in *Arabidopsis* seedlings, which have been treated with mild salt stress in the seedling stage, resulting in an increased tolerance upon an additional salt stress application [136]. At low resolution in primed seedlings, H3K4me2 and H3K4me3 most commonly consisted of higher peaks of pre-existing enriched histone modification domains, named islands, whereas H3K9me2 produced the least differences. By contrast, majority of differences in H3K27me3 resulted from a higher number of islands with lower genome coverage. At high resolution, changes in H3K27me3 were already detectable a few hours after salt addition, suggesting that demethylation of H3K27me3 operates at a speed that is comparable to that of transcriptional regulation. Interestingly, this effect fades over time; however, it is still clearly visible after a 10-day-growth period in control conditions. In response to a second stress treatment, genes with high responsiveness, such as *HKT1* (i.e. encoding a root-specific Na transporter) and *PIP2E* (i.e. encoding a plasma membrane aquaporin), experienced a decrease of H3K27me3, whereas genes with lower responsiveness, such as *GH3.1* and *GH3.3* (i.e. encoding auxin and JA-amino acid-conjugating enzymes, respectively), experienced an

increase of H3K27me₃. Conversely, another group reported that in plants that have experienced several exposures to dehydration stress no significant change in the level of H3K27me₃ could be detected on trainable and non-trainable genes, or during transcriptionally active/inactive gene states [137]. However, the high H3K27me₃ level present at inactive dehydration stress memory genes did not interfere with the transition to an active transcription and with the accumulation of H3K4me₃ [138]. Together, the function of H3K27me₃ in genes that dynamically change transcription seems to depend on the type of environmental stimuli.

In contrast to H3K27me₃, the higher level of H3K4me₃ retained at the trainable gene *RD29B*, when its transcription is low, further supports the idea that H3K4me₃ works as a 'memory' histone mark of a previously active state [138]. Generally, H3K4me₃ and H3K27me₃ play antagonistic roles in gene transcription and are therefore mutually repulsive at developmental genes [37, 139]. Seemingly, the presence of both marks, referred as 'bivalent domains', was first described in mammalian stem cells and was proposed to represent a pluripotent chromatin state that poises genes for activation upon appropriate developmental cues [140, 141]. Further work is required to determine whether H3K4me₃ and H3K27me₃ co-exist at certain genes whose expression is rapidly altered in response to environmental stimuli.

Because plant stress research has traditionally focused on single stresses, we separately described priming in response to abiotic or biotic environmental cues. However, in nature, plants are constantly exposed to mild environmental stresses during their lifetime. While testing how different environmental histories can affect the response of the plant to a subsequent biotic stress, Singh et al. [142] reported that *Arabidopsis* plants exposed to a recurrent abiotic stress (i.e. heat, cold or salt) were more resistant to *Pst* than plants grown in a more stable environment. This enhanced resistance was due to the priming of commonly used marker genes of pattern-triggered immunity (PTI; *WRKY53*, *FLG22-INDUCED RECEPTOR KINASE1* (*FRK1*) and *NDR1/HIN1-LIKE10* (*NHL10*)). Indeed, enrichment for epigenetic marks associated with transcriptional activation, such as H3K4me₂ and H3K4me₃, at PTI-responsive genes was observed after the exposure to recurrent stress, resulting in an enrichment of RNA polymerase II and a primed transcription in response to a subsequent bacterial infection. Collectively, these works on somatic stress memory promote the idea that, in plant, the environmental history can shape/modulate the response to stress, providing a mechanistic link between histone methylation and gene priming.

5.2. Transgenerational stress memory

The transgenerational stress memory refers to the transmittance of certain environmental responses from one generation to the next, thus providing the offspring of environmentally challenged plants with an adaptive advantage for better fitness (i.e. improve plant stress tolerance and impart developmental flexibility; [143]). Compared with DNA methylation and RNA interference (RNAi), very few studies suggest the involvement of histone methyltransferases and histone methylation changes in this process [144]. In *Arabidopsis*, changes in DNA methylation, histone modifications and gene expression were followed in the progeny of plants exposed to salt stress over one generation [145]. Although the DNA from the progeny of plants exposed to salt stress was globally hypomethylated, the majority of genes and promoters

causing methylation changes were hypermethylated and lowly expressed. In addition, DNA hypermethylation was correlated with an increased level of the repressive mark H3K9me2. Among these hypermethylated genes, a large number was encoding different histone methyltransferases, such as the *Arabidopsis* *SU(VAR)3-9* homologues *SUVH2*, *SUVH5*, *SUVH8*, involved in H3K9 methylation, or the PRC2 subunit *CURLY LEAF (CLF)*, involved in H3K27 methylation. Following this work, the progeny of heat-stressed plants was used to explore epigenetic variations under both normal and stressed conditions, in comparison to the progeny of control plants [146]. Similarly to salt stress, the progeny of plants exposed to heat stress had a global decrease of genomic DNA methylation and a reduced expression of several *SUVH* genes, which correlated with their enrichment in H3K9me2. Together, the hypermethylation of *SUVH* genes in the progeny of stressed plants may represent a protective mechanism against hypermethylation of the entire genome. Interestingly, in both works [145, 146], the transposon expression was elevated in the progeny of stressed plants. Because main targets of the *SUVH* pathway are transposable elements [24], the authors proposed that a decrease in the expression of *SUVH* genes might contribute to transposon activation, which at opportune times can create intragenomic potential upon transposition to facilitate adaptation in response to environmental changes [147]. In summary, these works suggested a role for histone methylation in the inheritance of stress memory; however, whether histone methylation changes are heritable through multiple generations and whether they sustain the acquisition of adaptive traits is still a matter of debate [148]. However, the evidence to date favours the view that stress-induced transgenerational changes in chromatin might increase the survival chances of the plant species, rather than each individual, by broadening the phenotypic plasticity and the genetic variation within the population [149, 150].

6. Discussion and perspectives

Recent advances, especially in *Arabidopsis*, have uncovered that chromatin remodelling through histone methylation changes are not only restricted to developmental needs but also an integral part of the very complex cascade of events that lead to abiotic/biotic stress tolerance, resistance and short-/long-term memory. Currently, a preliminary view is emerging, indicating that histone methylation changes, providing specific chromatin configurations, can be classified into several interrelated categories when involved in stress responses (Fig. 1): (i) histone methylation changes that are basally present on stress-related genes to establish a 'permissive' chromatin state that may either limit the spreading of repressive chromatin marks and/or potentiate a rapid transcriptional induction upon need; (ii) histone methylation changes that are transiently induced from an inactive or a permissive chromatin state by stress, to either facilitate the transcriptional initiation and/or reinforce transcription of stress-responding genes, and finally, histone methylation changes that are established in response to a stress; (iii) maintained for a certain time during the lifespan of an individual (i.e. somatic memory) or (iv) transmitted to one or more subsequent generations (i.e. transgenerational memory).

Figure 1. Hypothetical model depicting the role of histone methylation/demethylation in regulating plant stress responses. (A) Plants exposed either to biotic or abiotic stresses integrate the signal stress into the cell nuclei, where it affects the chromatin structure through histone methylation changes. According to our knowledge, these histone methylation changes can be classified into several interrelated categories. (B) The permissive state represents a more loosened chromatin state that will either offer a protection against repressive marks (represented with nucleosomes in red) and/or potentiates a rapid transcriptional induction upon stress induction. (C) The induced state represents histone methylation changes that are transiently induced by a stress signal. If methylation changes occur early during the stress-response process, they might participate to the transcriptional induction of stress-responding genes, while if they occur later, they might reinforce the transcription of stress-responding genes. Both permissive and the induced states can be maintained allowing a faster and/or stronger transcriptional induction of stress-responding genes upon a subsequent challenge. The memorized chromatin state can be maintained (D) for a certain time during the lifespan of an individual and referred as the somatic memory, or (E) transmitted to one or more subsequent generations and referred as the transgenerational memory.

Nonetheless, this emerging view is facing many gaps, inaccuracies and divergences, mainly related to numerous difficulties inherent to the study of such a dynamic and acute process. In this respect, plants in nature are usually challenged simultaneously by different kinds of stresses. Responses to these stress combinations are largely controlled by different signalling pathways that can interact in a non-additive manner, producing effects that could not have been predicted from the study of either stress individually [151, 152]. The occurrence of simultaneous biotic and abiotic stresses introduces an added degree of complexity that requires stresses to be imposed simultaneously and to treat each set of environmental conditions as an entirely new stress. For this reason and to clarify the mechanism behind the regulation of stress responses by histone methylation changes, there is a strong necessity to intensify our investigations. For instance, the correlation between histone methylation/demethylation and stress responses remains elusive and clarifications will require in-depth dynamic approaches based on comparative analyses of both epigenomes and transcriptomes during stress responses. In parallel, current knowledge about the corresponding histone-modifying enzymes is still largely missing. This lack of knowledge is pending on the identification of different stress-responsive histone modifiers and will require large-scale screens and genetic analyses for the sensitivity of different histone methyltransferases/demethylases mutants to various stresses, combined or not. Among other factors governing stress-induced chromatin changes, almost nothing is known about the specific reader/effector that will recognize particular histone methylation sites in order to determine their functional and structural outcome. An effort in this direction will most likely benefit the comprehensive understanding of the fundamental mechanisms connecting histone methylation changes with the modulation of transcription of stress-responsive genes, subsequently enabling plant to withstand stress. Higher-resolution chromatin studies are undoubtedly required to reveal the targeted stress-responsive genes and the specific sites of histone methylation/demethylation. Nevertheless, investigation of the direct effects of histone methylation/demethylation in plants is difficult. One reason is that plant genomes harbour high-copy number of histone genes (e.g. the *Arabidopsis* genome comprises 47 genes that encode 33 different core histone proteins; www.chromdb.org) and the incorporation/modification of such variants can result in the formation of chromatin with particular properties and functions [153–155]. Although ChIP assays have proven valuable in helping to identify histone methylation changes, many antibodies used to detect these changes have been so far unable to distinguish between different variants. New technologies (e.g. generation of mutants with point mutations targeting amino acid in the N-terminal tail of histone using the clustered regularly interspaced short palindromic repeats (CRISPR)/CRISPER-associated (Cas) system) [156] will need to be explored to unravel histone methylation changes of specific histone variants and their functions. Another challenge is that plants consist of many functionally specialized tissues and cell types, each with its own unique epigenome, transcriptome and proteome. Until now, histone methylation changes induced by stresses were exclusively addressed in entire plant or organs, meaning that the obtained profiles most likely reflect the consensus of multiple tissue- or cell-specific profiles that may differ. New methods allowing the mapping of chromatin features in specific tissue/cell types such as the one described by Wang and Deal [157] will be decisive for determining the cell-/tissue-specific chromatin alterations involved

in a particular stress response. Finally, as plants have finite resources that must be balanced between growth and defence against stresses, often resulting in a growth or yield penalty, histone methylation changes in response to stress should be integrated in a more global developmental view, taking into account the involvement of several histone methyltransferases/demethylases in various processes such as root growth, flowering time, floral organogenesis, gametophyte or embryo formation [33]. Finally, understanding such regulatory network is an essential step to provide both novel paradigms and potential tools for further exploitation towards sustainable agriculture.

Acknowledgements

This work was supported by the Centre National de la Recherche Scientifique, the Agence Nationale de la Recherche (ANR-12-BSV2-0013-02), the European Commission (FP7-PEOPLE-2013-ITN, grant number 607880), the National Council of Science and Technology of Mexico (CONACYT) and the Program 'Estancias Posdoctorales y Sabaticas al Extranjero' (Fellowship number 237557. BOBADILLA-LANDEY Roberto; CVU:162391).

Author details

Roberto Bobadilla and Alexandre Berr*

*Address all correspondence to: Alexandre.Berr@ibmp-cnrs.unistra.fr

Institut de Biologie Moléculaire des Plantes, CNRS UPR2357, Université de Strasbourg, Strasbourg, France

References

- [1] Osakabe Y, Yamaguchi-Shinozaki K, Shinozaki K and Tran LS. (2013) Sensing the environment: key roles of membrane-localized kinases in plant perception and response to abiotic stress. *J Exp Bot.* 64(2):445–458.
- [2] Pieterse CMJ, Leon-Reyes A, Van der Ent S and Van Wees SCM. (2009) Networking by small-molecule hormones in plant immunity. *Nat Chem Biol.* 5(5):308–316.
- [3] Nakashima K, Ito Y and Yamaguchi-Shinozaki K. (2009) Transcriptional regulatory networks in response to abiotic stresses in *Arabidopsis* and grasses. *Plant Physiol.* 149(1):88–95.
- [4] van Verk MC, Gatz C and Linthorst HJM. (2009) Transcriptional regulation of plant defense responses. *Adv Bot Res.* 51:397–438.

- [5] Riechmann JL and Ratcliffe OJ. (2000) A genomic perspective on plant transcription factors. *Curr Opin Plant Biol.* 3(5):423–434.
- [6] Moore JW, Loake GJ and Spoel SH. (2011) Transcription dynamics in plant immunity. *Plant Cell.* 23(8):2809–2820.
- [7] Luger K, Mäder AW, Richmond RK, Sargent DF and Richmond TJ. (1997) Crystal structure of the nucleosome core particle at 2.8 Å resolution. *Nature.* 389(6648):251–260.
- [8] Andrews AJ and Luger K. (2011) Nucleosome structure(s) and stability: variations on a theme. *Annu Rev Biophys.* 40:99–117.
- [9] Nelissen H, Boccardi TM, Himanen K and Van Lijsebettens M. (2007) Impact of core histone modifications on transcriptional regulation and plant growth. *Crit Rev Plant Sci.* 26(5-6):243–263.
- [10] Shen WH and Xu L. (2009) Chromatin remodeling in stem cell maintenance in *Arabidopsis thaliana*. *Mol Plant.* 2(4):600–609.
- [11] Berr A and Shen WH. (2010) Molecular mechanisms in epigenetic regulation of plant growth and development. In: Pua E. C. and Davey M. R. (Eds.) *Plant Dev Biol – Biotechnol Perspect.* 2:325–344.
- [12] Arents G and Moudrianakis EN. (1995) The histone fold: a ubiquitous architectural motif utilized in DNA compaction and protein dimerization. *Proc Natl Acad Sci USA.* 92(24):11170–11174.
- [13] Davey CA, Sargent DF, Luger K, Maeder AW and Richmond TJ. (2002) Solvent mediated interactions in the structure of the nucleosome core particle at 1.9 Å resolution. *J Mol Biol.* 319(5):1097–1113.
- [14] Marks P, Rifkind RA, Richon VM, Breslow R, Miller T and Kelly WK. (2001) Histone deacetylases and cancer: causes and therapies. *Nat Rev Cancer.* 1(3):194–202.
- [15] Mutskov V, Gerber D, Angelov D, Ausio J, Workman J and Dimitrov S. (1998) Persistent interactions of core histone tails with nucleosomal DNA following acetylation and transcription factor binding. *Mol Cell Biol.* 18(11):6293–6304.
- [16] Strahl BD and Allis CD. (2000) The language of covalent histone modifications. *Nature.* 403(6765):41–45.
- [17] Jaskiewicz M, Peterhansel C and Conrath U. (2011) Detection of histone modifications in plant leaves. *J Vis Exp.* (55).
- [18] Zhang X. 2012. Chromatin modifications in plants. In: Wendel JF, editor. *Plant genome diversity*. Vienna: Springer. p. 237–56.

- [19] Guccione E, Bassi C, Casadio F, Martinato F, Cesaroni M, Schuchlantz H, Lüscher Band Amati B. (2007) Methylation of histone H3R2 by PRMT6 and H3K4 by an MLL complex are mutually exclusive. *Nature*. 449(7164):933–937.
- [20] Wang X, Zhang Y, Ma Q, Zhang Z, Xue Y, Bao S and Chong K. (2007) SKB1-mediated symmetric dimethylation of histone H4R3 controls flowering time in *Arabidopsis*. *EMBO J*. 26(7):1934–1941.
- [21] Zhao Q, Rank G, Tan YT, Li H, Moritz RL, Simpson RJ, Cerruti L, Curtis DJ, Patel DJ, Allis CD, Cunningham JM and Jane SM. (2009) PRMT5-mediated methylation of histone H4R3 recruits DNMT3A, coupling histone and DNA methylation in gene silencing. *Nat Struct Mol Biol*. 16(3):304–311.
- [22] Wang H, Huang ZQ, Xia L, Feng Q, Erdjument-Bromage H, Strahl BD, Briggs SD, Allis CD, Wong J, Tempst P and Zhang Y. (2001) Methylation of histone H4 at arginine 3 facilitating transcriptional activation by nuclear hormone receptor. *Science*. 293(5531):853–857.
- [23] Strahl BD, Briggs SD, Brame CJ, Caldwell JA, Koh SS, Ma H, Cook RG, Shabanowitz J, Hunt DF, Stallcup MR and Allis CD. (2001) Methylation of histone H4 at arginine 3 occurs in vivo and is mediated by the nuclear receptor coactivator PRMT1. *Curr Biol*. 11(12):996–1000.
- [24] Thorstensen T, Grini PE and Aalen RB. (2011) SET domain proteins in plant development. *Biochim Biophys Acta*. 1809(8):407–420.
- [25] Di Lorenzo A and Bedford MT. (2011) Histone arginine methylation. *FEBS Lett*. 585(13):2024–2031.
- [26] Pedersen MT and Helin K. (2010) Histone demethylases in development and disease. *Trends in Cell Biol*. 20(11):662–671.
- [27] Springer NM, Napoli CA, Selinger DA, Pandey R, Cone KC, Chandler VL, Kaeppler HF and Kaeppler SM. (2003) Comparative analysis of SET domain proteins in maize and *Arabidopsis* reveals multiple duplications preceding the divergence of monocots and dicots. *Plant Physiol*. 132(2):907–925.
- [28] Ng DW, Wang T, Chandrasekharan MB, Aramayo R, Kertbundit S and Hall TC. (2007) Plant SET domain-containing proteins: structure, function and regulation. *Biochim Biophys Acta*. 1769(5-6):316–329.
- [29] Pontvianne F, Blevins T and Pikaard CS. (2010) *Arabidopsis* histone lysine methyltransferases. In: Delseny M. And Kader J. C. (eds) *Advances in botanical research*. London: Academic Press Ltd-Elsevier Science Ltd. 1–22.
- [30] Aquea F, Vega A, Timmermann T, Poupin MJ and Arce-Johnson P. (2011) Genome-wide analysis of the SET DOMAIN GROUP family in grapevine. *Plant Cell Rep*. 30(6):1087–1097.

- [31] Huang Y, Liu C, Shen WH and Ruan Y. (2011) Phylogenetic analysis and classification of the *Brassica rapa* SET-domain protein family. *BMC Plant Biol.* 11:175.
- [32] Aiese-Cigliano R, Sanseverino W, Cremona G, Ercolano MR, Conicella C and Consiglio FM. (2013) Genome-wide analysis of histone modifiers in tomato: gaining an insight into their developmental roles. *BMC Genomics.* 14:57.
- [33] Berr A, Shafiq S and Shen WH. (2011) Histone modifications in transcriptional activation during plant development. *Biochim Biophys Acta.* 1809(10):567–576.
- [34] Hammond-Kosack KE and Jones JDG. (2000) Response to plant pathogens. In: Buchannan B. Gruissem W. Jones R. (eds). *Biochemistry and molecular biology of plants*. Rockville, MD: American Society of Plant Physiologists. 1102–1157.
- [35] Jones JDG and Dangl JL. (2006) The plant immune system. *Nature.* 444(7117):323–329.
- [36] Mosher RA, Durrant WE, Wang D, Song J and Dong X. (2006) A comprehensive structure-function analysis of *Arabidopsis* SNI1 defines essential regions and transcriptional repressor activity. *Plant Cell.* 18(7):1750–1765.
- [37] Roudier F, Ahmed I, Bérard C, Sarazin A, Mary-Huard T, Cortijo S, Bouyer D, Cailieux E, Duvernois-Berthet E, Al-Shikhley L, Giraut L, Després B, Drevensek S, Barneche F, Dèrozier S, Brunaud V, Aubourg S, Schnittger A, Bowler C, Martin-Magniette M L, Robin S, Caboche M and Colot V. (2011). Integrative epigenomic mapping defines four main chromatin states in *Arabidopsis*. *EMBO J.* 30:1928–1938.
- [38] Alvarez-Venegas R, Al Abdallat A, Guo M, Alfano JR and Avramova Z (2007). Epigenetic control of a transcription factor at the cross section of two antagonistic pathways. *Epigenetics.* 2(2):106–113.
- [39] Gruner K, Griebel T, Návarová H, Attaran E and Zeier J. (2013) Reprogramming of plants during systemic acquired resistance. *Front Plant Sci.* 4:252.
- [40] Alvarez-Venegas R, Sadler M, Hlavacka A, Baluska F, Xia YN, Lu GQ, et al. (2006) The *Arabidopsis* homolog of trithorax, ATX1, binds phosphatidylinositol 5-phosphate, and the two regulate a common set of target genes. *Proc Natl Acad Sci USA.* 103(15): 6049–6054.
- [41] Li J, Brader G, Kariola T and Palva ET. (2006) WRKY70 modulates the selection of signaling pathways in plant defense. *Plant J.* 46(3):477–491.
- [42] Xu L, Zhao Z, Dong A, Soubigou-Taconnat L, Renou JP, Steinmetz A and Shen WH. (2008) Di- and tri- but not monomethylation on histone H3 lysine 36 marks active transcription of genes involved in flowering time regulation and other processes in *Arabidopsis thaliana*. *Mol Cell Biol.* 28(4):1348–1360.

- [43] Palma K, Thorgrimsen S, Malinovsky FG, Fiil BK, Nielsen HB, Brodersen P, et al. (2010) Autoimmunity in *Arabidopsis acd11* Is Mediated by Epigenetic Regulation of an Immune Receptor. *PLoS Pathog* 6(10): e1001137. doi:10.1371/journal.ppat.1001137.
- [44] De-La-Peña C, Rangel-Cano A and Alvarez-Venegas R. (2012) Regulation of disease-responsive genes mediated by epigenetic factors: interaction of *Arabidopsis-Pseudomonas*. *Mol Plant Pathol*. 13(4):388–398.
- [45] Tsuchiya T and Eulgem T. (2011) EMSY-like genes are required for full RPP7-mediated race-specific immunity and basal defense in *Arabidopsis*. *Mol Plant Microbe Interact*. 24(12):1573–1581.
- [46] Tsuchiya T and Eulgem T. (2013) An alternative polyadenylation mechanism coopted to the *Arabidopsis* RPP7 gene through intronic retrotransposon domestication. *Proc Natl Acad Sci U S A*. 110(37):E3535-E3543.
- [47] Musselman CA and Kutateladze TG. (2011) Handpicking epigenetic marks with PHD fingers. *Nucleic Acids Res*. 39(21):9061–9071.
- [48] Tsuchiya T and Eulgem T. (2013) Mutations in EDM2 selectively affect silencing states of transposons and induce plant developmental plasticity. *Sci Rep*. 3:1701.
- [49] Li T, Chen X, Zhong X, Zhao Y, Liu X, Zhou S, Cheng S and Zhou DX. (2013) Jumonji C domain protein JM705-mediated removal of histone H3 lysine 27 trimethylation is involved in defense-related gene activation in rice. *Plant Cell*. 25(11):4725–4736.
- [50] Pieterse CM, Poelman EH, Van Wees SC and Dicke M. (2013) Induced plant responses to microbes and insects. *Front Plant Sci*. 4:475.
- [51] Berr A, McCallum EJ, Alioua A, Heintz D, Heitz T and Shen WH. (2010). *Arabidopsis* histone methyltransferase SET DOMAIN GROUP8 mediates induction of the jasmonate/ethylene pathway genes in plant defense response to necrotrophic fungi. *Plant Physiol*. 154(3):1403–1414.
- [52] Berr A, Ménard R, Heitz T and Shen WH. (2012) Chromatin modification and remodelling: a regulatory landscape for the control of *Arabidopsis* defence responses upon pathogen attack. *Cell Microbiol*. 14(6):829–839.
- [53] Grossniklaus U and Paro R. (2014) Transcriptional silencing by polycomb-group proteins. *Cold Spring Harb Perspect Biol*. 6(11):a019331.
- [54] Wang W, Vinocur B and Altman A. (2007) Plant responses to drought, salinity and extreme temperatures towards genetic engineering for stress tolerance. *Planta*. 218(1):1–14.
- [55] Gao JP, Chao DY and Lin HX. (2007) Understanding abiotic stress tolerance mechanisms: recent studies on stress response in rice. *J Integr Plant Biol* 49(6):742–750.
- [56] Mittler R. (2006) Abiotic stress, the field environment and stress combination.” *Trends Plant Sci*. 11(1):15–19.

- [57] Santner A, Calderon-Villalobos LI and Estelle M. (2009) Plant hormones are versatile chemical regulators of plant growth. *Nat Chem Biol.* 5(5):301–307
- [58] Cutler SR, Rodriguez PL, Finkelstein RR and Abrams SR. (2010) Absciscic acid: emergence of a core signaling network. *Annu Rev Plant Biol.* 61:651–679.
- [59] Fujita M, Fujita Y, Noutoshi Y, Takahashi F, Narusaka Y, Yamaguchi-Shinozaki K and Shinozaki K. (2006) Crosstalk between abiotic and biotic stress responses: a current view from the points of convergence in the stress signaling networks. *Curr Opin Plant Biol.* 9(4):436–442.
- [60] Asselbergh B, De Vleeschauwer D and Hofte M. (2008) Global switches and fine-tuning – ABA modulates plant pathogen defense. *Mol Plant Microbe Interact.* 21(6): 709–719.
- [61] Yasuda M, Ishikawa A, Jikumaru Y, Seki M, Umezawa T, Asami T, Maruyama-Nakashita A, Kudo T, Shinozaki K, Yoshida S and Nakashita H. (2008) Antagonistic interaction between systemic acquired resistance and the abscisic acid-mediated abiotic stress response in *Arabidopsis*. *Plant Cell.* 20(6):1678–1692.
- [62] Atkinson NJ and Urwin PE. (2012) The interaction of plant biotic and abiotic stresses: from genes to the field. *J Exp Bot.* 63(10):3523–3543.
- [63] Rajjou L, Duval M, Gallardo K, Catusse J, Bally J, Job C and Job D. (2012) Seed germination and vigor. *Annu Rev Plant Biol.* 63:507–533.
- [64] Lopez-Molina L, Mongrand S and Chua NH. (2001) A postgermination developmental arrest checkpoint is mediated by abscisic acid and requires the ABI5 transcription factor in *Arabidopsis*. *Proc Natl Acad Sci USA.* 98(8):4782–4787.
- [65] Perruc E, Kinoshita N, Lopez-Molina L. (2007). The role of chromatin-remodeling factor PKL in balancing osmotic stress responses during *Arabidopsis* seed germination. *Plant J.* 52(5):927–936.
- [66] Raghavendra AS, Gonugunta VK, Christmann A and Grill E. (2010) ABA perception and signalling. *Trends Plant Sci.* 15(7):395–401.
- [67] Koornneef M, Leon-Kloosterziel K, Schwartz SH and Zeevaart JAD. (1998) The genetic and molecular dissection of abscisic acid biosynthesis and signal transduction in *Arabidopsis*. *Plant Physiol Biochem.* 36:83–89.
- [68] Ding Y, Avramova Z and Fromm M. (2011) The *Arabidopsis* trithorax-like factor ATX1 functions in dehydration stress responses via ABA-dependent and ABA-independent pathways. *Plant J.* 66(5):735–744.
- [69] Ramachandra RA, Chaitanya KV and Vivekanandan M. (2004) Drought induced responses of photosynthesis and antioxidant metabolism in higher plants. *J Plant Physiol.* 161(11):1189–1202.

- [70] Araus JL, Slafer GA, Royo C and Serret MD. (2008) Breeding for yield potential and stress adaptation in cereals. *Crit Rev Plant Sci.* 27:377–412.
- [71] Hirayama T and Shinozaki K. (2010) Research on plant abiotic stress responses in the post-genome era: past, present and future. *Plant J.* 61(6):1041–1052.
- [72] Fukao T and Xiong L. (2013) Genetic mechanisms conferring adaptation to submergence and drought in rice: simple or complex? *Curr Opin Plant Biol.* 16(2):196–204.
- [73] Huang D, Wu W, Abrams SR and Cutler AJ. (2008) The relationship of drought-related gene expression in *Arabidopsis thaliana* to hormonal and environmental factors. *J Exp Bot.* 59(11):2991–3007.
- [74] Kim JM, To TK, Ishida J, Morosawa T, Kawashima M, Matsui A, Toyoda T, Kimura H, Shinozaki K and Seki M. (2008). Alterations of lysine modifications on the histone H3 N-tail under drought stress conditions in *Arabidopsis thaliana*. *Plant Cell Physiol.* 49:1580–1588.
- [75] Kim JM, Sasaki T, Ueda M, Sako K and Seki M. (2015) Chromatin changes in response to drought, salinity, heat, and cold stresses in plants. *Front Plant Sci.* 6:114.
- [76] van Dijk K, Ding Y, Malkaram S, Riethoven JJ, Liu R, Yang J, Laczko P, Chen H, Xia Y, Ladunga I, Avramova Z and Fromm M. (2010) Dynamic changes in genome-wide histone H3 lysine 4 methylation patterns in response to dehydration stress in *Arabidopsis thaliana*. *BMC Plant Biol.* 10:238.
- [77] Papaefthimiou D and Tsaftaris A. (2012) Significant induction by drought of HvPKDM7-1, a gene encoding a jumonji-like histone demethylase homologue in barley (*H. Vulgare*). *Acta Physiol Plant.* 34(3):1187–1198.
- [78] Govind G, Vokkaliga Thammegowda H, Jayaker Kalaiarasi P, Iyer D, Muthappa S, Nese S and Makarla U. (2009) Identification and functional validation of a unique set of drought induced genes preferentially expressed in response to gradual water stress in peanut. *Mol Genet Genomics.* 281(6):591–605.
- [79] Rando OJ and Chang HY. (2009) Genome-wide views of chromatin structure. *Annu Rev Biochem.* 78:245–271.
- [80] Zong W, Zhong X, You J and Xiong L. (2013). Genome-wide profiling of histone H3K4-tri-methylation and gene expression in rice under drought stress. *Plant Mol Biol* 81(1-2):175–188.
- [81] Hsu FC and Shih MC. (2013) Plant defense after flooding. *Plant Signal Behav.* 8(11):e26922.
- [82] Tsuji H, Nakazono M, Saisho D, Tsutsumi N and Hirai A. (2000). Transcript levels of the nuclear-encoded respiratory genes in rice decrease by oxygen deprivation: evidence for involvement of calcium in expression of the alternative oxidase 1a gene. *FEBS Lett.* 471(2-3):201–204.

- [83] Tsuji H, Saika H, Tsutsumi N, Hirai A and Nakazono M. (2006) Dynamic and reversible changes in histone H3-Lys4 methylation and H3 acetylation occurring at submergence-inducible genes in rice. *Plant Cell Physiol.* 47(7):995–1003.
- [84] Carillo P, Annunziata MG, Pontecorvo G, Fuggi A and Woodrow P. (2011) Salinity stress and salt tolerance. In: Arun S, editor. *Abiotic Stress in Plants - Mechanisms and Adaptations*. Croatia: InTech. pp. 22–38.
- [85] Chen LT, Luo M, Wang YY and Wu K. (2010). Involvement of *Arabidopsis* histone deacetylase HDA6 in ABA and salt stress response. *J Exp Bot.* 61(2):3345–3353.
- [86] He G, Elling A and Deng X. (2011) The epigenome and plant development. *Rev Plant Biol.* 62(1):411.
- [87] Nightingale KP, Gendreizig S, White DA, Bradbury C, Hollfelder F and Turner BM. (2007) Cross-talk between histone modifications in response to histone deacetylase inhibitors: MLL4 links histone H3 acetylation and histone H3K4 methylation. *J Biol Chem.* 282(7):4408–4416.
- [88] Song Y1, Ji D, Li S, Wang P, Li Q and Xiang F. (2012) The dynamic changes of DNA methylation and histone modifications of salt responsive transcription factor genes in soybean. *PLoS One.* 7(7):e41274.
- [89] Deleris A, Greenberg MV, Ausin I, Law RW, Moissiard G, Schubert D and Jacobsen SE. (2010) Involvement of a Jumonji-C domain-containing histone demethylase in DRM2-mediated maintenance of DNA methylation. *EMBO Rep.* 11(12):950–955.
- [90] Lu F, Cui X, Zhang S, Liu C and Cao X. (2010) JMJ14 is an H3K4 demethylase regulating flowering time in *Arabidopsis*. *Cell Res.* 20(3):387–390.
- [91] Searle IR, Pontes O, Melnyk CW, Smith LM and Baulcombe DC. (2010) JMJ14, a JmjC domain protein, is required for RNA silencing and cell-to-cell movement of an RNA silencing signal in *Arabidopsis*. *Genes Dev.* 24(10):986–991.
- [92] Le Masson I, Jauvion V, Bouteiller N, Rivard M, Elmayan T and Vaucheret H. (2012) Mutations in the *Arabidopsis* H3K4me2/3 demethylase JMJ14 suppress posttranscriptional gene silencing by decreasing transgene transcription. *Plant Cell.* 24(9):3603–3612.
- [93] Yang H, Mo H, Fan D, Cao Y, Cui S and Ma L. (2012) Overexpression of a histone H3K4 demethylase, JMJ15, accelerates flowering time in *Arabidopsis*. *Plant Cell Rep.* 31(7):1297–1308.
- [94] Yang H, Han Z, Cao Y, Fan D, Li H, Mo H, Feng Y, Liu L, Wang Z, Yue Y, Cui S, Chen S, Chai J and Ma L. (2012) A companion cell-dominant and developmentally regulated H3K4 demethylase controls flowering time in *Arabidopsis* via the repression of FLC expression. *PLoS Genet.* 8(4):e1002664.

- [95] Cui X, Jin P, Cui X, Gu L, Lu Z, Xue Y, Wei L, Qi J, Song X, Luo M, An G and Cao X. (2013) Control of transposon activity by a histone H3K4 demethylase in rice. *Proc Natl Acad Sci USA*. 110(5):1953–1958.
- [96] Shen Y, Conde E, Silva N, Audonnet L, Servet C, Wei W and Zhou DX. (2014) Over-expression of histone H3K4 demethylase gene JM15 enhances salt tolerance in *Arabidopsis*. *Front Plant Sci*. 5:290.
- [97] Schmitz RJ, Sung S and Amasino RM. (2008) Histone arginine methylation is required for vernalization-induced epigenetic silencing of FLC in winter-annual *Arabidopsis thaliana*. *Proc Natl Acad Sci USA*. 105(2):411–416.
- [98] Zhang Z, Zhang S, Zhang Y, Wang X, Li D, Li Q, Yue M, Li Q, Zhang YE, Xu Y, Xue Y, Chong K and Bao S. (2011) *Arabidopsis* floral initiator SKB1 confers high salt tolerance by regulating transcription and pre-mRNA splicing through altering histone H4R3 and small nuclear ribonucleoprotein LSM4 methylation. *Plant Cell*. 23(1):396–411.
- [99] Thomashow MF. (1999) PLANT COLD ACCLIMATION: freezing tolerance genes and regulatory mechanisms. *Annu Rev Plant Physiol Plant Mol Biol*. 50:571–599.
- [100] Chinnusamy V, Zhu J and Zhu JK. (2007) Cold stress regulation of gene expression in plants. *Trends Plant Sci*. 12(10):444–451.
- [101] Bokszczanin KL and Fragkostefanakis S. (2013). Perspectives on deciphering mechanisms underlying plant heat stress response and thermotolerance. *Front Plant Sci*. 4:315.
- [102] Qu AL, Ding YF, Jiang Q and Zhu C. (2013). Molecular mechanisms of the plant heat stress response. *Biochem Biophys Res Commun*. 432(2):203–207.
- [103] Peng S, Huang J, Sheehy JE, Laza RC, Visperas RM, Zhong X, Centeno GS, Khush GS and Cassman KG. (2004) Rice yields decline with higher night temperature from global warming. *Proc Natl Acad Sci USA* 101(27):9971–9975.
- [104] Lobell DB, Schlenker W and Costa-Roberts J. (2011) Climate trends and global crop production since 1980. *Science*. 333:616–620.
- [105] Folsom JJ, Begcy K, Hao X, Wang D and Walia H. (2014) Rice fertilization-independent endosperm1 regulates seed size under heat stress by controlling early endosperm development. *Plant Physiol*. 165(1):238–248.
- [106] Strenkert D, Schmollinger S and Schroda M. (2013) Heat shock factor 1 counteracts epigenetic silencing of nuclear transgenes in *Chlamydomonas reinhardtii*. *Nucleic Acids Res*. 41(10):5273–5289.
- [107] Min L, Li Y, Hu Q, Zhu L, Gao W, Wu Y, Ding Y, Liu S, Yang X and Zhang X. (2014). Sugar and auxin signalling pathways respond to high-temperature stress during

anther development as revealed by transcript profiling analysis in cotton. *Plant Physiol.* 164(3):1293–1308.

- [108] Grandbastien MA. (1998) Activation of plant retrotransposons under stress conditions. *Trends Science.* 3(5):181–187. *Plants*
- [109] Pecinka A, Dinh HQ, Baubec T, Rosa M, Lettner N and Mittelsten Scheid O. (2010) Epigenetic regulation of repetitive elements is attenuated by prolonged heat stress in *Arabidopsis*. *Plant Cell.* 22(9):3118–3129.
- [110] Tittel-Elmer M, Bucher E, Broger L, Mathieu O, Paszkowski J and Vaillant I. (2010) Stress-induced activation of heterochromatic transcription. *PLoS Genet.* 6(10):e1001175.
- [111] Lippman Z, Gendrel AV, Black M, Vaughn MW, Dedhia N, McCombie WR, Lavine K, Mittal V, May B, Kasschau KD, Carrington JC, Doerge RW, Colot V and Martienssen R. (2004) Role of transposable elements in heterochromatin and epigenetic control. *Nature.* 430(6998):471–476.
- [112] Sung S and Amasino RM. (2005) Remembering winter: toward a molecular understanding of vernalization. *Annu Rev Plant Biol.* 56:491–508.
- [113] Kwon CS, Lee D, Choi G and Chung WI. (2009) Histone occupancy-dependent and -independent removal of H3K27 trimethylation at cold-responsive genes in *Arabidopsis*. *Plant J.* 60(1):112–121.
- [114] Hu Y, Zhang L, He S, Huang M, Tan J, Zhao L, Yan S, Li H, Zhou K, Liang Y and Li L. (2012) Cold stress selectively unsilences tandem repeats in heterochromatin associated with accumulation of H3K9ac. *Plant Cell Environ.* 35(12):2130–2142.
- [115] Durrant WE and Dong X. (2004) Systemic acquired resistance. *Annu Rev Phytopathol* 42:185–209.
- [116] Mishina TE and Zeier J. (2007) Pathogen-associated molecular pattern recognition rather than development of tissue necrosis contributes to bacterial induction of systemic acquired resistance in *Arabidopsis*. *Plant J.* 50(3):500–513.
- [117] Conrath U. (2011) Molecular aspects of defense priming. *Trends Plant Sci.* 16(10):524–531.
- [118] Heard E and Martienssen RA. (2014) Transgenerational epigenetic inheritance: myths and mechanisms. *Cell.* 157(1):95–109.
- [119] Jaskiewicz M, Conrath U and Peterhansel C. (2011) Chromatin modification acts as a memory for systemic acquired resistance in the plant stress response. *Embo Reports.* 12(1):50–55.
- [120] Luna E, Bruce TJ, Roberts MR, Flors V and Ton J. (2012) Next-generation systemic acquired resistance. *Plant Physiol.* 158(2):844–853.

- [121] Spoel SH, Koornneef A, Claessens SM, Korzelius JP, Van Pelt JA, Mueller MJ, Buchala AJ, Métraux JP, Brown R, Kazan K, Van Loon LC, Dong X and Pieterse CM. (2003) NPR1 modulates cross-talk between salicylate- and jasmonate-dependent defense pathways through a novel function in the cytosol. *Plant Cell*. 15(3):760–770.
- [122] Koornneef A and Pieterse CM. (2008) Cross talk in defense signaling. *Plant Physiol*. 146(3):839–844.
- [123] Kim JM, To TK, Ishida J, Matsui A, Kimura H and Seki M. (2012) Transition of chromatin status during the process of recovery from drought stress in *Arabidopsis thaliana*. *Plant Cell Physiol*. 53:847–856.
- [124] Ding Y, Fromm M and Avramova Z. (2012) Multiple exposures to drought ‘train’ transcriptional responses in *Arabidopsis*. *Nat Commun*. 3:740.
- [125] Ding Y, Lapko H, Ndamukong I, Xia Y, Al-Abdallat A, Lalithambika S, Sadler M, Saleh A, Fromm M, Riethoven JJ, Lu G and Avramova Z. (2009) The *Arabidopsis* chromatin modifier ATX1, the myotubularin-like AtMTM and the response to drought. *Plant Signal Behav*. 4(11):1049–1058.
- [126] Ding Y, Avramova Z and Fromm M. (2011) Two distinct roles of ARABIDOPSIS HOMOLOG OF TRITHORAX1 (ATX1) at promoters and within transcribed regions of ATX1-regulated genes. *Plant Cell*. 23(1):350–363.
- [127] Ding Y, Ndamukong I, Xu Z, Lapko H, Fromm M and Avramova Z. (2012) ATX1-generated H3K4me3 is required for efficient elongation of transcription, not initiation, at ATX1-regulated genes. *PLoS Genet*. 8(12):e1003111.
- [128] Saleh A, Alvarez-Venegas R, Yilmaz M, Le O, Hou G, Sadler M, Al-Abdallat A, Xia Y, Lu G, Ladunga I and Avramova Z. (2008) The highly similar *Arabidopsis* homologues of trithorax ATX1 and ATX2 encode proteins with divergent biochemical functions. *Plant Cell*. 20(3):568–579.
- [129] Avramova Z. (2009) Evolution and pleiotropy of TRITHORAX function in *Arabidopsis*. *Int J Dev Biol*. 53(2-3):371–381.
- [130] Berr A, Xu L, Gao J, Cognat V, Steinmetz A, Dong A and Shen WH. (2009) SET DOMAIN GROUP25 encodes a histone methyltransferase and is involved in FLOWERING LOCUS C activation and repression of flowering. *Plant Physiol*. 151(3):1476–1485.
- [131] Tamada Y, Yun JY, Woo SC and Amasino RM. (2009) ARABIDOPSIS TRITHORAX-RELATED7 is required for methylation of lysine 4 of histone H3 and for transcriptional activation of FLOWERING LOCUS C. *Plant Cell*. 21(10):3257–3269.
- [132] Berr A, McCallum EJ, Ménard R, Meyer D, Fuchs J, Dong A and Shen WH. (2010) *Arabidopsis* SET DOMAIN GROUP2 is required for H3K4 trimethylation and is crucial for both sporophyte and gametophyte development. *Plant Cell*. 22(10):3232–3248.

- [133] Guo L, Yu Y, Law JA and Zhang X. (2010) SET DOMAIN GROUP2 is the major histone H3 lysine [corrected] 4 trimethyltransferase in *Arabidopsis*. *Proc Natl Acad Sci USA*. 107(43):18557–18562.
- [134] Yun JY, Tamada Y, Kang YE and Amasino RM. (2012) *Arabidopsis* trithorax-related3/SET domain GROUP2 is required for the winter-annual habit of *Arabidopsis thaliana*. *Plant Cell Physiol*. 53(5):834–846.
- [135] Kim S, Lee J, Yang JY, Jung C and Chua NH. (2013) *Arabidopsis* histone methyltransferase SET DOMAIN GROUP2 is required for regulation of various hormone responsive genes. *J Plant Biol*. 56(1):39–48.
- [136] Sani E, Herzyk P, Perrella G, Colot V and Amtmann A. (2013). Hyperosmotic priming of *Arabidopsis* seedlings establishes a long-term somatic memory accompanied by specific changes of the epigenome. *Genome Biol*. 14(6):R59.
- [137] Liu N, Ding Y, Fromm M and Avramova Z. (2014) Different gene-specific mechanisms determine the 'revised-response' memory transcription patterns of a subset of *A. thaliana* dehydration stress responding genes. *Nucleic Acids Res*. 42(9):5556–5566.
- [138] Liu N, Fromm M and Avramova Z. (2014) H3K27me3 and H3K4me3 chromatin environment at super-induced dehydration stress memory genes of *Arabidopsis thaliana*. *Mol Plant*. 7(3):502–513.
- [139] Bouyer D, Roudier F, Heese M, Andersen ED, Gey D, Nowack MK, Goodrich J, Renou JP, Grini PE, Colot V and Schnittger A. (2011) Polycomb repressive complex 2 controls the embryo-to-seedling phase transition. *PLoS Genet*. 7(3):e1002014.
- [140] Azuara V, Perry P, Sauer S, Spivakov M, Jorgensen HF, John RM, Gouti M, Casanova M, Warnes G, Merckenschlager M and Fisher AG. (2006). Chromatin signatures of pluripotent cell lines. *Nat Cell Biol*. 8(5):532–538.
- [141] Bernstein BE, Mikkelsen TS, Xie X, Kamal M, Huebert DJ, Cuff J, Fry B, Meissner A, Wernig M, Plath K, Jaenisch R, Wagschal A, Feil R, Schreiber SL and Lander ES. (2006) A bivalent chromatin structure marks key developmental genes in embryonic stem cells. *Cell*. 125(2):315–326.
- [142] Singh P, Yekondi S, Chen PW, Tsai CH, Yu CW, Wu K, Zimmerli L. (2014) Environmental history modulates *Arabidopsis* pattern-triggered immunity in a HISTONE ACETYLTRANSFERASE1-Dependent manner. *Plant Cell*. 26(6):2676–2688.
- [143] Bruce TJA, Matthes MC, Napier JA and Pickett JA. (2007) Stressful 'memories' of plants: evidence and possible mechanisms. *Plant Sci*. 173(6):603–608.
- [144] Liu J, Feng L, Li J and He Z. (2015) Genetic and epigenetic control of plant heat responses. *Front Plant Sci*. 6:267.

- [145] Bilichak A, Ilnytskyi Y, Hollunder J and Kovalchuk I. (2012) The progeny of *Arabidopsis thaliana* plants exposed to salt exhibit changes in DNA methylation, histone modifications and gene expression. PLoS One. 7(1):e30515.
- [146] Migicovsky Z, Yao Y and Kovalchuk I. (2014) Transgenerational phenotypic and epigenetic changes in response to heat stress in *Arabidopsis thaliana*. Plant Signal Behav. 9(2):e27971.
- [147] Oliver KR, McComb JA and Greene WK. (2013) Transposable elements: powerful contributors to angiosperm evolution and diversity. Genome Biol Evol. 5(10):1886–1901.
- [148] Iwasaki M and Paszkowski J. (2014) Epigenetic memory in plants. EMBO J. 33(18):1987–1998.
- [149] Becker C and Weigel D. (2012) Epigenetic variation: origin and transgenerational inheritance. Curr Opin Plant Biol. 15(5):562–567.
- [150] Weigel D and Colot V. (2012) Epialleles in plant evolution. Genome Biol. 13(10):249.
- [151] Rizhsky L, Liang HJ, Shuman J, Shulaev V, Davletova S and Mittler R. (2004) When defense pathways collide. The response of *Arabidopsis* to a combination of drought and heat stress. Plant Physiol. 134(4):1683–1696.
- [152] Asselbergh B, Achuo AE, Hofte M and Van Gijsegem F. (2008) Absciscic acid deficiency leads to rapid activation of tomato defence responses upon infection with *Erwinia chrysanthemi*. Mol Plant Pathol. 9(1):11–24.
- [153] She W and Baroux C. (2014) Chromatin dynamics during plant sexual reproduction. Front Plant Sci. 5:354.
- [154] Talbert PB and Henikoff S. (2014) Environmental responses mediated by histone variants. Trends Cell Biol. 24(11):642–650.
- [155] Probst AV and Mittelsten Scheid O. (2015) Stress-induced structural changes in plant chromatin. Curr Opin Plant Biol. 27:8–16.
- [156] Sander JD and Joung JK. (2014) CRISPR-Cas systems for editing, regulating and targeting genomes. Nat Biotechnol. 32(4):347–355.
- [157] Wang D and Deal RB. (2015) Epigenome profiling of specific plant cell types using a streamlined INTACT protocol and ChIP-seq. Methods Mol Biol. 1284:3–25.