We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Rotifers as Models in Toxicity Screening of Chemicals and Environmental Samples

Roberto Rico-Martínez, Mario Alberto Arzate-Cárdenas,
Daniel Robles-Vargas, Ignacio Alejandro Pérez-Legaspi,
Alvarado-Flores Jesús and Gustavo Emilio Santos-Medrano

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/61771>

Abstract

An important objective of aquatic ecotoxicology is to determine the effects of toxic compounds in organisms that play a central role in aquatic communities where rotifers have a large impact on several important ecological processes. The contribution of the rotifers to secondary production in many aquatic communities is substantial as they are often the larger fraction of zooplankton biomass at certain times of the year. In addition to the importance of their ecological roles in aquatic communities, the rotifers are attractive organisms for ecotoxicological studies by its short life cycles and rapid reproduction, their small size, and little volumes needed for culture and toxicity assays. The main end points used in ecotoxicological studies are mortality, reproduction, behavior, and biomarkers. Such parameters are included in international regulations from all over the world, where different species are used to evaluate the effect of environmental samples or chemical compounds. The high diversity of rotifers is an important issue because it can modify their relative susceptibility to toxicants. Thus, more studies are needed to know the relations and mechanisms involved in clonal variation, sensitivity, and development, which can be all assessed by state-of-the-art procedures.

Keywords: aquatic toxicology, ecotoxicology, metal toxicity, acute toxicity, endocrine disruption

1. Introduction

The analytical equipment can identify and quantify a chemical substance but not its toxicity in the organisms or the environment, which can be evaluated only in life organisms [1]. Toxicity testing in water samples assesses the concentration and exposure time of the chemical

substances that produce an adverse effect in aquatic organisms, generating useful data for risk assessment. A toxicity test can be accepted by the scientific community if (a) it is capable to predict adverse effects for a variety of compounds in different organisms, (b) it must be replicable with statistical-based analysis, (c) its data must include adverse effects in a range of concentrations in real exposure times, (d) it must be useful to evaluate a risk, (e) it is economic and easy to perform, and (f) it is sensitive and realistic [2].

A model organism for a toxicity test must be abundant, native, or representative of the ecosystem that would be impacted, with ecological and economic importance; there must be a good knowledge of its basic biology that helps interpretation of data and be available for routine maintenance in the laboratory [2].

Toxicity tests have been developed for rotifers to assess many and diverse end points like (a) mortality, (b) reproduction, (c) production of amictic females, (d) cyst production, (e) probability of extinction, (f) behavior, (g) ingestion rates, (h) swimming activity, (i) *in vivo* enzymatic activity, and (j) genetic expression, among others [3–5].

Most of the toxicity tests in rotifers are lethal tests that rely in mortality as the end point to calculate the LC50 value typically at 24 or 48 h without feeding [5]. These tests once standardized are the base to develop monitoring protocols to assess water quality [4, 5]. On the other hand, chronic toxicity tests assess sublethal parameters like behavioral, physiological, or reproductive alterations as the first responses to toxic substances [5, 6], which show high sensitivity in shorter periods of time. The rotifer species *Brachionus calyciflorus* and *Brachionus plicatilis* are among the species most frequently used for both lethal and chronic tests [7, 8].

1.1. Phylum Rotifera

Rotifers are aquatic or semiaquatic microscopic invertebrates with nearly 1850 species; they are unsegmented, pseudocoelomate, and bilaterally symmetric [4, 9, 10]. Nowadays, two classes are recognized: Pararotatoria (with the single order Seisonacea) and Eurotatoria, with two subclasses, Bdelloidea and Monogononta. The size of rotifers ranges from 50 to 2000 μm in length. Most of the species are free swimmers, but some species are fixed to some substratum [4]. Their morphology is of saccate type that is cylindrical with three easily recognizable regions: corona, trunk, and foot (Figure 1) [11].

1.2. Ecological relevance

Rotifers are cosmopolitans. They inhabit aquatic environments of the three types: marine, freshwater, and estuarine [9]. Most rotifers are freshwater, littoral with a few species truly planktonic. The few species that comprise the order Seisonacea and nearly 100 species of the subclass Monogononta are exclusively marine [4]. Species of the subclass Bdelloidea are found in freshwater ecosystems, lakes, temporary pools, interstitial water, soil, moss, and lichens [4, 12]. Rotifers are important in freshwater environments due to having one of the highest reproductive rate among metazoans, thus obtaining high population densities in short times, being dominant in many zooplanktonic communities. They act as links between the microbial community and the higher trophic levels. Rotifers colonize habitats quickly and convert

Figure 1. Diagram of the freshwater rotifer *Brachionus calyciflorus* (Monogononta) [11].

primary production (algae and cyanobacteria) in a usable form for secondary consumers, making energy available for the next trophic levels. In interstitial water from swampy soils, they contribute to nutrient recycling [4]. Some bdelloids and Monogononta are abundant in wastewater treatment plants as part of activated sludge, in filtration systems, or residual lagoons feeding on the bacterial biomass [2, 13]. The population dynamics of rotifers is well characterized both in field as in the laboratory; for this reason, they are useful to investigate ecological and demographic principles. Besides, they are frequently used to assess aquatic toxicity from a population point of view. The intrinsic growth rate (r) is an end point commonly used and has proved to be highly sensitive [5, 14].

1.3. Culture

Rotifers can be obtained directly from a natural aquatic system or from laboratory cultures. Rotifers can be hatched from cysts, and today there are toxicity kits of two species: *B. calyciflorus* and *B. plicatilis* (Rotoxkits). The culture of rotifers has been developed for freshwater, estuarine, and marine organisms taken directly from the natural environment after a previous filtration, sterilization, and neutralization are performed. There is also synthetic hard like EPA medium [15] or synthetic marine water (Instant Ocean) that can be adjusted to the desired salinity. In freshwater species, usually the pH level is maintained in the physiological range (6.5–8) and temperature oscillates between 20 and 30°C. Marine species are maintained at salinities of 10 to 20 psu depending on a particular strain [16]. Rotifers are mainly filter feeders

on microalgae, bacteria, or detritus; a few species are predatory. The supply of fresh and optimal food is the main problem for culturing rotifers. However, numerous rotifer species have been kept in the laboratory routinely [4]. Among the main marine microalgae used to feed rotifers are as follows: *Nannochloropsis* sp., *Chaetoceros* sp., *Dunaliella* sp., *Pyramimonas*, *Isochrysis* sp., and *Tetraselmis* sp.; for freshwater: *Nannochloris* sp., *Nannochloropsis* sp., and *Chlorella* sp. [16]. Culture media have to be renewed 2–3 times a week, and the supply of food must be frequent. Some species are maintained on diluted suspensions of commercial fish food, grain extracts or infusions, manure, and soil [4]. Among the most cultured species are found those of the genus *Brachionus*: *B. plicatilis*, *B. rotundiformis*, and *B. calyciflorus* [17, 18]. Some species of the genus *Lecane* can be easily cultured [19]. The economic importance of these rotifer species that are mass cultured are mainly based in their use as live feed for fish larvae and crustaceans in aquaculture [18].

1.4. Reproduction

The life cycles of rotifers are shorter than many other animals [14]. The Bdelloidea are reproduced by exclusive parthenogenesis and males are unknown. In the subclass Monogononta, the life cycle is haplodiploid with cyclic parthenogenesis as the dominant phase (amictic females), whereas the sexual reproduction (mixis) involves mictic diploid females that produce haploid eggs that if fertilized produce diploid embryos or cysts (in old literature they are called “resting eggs”), which go through diapause before hatching into amictic females that go back to the asexual phase of the cycle (Figure 2). Mixis is triggered by specific environmental stimuli like high population density or photoperiod [18, 20]. Parthenogenesis eliminates the high cost of producing males resulting in a rapid growth population [18], which allows for the high production of clonal individuals that can be used for aquaculture or toxicity tests.

Figure 2. Reproductive cycle of *Brachionus calyciflorus*. Modified from Alvarado-Flores [21].

1.5. Cysts

Once the eggs are produced, they fall into the bottom and are deposited in the sediments [18]. The cysts are diploid and have a thick cover and can be viable for many years in dormancy. Cysts are very resistant to harsh environmental conditions like draught and freezing. Cysts can be dispersed through wind, water, or migratory animals. Under favorable conditions in a specific habitat, the cysts respond to a specific environmental clue (photoperiod, temperature, and salinity) and starts hatching producing an amictic diploid female [4]. Cysts can be stored for great periods of time without losing viability having rotifers in the desired period of time [22]. The use of rotifers hatched from cysts to develop toxicity tests was introduced by Snell and Persoone [23]. Nowadays, it is possible to obtain cysts from marine (*B. plicatilis*) and freshwater (*B. calyciflorus*) species in the market.

1.6. Rotifers as sentinels or bioindicators

Aquatic invertebrates are attractive model organisms in aquatic toxicology due to their short generation time compared with fishes besides their small size require small test volumes [14, 24]. Their small size, the fact that they are filter feeders, and the permeability of the integument made rotifers quite susceptible to chemical and physical changes [19, 25]. Due to the importance of rotifers in the aquatic trophic webs, they are useful as sentinel species to indicate toxicant exposure in affected ecosystems [4].

The knowledge of the (a) basic biology of rotifers, (b) sensitivity to contaminants at the physiological and demographic levels, (c) cosmopolitan distribution, (d) great ecological relevance, (e) high growth rate, (f) availability of neonates, (g) high ingestion rates, (h) ease of culture and handling, (i) transparency, (j) short life cycles, (k) ease to obtain clonal individuals, and (l) cyst production makes rotifers useful tools for assessment of aquatic ecosystems [4, 5, 8, 17, 18, 22, 24, 26].

2. Rotifers as model organism among invertebrates

Invertebrates are the most widely distributed organisms on the Earth and consist of a large and very diverse group, consisting in more than 30 phyla, several of which include more than 1000 different species [27]. The largest phylum within invertebrates is the Arthropoda, with more than one million species, in which insects and crustaceans are the two largest groups [28]. Aquatic crustaceans is a larger group in comparison to the aquatic insects, which a few of them have aquatic larvae [29]. Consequently, crustaceans are the most numerous and ecologically important group of invertebrates in marine and freshwater ecosystems, playing an important role in regulatory toxicity testing, whether in field or laboratory conditions [30, 31]. In addition, these methods are cost-effective because of the facility to obtain amictic eggs or in some cases resting eggs, which have been used to produce commercial kits for toxicity evaluation [32].

Current international protocols include several invertebrates' taxa for toxicity assessment (Table 1), in which cladocerans are the most used organisms and have more approved

protocols for both acute (lethal) and (sub)chronic (sublethal) water toxicity assays than other taxa. The genus *Daphnia* is the most studied taxon within cladocerans. For the acute toxicity test, three replicates for at least five toxicant or effluent concentrations are required, every one of them with 100 mL, 48 h for exposure, without feeding [33]. Moreover, the chronic toxicity assay needs almost the same conditions but with feeding and daily medium renewal for 21 days [34]. In comparison, the *Brachionus* or other rotifer bioassays need only 1 mL per replicate [23], and the results can be obtained within 2 days for acute toxicity evaluation, and in 5 days for the chronic ones. Hence, rotifers tests involve similar conditions to the cladocerans, but when time is a matter of concern, the former is the best option to know the toxicity of chemicals or field samples.

Rotifers are even more versatile than daphnids. In April 2010, after the Macondo oil spill in the Gulf of Mexico, the Environmental Protection Agency of the United States (USEPA) in order to assess the impact of petroleum and oil dispersants on the aquatic biota recommended using *B. plicatilis*, a species complex that included *B. manjavacas*, *B. plicatilis sensu stricto*, and *B. rotundiformis* among other that can exist [35]. This cost-effective test can be replicated hundreds of times in very short periods of time when necessary [36].

The Organization for the Economic Cooperation and Development (OECD) encourages developing test protocols that include crustaceans [37]. In this concern, the copepod *Eucyclops serrulatus* has been proposed as a model organism, with acute toxicity evaluations in 96 h [38] and chronic toxicity assessment in about 60 days [39]. These periods differ due to intrinsic characteristics of the species; copepods last longer to reach sexual maturity after hatching and require mating to obtain eggs for assays, while rotifer eggs are produced by parthenogenesis [40–42]. Nevertheless, with regard to the importance of copepods in aquatic ecosystems, there is not yet any approved protocol for toxicity evaluation as in rotifers [43, 44].

The cnidarian *Hydra attenuata* is another model organism included in standard methods [45]. This one is particularly interesting since lethal and sublethal end points are assessed in the same assay by quantifying the number of death animals and abnormalities found in the survivor hydras during 96 h for acute procedures, which are performed in 12-well microplates, with a test volume of 4 mL [46]. Clubbed tentacles appearance is the first sign of abnormalities due to toxicants exposure. From this point, abnormalities follow shortening tentacles and the tulip stage (all tentacles have disappeared), and finally the disintegration of animal bodies occurs. Therefore, lethal effects should be registered from the tulip stage [47].

It is worth to compare with some rotifers in which it is also possible to determine sublethal effects in the same time period. Alvarado-Flores et al. [48] found abnormal body shapes in *B. calyciflorus* after vinclozolin exposure, demonstrating this pesticide also acts as an endocrine disruptor. As result, both models can be used for sublethal toxicity screening in short periods of time, but the difference lies on the apparent ease to determine such effects, while in *Hydra attenuata* are very obvious. In *B. calyciflorus*, more than one thousand organisms were analyzed to find statistically abnormalities in this rotifer species.

Bivalves in ecotoxicology procedures refer the use of species such as *Lampsilis siliquoides*, *Corbicula fluminea*, *Leptodea fragilis*, *Ligumia subrostrata*, and *Megaloniais nervosa*, among others

[49, 50]. In the scientific literature, most of the bivalve specimens used in toxicological research and evaluations have been collected from environmental samples or commercial producers [51–54]. Glochida are dissected from adult bivalves and need a fish host to develop into juveniles [55–57]. Both immature stages, glochida and juveniles, are used for water quality assessment or chemicals released into aquatic systems [58]. In contrast, rotifer experimental organisms are produced in laboratory-controlled conditions, and animals are ready to use a few hours after hatching instead for several weeks to have juvenile unionids. Hence, rotifer bioassays are less time consuming than bivalves protocols.

In freshwater sediment, toxicity screening several protocols are currently carried out by international regulation agencies (Table 1). Nevertheless, rotifers are not included as test species, but it has been proved that the genus *Lecane* is suitable for those evaluations, representing a cost-effective and high reliable alternative to the standardized methods.

Class	Order	Family	Species
Freshwater			
Cladocera	Branchiopoda	Diplostraca	Daphniidae
			<i>Daphnia magna</i> [62-65]
			<i>D. pulex</i> [66-68]
			<i>D. similis</i> [69]
			<i>D. carinata</i> [70-71]
Copepoda	Branchiopoda	Anostraca	Ceriodaphnia dubia [62, 64, 66, [72]
			Thamnocephalidae
			<i>Thamnocephalus platyurus</i> [73-76]
	Maxillopoda	Calanoida	Cyclopidae
			<i>Eurytemora affinis</i> [77-80]
Rotifera	Monogononta	Ploima	<i>Eucyclops serrulatus</i> [31, 81-83]
			Brachionidae
			<i>Brachionus calyciflorus</i> [43, 84]
			<i>B. rubens</i> [23, 85-87]
			<i>B. patulus</i> [88-90]
			Lecanidae
			<i>Lecane quadridentata</i> [91-93]
Cnidaria	Hydrozoa	Anthoathecata	<i>L. hamata and L. luna</i> [94,95]
			<i>Euchlanis dilatata</i> [96, 97]
			Asplanchnidae
			<i>Asplanchna brightwelli</i> [98-101]
			Hydridae
Mollusca	Bivalvia	Unionoida	<i>Hydra attenuata</i> [45]
			<i>H. vulgaris</i> [102-104]
			Unionidae
			<i>Elliptio complanata</i> [58]
			<i>Lampsilis siliquoidea</i> [58]
			Veneroida
			Corbiculidae
			<i>Corbicula fluminea</i> [105-108]

	Class	Order	Family	Species
Freshwater sediments				
Crustaceans	Malacostraca	Amphipoda	Hyalellidae	<i>Hyallela azteca</i> [109, 112]
Annelida	Clitellata	Lumbriculida	Lumbriculidae	<i>Lumbriculus variegatus</i> [113-115]
				<i>Stylodrilus heringianus</i> [116, 117]
Rotifera	Bdelloidea		Philodinidae	<i>Philodina roseola</i> [118]
Arthropoda	Insecta	Diptera	Chironomidae	<i>Chironomus dilutus</i> [109, 110, 119-121]
				<i>C. riparius</i> [109, 110, 119-121]
Seawater				
Rotifera	Monogononta	Ploima	Brachionidae	<i>Brachionus plicatilis</i> [43]
Echinodermata	Echinoidea	Arbacioida	Arbaciidae	<i>Arbacia punctulata</i> [122]
Arthropoda	Malacostraca	Mysida	Mysidae	<i>Mysidopsis bahia</i> [123]
Copepoda	Maxillopoda	Calanoida	Acartiidae	<i>Acartia tonsa</i> [44, 124]
		Harpacticoida	Miraciidae	<i>Amphiascus tenuiremis</i> [44]
Molluscs	Bivalvia	Mytiloida	Mytilidae	<i>Mytilus edulis</i> [125, 126]
		Ostreoida	Osteridae	<i>Crassostrea gigas</i> [127]
Seawater sediments				
Annelida	Polychaeta	Capitellida	Arenicolidae	<i>Arenicola marina</i> [128]
Artropoda	Malocostraca	Amphipoda	Corophiidae	<i>Leptocheirus plumulosus</i> [110, 129, 130]

Note: Numbers in square brackets indicate reference number

Table 1. Toxicity evaluations protocols using different invertebrates for freshwater and freshwater sediments, seawater, and seawater sediments.

Differently to the protocols mentioned above, all those for seawater and seawater sediments are found in standardized probes approved by agencies such as USEPA, ASTM, and the OECD (Table 1). As previously described, *B. plicatilis* is a marine rotifer used in seawater toxicology evaluation, an although it is not included in the USEPA guidelines, the agency recommended its used based on the ASTM method [43]. For sediments, the prime option is the polychaetes such as *Arenicola marina*, mainly collected from beach sand in different locations around the world. The protocol involves an exposure time of about 28 days using artificial sediment and field sediments samples [59–61]. Moreover, *B. plicatilis* has been exposed to interstitial water from littoral ecosystems; although it is sensitive to bioavailable toxicants in pore water, it does not respond in the same manner that those organisms that live in the sediment.

Afterward, the use of rotifers in all the above-mentioned matrices is possible due to some advantages, such as their relative small size, their simple organization, their short life cycles that permit multigenerational studies in very short time periods, their reproduction through parthenogenesis, and their genetic homogeneity that leads to almost identical offspring [22]. Despite of these characteristics, there is still a need to continue researching the effect of toxicants and to improve protocols for rotifers in (eco)toxicology to better understand their physiology and how external factors alter rotifers normal responses, since studies with rotifers are not as numerous as other taxa (Figure 3).

Figure 3. Number of publications for the groups described in the text. The databases consulted were Elsevier (www.sciencedirect.com), Springer (www.link.springer.com), and Wiley Online Library (<http://onlinelibrary.wiley.com>). Query terms were the genera name and the word “toxicity”. Pie chart data represent percentages based on published material.

3. Rotifer sensitivity

The toxicity of a wide variety of chemicals including organic compounds, metals, and pharmaceuticals that can be found in water reservoirs as a result of human activity has been tested by performing toxicity tests using rotifers as bioindicators as these organisms play an important ecological role in aquatic environments [3, 131]. Rotifers of the genus *Brachionus* can be

found between the most commonly used test organisms [132]; moreover, several rotifers are included for comparison purposes.

A test organism with high sensitivity to detect adverse effects of xenobiotics is always desirable; nevertheless, differences in sensitivity among species can always be found. Generally, these are due to the biological characteristics of test organisms, the type of chemical, and its mode of action; thus, one species might be very sensitive to one toxic but not that sensitive to another compound [133]. This section focuses on the sensitivity of different rotifers, all of them from freshwater ecosystems, with exception of the marine species *B. plicatilis* and *Brachionus koreanus*, when testing the toxicity of pesticides, organic compounds, and metals attending their toxicological importance and the availability of literature.

Some examples of these variations on sensitivity are shown in Table 2. In relation to pesticides, malathion (organophosphorus) displays the lower toxicity and has a toxicity range from 33.72 to 45.5 mg/L, where the lower toxicity level is reported to *B. calyciflorus* and the higher to *B. plicatilis* [134, 135].

Lindane, an organochloride pesticide, and methyl parathion (organophosphorus) show moderate toxicity and have wider toxicity ranges in comparison to malathion. For the organochloride, toxicity range is given by *B. koreanus* and *B. plicatilis* (14 and 35.89 mg/L, respectively). In the case of methyl parathion, which shows higher acute toxicity, a range from 0.607 mg/L for *Euchlanis dilatata* to 29.19 mg/L for *B. calyciflorus* was found. Another organophosphorus of ecotoxicological relevance, chlorpyrifos, tends to affect significantly aquatic invertebrates due the LC50 values reported for *B. koreanus* (3.9 mg/L) and *B. calyciflorus* (11.85 mg/L) [96, 134, 136, 137].

The organochlorides endosulfan and pentachlorophenol (PCP) stand as highly toxic agrochemicals. For endosulfan, *B. koreanus* and *B. plicatilis* correspond to the reference organisms to establish the range because of their 24-h LC50 values found in literature [135, 136]. For PCP, the lowest and highest toxicity levels were noted for *B. calyciflorus* [138, 139].

On literature, little information is available about the toxicity of other organic compounds such as benzene, toluene, xylene, and hexane. Nevertheless, Ferrando and Andreu-Moliner [7] and Pérez-Legaspi et al. [95] have obtained some toxicological data (Table 2). For xylene, hexane, and toluene, *B. calyciflorus* and *B. plicatilis* are the reference organisms, where the first was most sensitive for the three chemicals [7, 95]. For benzene, rotifers from the genus *Lecane* resulted being more sensitive, but also a wide toxicity range is registered from 6.97 for *Lecane luna* to 3762 mg/L for *Lecane hamata* [95].

The lowest LC50 values, and thus, more toxic metals, were reported for zinc, copper, and mercury. Besides, a relatively not wide toxicity range can be observed for these three chemicals (Table 3). The most sensitive test organisms in these cases were *Lecane quadridentata*, *B. plicatilis*, and *B. calyciflorus*, respectively, while the more resistant were *Brachionus havanaensis* (Zn), *P. acuticornis* (Cu), and *L. hamata* (Hg) [93, 95, 135, 140, 143, 144].

Moreover, it is important to mention that there are physical factors that can alter the sensitivity of a test organism when performing a toxicity test. One example is given by Preston et al. [138],

Pollutant	Acute toxicity range in terms of 24h-LC50 (mg/L)
Pesticides	
Malathion	33.72[134] - 45.5[135]
Lindane	14[136] - 35.89[137]
Methyl parathion	*0.607[96] - 29.19[134]
Chlorpyrifos	3.9[136] - 11.85[137]
Endosulfan	4[136] - 6.6[135]
PCP	0.21[138] - 2.16[139]
Another organic compounds	
Xylene	252.7[7] - 495.9[7]
Hexane	68.3[7] - 154.3[7]
Acute toxicity range in terms of 24 and 48h-LC[50](mg/L)	
Benzene	6.97[95] - 3,762[95]
Toluene	113.3[7] - 552.6[7]

Note: Numbers in square brackets indicate reference number

*48h LC₅₀ value

In the case of inorganic toxicants (Table 3), lead appears as the “less” toxic and with the wider toxicity range from 0.035 for *E. dilatata* to 56.2 mg/L for *Philodina acuticornis* [97, 140]. For cadmium, *E. dilatata* shows a great sensitivity with a 48-h LC₅₀ of 0.014 mg/L, while *B. plicatilis* is more resistant (39 mg/L) [97, 141]. The most sensitive rotifer to chromium was *Lecane luna* (48-h LC₅₀ of 3.26 mg/L) [95], and the highest 24-h LC₅₀ of 17.4 mg/L was registered for *B. calyciflorus* [142].

Table 2. Toxicity range for some pesticides and organic compounds to rotifers used in bioassays.

Metal	Acute toxicity range in terms of 24 and 48h-LC[50] (mg/L)
Lead	0.035[97] - 56.2[140]
Cadmium	0.014[97] - 39[141]
Chromium	3.26[95] - 17.4[142]
Zinc	0.123[93] - 2.271[143]
Copper	0.01[135] - 1.9[140]
Mercury	0.06[144]- 1.37[95]

Note: Numbers in square brackets indicate reference number

Table 3. Toxicity range for different metals to rotifers used in bioassays.

where a decrease in LC₅₀ values for PCP and mercury using *B. calyciflorus* as bioindicator was recorded as UV-B exposure time was increased.

Other physical condition that might modify the sensitivity of a test organism is desiccation. Robles-Vargas and Snell [145] found a remarkable resistance of *Philodina* sp. emerged from desiccation to PCP, chlorpyrifos, and mercury in comparison to continuously hydrated rotifers when reproductive tests were implemented [145].

According to the information reviewed and in terms of acute toxicity, rotifers from the family Brachionidae tend to show a great sensitivity to different chemicals, including organic and inorganic compounds. Invertebrates from the family Euchlanidae and Lecanidae appear to be sensitive mainly to inorganic toxicants. Furthermore, rotifers belonging to the family Philodinidae exhibit a notorious resistance to chemicals.

4. Rotifer database for screening toxicants

Every chemical product that can reach ecosystems and, as a consequence, the human being needs to be characterized to avoid noxious effects on all organisms exposed. At first instance, toxicological studies were mainly anthropocentrically focused, but that point of view would not have been enough to protect our own existence at all if affecting organisms in different ecosystems would have continued. It is known that rotifers play an important role as secondary producers and in some freshwater ecosystems represent the larger fraction of biomass of zooplankton [146]. Therefore, these organisms are nowadays considered as model organisms for toxicological screening.

Currently, one can find in the scientific literature reports of almost every class of chemical compound used or produced by man, investigating their toxicological properties. These papers are relatively abundant in several websites specialized in gathering such kind of information, which is available to everyone who has access to the published material. For scientist, collecting this material is a daily activity that guarantees their studies to follow the right direction by not working in exactly the same topic and manner, which would represent the duplicity of information but at higher cost because of multiple double efforts and resources investment. Nonetheless, it must be pointed out that differences in susceptibility to toxicants can be found among clones or strains, which is very interesting to assess by multiclonal analyses [22, 147]. However, when it is not possible to access multiple clones or strains, previous reports (published material) from databases become more relevant as a source of comparison to improve the comprehension of the observed results.

The USEPA, in an attempt to ease the search for environmental toxicological data, has created the ECOTOX database, hosted in their website [150]. It provides a user-friendly interface and allows discrimination among taxonomic and chemical groups, aquatic or terrestrial environment, tests results or conditions, and type (year) of publication. Although it does not give access to the cited references, it returns a list of publications to be consulted elsewhere. In the ECOTOX results table, the data shown are species name, exposure type, chemical, media type, end points, bioconcentration factor, effects, and statistics, among others (Figure 4). A complete list of all abbreviations used in the ECOTOX database can be directly downloaded for the same host website.

The screenshot shows the ECOTOX Aquatic Report interface. At the top, it says "U.S. ENVIRONMENTAL PROTECTION AGENCY" and "ECOTOX Aquatic Report". Below this, there's a search bar and a table of results. The table has columns for Species, Test Name, Media Type, Test Type, Chemical Name, Concentration, Effect, and Date. The results are filtered for "Brachionus" and "chromium".

Species	Test Name	Media Type	Test Type	Chemical Name	Concentration	Effect	Date
Brachionus calyciflorus	ECOTOX	Water	Acute	Chromium (VI)	1.000 mg/L	100%	1989
Brachionus calyciflorus	ECOTOX	Water	Acute	Chromium (VI)	1.110 mg/L	100%	1989
Brachionus calyciflorus	ECOTOX	Water	Acute	Chromium (VI)	1.110 mg/L	100%	1989
Brachionus calyciflorus	ECOTOX	Water	Acute	Chromium (VI)	1.110 mg/L	100%	1989
Brachionus calyciflorus	ECOTOX	Water	Acute	Chromium (VI)	1.000 mg/L	100%	1989
Brachionus calyciflorus	ECOTOX	Water	Acute	Chromium (VI)	1.000 mg/L	100%	1989

Figure 4. Print screen of ECOTOX database results. Query terms were *Brachionus* and chromium, selected from the options displayed in the database [150].

Nowadays, toxicity assays with rotifers include very diverse toxicants, and a relatively high number of test have been assessed the effect of inorganic (metals and metalloids) and very versatile organic compounds like pesticides, solvents, colorants, detergents, and emergent toxicants, in which health and care products and pharmaceuticals form part of it (Table 4). Some of these toxicants are effective at concentrations as low as some nanograms per liter, altering rotifers population dynamics by different mechanisms like endocrine disruption.

Species	Chemical compound or metal	Species	Chemical compound or metal
Bc	1,1,1-trichloroethane [150]	Bc	Lauryl alcohol [150]
	1,3,5-Trinitrobenzene [150]	Ab, Ai, Bc, Ed Lh, Ll, Lq, Pa	Lead [91, 94, 97]
	1-Dodecanesulfonic acid, Sodium salt [150]	Ap, Bc, Br, Kq	Lindane [150]
	1-Octanol [150]		Linuron [150]
Bp	2,2',4,4'-Tetrabromodiphenyl ether (BDE-47) [151]	Bc	Lithium [157]
Bc, Kc	2,3,4,6-Tetrachlorophenol [150]		m- and p-Xylene [150]
Ba, Ka, Kc	2,4,5-T [150]		Malathion [150]
Bc	2,4,6-trichlorophenol [150]	Br	Malathion [150]
	2,4,6-Trinitrotoluene [150]	Pa	Malathion [150]

Species	Chemical compound or metal	Species	Chemical compound or metal
	2,4-Dichloro aniline [150]	Bc	Malathion [152]
	2,4-Dichlorophenoxyacetic [150]	Bc	Malathion [153]
	2,4-dinitrochlorobenzene [150]	Bc, Li, Lq	Manganese [150], [166], [169]
	2,4-xyleneol [150]	Bc	MARLON A390 [150]
	2,5-Dichloro aniline [150]	Bc, Bp, Ed, Lh, Ll, Lq, Pa	Mercury [91, 94, 97]
	2- [2- [2-(Dodecyloxy)ethoxy]ethoxy]ethanol, Hydrogen sulfate, Sodium salt [150]	Bc	Metallic silver [150]
	3,4-dichloro aniline [150]	Av	Methacrylamide [171]
	Acetaminophen [150]		Methacrylic acid [150]
Bc, Lh, Ll, Lq	Acetone [94, [150]]	Bc	Methanol [150]
Bc	Acetylsalicylic acid [152, 153]		Methoprene [150]
Av	Acrylamide [154]	As, Ba, Bc, Bp, Ed	Methyl parathion [97, 150]
	Acrylic acid [150]		Molinate [150]
	Aldrin [155]		N,N,N-Trimethyl-1-octanamonium chloride [150]
Bc	Alkyl* sodium benzene sulfonate [150]		N,N-Dimethyl-1-dodecanamine [150]
	Alkyl* trimethyl ammonium chloride [150]	Bc	N,N-Dimethyl-1-octanamine [150]
Ab, Bc, Li, Lq	Aluminum [93, 150, 156]		Naphthalenol [150]
Bc	Amitriptyline [152, 153, 157]		Naproxen [150]
Bc, Br	Ammonia [150]		n-Hexane [150]
Bc, Pa	Ammonium chloride [150]	Bc, Pa	Nickel [150]
Bc	Amphetamine sulfate [153]		Nicotine [152, 153]
Bc	Aroclor 1260 [150]		Nitric acid, Cadmium salt [150]
Bc, Pp, Pr,	Arsenic [150, 158>]	Bc	Nonyl phenol [150]
Bk	Atenolol [159]		Ofloxacin [161, 165]
Bc, Kq	Atrazine [150]		Orphenadrine [152, 153, 157]
Bc	Atropine [152, 153, 157]		Oxytetracycline [150]
Bc	Barium nitrate [150]	Bc, Bk	Oxytetracycline [159, 165]
Bc, Lh, Ll, Lq	Benzene [94, 150]	Bc	Paracetamol [152, 153]
Bc	Bezafibrate [160]		Para-chlorophenol [150]
Ab, Ai, Bc, Bh, Bp, Br, Bu, Ed, Lh, Ll, Lq, Pa	Cadmium [94, 143, 150, 160]	Bc	Para-dichlorobenzene [150]

Species	Chemical compound or metal	Species	Chemical compound or metal
Bc	Caffeine [150]		P-chloroaniline [150]
Pa	Calcium hypochlorite [150]	Ag, Bc, Br, Kq, Pa,	Pentachlorophenol PCP [150]
Bc, Bk	Carbamazepine [159, 161, 162]	Bc	Phenobarbital [152, 153, 157]
Ba, Hm, Mq, Pa	Carbaryl [150]	Bc, Br, Pa	Phenol [150]
Eb, Kq	Carbendazim [150]	Br	p-Nitrophenol [150]
	Carbofuran [150]	Bp	polybrominated diphenyl ethers [173]
Bc	Carbon tetrachloride [150]	Bc	Potassium chloride [150]
	Cetyl trimethyl ammonium chloride [150]	Bc, Pa	Potassium cyanide [150]
	Chloramphenicol [152, 153, 157]	Bc, Pa	Potassium dichromate [150]
Pa	Chlorine [150]		Prednisolone [174]
Bu	Chlornitrofen [150]		Prednisone [175]
	Chloroacetic acid [150]		Prococene [150]
Bc	Chloroform [150]		Propranolol [152, 153, 157, 176]
	Chloroquine [152, 153]	Bc	Quinidine [152]
Ab, Ba, Bb, Bc Bq, Bu, Eb, Ft, Hi, Hm, Kq, Mq, Pq, Tp	Chlorpyrifos [150, 163]		Ranitidine [177]
Bc, Bp, Lh, Ll, Lq	Chromium [94, 150, 163]		Selenium [150]
Bc	Clarithromycin [165]	Pa	Silver nitrate [150]
Pa	Cobalt chloride [150]	Br	Simetryn [150]
Ab, Bc, Bp, Br, Lh, Ll, Lq, Li, Pa, Pp	Copper [94, 150, 158, 164, 166, 167]	Bp	SLS [178]
Ab, Bc, Bp	DDT, p,p' [150, 169]	Bc	Sodium bromide [150]
	Diazepam [152]	Bc	Sodium chloride [150]
	Diazinon [150]	Pr	Sodium chromate [150]
Bc	Diclofenac [161, 162]	Bc	Sodium dodecylbenzene sulfonate [150]
	Dicofol [150]	Pa	Sodium fluoride [150]
	Digoxin [152]	Bc	Sodium hypochlorite [150]
Pa	Dimethoate [150]	Bc	Sodium laurate [150]
Bc	Dinitro cresol [150]	Bc, Br	Sodium lauryl sulfate [150]

Species	Chemical compound or metal	Species	Chemical compound or metal
	Diphenylhydantoin [152]		Sodium oxalate [150]
	Endosulfan [150]		Sodium selenate [150]
	Endothall [150]		Sodium tetradecyl sulfate [150]
	Erythromycin [165]		Sulfamethoxazole [159, 161, 165]
	Estradiol [150]	Bc	Testosterone [168]
	Ethinylestradiol [94, 168]		Thallium(I) sulfate [150]
Lh, Ll, Lq	Ethyl acetate [94, 168]		Theophylline [152, 153]
	Ethyl alcohol [150]		Thiobencarb[150]
	Ethyl methacrylate [150]		Thioridazine [153]
Bc	Ethylene glycol [150]	Li	Tin [166]
	Fenitrothion [150]	Lh, Ll, Lq	Titanium [94]
	Fenofibrate [160]	Bc, Lh, Ll, Lq	Toluene [94, 150]
Bc, Pa	Ferrous sulfate [150]		Tributyl phosphate [150]
Ab	Iron [156]	Bc	Tributylstannane [150]
	Flutamide [150]		Tributyltin chloride [150]
Bc	Furosemide [170]	Bc, Kq, Pa	Trichlorfon [150]
	Gemfibrozil [160]	Bk	Trimethoprim [159]
Bc, Lq	Glyphosate [150]	Bc	Verapamil [152, 153]
	Hexachlorophene [150]	Lh, Ll, Lq	Vinyl acetate [94]
Bc	Hydroquinone [150]		Warfarin [152]
Ab, Li, Lq	Iron [96, 156, 166]	Bc	Xylene [150]
	Isoniazid [152, 153]	Ab, Bc, Bh, Li, Lq, Pa	Zinc [93, 143, 150, 156, 166]
Bc	Isopropyl alcohol [150]		

Note: superscripts indicate reference number

Notes: Species are abbreviated with the first letters of their genus and species: Ad = *Adineta vaga*, Ab = *Asplanchna brightwellii*, Ag = *Asplanchna girodi*, Ai = *Asplanchna intermedia*, As = *Asplanchna sieboldi*, Ba = *Brachionus angularis*, Bb = *Brachionus bidentata*, Bc = *Brachionus calyciflorus*, Bh = *Brachionus havanaensis*, Bk = *Brachionus koreanus*, Bp = *Brachionus patulus*, Bp = *Brachionus plicatilis*, Bq = *Brachionus quadridentatus*, Br = *Brachionus rubens*, Bu = *Brachionus urceolaris*, Eb = *Epiphanes brachionus*, Ed = *Euchlanis dilatata*, Ft = *Filinia terminalis*, Hi = *Hexarthra intermedia*, Hm = *Hexarthra mira*, Ka = *Keratella americana*, Kc = *Keratella cochlearis*, Kq = *Keratella quadrata*, Lh = *Lecane hamata*, Li = *Lecane inermis*, Ll = *Lecane luna*, Lq = *Lecane quadridentata*, Mq = *Monostyla quadridentata*, Pa = *Philodina acuticornis*, Pr = *Philodina roséola*, Pp = *Platyonus patulus*, Pq = *Platyias quadricornis*, Tp = *Testudinella patina*.

Table 4. Relation of toxicants and rotifers species used in ecotoxicological studies

The available data about toxicity to rotifers show that organic compounds, as a complex group, is the most studied (32.4%) followed by the drugs group (24.58%), which could be due to the complexity of both groups, as they include a huge variety of toxicants. Inorganic compounds among other groups represent 17% out of the total number of tests included in this chapter, which can be split up into inorganic (nonmetallic compounds) and metals (Figure 5). The last one comprised those commonly named heavy metals [148]. Other groups might not be as numerous as the former ones, but this should not be misinterpreted as a lack of interest from the environmental toxicologists, but a challenge and a continuous research to find out how these chemicals are affecting the aquatic ecosystems and their inhabitants like rotifers species.

Figure 5. Comparison of toxicity tests between chemical groups conducted with different rotifers species.

Several rotifer species have been used to evaluate the effect and toxic mechanisms of the chemical groups aforementioned, where *B. calyciflorus* is the most abundant in the literature, with about 63% out of the total publications (Figure 6). However, there are other species that present advantages over brachionids, such as the littoral genus *Lecane* that might be used to assess the effect of sediment-associated toxicants, or the predator genus *Asplanchna*, used to evaluate toxicants biomagnification among rotifers and other aquatic invertebrates [149]. Therefore, rotifers represent a group of organisms that have not been completely studied and require more attention from toxicologists.

Red bars correspond to left *y*-axis and blue bars to right *y*-axis.

Figure 6. Number of ecotoxicological studies through by different rotifer species.

5. Rotifer tests already used worldwide and in certain regions

The ability to produce cysts has allowed the development of toxicity kits, called Rotoxkits, employed for acute/chronic marine and freshwater toxicity testing [179]. Cyst production is an outstanding characteristic that has enabled the development of several toxicity protocols using rotifers that have been used worldwide [180]. Rotifers are not directly represented in the legislation of several countries (as the cladocerans *Daphnia magna* and *Ceriodaphnia dubia* do). However, toxicity tests with rotifers have been published by official societies [43, 84, 181] Perhaps the most notorious participation of rotifer toxicity tests is when EPA asked BP plc (the former name of the company was British petroleum) to use the acute toxicity test with the euryhaline rotifer *B. plicatilis* to assess the toxicity generated after the 2010’s Gulf of Mexico oil spill [35]. The marine water rotifer toxicity test TK22 was used to analyze thousands of sites in the Gulf of Mexico; toxicity was analyzed for both oil and the oil dispersant used [36].

5.1. Europe

The Convention for the Protection of the Marine Environment of the North-East Atlantic (the “OSPAR Convention”) was established in 1992. The convention started working in 1998. It has been ratified by Belgium, Denmark, Finland, France, Germany, Iceland, Ireland, Luxembourg, Netherlands, Norway, Portugal, Sweden, Switzerland, and the United Kingdom and approved by the European Community and Spain. One of their goals was the development of directives for the analysis of toxicity both in marine and freshwater environments. In their 2007 publication “Practical Guidance Document on Whole Effluent Assessment” (OSPAR, 2007), they applied the protocol of determination of the chronic toxicity to *B. calyciflorus* in 48 h [84] and validated its use.

In the Port of Aveiro, Portugal, standardized acute toxicity test using the marine rotifer, *B. plicatilis*, on sediment elutriates revealed higher toxicity levels in cases where the European Union Water Framework Directive suggested that sediment quality was acceptable [183]. Isidori et al. [184] employing *B. plicatilis* in 24-h toxicity tests found that all samples of municipal solid waste landfills in southern Italy expressed acute toxicity.

In Poland, the toxicity of the leachates from sewage sludge were investigated using different toxicological kits: Microtox (*Vibrio fischeri*), microbial assay for toxic risk assessment (ten bacteria and one yeast), Protoxkit F™ (*Tetrahymena thermophila*), Rotoxkit F™ (*B. calyciflorus*), and Daphtoxkit F™ (*D. magna*). Differences were observed in the sensitivity of the test organisms to the presence of sewage sludge in the soil. The highest sensitivity was a characteristic of *B. calyciflorus* [185].

5.2. Oceania

The Resource Management Act 1991 of New Zealand [186] requires local authorities and industry to apply for consent to discharge effluent to water or land. Therefore, the requirement for whole effluent toxicity testing is now being included in these Resource Consents. The Ministry for the Environment (MfE) has encouraged research which evaluates technology used to monitor environmental parameters (e.g., effluent toxicity). Such research was done using the Rotoxkit F™ and Rotoxkit M™ (with *B. calyciflorus* and *B. plicatilis*, respectively). The study concluded that “while very cost effective and with very good precision (repeatability), were not as sensitive as the others, and therefore are not recommended for whole effluent toxicity screening” [187].

5.3. Latin America and the Caribbean

Sarma et al. [188] showed that Mexico City urban wastewater affects instantaneous growth rate of *Brachionus patulus*. Acute 48-h lethal effect measurements generated with *L. quadridentata* on municipal, industrial, and agricultural sites around the city of Aguascalientes, Mexico, indicated that most samples tested were toxic [189]. The municipal drinking water wells of Aguascalientes have also been assessed using acute toxicity tests with *L. quadridentata* [190]. *L. quadridentata* has been used to assess the status of the major wastewater treatment plants in the State of Aguascalientes [191, 192]. An ecotoxicological study of the most important river

in the State of Aguascalientes has also been performed [93]. *L. quadridentata* has also been used to assess toxicity in several rivers of the southern Huasteca area of Mexico with high concentrations of manganese (Mn) and the presence of DDT in the sediments and pore water. *L. quadridentata* was highly resistant to DDT and less susceptible to Mn than the cladocerans *D. magna* [165]. José de Paggi and Devercelli [193] examined the influence of watershed land use on microzooplankton around the city of Santa Fe in Argentina. Six rivers and a shallow lake located in rural and urban areas were sampled during 4 weeks. River microzooplankton abundance and rotifer species assemblages were found to be good indicators of land use. *Brachionus* spp. were associated with saline waters in rural areas and *Keratella* spp. (except *Keratella tropica*) with urban water bodies.

5.4. Asia

Many Asian countries have used rotifer toxicity tests for diverse monitoring and scientific purposes. Microcosm studies with rotifers have been used in India to evaluate tannery effluent [194]. *B. plicatilis* has been used to assess the toxicity of the various sewage sludge, one of the major ocean dumped materials in the Yellow Sea of Korea [195].

6. Perspectives for future studies regarding the importance of rotifers as models for toxicity screening of chemicals

Different rotifer species from all around the globe have been used to test the toxicity of a huge number of chemicals, both from freshwater (FW) and marine water (MW) ecosystems (see Rico-Martinez et al., 2013). Such species were initially collected from their natural habitats, a specific biogeographical zone, for their further acclimation to laboratory conditions and use as model organisms in toxicity evaluation protocols.

Nowadays, the rotifer species used as model organisms, due to their representativeness and wide natural distribution, include those of the genus *Brachionus* sp.: *B. calyciflorus* (FW), *B. plicatilis* (MW), *B. manjavacas* (MW), and *B. koreanus* (MW) (Lee et al.; Snell et al.), although some others like *B. rotundiformis* (MW) and *B. ibericus* (MW) could be used as model organisms (Pérez-Legaspi, 2015). Moreover, organisms within the genus *Lecane* sp. have been used for toxicological screening, demonstrating that *L. quadridentata* and other *Lecane* species are good indicators of water quality because of their high susceptibility to toxicants.

Despite all efforts to understand rotifers biology and their susceptibility to contaminants, there is still a need to conduct new studies with rotifers belonging to different habitats (biogeographical zones), climates, and niches. For such studies, researchers should take into consideration topics like clonal cultures obtaining, rotifers identification and classification through morphological and genetic (cytochrome oxidase rDNA, COI) characters, and the production of sexual eggs to preserve them in a resting eggs bank. Such eggs could be also a source for developing toxicity assessments kits, like those of Microbiotests Inc. [179], which mainly uses *Brachionus* sp. resting eggs in very efficient systems for their production.

Due to their immeasurable dispersion and diversity, annotating correctly the rotifers specimens' origin and recording their chemical backgrounds has become a very important issue because of the variable responses to toxicants observed within the same genus or even within the same species. For example, *Brachionus* sp. VER, isolated from the Gulf of Mexico, was the most tolerant to Macondo oil exposure (LC50 = 19.33%) in comparison to *B. rotundiformis* (LC50 = 11.02%) from Hawaii, *B. plicatilis* ss. (LC50 = 2.47%) from Tokyo, and *B. manjavacas* (LC50 = 5.43%) from Russia [36], which could be due to the presence of cryptic species within the taxon. Recently, researches and students from different countries participated in the workshop "Cryptic Speciation in *B. plicatilis*: A Workshop to Described Species within the Complex," and they estimated that there may be as many as 20 new species for this complex. In addition, the rotifers *B. calyciflorus* and *Lecane bulla* form a part of cryptic species [196, 197]. Thus, genetic and phylogeographic studies should be performed to assess how this species are distributed around the world.

In aquatic toxicology, currently there are methods that help elucidating the toxicity mechanisms for different sorts of chemicals. Therefore, they can be listed as follows:

- a. Standardized protocols for evaluation of acute, chronic, and sublethal toxicity. These methods are carried out by exposing neonates or resting eggs, for 24 to 48 h or more depending exposure concentrations, periods, and end points to measure. Acute and chronic toxic ratios are still in use despite all new technologies, as they are finally a reference point for further analysis. Mortality or immobility are the common responses observed in acute toxicity tests, but in chronic assays, population parameters are followed during the exposure period, such as the intrinsic rate of population increase (r) obtained from the life table analysis. Another possibility is assessing the hatching percentage, which represent the stability and health of cysts produced during stressful conditions, including abiotic factors like desiccation or the presence of contaminants. Alvarado-Flores et al. [48] evaluated the effect of 1.2 mg/L of the fungicide vinclozolin on the rotifer *B. calyciflorus*. Their findings demonstrated that there was no significant difference between exposed and nonexposed organisms. In addition, the population parameter r of rotifers hatching from VZ-promoted cysts was 1.21 ± 0.063 (mean $\pm$ SD), and for rotifers hatching from control cysts was 0.90 ± 0.064 .

It has been shown that multigenerational studies should be conducted as these could reproduce what happens in natural conditions when parthenogenetic females are exposed to toxicants and in their offspring is in certain way altered even before hatching [198], a phenomenon called the maternal effect, which could be for good while providing more energetic resources to deal with the stressful conditions in the medium, or negative through inheriting mutations that could bring deleterious effects in consecutive generations.

- b. *In vivo* enzyme activity assessment (esterases and phospholipase A2): This method has the potential to assess the adverse effects of contaminants for rotifers. In the rotifer *E. dilatata* (FW), a native species from Mexico, the inhibition of esterases and phospholipase A2 was assessed fluorometrically after *in vivo* exposure (30 min) to sublethal concentra-

tions of metals and pesticides in laboratory conditions. This study concluded that both enzymes are very sensitive to toxicants-induced alterations [96, 97].

- c. Stress granules (SGs): Eukaryotes share diverse mechanisms of adaptation and responses to the stress. In this matter, it has been shown in different insects, trypanosomids, yeast, mammal cells, and rotifers, in which they can sequester some proteins and mRNA into granules that protect cellular mRNA. Thus, SGs appear to be useful as biomarkers in rotifers [199].
- d. Bioconcentration factors (BCF): According to van der Oost et al. [200], biomarkers of exposure include the quantification of the toxicant or its metabolites. Therefore, BCF are very valuable tools to study exposure to a certain kind of compounds. Moreover, these assessments could help to trace toxicants in discharges to aquatic ecosystems, by monitoring both in laboratory and in natural conditions exposed animals [149, 201]
- e. Elemental composition using X-ray analysis on rotifers cuticles. This is an easy method that qualitatively determines different elements of interest, principally inorganic metals [48].
- f. Morphological analysis: These are changes induced by toxic exposure; although it could be controversial, it can be carried out by comparing morphological characters through image analysis. Because of rotifers phenotypic plasticity, the comparisons should be carefully performed to avoid misinterpretation of the results. However, it has been demonstrated, in *B. calyciflorus*, that morphological changes occurred after exposure to the fungicide vinclozolin and that abnormal and healthy animals are easily differentiated. However, the percentage of deformities is low, only 0.63% of 2868 organisms. Nevertheless, this is still significantly different to the control groups. Hence, morphometric analysis in rotifers could be a helpful tool to identify unrevealed targets of toxicants, and it might contribute to create a database for such effects and for several rotifers species to further comparisons among them, besides the likely identification process through image analysis [48].
- g. Aging in rotifers: because there is a great diversity in aging rates among species, geographical populations, and mutants within species, Smith and Snell [202] designed an experiment to follow rotifers longevity through 84 generations (about 1 year). Their results show that optimal growing conditions (e.g., constant food supply) altered life span and can reduce aging, which could be evolutionary adjustable, with selection working primarily on the length of the reproductive life span. Rotifers are considered good models to investigate the effect not only of toxicant on their life span but also other factors such as caloric restriction and the effect of vitamins. Thus, this represents a new field to incorporate studies with rotifers [203, 204].
- h. Hormones: Alvarado-Flores et al. [205] demonstrated the presence of some mammal-like proteins in the rotifer *B. calyciflorus* and concluded that it is necessary to generate more information about catecholaminergic and cholinergic systems in rotifers and the hormones related. Then it will be possible to assess their participation in mechanisms of detoxification and likely be used as toxicity models.

i. Genetic tools in rotifers and their applications

- RNA-seq: With mRNA-seq libraries for obligate parthenogenetic and cyclical parthenogenetic strains of the rotifer *B. calyciflorus*, it has been possible to identify genes specific to both modes of reproduction. Additionally, the studies performed by Hason et al. [206] allowed insights in the reproductive biology of obligated asexual bdelloid rotifers.
- Heat shock proteins (HSPs): The genes for these proteins synthesis are found from bacteria to higher eukaryotes and are related to functions like refolding denatured proteins due to stress that includes heat shock, reason for which they are called HSPs. Smith et al. [207] provided conclusive evidence that *hsp40*, *hsp60*, and *hsp70* are required for rotifer survival following heat stress, but that *hsp90* seems not to be essential for survival, at least with their data.
- Metallothionein (Mt): These are low-molecular weight and cysteine-rich proteins present in eukaryotes. They provide potent metal binding and some other functions are being discovered. Their presence in rotifers has been demonstrated as a consequence of chromium exposure [208]
- P-glycoprotein 8P-gp (Pg-p): This protein could be considered as the first line of defense against some chemicals, including pharmaceuticals and endocrine disruptors. This protein has been found and characterized in the rotifer *B. koreanus*. Specimens of this rotifer were exposed to several pharmaceuticals that retarded growth and promoted the overexpression of Pg-p [209].
- Epigenetics: Germ cells can be specified early in embryogenesis by maternal determinants inherited in the cytoplasm of the oocyte or they can be selected later in the embryonic development from undifferentiated precursors by a localized inductive signal that is called epigenesis [210]. Epigenetic processes were found in the ovary of *B. plicatilis*, describing the participation of *vasa* and *nos* genes. As the first description of its kind, it opens the possibilities to explore and perform embryo development within the phylum.
- Cell-penetrating peptides (CPPs): Liu et al. (2013) demonstrated that some little peptides can penetrate cell membranes and delivered their cargoes; then their function (if cargoes have it) can be assessed as the functional HR9-delivered plasmid DNAs and RFP coding sequences that could be actively expressed in rotifers. This method provides a tool not only for genetic material but also for nanoparticles and proteins, which in the future could facilitate studying the effect of chemicals within rotifer cells.

j. Innate immunity in rotifers: In invertebrates, the nomenclature, annotation, and reports of cytokines could be controversial. Nonetheless, there is a continuous and increasing knowledge about cytokine-mediated immune regulation, although adaptive immune responses are likely absent in invertebrates, including rotifers. In this field, Jeong et al. [211] identified three genes of lipopolysaccharide-induced TNF- α factor (LITAF) in *B. koreanus*. The *in silico* analysis showed that these genes could be involved in innate

immunity in primitive rotifers. In addition, exposure to lipopolysaccharide caused the overexpression of *LITAF1* and *LITAF2* but depleted glutathione concentration. Thereafter, *LITAF* genes have potential sensitivities to immune stimulator-triggered oxidative stress.

In conclusion, rotifers as models for ecotoxicological tests present several advantages, including a relative short life cycle that allows multigenerational studies and epigenetic research to unveil functions and processes in mictic and amictic rotifers, the simplicity of their body structure that ease the permeability of dyes for *in vivo* examination to quantify toxicant concentrations (e.g., Leadmium green®) or for systems descriptions like both the cholinergic and catecholaminergic systems, and their easy culture conditions and supply of resting eggs from different sources like some commercial brands. Nowadays, there are several protocols that describe the use of rotifers as indicators of water quality and safety as even thousands of probes can be performed in very short periods. Furthermore, every year, new technologies are becoming available to explore in deep detail the effect, the mechanisms, and the targets of toxicants. Hence, rotifer studies cannot be the exception, opening new possibilities to explore and describe more accurately the interaction of toxicants with the aquatic biota.

Author details

Roberto Rico-Martínez^{1*}, Mario Alberto Arzate-Cárdenas¹, Daniel Robles-Vargas¹, Ignacio Alejandro Pérez-Legaspi², Alvarado-Flores Jesús³ and Gustavo Emilio Santos-Medrano¹

*Address all correspondence to: rrico@correo.uaa.mx

¹ Departamento de Química, Centro de Ciencias Básicas, Universidad Autónoma de Aguascalientes, Aguascalientes, AGS, Mexico

² División de Estudios de Posgrado e Investigación, Instituto Tecnológico de Boca del Río, Boca del Río, Veracruz, Mexico

³ Catedrático CONACYT, Centro de Investigación Científica de Yucatán, Unidad de Ciencias del Agua, Cancún, Quintana Roo, México, Unidad de Ciencias del Agua, Cancún, Quintana Roo, Mexico

References

- [1] Shane, B. Introduction to Ecotoxicology. In: Cockerham LG, Shane BS, editors. Basic Environmental Toxicology. Boca Raton, Florida: CRC Press; 1994. pp. 3–10.
- [2] Rand, G.M. and Petrocelli, S.R. Fundamentals of Aquatic Toxicology. Hemisphere Publishing Corporation, New York. 1985. p. 666.

- [3] Snell, T.W. and Janssen C.R. Rotifers in ecotoxicology: a review. *Hydrobiologia*. 1995. 313/314: 231–247. DOI: 10.1007/978-94-009-1583-1_32
- [4] Wallace, R.L., Snell, T.W., Ricci, C. and Nogrady, T. Rotifera. 1: biology, ecology and systematics. In: Dumont HJF, editor. *Guides to the Identification of the Microinvertebrates of the World*. 2nd ed. The Netherlands: Kenobi Productions and Backhuys Publishers; 2006. p. 299.
- [5] Pérez-Legaspi, I.A., Quintanar, J.L. and Rico-Martínez, R. Comparing toxicity endpoints on *Lecane quadridentata* (Rotifera: Monogononta) exposed to two anticholinesterases pesticides. *Environment Toxicology*. 2012; 27, 518–525. DOI: 10.1002/tox.20668
- [6] Burbank, S. and Snell, T.W. Rapid toxicity assessment using esterase biomarkers in *Brachionus calyciflorus* (Rotifera). *Environment Toxicology and Water Quality*. 1994. 9: 171–178. DOI: 10.1002/tox.2530090303
- [7] Ferrando, M.D. and Andreu-Moliner E. Acute toxicity of toluene, hexane, xylene and benzene to the rotifers *Brachionus calyciflorus* and *Brachionus plicatilis*. *Bulletin of Environmental Contamination and Toxicology*. 1992. 49: 266 –271. DOI: 10.1007/BF00191765
- [8] Snell, T.W. and Janssen, C.R. Microscale toxicity testing with rotifers. In: Wells PG, Lee K, Blaise C, editors. *Microscale Testing in Aquatic Toxicology, Advances, Techniques and Practice*. Boca Raton, Florida: CRC Press. 1998. p 409–422.
- [9] Pennak, R.W. *Freshwater invertebrates of the United States*. 3rd ed. New York: Wiley; 1989. 628 p.
- [10] Segers, H. The nomenclature of the Rotifera: Annotated checklist of valid family-and genus-group names. *Journal of Natural History*. 2002. 36: 631–640. DOI: 10.1080/00222930010024573
- [11] Pérez-Legaspi, I.A. Estudio de los efectos de dos pesticidas en el sistema colinérgico en especies selectas de rotíferos dulceacuícolas [thesis]. Aguascalientes: Universidad Autónoma de Aguascalientes; 2008.
- [12] Clemént, P. and Wurdak, E. Rotifera. In: Harrison FW, Ruppert E, editors. *Microscopy Anatomy of Invertebrates*. Vol. 4: Aschelminthes. New York: Wiley-Liss; 1991. pp. 219–297.
- [13] Klimek, B., Fialkowska, E., Kocerba-Soroka, W., Fida, J., Sobczyk, M. and Pajdak-Stós, A. The toxicity of selected trace metals to *Lecane inermis* rotifers isolated from activated sludge. *Bulletin of Environmental Contamination and Toxicology*. 2013. 91: 330–333. DOI: 10.1007/s00128-013-1062-z
- [14] Snell, T.W. and Moffat, B.D. A 2-d life cycle test with the rotifer *Brachionus calyciflorus*. *Environmental Toxicology and Chemistry*. 1992. 11: 1249–1257. DOI: 10.1016/0147-6513(91)90070-6

- [15] Peltier, W.H. and Weber, C.I., editors. Methods for measuring the acute toxicity of effluents to freshwater and marine organisms. EPA-600/4-85-013. Cincinnati, Ohio: US Environmental Protection Agency; 1985. 231 p.
- [16] Hoff, F.H. and Snell T.W. Plankton culture manual. 3rd ed. Dade City, Florida, USA: Florida Aqua Farms; 2008. 186 p.
- [17] Yúfera, M. Studies on *Brachionus* (Rotifera): an example of interaction between fundamental and applied research. *Hydrobiologia*. 2001. 446/447: 383–392. DOI: 10.1023/A:1017583729646
- [18] Kostopoulou, V., Carmona, M.J. and Divanach P. The rotifer *Brachionus plicatilis*: an emerging bio-tool for numerous applications. *Journal of Biological Research–Thessaloniki*. 2012. 17: 97–112.
- [19] Pérez-Legaspi, I.A. and Rico-Martínez R. Effect of temperature and food concentration in two species of littoral rotifers. *Hydrobiologia*. 1998. 387/388: 341–348. DOI: 10.1023/A:1017099906853
- [20] Wallace, R.L. Rotifers: Exquisite metazoans. *Integrative and Comparative Biology*. 2002. 42: 660–667. DOI: 10.1093/icb/42.3.660
- [21] Alvarado-Flores, J. Detección del efecto de disruptores endocrinos y metales en la reproducción de rotíferos dulceacuícolas. [thesis]. Aguascalientes: Universidad Autónoma de Aguascalientes; 2014.
- [22] Dahms, H.U., Hagiwara, A. and Lee, J.S. Ecotoxicology, ecophysiology, and mechanistic studies with rotifers. *Aquatic Toxicology*. 2011. 101 (1): 1–12. DOI: 10.1016/j.aquatox.2010.09.006
- [23] Snell, T.W. and Persoone, G. 1989. Acute toxicity bioassays using rotifers. II. A freshwater test with *Brachionus rubens*. *Aquatic Toxicology*. 14 (1): 81–92. DOI: 10.1016/0166-445X(89)90056-8
- [24] Janssen, C.R., Rodrigo, M.D.F. and Persoone, G. Ecotoxicological studies with the freshwater rotifer *Brachionus calyciflorus*. I. Conceptual framework and applications. *Hydrobiologia*. 1993. 255/256: 21–32. DOI: 10.1007/BF00025817
- [25] Nogrady, T., Wallace, R.L. and Snell, T.W. Rotifera. In: Dumont, editor. *Biology, Ecology and Systematics*. Volume 4: Guides to the identification of the microinvertebrates of the continental waters of the world. The Netherlands: SPB Academic Publishing; 1993. p. 142.
- [26] Marcial, H.S., Hagiwara, A. and Snell, T.W. Effect of some pesticides on reproduction of rotifer *Brachionus plicatilis* Müller. *Hydrobiologia*. 2005. 546: 569–575.
- [27] Ruppert, E.E., Fox, R.S. and Barnes, R.D. *Invertebrate zoology: a functional evolutionary approach*, 7th ed. Thomson-Brooks/Cole, Belmont. 2004. pp. 617–625, 784–789

- [28] Thorp, J.H. and Rogers, D.C. Field Guide to Freshwater Invertebrates of North America. Academic Press. Elsevier. San Diego, California. USA. 2010. pp. 109–119
- [29] De Walt, R.W., Resh, V.H. and Hilsenhoff, W.L. Diversity and Classification of Insects and Collembola. In: Thorp JH. Ecology and Classification of North American Freshwater Invertebrates. Academic Press. Elsevier. San Diego. California. USA. 2010. pp. 587–593
- [30] Hutchinson, T.H. Reproductive and developmental effects of endocrine disruptors in invertebrates: in vitro and in vivo approaches. Toxicology Letters. 2002. 131: 75–81. DOI:10.1016/S0378-4274(02)00046-2
- [31] Baun, A., Hartmann, N.B., Grieger, K. and Kusk, K.O. Ecotoxicity of engineered nanoparticles to aquatic invertebrates: a brief review and recommendations for future toxicity testing. Ecotoxicology. 2008. 17(5):387–95. 10.1007/s10646-008-0208-y
- [32] Cattaneo, A.G., Gornati, R., Chiriva-Internati, M. and Bernardini, G. Ecotoxicology of nanomaterials: the role of invertebrate testing. Invertebrate Survival Journal. 2009. 6: 78–97.
- [33] U.S. EPA-821-R-02-012. Methods for measuring the acute toxicity of effluents and receiving waters to freshwater and marine organisms. 5th ed. 2002. http://water.epa.gov/scitech/methods/cwa/wet/upload/2007_07_10_methods_wet_disk2_atx.pdf
- [34] U.S. EPA-821-R-02-013. Short-Term Methods for Estimating the Chronic Toxicity of Effluents and Receiving Water to Freshwater Organisms. 4th ed. 2002. http://water.epa.gov/scitech/methods/cwa/wet/upload/2007_07_10_methods_wet_disk3_ctf.pdf
- [35] U.S. Environmental Protection Agency. Dispersant Monitoring and Assessment Final Directive for Subsurface Dispersant Application [Internet]. 2010. Available from: <http://archive.epa.gov/bpspill/web/pdf/subsurface-dispersant-directive-final.pdf>. [Accessed: 2015-09-03]
- [36] Rico-Martínez, R., Snell, T.W. and Shearer, T.L. Synergistic toxicity of Macondo crude oil and dispersant Corexit 9500A(®) to the *Brachionus plicatilis* species complex (Rotifera). Environmental Pollution. 2013. 173: 5–10. DOI:10.1016/j.envpol.2012.09.024
- [37] Gourmelon, A. and Ahtiainen, J. Developing Test Guidelines on invertebrate development and reproduction for the assessment of chemicals, including potential endocrine active substances—the OECD perspective. Ecotoxicology. 2007. 16: 161–167. DOI: 10.1007/s10646-006-0105-1
- [38] Di Lorenzo, T., Di Marzio, W.D., Sáenz, M.E., Baratti, M., Dedonno, A.A., Iannucci, A., Cannicci, S., Messana, G. and Galassi, D.M. Sensitivity of hypogean and epigean freshwater copepods to agricultural pollutants. Environmental Science and Pollution Research International. Mar. 2014. 21 (6): 4643–455. DOI: 10.1007/s11356-013-2390-6

- [39] Cifoni, M., Di Lorenzo, T., Faraloni, C. and Galassi D.M.P. A protocol for a development, reproduction and population growth test with freshwater copepods. Conference Paper. European Project AQUALIFE (LIFE12 BIO/IT/000231). 2015. International Conference on Groundwater in Karst. <http://www.birmingham.ac.uk/generic/kgatb/index.aspx>
- [40] Gilbert, J.J. and Birky, C.W. Sensitivity and specificity of the *Asplanchna* response to dietary alpha-tocopherol. *Journal of Nutrition*. 1971. 101 (1): 113–126.
- [41] Stelzer, C.P. Population regulation in sexual and asexual rotifers: an eco-evolutionary feedback to population size? *Functional Ecology*. 2012. 26: 180–188. DOI:10.1111/j.1365-2435.2011.01918.x
- [42] Wallace., R.L. and Smith, H.A. Rotifera. In: eLS. John Wiley & Sons, Ltd: Chichester. 2013. DOI: 10.1002/9780470015902.a0001588.pub2
- [43] ASTM E1440-91. Standard Guide for Acute Toxicity Test with the Rotifer *Brachionus*, ASTM International, West Conshohocken, PA. 2012. DOI: 10.1520/E1440-91R12. www.astm.org
- [44] ASTM E2317-04. Standard Guide for Conducting Renewal Microplate-Based Life-Cycle Toxicity Tests with a Marine Meiobenthic Copepod, ASTM International, West Conshohocken, PA, 2012. DOI: 10.1520/E2317-04R12. www.astm.org
- [45] ASTM STP921-EB. Aquatic Toxicology and Environmental Fate: Ninth Volume. 1986. DOI: 10.1520/STP921-EB. www.astm.org
- [46] Quinn, B., Gagne, F. and Blaise, C. Validation of a multi-well plate toxicity test to assess feeding behavior of the cnidarian, *Hydra attenuata*. *Fresenius Environmental Bulletin*. 2007. 16: 1100–1107.
- [47] Santos, M.A.P.F., Vicensotti, J. and Monteiro, R.T.R. Sensitivity of four test organisms (*Chironomus xanthus*, *Daphnia magna*, *Hydra attenuata* and *Pseudokirchneriella subcapitata*) to NaCl: an alternative reference toxicant. *Journal of the Brazilian Society of Ecotoxicology*. 2007. 2 (3): 229–236. DOI: 10.5132/jbse.2007.03.004
- [48] Alvarado-Flores, J., Rico-Martínez, R., Adabache-Ortiz, A. and Silva-Briano, M. Morphological alterations in the freshwater rotifer *Brachionus calyciflorus* Pallas 1766 (Rotifera: Monogononta) caused by vinclozolin chronic exposure. *Ecotoxicology*. 2015. 24 (4): 915–925. DOI: 10.1007/s10646-015-1434-8
- [49] Milam, C.D., Farris, J.L., Dwyer, F.J. and Hardesty, D.K. Acute toxicity of six freshwater mussel species (Glochidia) to six chemicals: implications for Daphnids and *Uterbackia imbecillis* as surrogates for protection of freshwater mussels (Unionidae). *Archives of Environmental Contamination and Toxicology*. 2005. 48 (2): 166–173. DOI: 10.1007/s00244-003-3125-3
- [50] Besser, J.M., Ingersoll, C.G., Brumbaugh, W.G., Kemble, N.E., May, T.W., Wang, N., MacDonald, D.D. and Roberts, A.D. Toxicity of sediments from lead-zinc mining

- areas to juvenile freshwater mussels (*Lampsilis siliquoidea*) compared to standard test organisms. *Environmental Toxicology and Chemistry*. 2015. 34 (3): 626–639. DOI: 10.1002/etc.2849
- [51] Ringwood, A.H. Comparative sensitivity of gametes and early developmental stages of a sea urchin species (*Echinometra mathaei*) and a bivalve species (*Isognomon californicum*) during metal exposures. *Archives of Environmental Contamination and Toxicology*. 1992. 22 (3): 288–295. DOI: 10.1007/BF00212088
- [52] Golding, C., Krassoi, R. and Baker, E. The development and application of a marine toxicity identification evaluation (TIE) protocol for use with an Australian bivalve. *Australasian Journal of Ecotoxicology*. 2006. 12: 37–44.
- [53] Cheung, V.V., Depledge, M.H. and Jha, A.N. An evaluation of the relative sensitivity of two marine bivalve mollusc species using the Comet assay. *Marine Environmental Research*. 2006. 62 Suppl:S301-5. DOI: 10.1016/j.marenvres.2006.04.053
- [54] Gerdol, M., De Moro, G., Manfrin, C., Milandri, A., Riccardi, E., Beran, A., Venier, P. and Pallavicini, A. RNA sequencing and *de novo* assembly of the digestive gland transcriptome in *Mytilus galloprovincialis* fed with toxinogenic and non-toxic strains of *Alexandrium minutum*. *BMC Research Notes*. 2014. 7: 722. DOI: 10.1186/1756-0500-7-722
- [55] Akiyama, Y. and Iwakuma, T. Survival of glochidial larvae of the freshwater pearl mussel, *Margaritifera laevis* (Bivalvia: Unionoida), at different temperatures: a comparison between two populations with and without recruitment. *Zoological Science*. 2007. 24 (9): 890–893. DOI: 10.2108/zsj.24.890
- [56] Dodd, B.J., Barnhart, M.C., Rogers-Lowery, C.L., Fobian, T.B. and Dimock, R.V. Jr. Persistence of host response against glochidia larvae in *Micropterus salmoides*. *Fish and Shellfish Immunology*. 2006. 21 (5): 473–484. DOI: 10.1016/j.fsi.2006.02.002
- [57] Blakeslee, C.J., Galbraith, H.S., Robertson, L.S., and St John White, B. The effects of salinity exposure on multiple life stages of a common freshwater mussel, *Elliptio complanata*. *Environmental Toxicology and Chemistry*. 2013. 32 (12): 2849–2854. DOI: 10.1002/etc.2381
- [58] ASTM E2455-06. Standard Guide for Conducting Laboratory Toxicity Tests with Freshwater Mussels. ASTM International, West Conshohocken, PA. 2013. DOI: 10.1520/E2455-06R13. www.astm.org
- [59] Bat, L. and Raffaelli, D. Sediment toxicity testing: a bioassay approach using the amphipod *Corophium volutator* and the polychaete *Arenicola marina*. *Journal of Experimental Marine Biology and Ecology*. 1998. 226: 217–239. DOI: 10.1016/S0022-0981(97)00249-9

- [60] Lewis, C., Pook, C. and Galloway, T. Reproductive toxicity of the water accommodated fraction (WAF) of crude oil in the polychaetes *Arenicola marina* (L.) and *Nereis virens* (Sars). *Aquatic Toxicology*. 2008. 90: 73–81. DOI: 10.1016/j.aquatox.2008.08.001.
- [61] Besseling, E., Wegner, A., Foekema, E.M., van den Heuvel-Greve, M.J., Koelmans, A.A. Effects of microplastic on fitness and PCB bioaccumulation by the lugworm *Arenicola marina* (L.). *Environmental Science and Technology*. 2013. 47 (1): 593–600. DOI: 10.1021/es302763x.
- [62] EPA 850.1010. Aquatic Invertebrate Acute Toxicity Test, Freshwater Daphnids. http://www.epa.gov/ocspp/pubs/frs/publications/OPPTS_Harmonized/850_Ecological_Effects_Test_Guidelines/Drafts/850-1010.pdf
- [63] OECD. Test No. 202: *Daphnia* sp. Acute Immobilisation Test, OECD Guidelines for the Testing of Chemicals, Section 2, OECD Publishing, Paris. 2004. DOI: 10.1787/9789264069947-en
- [64] ASTM E1295-01. Standard Guide for Conducting Three-Brood, Renewal Toxicity Tests with *Ceriodaphnia dubia*. ASTM International, West Conshohocken, PA. 2013. DOI: 10.1520/E1295-01R13 www.astm.org
- [65] ISO 6341:2012. Water Quality. Determination of the Inhibition of the Mobility of *Daphnia magna* Straus (Cladocera, Crustacea). Acute Toxicity Test. International Organization for Standardization. Geneva, Switzerland. 2012. http://www.iso.org/iso/iso_catalogue/catalogue_tc/catalogue_detail.htm?csnumber=54614
- [66] EPA 850.1300. Short-Term Methods for Estimating the Chronic Toxicity of Effluents and Receiving Waters to Freshwater Organisms. 4th ed. 2002. http://water.epa.gov/scitech/methods/cwa/wet/upload/2007_07_10_methods_wet_disk3_ctf.pdf
- [67] OECD. Test No. 211: *Daphnia magna* Reproduction Test, OECD Guidelines for the Testing of Chemicals, Section 2. OECD Publishing, Paris. 2012. DOI: 10.1787/9789264185203-en
- [68] ASTM E1193-97. Standard Guide for Conducting *Daphnia magna* Life-Cycle Toxicity Tests. ASTM International, West Conshohocken, PA. 2012. DOI: 10.1520/E1193-97R12. www.astm.org
- [69] CETESB. Métodos de avaliação da toxicidade de poluentes a organismos aquáticos. São Paulo. 1992.
- [70] NSW EPA. Cladoceran toxicity testing. Laboratory Procedures Manual. 1999. 5: 17.
- [71] NSW EPA. Chronic reproduction impairment test. Laboratory Procedures Manual. 2003. 6: 16
- [72] ISO 20665. Water quality. Determination of chronic toxicity to *Ceriodaphnia dubia*. International Organization for Standardization. Geneva, Switzerland. 2008. http://www.iso.org/iso/catalogue_detail.htm?csnumber=39673

- [73] Sepic, E., Bricelj, M. and Leskovsek, H. Toxicity of fluoranthene and its biodegradation metabolites to aquatic organisms. *Chemosphere*. 2003. 52 (7): 1125–1133. DOI: 10.1016/S0045-6535(03)00321-7
- [74] Heinlaan, M., Ivask, A., Blinova, I., Dubourguier, H.C. and Kahru, A. Toxicity of nanosized and bulk ZnO, CuO and TiO₂ to bacteria *Vibrio fischeri* and crustaceans *Daphnia magna* and *Thamnocephalus platyurus*. *Chemosphere*. 2008. 71 (7): 1308–1316. DOI: 10.1016/j.chemosphere.2007.11.047.
- [75] Brausch, J.M. and Smith, P.N. Mechanisms of resistance and cross-resistance to agrochemicals in the fairy shrimp *Thamnocephalus platyurus* (Crustacea: Anostraca). *Aquatic Toxicology*. 2009. 92 (3): 140–145. DOI: 10.1016/j.aquatox.2009.02.002
- [76] Blinova, I., Niskanen, J., Kajankari, P., Kanarbik, L., Käkinen, A., Tenhu, H., Penttinen, O.P. and Kahru, A. Toxicity of two types of silver nanoparticles to aquatic crustaceans *Daphnia magna* and *Thamnocephalus platyurus*. *Environmental Science and Pollution Research International*. 2013. 20 (5): 3456–3463. DOI: 10.1007/s11356-012-1290-5.
- [77] Ger KA, Teh SJ, Goldman CR. Microcystin-LR toxicity on dominant copepods *Eurytemora affinis* and *Pseudodiaptomus forbesi* of the upper San Francisco Estuary. *Science of the Total Environment*. 2009. 407 (17): 4852–4857. DOI: 10.1016/j.scitotenv.2009.05.043
- [78] Cailleaud, K., Budzinski, H., Lardy, S., Augagneur, S., Barka, S., Souissi, S. and Forget-Leray, J. Uptake and elimination, and effect of estrogen-like contaminants in estuarine copepods: an experimental study. *Environmental Science and Pollution Research International*. 2011. 18 (2): 226–236. DOI: 10.1007/s11356-010-0355-6
- [79] Michalec, F.G., Holzner, M., Menu, D., Hwang, J.S. and Souissi, S. Behavioral responses of the estuarine calanoid copepod *Eurytemora affinis* to sub-lethal concentrations of waterborne pollutants. *Aquatic Toxicology*. 2013. 15: 138–139:129–138. DOI: 10.1016/j.aquatox.2013.05.007
- [80] Leseur, T., Boulangé-Lecomte, C., Restoux, G., Deloffre, J., Xuereb, B., Le Menach, K., Budzinski, H., Petrucciani, N., Marie, S., Petit, F. and Forget-Leray, J. Toxicity of sediment-bound pollutants in the Seine estuary, France, using a *Eurytemora affinis* larval bioassay. *Ecotoxicology Environmental Safety*. 2015. 113:169–175. DOI: 10.1016/j.ecoenv.2014.11.033
- [81] Sánchez-Bayo, F. Comparative acute toxicity of organic pollutants and reference values for crustaceans. I. Branchiopoda, Copepoda and Ostracoda. *Environmental Pollution*. 2006. 139 (3): 385–420. DOI: 10.1016/j.envpol.2005.06.016
- [82] Nandini, S. and Sarma, S.S.S. Effect of algal and animal diets on life history of the freshwater copepod *Eucyclops serrulatus* (Fischer, 1851). *Aquatic Ecology*. 2007. 41 (1): 75–84. DOI: 10.1007/s10452-006-9051-5
- [83] Di Lorenzo, T., Di Marzio, W.D., Spigoli, D., Baratti, M., Messina, G., Cannicci, S. and Galassi, D.M.P. Metabolic rates of a hypogean and an epigean species of cope-

- pod in an alluvial aquifer. *Freshwater Biology*. 2015. 60: 426–435. DOI: 10.1111/fwb.12509
- [84] ISO 20666. Water quality. Determination of the chronic toxicity to *Brachionus calyciflorus* in 48 h. International Organization for Standardization. Geneva, Switzerland. 2008. http://www.iso.org/iso/catalogue_detail.htm?csnumber=39674
- [85] Arauzo, M. and Valladolid, M. Short-term harmful effects of unionised ammonia on natural populations of *Moina micrura* and *Brachionus rubens* in a deep waste treatment pond. *Water Research*. 2003. 37 (11): 2547–2554. DOI: 10.1016/S0043-1354(03)00023-X
- [86] Azuara-García, R., Sarma, S.S.S. and Nandini S. The combined effects of zinc and alga on the life table demography of *Anuraeopsis fissa* and *Brachionus rubens* (Rotifera). *Journal of Environmental Science and Health. Part A, Toxic/Hazardous Substances and Environmental Engineering*. 2006. 41 (4): 559–572. DOI: 10.1080/10934520600564394
- [87] Geng, H. and Xie, P. Experimental studies on the effects of toxic *Microcystis aeruginosa* PCC7820 on the survival and reproduction of two freshwater rotifers *Brachionus calyciflorus* and *Brachionus rubens*. *Ecotoxicology*. 2008. 17 (8): 709–715. DOI: 10.1007/s10646-008-0219-8
- [88] Sarma, S.S.S., Ramírez-Pérez, T. and Nandini, S. Comparison of the sensitivity of *Brachionus calyciflorus* and *Brachionus patulus* (Rotifera) to selected heavy metals under low and high food (*Chlorella vulgaris*) levels. *Bulletin of Environmental Contamination and Toxicology*. 2000. 64 (5): 735–739. DOI: 10.1007/s001280000065
- [89] Sarma, S.S.S., Nandini, S. and Pérez, T.R. Combined effects of mercury and algal food density on the population dynamics of *Brachionus patulus* (Rotifera). *Bulletin of Environmental Contamination and Toxicology*. 2001. 67 (6): 841–847. DOI: 10.1007/s001280199
- [90] Peredo-Alvarez, V.M., Sarma, S.S. and Nandini, S. Combined effect of concentrations of algal food (*Chlorella vulgaris*) and salt (sodium chloride) on the population growth of *Brachionus calyciflorus* and *Brachionus patulus* (Rotifera). *Revista de Biología Tropical*. 2003. 51 (2): 399–407.
- [91] Hernández-Flores, S. and Rico-Martínez, R. Study of the effects of Pb and Hg toxicity using a chronic toxicity reproductive 5-day test with the freshwater rotifer *Lecane quadridentata*. *Environmental Toxicology*. 2006. 21 (5): 533–540. DOI: 10.1002/tox.20218
- [92] Domínguez-Cortinas, G., Mejía-Saavedra, J., Santos-Medrano, G.E. and Rico-Martínez, R. Analysis of the toxicity of glyphosate and Faena® using the freshwater invertebrates *Daphnia magna* and *Lecane quadridentata*. *Toxicological and Environmental Chemistry*. 2008. 90 (2): 377–384. DOI: 10.1080/02772240701529038
- [93] Torres-Guzmán, F., Avelar-González, F.J. and Rico-Martínez, R. Implementing *Lecane quadridentata* acute toxicity tests to assess the toxic effects of selected metals (Al, Fe

- p>and Zn).
- Ecotoxicology and Environmental Safety*
- . 2010. 73 (3): 287–295. DOI: 10.1016/j.ecoenv.2009.10.006
- [94] Pérez-Legaspi, I.A. and Rico-Martínez, R. Acute toxicity tests on three species of the genus *Lecane* (Rotifera: Monogononta). *Hydrobiologia*. 2001. 446/447 (1): 375–381. DOI: 10.1023/A:1017531712808
- [95] Pérez-Legaspi, I.A., Rico-Martínez, R. and Pineda-Rosas, A. Toxicity testing using esterase inhibition as a biomarker in three species of the genus *Lecane* (Rotifera). *Environmental Toxicology and Chemistry*. 2002. 21 (4): 776–782. DOI: 10.1002/etc.5620210413
- [96] Arias-Almeida, J.C., Rico-Martínez, R. Inhibition of two enzyme systems in *Euchlanis dilatata* (Rotifera: Monogononta) as biomarker of effect of metals and pesticides. *Biomarkers*. 2011. 16 (1): 12–9. DOI: 10.3109/1354750X.2010.520336
- [97] Arias-Almeida, J.C. and Rico-Martínez, R. Toxicity of cadmium, lead, mercury and methyl parathion on *Euchlanis dilatata* Ehrenberg 1832 (Rotifera: Monogononta). *Bulletin of Environmental Contamination and Toxicology*. 2011. 87 (2): 138–1342. DOI: 10.1007/s00128-011-0308-x
- [98] Gómez, A., Cecchine, G. and Snell, T.W. Effect of pentachlorophenol on predator-prey interaction of two rotifers. *Aquatic Toxicology*. 1997. 37 (4): 271–282. DOI: 10.1016/S0166-445X(96)00833-8
- [99] Gama-Flores, J.L., Ferrara-Guerrero, M.J., Sarma, S.S. and Nandini, S. Prey (*Brachionus calyciflorus* and *Brachionus havanaensis*) exposed to heavy metals (Cu and Cd) for different durations and concentrations affect predator's (*Asplanchna brightwellii*) population growth. *Journal of Environmental Science and Health. Part A, Toxic/Hazardous Substances and Environmental Engineering*. 2007. 42 (10): 1483–1488. DOI: 10.1080/10934520701480912
- [100] Guo, R., Snell, T.W. and Yang, J. Studies of the effect of environmental factors on the rotifer predator–prey system in freshwater. *Hydrobiologia*. 2010. 655 (1): 49–60. DOI: 10.1007/s10750-010-0403-8
- [101] Chen, J., Wang, Z., Li, G. and Guo, R. The swimming speed alteration of two freshwater rotifers *Brachionus calyciflorus* and *Asplanchna brightwelli* under dimethoate stress. *Chemosphere*. 2014. 95: 256–260. DOI: 10.1016/j.chemosphere.2013.08.086
- [102] Karntanut, W., Pascoe, D. A comparison of methods for measuring acute toxicity to *Hydra vulgaris*. *Chemosphere*. 2000. 41 (10): 1543–1548. DOI: 10.1016/S0045-6535(00)00068-0
- [103] Pascoe, D., Carroll, K., Karntanut, W. and Watts, M.M. Toxicity of 17alpha-ethinylenestradiol and bisphenol A to the freshwater Cnidarian *Hydra vulgaris*. *Archives of Environmental Contamination and Toxicology*. 2002. 43 (1): 56–63. DOI: 10.1007/s00244-001-0016-3

- [104] Pascoe, D., Karntanut, W. and Müller, C.T. Do pharmaceuticals affect freshwater invertebrates? A study with the cnidarian *Hydra vulgaris*. *Chemosphere*. 2003. 51 (6): 521–528. DOI: 10.1016/S0045-6535(02)00860-3
- [105] Cataldo, D., Colombo, J.C., Boltovskoy, D., Bilos, C. and Landoni, P. Environmental toxicity assessment in the Paraná river delta (Argentina): simultaneous evaluation of selected pollutants and mortality rates of *Corbicula fluminea* (Bivalvia) early juveniles. *Environmental Pollution*. 2001. 112 (3): 379–389. DOI: 10.1016/S0269-7491(00)00145-7
- [106] Achard, M., Baudrimont, M., Boudou, A., Bourdineaud, J.P. Induction of a multixenobiotic resistance protein (MXR) in the Asiatic clam *Corbicula fluminea* after heavy metals exposure. *Aquatic Toxicology*. 2004. 67 (4): 347–357. DOI: 10.1016/j.aquatox.2004.01.014
- [107] Cherry, D.S., Scheller, J.L., Cooper, N.L. and Bidwell, J.R. Potential effects of Asian clam (*Corbicula fluminea*) die-offs on native freshwater mussels (Unionidae) I: water-column ammonia levels and ammonia toxicity. *Journal of the North American Benthological Society*. 2005. 24 (2): 369–380. DOI: 10.1899/04-073.1
- [108] Liao, C.M., Jou, L.J., Lin, C.M., Chiang, K.C., Yeh, C.H. and Chou, B.Y. Predicting acute copper toxicity to valve closure behavior in the freshwater clam *Corbicula fluminea* supports the biotic ligand model. *Environmental Toxicology*. 2007. 22 (3): 295–307. DOI: 10.1002/tox.20263
- [109] ASTM E1706-05. Standard Test Method for Measuring the Toxicity of Sediment-Associated Contaminants with Freshwater Invertebrates. ASTM International, West Conshohocken, PA. 2010. DOI: 10.1520/E1706-05R10. www.astm.org
- [110] EPA OPPTS 850.1735. Whole Sediment Acute Toxicity Invertebrates, Freshwater. 1996. http://www.epa.gov/ocspp/pubs/frs/publications/OPPTS_Harmonized/850_Ecological_Effects_Test_Guidelines/Drafts/850-1735.pdf
- [111] EPA 600/R99.064 Methods for Measuring the Toxicity and Bioaccumulation of Sediment-associated Contaminants with Freshwater Invertebrates. 2000. <http://water.epa.gov/polwaste/sediments/cs/freshfact.cfm>
- [112] EPS 1/RM/33. Environment Canada. Biological Test Method: Test for Survival and Growth in Sediment Using the Freshwater Amphipod *Hyaella azteca*. 2012. 184 p. <http://ec.gc.ca/Publications/4BA9E8C1-30EA-48AE-B593-F8F763EED8F0/RM33---accessible.PDF>
- [113] ASTM E1688-10. Standard Guide for Determination of the Bioaccumulation of Sediment-Associated Contaminants by Benthic Invertebrates, ASTM International, West Conshohocken, PA, 2010. DOI: 10.1520/E1688-10. www.astm.org
- [114] EPA 823-f-00-002. Methods for Measuring the Toxicity and Bioaccumulation of Sediment-associated Contaminants with Freshwater Invertebrates—Second Edition Fact Sheet. 2000. 2 p. <http://www.epa.gov/ostwater/cs/freshfact.html>

- [115] OECD Test No. 225: Sediment–Water *Lumbriculus Toxicity* Test Using Spiked Sediment, OECD Guidelines for the Testing of Chemicals. Section 2. OECD Publishing. Paris. 2007. DOI: 10.1787/9789264067356-en
- [116] White, D.S. and Keilty, T.J. Burrowing avoidance assays of contaminated Detroit River sediments, using the freshwater Oligochaete *Stylodrilus heringianus* (Lumbriculidae). Archives of Environmental Contamination and Toxicology. 1998. 17 (5): 673–681. DOI: 10.1007/BF01055837
- [117] Keilty, T.J., White, D.S. and Landrum, P.F. Sublethal responses to endrin in sediment by *Stylodrilus heringianus* (Lumbriculidae) as measured by a 137cesium marker layer technique. Aquatic Toxicology. 1988. 13 (3): 251–269. DOI: 10.1016/0166-445X(88)90056-2
- [118] Moreira, R.A., da Silva Mansano, A. and Rocha, O. 2015. The toxicity of carbofuran to the freshwater rotifer, *Philodina roseola*. Ecotoxicology. 24 (3): 604–615. DOI: 10.1007/s10646-014-1408-2
- [119] EPA OPPTS 850.1790. Chironomid sediment toxicity test. Ecological Effects Test Guidelines. 1996. http://www.epa.gov/ocspp/pubs/frs/publications/OPPTS_Harmonized/850_Ecological_Effects_Test_Guidelines/Drafts/850-1790.pdf
- [120] OECD. Test No. 235: *Chironomus* sp., Acute Immobilisation Test. OECD Guidelines for the Testing of Chemicals, Section 2. OECD Publishing, Paris. 2011. DOI: 10.1787/9789264122383-en
- [121] EPS 1/RM/32. Environment Canada. Biological Test Method: Test for Survival and Growth in Sediment using the Larvae of Freshwater Midges *Chironomus tentans* or *Chironomus riparius*. 1997. 150 p. <http://publications.gc.ca/site/eng/460451/publication.html>
- [122] EPA 821-R-02-014 Short-term Methods for Estimating the Chronic Toxicity of Effluents and Receiving Waters to Marine and Estuarine Organisms. 3rd ed. 486 p. http://water.epa.gov/scitech/methods/cwa/wet/upload/2007_07_10_methods_wet_disk1_ctm.pdf
- [123] EPA/600/R-09/121 *Americamysis bahia* Stochastic Matrix Population Model for Laboratory Populations: Technical Documentation. 2009. <http://nepis.epa.gov/Exe/ZyPDF.cgi/P1005OBH.PDF?Dockkey=P1005OBH.PDF>
- [124] ISO 16778:2015. Water quality. Calanoid copepod early-life stage test with *Acartia tonsa*. International Organization for Standardization. Geneva, Switzerland. 2015. http://www.iso.org/iso/catalogue_detail.htm?csnumber=57698
- [125] ASTM E2122-02. Standard Guide for Conducting In-situ Field Bioassays with Caged Bivalves. ASTM International, West Conshohocken, PA, 2013. DOI: 10.1520/E2122-02R13. www.astm.org

- [126] ASTM E724-98. Standard Guide for Conducting Static Acute Toxicity Tests Starting with Embryos of Four Species of Saltwater Bivalve Molluscs. ASTM International, West Conshohocken, PA. 2012. DOI: 10.1520/E0724-98R12. www.astm.org
- [127] EPA 850.1055. Bivalve acute toxicity test (embryo larval). Ecological Effects Test. 1996. 7p. http://www.epa.gov/ocspp/pubs/frs/publications/OPPTS_Harmonized/850_Ecological_Effects_Test_Guidelines/Drafts/850-1055.pdf
- [128] Thain, J. and Bifield, S. Biological effects of contaminants: Sediment bioassay using the polychaete *Arenicola marina*. ICES Techniques in Marine Environmental Sciences, 2001. 29: 16 pp.
- [129] ASTM E1367-03. Standard Test Method for Measuring the Toxicity of Sediment-Associated Contaminants with Estuarine and Marine Invertebrates. ASTM International, West Conshohocken, PA. 2014. DOI: 10.1520/E1367-03R14. www.astm.org
- [130] EPA 600-R-01-020. Methods for Assessing the Chronic Toxicity of Marine and Estuarine Sediment-associated Contaminants with the Amphipod *Leptocheirus plumulosus*. 1st ed. 2001. 121 pp. <http://water.epa.gov/polwaste/sediments/cs/upload/guidance-manual.pdf>
- [131] Zadorozhnaya, O., Kirsanov, D., Buzhinsky, I., Tsarev, F., Abramova, N., Bratov, A., Javier-Muñoz, F., Ribó, J., Bori, J., Carmen-Riva, M. and Legin, A. Water pollution monitoring by an artificial sensory system performing in terms of *Vibrio fischeri* bacteria. Sensors and Actuators B. 2015. 207: 1069–1075. DOI: 10.1016/j.snb.2014.08.056
- [132] Janssen, C. R., Ferrando-Rodrigo, M. D. and Persoone, G. Ecotoxicological studies with the freshwater rotifer *Brachionus calyciflorus*. I. Conceptual framework and applications. Hydrobiologia. 1993. 255/256: 21–32. DOI: 10.1007/978-94-011-1606-0_4
- [133] Vaal, M., van der Wal, J. T., Hermens, J. and Hoekstra, J. Pattern analysis of the variation in the sensitivity of aquatic species to toxicants. Pergamon. 1997. 35 (6): 1291–1309. DOI: 10.1016/S0045-6535(97)00166-5
- [134] Fernández-Casalderrey, A., Ferrando, M. D. and Andreu-Moliner, E. Acute toxicity of several pesticides to rotifer (*Brachionus calyciflorus*). Bulletin of Environmental Contamination and Toxicology. 1992a. 48: 14–17. DOI: 10.1007/BF00197477
- [135] PAN Pesticides Database. Chemical Toxicity Studies on Aquatic Organisms. Aquatic toxicity. Available from: <http://www.pesticideinfo.org/> [Accessed: 2015-08-10]
- [136] Kim, B. M., Lee, J. W., Seo, J. S., Shin, K. H., Rhee, J. S. and Lee J. S. Modulated expression and enzymatic activity of the monogonont rotifer *Brachionus koreanus* Cu/Zn- and Mn-superoxide dismutase (SOD) in response to environmental biocides. Chemosphere. 2015. 120: 470–478. DOI: 10.1016/j.chemosphere.2014.08.042
- [137] Ferrando, M. D. and Andreu-Moliner, E. Acute lethal toxicity of some pesticides to *Brachionus calyciflorus* and *Brachionus plicatilis*. Bulletin of Environmental Contamination and Toxicology. 1991. 47: 479–484. DOI: 10.1007/BF01702214

- [138] Preston, B. L., Snell, T. W. and Kneisel, R. UV-B exposure increases acute toxicity of pentachlorophenol and mercury to the rotifer *Brachionus calyciflorus*. Environmental Pollution. 1999. 106: 23–31. DOI: 10.1016/S0269-7491(99)00065-2
- [139] Ferrando, M. D., Andreu-Moliner, E. and Fernández-Casalderrey, A. Relative sensitivity of *Daphnia magna* and *Brachionus calyciflorus* to five pesticides. Journal of Environmental Science and Health. Part B. 1992. 27 (5): 511–522. DOI: 10.1080/03601239209372798
- [140] Buikema, A. L., Cairns, J. J. and Sullivan, G. W. Evaluation of *Philodina acuticornis* (Rotifera) as a bioassay organism for heavy metals. Water Resources Bulletin. 1974. 10 (4): 648–661. DOI: 10.1111/j.1752-1688.1974.tb05624.x
- [141] Snell, T. W., Moffat, B. D., Janssen C. and Persoone, G. Acute toxicity tests using rotifers. III. Effects of temperature, strain and exposure time on the sensitivity of *Brachionus plicatilis*. Environmental Toxicology and Water Quality. 1991. 6 (1): 63–75. DOI: 10.1002/tox.2530060106
- [142] Sarma, S. S. S., Martínez-Jerónimo, F., Ramírez-Pérez, T. and Nandini, S. Effect of cadmium and chromium toxicity on the demography and population growth of *Brachionus calyciflorus* and *Brachionus patulus* (Rotifera). Journal of Environmental Science and Health. Part A. 2006. 41: 543–558. DOI: 10.1080/10934520600564311
- [143] Juárez-Franco, M. F., Sarma, S. S. S. and Nandini, S. Effect of cadmium and zinc on the population growth of *Brachionus havanaensis* (Rotifera: Brachionidae). Journal of Environmental Science and Health. Part A. 2007. 42: 1489–1493. DOI: 10.1080/10934520701480938
- [144] Snell, T. W., Moffat, B. D., Janssen, C. and Persoone, G. Acute toxicity tests using rotifers: IV. Effects of cyst age, temperature and salinity on the sensitivity of *Brachionus calyciflorus*. Ecotoxicology and Environmental Safety. 1991. 21(3): 308–317. DOI: 10.1016/0147-6513(91)90070-6
- [145] Robles-Vargas, D. and Snell, T. W. Effects of desiccation on the toxicant sensitivity of rotifers. Hydrobiologia. 2010. 652: 185–193. DOI: 10.1007/s10750-010-0330-8
- [146] Makarewicz, J. C. and Likens, G. E. 1979. Structure and function of the zooplankton community of Mirror Lake, New Hampshire. Ecological Monographs. 49: 109–127.
- [147] Sun, D., Ma, R., Liu, W. and Niu, C. Sexual reproduction and short-term fitness advantage in the rotifer *Brachionus calyciflorus*: implications for the coexistence of sympatric clones. Chinese Journal of Oceanology and Limnology. 2013. 31 (5): 987–993. DOI: 10.1007/s00343-013-2271-3
- [148] Chapman, P. Heavy metal-music, not science. Environmental Science and Technology. 2007. 41 (12): 6C.

- [149] Rubio-Franchini, I. and Rico-Martínez, R. Evidence of lead biomagnification in invertebrate predators from laboratory and field experiments. *Environmental Pollution*. 2011. 159 (7): 1831–1835. DOI: 10.1016/j.envpol.2011.03.021.
- [150] ECOTOX Database Query. Advanced Database Query. 2015. Available from: http://cfpub.epa.gov/ecotox/advanced_query.htm [Accessed: 2015-09-03].
- [151] Wang, H., Tang, X., Sha, J., Chen, H., Sun, T, and Wang, Y. The reproductive toxicity on the rotifer *Brachionus plicatilis* induced by BDE-47 and studies on the effective mechanism based on antioxidant defense system changes. *Chemosphere*. 2015. 135: 129–37. DOI: 10.1016/j.chemosphere.2015.03.090
- [152] Calleja, M., C., Personne, G. and Geladi, P. Comparative acute toxicity of the first 50 multicentre evaluation of in vitro cytotoxicity chemicals to aquatic non-vertebrates. *Archives of Environmental Contamination and Toxicology*. 1994. 26:69–78. DOI: 10.1007/BF00212796
- [153] Calleja, M., C., Personne, G and Geladi, P. Human acute toxicity prediction of the first 50 MEIC chemicals by a battery of ecotoxicological tests and physicochemical properties. *Food and Chemical Toxicology*. 1994. 32 (2): 173–187. DOI: 10.1016/0278-6915(94)90180-5
- [154] Orstan, A. Toxicity of Acrylamide Derivatives to Embryos of the Rotifer *Adineta vaga*. *Bulletin of Environmental Contamination and Toxicology*. 1992. 48 (6): 901–906. DOI: 10.1007/BF00201152
- [155] Huang, L., Xi, Y., Zha, W. and Wen, X. Effect of aldrin on life history characteristics of rotifer *Brachionus calyciflorus* Pallas. *Bulletin of Environmental Contamination and Toxicology*. 2007. 79: 524–528. DOI: org/10.1051/limn/2013038
- [156] Santos-Medrano, G.E., and Rico-Martinez, R. Lethal effects of five metals on the freshwater rotifers *Asplanchna brightwellii* and *Brachionus calyciflorus*. *Hidrobiológica*. 2013. 23 (1): 82–86.
- [157] Calleja, M.C., Persoone, G. and Geladi, P. The predictive potential of a battery of ecotoxicological tests for human acute toxicity as evaluated with the first 50 MEIC chemicals. *ATLA Alternative to Laboratory Animals*. 1993. 21 (3): 330–349.
- [158] Rios-Arana, V., J., Gardea-Torresdey, J., L., Webb, R. and Walsh, E., J. Heat shock protein 60 (HSP60) response of *Platyonus patulus* (Rotifera: Monogononta) to combined exposures of arsenic and heavy metals.. *Hydrobiologia*. 2005. 546: 577–586. DOI: 10.1007/1-4020-4408-9_58
- [159] Rhee, J., Jeong, C., B., Kin, B., M. and Lee, J., L. P-glycoprotein (P-gp) in the monogonont rotifer, *Brachionus koreanus*: molecular characterization and expression in response to pharmaceuticals. *Aquatic Toxicology* 2012. 114–115: 104–118. DOI: 10.1016/j.aquatox.2012.02.009

- [160] Isidori, M., Nardelli, A., Pascarella, L., Rubionp. M. and Parrella. A. Toxic and genotoxic impact of fibrates and their photoproducts on non-target organisms. *Environmental International*. 2007. 33: 635–641. DOI: 10.1016/j.envint.2007.01.006
- [161] Ferrari, B., Paxéus, N., Lo Giudice, N., Pollio, A. and Garric, J. Ecotoxicological impact of pharmaceuticals found in treated wastewaters: study of carbamazepine, clofibric acid, and diclofenac. *Ecotoxicology and Environmental Safety*. 2003. 55: 359–370. DOI: 10.1016/S0147-6513(02)00082-9
- [162] Ferrari, B., Mons, R., Vollat, B., Fraysse, B., Paxéus, N., Lo Giudice, R., Pollio, A., Garric, J. Environmental risk assessment of six human pharmaceuticals: are the current environmental risk assessment procedures sufficient for the protection of the aquatic environment? *Environmental Toxicology and Chemistry*. 2004. 23 (5): 1344–1354. DOI: 10.1897/03-246
- [163] Hurlbert, S.H., M.S. Mulla, and H.R. Willson. Effects of an organophosphorus insecticide on the phytoplankton, zooplankton, and insect populations of fresh-water ponds. *Ecological Monographs*. 1972. 42(3): 269–299. DOI: 10.2307/1942211
- [164] Filenko, O. F. and. Samoylova, T. A. Effect of heavy metals on samples and model population of rotifers *Brachionus plicatilis* Muller at different salinity. *Moscow University Biological Sciences Bulletin*. 2008. 63 (2): 80–83. DOI: 10.3103/S0096392508020077
- [165] Isidori, M., Lavorgna, M., Nardelli, A., Pascarella, L. and Parrella, A. Toxic and genotoxic evaluation of six antibiotics on non-target organisms. *Science of the Total Environment*. 2005. 346: 87–98. DOI: 10.1016/j.scitotenv.2004.11.017
- [166] Klimek, B., Fiałkowska, E., Kocerba-Soroka, W., Fyda, J., Sobczyk, M. and Pajdak-Stós, A. The toxicity of selected trace metals to *Lecane inermis* rotifers isolated from activated sludge. *Bulletin of Environmental Contamination and Toxicology*. 2013. 91:330–333. DOI: 10.1007/s00128-013-1062-z
- [167] Enesco, H.E., Wolanskyj, A. and Sawada, M. Effect of copper on lifespan and lipid peroxidation in rotifers. *Age*. 1989. 12 (1): 19–23. DOI: 10.1007/BF02432514
- [168] Radix, P., Severin, G., Schramm, K-W., and Kettrup, A. Reproduction disturbances of *Brachionus calyciflorus* (rotifer) for the screening of environmental endocrine disrupters. *Chemosphere*. 2002. 47: 1097–1101. DOI: 10.1016/S0045-6535(01)00335-6
- [169] Mejía-Saavedra, J., Sánchez-Armass, S., Santos-Medrano, G., E., Gonzáles-Amaro, R., Razo-Soto, I., Rico-Martínez, R. and Díaz-Barriga, F. Effect of coexposure to ddt and manganese on freshwater invertebrates: pore water from contaminated rivers and laboratory studies. *Environmental Toxicology and Chemistry*. 2005. 24 (8): 2037–2044. DOI: 0.1897/04-438R

- [170] Isidori, M., Nardelli, A., Parrella, A., Pascarella, L. and Previtera, L. A multispecies study to assess the toxic and genotoxic effect of pharmaceuticals: furosemide and its photoproduct. *Chemosphere*. 2006. 63: 785–793.
- [171] Orstan, A. Toxicity of acrylamide derivatives to embryos of the rotifer *Adineta vaga*. *Bulletin of Environmental Contamination and Toxicology*. 1992. 48 (6): 901–906.
- [172] Isidori, M., Lavorgna, M., Nardelli, A., Parella, A., Previtera, L. and Rubino, M. Ecotoxicity of naproxen and its phototransformation products. *Science of the Total Environment*. 2005. 348: 93–101. DOI: 10.1016/j.scitotenv.2004.12.068
- [173] Sha, J., Wang, Y., Chen, H., Wang, M., Wang, H., Li, X., Qi L., and Tang, X. Using population demographic parameters to assess impacts of two polybrominated diphenyl ethers (BDE-47, BDE-209) on the rotifer *Brachionus plicatilis*. *Ecotoxicology and Environmental Safety*. 2015. 119: 106–115. DOI: 10.1016/j.ecoenv.2015.05.009
- [174] DellaGreca, M., Fiorentino, A., Isidori, M., Lavorgna, M., Previtera, L., Rubino, M. and Temussi, F. Toxicity of prednisolone, dexamethasone and their photochemical derivatives on aquatic organisms. *Chemosphere*. 2004. 54: 629–637. DOI:10.1016/j.chemosphere.2003.09.008
- [175] DellaGreca, M., Fiorentino, A., Lesce, M.m R., Isidori, M., Nardelli, A., Previtera, L. and Temussi, F. Identification of phototransformation products of prednisone by sunlight: toxicity of the drug and its derivatives on aquatic organisms. *Environmental Toxicology and Chemistry*. 2003. 22 (3): 534–539. DOI: 10.1002/etc.5620220310
- [176] Liu, Q.T., Williams, T.D., Cumming, R.I., Holm, G., Hetheridge, M.J. and Murray-Smith, R. Comparative aquatic toxicity of propranolol and its photodegraded mixtures; algae and rotifer screening. *Environmental Toxicology and Chemistry*. 2009. 28: 2622–2631. DOI: 10.1897/09-071.1
- [177] Isidori, M., Parrella, A., Pistillo, P. and Temussi, F. Effects of ranitidine and its photo-derivatives in the aquatic environment. *Environment International*. 2009. 35:821–825. DOI: 10.1016/j.envint.2008.12.002
- [178] Persoone, G., Van de Vel, A., Van Steertegem, M. and De Nayer, B. Predictive value of laboratory tests with aquatic invertebrates: influences of experimental conditions. *Aquatic Toxicology*. 1989. 14: 149–166. DOI: 10.1016/0166-445X(89)90025-8
- [179] MicroBioTests Inc. [Internet]. 2015. Available: <http://www.microbiotests.be> [Accessed 2015-09-03]
- [180] Rico-Martínez, R., Pérez-Legaspi, I.A., Arias-Almeida, J.C and Santos-Medrano, G.E. 2013. Rotifers in ecotoxicology. In: *Encyclopedia of Aquatic Ecotoxicology* (Férard J. F., y Blaise C., Eds.). Springer, Berlin, Germany, 973–996. DOI: 10.1007/SpringerReference_380004
- [181] APHA. *Standard Methods for the Examination of Water and Wastewater*. 20th ed. American Public Health Association, Washington, DC. 1998.

- [182] OSPAR Commission. Practical Guidance Document on Whole Effluent Assessment. Publication number 316/2007. OSPAR Commission. 2007.
- [183] Gonçalves, S.F., Calado, R., Gomes, N.C.M., Soares, A.M.V.M. and Loureiro, S. An ecotoxicological analysis of the sediment quality in a European Atlantic harbor emphasizes the current limitations of the Water Framework Directive. *Marine Pollution Bulletin*. 2013. 72 (1): 197–204. DOI: 10.1016/j.marpolbul.2013.04.003
- [184] Isidori, M., Lavorgna, M., Nardelli, A., Parrella, A. Toxicity identification evaluation of leachates from municipal solid waste landfills: a multispecies approach. *Chemosphere*. 2003. 52 (1): 85–94. DOI: 10.1016/S0045-6535(03)00298-4
- [185] Malara A. and Oleszczuk, P. Application of a battery of biotests for the determination of leachate toxicity to bacteria and invertebrates from sewage sludge-amended soil. *Environmental Science and Pollution Research International*. 2013. 20 (5): 3435–3446. DOI: 10.1007/s11356-012-1268-3
- [186] Resource Management Act. Public Act 1991 No 69. New Zealand Legislation. Parliamentary Counsel Office. [Internet]. 1991. Available from: <http://www.legislation.govt.nz/act/public/1991/0069/latest/whole.html#DLM230265> [Accessed: 2015-08-20].
- [187] Ruck, J.G., Martin, M. and Mabon, M. 2000. Evaluation of Toxkits as methods for monitoring water quality in New Zealand. In: *New Microbiotests for Routine Toxicity Screening and Biomonitoring*. Ed. Persoone, G., Janssen, C. and De Coen, W. 103–119 p. DOI: 10.1007/978-1-4615-4289-6_9
- [188] Sarma, S.S.S., Trujillo-Hernández, H.E. and Nandini, S. Population growth of herbivorous rotifers and their predator (*Asplanchna*) on urban wastewaters. *Aquatic Ecology*. 2003. 37: 243–250. DOI: 10.1023/A:1025896703470
- [189] Santos-Medrano, G.E., Ramírez-López, E.M., Hernández-Flores S. Azuara-Medina, P.M. and Rico-Martínez, R. Determination of toxicity levels in the San Pedro River watershed, Aguascalientes, Mexico. *Journal of Environmental Science and Health. Part A. Toxic/Hazardous Substances and Environmental Engineering*. 2007. 42: 1403–1410. DOI: 10.1080/10934520701480557
- [190] Rico-Martínez R., Velázquez-Rojas, C.A., Pérez-Legaspi, I.A. and Santos-Medrano, G.E. 2001. The use of aquatic invertebrate toxicity tests and invertebrate enzyme biomarkers to assess toxicity in the states of Aguascalientes and Jalisco, Mexico. In: *Butterworth, F.M., Gunatilake, A. and Gonsebatt-Bonaparte M.E. Eds. Biomonitoring and Biomarkers as Indicators of Environmental Change, Volume 2*. Plenum Press, Nueva York. pp. 427–438.
- [191] Torres-Guzmán, F., Avelar-González, F.J., y R. Rico-Martínez. An assessment of chemical and physical parameters, several contaminants including metals, and toxicity in the seven major wastewater treatment plants in the state of Aguascalientes, Mexico. *Journal of Environmental Science and Health. Part A. Toxic/Hazardous Sub-*

- stances and Environmental Engineering. 2010. 45 (1): 2–13. DOI: 10.1080/10934520903388517
- [192] Robles-Vargas D., S. M. Montoya-Castillo, F. J. Avelar-González, J. Jauregui-Rincón, F. J. Rodríguez-Valadez, & R. Rico-Martínez. Assessment of the quality and toxicity of the discharges of a wastewater treatment plant and alternatives to improve its operation. Journal of Environmental Health and Science. Part A. Toxic/Hazardous Substances and Environmental Engineering. 2012. 47 (4): 589–597. DOI: 10.1080/10934529.2012.650578
- [193] José de Paggi, S., and Devercelli, M. Land use and basin characteristics determine the composition and abundance of the microzooplankton. Water, Air and Soil Pollution. 2011. 218 (1/4): 93–108. DOI: 10.1007/s11270-010-0626-3
- [194] Koteswari, Y.N. and Ramanibai, R. Evaluation of toxicity of tannery effluent on plankton community structure: a multispecies microcosm study II. Turkish Journal of Biology. 2004. 28: 55–63.
- [195] Park, G., S., Lee, S.H., Park, S.Y., Yoon, S.J., Lee, S.M., Chung, C.S., Hong, G.H. and Kim S. H. Ecotoxicological evaluation of sewage sludge using bioluminescent marine bacteria and rotifer. Ocean Science Journal. 2005. 40 (2): 91–100. DOI: 10.1007/BF03028589
- [196] Gilbert, J.J. and Walsh, E.J. *Brachionus calyciflorus* is a species complex: mating behavior and genetic differentiation among four geographically isolated strains. Hydrobiologia. 2005. 546: 257–265. DOI: 10.1007/s10750-005-4205-3
- [197] Walsh, E.J., Schröder, T., Wallace, R.L. and Rico-Martínez, R. Cryptic speciation in *Lecane bulla* (Monogononta: Rotifera) in Chihuahuan Desert waters. Verhandlungen des Internationalen Verein Limnologie. 2009. 30 (7): 1046–1050.
- [198] Guo, R. and Chen, J. 2015. Assessing the impact of dimethoate on rotifers reproduction through the pre-exposure history. Ecotoxicology and Environmental Safety. 111: 199–205. DOI: 10.1016/j.ecoenv.2014.10.023
- [199] Jones, B.L., VanLoosen, J., Kim, M.H., Miles, S.J., Dunham, C.M., Williams, L.D. and Snell TW. Stress granules form in *Brachionus manjavacas* (Rotifera) in response to a variety of stressors. Comparative Biochemistry and Physiology. Part A. Molecular and Integrative Physiology. 2013. 166 (2): 375–384. DOI: 10.1016/j.cbpa.2013.07.009
- [200] van der Oost, R., Beyer, J. and Vermeulen, N.P. Fish bioaccumulation and biomarkers in environmental risk assessment: a review. Environmental Toxicology and Pharmacology. 2003. 13 (2): 57–149. DOI: 10.1016/S1382-6689(02)00126-6
- [201] Alvarado-Flores, J., Rico-Martínez, R., Ventura-Juárez, J., Silva-Briano, M. and Rubio-Franchini, I. Bioconcentration and localization of lead in the freshwater rotifer *Brachionus calyciflorus* Pallas 1677 (Rotifera: Monogononta). Aquatic Toxicology. 2012. 109: 127–132.

- [202] Smith, H.A. and Snell, T.W. Differential evolution of asexual and sexual females in a benign culture environment. *International Review of Hydrobiology*. 2014. 99 (1–2): 117–124. DOI: 10.1002/iroh.201301711
- [203] Snell, T.W., Johnston, R.K., Gribble, K.E. and Mark Welch, D.B. Rotifers as experimental tools for investigating aging. *Invertebrate Reproduction and Development*. 2015. 59 (suppl 1): 5–10. DOI: 10.1080/07924259.2014.925516
- [204] Snell, T.W. and Johnston, R.K. Glycerol extends lifespan of *Brachionus manjavacas* (Rotifera) and protects against stressors. *Experimental Gerontology*. 0: 47–56. DOI: 10.1016/j.exger.2014.05.005
- [205] Alvarado-Flores, J., Montoya-García, M. Del R., Ventura-Juárez, J. and Rico-Martínez, R. Immunodetection of luteinizing hormone (LH), follicle stimulating hormone (FSH), thyroid stimulating hormone (TSH) and prolactin (PRL) in *Brachionus calyciflorus* (Rotifera: Monogononta). 2009. *Revista de Biología Tropical*. 2009. 57 (4): 1049–1058.
- [206] Hanson, S.J., Stelzer, C.P., Mark Welch, D.B. and Logsdon, J.M. Comparative transcriptome analysis of obligately asexual and cyclically sexual rotifers reveals genes with putative functions in sexual reproduction, dormancy, and asexual egg production. *BMC Genomics*. 2013. 14: 412. DOI: 10.1186/1471-2164-14-412
- [207] Smith, H.A., Burns, A.R., Shearer, T.L. and Snell, T.W. 2012. Three heat shock proteins are essential for rotifer thermotolerance. *Journal of Experimental Marine Biology and Ecology*. 413: 1–6
- [208] Hernández-Ruiz, E., Alvarado-Flores, J., Rubio-Franchini, I., Ventura-Juárez, J. and Rico-Martínez, R. Adverse effects and bioconcentration of chromium in two freshwater rotifer species. [In preparation].
- [209] Rhee, J.S., Jeong, C.B., Kim, B.M. and Lee, J.S. P-glycoprotein (P-gp) in the monogonot rotifer, *Brachionus koreanus*: molecular characterization and expression in response to pharmaceutical. *Aquatic Toxicology*. 2012. 114–115: 10–118. DOI: 10.1016/j.aquatox.2012.02.009
- [210] Liu BR, Liou JS, Chen YJ, Huang YW, Lee HJ. Delivery of nucleic acids, proteins and nanoparticles by arginine-rich cell-penetrating peptides in rotifers. *Marine Biotechnology*. 2013. 15: 584–595. DOI: 10.1007/s10126-013-9509-0
- [211] Jeong, C.B., Lee, J.H., Lee, J.S. and Rhee, J.S. Early expansion and expression of the lipopolysaccharide (LPS)-induced TNF- α factor (LITAF) gene family in the LPS-exposed monogonont rotifer *Brachionus koreanus*. *Comparative Biochemistry and Physiology Part B Biochemistry and Molecular Biology*. 2015. 188: 15–23. DOI: 10.1016/j.cbpb.2015.06.001