We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

The Role of Glutathione Transferases in the Development of Insecticide Resistance

Zazali Alias

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/61972>

Abstract

Glutathione transferases are multifunctional enzymes. Some of the known functions of the enzymes are biotransformation of xenobiotics, countering oxidative stress and participating in cell regulatory functions. As the isoforms present in number of classes the purification of a particular isoform for characterization is a challenging task. In insect, the study of GSTs is focusing on their roles in development of insecticide resistance. There were evident that certain classes of the enzymes are reactive towards conjugating the pesticides. This makes GSTs one of the enzymes of intention in the discipline of pesticide control management.

Keywords: Glutathione transferases, detoxification, insecticide resistance

1. Introduction

This chapter will review literatures concerning glutathione S-transferases from a broad point of view but with an emphasis on their properties, functions, and purification strategies and its challenges. The focus will be on the occurrence of GSTs in insects and the understanding of their role in insecticide resistance. The intention of this review will also be to look into the relationship of particular isoforms of the GSTs to responses to most used insecticides in agriculture.

2. Glutathione-dependent enzymes

Glutathione (GSH, γ -glutamylcysteinylglycine) is a low molecular weight sulfhydryl compound. It is a tripeptide with the sequence glutamic acid, cysteine, and glycine. GSH is a

crystalline solid with a melting point of 192–195°C and molecular weight of 307.33. It dissolves readily in water. There are two peptide bonds, two carboxylic acid groups ($pK_{\text{COOH}} = 3.53$ and 2.12), one amino group ($pK_{\text{NH}_3^+} = 8.66$), and one thiol group ($pH_{\text{SH}} = 9.66$). GSH is found intracellularly in all mammalian tissues and is the major nonprotein thiol compound present in the cell, with concentrations ranging from 0.1 to 10 mM. It is involved in a variety of metabolic processes, for instance, detoxication of xenobiotics, reduction of hydroperoxides, synthesis of leukotrienes and prostaglandins, maintenance of protein and membrane structures, and regulation of numerous enzyme activities. This functional diversity is due to the properties of the thiol group that participates in redox transitions, thiol exchange reactions, thioether formation, and radical scavenging.

A variety of different enzymes utilize glutathione in a variety of biotransformations [1]. Glutathione reductase (GR) catalyses the reduction of GSSG (oxidized glutathione) using NADPH as a reductant. GR is important in maintaining the high cellular reduction potential.

Selenium-dependent glutathione peroxidase (GPOX) is another type of GSH-requiring enzyme that catalyses the reduction of peroxides using GSH as the reducing agent. There are the glutathione S-transferases (GSTs) that are also GSH dependent enzymes with many catalytic activities including the conjugation of GSH to xenobiotics [2,3].

3. Glutathione S-transferases (GSTs, E.C. 2.5. 1.18) with diverse functions

GSTs are found in almost every species, including plants [4] microorganisms [5,6], and animals [7]. GSTs are divided into classes based on their amino acid sequence, immunological, kinetic, and structural properties. In mammals, at least nine classes of GSTs have been identified, namely, Alpha, Mu, Pi, Theta, Omega, Sigma, Zeta, Kappa, and a microsomal class. Human GSTs have been reviewed in reference [8, 9], and [10]. The majority of GSTs are found mainly in the cytosol. Each class consists of one or more protein isoforms. The classes are defined such that the amino acid identity between two isoforms of the same class is more than 50% but more than 30% if they are in different classes [11]. Human cytosolic GSTs are not only in cytoplasm but may also be localized in the mitochondria or the nucleus [12]. The microsomal family of membrane-bound GSTs is also reported and is different from cytosolic GSTs in molecular weight, subunit structure, and immunological reactivity [13-15]. The microsomal GSTs are trimeric, membrane-bound proteins. Mitochondrial GST 13-13 previously purified from rat liver [16] has been later characterized as GSTK1-1 of a Kappa class GST [17]. The Kappa class GSTs are located in mammalian mitochondria and peroxisomes [18,19] and are structurally distinct from the microsomal and cytosolic GSTs [20].

GSTs have a broad and overlapping specificity. Among the reactions catalyzed by GSTs are the substitution of halogens in halogenohydrocarbon, the addition to double bonds, the cleavage of epoxides, and the reduction of organic peroxides. 1-Chloro-2,4-dinitrobenzene (CDNB) is the most common substrate used to assay GSTs in the laboratory as most, but not all, GSTs show catalytic activity with it. Other substrates that have been commonly used to characterize the enzymes are 1,2-dichloro-4-nitrobenzene (DCNB), trans-4-phenyl-3-butene-2-

one (PBO), ethacrynic acid (EA), 1,2-epoxy-3-nitrophenoxypropane (EPNP), *p*-nitrobenzyl chloride (NBC), and sulfobromophthalein (BSP) (Figure 1).

GSTs catalyze the nucleophilic attack by the thiol group of reduced glutathione (GSH) on a wide range of electrophilic substrates. GSTs play important roles in the development of resistance to a variety of exogenous xenobiotics, such as chemotherapeutic drugs [21], chemical carcinogens [22], herbicides [4], and insecticides [7].

3.1. Conjugation of exogenous toxins (Biotransformation)

GSTs play important roles in the protection of macromolecules from attack by reactive electrophiles. The enzymes generally exist in dimeric forms with a subunit molecular weight of approximately 26 kDa. GSTs occur both as homo- and heterodimers. The cytosolic isoenzymes have two active sites per dimer that behave independently of one another [23]. Each active site consists of at least two ligand-binding regions, namely, the GSH binding site (hydrophilic G-site), which is specific for GSH, and the electrophile-binding site (hydrophobic H-site), which is less specific and thus enables GSTs to react with a wide variety of xenobiotics [24]. A review in reference [25] listed xenobiotics that could be conjugated by GSTs. These include halogenonitrobenzenes, organophosphorus compounds, steroids, $\alpha\beta$ -unsaturated carbonyl compounds, aryl halides epoxides, quinones, isothiocyanates, and aryl nitro compounds.

The conjugations catalyzed by the GSTs occur between the nucleophilic GSH and the compounds possessing a sufficiently electrophilic center [25]. The GSTs function by decreasing the pK_a of GSH, thereby allowing its deprotonation and the formation of a more reactive thiolate anion. In most GST classes (Pi, Mu, Alpha, and Sigma), a tyrosine residue in the N-terminal region interacts with GSH to stabilize the thiolate anion. In Theta and Omega classes, this role is carried out by serine and cysteine residues, respectively [9].

This GSH conjugation has been shown to occur in mammals, birds, reptiles, amphibians, fish, insects, and other invertebrates [26], and it is the first step of mercapturic acid formation that is one of the metabolic pathways for detoxication of xenobiotics *in vivo*. The glutathione conjugates, which are water soluble and generally nontoxic, may be converted to the corresponding cysteine conjugate following sequential removal of glutamate and glycine. The cysteine conjugate is either N-acetylated to be excreted as a mercapturic acid or cleaved to a mercaptan by β -lyase. The thiol can be then further metabolized, for example, to be excreted as a glucuronide [26, 27].

Prostaglandin D-synthase, the enzyme involved in production of the D and J series of prostanooids, was characterized as belonging to the Sigma class of GSTs [28]. GSTs also participate in the isomerization of biologically active molecules. A prostaglandin-H E-isomerase of *Ascaridia galli* [29] and prostaglandin-H D-isomerase of rat spleen [30] were identified as Sigma class GSTs. GSTs can also catalyze *cis-trans* isomerizations, for example, the isomerization of maleylacetoacetic acid to fumarylacetoacetic acid. The maleylacetoacetate isomerase activity of hGSTZ1-1 (human GST Zeta 1-1) has been investigated [31] using a spectrophotometric assay with ($\pm$)-2-bromo-3-(4-nitrophenyl)propanoic acid (BNPP) as substrate. Some GSTs

Figure 1. Enzymatic conjugation of common substrates by GST in the presence of GSH. (A) CDNB, (B) DCNB, (C) PBO, (D) EA, (E)EPNP, and (F) BSP.

possess keto-steroid isomerase activity and catalyze the conversion of Δ^5 -3-androstene-3,17-dione to Δ^4 -3-androstene-3,17-dione [21,32].

Human GST A3-3 was shown to efficiently catalyze the double-bond isomerization of Δ^5 -androstene-3,17-dione and Δ^5 -pregnene-3,20-dione [12]. Human GSTs were reported to act as retinoic isomerases that catalyze the steric conversion of 13-*cis*-retinoic acid (13-cRA) to all-*trans*-retinoic acid (t-RA) [33].

Figure 2. Some examples of GSTs with isomerase and synthase functions. (adapted from [9])

3.2. Participation in countering oxidative stress

Reactive oxygen species (ROS) such as superoxide anion, hydrogen peroxide, and the hydroxyl radical are constantly produced during normal aerobic metabolism. The free radicals may attack the polyunsaturated fatty acyl moieties and lead to the peroxidation of lipid bio-membranes. The cleavage of polyunsaturated fatty acids is known to be associated with formation of organic hydroperoxides and reactive aldehydes. These include alkanals, alkenals, and malondialdehyde. These may interfere with several biological processes such as DNA and protein synthesis by inhibition of specific enzyme reactions [34,35].

Peroxidized lipids, produced during oxidative stress, are substrates for GSTs. 4-Hydroxyal-kenals are derived from membrane-bound phospholipid hydroperoxides. They were shown

to be efficiently conjugated to GSH by GST A4-4 [36,37]. A membrane-bound mouse GSTA4-4 [38] and a rat GSTA8-8 [39,40] were also shown efficiently conjugating 4-hydroxyalkenals. It was reported that transfected mGSTA4 protects HL 60 [41] and K562 human erythroleukemia cells [42] against 4-hydroxynonenal-induced apoptosis.

Free radicals can also cause DNA peroxidation. The toxicity of thymine propenal, which is generated by oxidative damage to DNA, was shown to be substantially reduced when HeLa cells received GSTP1-1 and GSH [43]. The rat GST6-6 had previously shown active toward thymine hydroperoxide [44].

Within the cells, peroxides occur either as hydrogen peroxide (H_2O_2) or organic hydroperoxides, such as fatty acid and phospholipid hydroperoxides. GSTs protect tissues from endogenous organic hydroperoxides produced during oxidative stress [35,45]. Some GSTs have shown selenium-independent H_2O_2 and organic hydroperoxidase activity, which are involved in free radical reactions during oxidative stress [46]. A microsomal GST A1-1 of sheep liver exhibited peroxidase activity toward fatty acid hydroperoxides such as linoleic and arachidonic acid hydroperoxides [47]. Human GSTs, such as hGSTA1-1 and hGSTA2-2 [48,49], also exhibited glutathione peroxidase activity toward phospholipid hydroperoxide [50]. Other workers observed elevated GSTs in *Nilaparvata lugens* when treated with pyrethroids, which induces oxidative stress and lipid peroxidation in insects [51].

3.3. Involve in cells regulatory functions

Recent studies of GSTs have demonstrated that a Pi class GST is involved in regulation of c-Jun N-terminal kinase (JNK) signaling in mammals. GSTP interacts with c-Jun N-terminal kinase 1 (JNK1) suppressing the basal kinase activity [52,53]. A model of GST inhibition of JNK signaling was proposed [52]. Under a nonstressed condition, GSTp can exist as free dimeric enzyme or complexed with Jun-JNK thus inhibiting JNK. Upon stress, GSTP forms larger aggregates, which are unable to associate with the Jun-JNK complex, thus enabling the JNK phosphorylation of c-Jun. Phosphorylated Jun can act as a stable and active transcription factor. The accumulation of ROS in response to oxidative stress results in the activation of multiple stress kinase cascades and an elevated level of GSTp expression [54].

Apoptosis signal-regulating kinase 1 (ASK 1) can activate the JNK and the p38 signaling pathways. It plays important role in stress-induced apoptosis. Mouse GSTM1-1 was shown to physically interact with ASK1 and repress ASK1-mediated signaling [55,56].

It has also been reported that human GST class Omega, GSTO1-1, modulates calcium channel (ryanodine receptors, RyRs) protecting mammalian cells from apoptosis induced by calcium (Ca^{2+}) mobilization [57]. It was suggested that RyRs has two binding sites for GSTO1-1. The mammalian protein Bax (21 kDa) is an inducer of apoptosis. A study [58] has reported a plant GST (Theta class) as one of the Bax-inhibiting plant proteins, which prevent apoptosis in plants. GSTP1-1 was proposed [59] interacting with physiological nitric oxide (NO) carriers such as S-nitrosoglutathione (GSNO) and dinitrosyl-diglutathionyl iron complex (DNDGIC). In the absence of GSH, GSNO interacts with and modifies Cys⁴⁷ and Cys¹⁰¹ residues of GSTP1-1 by an S-nitrosylation reaction. Thus, in the cellular depletion of GSH, GSTP1-1 acts directly as an

NO carrier without losing its detoxication activity. The expression of 'tissue' transglutaminase (tTG) is induced in cells programmed to die such as in cells undergoing apoptosis. It was reported that [60] the overexpression of tTG in human neuroblastoma cells increases apoptosis. The study showed that tTG interacts with β -tubulin, histone H2B, and GSTP1-1 to form a protein complex. It was proposed that the interaction with tTG resulted in oligomerization of GSTP1-1. The formation of multimers of GSTP1-1 leads to inactivation of the enzyme toward ROS.

4. GSTs in insects

In insects, GSTs are classified into two groups, class I and class II GSTs [61]. According to a recent proposed classification [62], an insect-specific Delta class GST is classified as a class I GST. This includes those from *Drosophila melanogaster*, DmGSTD1 to DmGST10 [62,63]; *Musca domestica*, MdGSTD1 to MdGSTD5 [49,61] and *Anopheles gambiae*, AgGSTD1 to AgGSTD6 [64,65]. Class II consists primarily of Sigma class GSTs as identified in *D. melanogaster*, DmGSTS1 [66] and *A. gambiae*, AgGSTS1 [67]. A third proposed class of insect's GSTs (class III) [68] that comprises GSTs classified as the Epsilon class in *Drosophila* and the AgGST3-1 and AgGST3-2 of *Anopheles gambiae*.

Studies on insect GSTs were reviewed in detail [7,23,69]. GSTs have been detected in Lepidoptera, Diptera, Coleoptera, Dictyoptera, and Hymenoptera [23]. At present, only few insect GST structures are known. These include that from the Australian sheep blowfly, *L. cuprina* (Theta classes) [66], mosquito, *Anopheles dirus* (isoenzymes 1–3 and 1–4 by [70,71] and fruit fly, *D. melanogaster* (GSTS1) [72,73].

4.1. Characterization of GSTs and its challenges

A problem faced during the extraction of insect GSTs is the possible presence of endogenous inhibitors [74]. Quinones and catecholamines released during homogenization can inhibit the GSTs' activity [75]. The last-named authors suggested the inclusion of GSH in the homogenizing buffer to protect the GSTs from inhibition. Polyphenol pigments have also been shown to inactivate the GSTs. The inclusion of 5–10 mM cysteine in the homogenate prevents the formation of polyphenol. The endogenous inhibitors can also be removed by initial loading of the crude homogenate through an ion exchange or gel filtration column [7]. There are many different strategies employed to purify the GSTs from insects. One of those is by using affinity chromatography with several different affinity matrices. A ligand, sulfobromophthalein-glutathione conjugate (BSP), has been immobilized to an agarose matrix by using either cyanogen bromide [76] or epichlorohydrin activation [77]. The enzymes bind selectively to the resulting matrices when a crude homogenate is applied directly to the columns. The bound enzymes can be eluted by using 0–5 mM of BSP [77] or 1–5 mM GSH solution [76]. The matrix has been used to purify a GST to homogeneity from *Galleria mellonella* [76,78], *Costelytra zealandica* [79], *Musca domestica* [77,80,81], *Drosophila melanogaster* [82] and *Wiseana cervinata* [83]. Another form of affinity matrix that has been used widely is the immobilized GSH-

agarose matrix [84]. The bound enzymes are best eluted from this support with GSH solution. This technique has been used to purify GSTs from a number of insects. These include *Drosophila* [85]. In another instance, a study [86] had used another form of affinity column, namely immobilized S-hexylglutathione [87] to purify GSTs from *Drosophila*.

Investigation of multiple forms of GSTs with different isoelectric points could be performed by using isoelectrofocusing [78,88,89] or chromatofocusing [90]. Purification by affinity chromatography followed by isoelectrofocusing revealed the existence of multiple forms of GSTs [91], in house fly strains Rutgers, Cornell R, and Hirokawa. The presence of multiple isoenzymes of GSTs have also been reported in other species, such as *Aedes aegypti* [92], *G. mellonella* [78], *Plutella xylostella* [92], *C. zealandica* [79] and *Tenebrio molitor* [94].

4.2. GSTs and insecticide resistance

The majority of studies on insect GSTs have focused on their role in conferring insecticide resistance. A review [95] has indicated the importance of genetic and biochemical mechanisms in *Drosophila* in countering toxins and thus developing resistance. There are two types of biochemical mechanism of insecticide resistance outlined [96,97]. The first is by selection of an altered target molecule so that no interaction between the molecules and the toxins occurs, resulting in target site resistance. The target sites of insecticides in insects include the GABA receptor, sodium channel, acetylcholine esterase, acetylcholine receptor, and juvenile hormone receptor. Second, through the process of metabolism, these toxins may be converted to less toxic and more easily excretable derivatives before reaching the target sites. This is a metabolic resistance. Several enzymes play important roles in these types of resistance, such as hydrolases, mixed function oxidases, and glutathione S-transferases and a variety of other conjugating (Phase II) enzymes.

High levels of GSTs have consistently been observed in resistant insect strains and play a major role in insecticide resistance [77,98,99]. Increased activity of GSTs in housefly was found to be associated with resistance to azinphosmethyl [100], parathion [98], pyrethroids [101], tetrachlorovinphos [102] and malathion [103]. *D. melanogaster* develops resistance toward several insecticides. These include malathion [104,105] and 1,1,1-trichloro-2,2-bis-(p-chlorophenyl)ethane (DDT) dehydrochlorinase [106]. Resistance to diazinon, chlorpyrifos, propanthos, and dichlofenthion and their correlation to increased GST activity has also been reported in larval *L. cuprina* [107].

GSTs have been shown to play an important role in insecticide resistance. The catalysis of conjugation of insecticides, such as organophosphorus compounds, chlorinated hydrocarbons, and carbamate insecticides is shown in Figure 3. It was classified three types of reactions catalyzed by GSTs in metabolism of organophosphorus insecticides [108]. The detoxification of organophosphates (OP) occurs by the conjugation of GSH to OP via an O-dealkylation or O-dearylation conjugation, which later forms O-alkyl, O-aryl, and phosphonate conjugates which are all less toxic derivatives.

For the organochlorine insecticides the process involves dehydrochlorination and the GSH conjugation to the parent molecules [97]. Pyrethroids, however, trigger oxidative damage in

cell. Therefore, GSTs role has been detoxifying the lipid peroxidation products resulted by the insecticide [51]. It is well known that some classes of GSTs have shown peroxidases activities. For example, a Delta class GST of *N. lugens* [51] and Epsilon class GST of *A. aegypti* mosquitoes [109,110] and Sigma class GST of *D. melanogaster* [111,112]. There was however report to suggest that protection against pyrethroids can be achieved through a passive sequestration process in which GSTs are capable to bind to various pyrethroids [113,114].

Figure 3. Structures of some (A) halogenated hydrocarbons, (B) organophosphorus, (C) carbamates, and (D) pyrethroids.

5. Summary

GSTs are enzymes of multi-functional roles. Studies in insect have always directed the role of GSTs in conferring resistance toward insecticides resistance. Several classes of GSTs have been

shown to counter the insecticides through direct GSH conjugation process and also their ability to react against lipid peroxidation products. This is due to the fact that some insecticides cause oxidative damage. Direct isolation of responsible GST from insect has been of a challenge due to limited ability of available affinity matrix to capture all classes of GSTs. The characterization of recombinant GSTs could have led to a better understanding of the mechanism of action and thus the regulation of the GSTs upon exposure to insecticides. The availability of fully sequence genomes in model insect such as *D. melanogaster* [115] and recently of *M. domestica* [116] could be of advantage for further studies in glutathione transferase-related insecticides resistance.

Acknowledgements

The author is grateful to Dr A.G. Clark (Victoria University of Wellington) and the Ministry of Higher Education (FRGS: FP052-2014A) for financial assistance.

Author details

Zazali Alias*

Address all correspondence to: alias@um.edu.my

Institute of Biological Sciences, Faculty of Science, University of Malaya, Kuala Lumpur, Malaysia

References

- [1] Fukami J-I. Metabolism of several insecticides by glutathione S-transferase. *Pharmacol. Ther.* 1979;10: 473–514.
- [2] Mannervik B. Glutathione and evolution of enzymes for detoxication of products of oxygen metabolism. *Chemica Scripta.* 1986;26: 281–284
- [3] van Bladere PJ. Glutathione conjugation as a bioactivation reaction. *Chem. Biol. Interact.* 2000;129: 61–76
- [4] Edwards R, Dixon P, Walbot V. Plant glutathione S-transferases: Enzymes with multiple functions in sickness and health. *Trends Plant Sci.* 2000;5: 193–198
- [5] Sheehan D, Casey JP. Microbial glutathione S-transferases. *Comp. Biochem. Physiol.* 1993;104: 1–6
- [6] Vuilleumier S. Bacterial glutathione S-transferases: What are they good for? *J. Bacteriol.* 1997;179: 1431–1441

- [7] Clark AG. The comparative enzymology of the glutathione S-transferases from non-vertebrate organisms. *Comp. Biochem. Physiol.* 1989;92: 419–446
- [8] Ketterer B. A bird's view of the glutathione transferase field. *Chem. Biol. Interact.* 2001;138: 27–42
- [9] Sheeha, D, Meade G, Foley VM, Dowd CA. Structure, function and evolution of glutathione transferases: implications for classification of non-mammalian members of an ancient enzyme superfamily. *Biochem J.* 2001;360: 1–16
- [10] Hayes JD, Flanagan JU, Jowsey IR. Glutathione transferases. *Annu. Rev. Pharmacol. Toxicol.* 2005;45: 51–88
- [11] Mannervik B, Awasthi YC, Board PG, Hayes JD, Di Ilio C, Ketterer B, Litowsky I, Morgenstern R, Muramatsu M, Pearson WR., Picket CB, Sat K, Widersten M, Wolf C.R. Nomenclature for human glutathione transferases. *Biochem. J.* 1992;282: 305–306
- [12] Johansson AS, and Mannervik, B. Human glutathione transferase A3-3, a highly efficient catalyst of double-bond isomerisation in the biosynthetic pathway of steroid hormones. *J. Biol. Chem.* 2001;276: 33061–33065
- [13] Morgenstern R, Meijer J, DePierre JW, Ernster, L. Characterization of rat-liver microsomal glutathione S-transferase activity. *Eur. J. Biochem.* 1980;104: 167–174
- [14] Morgenstern R, Guthenberg C, Depierre JW. Microsomal glutathione S-transferase. Purification, initial characterisation and demonstration that it is not identical to cytosolic glutathione S-transferase A, B, C. *Eur. J. Biochem.* 1982;128: 243–248
- [15] Morgenstern R, DePierre JW. Microsomal glutathione transferase : Purification in un-activated form and further characterization of the activation process, substrate specificity and amino acid composition. *Eur. J. Biochem.* 1983;134: 5911–597
- [16] Harri, JM, Meye DJ, Coles B, Ketterer, B. A novel glutathione transferase (13-13) isolated from the matrix of rat liver mitochondria having structural similarity to class theta enzymes. *Biochem. J.* 1991;278: 137–141
- [17] Pemble SE, Wardle AT, Taylor JB. Glutathione S-transferase class Kappa : characterisation by the cloning of rat mitochondrial GST and identification of a human homologue. *Biochem J.* 1996;319: 749–754
- [18] Morel F, Rauch C, Petit E, Piton, Theret N, Coles B, Guillouzo A. Gene and protein characterization of the human glutathione S-transferase Kappa and evidence for peroxisomal localization. *J. Biol. Chem.* 2004;279: 16246–16253.
- [19] Landler JE, Parsons JF, Rife C, Gilliland GL, Armstrong RN. Parallel evolutionary pathways for glutathione transferases: structure and mechanisms of the mitochondrial class Kappa rGSTK1-1. *Biochemistry.* 2004;43: 352–361
- [20] Robinson A, Huttley GA, Booth HS, Board, PG. Modelling and bioinformatics studies of the human Kappa-class glutathione transferase predict a novel third glutathione transferase.

- thione transferase family with similarity to prokaryotic 2-hydroxychromene-2-carboxylate isomerase. *Biochem J.* 2004;379: 541–552.
- [21] Haye, JD, Pulford DJ. The glutathione S-transferase supergene family: Regulation of GST and the contribution of the isoenzymes to cancer chemoprotection and drug resistance. *Crit. Rev. Biochem. Mol. Biol.* 1995;30: 445–600
- [22] Coles B, Ketterer B. The role of glutathione and glutathione S-transferases in chemical carcinogenesis. *Biochem. Mol. Biol.* 1990;25: 47–70
- [23] Danielson UH, Mannervik B. Kinetic independence of the subunits of cytosolic glutathione S-transferase from the rat. *Biochem. J.* 1985;231: 263–267
- [24] Dir H, Reinemer P, Huber R. X-ray crystal structure of cytosolic glutathione S-transferases: implication for protein architecture, substrate recognition and catalytic function. *Eur. J. Biochem.* 1994;220: 645–661
- [25] Chasseaud LF. The role of glutathione and glutathione S-transferases in the metabolism of chemical carcinogens and other electrophilic agents. *Adv. Cancer Res.* 1979;29: 175–274
- [26] Boyland E, Chasseaud LF. The role of glutathione and glutathione S-transferases in mercapturic acid biosynthesis. *Adv. Enzymol.* 1969;32: 173–219
- [27] Picket CB, Lu AYH, Glutathione S-transferases: gene structure, regulation and biological function. *Annu. Rev. Biochem.* 1989;58: 743–764
- [28] Kanaoka Y, Fujimori K, Kikuno R, Sakaguchi, Y, Urade, Y, and Hyaishi, O. Structure and chromosomal localization of human and mouse genes for hematopoietic prostaglandin D synthase. *Eur. J. Biochem.* 2000;267: 3315–3322
- [29] Meyer DJ, Muimo R, Thomas M, Coates D. and Isaac, R.E. Purification and characterization of prostaglandin-H E-isomerase, a sigma-class glutathione S-transferase, from *Ascaridia galli*. *Biochem. J.* 1996;313: 223–227
- [30] Meyer DJ, Thomas M. Characterization of rat spleen prostaglandin H D-isomerase as a sigma-class GSH transferase. *Biochem. J.* 1995;311: 739–742
- [31] Board PG, Taylor MC, Coggan M, Parke MW, Lantum HB, Anders MW. Clarification of the role of key active site residues of glutathione transferase zeta/maleylacetate isomerase by a new spectrometric technique. *Biochem. J.* 2003;374: 731–737
- [32] Jakoby WB, Habig WH, 1980. Glutathione S-transferases. In *Enzymatic Basis of Detoxification Vol II*. Academic Press Inc.
- [33] Chen H, Juchau MR. Recombinant human glutathione S-transferases catalyze enzymic isomerization of 13-*cis*-retinoic acid to all-*trans*-retinoic acid *in vitro*. *Biochem. J.* 1998;336: 223–226

- [34] Esterbauer H, Cheeseman KH, Dianzani MU, Poli G, and Slater TF. Separation and characterisation of the aldehydic products of lipid peroxidation stimulated by ADP-Fe²⁺ in rat liver microsomes. *Biochem. J.* 1982;208: 129–140
- [35] Ketterer B, Meyer DJ, Taylor JB, Pemble S, Coles B, Fraser G. GSTs and protection against oxidative stress. In : *Proceedings of 3rd International GST conference*, Scotland : Taylor and Francis, London; 1989. p. 97–109.
- [36] Cheng JZ, Yang Y, Singh SP, Singhal SS, Awasthi S, Pan SS, Singh SV, Awasthi YC. Two distinct 4-hydroxynonenal metabolising glutathione S-transferase isozymes are differentially expressed in human tissues. *Biochem. Biophys. Res. Commun.* 2001;282: 1268–1274
- [37] Hubatsch I, Ridderström M, Mannervik B. Human glutathione transferase A4-4: an alpha class enzyme with high catalytic efficiency in the conjugation of 4-hydroxynonenal and other genotoxic products of lipid peroxidation. *Biochem. J.* 1998;330: 175–179
- [38] Singh SP, Janecki AJ, Srivastava SK, Awasthi S, Awasthi YC, Xia SJ, Zimniak P. Membrane association of glutathione S-transferase mGSTA4-4, and enzyme that metabolizes lipid peroxidation products. *J. Biol. Chem.* 2002;277: 4232–4239
- [39] Jensson H, Guthenber, C, Ålin P, Mannervik B. Rat glutathione transferase 8-8, an enzyme efficiently detoxifying 4-hydroxyalk-2-enals. *FEBS Lett.* 1986;203: 207–209
- [40] Stenberg G, Ridderström M, Engström Å, Pemble SE, Mannervik B. Cloning and heterologous expression of cDNA encoding class alpha rat glutathione transferase 8-8, an enzyme with high catalytic activity towards genotoxic α,β -unsaturated carbonyl compounds. *Biochem. J.* 1992;284: 313–319
- [41] Cheng JZ, Singhal SS, Sharma A, Saini M, Yang Y, Awasthi S, Zimniak P, Awasthi YC. Transfection of mGSTA4 in HL-60 cells protects against 4-hydroxynonenal-induced apoptosis by inhibiting JNK-mediated signalling. *Arch. Biochem. Biophys.* 2001;392:197–207
- [42] Cheng JZ, Singhal SS, Saini M, Singhal J, Piper JT, van Kuijk FJ, , Zimniak GM, Awasthi, YC, Awasthi S. Effects of mGST A4 transfection on 4-hydroxynonenal-mediated apoptosis and differentiation of K562 human erythroleukimia cells. *Arch. Biochem. Biophys.* 1999;372: 29–36
- [43] Berhane K, Widersten M, Engstrom A, Kozarich JW, Mannervik B. Detoxication of base propenals and other $\alpha\beta$ -unsaturated aldehyde products of radical reaction and lipid peroxidation by human GSTs. *Proc. Natl. Acad. Sci. U. S. A.* 1994;91: 1480–1484
- [44] Tan KH, Meyer DJ, Ketterer B. Thymine hydroperoxide, a substrate for rat Se-dependent glutathione peroxidase and glutathione transferase isoenzymes. *FEBS Lett.* 1986;207: 231–233

- [45] Storey KB.. Oxidative Stress : Animal adaptations in nature. *Braz. J. Med. Biol. Res.* 1996;29: 1715-1733
- [46] Wilce MCJ, Parker MW. Structure and function of glutathione S-transferases. *Biochem. Biophys. Acta.* 1994;205:1-18
- [47] Prabhu KS, Reddy PV, Gumprich E, Hildenbrandt GR, Scholz RW, Sordillo LM, Reddy CC. Microsomal glutathione S-transferase A1-1 with glutathione peroxidase activity from sheep liver: molecular cloning, expression and characterisation. *Biochem. J.* 2001;360: 345-354
- [48] Yang Y, Cheng JZ, Singhal SS, Sain, M, Pandya U, Awasthi S, Awasthi YC. Role of glutathione S-transferases in protection against lipid peroxidation. *J. Biol. Chem.* 2001;276: 19220-19230
- [49] Zhou ZH, Syvanen M. A complex glutathione transferase gene family in the housefly *Musca domestica*. *Mol. Gen. Genet.* 1997;256: 187-194
- [50] Hurst R, Bao Y, Jemth P, Mannervik B, Williamson G. Phospholipid hydroperoxide glutathione peroxidase activity of human glutathione transferase. *Biochem. J.* 1998;332: 97-100
- [51] Vontas JG, Small GJ, Hemingway J. Glutathione S-transferases as antioxidant defence agents confer pyrethroid resistance in *Nilaparvata lugens*. *Biocem. J.* 2001;357: 65-72
- [52] Adler V, Yin Z, Fuchs SY, Benezra M, Rosario L, Tew KD, Pincus MR, Sardana M, Henderson CJ, Wolf CR, Davis RJ, Ronai Z. Regulation of JNK signalling by GSTp. *EMBO J.* 1999;18: 1321-1334
- [53] Wang T, Arifoglu P, Ronai Z, Tew KD. Glutathione S-transferase P1-1 (GSTP1-1) inhibits c-Jun N-terminal Kinase (JNK1) signaling through interaction with C-terminus. *J. Biol. Chem.* 2001;276: 20999-21003
- [54] Ainbinder E, Bergelson S, Pinkus R, Daniel V. Regulatory mechanisms involved in activator-protein-1 (AP-1)-mediated activation of glutathione S-transferase gene expression by chemical agents. *Eur. J. Biochem.* 1997;243: 49-57
- [55] Cho SG, Lee YH, Park HS, Ryoo K, Kang KW, Park J, Eom SJ, Kim MJ, Chang TS, Choi, SY, Shim J, Kim Y, Dong MS, Lee MJ, Kim SG, Ichijo H, Choi EJ. Glutathione S-transferase Mu modulates the stress-activated signals by suppressing apoptosis signal-regulating kinase 1. *J. Biol. Chem.* 2001;276: 12749-12755
- [56] Gilot D, Loyer P, Corlu A, Glaise D, Lagadic-Gossmann D, Atfi A, Morel F, Ichijo H, Guguen-Guillouzo C. Liver protection from apoptosis requires both blockage of initiator caspase activities and inhibition of ASK1/JNK pathway via glutathione S-transferase regulation. *J. Biol. Chem.* 2002;277: 49220-49229

- [57] Dulhunty A, Gage P, Curtis S, Chelvanayagam G, Board P. The glutathione transferase structural family includes a nuclear chloride channel and a ryanodine receptor calcium release channel modulator. *J. Biol. Chem.* 2001;276: 3319–3323
- [58] Kampranis SC, Damianova R, Atallah M, Toby G, Kondi G, Tsihchlis PN, Makris AM. A novel plant glutathione S-transferase/peroxidase suppresses Bax lethality in yeast. *J. Biol. Chem.* 2000;275: 29207–29297
- [59] Lo Bello M, Nuccetelli M, Caccuri AM, Stella L, Parker MW, Rossjohn J, McKinstry WJ, Mozzi AF, Federici G, Polizio F, Pederson JZ, Ricci G. Human glutathione transferase P1-1 and nitric oxide carriers. *J. Biol. Chem.* 2001;276: 42138–42145
- [60] Piredda L, Farrace MG, Belloo ML, Malorni W, Melino G, Petruzzeli R, Piacetini M. Identification of ‘tissue’ transglutaminase binding proteins in neural cells committed to apoptosis. *FASEB J.* 1999;13: 335–364
- [61] Fournier D, Bride JM, Poirie M, Berge JB, Flapp Jr. FW. Insect glutathione S-transferases: biochemical characteristics of the major forms from houseflies susceptible and resistant to insecticides. *J. Biol. Chem.* 1991;267: 1840–1845
- [62] Chelvanayagam G, Parker MW, Board PG. Fly fishing for GSTs: a unified nomenclature for mammalian and insect glutathione transferases. *Chem. Biol. Interact.* 2001;133: 256–260
- [63] Toung YPS, Hsieh TS, Tu CPD. The glutathione S-transferase D genes: a divergently organised, intronless gene family in *Drosophila melanogaster*. *J. Biol. Chem.* 1993;268: 9737–9746
- [64] Ranson HR, Cornel AJ, Fournier D, Vaughan A, Collins FH, Hemingway J. Cloning and localization of a glutathione S-transferase class I gene *Anopheles gambiae*. *J. Biol. Chem.* 1997;272: 5464–5468
- [65] Ranson H, Collins J, Hemingway J. The role of alternative mRNA splicing in generating heterogeneity within the *Anopheles gambiae* class I glutathione S-transferase family. *Proc. Natl. Acad. Sci. U. S. A.* 1998;95: 14284–14289
- [66] Beall C, Fryberg C, Song S, Fryberg E. Isolation of *Drosophila* gene encoding glutathione S-transferase. *Biochem. Genet.* 1992;30: 515–527
- [67] Reiss RA, James AA. A glutathione S-transferase gene of the vector mosquito, *Anopheles gambiae*. *Insect Mol. Biol.* 1993;2: 25–32
- [68] Ranson H, Rossiter L, Ortellì F, Jensen B, Wang X, Rothe CW, Collins FH, Hemingway J. Identification of a novel class of insect glutathione S-transferases involved in resistance to DDT in the malaria vector *Anopheles gambiae* *Biochem J.* 2001;359: 295–304
- [69] Enayati AA, Ranson H, Hemingway J. Insect glutathione transferases and insecticides resistance. *Insect Mol. Biol.* 2005;14: 3–8

- [70] Oakley AJ, Harnnoi T, Udomsinprasert R, Jirajaroenrat K, Kettermann AJ, Wilce MC. The crystal structures of glutathione S-transferases isoenzymes 1–3 and 1–4 from *Anopheles dirus* species B. *Protein Sci.* 2001;10: 2176–2185
- [71] Udomsinprasert R, Pongjaroenkit S, Wongsantichon J, Oakley AJ, Prapantadhara LA, Wilce MCJ, Kettermann AJ. Identification, characterization and structure of a new Delta class glutathione transferase isoenzyme. *Biochem. J.* 2005; 388: 763–771
- [72] Agianian B, Clayton JD, Leonard K, Tucker P, Bullard B, Gros P. Crystallization and preliminary X-ray analysis of *Drosophila* glutathione S-transferase-2. *Acta Crystallogr. D. Biol. Crystallogr.* 2000;57: 725–727
- [73] Agianian B, Tucker PA, Schouten A, Leonard K, Bullard B, Gros P. Structure of a *Drosophila* sigma class glutathione S-transferase reveals a novel active site topography suited for lipid peroxidation products. *J. Mol. Biol.* 2003;326: 151–165
- [74] Motoyama N, Kulkarni AP, Hodgson E, Dauterman WC. Endogenous inhibitors of glutathione S-transferase in houseflies. *Pestic. Biochem. Physiol.* 1978;9: 255–262
- [75] Kulkarni AP, Motoyama N, Dauterman WC, Hodgson E. Inhibition of housefly glutathione S-transferase by catecholamines and quinones. *Bull. Environ. Contam. Toxicol.* 1978;20: 221–232
- [76] Clark AG, Letoa M, Ting WS. The purification by affinity chromatography of a glutathione S-transferase from larvae of *Galleria mellonella*. *Life Sci.* 1977;20: 141–148
- [77] Clark, A.G. and Dauterman, W.C. The characterisation by affinity chromatography of glutathione S-transferases from different strains of housefly. *Pestic. Biochem. Physiol.* 1982;117: 307–314
- [78] Baker WL, Clark AG, Faulds G, Nielson JS. Multiple glutathione S-transferases in *Galleria mellonella* ; their detection with fluorogenic substrates. *Insect Biochem. Molec. Biol.* 1994;24: 301–307
- [79] Clark AG, Dick GL, Martindal SM, Smith JN. Glutathione S-transferases from New Zealand grass grub, *Costelytera zealandica*. *Insect Biochem.* 1985;15: 35–44
- [80] Clark AG, Shamaan NA. Evidence that DDT-dehydrochlorinase from the housefly is a glutathione S-transferase. *Pestic. Biochem. Physiol.* 1984;22: 249–261
- [81] Clark AG, Marshall, SN, Qureshi AR. Synthesis and use of an isoform-specific affinity matrix in the purification of glutathione S-transferases from the house fly, *Musca domestica* (L.). *Protein Expr. Purif.* 1990;1: 121–126.
- [82] Alias Z, Clark AG. Studies on the glutathione S-transferase proteome of adult *Drosophila melanogaster*: responsiveness to chemical challenge. *Proteomics.* 2007;7: 3618–3628.
- [83] Clark AG, Drake B. Purification and properties of glutathione S-transferases from larvae of *Wiseana cervinata*. *Biochem. J.* 1984;217: 41–50

- [84] Simon PC, van der Jagt DL. Purification of glutathione S-transferases from human liver by glutathione-affinity chromatography. *Anal. Biochem.* 1977;82: 334–341
- [85] Cochrane BJ, Morrissey JJ, LeBlanc GA. The genetics of xenobiotic metabolism in *Drosophila*. IV. Purification and characterisation of the major glutathione S-transferase. *Insect Biochem.* 1987;17: 731–738
- [86] Toung YP, Hsieh TS, Tu CPD. *Drosophila* glutathione S-transferase 1-1 shares a region of sequence homology with the maize glutathione S-transferases III. *Proc. Natl. Acad. Sci. U. S. A.* 1990;87: 31–35
- [87] Mannervik B, Guthenberg C. Glutathione transferase (human placenta). *Methods Enzymol.* 1981;77: 231–235
- [88] Clark AG, Smith JN, Speir TW. Cross specificity in some vertebrate and insect glutathione S-transferases with methylparathion (dimethyl p-nitrophenyl phosphorothionate), 1-chloro-2,4-dinitrobenzene and *N*-crotonyl-*N*-acetyl cysteamine as substrates. *Biochem. J.* 1973;135: 285–392
- [89] Yawetz A, Koren B. Purification and properties of the Mediterranean fruit fly *Ceratitis capitata* W. glutathione S-transferase. *Insect Biochem.* 1984;14: 663–670
- [90] Keeran WS, Lee RF. The purification and characterisation of glutathione S-transferase from the hepatopancreas of the Blue Crab, *Callinectes sapidus*. *Arch. Biochem. Biophys.* 1975;25: 8670–8673
- [91] Clark, A.G. and Dauterman, W.C. The characterisation by affinity chromatography of glutathione S-transferases from different strains of housefly. *Pestic. Biochem. Physiol.* 1982;117: 307–314.
- [92] Grant DF, Dietze EC, Hammock BD. Glutathione S-transferases isoenzymes in: purification, characterisation and isoenzyme-specific regulation. *Insect Biochem.* 1991;21: 421–433
- [93] Chiang FM, Sun CN. Glutathione transferase isoenzymes of diamondback moth larvae and their role in the degradation of some organophosphorus insecticides. *Pestic. Biochem. Physiol.* 1993;45: 7–14
- [94] Kostaropoulos I, Mantzari AE, Papadopoulos AI. Alterations of some glutathione S-transferase characteristics during the development of *Tenebrio molitor* (Insecta: Coleoptera). *Insect Biochem. Mol. Biol.* 1996;26: 963–969
- [95] Wilson TG. Resistance of *Drosophila* to toxins. *Annu. Rev. Entomol.* 2001;46: 545–571.
- [96] Morton RA. Evolution of *Drosophila* insecticide resistance. *Genome.* 1992;36: 1–7
- [97] Oppenoorth FJ. Biochemistry of insecticide resistance. *Pestic. Biochem. Physiol.* 14;22: 187–193

- [98] Oppenoorth FJ, Rupes V, El-Bashir S, Houx NWH, Voerman S. Glutathione dependent degradation of parathion and its significance for resistance in housefly. *Pestic. Biochem. Physiol.* 1972;2: 262–269
- [99] Motoyama N, Dauterman WC. Interstrain comparison of glutathione-dependent reactions in susceptible and resistant houseflies. *Pestic. Biochem. Physiol.* 1975;5: 489–495
- [100] Motoyama N., Dauterman WC. *In vitro* metabolism of azinphosmethyl in susceptible and resistant house flies. *Pestic. Biochem. Physiol.* 1972;2: 113–122
- [101] Papadoplus AI., Boukouvala E, Kakaliouras G, Kostaropoulos J, Papadopoulou-Mourkidou. Effect of organophosphate and pyrethroid insecticides on the expression of GST from *Tenebrio molitor* pupae. *Pestic. Biochem. Physiol.* 2000;68: 26–33
- [102] Oppenoorth FJ, van der Pas LJT, Houx NWH. Glutathione S-transferase and hydrolytic activity in a tetrachlorvinphos-resistant strain of housefly and their influence on resistance. *Pestic. Biochem. Physiol.* 1979;11: 176–188
- [103] Niwa Y, Miyata T, Saito T. In vitro metabolism of malathion by malathion resistant and susceptible strains of houseflies, *Musca domestica*. *J. Pestic. Sci.* 1977;2: 151–157
- [104] Houpt DR, Pursey JC, Morton RA. Genes controlling malathion resistance in laboratory-selected population of *Drosophila melanogaster*. *Genome.* 1988;30: 844–853
- [105] Cochrane JB, LeBlanc GA. Genetic of metabolism in *Drosophila*. I. Genetic and environmental factors affecting glutathione S-transferase in larvae. *Biochem. Pharmacol.* 1986;35: 1679–1684
- [106] Shepanski MC, Glover TJ, Kuhr RJ. Resistance of *Drosophila* to DDT. *J. Econ. Entomol.* 1977;70: 539–543
- [107] Wilson JA, Clark AG. The role of E3 esterase, glutathione S-transferases and other nonoxidative mechanisms in resistance to diazinon and other organophosphate insecticides in *Lucilia cuprina*. *Pestic. Biochem. Physiol.* 1996;54: 85–95
- [108] Motoyama N, Dauterman WC. Glutathione S-transferases: Their role in the metabolism of organophosphorus insecticides. *Rev. Biochem. Toxicol.* 1980;2: 49–69
- [109] Lumjuan N, McCarroll L, Prapanthadara L, Hemingway J, Ranson H. Elevated activity of an Epsilon class glutathione transferase confers DDT resistance in the dengue vector, *Aedes aegypti*. *Insect Biochem. Mol. Biol.* 2005;35: 861–871
- [110] Ding Y, Hawkes N, Meredith J, Eggleston P, Hemingway, J, Ranson, H. Characterization of the promoters of Epsilon glutathione transferases in the mosquito *Anopheles gambiae* and their response to oxidative stress. *Biochem J.* 2005;387: 879–888.
- [111] Sawicki R , Singh S P, Mondal, A K, Benes H, Zimniak P. Cloning, expression and biochemical characterization of one Epsilon-class (GST-3) and ten Delta-class (GST-1)

glutathione S-transferases from *Drosophila melanogaster*, and identification of additional nine members of the Epsilon class. *Biochem. J.* 2003;370: 661–669

- [112] Singh SP, Coronella JA, Benes H, Cochrane BJ, Zimniak, P. Catalytic function of *Drosophila melanogaster* glutathione S-transferase DmGSTS1-1 (GST-2) in conjugation of lipid peroxidation end products. *Eur. J. Biochem.* 2001;268: 2912–2923
- [113] Udomsinprasert R, Kettermann AJ. Expression and characterisation of a novel class of glutathione S-transferase from *Anopheles dirus*. *Insect Biochem. Mol. Biol.* 2002;32: 425–433
- [114] Jirajaroenrat K, Pongjaroenkit S, Krittanai C, Prapanthadara L, Kettermann AJ. Heterologous expression and characterization of alternatively spliced glutathione-S-transferases from a single *Anopheles* gene. *Insect Biochem. Mol. Biol.* 2001;31: 867–875
- [115] Wilson TG. *Drosophila melanogaster* (Diptera: Drosophilidae): a model insect for insecticide resistance studies. *J. Econ. Entomol.* 1988;81: 22–27
- [116] Scott JG, Warren WC, Beukeboom LW, Bopp D, Clark AG, Giers SD, Hediger M, Jones AK, Kasai S, Leichter CA, Li M, Meisel RP, Minx P, Murphy TD, Nelson, DR, Reid WR, Rinkevich FD, Robertson, HM, Sackton TB, Sattelle DB, Thibaud-Nissen, F, Tomlinson C, van de Zande L, Walden KKO, Wilson RK, Liu N. Genome of the house fly, *Musca domestica* L., a global vector of diseases with adaptations to a septic environment. *Genome Biol.* 2014;15: 466–482