We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Modelling in Metal Risk Assessment

T.T. Yen Le

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/62911>

Abstract

Unique properties of metals that are largely different from the characteristics of organic substances should be considered in risk assessment. The bioavailability and toxicity of metals depend on their chemical speciation, that is, physical-chemical forms, in the environment, which is largely influenced by the environmental chemistry. Since metals in the environment are not always available while organisms have developed different processes to actively regulate the body burden, assessment of metal bioaccumulation might provide a better understanding of potential risks. Metal bioaccumulation is a prerequisite for metal toxicity, but is not the only determinant of metal toxicity. In addition to metal accumulation, metal toxicity is influenced by the subcellular partitioning of metals, which is controlled by the capacity of organisms to sequester and to detoxify metals. Different modelling approaches have been developed to address some of these factors. Both empirical and mechanistic equilibrium models have been developed and applied for characterising metal speciation in the environment. Metal bioaccumulation has been predicted by biodynamic models. The ability of organisms to detoxify metals has been taken into account in assessment based on the induction of metallothionein (MT) or subcellular partitioning. In addition, the interactions between organisms and metal ions have been taken into consideration in assessment of metal toxicity based on the accumulation of metal ions at biological surfaces.

Keywords: Metal, modelling, risk assessment, speciation, bioaccumulation, subcellular partitioning, toxicity

1. Introduction

Environmental pollution with metals is a serious problem in many areas in the world, and assessment of metal bioaccumulation and toxicity is of high concern. In such assessment, unique properties of metals that are largely different from the characteristics of organic substances

should be considered. The bioavailability and toxicity of metals are controlled by their chemical speciation, which is heavily influenced by environmental chemistry [1–5]. Metals are an intrinsic component of the environment, naturally occurring with varying background concentrations [6, 7]. Several metals are essential elements, that is, the elements that are present in living organisms and able to interact with the living system, and an insufficient amount of these elements leads to preventable or reversible effects on certain biological function, which do not occur at a physiological amount of the elements [8]. There is a limit to the regulation by the homeostasis mechanisms. Biological functions of organisms are affected by either the deficiency (below the limit) or toxicity (above the limit) of the essential metals. Because of these unique properties, organisms have developed various regulatory mechanisms for uptake, metabolism, distribution, storage, and elimination in response to metal exposure and to maintain metal homeostasis. The responses of organisms depend on a number of factors, from environmental conditions, bioaccumulation, to internal detoxification [9–11]. Metal bioaccumulation, a prerequisite for metal toxicity, is determined by the chemical speciation of metals in the environment, which depends on environmental conditions. In addition to metal accumulation, metal toxicity is influenced by metal subcellular partitioning. Different modelling approaches have been developed to address some of these factors as described in more detail in the following sections.

2. Metal availability in the environment

The bioavailability and toxicity of metals depend to a large extent on the chemical speciation of metals [12–15]. A widely recognised definition of the chemical speciation is that the chemical speciation of an element is the identification and quantification of the different, defined species, forms, or phases in which the element is present [16, 17]. Each form of the element is defined by its isotopic composition, electronic or oxidation state, or molecular structure [18, 19]. The mobility of metals in soil depends on the partition of metals between the solid and solution phases [20], which is determined by properties of the metals as well as composition of both solid and solution phases [21]. The most important soil properties include the total metal content, pH, cation exchange capacity (CEC), contents of soil organic matter (SOM), clay, oxides, and exchangeable Ca and Mg [22–26]. Besides the chemical extraction for direct determination, the solid-solution partitioning of metals in soils can be characterised by equilibrium models, which allow for predicting the speciation of a metal based on soil properties. Generally, two main approaches have been applied for modelling the solid-solution partitioning of metals in soils. They are based on: (1) empirical relations or transfer functions between the concentration of metals in soil solution or in a specific form and soil and solution properties [22, 24, 26, 27] or (2) complex process-based (mechanistic) multi-surface complexation models incorporating a detailed simulation of soil surface complexation and solution chemistry [23, 26, 28–31].

2.1. Transfer functions

The partition of metals between the solid and solution phases in soil has been expressed by adsorption isotherms or a distribution coefficient (K_d ; L/kg):

$$K_d = \frac{C}{C_{ss}} \quad (1)$$

where K_d (L/kg) is the distribution coefficient; C (mg/kg or mol/kg) is the total or reactive concentration of the metal in soil; C_{ss} (mg/L or mol/L) is the solution concentration.

In aerobic conditions, metals occluded in mineral particles and organic matter that are not readily available are included in the total pool, whilst the reactive fraction (extracted with 0.43M HNO_3) represents the metals available for exchange with soil solution with a particular time span [32]. Therefore, the reactive concentration has been used to substitute the total concentration in the above equation [32, 33]. The partitioning of metals in soil is usually expressed by sorption isotherms as described by a Freundlich equation:

$$C = k \cdot C_{ss}^n \quad (2)$$

where k and n are the Freundlich parameters. The exponent n reflects the variation in the binding strength with varying total pools. Equation 2 can be rewritten and extended to take into account the influence of soil properties [22, 32, 34–36]:

$$\log C = \log k + n \cdot \log C_{ss} + \sum a_i \cdot \log(X_i) \quad (3)$$

where a_i is the coefficient that reflects the influence of soil parameter X_i on the partitioning of metals between the solid and solution phases. The distribution coefficient has also been related to soil properties:

$$\log K_d = a_0 + a \cdot \log C_{ss} + \sum a_i \cdot \log(X_i) \quad (4)$$

Empirical relationships have been established between the distribution coefficient or the dissolved concentration, on the one hand, and the total/reactive concentration and soil properties, on the other. Different equations have been derived, depending on the empirical data that they have been based on, whilst a full description of soil properties is usually not available. Among soil properties, pH is the most commonly integrated parameter in transfer functions, followed by soil organic carbon (SOC), while other soil properties are rarely included in. This reflects the importance of pH and organic matter in determining metal speciation as well as the partitioning of metals between the solid and solution phases.

2.2. Mechanistic multi-surface complexation models

Substantial progress has been obtained in simulating the sorption of metal ions to soil and dissolved organic matter (DOM) with the development of different models, for example, MINEQL [37], GEOCHEM [38], NICADonnan [39], and WHAM V and VI [40, 41]. Metal speciation in soil has been characterised by various models calculating metal movement and available fractions and simulating impacts of physicochemical properties on metal dynamics in soil. The movement of different metal species in an environment is characterised by fate and transport models, while the partitioning of metals into dissolved, colloidal, and particulate phases is predicted by speciation/complexation models. Assemblage models have been formed by incorporating various surface complexation models. These process-based models provide a detailed description of metal complexation on different sorption surfaces [39, 42]. Surface complexation models based on thermodynamic processes have been developed for describing reactions of metals on the surface of soil reactive phases, such as SOM [39, 41], oxides [42, 43], and clay minerals [44]. Multi-surface complexation models have been formed from individual surface complexation models, assuming that the adsorption properties of soil can be characterised by the sum of the individual adsorption properties of individual components. Metal concentrations in soil solution have been predicted by using mechanistic geochemical models accounting for interactions of metals with various reactive surfaces [23, 24, 26, 29, 30].

Originally developed to characterise metal speciation in oxic waters [40, 41], WHAM has been applied to simulate the solid-solution partitioning [23, 45]. This model consists of several sub-models: Humic Ion-Binding Model V and models of inorganic solution chemistry, adsorption-desorption reactions of fulvic acids, precipitation of oxides, and cation exchange reactions on clays. In the Humic Ion-Binding Model V, humic substances are represented by molecules containing proton-dissociating groups that can bind to metal ions [40]. The binding of metal ions to humic substances is described in terms of complexation at discrete sites while taking into account both electrostatic interactions and competition among ions. In the application to soil systems, a fixed charge mineral cation exchanger was included to describe the presence of clays [46].

Weng et al. [29, 47] have proposed another multi-surface or mechanistic model for characterizing metal speciation in which soil is considered a set of independent sorption surfaces, that is, organic matter, clay silicates, and iron (hydr)oxides. Metal binding to these surfaces is simulated using advanced adsorption, cation exchange models, and default binding parameters. Humic acid is considered an analogue for SOM. The site density of SOM is assumed to be equal to 31% of the density of humic acid [47], while metal binding to SOM is described by using the non-ideal competitive consistent Adsorption (NICA)-Donnan model [39]. Clays are represented by illite because of their importance in Dutch soils [48]. The average charge of illite is assumed to be pH dependent (0.25 mol/kg; [49]). An electrostatic Donnan model is used to describe metal binding to clay. For amorphous iron (hydr)oxides, their amount is predicted as the oxalate-extractable Fe, and their surface area is considered similar to that of hydrous ferric oxides (HFO; 600 m²/g) [42]. For crystalline iron (hydr)oxides, their surface area is assumed to be the same as that of goethite (100 m²/g), and their amount is calculated as the difference between total (aqua regia) and oxalate-extractable Fe. Metal binding to the oxides is simulated

using the two-site surface complexation diffuse double layer model [42]. The complexation of metals with carbonate and dissolved organic matter (DOM) is also taken into account in the model. DOM is assumed to consist of 30% humic acid and 30% fulvic acid [29].

The ORCHESTRA (Objective Representing CHEmical Speciation and TRANsport) modelling framework [50, 51] is a combination of different sub-models. Thermodynamic data from the MINTEQA2 [52] were used to calculate solution speciation. The sorption of ions to SOM and DOM was simulated with the NICA-Donnan model [39, 53, 54] using generic binding parameters and constants derived by Milne et al. [55, 56]. When data on DOM were not available, DOM was assumed to consist of 50% reactive humic substances. In addition, SOM and DOM were represented by humic acid assuming that humic acid consists of 50% carbon [56]. The sorption of ions on the surface of (hydr)oxides was described by the generalized two layer model (GTLM) of Dzombak and Morel [42]. Site densities of the amorphous iron and aluminium (hydr)oxides were calculated from the assumed specific surface area of 600 m²/g for hydrous ferric oxide [42]. Crystalline iron (hydr)oxides were represented by the hydrous ferric oxide, assuming the same reactivity. Moreover, a specific surface area of 100 m²/g was assumed for these (hydr)oxides [43]. The nonspecific sorption of ions to permanently charged clay surfaces was modelled with a Donnan model. The illitic clay mineral with an average charge density of 0.25 eq/kg and a fixed Donnan model of 1 L/kg were used to represent the clay mineral [49].

3. Metal bioaccumulation

Since metals in the environment are not always available, assessing metal accumulation in sentinel species is one of the most effective methods for evaluating effects of metals on biota. The tendency of chemicals to be accumulated in organisms is usually expressed by the bioaccumulation factor (BAF) and the bioconcentration factor (BCF). These factors represent the ratio of the residue in the organisms versus the concentration in water at equilibrium, including or excluding the uptake from dietary sources, respectively. For metals, BCF and BAF vary widely and are inversely related to the external water concentration [57–59]. These characteristics of the accumulation factors for metals are attributed to the complex mechanisms of metal uptake and accumulation [60]. In a number of studies, uptake via saturable kinetics-exhibiting mechanisms has been shown to be more common and toxicologically relevant than passive diffusion [61–65]. The concentration of metals accumulated in organisms is further influenced by different physiological and anatomical mechanisms in response to metal exposure [6]. For essential metals, organisms are able to actively regulate metal bioaccumulation and maintain homeostasis over a range of exposure via exclusion or increased elimination [11, 57, 65]. Organisms have also developed different mechanisms to sequester, detoxify, or store excess metals [11, 62, 66]. The dependence of BCF and BAF on exposure conditions as well their inverse relationship with the exposure concentration invalidates the use of these factors in risk assessment [6, 57, 59].

Biodynamic models have been recommended as an alternative to single and generic values of BCF or BAF in the assessment of metal bioaccumulation [67]. Such models allow for integrating

and distinguishing different exposure routes (e.g., water vs. diet) and the dynamic nature of bioaccumulation processes [6, 68, 69]. The models provide a better understanding of the extent of and the contributors to the variability in the bioaccumulation among metals in various species and at different conditions [67, 70]. The biodynamic models are based on the concept of biodynamics that the accumulation of chemicals occurs as a result of a balance of different fluxes. The models assume that the rate of the fluxes can be realistically determined in controlled experiments with varying concentrations and conditions [67]. Uncertainties are inherent in the application of these empirical rates to different conditions, especially those that are extremely different from the conditions in which the rates have been derived. This limitation can be eliminated by integrating inverse phenomena into unifying concepts. The biokinetic model based on a unifying concept is considered 'mechanistic' from the perspective that physiological rates are estimated from chemical-specific properties of substances and species-specific physiological characteristics of organisms [71, 72]. This method facilitates extrapolation to a wide range of chemicals, organisms, and environmental conditions, without the necessity for case-specific calibration. Quantitative relations between uptake and elimination rate constants, on the one hand, and metal-specific properties and species-specific physiological characteristics, on the other hand, are required for the development of a mechanistic model. Some potential for meeting this requirement has been revealed from the findings during the last decades. Specifically, metal absorption and elimination rates have been related to the filtration rate and species weight, respectively [71, 73]. The affinity of metals for proteins, which are intensively involved in trafficking processes of metals, may explain the uptake rate constant from the dissolved phase [72]. The affinity of metals for the proteins depends on metal charge and atomic radius and reflects the preferences for coordination and ligands [72, 74, 75]. The suggestion on a potential correlation between uptake kinetics and the affinity of metals for protein has been substantiated by the reported relationship between the absorption efficiency and the binding of metals to membrane transport proteins [76]. Efforts have been put in developing bioaccumulation and toxicity models based on the affinity of metals for biological ligands [69, 72, 77]. Different indicators of metal binding to biological ligands and bioaccumulation have been related to various chemical properties of metals [78–82].

The accumulation of metals in organisms occurs as a result of a balance of the uptake from food as well as water and losses. Moreover, metal concentrations in organisms are affected by the growth dilution. Taking these factors into consideration, the concentration of metals accumulated in organisms (C ; $\mu\text{g/g dw}$) can be expressed by the following equation:

$$\frac{dC}{dt} = (k_u \times C_w) + (IR \times AE \times C_f) - (k_{ew} + k_{ef} + g) \times C \quad (5)$$

where k_u (L/g dw/d) is the absorption rate; C_w ($\mu\text{g/L}$) is the dissolved metal concentration; IR (g/g dw/d) is the ingestion rate; AE (/) is the assimilation efficiency; C_f ($\mu\text{g/g}$) is the metal concentration in food; k_{ew} (1/d) is the excretion rate; k_{ef} (1/d) is the egestion rate; and g (1/d) is

the growth rate. The instant concentration of metals in organisms can be solved from Equation 5:

$$C = \frac{(k_u \times C_w) + (IR \times AE \times C_f)}{k_{ew} + k_{ef} + g} \times \left(1 - e^{-(k_{ew} + k_{ef} + g) \times t}\right) \quad (6)$$

The first factor in Equation 5 represents the uptake from the dissolved phase and can be further elaborated based on the absorption efficiency and the filtration rate [69]. In the study of Le et al. [69], the filtration, ingestion, and growth rates were related to the species weight while the absorption and assimilation efficiency was considered metal specific. In addition, elimination rates were parameterised based on both chemical properties of metals and mussel size.

Such a mechanistic model has shown good potential for predicting metal accumulation in the zebra mussel with different size and from various sites. A mechanistic model, which is based on chemical properties of metals and physiological characteristics of organisms while taking site-specific contamination levels into consideration, facilitates a wide extrapolation to different conditions and metals. However, a number of disadvantages are inherent in the current model. The modelling is based on the dissolved metal concentrations without a specification of chemical species and forms of metals in the environment. Although different uptake pathways are included in the model, a distinction of tissue-specific accumulation of metals is not accounted for. Another disadvantage of the model is related to the derivation of the relationship between the rate of physiological processes and the chemical properties of metals. Despite the potential correlation between the uptake kinetics and some chemical properties of metals shown recently, quantitative relationships established between these two factors have hardly been validated. Moreover, relationships between uptake kinetics and chemical properties of metals have usually been derived based on limited experimental data, thereby leading to intrinsic uncertainties in the application of the derived relationships.

4. Metal subcellular fates and partitioning

4.1. Subcellular partitioning

Bioaccumulation is a prerequisite, but not necessarily a reliable indicator of metal toxicity due to the species-specific capacity of organisms to detoxify the metals accumulated [9, 59, 82]. The detoxifying mechanisms have been suggested to account for the tolerance of organisms [83]. Besides metal uptake, metal toxicity is determined by the subcellular partitioning of metals in organisms [83]. Metals accumulated are distributed to different cellular components and sequestered by binding to proteins or peptides (e.g., metallothionein and glutathione) and granules [10, 83–86]. Therefore, only parts of metals are accumulated in sensitive cellular fractions [9]. From a toxicology perspective, metals accumulated have been divided into two fractions, that is, metal-sensitive fraction (MSF) and biologically detoxified metal (BDM) [83, 87, 88]. The former consists of metals in mitochondria, associated with heat-denaturable

proteins (HDPs), lysosomes, and microsomes. The latter includes metals in association with heat-stable proteins (HSPs) or metallothionein (MT)-like proteins and metal-rich granules.

It has been suggested that sub-lethal toxicity is accompanied with changes in subcellular partitioning, especially when the threshold is exceeded, that is, saturation of detoxification mechanisms [83]. Eyckmans et al. [89] showed the relationship between the tolerance of three freshwater fish species to excess Cu and the subcellular partitioning of Cu. The subcellular partitioning of metals may divulge potential mechanisms of toxicity as well as the fate of accumulated metals [90]. For instance, the binding of metals to cytosolic proteins such as MT has widely been demonstrated to alleviate toxic effects [91]. In contrast, the association of metals such as Cd to mitochondria modulates oxidative phosphorylation, followed by decreases in ATP production [92]. Similarly, the binding of Cd to the nucleus is potentially toxic because it can lead to DNA damage and stimulate mutagenesis [93, 94]. Different ligands are included in the HSP fraction, that is, amino acids, glutathione, and metallothioneins [95]. This fraction determines the tolerance and resistance of organisms to metals [83, 84, 87]. The relationships between subcellular partitioning and metal toxicity are complicated because of the different physiological functions of each fraction. The accumulation of metals in the lysosomes and microsomes might reflect the storage for eventual elimination if metals are mainly in the lysosomal fraction [85, 96]. In contrast, metals associated with microsomes can indicate toxicity because of the presence of fragmented endoplasmic reticulum, which is involved in the synthesis and transport of proteins [85, 87]. In addition, mitochondria is the most sensitive fraction [85]. Because of these factors, subcellular partitioning has recently been included in the assessment of metal toxicity.

4.2. Assessment based on metallothionein induction

The induction of MT has been integrated in various monitoring programmes and ecotoxicological assessment. However, recent reviews on the use of MT induction as biomarkers of metal exposure and toxicity raise questions on the validity of this method [11, 97]. The main concern comes from the lack of well-described time- and dose-dependent MT induction, while the induction of MT widely varies, depending on metals, species, and environmental conditions [97]. Further concern results from the lack of well-described relationships between MT induction and metal exposure, metal accumulation, and biological effects [11]. The validity of the use of MT induction as a biomarker is questionable because of the lack of enhancement in MT induction in response to metal exposure and/or the insignificant relationships between metal and MT concentrations [11]. As explained in the reviews by these authors, these results are attributed to different factors determining the cytosolic free metal ion concentrations, the presence of different MT isoforms in different tissues with various physiological functions, and turnover kinetics of MT as well as the high variability of MT induction and the basal level [11]. These factors should be taken into consideration in the assessment of metal exposure and toxicity based on MT induction. Another concern is related to the selection of organs/tissues as the target in the assessment. The digestive gland has widely been used as the target tissue [98, 99] from the point of view that this is the long-term storage organ. In addition, this organ plays an important role in different processes such as immune defence, homeostasis, xenobiotic

mechanisms, elimination, and detoxification [100–103]. Another reason for the use of the digestive gland is that this organ has the highest amount of proteins, including MT [104]. However, the suitability of the digestive ligand is doubtful because of the influence of natural factors and physiological changes on the induction of MT in this organ [104, 105]. As a major tissue for metal uptake, the induction of MT in gills is expected to represent the response of organisms to the ongoing exposure or accidental pollution [106]. Because of these issues, a single value of MT concentrations in the whole soft tissue or in a specific tissue is not always good indicator of metal exposure or effects on biota. As suggested by Le et al. [11], the induction of MT needs to be considered in relation to a number of factors as briefly mentioned above.

4.3. Assessment based on subcellular partitioning

Binding to MT is not the only mechanism for organisms to detoxify and to sequester excess metals. Metals can be associated with insoluble complexes in granules or lysosomes as mentioned above. The proportion of metals bound to MT is not always the dominant fraction and so MT induction is not necessarily a reliable indicator of metal exposure and biological effects. According to Adams et al. [107], the kinetic partitioning of metals into MSF and BDM is the key factor for an understanding of the exposure-toxicity relationship. This has been demonstrated by the shift of Cu accumulated in gills of common carp from MSF to BDM in the first period of the Cu exposure [89]. The ‘Spillover’ hypothesis has been used to simulate the relationship between metal bioaccumulation and toxicity. This term indicates the situation when the capacity of organisms to detoxify excess metals is overwhelmed, thereby resulting in adverse effects [66]. From a broad perspective, spillover is assumed to occur when the uptake rate exceeds a combination of the detoxification and elimination rates, leading to the accumulation of metals in the MSF fractions [9, 108]. This hypothesis has been used in some approaches for assessing metal toxicity.

The first approach is based on the ratio between MSF and MDP fractions or the relative contribution of the MSF and MDF fractions in other words [89, 90]. This approach is simplistic and does not provide a full description of toxicity [90]. There is not constant partitioning between MSF and BDM. Metals can be exchanged between the two fractions, and only a small fraction of metals occurs as free ions [90]. The use of the relative distribution between these two pools may lead to misinterpretation of spillover [90, 92, 109]. For instance, the increase in the amount of metals bound to MSF might be hidden by the lack of changes in the relative proportion, while the accumulation in all compartments increases. This has been illustrated by increases in metal concentrations in both BDM and MSF fractions in yellow eels [110], mummichogs [111], or yellow perch [85].

The second approach is to develop a relationship between MSF and toxic effects [112], assuming that adverse impacts on organisms increase with increasing amounts of metals in MSF. This method is based on the hypothesis that spillover occurs when certain detoxification mechanisms are overwhelmed and excess metals are accumulated in MSF, leading to adverse effects.

The third approach is based on the kinetics of metal accumulation in MSF [108, 113, 114]. The accumulation of metals in MSF can be described as a balance of uptake (e.g., from the dis-

solved), losses via elimination, and the partitioning of the metals to BDM, expressed by the detoxification rate:

$$MIT = (k_u \times C_w) = (k_{detox} + k_e) \times C_{IT} \quad (7)$$

where MIT (mol/g/d) is the metal influx threshold ; k_u (L/g/d) is the uptake rate constant; C_w (mol/L) is the dissolved metal concentration; k_{detox} (1/d) is the detoxification rate constant; k_e (1/d) is the elimination rate constant; C_{IT} (mol/g) is the metal accumulation in MSF at the influx threshold.

This approach is supported by a negative correlation between elimination and detoxification [113, 115]. However, disadvantages are inherent in the assumption of this method that spillover occurs when the uptake rate exceeds combination of the elimination and detoxification rates, that is, no metal is accumulated in MSF below the threshold. Previous studies have indicated that spillover does not happen at low-exposure concentrations. In other words, there is no threshold below which the accumulation of metals in the sensitive fractions does not occur [84, 85].

5. Biological responses to metal exposure

Available approaches based on the induction of MT or taking into account subcellular partitioning of metals have shown some limitations in predicting metal toxicity as mentioned in the previous section. Moreover, the target organ or tissue is species specific, depending on the kinetics of metal accumulation, that is, the differences in absorption, distribution, and excretion [6]. A method that has been demonstrated to be more applicable to different species is to predict metal toxicity based on the accumulation of metals at biological surfaces, which allows for taking into account interactions between organisms and metals at biological surfaces.

5.1. Biotic Ligand Model

Interactions at the water-organism interface have been integrated in the Biotic Ligand Model (BLM). The conceptual framework of the BLM originates from two models: the gill surface interaction model and the free ion activity model (FIAM) [116–118]. The FIAM model assumes that free ions are the main reactive species of metals, determining metal bioavailability and toxicity [116, 117]. The FIAM model has then been extended to take into account the interactions of metals with organisms. One example is the fish gill surface interaction model [119], which has been developed by integrating conditional metal-gill surface binding constants to a geochemical speciation model. On the basis of the fish gill surface interaction model, the BLM has been developed to facilitate the application to various species, metals, and exposure conditions as well. According to the concept of the BLM, environmental geochemistry as well as toxicology principles are taken into consideration in determining the fraction of metals that

provokes effects on biota [119–121]. Initial toxicology bases of the model were effects of trace metals on ionoregulation. Apical (e.g., Na^+ and Ca^{2+} channels) and basolateral (e.g., Na^+/K^+ - and Ca^{2+} -ATPase) transport channels, exchangers, and enzymes are negatively charged proteins that potentially bind to metal cations [122].

With the principles mentioned above, the BLM has shown substantial advantages in assessment of metal bioavailability. By including interactions of metal ions with both biotic and abiotic ligands, the BLM might provide more reliable estimates of metal bioavailability and toxicity. Moreover, the accumulation of metal ions at the physiological active sites of toxic actions, which determines toxic effect, is, in principle, distinguished from the total body burden [6]. Another advantage of the BLM is the inclusion of environmental conditions in estimating metal bioavailability and subsequent application to site-specific assessment [6]. Furthermore, the BLM provides a mechanistic understanding of metal-organism interactions as well as a mechanistic interpretation of metal toxicity, as illustrated by the results in the study of Le et al. [123]. The BLM has potential for assessing the toxicity of metal mixtures [123, 124]. Norwood et al. [124] suggested that metal-metal interactions can be predicted based on known stability constants. If two metals compete for the same binding site, the total amount of the metals bound to the biotic ligands determines combined effects [123]. Alternatively, the accumulation of individual metals at the biotic ligands can be used as inputs to the response addition model for estimating toxicity of metal mixtures [124, 125]. This suggestion has been demonstrated by the increasing application of the BLM for predicting joint toxicity of metals. Liu et al. [126] predicted the toxicity of metal mixtures based on the simple sum of the fraction of biotic ligands occupied by individual metals. This approach allows for taking into account the metal-specific affinity for binding sites of biotic ligands, but not the metal-specific toxic potency. In other studies, the BLM concepts are usually combined with conventional concepts of mixture toxicity such as concentration addition. As such, the specific toxic potency of metals is taken into account in the estimates. For instance, the toxicity of metal mixtures has been related to the toxic equivalent quotient (TEQ), which is based on the accumulation of metal ions at the binding sites of biotic ligands while giving consideration to the metal-specific toxic potency [123, 126]. The toxic unit (TU) is the concept mostly integrated in the BLM-based approaches for estimating joint toxicity of metals.

5.2. The electrostatic model

The electrostatic model originates from the reported differences between the ion concentration at the root plasma membrane surface and the ion concentration in the external medium, which is induced by the negative charge at the plasma membrane surface [127, 128]. The surface potential affects the activity of ions at the plasma membrane surface via electrostatic attraction or repulsion. In addition, the potential influences the difference in the electrical potential across the membrane, which stimulates the transport of ions through the membrane. A number of studies have shown the effects of electrical potential at the plasma membrane surface on cation uptake [127, 128]. The principle effect of the electrical potential at the plasma membrane surface is to control the ion activity at the surface [129].

The model allows for integrating plant-ion interactions in predicting metal toxicity [130–132]. In addition, the model gives consideration to the interactions among ions while estimating metal toxicity [132]. The electrostatic model has been applied to simulate effects of major cations on the toxicity of trace metal ions [131–134]. According to the principle of the electrostatic model, additions of cations to the bulk medium reduce the negativity of the electrical potential at the plasma membrane surface, thereby leading to decreases in the negativity of the electrical potential at the plasma membrane surface accompanied by reduced accumulation of trace metal ions at the plasma membrane surface and subsequent alleviation of toxic effects [135]. The electrostatic approach therefore might provide additional explanation for interpreting ion-ion interactions as well as effects of major cations on the toxicity of trace metal ions besides the competitive binding assumed in the BLM. According to the electrostatic principles, three mechanisms have been suggested to account for ameliorative effects of Ca^{2+} on metal toxicity [128]. The first mechanism is the electrostatic displacement of trace metal ions at the plasma membrane surface by Ca^{2+} . The second is the restoration of Ca^{2+} at the cell surface in response to low levels of surface Ca^{2+} . This mechanism is mediated in order to avoid the inhibition of the plant growth induced by the low contents of Ca^{2+} at the cell surface. The third mechanism covers other types of interactions between Ca^{2+} and trace metal ions, such as the blockade of the ion channel [136]. Despite the toxicant- and major cation-specific relative importance of these mechanisms, mechanism 1 generally occurs in all cases while the contribution of mechanism 2 is minor [128]. The significance of mechanism 3 is trace metal ion specific. The relative influence of major cations such as Ca^{2+} and Mg^{2+} on the toxicity of trace metal ions varies, depending on the trace metal ions and species [128]. For instance, Ca^{2+} has stronger effects on the toxicity of Al^{3+} to soybean than Mg^{2+} [137, 138]. In wheat, Mg^{2+} is more effective than Ca^{2+} in inhibiting Zn^{2+} toxicity, whereas Mg^{2+} does not have effects on the toxicity of Al^{3+} [128].

5.3. WHAM-based bioavailability model

Originally developed for determining and quantifying chemical forms and species of metals in different phases in the environment, speciation models such as WHAM have recently been used for estimating metal bioavailability and toxicity (hereafter referred to as the WHAM-based model). In this method, humic acid is considered a surrogate to biological surfaces [139–144]. As such, the amount of metals bound to humic acid computed by WHAM is used for estimating metal toxicity for different species. In the application of the WHAM-based model to plants, humic acid is used as a surrogate for the root surface [144]. In other words, the interactions of metal ions and the root surface are simulated by the interactions with humic acid.

The WHAM-based model was initiated from the similarities in terms of chemical structure between the root surface and humic acid. The root surface is a heterogeneous mixture of various metal-binding functional groups [145–147]. Similarly, humic acid possesses a heterogeneous mixture of functional groups, mainly carboxylic and phenolic acids [141]. Furthermore, the ratio of 1:2 between the site densities of phenolic and carboxylic groups assumed in WHAM [40, 41] lies in the range from 0.49:1 to 1:1 reported for root cell walls of different plant

species [148]. Because of this similar structure of the root surface and humic acid, metal accumulation on the root surface and metal binding to humic acids are influenced by similar factors. These factors include electrostatic interactions and chemical heterogeneity [127, 128, 149, 150]. The relevance of using metal binding to humic acid to represent metal accumulation at biological surfaces is further substantiated by the nature of the sorption of metal cations to the biological surfaces [151].

The WHAM-based model allows for the interactions between metal ions to be integrated in modelling metal toxicity [144]. Specifically, interactions between ions are considered in modelling metal binding to humic acid in WHAM and therefore accounted for in estimating metal accumulation at the biological surfaces. Another advantage of the WHAM-based model is related to the availability of binding constants in WHAM, which facilitates a wide application to different metals [144]. Moreover, previous studies have shown small variations in the binding constants among different species [145, 147, 152, 153]. For instance, the logarithm of the binding constants of Cd to the carboxylic group on the membrane surface of bacteria, fungi, and plants were in the range 3.3–3.5. Moreover, the cell walls of algae, maize, soybeans, and higher plants have similar titration curves [154–156]. These results indicate the applicability of a single set of binding constants to different organisms.

In the study of Le et al. [144], the amount of metals bound to humic acid was computed with WHAM in which metal sorption to humic substances is simulated by using a structured formulation of discrete, chemically plausible binding sites for protons and metals. This allows the creation of regular arrays of bidentate and tridentate binding sites for metals. Metal aquo ions compete with their first hydrolysis products, protons, and other metals for binding sites. In addition to the intrinsic metal binding strength to uncharged molecules, electrostatic effects were taken into consideration while predicting metal binding to humic acid. Le et al. [144] showed the potential of the WHAM-HA model for predicting both total and internalised metal concentrations in roots. Another example of the approaches based on the geochemical equilibrium in WHAM is the WHAM- F_{TOX} model for estimating metal toxicity [140, 143]. In this approach, mixture toxicity was simulated as a function of the metal-specific toxic potency and the amount of metal ions bound to the biological surfaces computed with WHAM.

Author details

T.T. Yen Le

Address all correspondence to: yen.le@uni-due.de

Department of Aquatic Ecology, University of Duisburg-Essen, Germany

References

- [1] Del Castillo P, Chardon WJ, Salomons W. Influence of cattle-manure slurry application on the solubility of cadmium, copper, and zinc in a manured acidic, loamy-sand soil. *Journal of Environmental Quality*. 1993;22:689–997.
- [2] Janssen RPT. Speciation of heavy metal ions as influenced by interactions with montmorillonite, Al hydroxide polymers and citrate [thesis]. Landbouwwuniversiteit Wageningen, ISBN 90 – 5485-399-9; 1995.
- [3] John DA, Leventhal JS. Bioavailability of metals. In: Du Bray EA, editor. Preliminary Compilation of Descriptive Geoenvironmental Mineral Deposit Models. U.S. Department of the Interior. U.S. Geological survey; 1995.
- [4] Goody DC, Shand P, Kinniburgh DG, Van Riemsdijk WH. Field-based partition coefficients for trace elements in soil solutions. *European Journal of Soil Science*. 1995;46:265–285.
- [5] Landner L, Reuther R. Metals in society and in the environment: a critical review of current knowledge on fluxes, speciation, bioavailability and risk for adverse effects of copper, chromium, nickel and zinc. Dordrecht, the Netherlands: Kluwer Academic Publishers; 2004.
- [6] Fairbrother A, Wenstel R, Sappinton K, Wood W. Framework for metals risk assessment. *Ecotoxicology and Environmental Safety*. 2007;68:145–227.
- [7] Tchounwou PB, Yedjou CG, Patlolla AK, Sutton D. Heavy metals toxicity and the environment. *EXS*. 2012;101:133–164.
- [8] Mertz W. Newer essential trace-elements, chromium, tin, vanadium, nickel and silicon. *Proceedings of the Nutrition Society*; 1974;33:307–313.
- [9] Rainbow PS. Trace metal concentrations in aquatic invertebrates: why and so what? *Environmental Pollution*. 2002;120:497–507.
- [10] Vijver MG, Van Gestel CAM, Lanno RP, Van Straalen NM, Peijnenburg WJGM. Internal metal sequestration and its ecotoxicological relevance: a review. *Environmental Science and Technology*. 2004;38:4705–4712.
- [11] Le TTY, Zimmermann S, Sures B. How does the metallothionein induction in bivalves meet the criteria for biomarkers of metal exposure? *Environmental Pollution*. 2016;212:257–268.
- [12] Sunda W, Guillard RRL. The relationship between cupric ion activity and the toxicity of copper to phytoplankton. *Journal of Marine Research*. 1976;34:511–529.
- [13] Sunda W, Guillard RRL. The relationship between cupric ion activity and the toxicity of copper to phytoplankton. *Journal of Marine Research*. 1976;34:511–529.

- [14] Senesi N. 1992. Metal-humic substance complexes in the environment. Molecular and mechanistic aspects by multiple spectroscopic approach. In: Adriano DC, editor. *Biogeochemistry of Trace Elements*. Boca Raton, FL: Lewis; 1992. pp. 429–451.
- [15] Uzu G, Sobanska S, Aliouane Y, Pradere P, Dumat C. Study of lead phytoavailability for atmospheric industrial micronic and sub-micronic particles in relation with lead speciation. *Environmental Pollution*. 2009;157:1178–1185.
- [16] Ure AM. Trace elements in soil: their determination and speciation. *Fresenius Journal of Analytical Chemistry*. 1990;337:577–581.
- [17] Tack FMG, Verloo MG. Chemical speciation and fractionation in soil and sediment heavy metal analysis: a review. *International Journal of Environmental Analytical Chemistry*. 1995;59:225–238.
- [18] Carbonaro RF, Stone AT. Speciation of chromium(III) and cobalt(III) (amino)carboxylate complexes using capillary electrophoresis. *Analytical Chemistry*. 2005;77:155–164.
- [19] Clough R, Lindsay R, Drennar H, Harrinton CF, Hill SJ, Tyson JF. Atomic spectrometry update: elemental speciation. *Journal of Analytical Atomic Spectrometry*. 2012;27:1185–1224.
- [20] Rieuwerts J, Thornton I, Farago M, Ashmore M. Quantifying the influence of soil properties on the solubility of metals by predictive modeling of secondary data. *Chemical Speciation and Bioavailability*. 1998;10:83–94.
- [21] Bonten LTC, Groenenberg JE, Weng L, Van Riemsdijk WH. Use of speciation and complexation models to estimate heavy metal sorption in soils. *Geoderma*. 2008;146:303–310.
- [22] Sauve S, Hendershot W, Allen HE. Solid-solution partitioning of metals in contaminated soils: dependence on pH, total metal burden, and organic matter. *Environmental Science and Technology*. 2000a;34:1125–1131.
- [23] Cances B, Ponthieu M, Castre Rouelle M, Aubry E, Benedetti MF. Metal ions speciation in a soil and its solution: experimental data and model results. *Geoderma*. 2003;113:341–355.
- [24] Tipping E, Rieuwerts J, Pan G, Ashmore MR, Loftis S, Hill MTR, Farago ME, Thornton I. The solid-solution partitioning of heavy metals (Cu, Zn, Cd, Pb) in upland soils of England and Wales. *Environmental Pollution*. 2003;125:213–225.
- [25] Smolders E, Buekers J, Oliver I, McLaughlin MJ. Soil properties affecting toxicity of zinc to soil microbial properties in laboratory-spiked and field-contaminated soils. *Environmental Toxicology and Chemistry*. 2004;23:2633–2640.
- [26] Le TTY, Hendriks AJ. Uncertainties associated with lacking data for predictions of solid-solution partitioning of metals in soil. *Science of the Total Environment*. 2014;490:44–49.

- [27] Voegelin A, Kretzschmar R. Modelling sorption and mobility of cadmium and zinc in soils with scaled exchange coefficients. *European Journal of Soil Science*. 2003;54:387–400.
- [28] Lofts S, Tipping E. An assemblage model for cation binding by natural particulate matter. *Geochim Cosmochim Acta*. 1998;62:2609–2625.
- [29] Weng LP, Temminghoff EJ, Lofts S, Tipping E, Van Riemsdijk WH. *Environmental Science and Technology*. 2002;36:4804–4810.
- [30] Gustafsson JP, Pechova P, Berggren D. Modeling metal binding to soils: the role of natural organic matter. *Environmental Science and Technology*. 2003;37:2767–2774.
- [31] Dijkstra JJ, Meeussen JCL, Comans RNJ. Leaching of heavy metals from contaminated soils: an experimental and modeling study. *Environmental Science and Technology*. 2004;38:4390–4395.
- [32] Rodrigues SM, Henriques B, da Silva EF, Pereira ME, Duarte AC, Groenenberg JE, Romkens PFAM. Evaluation of an approach for the characterization of reactive and available pools of 20 potentially toxic elements in soils: Part II – Solid-solution partition relationships and ion activity in soil solutions. *Chemosphere*. 2010;81:1560–1570.
- [33] Groenenberg JE, Dijkstra JJ, Bonten LTC, de Vries W, Comans RNJ. 2012. Evaluation of the performance and limitations of empirical partition-relations and process based multisurface models to predict trace element solubility in soils. *Environmental Pollution*. 2012;166:98–107.
- [34] Sauve S, Norvell WA, McBride M, Hendershot W. Speciation and complexation of cadmium in extracted soil solutions. *Environmental Science and Technology*. 2000b;34:291–296.
- [35] Degryse F, Smolders E, Parker DR. Partitioning of metals (Cd, Co, Cu, Ni, Pb, Zn) in soils: concepts, methodologies, prediction and applications – a review. *European Journal of Soil Science*. 2009;60:590–612.
- [36] Groenenberg JE, Romkens PFAM, Comans RNJ, Luster J, Pampura T, Shotbolt L, Tipping E, de Vries W. Transfer functions for solid-solution partitioning of cadmium, copper, nickel, lead and zinc in soils: derivation of relationships for free metal ion activities and validation with independent data. *European Journal of Soil Science*. 2010;61:58–73.
- [37] Westall JC, Zachary JL, Morel FMM. MINEQL: A Computer Program for the Calculation of the Chemical Equilibrium Composition of Aqueous Systems. Tech Note No. 18, Ralph M. Parsons Laboratory, Massachusetts Institute of Technology, Cambridge, MA; 1976.
- [38] Sposito G, Mattigod SV. GEOCHEM: a computer program for calculation of chemical equilibria in soil solutions and other natural water systems. Kearny Foundation of Soil Science, University of California, Riverside; 1980.

- [39] Kinniburgh DG, Van Riemsdijk WH, Koopal LK, Borkovic M, Benedetti MF, Avena MJ. Ion binding to natural organic matter: competition, heterogeneity, stoichiometry and thermodynamic consistency. *Colloids and Surfaces A*. 1999;151:147–166.
- [40] Tipping E. WHAM – a chemical equilibrium model and computer code for waters, sediments, and soils incorporating a discrete site/electrostatic model of ion-binding by humic substances. *Computers & Geosciences*. 1994;20:973–1023.
- [41] Tipping E. Humic ion-binding Model VI: an improved description of the interactions of protons and metal ions with humic substances. *Aquatic Geochemistry*. 1998;4:3–48.
- [42] Dzombak DA, Morel FMM. *Surface Complexation Modeling: Hydrous Ferric Oxide*. New York: John Wiley & Sons; 1990.
- [43] Hiemstra T, de Wit JCM, Van Riemsdijk WH. Multisite proton adsorption modeling at the solid-solution interface of (hydr)oxides: a new approach: II. Application to various important (hydr)oxides. *Journal of Colloid and Interface Science*. 1989;133:105–117.
- [44] Evans LJ, Barabash SJ, Lumsdon DG, Gu X. Application of chemical speciation modelling to studies on toxic element behaviour in soils. In: Hooda PS, editor. *Toxic Elements in Soils*. Chichester, England: John Wiley & Sons; 2010. pp. 199–226.
- [45] Meers E, Unamuno V, Vandegheuchte M, Vanbroekhoven K, Geebelen W, Samson R, Vangronsveld J, Diels L, Ruttens A, Du Laing G, Tack F. Soil-solution speciation of Cd as affected by soil characteristics in unpolluted and polluted soils. *Environmental Toxicology and Chemistry*. 2005;24:499–509.
- [46] Tipping E, Berggren D, Mulder J, Woof C. Modelling the solid-solution distributions of protons, aluminium, base cations and humic substances in acid soils. *European Journal of Soil Science*. 1995;46:77–94.
- [47] Weng L, Temminghoff EJM, Van Riemsdijk WH. Contribution of individual sorbents to the control of heavy metal activity in sandy soil. *Environmental Science and Technology*. 2001;35:4436–4443.
- [48] Kuipers SF. *Bodemkunde*. Culumborg, the Netherlands; 1984.
- [49] McBride MB. *Environmental Chemistry of Soils*. New York: Oxford University Press; 1994.
- [50] Meeussen JCL. ORCHESTRA: an object-oriented framework for implementing chemical equilibrium models. *Environmental Science and Technology*. 2003;37:1175–1182.
- [51] Dijkstra JJ, Meeussen JCL, Comans RNJ. Evaluation of a generic multisurface sorption model for inorganic soil contaminants. *Environmental Science and Technology*. 2009;43:6196–6201.

- [52] Allison JD, Brown DS, Novo-gradac KJ. MINTEQA2/PRODEFA2, Geochemical assessment model for environmental systems: version 3.11 databases and version 3.0 user's manual Environmental Research Laboratory, US EPA, Athens, GA; 1991.
- [53] Benedetti MF, Milne CJ, Kinniburgh DG, Van Riemsdijk WH, Koopal LK. Metal ion binding to humic substances: application of the non-ideal competitive adsorption model. *Environmental Science and Technology*. 1995;29:446–457.
- [54] Kinniburgh DG, Milne CJ, Benedetti MF, Pinheiro JP, Filius J, Koopal LK, Van Riemsdijk W. Metal ion binding by humic acid: application of the NICA-Donnan Model. *Environmental Science and Technology*. 1996;30:1687–1698.
- [55] Milne CJ, Kinniburgh DG, Tipping E. Generic NICA-Donnan model parameters for proton binding by humic substances. *Environmental Science and Technology*. 2001;35:2049–2059.
- [56] Milne CJ, Kinniburgh DG, Van Riemsdijk WH, Tipping E. Generic NICA-Donnan model parameters for metal-ion binding by humic substances. *Environmental Science and Technology*. 2003;37:958–971.
- [57] Chapman PM, Allen HE, Godtfredsen K, Z'Graggen MN. Evaluation of bioaccumulation factors in regulating metals. *Environmental Science and Technology*. 1996;30:448–452.
- [58] McGeer J, Brix KV, Skeaff JM, DeForest DK. The use of bioaccumulation criteria for hazard identification of metals. ICMF Fact Sheet on Environmental Risk Assessment. Published by the International Council on Mining and Metals (ICMM), London; 2002.
- [59] McGeer JC, Brix KV, Skeaff JM, DeForest DK, Brigham SI, Adams WJ, Green A. Inverse relationship between bioconcentration factor and exposure concentration for metals: implications for hazard assessment of metals in the aquatic environment. *Environmental Toxicology and Chemistry*. 2003;22:1017–1037.
- [60] Le TTY. Modelling bioaccumulation and toxicity of metal mixtures [thesis]. Radboud University Nijmegen, the Netherlands. ISBN: 978-94-91066-06-1; 2012. <http://repository.ubn.ru.nl/bitstream/2066/100831/1/100831.pdf>
- [61] Simkiss K, Taylor MG. Metal fluxes across membranes of aquatic organisms. *Reviews in Aquatic Science*. 1989;1:173–188.
- [62] McDonald DG, Wood CM. Branchial mechanisms of acclimation to metals in freshwater fish. In: Rankin JC, Jensen FB, editors. *Fish Ecophysiology*. London, UK: Chapman & Hall; 1993. p. 297–321.
- [63] McKim JM. Physiological and biochemical mechanisms that regulate the accumulation and toxicity of environmental chemicals in fish. In: Hamelink JL, Landrum PF, Bergman HL, Bensen WH, editors. *Bioavailability: Physical, Chemical and Biological Interactions*. Boca Raton, FL, US: CRC; 1994. pp. 179–201.

- [64] Newman MC. Bioaccumulation in Quantitative Methods in Aquatic Ecotoxicology. Boca Raton, FL, US: CRC; 1995.
- [65] Wood CM. Toxic responses of the gill. In: Schlenk DW, Benson WH, editors. Target Organ Toxicity in Marine and Freshwater Teleosts. Washington, DC, US: Taylor & Francis; Vol. 1: Organs; 2001.
- [66] Mason AZ, Jenkins KD. Metal detoxification in aquatic organisms. In: Tessier A, Turner DR, editors. Metal Speciation and Bioavailability in Aquatic Systems. Chichester, UK: John Wiley & Sons; 1995. pp. 479–608.
- [67] Luoma SN, Rainbow PS. Why is metal bioaccumulation so variable? Biodynamics as a unifying concept. *Environmental Science and Technology*. 2005;39:1921–1931.
- [68] Ahlf W, Drost W, Heise S. Incorporation of metal bioavailability into regulatory frameworks – metal exposure in water and sediment. *Journal of Soils and Sediments*. 2009;9:411–419.
- [69] Le TTY, Leuven RSEW, Hendriks AJ. Modeling metal bioaccumulation in the invasive mussels *Dreissena polymorpha* and *Dreissena rostriformis bugensis*. *Environmental Toxicology and Chemistry*. 2011;30:2825–2830.
- [70] Wang W-X, Fisher NS. Delineating metal accumulation pathways for marine invertebrates. *Science of the Total Environment*. 1999;237/238:459–472.
- [71] Hendriks AJ, Heikens A. The power of size. 2. Rate constants and equilibrium ratios for accumulation of inorganic substances related to species weight. *Environmental Toxicology and Chemistry*. 2001;20:1421–1437.
- [72] Veltman K, Huijbregts MAJ, Van Kolck M, Wang W-X, Hendriks AJ. Metal bioaccumulation in aquatic species: qualification of uptake and elimination rate constants using physicochemical properties of metals and physiological characteristics of species. *Environmental Science and Technology*. 2008;42:852–858.
- [73] Baines SB, Fisher NS, Kinney EL. Effects of temperature on uptake of aqueous metals by blue mussels *Mytilus edulis* from Arctic and temperate waters. *Marine Ecology Progress Series*. 2006;308:117–128.
- [74] Bell RA, Ogden N, Kramer JR. The biotic ligand model and a cellular approach to class B metal aquatic toxicity. *Comparative Biochemistry and Physiology C*. 2002;133:175–188.
- [75] Handy RD, Eddy FB. Transport of solutes across biological membranes in eukaryotes: an environmental perspective. In: van Leeuwen HP, Koster W, editors. Physicochemical Kinetics and Transport at Biointerfaces. IUPAC Series on Analytical and Physical Chemistry of Environmental Systems. New York: John Wiley & Sons, Vol. 9; 2004.

- [76] Bryan GW. Pollution due to heavy metals and their compounds. In: Kinne O, editor. Marine Ecology. West Sussex, UK: John Wiley & Sons; 1984.
- [77] Newman MC, McCloskey JT. Predicting relative toxicity and interactions of divalent metal ions: microtox bioluminescence assay. Environmental Toxicology and Chemistry. 1996;15:275–281.
- [78] Van Kolck M, Huijbregts MAJ, Veltman K, Hendriks AJ. Estimating bioconcentration factors, lethal concentrations and critical body residues of metals in the mollusks *Perna viridis* and *Mytilus edulis* using ion characteristics. Environmental Toxicology and Chemistry. 2008;27:272–276.
- [79] Zhou D-M, Li L-Z, Peijnenburg WJGM, Ownby DR, Hendriks AJ, Wang P, Li D-D. A QICAR approach for quantifying binding constants for metal-ligand complexes. Ecotoxicology and Environmental Safety. 2011;74:1036–1042.
- [80] McCloskey JT, Newman MC, Clark SB. Predicting the relative toxicity of metal ions using ion characteristics: microtox bioluminescence assay. Environmental Toxicology and Chemistry. 1996;15:1730–1737.
- [81] Le TTY, Hendriks AJ. Relationships between absorption efficiency of elements in mammals and chemical properties. Critical Reviews in Toxicology. 2013;43:800–809.
- [82] Rosen G, Rivera-Duarte I, Chadwick DB, Ryan A, Santore RC, Paquin PR. Critical tissue copper residues for marine bivalve (*Mytilus galloprovincialis*) and echinoderm (*Strongylocentrotus purpuratus*) embryonic development: conceptual, regulatory and environmental implications. Marine Environmental Research. 2008;66:327–336.
- [83] Wallace WG, Lee B-G, Luoma SN. Subcellular compartmentalization of Cd and Zn in two bivalves. I. Significance of metal-sensitive fractions (MSF) and biologically detoxified metal (BDM). Marine Ecology Progress Series. 2003;249:183–197.
- [84] Campbell PGC, Guguere A, Bonneris E, Hare L. Cadmium-handling strategies in two chronically exposed indigenous freshwater organisms-the yellow perch (*Ferca flavescenes*) and the floater mollusc (*Pyganodon grandis*). Aquatic Toxicology. 2005;72:83–97.
- [85] Giguere A, Campbell PGC, Hare L, Couture P. Sub-cellular partitioning of cadmium, copper, nickel and zinc in indigenous yellow perch (*Perca flavescenes*) sampled along a polymetallic gradient. Aquatic Toxicology. 2006;77:178–189.
- [86] Voets J, Redeker ES, Blust R, Bervoets L. Differences in metal sequestration between zebra mussels from clean and polluted filed locations. Aquatic Toxicology. 2009;93:53–60.
- [87] Bonneris E, Perceval O, Masson S, Hare L, Campbell P. Sub-cellular partitioning of Cd, Cu and Zn in tissues of indigenous unionid bivalves living along a metal exposed gradient and links a metal-induced effects. Environmental Pollution. 2005;135:195–208.
- [88] Wadige CPMM, Taylor AM, Maher WA, Krikowa F. Bioavailability and toxicity of zinc from contaminated freshwater sediments: linking exposure-dose-response relation-

ships of the freshwater bivalve *Hyridella australis* to zinc-spiked sediments. *Aquatic Toxicology*. 2014;156:179–190.

- [89] Eyckmans M, Blust R, De Boeck G. Subcellular differences in handling Cu excess in three freshwater fish species contributes greatly to their differences in sensitivity to Cu. *Aquatic Toxicology*. 2012;118–119:97–107.
- [90] Kamunde C. Early subcellular partitioning of cadmium in gill and liver of rainbow trout (*Oncorhynchus mykiss*) following low-to-near-lethal waterborne cadmium exposure. *Aquatic Toxicology*. 2009;91:291–301.
- [91] Roesijadi G. Metallothionein and its role in toxic metal regulation. *Comparative Biochemistry and Physiology C*. 1996;113:117–123.
- [92] Sokolova IM, Ringwood AH, Johnson, C. Tissue-specific accumulation of cadmium in subcellular compartments of eastern oysters *Crassostrea virginica* Gmelin (Bivalvia: Ostreidae). *Aquatic Toxicology*. 2005;74:218–228.
- [93] Risso-de Faverney C, Devaux A, Lafaurie M, Girar JP, Bailly B, Rahmani R. Cadmium induces apoptosis and genotoxicity in rainbow trout hepatocytes through generation of reactive oxygen species. *Aquatic Toxicology*. 2001;53:65–76.
- [94] Pruski AM, Dixon DR. Effects of cadmium on nuclear integrity and DNA repair efficiency in the gill cells of *Mytilus edulis* L. *Aquatic Toxicology*. 2002;57:127–137.
- [95] Kraemer LD, Campbell PGC, Hare L. A field study examining metal elimination kinetics in juvenile yellow perch (*Perca flavescens*). *Aquatic Toxicology*. 2005;75:108–126.
- [96] Bustamante P, Cosson R, Gallien I, Caurant F, Miramand P. Cadmium detoxification processes in the digestive gland of cephalopods in relation to accumulated camium concentrations. *Marine Environmental Research*. 2002;53:227–241.
- [97] Amiard JC, Amiard-Triquet C, Barka S, Pellerin J, Rainbow PS. Metallothioneins in aquatic invertebrates: their role in metal detoxification and their use as biomarkers. *Aquatic Toxicology*. 2006;76:160–202.
- [98] Lionetto MG, Giordano ME, Caricato R, Pascariello MF, Marinosci L, Schettino T. Biomonitoring of heavy metal contamination along the Salento coast (Italy) by metallothionein evaluation in *Mytilus galloprovincialis* and *Mullus barbatus*. *Aquatic Conservation Marine and Freshwater Ecosystems*. 2001;11:305–310.
- [99] Andreani G, Carpena E, Capranico G, Isani G. Metallothionein cDNA cloning, metallothionein expression and heavy metals in *Scapharca inaequivalvis* along the Northern Adriatic coast of Italy. *Ecotoxicology and Environmental Safety*. 2011;74:366–372.
- [100] Marigomez I, Soto M, Cajaraville MP, Angulo E, Giamberini L. Cellular and subcellular distribution of metals in molluscs. *Microscopy Research and Technique*. 2002;56:358–392.

- [101] Moore MN, Allen JI. A computational model of the digestive gland epithelial cells of marine mussels and its simulated responses to oil-derived aromatic hydrocarbons. *Marine Environmental Research*. 2002;54:579–584.
- [102] McVeigh A, Allen JI, Moore MN, Dyke P, Noble D. A carbon and nitrogen flux model of mussel digestive epithelial cells and their simulated response to pollutants. *Marine Environmental Research*. 2004;58:821–827.
- [103] Rajalakshmi S, Mohandas A. Copper-induced changes in tissue enzyme activity in a freshwater mussel. *Ecotoxicology and Environmental Safety*. 2005;62:140–143.
- [104] Raspor B, Dragun Z, Erk M, Ivankovic D, Pavicic J. Is the digestive gland of *Mytilus galloprovincialis* a tissue of choice for estimating cadmium exposure by means of metallothioneins? *Science of the Total Environment*. 2004;333:99–108.
- [105] Trinchella F, Esposito MG, Simoniello P, Scudiero R. Cadmium, lead and metallothionein contents in cultivated mussels (*Mytilus galloprovincialis*) from the Gulf of Naples (Southern Italy). *Aquaculture Research*. 2013;44:1076–1084.
- [106] Cooper S, Hare L, Campbell PGC. Subcellular partitioning of cadmium in the freshwater bivalve, *Pyganodon grandis*, after separate short-term exposures to waterborne or diet-borne metal. *Aquatic Toxicology*. 2010;100:303–312.
- [107] Adams WJ, Blust R, Borgmann U, Brix KV, DeForst DK, Green AS, Meyer JS, McGeer JC, Paquin PR, Rainbow PS, Wood CM. Utility of tissue residues for predicting effects of metals on aquatic organisms. *Integrated Environmental Assessment and Management*. 2011;7:75–98.
- [108] Croteau M-N, Luoma SN. Predicting dietborne metal toxicity from metal influxes. *Environmental Science and Technology*. 2009;43:4915–4921.
- [109] Wicklund Glynn A. The concentration dependency of branchial intracellular cadmium distribution and influx in the zebrafish (*Brachydanio rerio*). *Aquatic Toxicology*. 1996;35:47–58.
- [110] Rosabal M, Pierron F, Counture P, Baudrimont M, Hare L, Campbell PGC. Subcellular partitioning of non-essential trace metals (Ag, As, Cd, Ni, Pb, and Tl) in livers of American (*Anguilla rostrata*) and European (*Anguilla anguilla*) yellow eels. *Aquatic Toxicology*. 2015;160:128–141.
- [111] Goto D, Wallace WG. Metal intracellular partitioning as a detoxification mechanism for mummichogs (*Fundulus heteroclitus*) living in metal-polluted salt marshes. *Marine Environmental Research*. 2010;69:163–171.
- [112] Dandan L, Dongmei Z, Peng W, Nanyan W, Xiangdong Z. Subcellular Cd distribution and its correlation with antioxidant enzymatic activities in wheat (*Triticum aestivum*) roots. *Ecotoxicology and Environmental Safety*. 2011;74:874–881.

- [113] Chen B-C, Chen W-Y, Ju Y-R, Tsai J-W, Jou L-J, Singh S, Liao C-M. Combining bioaccumulation and coping mechanism to enhance long-term site-specific risk assessment for zinc susceptibility of bivalves. *Chemosphere*. 2011;84:707–715.
- [114] Liao C-M, Ju Y-R, Chen W-Y. Subcellular partitioning links BLM-based toxicokinetics for assessing cadmium toxicity to rainbow trout. *Environmental Toxicology*. 2011;26:600–609.
- [115] Buchwalter DB, Cain DJ, Martin CA, Xie L, Luoma SN, Garland Jr T. Aquatic insect ecophysiological traits reveal phylogenetically based differences in dissolved cadmium susceptibility. *Proceedings of the National Academy of Sciences of the United States of America*. 2008;105:8321–8326.
- [116] Morel FMM. *Principles of Aquatic Chemistry*. New York, US: Wiley-Interscience; 1983.
- [117] Pagenkopf GK. Gill surface interaction model for trace metal toxicity to fishes: role of complexation, pH and water hardness. *Environmental Science and Technology*. 1983;17:342–347.
- [118] Campbell PGC. Interactions between trace metals and aquatic organism: a critique of the free-ion activity model. In: Tessier A, Turner DR, editors. *Metal Speciation and Bioavailability in Aquatic Systems*. Chichester, UK: John Wiley & Sons; 1995. pp. 45–102.
- [119] Playle RC. Modelling metal interactions at fish gills. *Science of the Total Environment*. 1998;219:147–163.
- [120] McGeer JC, Playle RC, Wood CM, Galvez F. A physiologically based biotic ligand model for predicting the acute toxicity of waterborne silver to rainbow trout in freshwaters. *Environmental Science and Technology*. 2000;34:4199–4207.
- [121] Di Toro DM, Allen HE, Bergman HL, Meyer JS, Paquin PR, Santore RC. Biotic ligand model of the acute toxicity of metals. 1. Technical basis. *Environmental Toxicology and Chemistry*. 2001;20:2382–2393.
- [122] Niyogi S, Wood CM. Effects of chronic waterborne and dietary metal exposures on gill metal-binding: implications for the biotic ligand model. *Human and Ecological Risk Assessment*. 2003;4:813–846.
- [123] Le TTY, Vijver MG, Hendriks AJ, Peijnenburg WJGM. Modeling toxicity of binary metal mixtures (Cu^{2+} - Ag^{+} , Cu^{2+} - Zn^{2+}) to lettuce, *Lactuca sativa*, with the biotic ligand model. *Environmental Toxicology and Chemistry*. 2013;32:137–143.
- [124] Norwood WP, Borgmann U, Dixon DG, Wallance A. Effects of metal mixtures on aquatic biota: a review of observations and methods. *Human and Ecological Risk Assessment*. 2003;4:795–811.
- [125] Borgmann U, Norwood WP, Dixon DG. Modelling bioaccumulation and toxicity of metal mixtures. *Human and Ecological Risk Assessment*. 2008;14:266–289.

- [126] Liu Y, Vijver MG, Peijnenburg WJGM. Comparing three approaches in extending biotic ligand models to predict the toxicity of binary metal mixtures (Cu-Ni, Cu-Zn and Cu-Ag) to lettuce (*Lactuca sativa* L.). *Chemosphere*. 2014;112:282–288.
- [127] Wagatsuma T, Akiba R. Low surface negativity of root protoplasts from aluminium-tolerant plant species. *Soil Science and Plant Nutrition*. 1989;35:443–452.
- [128] Kinraide TB. Three mechanisms for the calcium alleviation of mineral toxicities. *Plant Physiology*. 1998;118:513–520.
- [129] Nobel PS. *Physicochemical and Environmental Plant Physiology*. San Diego, CA: Academic; 1961.
- [130] Wang P, Zhou D-M, Peijnenburg WJGM, Li L-Z, Weng N. Evaluating mechanisms for plant-ion (Ca^{2+} , Cu^{2+} , Cd^{2+} or Ni^{2+}) interactions and their effectiveness on rhizotoxicity. *Plant and Soil*. 2010;334:277–288.
- [131] Kopittke PM, Blamey FPC, Wang P, Menzies NW. Calculated activity of Mn^{2+} at the outer surface of the root cell plasma membrane governs Mn nutrition of cowpea seedlings. *Journal of Experimental Botany*. 2011;62:3393–4001.
- [132] Le TTY, Wang P, Vijver MG, Kinraide TB, Hendriks AJ, Peijnenburg WJGM. Delineating ion-ion interactions by electrostatic modeling for predicting rhizotoxicity of metal mixtures to lettuce *Lactuca sativa*. *Environmental Toxicology and Chemistry*. 2014;33:1988–1995.
- [133] Wang P, Kopittke PM, De Schamphelaere KAC, Zhao F-J, Zou D-M, Lock K, Ma Y-B, Peijnenburg WJGM, McGrath SP. Evaluation of an electrostatic toxicity model for predicting Ni^{2+} toxicity to barley root elongation in hydroponic cultures and in soils. *New Phytologist*. 2011;192:414–427.
- [134] Wang Y-M, Kinraide TB, Wang P, Hao X-Z, Zhou D-M. Surface electrical potentials of root cell plasma membranes: implications for ion interactions, rhizotoxicity, and uptake. *International Journal of Molecular Sciences*. 2014;15:22661–22677.
- [135] Kinraide TB. Plasma membrane surface potential (Ψ_{PM}) as a determinant of ion bioavailability: a critical analysis of new and published toxicological studies and a simplified method for the computation of plant Ψ_{PM} . *Environmental Toxicology and Chemistry*. 2006;25:3188–3198.
- [136] Tyerman SD, Skerrett M, Garrill A, Findlay GP, Leigh RA. Pathways for the permeation of Na^{+} and Cl^{-} into protoplasts derived from the cortex of wheat roots. *Journal of Experimental Botany*. 1997;48:459–480.
- [137] Silva IR, Smyth TJ, Israel DW, Raper CD, Rufty T. Magnesium ameliorates aluminum rhizotoxicity in soybean by increasing citric acid production and exudation by roots. *Plant and Cell Physiology*. 2001a;42:546–554.
- [138] Silva IR, Smyth TJ, Israel DW, Raper CD, Rufty TW. Magnesium is more efficient than calcium in alleviating aluminum rhizotoxicity in soybean and its ameliorative effect is

not explained by the Gouy-Chapman-Stern model. *Plant and Cell Physiology*. 2001b; 42:538–545.

- [139] Tipping E, Vincent CD, Lawlor AJ, Lofts S. Metal accumulation by stream bryophytes, related to chemical speciation. *Environmental Pollution*. 2008;156:936–943.
- [140] Stockdale A, Tipping E, Lofts S, Ormerod S, Clements W, Blust R. Toxicity of proton-metal mixtures in the field: linking stream macroinvertebrate species diversity to chemical speciation and bioavailability. *Aquatic Toxicology*. 2010;100:112–119.
- [141] Antunes PMC, Scornaienchi ML, Roshon HD. Copper toxicity to *Lemna minor* modelled using humic acid as a surrogate to the plant root. *Chemosphere*. 2012;88:389–394.
- [142] Iwasaki Y, Cadmus P, Clements WH. Comparison of different predictors of exposure for modeling impacts of metal mixtures on macroinvertebrates in stream microcosms. *Aquatic Toxicology*. 2013;132–133:151–156.
- [143] Tipping E, Lofts S. Metal mixture toxicity to aquatic biota in laboratory experiments: application of the WHAM- F_{TOX} model. *Aquatic Toxicology*. 2013;142–143:114–122.
- [144] Le TTY, Swartjes F, Romkens P, Groenenberg JE, Wang P, Lofts S, Hendriks AJ. Modelling metal accumulation using humic acid as a surrogate for plant roots. *Chemosphere*. 2015;124:61–69.
- [145] Fein JB, Daughney CJ, Yee N, Davis T. A chemical equilibrium model for metal adsorption onto bacterial surfaces. *Geochim Cosmochim Acta*. 1997;61:3319–3328.
- [146] Cox JS, Smith DS, Warren LA, Ferris FG. Characterizing heterogeneous bacterial surface functional groups using discrete affinity spectra for proton binding. *Environmental Science and Technology*. 1999; 33:4514–4521.
- [147] Ginn BR, Szymanowski JS, Fein JB. Metal and proton binding onto the roots of *Escue rubra*. *Chemical Geology*. 2008;253:130–135.
- [148] Meychik NR, Yermakov IP. Ion exchange properties of plant root cell walls. *Plant and Soil*. 2001;234:181–193.
- [149] Milne CJ, Kinniburgh DG, de Wit JCM, van Riemsdijk WH, Koopal LK. Analysis of proton binding by a peat humic acid using a simple electrostatic model. *Geochim Cosmochim Acta*. 1995;59:1101–1112.
- [150] Lindberg S, Landberg T, Greger M. A new method to detect cadmium uptake in protoplasts. *Planta*. 2004;214:526–532.
- [151] Postma JWM, Keltjens WG, van Riemsdijk WH. Calcium-(organo)aluminium-proton competition for adsorption to tomato root cell walls: experimental data and exchange model calculations. *Environmental Science and Technology*. 2005;39:5247–5254.

- [152] Kaulbach ES, Szymanowski JES, Fein JB. Surface complexation modeling of proton and Cd adsorption onto an algal cell wall. *Environmental Science and Technology*; 2005;39:4060–4065.
- [153] Naeem A, Woertz JR, Fein JB. Experimental measurement of proton, Cd, Pb, Sr, and Zn adsorption onto the fungal species *Saccharomyces cerevisiae*. *Environmental Science and Technology*. 2006;40:5724–5729.
- [154] Morvan C, Demarty M, Thellier M. Titration of isolated cell walls of *Lemna minor* L. *Plant Physiology*. 1979;63:1117–1122.
- [155] Dufey JE, Amory DE, Braun R. Proprietes electriques et selectivite d'echange ionique des racines. *Pedologie*. 1985;35:231–249.
- [156] Allan DL, Jarrell WM. Proton and copper adsorption to maize and soybean root cell walls. *Plant Physiology*. 1989;89:823–832.