We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

The Role of Prostate-Specific Antigen (PSA) and PSA Kinetics in the Management of Advanced Prostate Cancer

Jeremy Teoh and Ming-Kwong Yiu

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/64180>

Abstract

Prostate-specific antigen (PSA) plays an important role in the diagnosis and management of prostate cancer. The utility of PSA has been extended to a number of parameters which may guide clinical decision-making in subsequent treatment. This book chapter systematically reviewed the current evidence of PSA and PSA kinetics in the management of advanced prostate cancer. Results showed that the prognostic significance of pre-treatment PSA level is uncertain. PSA nadir predicts survival outcomes but may be confounded by the pre-treatment PSA level, and the PSA nadir may only be known after there is a PSA rise in subsequent follow-up. Time to PSA nadir has some prognostic significance but is limited by the potential immortal bias. Evidence on the use of PSA doubling time is limited and the different calculation methodologies render difficulties in generalization of such parameter. PSA progression is the best surrogate marker of survival and can be considered as the primary endpoint in future clinical trials. PSA response predicts survival but has not been shown prospectively to be a surrogate of clinical benefit. PSA and its kinetics should play an important role in the management of advanced prostate cancer and should be utilized in a more standardized manner.

Keywords: prostate cancer, prostate-specific antigen, prostate-specific antigen nadir, time to prostate-specific antigen nadir, prostatic-specific antigen doubling time, prostate-specific antigen progression, prostate-specific antigen response

1. Introduction

The first study investigating tissue-specific antibodies in the human prostate can be traced back to 1969 by Ablin et al. [1]. Nadji et al. [2] later characterized prostate-specific antigen (PSA) as a potential immunohistologic marker for prostatic neoplasms. The landmark article by Stamey et al. [3] showed that serum PSA has a much better performance than prostatic acid phosphatase in the detection of prostate cancer, and appeared to be useful in detecting residual or early recurrence of tumour, and in monitoring response to primary treatment. It has led to extensive researches in this area, and the discovery of PSA has revolutionized the management of prostate cancer, from early detection to definitive treatment and monitoring of the disease. The utility of PSA has been extended to a number of parameters that may have prognostic significance in prostate cancer and hence has gained wide interest in the past two decades.

2. Objectives

In this chapter, we systemically reviewed and appraised the current role of PSA and PSA kinetics in the management of advanced prostate cancer. We further discussed the potential benefits and controversies of the different PSA-related parameters.

3. Methods

A systematic search was conducted in the PubMed database through November 2015 using the following terms: 'prostate cancer', 'prostate-specific antigen', 'prostate-specific antigen nadir', 'time to prostate-specific antigen nadir', 'prostatic-specific antigen doubling time', 'prostate-specific antigen progression' and 'prostate-specific antigen response'. Only original full research articles published in English with full-length text available were reviewed. A manual search using the Web-based search engine Google Scholar was also performed. Reference lists of the retrieved articles were reviewed for other relevant studies.

4. Results

4.1. Pre-treatment PSA level

To a certain extent, the pre-treatment PSA level may reflect the volume of cancer cells, and hence, it is a parameter of interest in predicting disease prognosis. However, it does not reflect the sensitivity of cancer cells in response to subsequent therapy, in particular hormonal therapy in the context of metastatic disease. The prognostic significance of pre-treatment PSA level is uncertain. Some studies showed that higher pre-treatment PSA level was associated with disease progression, cancer-specific mortality and all-cause mortality [4–14], while other

studies either showed no association or failed to demonstrate statistical significance upon multivariate analyses [15–32]. The wide range of pre-treatment PSA level in metastatic disease also limited its clinical application.

4.2. PSA nadir

PSA nadir was defined as the lowest PSA level achieved after the initiation of treatment. An undetectable PSA nadir level reflects that most if not all of the prostate cancer cells are androgen-sensitive, while any detectable PSA level reflects the presence of androgen-insensitive prostate cancer cells. This was supported by a study which showed that patients who had biochemical relapse following 3 months of neoadjuvant androgen deprivation therapy and radical prostatectomy had greater PSA mRNA levels and more intense PSA immunostaining despite castrate levels of testosterone, than patients who did not relapse, yet they had similar levels of androgen receptor gene expression and protein staining [33]. The majority of the literature showed that PSA nadir is consistent in predicting disease prognosis. A higher PSA nadir level has been shown to be associated with biochemical or disease progression [4, 15, 19, 25, 31, 32, 34–37], prostate cancer-specific mortality [6, 7, 16, 22, 25, 38–40] and all-cause mortality [6, 15, 16, 24, 26, 29, 30, 41–44]. However, there is no absolute threshold level for PSA nadir being recognized by any regulatory agency, and cut-off values at 0.2 ng/mL, 1.0 ng/mL and 4.0 ng/mL have been proposed in various studies. In particular, the drop of PSA to < 4.0 ng/mL has commonly been recognized as PSA normalization, and similar to PSA nadir, PSA normalization was associated with better progression-free survival, cancer-specific survival and overall survival [35, 39, 43, 44]. However, PSA nadir may be affected by the pre-treatment PSA level, rendering difficulty in clinical application, and the PSA nadir may only be known after there is a PSA rise in subsequent follow-up.

4.3. Time to PSA nadir

Time to PSA nadir was defined as the duration needed for the PSA level to reach its nadir after the initiation of treatment. Upon hormonal therapy, one may expect the PSA level to drop to its nadir within a shorter period of time in case of hormone-sensitive prostate cancer, but the ability to have sustained continuous suppression over a longer period of time may be as important. The majority of the studies showed that a longer time to PSA nadir was associated with better outcomes including biochemical or disease progression, cancer-specific survival and overall survival [15, 16, 26, 29, 34, 36]. The other studies either showed the contrary or did not detect any associations between them [6, 22, 23, 31]. Due to the potential immortal time bias, the relationship between time to PSA nadir and survival has to be interpreted with caution [45]. For example, one must have survived 12 months in order to have a time to PSA nadir of 12 months. Hence, this immortal time bias favours a positive correlation between time to PSA nadir and survival outcomes. In order to minimize this potential bias, one study attempted to investigate the prognostic significance of time to PSA nadir using survival beyond time to PSA nadir as an alternative outcome measurement. It has been shown that a longer time to PSA nadir was associated with better survival beyond time to PSA nadir [45]. A longer time to PSA

nadir was also shown to be associated with a lower PSA velocity after progression, but whether PSA velocity after progression can be a surrogate for survival is doubtful [46].

4.4. PSA doubling time

PSA doubling time can generally be interpreted as the time needed for the PSA level to double itself. It assumes an exponential increase in serum PSA and first-order kinetics and can be calculated by natural logarithm of 2 divided by the slope of the relationship between the logarithm of PSA and time of PSA measurement [47]. However, several other calculation models have been proposed, and there is no standardization in the calculation of PSA doubling time. A shorter PSA doubling time has been shown to predict metastasis after prior radical prostatectomy [27, 28, 47], disease progression [32, 48], prostate cancer-specific mortality [6, 7, 22, 23, 25, 40, 49–51] and all-cause mortality [6, 9, 24, 27, 38, 52–54]. The utility of PSA doubling time has been widespread, yet the inconsistencies in the methodologies in calculating PSA doubling time [55] and the complicated logarithm calculations involved limited its use in clinical practice. Small deviations from the different methods of calculations may also lead to wide variations in the calculated PSA doubling time [55]. In 2008, the Prostate Cancer Clinical Trials Working Group (PCWG2) discourages the use of PSA doubling time as the primary endpoint in clinical trials because its significance is uncertain [56]. A subsequent systematic review also concluded that the evidence on PSA doubling time is limited and there is no justification for the use of PSA doubling time to guide decision-making in subsequent treatment [57].

4.5. PSA progression

PSA progression is commonly used as an endpoint in clinical trials, and it was generally thought to represent disease progression and hence reflect the survival outcomes. However, in particular for metastatic disease, multiple definitions of PSA progression have been proposed; they rendered difficulties comparing the results between different studies and limited the generalization of the utility of PSA progression. In 1999, Prostate-specific Antigen Working Group (PCWG1) made consensus recommendations for different outcome measures in clinical trials in prostate cancer [58]. PCWG1 defined PSA progression as a >50% increase from nadir and an increase of at least 5 ng/mL, or back to baseline, whichever was lowest. In 2008, PCWG2 proposed another definition for PSA progression, recognizing that early changes in PSA should not be used for clinical decision-making [56]. For those with PSA decline from baseline, PSA progression was defined as an increase in PSA by $\geq 25\%$ and ≥ 2 ng/mL above the nadir, which should be confirmed by a second value 3 or more weeks later; for those with no PSA decline, PSA progression was defined as an increase in PSA $\geq 25\%$ and ≥ 2 ng/mL after 12 weeks. Hussain et al. [44] reviewed the data from two large-scale clinical trials, namely the Southwest Oncology Group (SWOG) 9346 trial on intermittent ADT and the SWOG 9916 trial on docetaxel. It was shown that both PCWG1 and PCWG2 definitions of PSA progression predicted a 2.4-fold increase in risk of death and a more than 4-fold increase in the risk of death if PSA progression occurred in the first 7 months. This important study demonstrated that PSA progression is a significant predictor of survival in patients who have newly diagnosed

metastatic hormone-sensitive prostate cancer as well as in those with castration resistant prostate cancer treated with chemotherapy. The authors suggested that the PCWG2 definition might be more appealing as patients are identified with progression relatively earlier on. Pooling data from 9 cancer and leukaemia Group B trials [11], both PCWG1 and PCWG2 definitions of PSA progression were shown to be significant predictors of overall survival with hazard ratios of 1.44 (95% CI 1.28–1.62, $P < 0.001$) and 1.43 (95% CI 1.27–1.61, $P < 0.001$), respectively. The above evidence formed the basis of using PSA progression as the primary endpoint in various clinical trials.

4.6. PSA response

PSA response was determined by the degree of decline from its pre-treatment level. The PCWG1 [58] defined PSA response as a decline of $>50\%$ from baseline, measured twice 3–4 weeks apart. Several studies have shown that a PSA decline of $>50\%$ was associated with better cancer-specific survival [39, 59] and overall survival [17, 18, 60–62]. However, post hoc analyses in both SWOG 9916 [63] and TAX 327 [43] trials on the use of docetaxel showed that a PSA decline of $>30\%$ might be a better surrogate marker for survival than a PSA decline of $>50\%$ based on the proportion of treatment effect and the proportion of variation. A subsequent combined analysis on the SWOG 9346 and SWOG 9916 trials [44] showed that a PSA decline of $>30\%$ was associated with better overall survival. However, PCWG2 [56] advised against reporting PSA response rates in clinical trials. Concerns were raised about the strength of association between PSA decline and survival, and no criterion, be it $>50\%$ or $>30\%$ decline in PSA, has been shown prospectively to be a surrogate of clinical benefit [64]. Instead, PCWG2 recommended the use of waterfall plot to provide a broader and more sensitive display of data. On the other hand, following the discovery of AR-V7 splice variant [65], it was proposed that the lack of PSA response after initial hormonal manipulation might represent primary resistance to hormonal therapy. This is particularly important, as other non-hormonal treatment such as chemotherapy should be considered early on, based on the prediction of poor response to further hormonal manipulation.

5. Conclusions

PSA and PSA kinetics may provide additional information about the biological behaviour of prostate cancer and may aid the treatment decision in an individualized approach. One should be aware of the pros and cons of the different PSA-related parameters and should be cautious when interpreting the results from different studies. PSA and its kinetics should play an important role in the management of advanced prostate cancer, and generalization can only be achieved if definitions of the different parameters can be utilized in a more standardized manner. Among the different parameters discussed, PSA progression appeared to be the most consistent and reliable surrogate marker of survival and can serve as the primary endpoint in future clinical trials.

Author details

Jeremy Teoh^{1,2*} and Ming-Kwong Yiu^{1,2}

*Address all correspondence to: jeremyteoh@surgery.cuhk.edu.hk

1 Division of Urology, Department of Surgery, Prince of Wales Hospital, The Chinese University of Hong Kong, Hong Kong, China

2 Division of Urology, Department of Surgery, Queen Mary Hospital, The University of Hong Kong, Hong Kong, China

References

- [1] Ablin RJ, Pfeiffer L, Gonder MJ, Soanes WA. Precipitating antibody in the sera of patients treated cryosurgically for carcinoma of the prostate. *Exp Med Surg* 1969; 27:406–10.
- [2] Nadji M, Tabei SZ, Castro A, Chu TM, Murphy GP, Wang MC, et al. Prostatic-specific antigen: an immunohistologic marker for prostatic neoplasms. *Cancer* 1981; 48:1229–32.
- [3] Stamey TA, Yang N, Hay AR, McNeal JE, Freiha FS, Redwine E. Prostate-specific antigen as a serum marker for adenocarcinoma of the prostate. *N Engl J Med* 1987; 317:909–16.
- [4] Soga N, Arima K, Sugimura Y. Undetectable level of prostate specific antigen (PSA) nadir predicts PSA biochemical failure in local prostate cancer with delayed-combined androgen blockade. *Jpn J Clin Oncol* 2008; 38:617–22.
- [5] Ross RW, Xie W, Regan MM, Pomerantz M, Nakabayashi M, Daskivich TJ, et al. Efficacy of androgen deprivation therapy (ADT) in patients with advanced prostate cancer: association between Gleason score, prostate-specific antigen level, and prior ADT exposure with duration of ADT effect. *Cancer* 2008; 112:1247–53.
- [6] Chung CS, Chen MH, Cullen J, McLeod D, Carroll P, D'Amico AV. Time to prostate-specific antigen nadir after androgen suppression therapy for postoperative or postradiation PSA failure and risk of prostate cancer-specific mortality. *Urology* 2008; 71:136–40.
- [7] Stewart AJ, Scher HI, Chen MH, McLeod DG, Carroll PR, Moul JW, et al. Prostate-specific antigen nadir and cancer-specific mortality following hormonal therapy for prostate-specific antigen failure. *J Clin Oncol* 2005; 23:6556–60.

- [8] Kuriyama M, Wang MC, Lee CI, Papsidero LD, Killian CS, Inaji H, et al. Use of human prostate-specific antigen in monitoring prostate cancer. *Cancer Res* 1981; 41:3874–6.
- [9] Armstrong AJ, Garrett-Mayer ES, Yang YC, de Wit R, Tannock IF, Eisenberger M. A contemporary prognostic nomogram for men with hormone-refractory metastatic prostate cancer: a TAX327 study analysis. *Clin Cancer Res* 2007; 13:6396–403.
- [10] Berthold DR, Pond GR, Soban F, de Wit R, Eisenberger M, Tannock IF. Docetaxel plus prednisone or mitoxantrone plus prednisone for advanced prostate cancer: updated survival in the TAX 327 study. *J Clin Oncol* 2008; 26:242–5.
- [11] Halabi S, Vogelzang NJ, Ou SS, Owzar K, Archer L, Small EJ. Progression-free survival as a predictor of overall survival in men with castrate-resistant prostate cancer. *J Clin Oncol* 2009; 27:2766–71.
- [12] Smith MR, Cook R, Lee KA, Nelson JB. Disease and host characteristics as predictors of time to first bone metastasis and death in men with progressive castration-resistant nonmetastatic prostate cancer. *Cancer* 2011; 117:2077–85.
- [13] Crook JM, O'Callaghan CJ, Duncan G, Dearnaley DP, Higano CS, Horwitz EM, et al. Intermittent androgen suppression for rising PSA level after radiotherapy. *N Engl J Med* 2012; 367:895–903.
- [14] Saad F, Segal S, Eastham J. Prostate-specific antigen kinetics and outcomes in patients with bone metastases from castration-resistant prostate cancer treated with or without zoledronic acid. *Eur Urol* 2014; 65:146–53.
- [15] Huang SP, Bao BY, Wu MT, Choueiri TK, Goggins WB, Huang CY, et al. Impact of prostate-specific antigen (PSA) nadir and time to PSA nadir on disease progression in prostate cancer treated with androgen-deprivation therapy. *Prostate* 2011; 71:1189–97.
- [16] Huang SP, Bao BY, Wu MT, Choueiri TK, Goggins WB, Liu CC, et al. Significant associations of prostate-specific antigen nadir and time to prostate-specific antigen nadir with survival in prostate cancer patients treated with androgen-deprivation therapy. *Aging Male* 2012; 15:34–41.
- [17] Kelly WK, Scher HI, Mazumdar M, Vlamis V, Schwartz M, Fossa SD. Prostate-specific antigen as a measure of disease outcome in metastatic hormone-refractory prostate cancer. *J Clin Oncol* 1993; 11:607–15.
- [18] Dowling AJ, Czaykowski PM, Krahn MD, Moore MJ, Tannock IF. Prostate specific antigen response to mitoxantrone and prednisone in patients with refractory prostate cancer: prognostic factors and generalizability of a multicenter trial to clinical practice. *J Urol* 2000; 163:1481–5.
- [19] Kwak C, Jeong SJ, Park MS, Lee E, Lee SE. Prognostic significance of the nadir prostate specific antigen level after hormone therapy for prostate cancer. *J Urol* 2002; 168:995–1000.

- [20] Petrylak DP, Tangen CM, Hussain MH, Lara PN, Jr., Jones JA, Taplin ME, et al. Docetaxel and estramustine compared with mitoxantrone and prednisone for advanced refractory prostate cancer. *N Engl J Med* 2004; 351:1513–20.
- [21] Berruti A, Mosca A, Tucci M, Terrone C, Torta M, Tarabuzzi R, et al. Independent prognostic role of circulating chromogranin A in prostate cancer patients with hormone-refractory disease. *Endocr Relat Cancer* 2005; 12:109–17.
- [22] Rodrigues NA, Chen MH, Catalona WJ, Roehl KA, Richie JP, D'Amico AV. Predictors of mortality after androgen-deprivation therapy in patients with rapidly rising prostate-specific antigen levels after local therapy for prostate cancer. *Cancer* 2006; 107:514–20.
- [23] D'Amico AV, McLeod DG, Carroll PR, Cullen J, Chen MH. Time to an undetectable prostate-specific antigen (PSA) after androgen suppression therapy for postoperative or postradiation PSA recurrence and prostate cancer-specific mortality. *Cancer* 2007; 109:1290–5.
- [24] Daskivich TJ, Regan MM, Oh WK. Distinct prognostic role of prostate-specific antigen doubling time and velocity at emergence of androgen independence in patients treated with chemotherapy. *Urology* 2007; 70:527–31.
- [25] Scholz M, Lam R, Strum S, Jennrich R, Johnson H, Trilling T. Prostate-cancer-specific survival and clinical progression-free survival in men with prostate cancer treated intermittently with testosterone-inactivating pharmaceuticals. *Urology* 2007; 70:506–10.
- [26] Choueiri TK, Xie W, D'Amico AV, Ross RW, Hu JC, Pomerantz M, et al. Time to prostate-specific antigen nadir independently predicts overall survival in patients who have metastatic hormone-sensitive prostate cancer treated with androgen-deprivation therapy. *Cancer* 2009; 115:981–7.
- [27] Antonarakis ES, Chen Y, Elsamanoudi SI, Brassell SA, Da Rocha MV, Eisenberger MA, et al. Long-term overall survival and metastasis-free survival for men with prostate-specific antigen-recurrent prostate cancer after prostatectomy: analysis of the Center for Prostate Disease Research National Database. *BJU Int* 2011; 108:378–85.
- [28] Antonarakis ES, Feng Z, Trock BJ, Humphreys EB, Carducci MA, Partin AW, et al. The natural history of metastatic progression in men with prostate-specific antigen recurrence after radical prostatectomy: long-term follow-up. *BJU Int* 2012; 109:32–9.
- [29] Sasaki T, Onishi T, Hoshina A. Nadir PSA level and time to PSA nadir following primary androgen deprivation therapy are the early survival predictors for prostate cancer patients with bone metastasis. *Prostate Cancer Prostatic Dis* 2011; 14:248–52.
- [30] Miyamoto S, Ito K, Miyakubo M, Suzuki R, Yamamoto T, Suzuki K, et al. Impact of pretreatment factors, biopsy Gleason grade volume indices and post-treatment nadir

PSA on overall survival in patients with metastatic prostate cancer treated with step-up hormonal therapy. *Prostate Cancer Prostatic Dis* 2012; 15:75–86.

- [31] Benaim EA, Pace CM, Lam PM, Roehrborn CG. Nadir prostate-specific antigen as a predictor of progression to androgen-independent prostate cancer. *Urology* 2002; 59:73–8.
- [32] Keizman D, Huang P, Antonarakis ES, Sinibaldi V, Carducci MA, Denmeade S, et al. The change of PSA doubling time and its association with disease progression in patients with biochemically relapsed prostate cancer treated with intermittent androgen deprivation. *Prostate* 2011; 71:1608–15.
- [33] Ryan CJ, Smith A, Lal P, Satagopan J, Reuter V, Scardino P, et al. Persistent prostate-specific antigen expression after neoadjuvant androgen depletion: an early predictor of relapse or incomplete androgen suppression. *Urology* 2006; 68:834–9.
- [34] Hori S, Jabbar T, Kachroo N, Vasconcelos JC, Robson CN, Gnanapragasam VJ. Outcomes and predictive factors for biochemical relapse following primary androgen deprivation therapy in men with bone scan negative prostate cancer. *J Cancer Res Clin Oncol* 2011; 137:235–41.
- [35] Oosterlinck W, Mattelaer J, Casselman J, Van Velthoven R, Derde MP, Kaufman L. PSA evolution: a prognostic factor during treatment of advanced prostatic carcinoma with total androgen blockade. Data from a Belgian multicentric study of 546 patients. *Acta Urol Belg* 1997; 65:63–71.
- [36] Morote J, Trilla E, Esquena S, Abascal JM, Reventos J. Nadir prostate-specific antigen best predicts the progression to androgen-independent prostate cancer. *Int J Cancer* 2004; 108:877–81.
- [37] Park SC, Rim JS, Choi HY, Kim CS, Hong SJ, Kim WJ, et al. Failing to achieve a nadir prostate-specific antigen after combined androgen blockade: predictive factors. *Int J Urol* 2009; 16:670–5.
- [38] D'Amico AV, Chen MH, de Castro M, Loffredo M, Lamb DS, Steigler A, et al. Surrogate endpoints for prostate cancer-specific mortality after radiotherapy and androgen suppression therapy in men with localised or locally advanced prostate cancer: an analysis of two randomised trials. *Lancet Oncol* 2012; 13:189–95.
- [39] Suzuki H, Okihara K, Miyake H, Fujisawa M, Miyoshi S, Matsumoto T, et al. Alternative nonsteroidal antiandrogen therapy for advanced prostate cancer that relapsed after initial maximum androgen blockade. *J Urol* 2008; 180:921–7.
- [40] Park YH, Hwang IS, Jeong CW, Kim HH, Lee SE, Kwak C. Prostate specific antigen half-time and prostate specific antigen doubling time as predictors of response to androgen deprivation therapy for metastatic prostate cancer. *J Urol* 2009; 181:2520–4; discussion 5.
- [41] Teoh JY, Tsu JH, Yuen SK, Chan SY, Chiu PK, Wong KW, et al. Survival outcomes of Chinese metastatic prostate cancer patients following primary androgen deprivation

therapy in relation to prostate-specific antigen nadir level. *Asia Pac J Clin Oncol* 2014. (Epub ahead of print).

- [42] Hussain M, Tangen CM, Higano C, Schelhammer PF, Faulkner J, Crawford ED, et al. Absolute prostate-specific antigen value after androgen deprivation is a strong independent predictor of survival in new metastatic prostate cancer: data from Southwest Oncology Group Trial 9346 (INT-0162). *J Clin Oncol* 2006; 24:3984–90.
- [43] Armstrong AJ, Garrett-Mayer E, Ou Yang YC, Carducci MA, Tannock I, de Wit R, et al. Prostate-specific antigen and pain surrogacy analysis in metastatic hormone-refractory prostate cancer. *J Clin Oncol* 2007; 25:3965–70.
- [44] Hussain M, Goldman B, Tangen C, Higano CS, Petrylak DP, Wilding G, et al. Prostate-specific antigen progression predicts overall survival in patients with metastatic prostate cancer: data from Southwest Oncology Group Trials 9346 (Intergroup Study 0162) and 9916. *J Clin Oncol* 2009; 27:2450–6.
- [45] Teoh JY, Tsu JH, Yuen SK, Chan SY, Chiu PK, Lee WM, et al. Prognostic significance of time to prostate-specific antigen (PSA) nadir and its relationship to survival beyond time to PSA nadir for prostate cancer patients with bone metastases after primary androgen deprivation therapy. *Ann Surg Oncol* 2015; 22:1385–91.
- [46] Teoh JY, Tsu JH, Yuen SK, Chiu PK, Chan SY, Wong KW, et al. Association of time to prostate-specific antigen nadir and logarithm of prostate-specific antigen velocity after progression in metastatic prostate cancer with prior primary androgen deprivation therapy. *Asian J Androl* 2015. (Epub ahead of print).
- [47] Pound CR, Partin AW, Eisenberger MA, Chan DW, Pearson JD, Walsh PC. Natural history of progression after PSA elevation following radical prostatectomy. *JAMA* 1999; 281:1591–7.
- [48] Keizman D, Huang P, Carducci MA, Eisenberger MA. Contemporary experience with ketoconazole in patients with metastatic castration-resistant prostate cancer: clinical factors associated with PSA response and disease progression. *Prostate* 2012; 72:461–7.
- [49] D'Amico AV, Halabi S, Tempany C, Titelbaum D, Philips GK, Loffredo M, et al. Tumor volume changes on 1.5 tesla endorectal MRI during neoadjuvant androgen suppression therapy for higher-risk prostate cancer and recurrence in men treated using radiation therapy results of the phase II CALGB 9682 study. *Int J Radiat Oncol Biol Phys* 2008; 71:9–15.
- [50] Freedland SJ, Humphreys EB, Mangold LA, Eisenberger M, Dorey FJ, Walsh PC, et al. Risk of prostate cancer-specific mortality following biochemical recurrence after radical prostatectomy. *JAMA* 2005; 294:433–9.
- [51] Svatek RS, Shulman M, Choudhary PK, Benaim E. Critical analysis of prostate-specific antigen doubling time calculation methodology. *Cancer* 2006; 106:1047–53.

- [52] Semeniuk RC, Venner PM, North S. Prostate-specific antigen doubling time is associated with survival in men with hormone-refractory prostate cancer. *Urology* 2006; 68:565–9.
- [53] Oudard S, Banu E, Scotte F, Banu A, Medioni J, Beuzeboc P, et al. Prostate-specific antigen doubling time before onset of chemotherapy as a predictor of survival for hormone-refractory prostate cancer patients. *Ann Oncol* 2007; 18:1828–33.
- [54] Oudard S, Banu E, Medioni J, Scotte F, Banu A, Levy E, et al. What is the real impact of bone pain on survival in patients with metastatic hormone-refractory prostate cancer treated with docetaxel? *BJU Int* 2009; 103:1641–6.
- [55] Daskivich TJ, Regan MM, Oh WK. Prostate specific antigen doubling time calculation: not as easy as 1, 2, 4. *J Urol* 2006; 176:1927–37.
- [56] Scher HI, Halabi S, Tannock I, Morris M, Sternberg CN, Carducci MA, et al. Design and end points of clinical trials for patients with progressive prostate cancer and castrate levels of testosterone: recommendations of the Prostate Cancer Clinical Trials Working Group. *J Clin Oncol* 2008; 26:1148–59.
- [57] Vickers AJ, Savage C, O'Brien MF, Lilja H. Systematic review of pretreatment prostate-specific antigen velocity and doubling time as predictors for prostate cancer. *J Clin Oncol* 2009; 27:398–403.
- [58] Bubley GJ, Carducci M, Dahut W, Dawson N, Daliani D, Eisenberger M, et al. Eligibility and response guidelines for phase II clinical trials in androgen-independent prostate cancer: recommendations from the Prostate-Specific Antigen Working Group. *J Clin Oncol* 1999; 17:3461–7.
- [59] Okihara K, Ukimura O, Kanemitsu N, Mizutani Y, Kawauchi A, Miki T, et al. Clinical efficacy of alternative antiandrogen therapy in Japanese men with relapsed prostate cancer after first-line hormonal therapy. *Int J Urol* 2007; 14:128–32.
- [60] Smith DC, Dunn RL, Strawderman MS, Pienta KJ. Change in serum prostate-specific antigen as a marker of response to cytotoxic therapy for hormone-refractory prostate cancer. *J Clin Oncol* 1998; 16:1835–43.
- [61] Small EJ, McMillan A, Meyer M, Chen L, Slichenmyer WJ, Lenehan PF, et al. Serum prostate-specific antigen decline as a marker of clinical outcome in hormone-refractory prostate cancer patients: association with progression-free survival, pain end points, and survival. *J Clin Oncol* 2001; 19:1304–11.
- [62] Berthold DR, Pond GR, Roessner M, de Wit R, Eisenberger M, Tannock AI, et al. Treatment of hormone-refractory prostate cancer with docetaxel or mitoxantrone: relationships between prostate-specific antigen, pain, and quality of life response and survival in the TAX-327 study. *Clin Cancer Res* 2008; 14:2763–7.

- [63] Petrylak DP, Ankerst DP, Jiang CS, Tangen CM, Hussain MH, Lara PN, Jr., et al. Evaluation of prostate-specific antigen declines for surrogacy in patients treated on SWOG 99-16. *J Natl Cancer Inst* 2006; 98:516–21.
- [64] Scher HI, Morris MJ, Basch E, Heller G. End points and outcomes in castration-resistant prostate cancer: from clinical trials to clinical practice. *J Clin Oncol* 2011; 29:3695–704.
- [65] Antonarakis ES, Lu C, Wang H, Luber B, Nakazawa M, Roeser JC, et al. AR-V7 and resistance to enzalutamide and abiraterone in prostate cancer. *N Engl J Med* 2014; 371:1028–38.