We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Non-Bone Effects of Vitamin D in Children, Adolescents, and Young Adults

Mohsin Ali Cheema and Khalid Parvez Lone

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/65079>

Abstract

Vitamin D, also known as “sunshine vitamin”, has long been established as an essential component for the maintenance of adequate bone health. Large number of studies are available which demonstrate the various biochemical pathways of vitamin D in bone physiology and its important role in musculoskeletal health. In last five decades, data regarding the non-bone effects of vitamin D have started to emerge, and now many important non-bone physiological processes are explained by the biochemical pathways and functions of vitamin D. However, majority of the data regarding extra-skeletal effects of vitamin D are available regarding adult population. In this chapter, we try to focus on the role of vitamin D in aging and various diseases which are frequently seen in children, adolescents, and young adults such as cancer, type 1 diabetes mellitus, allergies, asthma, and various autoimmune diseases.

Keywords: vitamin D, autoimmune disease, aging, diabetes, cancer, children

1. Introduction

Vitamin D, also known as “sunshine vitamin”, is considered essential for maintenance of bone health along with many other key roles in metabolic processes of the body. About 10,000–20,000 IU of vitamin D₃ is produced in our skin from the sun's ultraviolet light (UVB) after full body exposure for 15 min [1]. Complexion or skin phototype, use of sunblock, smog, cloud cover, latitude, time of day (10:00 am to 02:00 pm), and season are some of the factors that affect our body's ability to produce vitamin D [2]. Latitudes from 0 to 35° north or south allow yearlong production of vitamin D in our body, but as latitude increases the amount of vitamin

D production will decrease [3]. The latitude of Pakistan is 32.0162°N which is suitable for yearlong production of vitamin D.

Recommended daily intake of vitamin D in the first year of life is 400 IU and 600 IU for everyone above 1 year of age [4]. Vitamin D is essential for proper bone growth and bone remodeling by the action of osteoblasts and osteoclasts [5]. Vitamin D insufficiency or deficiency can lead to osteoporosis by decreasing calcium absorption from intestine and kidney [6]. Harinarayan et al. have showed that normal homeostasis of bone has been seriously affected by decreased intake of calcium and vitamin D in diet [7]. In children, inadequate mineralization of the growing skeleton leading to rickets is an established fact [8]. Välimäki et al. discussed that vitamin D deficiency is common in young men, especially in winter, and this deficiency may have detrimental effects on the acquisition of maximal peak bone mass [9]. Gait disturbances and myopathy have also been associated with severe vitamin D deficiency [10]. Discovery of the vitamin D receptor (VDR) expressed in the cell nuclei of muscle cells and its association with muscle cell contractility has been documented by Bischoff-Ferrari et al [11]. Meta-analysis done by Papadimitropoulos et al. concluded that adequate levels of vitamin D decrease the rate of vertebral and non-vertebral fractures [12].

1.1. Prevalence of vitamin D deficiency

Studies from around the world have shown pandemic of vitamin D deficiency or insufficiency [13]. In last decade, various studies from Pakistan have found critically low levels of vitamin D in diverse groups of population. Work done by Siddiqui and Rai [14] in Hazara division of Pakistan showed that nutritional rickets resulting from vitamin D deficient diet is a predisposing factor for different childhood illnesses such as pneumonia, diarrhea, and delayed motor milestones. They attributed lack of sun exposure, malnutrition, and various antenatal factors as important causes for vitamin D deficiency. Similarly, Anwar et al. [15] found that 99.5% of women and 97.3% of neonates from urban population sample and 89% of women and 82% of neonates from rural population sample had below 50 nmol/l levels of vitamin D. Jamal et al. [16] and Qamar et al. [17] have also found insufficient/deficient vitamin D levels in more than 90% children included in their studies. A study done in our laboratory including 5- to 11-year-old children with intellectual disability also showed extremely low levels of vitamin D (12.08 ± 9.06 ng/ml) in 85% of study sample. These numbers are consistent with another population-based study done in Karachi by Iqbal et al. [18] who found that vitamin D was deficient in 73.7% and insufficient in 13.8% study subjects.

In last two decades, lot of studies came out from around the world reporting lower or deficient levels of vitamin D; however, few questions remained ambiguous: How much vitamin D is needed to achieve desirable health outcomes and how much of vitamin D is too much? In 2011, a new public health report on dietary intake requirements for calcium and vitamin D authored by the committee from the Institute of Medicine (IOM) gave their recommendations regarding above queries. The committee concluded that serum 25OHD levels of 16 ng/ml (40 nmol/l) cover the requirements of approximately half of the population, and levels of 20 ng/ml (50 nmol/l) cover the requirements of at least 97.5% of the population. The IOM committee also highlighted the fact that data for the upper cutoff limit were scarce and long-term effects of

chronically high concentrations of vitamin D, and a margin of safety for public health recommendations was prudent. Thus, serum 25OHD levels above 50 ng/ml (125 nmol/l) should raise concerns among clinicians about potential adverse effects [4].

1.2. Vitamin D status and skin complexion

Complexion or skin phototype is another important factor that determines amount of vitamin D production in our body. Individuals with darker complexion have higher chances of developing vitamin D deficiency than fair-skinned people [19]. The deficiency in circulating levels of 25-hydroxyvitamin D in individuals with darker skin complexion is mediated largely by melanin which protects the skin from ultraviolet light rays. By blocking the sun's ultra violet rays, melanin greatly reduces the skin's ability to convert 7-dehydrocholesterol to cholecalciferol, the precursor of 25(OH)D, in the skin [20]. In 2011, Signorello et al. investigated the association between 94 single nucleotide polymorphisms (SNPs) in five vitamin D pathway genes (*GC*, *VDR*, *CYP2R1*, *CYP24A1*, *CYP27B1*) and serum 25-hydroxyvitamin D (25(OH)D) levels among 379 African American and 379 Caucasian participants, and they found that common variation in vitamin D pathway genes predicts circulating 25-hydroxyvitamin D levels among African Americans having darker complexion [21].

Vitamin D supplementation is considered as an important factor in improving the poor bone health of people with both fair and dark skin complexions. Studies have shown that people with both dark and white skin color appear to have similar capacities to synthesize vitamin D in the skin, but vitamin D synthesis is less efficient among blacks at usual levels of sun exposure as compared to white people [22]. Recently, a study done by Gallagher et al. demonstrated that despite having lower levels of vitamin D at baseline, the increase in serum vitamin D after supplementation in African American women was similar to that seen in the Caucasian women. Furthermore, 97.5% of both dark- and white-colored women on 800 IU/daily for 12 months reached a level of 20 ng/ml. They also concluded that because absorption and metabolism of oral vitamin D are similar in both dark and fair complexioned people, lower levels of serum 25OHD in African Americans must be due to lower production of vitamin D in skin [23].

1.3. Is Vitamin D synthesis gender specific?

Data regarding the association of gender with vitamin D synthesis are inconsistent and scarce. There are number of conflicting hypotheses which indicate that differences in the amount of adipose tissue in males and females and in their skins may lead to the variation in vitamin D synthesis. Number of studies has demonstrated decreased bioavailability of fat soluble vitamin D3 from cutaneous and dietary sources because of its deposition in body fat [24–27]. This may point toward decreased vitamin D levels in women because of their excessive adiposity as compared to men. On the other hand, it has also been hypothesized that the lighter color of female skin permits synthesis of relatively higher amounts of vitamin D3 during pregnancy and lactation [28]. On the contrary, data from a recent study in mice have demonstrated that androgens decrease the synthesis of vitamin D induced by ultraviolet rays in the skin of a male mice by an enzymatic mechanism [29].

1.4. Physiology of non-bone effects of vitamin D

Initially, it was thought that vitamin D has a restricted role in calcium homeostasis, but it is only now that the pleiotropic actions of vitamin D with their clinical significance are becoming apparent. Recently, 2776 genomic positions occupied by the vitamin D receptor (VDR) and 229 genes showing significant changes in expression in response to vitamin D have been identified [30]. VDR has been identified in wide variety of tissues other than small intestine, kidneys, and bone such as brain, heart, stomach, pancreas, activated T and B lymphocytes, skin and gonads [31]. One hydroxylase activity has also been identified in cultured cells from skin, colon, prostate, breast, lung, and brain [1, 32, 33]. Addition of vitamin D in our diet in the form of supplementation or food fortification can reduce the risk of asthma, influenza, respiratory tract infections, autoimmune diseases, diabetes, and cancer and improve bone and overall health [2]. This evidence points to the diverse functions of vitamin D and its implicated role in various diseases related to wide distribution of vitamin D receptors in many tissues.

In the following sections of this chapter, recent review of literature regarding physiological role of vitamin D in various non-bone effects is given.

2. Vitamin D and aging

It has been long known that aging decreases the capacity of human skin to produce vitamin D₃ (7-dehydrocholesterol). This is evident from the classic study done by McLaughlin and Holick in 1985 in which they exposed the skin samples from various age groups to ultraviolet rays and compared the amount of previtamin D₃ produced in the skin samples from 8- to 18-year-old subjects with the amount produced in the skin samples from 77- to 82-year-old subjects. The results revealed that aging can decrease the capacity of the skin to produce previtamin D₃ by greater than twofold [34].

In humans, aging represents the accumulation of changes in a human being over time, encompassing physical, psychological, and social change. Telomeres, the DNA-protein structures located at the ends of chromosomes, have been proposed to act as a biomarker of aging [35]. A cross-sectional analysis of the Nurses' Health Study (NHS) data done by Liu et al. [36] demonstrated the association between vitamin D and telomere length in peripheral blood leukocytes by using plasma biomarkers of both 25(OH)D and 1,25(OH)₂D. They found that higher plasma 25(OH)D levels are associated with longer telomeres, and this association may be modified by calcium intake. The difference in telomere length between those with high (vitamin D sufficient) and low (vitamin D insufficient) levels of vitamin D corresponded to 5 years of aging. The shortening of telomeres is thought to be caused by decreasing inflammatory mediators and cell proliferation [37]. Changes in the expression of VDR leading to vitamin D resistance with aging have also been elaborated in recent literature. Several studies have reported association of aging with decrease in the expression of VDR in bone, intestine, and muscle tissues [38–40]. Various factors that decrease with aging have been identified to influence VDR such as estrogen, growth hormones, and vitamin D itself [11, 41]. On the other hand, the expression of TNF alpha which increases with aging has been shown to downregu-

late the expression of VDR [42, 43]. An animal study has also demonstrated a decrease in number of vitamin D receptors in addition to decreased binding of active metabolite of vitamin D with VDR in aged rat intestinal subcellular fractions [44]. Association of premature aging phenotype with too high and too low vitamin D levels has also been demonstrated in mice [45]. A recent study found that individuals in the highest quartile of serum vitamin D had the longest lifespan compared to those in the lowest quartile [46]. Above evidence and already established associations of several common aging-associated diseases such as osteoporosis, hypertension, and diabetes with vitamin D deficiency point us to consider vitamin D levels as a biomarker for aging.

3. Cancer in children

Although cancer in children is rare, it is the leading cause of death by disease past infancy among children in the United States. The most common types of cancer diagnosed in children and adolescents are leukemia, brain and other central nervous system tumors, lymphoma, rhabdomyosarcoma, neuroblastoma, Wilms tumor, bone cancer, and gonadal (testicular and ovarian) germ cell tumors [47]. Pakistan is a developing country with a population of about 180 million people out of which 39% are under 15 years of age. Although malnutrition and communicable diseases are still the major killers, cancer is becoming an important cause of morbidity and mortality in children. Cancer among children is fortunately less common than in adult accounting for 3–5% of all cancers. Based on Karachi Cancer Registry, it is estimated that about 7000–7500 children get cancer every year in Pakistan [48]. A limited number of small studies have examined the vitamin D status of pediatric oncology patients, and the results indicate an increased prevalence of hypovitaminosis D [49, 50].

3.1. Role of vitamin D in cancer

Specific vitamin D receptors found in nearly all tissues produce 1,25(OH)₂D in the presence of 25(OH)D. This 1,25(OH)₂D is recognized to cause activation of VDR resulting in heterodimerization with the retinoid X receptor and binding to cognate vitamin D response elements (VDREs) in target genes involved in cellular differentiation, cell growth, apoptosis, inflammation, and immune modulation [51, 52]. Formation of 1 alpha hydroxylase enzyme in cancer cells has also been demonstrated [53, 54]. In last decade of twentieth century, number of studies came out that highlighted the association of decreased vitamin D levels in individuals living at high latitudes and increased risk of dying of colon, breast, prostate, and ovarian cancer [55–57].

Polymorphism of VDR gene has been associated with high risk of cancer. Women with mutations of VDR gene have higher risk of breast cancer [58]. Results of a hospital-based case-control study conducted by our department demonstrated that the *BsmI* polymorphism in the VDR gene may be associated with an increased breast cancer risk in Pakistani women negative for *BRCA1/2* germline mutations [59].

Vitamin D-liganded VDR displays antiproliferative activities in many tumor types, as do activated members of the p53 family, through the induction of cell cycle arrest, senescence, differentiation, and apoptosis [60]. Several metabolites of vitamin D which do not cause hypercalcemia show antitumor activity in a subset of cancer patients with high VDR expression and are associated with good prognosis [61]. Some transcription factors repress VDR gene expression in human colon cancer cells. In human colon cancers, elevated expression of transcription factors correlates with downregulation of VDR [62]. The malignant cells are capable of annulling the antiproliferative activity of vitamin D by increasing the expression of transcriptional factors.

Several preclinical trials have demonstrated the antiproliferative effects of active form of vitamin D in various tumor types. The mechanisms by which vitamin D can exert antitumor effects may include inhibition of tumor angiogenesis, induction of apoptosis, regulation of different signaling pathways in tumor cells, and arrest of tumor cells in G0/G1 phase of cell cycle. Preclinical data indicate that maximal antitumor effects are seen with pharmacological doses of $1\alpha,25(\text{OH})_2\text{D}_3$ and can be safely achieved in animals using a high-dose, intermittent schedule of administration [60]. Some clinical trial data indicate that $1\alpha,25(\text{OH})_2\text{D}_3$ is well tolerated in cancer patients within a proper dosing schedule. Data support the hypothesis that vitamin D compounds may have an important role in cancer therapy and prevention, and merit further investigation [63].

Despite the presence of wealthy data regarding vitamin D and cancer, only a handful of studies have assessed vitamin D status in pediatric patients with malignancy, and all included relatively small numbers of patients [50]. This may point to the need of continuing elaborative research to identify the role of vitamin D in cancer, especially in populations with younger age groups.

4. Diabetes and vitamin D

Vitamin D deficiency and diabetes have one major trait in common: both are pandemic. The International Diabetes Federation estimated the number of people with diabetes worldwide to be nearly 415 million out of which three-quarters (75%) live in low- and middle-income countries. In Pakistan, prevalence of diabetes is 8.1%, whereas number of new cases of type 1 diabetes per 100,000 children per year is 0.5 [64]. Studies from different populations throughout the globe have reported high rates of vitamin D deficiency in children with type 1 diabetes [65–70]. Also several studies suggest that vitamin D deficiency correlates with the severity and frequency of type 1 diabetes in children as well as that vitamin D supplementation may reduce the risk of developing type 1 diabetes in younger age groups [71–73]. An analytical cross-sectional study conducted in our department demonstrated that offsprings of type 2 diabetics were severely deficient in vitamin D and its levels were inversely correlated with most of the components of metabolic syndrome [74]. Researchers have also explored the geographical variation in childhood diabetes. In study done by Mohr et al. in 2008, incidence rates of type 1 diabetes in children aged <14 years during 1990–1994 in 51 regions worldwide were assessed

by using multiple regression. This study found an association between low ultraviolet B irradiance and high incidence rates of type 1 childhood diabetes after controlling for per capita health expenditure. Incidence rates of type 1 diabetes approached zero in regions worldwide with high UVB irradiance, adding new support to the concept of a role of vitamin D in reducing the risk of the disease [75]. Similarly, data from sixth edition of diabetes atlas showed that incidence of type 1 diabetes is higher in countries that are located furthest from the equator as compared to countries located closest to equator. Staples et al. in their another study, which specifically examined the latitude gradient within Australia's territories, found childhood diabetes to be positively related to latitude. There was a strong threefold increase in prevalence of type 1 diabetes moving from the most northern territory to the most southern territory of Australia [76]. The above discussion positively points toward the association of incidence of childhood type 1 diabetes with latitudinal location.

Type 1 diabetes mellitus results from a cellular-mediated autoimmune destruction of the beta cells of the pancreas [77]. Studies on mice suggest that vitamin D inhibits IL-12 production and pancreatic infiltration of T helper cells along with the increase in number of CD4 positive and d CD25 positive regulatory T cells in pancreatic lymph nodes which may help in limiting the immunological progression and preventing the clinical onset of type 1 diabetes [78, 79]. There is evidence that vitamin D is important in the prevention of islet cell death and might be useful in improving the survival of islet cell grafts [80]. The presence of VDR and vitamin D-dependant calcium binding protein on beta cells of pancreas has also been reported. The effects of vitamin D on beta cells may be by its regulation of extracellular calcium and calcium flux through the beta cell or through calcium-independent pathways. Vitamin D deficiency may also impair insulin secretion through its associated increase in parathormone levels. It may reduce insulin resistance by its immunomodulatory and antiinflammatory effects [61]. Above evidence suggests that vitamin D may play a role in the prevention and treatment of type 1 diabetes mellitus in children; however, definitive and conclusive evidence regarding the potential role of vitamin D in alleviating the increasing menace of diabetes needs to be further studied through its action on systemic inflammation, insulin secretion, and resistance.

5. Allergies and asthma in children

Asthma and allergies are common chronic diseases in developed world [81–83]. CDC reports from national surveys in United States show the prevalence of asthma in children as 8.3% in 2013. The highest rates were observed among children aged 5–14 years, boys of less than 18 years, and people below 100% of poverty level [84]. Epidemiology of allergies and asthma is not well documented in Pakistan. A survey conducted in the city of Karachi among school children of age 3–16 years reported the prevalence of asthma at 15.8% and that of allergic rhinitis at 28.50% [85]. Another two-stage community-based representative cross-sectional survey conducted in Karachi from March 2012 to April 2013 reported overall prevalence of asthma among study participants as 10.2% [86].

5.1. Role of vitamin D in allergies and asthma in children

Many potential reasons have been reported in literature that could account for the pattern of increased burden of allergic diseases and asthma in developing as well as developed world. In last couple of decades, several studies have proposed pandemic of vitamin D deficiency as an important candidate that could explain a significant proportion of increased prevalence of allergic disease and asthma [87]. Potential mechanisms of how vitamin D can affect the risk of developing asthma and allergies include genetic pathways, role of vitamin D in prevention of bacterial infections, immune system effects, and effects on lung development and functions [88]. Significant associations between polymorphisms in the VDR gene and asthma have been reported by number of studies [89–91]. It has been proposed that asthmatics with bacterial and viral infections are at higher risk of more severe symptoms [92, 93]. Vitamin D induces the production of the antimicrobial polypeptide cathelicidin which has both antibacterial and antiviral effects [94, 95], and supplementation with vitamin D in asthmatics demonstrated decreased incidence of cold or influenza symptoms [96]. Studies done in mice have shown that vitamin D deficiency *in utero* leads to decreased lung volumes by affecting lung development [97]. Similarly, vitamin D also appears to affect *in utero* immune system development that begins to exert its effects in early life. Work done by Chi et al. [98] showed that cord blood vitamin D levels were inversely associated with the proportions of CD25(+), CD25(Bright), and CD25(+) FoxP3 cells to total CD4(+) T cells. While the clinical consequence of this inverse association remains unclear, it supports the notion that *in utero* vitamin D levels affect immune development and may influence immune regulation early in life. The number of studies has shown that maternal supplementation with vitamin D during pregnancy may eventually decrease the risk of allergies and wheezing in their children [99–101]. Other studies have investigated the association between vitamin D levels and asthma and allergies in the post-partum period. Several case-control studies from different parts of the world have found greater prevalence of vitamin D deficiency among asthmatic children than in controls [102–104]. In a recent work done in our department, an inverse relation of vitamin D levels with severity of asthma has been found (Lone Unpublished data).

Although more studies have shown a beneficial effect of vitamin D on asthma and allergies than studies showing negative results, definitive clinical trials are lacking and the optimal dose and level of vitamin D for decreasing the burden of asthma and allergies in children remain unknown.

6. Autoimmune diseases in children and vitamin D

There are at least 80 recognized human autoimmune diseases with new diseases frequently added to the list [105]. Autoimmune diseases are generally rare in children; however, when they occur, they can be challenging to diagnose and difficult to treat because most of the autoimmune diseases that are common in children have not cured yet. Besides type 1 diabetes mellitus, other autoimmune diseases which commonly occur in children include celiac disease,

lupus (SLE), juvenile dermatomyositis, scleroderma, juvenile idiopathic arthritis (JIA), and multiple sclerosis (MS) [106].

The effects of vitamin D in the immune system translate into an enhancement of innate immunity associated with a multifaceted regulation of acquired immunity [107]. Vitamin D supplementation is considered as an appealing therapy in pediatric autoimmune diseases. The diverse effects of vitamin D may minimize disease-related comorbidities resulting from bone weakening and infections in addition to attenuation of immune hyperactivation that is characteristic of pediatric SLE [108]. The SLE Disease Activity Index scores were found to be significantly higher in children who had 25(OH)D level less than 20 ng/ml [109]. Work done by Robinson et al. [110] demonstrated that low serum 25(OH)D level in children with SLE is associated with proteinuria and urinary vitamin D binding protein. Animal models have also shown that vitamin D and calcium supplementation can inhibit lupus activity [111, 112]. An incidental finding of an association between vitamin D deficiency and SLE nephritis has also been reported [113], but studies evaluating this relationship are lacking. Juvenile dermatomyositis (JDM) is a photosensitive rheumatic disease, treated commonly with sun avoidance, corticosteroids, methotrexate, and hydroxychloroquine. JDM shares many similarities with pediatric SLE in photosensitivity and methods of treatment, but it is not associated with proteinuria, making this an ideal comparison group for SLE. Robinson et al. [110] also reported that vitamin D deficiency is associated with disease activity in children with JDM.

Multiple sclerosis is an autoimmune disease of the central nervous system characterized by inadequate recognition of autoepitopes in myelinated nerve fibers by cells of the acquired immune system, generating an inflammatory immune response mediated by lymphocytes and macrophages, resulting in localized areas of inflammation and demyelination [114]. Some studies have also demonstrated the association of vitamin D deficiency and MS and its role not only in the reduction of relapse rates, but also in the prevention of its development [115, 116]. In MS, allelic variation in the MHC class II region exerts the single strongest effect on genetic risk. Environmental factors act at a population level. Sunlight or vitamin D is a key environmental factor in etiology and might interact with inherited factors in the MHC class II region [61]. A single MHC vitamin D response element (VDRE) gene has been identified as *HLA-DRB1*15* haplotypes. In a subgroup of individuals genetically predisposed to multiple sclerosis, deficiency of vitamin D may cause non-activation of histocompatibility genes necessary for differentiating between self and foreign proteins [117].

Despite the compelling evidence of low levels of vitamin D and its association with various autoimmune diseases, recently studies are coming out that challenge the assumption that serum levels of 25OH vitamin D are a sensitive marker of the autoimmune disease state [118]. Now many commentators are also advocating clinicians to stop costly measurements of 25OH vitamin D in asymptomatic patients [119–121]. The consideration of vitamin D as a marker for autoimmune diseases requires further sensitive evidence from randomized controlled clinical trials. The results then may elaborate the status of vitamin D as a marker for autoimmune diseases in future. Although more epidemiologic studies are needed to better understand the theory of vitamin D deficiency and its association with various auto immune diseases in

children, the compelling data pointing to a role for vitamin D in immune regulation suggest that special attention should be paid to these at-risk populations.

6.1. Role of sunlight in autoimmune diseases

The ultraviolet radiation in sunlight can induce the onset of, or exacerbate, the symptoms of certain autoimmune diseases. Work done by Fraser et al. suggested that ultraviolet light exposure triggers production of reactive oxygen species as normal by-products. If the cell does not quickly eliminate the reactive oxygen species, however, the buildup can cause cellular and DNA damage. The *GSTM1* gene normally codes for an enzyme, glutathione S-transferase, which rids the body of reactive oxygen species. Individuals who have the *GSTM1* null genotype and are missing the enzyme may, therefore, be at an increased risk of DNA damage and can induce the onset of lupus [122]. Work done by Love et al. to study the distribution of myositis phenotypes and ultraviolet radiation exposure in the United States showed that ultraviolet radiation may modulate the clinical and immunological expression of autoimmune disease in women [123]. The above evidence points toward the stimulation and exacerbation of autoimmune process in specific disease, whereas, on the other hand, ultraviolet radiations can also prevent or reduce the symptoms of other autoimmune diseases by vitamin D formation in skin as elaborated in various parts of this chapter.

7. Conclusion

In last decade, the number of studies that have investigated the non-bone effects of vitamin D has increased tremendously. Many studies from around the globe have measured circulating 25-hydroxyvitamin D as a determinant of vitamin D status. However, several queries regarding the definitive place of 25OH vitamin D as a biomarker of vitamin D status and the exact level of 25OHD that determines optimal vitamin D status in children for majority of non-bone effects of vitamin D still remain elusive. Although the committee from Institute of Medicine has recommended that a 25OHD level of 50 nmol/l (20 ng/ml) should be considered sufficient, this recommendation was mainly based on studies of bone health and the committee acknowledged that studies in other disease states are sorely lacking [4]. There are number of authors who have dissenting opinion regarding the IOM recommendations of optimal vitamin D levels [124]. Furthermore, there are data that suggest that optimal circulating levels regarding adequate bone health as well as other non-bone effects are much higher than the current IOM recommendations [125]. Most of the studies done in children have only measured vitamin D level once at one point in time. It is known that vitamin D levels vary over seasons and likely over time. Future studies need to measure 25OHD at multiple time points in relation to the outcome of interest. Finally, there is a continuing need for further conclusive studies to define appropriate levels of vitamin D status and recommended daily allowance regarding adequate bone health and other non-bone effects of vitamin D, especially in children, adolescents, and young adults.

Author details

Mohsin Ali Cheema and Khalid Parvez Lone*

*Address all correspondence to: khalid.lone@gmail.com

Department of Physiology and Cell Biology, and Centre for Research in Endocrinology and Reproductive Sciences (CRERS), University of Health Sciences, Lahore, Pakistan

References

- [1] Holick MF. Vitamin D deficiency. *N Engl J Med*. 2007;357(3):266–81.
- [2] Watson L. (2012). 10 things you need to know about vitamin D. [online] Available at: <http://www.drlisawatson.com/10-things-you-need-to-know-about-vitamin-d> [Accessed 10 January 2016].
- [3] Pettifor JM, Moodley GP, Hough FS, Koch H, Chen T, Lu Z, Holick MF. The effect of season and latitude on *in vitro* vitamin D formation by sunlight in South Africa. *S Afr Med J*. 1996;86(10):1270–2.
- [4] Ross AC, Manson JE, Abrams SA, Aloia JF, Brannon PM, Clinton SK, Durazo-Arvizu RA, Gallagher JC, Gallo RL, Jones G, Kovacs CS, Mayne ST, Rosen CJ, Shapses SA. The 2011 report on dietary reference intakes for calcium and vitamin D from the Institute of Medicine: what clinicians need to know. *J Clin Endocrinol Metab*. 2011;96(1):53–8.
- [5] Cranney A, Horsley T, O'Donnell S, Weiler H, Puil L, Ooi D, Atkinson S, Ward L, Moher D, Hanley D, Fang M, Yazdi F, Garritty C, Sampson M, Barrowman N, Tsertsvadze A, Mamaladze V. Effectiveness and safety of vitamin D in relation to bone health. *Evid Rep Technol Assess (Full Rep)*. 2007;(158):1–235.
- [6] Heaney RP. Advances in therapy for osteoporosis. *Clin Med Res*. 2003;1(2):93–9. Review.
- [7] Harinarayan CV, Ramalakshmi T, Prasad UV, Sudhakar D. Vitamin D status in Andhra Pradesh: a population based study. *Indian J Med Res*. 2008;127(3):211–8.
- [8] Goldring SR, Krane SM, Aviolo LV. Disorders of calcification: osteomalacia and rickets. In: LJD, editor. *Endocrinology*, 3rd edition. Philadelphia (PA): WB Saunders, pp. 1204–27, 1995.
- [9] Välimäki VV, Alftan H, Lehmuskallio E, Löyttyniemi E, Sahi T, Stenman UH, Suominen H, Välimäki MJ. Vitamin D status as a determinant of peak bone mass in young Finnish men. *J Clin Endocrinol Metab*. 2004;89(1):76–80.

- [10] Glerup H, Mikkelsen K, Poulsen L, Hass E, Overbeck S, Andersen H, Charles P, Eriksen EF. Hypovitaminosis D myopathy without biochemical signs of osteomalacic bone involvement. *Calcif Tissue Int.* 2000;66(6):419–424.
- [11] Bischoff-Ferrari HA, Borchers M, Gudat F, Dürmüller U, Stähelin HB, Dick W. Vitamin D receptor expression in human muscle tissue decreases with age. *J Bone Miner Res.* 2004;19(2):265–9.
- [12] Papadimitropoulos E, Wells G, Shea B, Gillespie W, Weaver B, Zytaruk N, Cranney A, Adachi J, Tugwell P, Josse R, Greenwood C, Guyatt G, Osteoporosis Methodology Group and The Osteoporosis Research Advisory Group. Meta-analyses of therapies for postmenopausal osteoporosis. VIII: meta-analysis of the efficacy of vitamin D treatment in preventing osteoporosis in postmenopausal women. *Endocr Rev.* 2002;23(4):560–9.
- [13] Holick MF. Resurrection of vitamin D deficiency and rickets. *J Clin Invest.* 2006;116:2062–72.
- [14] Siddiqui TS, Rai MI. Presentation and predisposing factors of nutritional rickets in children of Hazara Division. *J Ayub Med Coll Abbottabad.* 2005;17(3):29–32.
- [15] Anwar S, Iqbal MP, Azam I, Habib A, Bhutta S, Soofi SB, Bhutta ZA. Urban and rural comparison of vitamin D status in Pakistani pregnant women and neonates. *J Obstet Gynaecol.* 2016;36(3):318–23. doi:10.3109/01443615.2015.1050647. Epub 2015 Oct 14.
- [16] Jamal A, Khanani MR, Billo G, Asghar A, Jafri Z. Rickets in a slum of Karachi. *Pak J Med Sci.* 1996;12:247–50.
- [17] Qamar S, Akbani S, Shamim S, Khan G. Vitamin D levels in children with growing pains. *J Coll Phys Surg Pak.* 2011;21(5):284–7.
- [18] Iqbal R, Jafri L, Haroon A, Habib Khan A. Illuminating the dark side—vitamin D status in different localities of Karachi. *J Coll Phys Surg Pak.* 2013;23(8):604–6. doi:08.2013/JCPSP.604606.
- [19] Signorello LB, Williams SM, Zheng W, Smith JR, Long J, Cai Q, Hargreaves MK, Hollis BW, Blot WJ. Blood vitamin D levels in relation to genetic estimation of African ancestry. *Cancer Epidemiol Biomarkers Prev.* 2010;19(9):2325–31. doi:10.1158/1055-9965.EPI-10-0482. Epub 2010 Jul 20.
- [20] Shoenfeld N, Amital H, Shoenfeld Y. The effect of melanism and vitamin D synthesis on the incidence of autoimmune disease. *Nat Clin Pract Rheumatol.* 2009;5(2):99–105. doi:10.1038/ncprheum0989.
- [21] Signorello LB, Shi J, Cai Q, Zheng W, Williams SM, Long J, Cohen SS, Li G, Hollis BW, Smith JR, Blot WJ. Common variation in vitamin D pathway genes predicts circulating 25-hydroxyvitamin D levels among African Americans. *PLoS One.* 2011;6(12):e28623. doi:10.1371/journal.pone.0028623. Epub 2011 Dec 21.

- [22] Dawson-Hughes B. Racial/ethnic considerations in making recommendations for vitamin D for adult and elderly men and women. *Am J Clin Nutr.* 2004;80(6 Suppl): 1763S–6S.
- [23] Gallagher JC, Peacock M, Yalamanchili V, Smith LM. Effects of vitamin D supplementation in older African American women. *J Clin Endocrinol Metab.* 2013;98(3):1137–46. doi:10.1210/jc.2012-3106. Epub 2013 Feb 5.
- [24] Wortsman J, Matsuoka LY, Chen TC, Lu Z, Holick MF. Decreased bioavailability of vitamin D in obesity. *Am J Clin Nutr.* 2000;72(3):690–3.
- [25] *Arthritis Rheum.* 2009 Aug;60(8):2499–504. doi: 10.1002/art.24702. Ultraviolet radiation intensity predicts the relative distribution of dermatomyositis and anti-Mi-2 autoantibodies in women. Love LA, Weinberg CR, McConnaughey DR, Oddis CV, Medsger TA Jr, Reveille JD, Arnett FC, Targoff IN, Miller FW.
- [26] Lagunova Z, Porojnicu AC, Lindberg F, Hexeberg S, Moan J. The dependency of vitamin D status on body mass index, gender, age and season. *Anticancer Res.* 2009;29(9):3713–20.
- [27] Verdoia M, Schaffer A, Barbieri L, Di Giovine G, Marino P, Suryapranata H, De Luca G; Novara Atherosclerosis Study Group (NAS). Impact of gender difference on vitamin D status and its relationship with the extent of coronary artery disease. *Nutr Metab Cardiovasc Dis.* 2015;25(5):464–70. doi:10.1016/j.numecd.2015.01.009. Epub 2015 Feb 7.
- [28] Jablonski NG, Chaplin G. The evolution of human skin coloration. *J Hum Evol.* 2000;39(1):57–106.
- [29] Xue Y, Ying L, Horst RL, Watson G, Goltzman D. Androgens attenuate Vitamin D production induced by UVB irradiation of the skin of male mice by an enzymatic mechanism. *J Invest Dermatol.* 2015;135(12):3125–32. doi:10.1038/jid.2015.297. Epub 2015 Jul 27.
- [30] Ramagopalan SV, Heger A, Berlanga AJ, Maugeri NJ, Lincoln MR, Burrell A, Handunnetthi L, Handel AE, Disanto G, Orton SM, Watson CT, Morahan JM, Giovannoni G, Ponting CP, Ebers GC, Knight JC. A ChIP-seq defined genome-wide map of vitamin D receptor binding: associations with disease and evolution. *Genome Res.* 2010;20(10): 1352–60. doi:10.1101/gr.107920.110. Epub 2010 Aug 24.
- [31] Holick MF. Vitamin D: a millennium perspective. *J Cell Biochem.* 2003;88(2):296–307.
- [32] Colston K, Colston MJ, Feldman D. 1,25-Dihydroxyvitamin D3 and malignant melanoma: the presence of receptors and inhibition of cell growth in culture. *Endocrinology.* 1981;108(3):1083–6.
- [33] Cross HS, Bareis P, Hofer H, Bischof MG, Bajna E, Kriwanek S, Bonner E, Peterlik M. 25-Hydroxyvitamin D(3)-1alpha-hydroxylase and vitamin D receptor gene expression in human colonic mucosa is elevated during early cancerogenesis. *Steroids.* 2001;66(3–5):287–92.

- [34] MacLaughlin J, Holick MF. Aging decreases the capacity of human skin to produce vitamin D3. *J Clin Invest.* 1985;76(4):1536–8.
- [35] Mather KA, Jorm AF, Parslow RA, Christensen H. Is telomere length a biomarker of aging? A review. *J Gerontol A Biol Sci Med Sci.* 2011;66(2):202–13. doi:10.1093/gerona/gdq180. Epub 2010 Oct 28.
- [36] Liu JJ, Prescott J, Giovannucci E, Hankinson SE, Rosner B, Han J, De Vivo I. Plasma vitamin D biomarkers and leukocyte telomere length. *Am J Epidemiol.* 2013;177(12):1411–7. doi:10.1093/aje/kws435. Epub 2013 May 9.
- [37] Tuohimaa P. Vitamin D and aging. *J Steroid Biochem Mol Biol.* 2009;114(1–2):78–84.
- [38] Duque G, El Abdaimi K, Macoritto M, Miller MM, Kremer R. Estrogens (E2) regulate expression and response of 1,25-dihydroxyvitamin D3 receptors in bone cells: changes with aging and hormone deprivation. *Biochem Biophys Res Commun.* 2002;299(3):446–54.
- [39] Walters JR, Balesaria S, Chavele KM, Taylor V, Berry JL, Khair U, Barley NF, van Heel DA, Field J, Hayat JO, Bhattacharjee A, Jeffery R, Poulson R. Calcium channel TRPV6 expression in human duodenum: different relationships to the vitamin D system and aging in men and women. *J Bone Miner Res.* 2006;21(11):1770–7.
- [40] Welsh J, Wietzke JA, Zinser GM, Smyczek S, Romu S, Tribble E, Welsh JC, Byrne B, Narvaez CJ. Impact of the Vitamin D3 receptor on growth-regulatory pathways in mammary gland and breast cancer. *J Steroid Biochem Mol Biol.* 2002;83(1–5):85–92.
- [41] Klaus G, Weber L, Rodríguez J, Fernández P, Klein T, Grulich-Henn J, Hügel U, Ritz E, Mehls P. Interaction of IGF-I and 1 alpha, 25(OH)2D3 on receptor expression and growth stimulation in rat growth plate chondrocytes. *Kidney Int.* 1998;53(5):1152–61.
- [42] Andress D. Nonclassical aspects of differential vitamin D receptor activation: implications for survival in patients with chronic kidney disease. *Drugs.* 2007;67(14):1999–2012.
- [43] Fernandez-Martin JL, Kurian S, Farmer P, Nanes MS. Tumor necrosis factor activates a nuclear inhibitor of vitamin D and retinoid-X receptors. *Mol Cell Endocrinol.* 1998;141(1–2):65–72.
- [44] González Pardo V, Boland R, de Boland AR. Vitamin D receptor levels and binding are reduced in aged rat intestinal subcellular fractions. *Biogerontology.* 2008;9(2):109–18. Epub 2007 Dec 1.
- [45] Keisala T, Minasyan A, Lou YR, Zou J, Kalueff AV, Pyykkö I, Tuohimaa P. Premature aging in vitamin D receptor mutant mice. *J Steroid Biochem Mol Biol.* 2009;115(3–5):91–7. doi:10.1016/j.jsbmb.2009.03.007. Epub 2009 Mar 31.
- [46] Smit E, Crespo CJ, Michael Y, Ramirez-Marrero FA, Brodowicz GR, Bartlett S, Andersen RE. The effect of vitamin D and frailty on mortality among non-institutionalized US older adults. *Eur J Clin Nutr.* 2012;66(9):1024–8. doi:10.1038/ejcn.2012.67. Epub 2012 Jun 13.

- [47] Online available at: <http://www.cancer.gov/types/childhood-cancers/child-adolescent-cancers-fact-sheet>
- [48] Online available at: <http://ccfpakistan.org/2011/05/23/childhood-cancer/>
- [49] Helou M, Ning Y, Yang S, Irvine P, Bachmann LM, Godder K, Massey G. Vitamin D deficiency in children with cancer. *J Pediatr Hematol Oncol.* 2014;36(3):212–7. doi: 10.1097/MPH.0b013e31829f3754.
- [50] Modan-Moses D, Pinhas-Hamiel O, Munitz-Shenkar D, Temam V, Kanety H, Toren A. Vitamin D status in pediatric patients with a history of malignancy. *Pediatr Res.* 2012;72(6):620–4. doi:10.1038/pr.2012.131. Epub 2012 Oct 4.
- [51] Lin R, White JH. The pleiotropic actions of vitamin D. *Bioessays.* 2004;26(1):21–8. pmid: 14696037 doi:10.1002/bies.10368
- [52] Jones G, Strugnell SA, DeLuca HF. Current understanding of the molecular actions of vitamin D. *Physiol Rev.* 1998;78(4):1193–231. pmid:9790574
- [53] Tangpricha V, Flanagan JN, Whitlatch LW, Tseng CC, Chen TC, Holt PR, Lipkin MS, Holick MF. 25-hydroxyvitamin D-1 α -hydroxylase in normal and malignant colon tissue. *Lancet.* 2001;357(9269):1673–4.
- [54] Mawer EB, Hayes ME, Heys SE, Davies M, White A, Stewart MF, Smith GN. Constitutive synthesis of 1,25-dihydroxyvitamin D₃ by a human small cell lung cancer cell line. *J Clin Endocrinol Metab.* 1994;79(2):554–60.
- [55] Garland CF, Garland FC, Gorham ED. Can colon cancer incidence and death rates be reduced with calcium and vitamin D? *Am J Clin Nutr.* 1991;54(1 Suppl):193S–201S.
- [56] Garland CF, Comstock GW, Garland FC, Helsing KJ, Shaw EK, Gorham ED. Serum 25-hydroxyvitamin D and colon cancer: eight-year prospective study. *Lancet.* 1989;2(8673): 1176–8.
- [57] Hanchette CL, Schwartz GG. Geographic patterns of prostate cancer mortality. Evidence for a protective effect of ultraviolet radiation. *Cancer.* 1992;70(12):2861–9.
- [58] Guy M, Lowe LC, Bretherton-Watt D, Mansi JL, Peckitt C, Bliss J, Wilson RG, Thomas V, Colston KW. Vitamin D receptor gene polymorphisms and breast cancer risk. *Clin Cancer Res.* 2004;10(16):5472–81.
- [59] Rashid MU, Muzaffar M, Khan FA, Kabisch M, Muhammad N, Faiz S, Loya A, Hamann U. Association between the BsmI polymorphism in the vitamin D receptor gene and breast cancer risk: results from a Pakistani Case-Control Study. *PLoS One.* 2015;10(10):e0141562. doi:10.1371/journal.pone.0141562. eCollection 2015.
- [60] Feldman D, Krishnan AV, Swami S, Giovannucci E, Feldman BJ. The role of vitamin D in reducing cancer risk and progression. *Nat Rev Cancer.* 2014;14(5):342–57. doi: 10.1038/nrc3691. pmid:24705652. Epub 2014 Apr 4.

- [61] Visweswaran RK, Lekha H. Extraskelatal effects and manifestations of Vitamin D deficiency. *Indian J Endocrinol Metab.* 2013;17(4):602–10. doi:10.4103/2230-8210.113750.
- [62] Pálmer HG, Larriba MJ, García JM, Ordóñez-Morán P, Peña C, Peiró S, Puig I, Rodríguez R, de la Fuente R, Bernad A, Pollán M, Bonilla F, Gamallo C, de Herreros AG, Muñoz A. The transcription factor SNAIL represses vitamin D receptor expression and responsiveness in human colon cancer. *Nat Med.* 2004;10(9):917–9.
- [63] Deeb KK, Trump DL, Johnson CS. Vitamin D signalling pathways in cancer: potential for anticancer therapeutics. *Nat Rev Cancer.* 2007;7(9):684–700.
- [64] Online available at: <http://www.diabetesatlas.org/>
- [65] Pozzilli P, Manfrini S, Crino A, Picardi A, Leomanni C, Cherubini V, Valente L, Khazrai M, Visalli N; IMDIAB group. Low levels of 25-hydroxyvitamin D3 and 1,25-dihydroxyvitamin D3 in patients with newly diagnosed type 1 diabetes. *Horm Metab Res.* 2005;37:680–3. [PubMed].
- [66] Littorin B, Blom P, Scholin A, Arnqvist HJ, Blohme G, Bolinder J, Ekbom-Schnell A, Eriksson JW, Gudbjörnsdottir S, Nyström L, Ostman J, Sundkvist G. Lower levels of plasma 25-hydroxyvitamin D among young adults at diagnosis of autoimmune T1D compared with control subjects: results from the nationwide Diabetes Incidence Study in Sweden (DISS) Diabetologia. 2006;49:2847–52. [PubMed].
- [67] Greer RM, Rogers MA, Bowling FG, Buntain HM, Harris M, Leong GM, Cotterill AM. Australian children and adolescents with T1D have low vitamin D levels. *Med J Aust.* 2007;187:59–60. [PubMed].
- [68] Svoren BM, Volkening LK, Wood JR, Laffel LM. Significant vitamin D deficiency in youth with T1D mellitus. *J Pediatr.* 2009;154:132–4. [PMC free article] [PubMed].
- [69] Borkar VV, Devidayal, Verma S, Bhalla AK. Low levels of vitamin D in North Indian children with newly diagnosed type 1 diabetes. *Pediatr Diabetes.* 2010;11:345–50. [PubMed].
- [70] Bener A, Alsaied A, Al-Ali M, Al-Kubaisi A, Basha B, Abraham A, Guiter G, Mian M. High prevalence of vitamin D deficiency in T1D mellitus and healthy children. *Acta Diabetol.* 2009;46:183–9. [PubMed].
- [71] Zipitis CS, Akobeng AK. Vitamin D supplementation in early childhood and risk of type 1 diabetes: a systematic review and meta-analysis. *Arch Dis Child.* 2008;93:512–7.
- [72] Hyppönen E, Läärä E, Reunanen A, Järvelin MR, Virtanen SM. Intake of vitamin D and risk of type 1 diabetes: a birth-cohort study. *Lancet.* 2001;358:1500–3.
- [73] Stene LC, Joner G. Use of cod liver oil during the first year of life is associated with lower risk of childhood onset type 1 diabetes: a large population-based, case control study. *Am J Clin Nutr.* 2003;78:1128–34.

- [74] Roomi MA, Lone KP, Madassar A. Vitamin D and cardiometabolic risk factors in adult non-diabetic offspring of type 2 diabetic parents. *J Pak Med Assoc.* 2014;64(11):1229–34.
- [75] Mohr SB, Garland CF, Gorham ED, Garland FC. The association between ultraviolet B irradiance, vitamin D status and incidence rates of type 1 diabetes in 51 regions worldwide. *Diabetologia.* 2008;51(8):1391–8. doi:10.1007/s00125-008-1061-5. Epub 2008 Jun 12.
- [76] Staples JA, Ponsonby AL, Lim LL, McMichael AJ. Ecologic analysis of some immune-related disorders, including type 1 diabetes, in Australia: latitude, regional ultraviolet radiation, and disease prevalence. *Environ Health Perspect.* 2003;111(4):518–23.
- [77] American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care.* 2009;32(Suppl 1):S62–7.
- [78] Gregori S, Giarratana N, Smirolto S, Uskokovic M, Adorini L. A 1 α ,25-dihydroxyvitamin D(3) analog enhances regulatory T-cells and arrests autoimmune diabetes in NOD mice. *Diabetes.* 2002;51(5):1367–74.
- [79] Mathieu C1, Waer M, Casteels K, Laureys J, Bouillon R. Prevention of type I diabetes in NOD mice by nonhypercalcemic doses of a new structural analog of 1,25-dihydroxyvitamin D3, KH1060. *Endocrinology.* 1995;136(3):866–72.
- [80] Aljabri KS, Bokhari SA, and Khan MJ. Glycemic changes after vitamin D supplementation in patients with type 1 diabetes mellitus and vitamin D deficiency. *Ann Saudi Med.* 2010;30(6):454–8.
- [81] Masoli M, Fabian D, Holt S, Beasley R. The global burden of asthma: executive summary of the GINA Dissemination Committee report. *Allergy.* 2004;59:469–78.
- [82] Mannino DM, Homa DM, Akinbami LJ, et al. Surveillance for asthma: United States, 1980–1999. *MMWR Surveill Summ.* 2002;51:1–13.
- [83] Devereux G. The increase in allergic disease: environment and susceptibility. Proceedings of a symposium held at the Royal Society of Edinburgh, 4th June 2002. *Clin Exp Allergy.* 2003;33:394–406.
- [84] (CDC) CfDCaP. Vital signs: asthma prevalence, disease characteristics, and self-management education: United States, 2001–2009. *Morb Mortal Wkly Rep.* 2011;60:547–52.
- [85] Hasnain SM, Khan M, Saleem A, Waqar MA. Prevalence of asthma and allergic rhinitis among school children of Karachi, Pakistan, 2007. *J Asthma.* 2009;46(1):86–90. doi:10.1080/02770900802513023.
- [86] Khan AA, Tanzil S, Jamali T, Shahid A, Naeem S, Sahito A, Siddiqui FA, Nafees AA, Fatmi Z. Burden of asthma among children in a developing megacity: childhood

- asthma study, Pakistan. *J Asthma*. 2014;51(9):891–9. doi:10.3109/02770903.2014.930882. Epub 2014 Jul 3.
- [87] Litonjua AA. Vitamin D deficiency as a risk factor for childhood allergic disease and asthma. *Curr Opin Allergy Clin Immunol*. 2012;12(2):179–85. doi:10.1097/ACI.0b013e3283507927.
- [88] Litonjua AA. Childhood asthma may be a consequence of vitamin D deficiency. *Curr Opin Allergy Clin Immunol*. 2009;9(3):202–7. doi:10.1097/ACI.0b013e32832b36cd.
- [89] Raby BA, Lazarus R, Silverman EK, Lake S, Lange C, Wjst M, Weiss ST. Association of vitamin D receptor gene polymorphisms with childhood and adult asthma. *Am J Respir Crit Care Med*. 2004;170(10):1057–65. Epub 2004 Jul 28.
- [90] Poon AH, Laprise C, Lemire M, Montpetit A, Sinnett D, Schurr E, Hudson TJ. Association of vitamin D receptor genetic variants with susceptibility to asthma and atopy. *Am J Respir Crit Care Med*. 2004;170(9):967–73. Epub 2004 Jul 28.
- [91] Wjst M, Altmüller J, Faus-Kessler T, Braig C, Bahnweg M, André E. Asthma families show transmission disequilibrium of gene variants in the vitamin D metabolism and signalling pathway. *Respir Res*. 2006;7:60.
- [92] Ramsey CD, Celedón JC. The hygiene hypothesis and asthma. *Curr Opin Pulm Med*. 2005;11(1):14–20.
- [93] Corne JM, Marshall C, Smith S, Schreiber J, Sanderson G, Holgate ST, Johnston SL. Frequency, severity, and duration of rhinovirus infections in asthmatic and non-asthmatic individuals: a longitudinal cohort study. *Lancet*. 2002;359(9309):831–4.
- [94] Liu PT, Stenger S, Tang DH, Modlin RL. Cutting edge: vitamin D-mediated human antimicrobial activity against *Mycobacterium tuberculosis* is dependent on the induction of cathelicidin. *J Immunol*. 2007;179(4):2060–3.
- [95] Herr C, Shaykhiev R, Bals R. The role of cathelicidin and defensins in pulmonary inflammatory diseases. *Expert Opin Biol Ther*. 2007;7(9):1449–61.
- [96] Ginde AA, Mansbach JM, Camargo CA Jr. Association between serum 25-hydroxyvitamin D level and upper respiratory tract infection in the Third National Health and Nutrition Examination Survey. *Arch Intern Med*. 2009;169(4):384–90. doi:10.1001/archinternmed.2008.560.
- [97] Zosky GR, Berry LJ, Elliot JG, James AL, Gorman S, Hart PH. Vitamin D deficiency causes deficits in lung function and alters lung structure. *Am J Respir Crit Care Med*. 2011;183(10):1336–43.
- [98] Chi A, Wildfire J, McLoughlin R, Wood RA, Bloomberg GR, Kattan M, Gergen P, Gold DR, Witter F, Chen T, Holick M, Visness C, Gern J, O'Connor GT. Umbilical cord plasma 25-hydroxyvitamin D concentration and immune function at birth: the Urban Envi-

- ronment and Childhood Asthma study. *Clin Exp Allergy*. 2011;41(6):842–50. doi: 10.1111/j.1365-2222.2011.03712.x. Epub 2011 Apr 11.
- [99] Miyake Y, Sasaki S, Tanaka K, Hirota Y. Dairy food, calcium and vitamin D intake in pregnancy, and wheeze and eczema in infants. *Eur Respir J*. 2010;35(6):1228–34.
- [100] Rothers J, Wright AL, Stern DA, Halonen M, Camargo CA Jr. Cord blood 25-hydroxy-vitamin D levels are associated with aeroallergen sensitization in children from Tucson, Arizona. *J Allergy Clin Immunol*. 2011;128(5):1093–9.e1–5.
- [101] Camargo CA Jr, Ingham T, Wickens K, Thadhani R, Silvers KM, Epton MJ, Town GI, Pattemore PK, Espinola JA, Crane J, New Zealand Asthma and Allergy Cohort Study Group. Cord-blood 25-hydroxyvitamin D levels and risk of respiratory infection, wheezing, and asthma. *Pediatrics*. 2011;127(1):e180–7.
- [102] Freishtat RJ, Iqbal SF, Pillai DK, Klein CJ, Ryan LM, Benton AS, Teach SJ. High prevalence of vitamin D deficiency among inner-city African American youth with asthma in Washington, DC. *J Pediatr*. 2010;156(6):948–52.
- [103] Bener A, Ehlayel MS, Tulic MK, Hamid Q. Vitamin D deficiency as a strong predictor of asthma in children. *Int Arch Allergy Immunol*. 2012;157(2):168–75.
- [104] Alyasin S, Momen T, Kashef S, Alipour A, Amin R. The relationship between serum 25 hydroxy vitamin D levels and asthma in children. *Allergy Asthma Immunol Res*. 2011;3(4):251–5.
- [105] Available at: <http://autoimmune.pathology.jhmi.edu/faqs.cfm>. [Accessed 23 March 2016].
- [106] Available at: <http://www.childrenshospital.org/conditions-and-treatments/conditions/autoimmune-diseases> 15th April, 2016
- [107] Adorini A, Penna G. Control of autoimmune diseases by the vitamin D endocrine system. *Nat Clin Pract Rheumatol*. 2008;4:404–12
- [108] Bultink IE, Lems WF, Kostense PJ, Dijkmans BA, Voskuyl AE. Prevalence of and risk factors for low bone mineral density and vertebral fractures in patients with systemic lupus erythematosus. *Arthritis Rheum*. 2005;52(7):2044–50.
- [109] Wright TB, Shults J, Leonard MB, Zemel BS, Burnham JM. Hypovitaminosis D is associated with greater body mass index and disease activity in pediatric systemic lupus erythematosus. *J Pediatr*. 2009;155(2):260–5.
- [110] Robinson AB, Thierry-Palmer M, Gibson KL, Rabinovich CE. Disease activity, proteinuria, and vitamin D status in children with systemic lupus erythematosus and juvenile dermatomyositis. *J Pediatr*. 2012;160(2):297–302. doi:10.1016/j.jpeds.2011.08.011. Epub 2011 Sep 15.

- [111] Abe J, Nakamura K, Takita Y, Nakano T, Irie H, Nishii Y. Prevention of immunological disorders in MRL/l mice by a new synthetic analogue of vitamin D3: 22-oxa-1 alpha, 25-dihydroxyvitamin D3. *J Nutr Sci Vitaminol (Tokyo)*. 1990;36(1):21–31
- [112] Lemire JM, Ince A, Takashima M. 1,25-Dihydroxyvitamin D3 attenuates the expression of experimental murine lupus of MRL/l mice. *Autoimmunity*. 1992;12(2):143–8.
- [113] Kamen DL, Cooper GS, Bouali H, Shaftman SR, Hollis BW, Gilkeson GS. Vitamin D deficiency in systemic lupus erythematosus. *Autoimmun Rev*. 2006;5(2):114–7
- [114] Nagpal S, Na S, Rathnachalam R. Noncalcemic actions of vitamin D receptor ligands. *Endocr Rev*. 2005;26(5):662–87.
- [115] Munger KL, Zhang SM, O'Reilly E, Hernán MA, Olek MJ, Willett WC. Vitamin D intake and incidence of multiple sclerosis. *Neurology*. 2004;62:60–5.
- [116] Goldberg P, Fleming MC, Picard EH. Multiple sclerosis: decreased relapse rate through dietary supplementation with calcium, magnesium and vitamin D. *Med Hypotheses*. 1986;21:193–200.
- [117] Ramagopalan SV, Maugeri NJ, Handunnetthi L, Lincoln MR, Orton SM, Dymment DA, Deluca GC, Herrera BM, Chao MJ, Sadovnick AD, Ebers GC, Knight JC. Expression of the multiple sclerosis-associated MHC class II Allele HLA-DRB1*1501 is regulated by vitamin D. *PLoS Genet*. 2009;5(2):e1000369.
- [118] Blaney GP, Albert PJ, Proal AD. Vitamin D metabolites as clinical markers in autoimmune and chronic disease. *Ann N Y Acad Sci*. 2009;1173:384–90. doi:10.1111/j.1749-6632.2009.04875.x.
- [119] Welsh P, Sattar N. Vitamin D and chronic disease prevention. *BMJ*. 2014;348:g2280. doi:10.1136/bmj.g2280.
- [120] Sattar N, Welsh P, Panarelli M, Forouhi NG. Increasing requests for vitamin D measurement: costly, confusing, and without credibility. *Lancet*. 2012;379(9811):95–6. doi:10.1016/S0140-6736(11)61816-3.
- [121] Fletcher RH. Review: vitamin D3 supplementation may reduce mortality in adults; vitamin D2 does not. *Ann Intern Med*. 2014;161(2):JC5. doi:10.7326/0003-4819-161-2-201407150-02005.
- [122] Fraser PA, Ding WZ, Mohseni M, Treadwell EL, Dooley MA, St Clair EW, Gilkeson GS, Cooper GS. Glutathione S-transferase M null homozygosity and risk of systemic lupus erythematosus associated with sun exposure: a possible gene-environment interaction for autoimmunity. *J Rheumatol*. 2003;30(2):276–82.
- [123] Love LA, Weinberg CR, McConnaughey DR, Oddis CV, Medsger TA Jr, Reveille JD, Arnett FC, Targoff IN, Miller FW. Ultraviolet radiation intensity predicts the relative distribution of dermatomyositis and anti-Mi-2 autoantibodies in women. *Arthritis Rheum*. 2009;60(8):2499–504. doi:10.1002/art.24702.

- [124] Heaney RP, Holick MF. Why the IOM recommendations for vitamin D are deficient. *J Bone Miner Res.* 2011;26(3):455–7. doi:10.1002/jbmr.328.
- [125] Heaney RP, Armas LA, Shary JR, Bell NH, Binkley N, Hollis BW. 25-Hydroxylation of vitamin D3: relation to circulating vitamin D3 under various input conditions. *Am J Clin Nutr.* 2008;87(6):1738–42.

IntechOpen

IntechOpen