We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Telomere and Telomerase in Cancer

Angayarkanni Jayaraman,
Kalarikkal Gopikrishnan Kiran and
Palaniswamy Muthusamy

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/64721>

Abstract

The linear ends of chromosome are protected by specialized ribonucleoprotein (RNP) termed as telomere. These specialized terminal elements with tandem repeated sequence are the protective cap that alleviate end replication problem and cell senescence. The telomere length maintenance is essential to avoid cell death and apoptosis. Telomere shortening has been related to chronic stress due to several factors, which include not only psychological stress but also diseases such as cardiovascular diseases and cancer. Telomerase enzyme which maintains telomere length is the major factor responsible for evading cell death. Telomere length maintenance and telomerase expression put together are the prerequisite for immortality, an essential character for cancer cells. Understanding the mechanism of telomere and telomerase functions paves way for eradicating the diseases such as cancer.

Keywords: telomere length, telomerase, tumor progression, tumor immortality

1. Introduction

Telomeres are specialized DNA structures consisting of tandem arrays of hexa-nucleotide (TTAGGG) DNA sequences that cap the end of chromosomes. These structures residing near the ends of chromosomes play a very vital role in maintaining the integrity of the whole genome and prevent the loss of genetic material. As a consequence of cell division, short stretches of DNA will be lost from telomeric region and eventually lead to cellular senescence and death. There are a lot of evidences to suggest that telomere length is a better biomarker for overall health status, compared to the currently used biomarkers for the same. Few of the phenomena regarding the telomere length are unanswered till now. Till date it is unclear about the mechanistic

regarding the telomere length are unanswered till now. Till date it is unclear about the mechanistic correlation between telomere size and life span of various species. To prevent senescence of cell, the preservation of telomere length is essential, which is maintained by telomerase enzyme. Telomerase is a reverse transcriptase enzyme which has recently emerged as an attractive target for cancer as it is a crucial factor required for the tumor immortalization of a subset of cells, including cancer stem cells. Studies have proved that 80–85% of the tumor cells express telomerase whereas somatic cells lack the expression of telomerase [1]. The important paradox is that telomerase-negative normal cells have lengthier telomeres than telomerase-positive cancer cells [2]. Thus difference in telomere length and cell kinetics between normal and cancerous cells shows that targeting telomerase is a more effective way to target cancer cells. Owing to the significant role of telomere and telomerase in tumor immortalization understanding their amassed influence on cancer is crucial for cancer therapy. In this context, this book chapter is focused on disseminating the integrated impact of telomere and telomerase on cancer progression.

2. Telomere construction

The basic structure of telomere is conserved among eukaryotes and consists of short tandem DNA repeats, with G-rich sequence at the three-end referred to as the G-strand and the complementary strand is called the C-strand at the five-end (**Figure 1**) [3]. The length of telomeric DNA varies from 2 to 20 kilobase pairs, depending on factors such as tissue type and human age [4]. The telomere is conspicuous owing to the presence of G-overhang which extends beyond its complementary C-rich strand to form a single-stranded overhang, termed the G-tail. It has been proposed that the 3' G-overhang can be sequestered into a lasso-like structure known as the T-loop (**Figure 2**) [5, 6]. The single-stranded G-overhang invades the double-stranded telomeric DNA, which displaces the bound G-strand base-pairing with the C-strand. As this displaced binding takes place at a distance from the physical end of the telomere, it generates a large duplex structure called the T-loop (**Figure 2**) [3, 5].

Figure 1. Telomeric DNA.

Telomeres can also fold into G-quadruplex DNA, an unusual DNA conformation that is based on a guanine quartet [7]. The repetitive and GC-rich nature of telomeric DNA endows it with the capability to form the higher-order DNA conformation, G-quadruplexes [8, 9]. To maintain such unusual structure of telomeres, a set of telomeric protein complex has evolved, termed shelterin. The shelterin complex consists of six individual proteins, telomeric repeat binding factor 1 (TRF1), TRF2, repressor/activator protein 1 (RAP1), TIN2 (TRF1 interacting protein 2),

TPP1 (TINT1/PIP1/PTOP 1), and protection of telomeres 1 (POT1) [6, 10]. The proteins TRF1 and TRF2 attach to double-stranded telomeric repeats facilitating the anchoring of the complex along the length of telomeres [11–14], whereas POT1 binds to the single-stranded overhang. TPP1 and TIN2 act as bridging proteins between the above DNA-binding modules and are crucial for chromosome end protection and telomere length regulation. TRF1 and TRF2 with the help of TIN2 [15] bind with POT1 via TPP1 [16–19]. TIN2 also connects TRF1 to TRF2, and this interaction contributes to the stabilization of TRF2 on telomeres [17–19]. Besides this shelterin complex, other proteins like TEN1 and Pinx1, which are not telomere specific, are also present at telomeres and carry out important functions in recruitment of telomerase [20].

Figure 2. T-loop formation.

Figure 3. Telomere-telomerase assembly.

3. Telomere: a knight cap

During the evolution of linear genome, the natural ends of linear chromosomes resemble DNA breaks and tend to induce DNA damage response (DDR). These natural linear ends are protected by the sequestration of the ribonucleoprotein (RNP) sequence, telomeres which mask the ends from continuous exposure to the DNA damage response (DDR). Telomeres serve as protective caps, preventing the chromosomal ends from being recognized as double-strand breaks by the DNA damage repair system and the activation of the p53 or p16INK4a pathway and the start of senescence or apoptosis. If the telomere cap is removed, genome instability is induced. Telomeres with its tightly regulated complexes consisting of repetitive G-rich DNA and specialized proteins accomplish the task of not only concealing the linear chromosome ends from detection and undesired repair, but also protect from checkpoints, homologous recombination, end-to-end fusions, or other events that normally promote repair of intra-chromosomal DNA breaks acts [21]. Telomeric proteins and their interacting factors create an environment at chromosome ends that inhibits DNA repair at that point; however, the repair machinery is also essential for proper telomere function.

The closed configuration of the T-loop of telomeric region provides a protective cap that defines the natural end of the chromosome and masks the telomere from the DDR machinery (**Figure 2**) [6]. In particular, T-loops could provide an architectural solution to the repression of the ataxia telangiectasia mutated (ATM) kinase pathway, which relies on a sensor (the MRN (Mre11/Rad50/Nbs1) complex) with DNA end-binding activity. In addition, T-loops could prevent the Ku70/80 heterodimer, a DNA repair factor that binds to DNA ends, from loading onto the telomere terminus, thereby blocking the initiation of the non-homologous end-joining (NHEJ) pathway (**Figure 4**).

Figure 4. DNA damage response pathway.

Among the DNA-binding proteins, TRF1 has DNA remodeling activity [5, 22] and also shown to promote efficient replication of telomeres [23, 24]. On the contrary, TRF2 engages in chromosome end protection by inducing topological changes in telomeric DNA [25], T-loop assembly [26, 27] and by suppression of ATM dependent DDR and NHEJ (**Figure 4**) [28, 29]. Besides that, TRF2 plays a critical role in chromatin assembly, which was demonstrated by the observation that overexpression of TRF2 resulted in aberrant nucleosome spacing and decreased abundance of the core histones H3 and H4 at chromosome ends [30]. TRF2 lacking cells are reported to be growth arrested because of up-regulation of p53 and also show other hallmarks of ATM signaling, including the phosphorylation of ATM and Chk2 [31, 32]. Among the two DDR, the ATM kinase pathway at telomeres is repressed by TRF2 subunit, whereas POT1 is responsible for protection of telomere by suppression of ATR (ATM Rad3-related protein)-dependant DDR pathways [28, 33].

The high affinity of POT1 for single-stranded telomeric DNA makes it a G-strand binding component displaced from the T-loop and forms a closed configuration locking in the structure (**Figure 2**). Earlier report models suggest that POT1 and TPP1 compete with telomerase for access to the overhang [33]. Contrarily, direct interaction between TPP1 and telomerase bolsters telomerase processivity [34, 35] whereas increased loading of POT1 along the overhang block telomerase accessibility to the 3'-OH substrate. The role of RAP1 is obscure and its function has recently been elucidated. The presence of RAP1 at telomeres appears as a backup mechanism to prevent NHEJ when topology-mediated telomere protection is impaired [36]. In case of mutation in TRF2 which wraps the DNA, RAP1 has been implicated in the inhibition of NHEJ [37, 38].

Thus physically taken together, the shelterin complex and G-overhang of telomere are the protective cap that have an immensely complex role in convergence of end protection and telomere length maintenance mechanisms.

4. Telomerase activity

With each cell division, telomere length is reduced by ~50 to 200 bp [39] primarily because the lagging strand of DNA synthesis is unable to replicate the extreme 3' end of the chromosome which is denoted as end replication problem [40, 41]. When telomeres become sufficiently short, cells enter an irreversible growth arrest called cellular senescence. In most eukaryotes, telomeres are stabilized, and the shortening telomeric DNA is replenished, by the action of the RNP reverse transcriptase telomerase. Progressive telomere loss has been experimentally demonstrated using non-immortalized cells in culture that lack detectable telomerase [42, 43].

In cells with active telomerase, such as cancer cells, the telomere length is continually being built up and shortened in a regulated way that maintains telomere length homeostasis and retains telomere functionality. Shortening of telomeres occur due to nucleolytic degradation and incomplete DNA replication. On the contrary, lengthening is primarily accomplished by the action of a specialized reverse transcriptase called telomerase [44] and occasionally by homologous recombination (HR) [45]. Telomerase uses the 3' G-rich strand of a chromosome

as primer to elongate chromosome end by reverse-transcribing the template region of its tightly associated RNA moiety and coordinative action with the DNA replication machinery [44, 46]. For lengthening activity, telomerase requires not only hTERT catalytic subunit and RNA template (hTR) but also other factors [47, 48].

The 3' half of the hTR resembling the box H/ACA family of small nucleolar RNAs (snoRNAs) [49, 50] is essential for proper 3'-end processing, stability and nucleolar targeting *in vivo* [44]. The 5' end of hTR not only acts as template for the telomere extension at chromosome ends [5, 51] but also serves as a pseudoknot that is likely to be important for telomerase function (5, 49). A 6 bp U-rich sequence at the 5' end of hTR also interacts directly with hnRNPs C1 and C2 (**Figure 3**) [52]. Even though hTR is highly expressed in all tissues regardless of telomerase activity [53], in cancer cells hTR is generally expressed fivefold higher than normal cells [54]. However, the expression (mRNA) of the telomeric catalytic component hTERT which is closely associated with telomerase activity is estimated to be less than one to five copies per cell [54]. hTERT is generally repressed in normal cells and up-regulated in immortal cells, suggesting that hTERT is the primary determinant for the enzyme activity.

Figure 5. Telomerase complex.

It has been suggested that in addition to telomere elongation another aspect of telomerase RNP function is to allow even short telomeres to remain functional, which in the absence of telomerase would have caused cells to stop dividing or led to telomere–telomere fusions [55].

In other words, telomerase permits cell proliferation by stabilizing short telomeres that would be unstable in the absence of functional telomerase. In recent years, evidence has accumulated that telomerase, and in particular its catalytic subunit TERT, is involved in various non-telomere-related functions such as regulation of gene expression, growth factors and cell proliferation [56–61]. It has been reported that the telomerase has a role in modulation of Wnt/ β -catenin pathway [60]. TERT has been demonstrated to bind to TBE-containing promoter elements, the specific chromatin sites of Wnt/ β -catenin target genes, forming a part of the β -catenin transcriptional complex, which was facilitated by interaction with BRG1. These data endorsed the precipitous role for telomerase as a transcriptional modulator of Wnt/ β -catenin signaling pathway involved in progenitor cell regulation.

In addition, various groups have shown that TERT shuttles from the nucleus and translocates to mitochondria upon exogenous stress [62–67]. Singhapol and his coworkers have demonstrated that mitochondrial telomerase localization specifically decreases mitochondrial ROS generation and cellular oxidative stress after induction of exogenous stress generated by H_2O_2 or irradiation in cancer cells and might thereby prevent damage to nuclear DNA [68]. Thus the presence of telomerase not only maintains telomere length imparting immortality but also play multifarious role in tumorigenesis via non-telomere-dependent mechanism which demonstrated the imperative ubiquity of telomerase in cancer cells.

5. Skewed expression of telomerase

Telomerase, the RNA-dependent DNA polymerase by preventing the shortening of telomeric DNA sequences, accouters unlimited proliferation. As per the telomere hypothesis of cancer cell immortalization, telomere shortening limits the life span of telomerase-negative normal cells, whereas telomerase activation in cancer cells extends their life span [4]. In normal human cells, telomerase activity is quenched during embryonic differentiation [69]. On the contrary in some tissues, like male germ cells, activated lymphocytes, and certain types of stem cell populations, the telomerase activity is induced [15, 70]. Owing to its diverse activity, the telomerase [71] which was established to be absent in most of the normal human somatic cells is recorded to be expressed in more than 90% of cancerous cells and in vitro-immortalized cells [15, 70]. A study showed that while most of the glioma tissues possess increased telomerase activity, only few (10%) anaplastic astrocytomas are reported to be telomerase positive [72–74]. In contrast to most cancerous cells, the telomerase expression is present in only 50% of glioblastoma and retinoblastoma samples, and activity is even rarely found in meningiomas and astrocytomas [75, 76].

Induction of telomerase activity in primary human keratinocytes and mammary epithelial cells has been attributed to the effect of human papillomavirus 16 E6 protein [77]. Similarly, during the menstrual cycle involving the proliferation of endometrial cells, telomerase activity is detected in normal human endometrium [78, 79]. These reports emphasis that telomerase might be the reason for tumorigenesis in hormone-dependent cancers.

It has been suggested that up-regulated expression of telomerase is contributed by the increased copy number of hTERT which was demonstrated by the report that while hTERT protein expression was strongly positive in tumor cells, the expression of hTERT in non-neoplastic mucosal cells as well as stromal elements (except lymphocytes) was weak or negative [80]. In most cases, hTERT expression is closely correlated not only with telomerase activity but also with cancer initiation and progression. In head and neck squamous cell carcinoma and human glioma cell lines, there was decrease in telomerase activity which has been correlated with overexpression of p53, E2F, p16, p21, and p15 individually [81, 82]. In malignant and nonmalignant human hematopoietic cell lines, primary leukemic cells, and normal T lymphocytes, IFN- α is reported to inhibit telomerase activity by suppressing hTERT transcription [83]. In addition to growth and differentiation-related regulation, telomerase activity is subject to regulation by other external and intracellular factors such as UV irradiation [84]. The telomerase having influence over several signaling pathways that determine cell proliferative or death responses when overexpressed might abrogate anti-proliferative or cell death signals. Thus cancer cells with high levels of telomerase might gain a selective growth advantage.

6. Telomerase as biomarker of cancer

Advent of latest cancer biomarkers has increased opportunities for improving cancer diagnostics by enhancing the quickness of detection and efficacy of treatment. In relation to the practice of new therapeutic interventions, proficient biomarkers are helpful in detection and prediction of remission or relapse of cancer at both gross and molecular levels. Telomerase activity is a hallmark of most cancer biopsies, but not generally detected in premalignant lesions and in normal tissue samples except germ cells and hematopoietic stem cells. Thus telomerase activity can be a promising biomarker for diagnosis of malignancies and a target for chemotherapy or gene therapy. Extent of telomerase activity in tumor tissues may be prognostic indicators of patient outcome. Thus, at the present time telomerase is being studied in anticipation of clinical usage. Many clinical trials for telomerase assay in cancer diagnosis are under trial. Fresh or fresh-frozen biopsies, fluids, and secretions are subject for these trials.

Other components of telomerase enzyme complex have also been utilized as biomarkers for telomerase activity. The expression of the RNA subunit of the telomerase complex (hTR) is also regarded as a diagnostic marker [85]. But the expression of hTR does not always correlate with telomerase protein expression in that particular cell type. hTR can be constitutively expressed in certain cell types in which even telomerase activity is not present [86]. Apart from this, mutation in genes of telomerase and associated proteins are considered as a diagnostic and prognostic marker for many genetic abnormalities collectively termed as telomeropathies. Early-onset melanoma tumor syndrome with multiple co-morbid cancers can be predicted from telomerase gene promoter mutation analysis. In this disorder, the mutation in promoter of telomerase gene introduces an erythroblast transformation-specific transcription factor-binding site, resulting in approximately twofold up-regulation of telomerase [87].

Introduction of telomeric repeat amplification protocol (TRAP) assay has facilitated the detection of telomerase activity in tumor biopsy samples as well as cell lines [88]. Specificity of telomerase activity in malignant phenotype further enforces the reliability of this assay. The most important advantage of TRAP assay is its low detectable limit. TRAP assay has allowed the analysis of minimal tissue samples, such as fine-needle aspirates of the breast and thyroid, cervical smears, oral washings, and urine [89, 90]. Telomerase also has been used to detect circulating tumor cells also [85]. Newly emerged technique, droplet digital TRAP assay can detect telomerase activity even in a single cell [91]. However, the positive ratios of detection of telomerase vary in sedimented cells obtained from secretion, washing, brushing samples, etc. Electrochemical telomerase assay (ECTA) is another newly emerged technique to detect telomerase activity in biological samples [92]. It is comparatively simple and rapid PCR-free method. ECTA consists of a TS primer-immobilized electrode and ferrocenyl naphthalene diimide derivative as a tetraplex binder. This method has shown a high efficiency of telomerase detection in oral cancer biopsies [93]. Taken in account of all these reports, telomerase and its functionality can be utilized as a promising diagnostic and prognostic method in cancer.

7. Telomeres in prognosis

Better understanding of telomere structure and its dynamics focused the research on telomeres as biomarkers for several diseases especially in early detection and prognosis of cancers. A reduced telomere length in human hematopoietic tumors predicts a reduced survival time in patients suffering from myeloid leucaemia [94], chronic lymphocytic leucaemia [95], and myelodysplastic syndromes [96]. Although telomere length in solid tumors is suggested as a potential prognostic marker, patient survival rates vary with different cancer types. For example, a short telomere length in prostate cancer correlates with short disease-free interval and shorter overall survival time [97]. Analysis of telomere length of blood cells is also considered as predictive markers for pulmonary and esophageal neoplasia as well as of lymphoma in humans [98]. It has been suggested that the reduced TL in these patients reflects the effects of increased oxidative stress which correlates with cancer risk. Advent of latest technologies to measure relative and absolute telomere lengths has paved the way to use telomere length as diagnostic and prognostic markers. Telomere restriction fragment assay, qFISH, flow FISH, qPCR assay, single telomere length analysis (STELA), and dot-blot telomere assay are the currently available assay methods for telomere length [99].

8. Telomerase as drug target

Telomerase enzyme has recently emerged as an attractive target for cancer as it is a crucial factor required for the tumor immortalization of a subset of cells, including cancer stem cells. Studies have shown that 80–85% of the tumor cells express telomerase, whereas somatic cells lack the expression of telomerase [1]. In the present scenario, the major concern about the

chemotherapeutic approaches is the specificity of action and side effects of drugs on normal cells. Difference in telomere length and cell kinetics between normal and cancerous cells shows that targeting telomerase is an effective system to target cancer cells specifically [100]. Although telomerase is not considered as an oncogene, the expression of telomerase is the major reason for the transformation of a normal cell to cancer cell [80]. Compared to most other cancer targets, telomerase antagonists are advantageous due to the wide expression of this enzyme in cancer types [1]. Telomerase also possesses extra telomeric functions which are very crucial for tumor survival and homeostasis [101]. Studies have shown that telomerase-based cancer therapies are less likely to develop resistance against the drugs compared to drugs which target growth factor receptors or signal-transducing enzymes in cancer cells [2]. This ensures that cancer drugs based on telomerase inhibition are non-cytotoxic anticancer approach and have a broad therapeutic value.

9. Conclusion

Maintenance of telomere length in cancer cells is a critical factor in imparting the ability to undergo uncontrolled multiplication and thus immortality to the cells. It is imperative to discern the factors involved in telomere length preservation. Understanding the influence of telomerase and other factors in sustaining telomere length in cancer cells paves way for perceiving the theranostic role of telomere and telomerase in cancer treatment. Besides the theranostic effect, the possible side effects could be determined leading to precautionary methods to nullify the negative impact.

Author details

Angayarkanni Jayaraman^{1*}, Kalarikkal Gopikrishnan Kiran¹ and Palaniswamy Muthusamy²

*Address all correspondence to: angaibiotech@buc.edu.in

1 Department of Microbial Biotechnology, Bharathiar University, Coimbatore, India

2 Department of Microbiology, Karpagam University, Coimbatore, India

References

- [1] Flores I, Benetti R, Blasco MA. Telomerase regulation and stem cell behaviour. *Current Opinion in Cell Biology*. 2006;18:254–60. doi:10.1016/j.ceb.2006.03.003
- [2] Harley CB. Telomerase and cancer therapeutics. *Nature Reviews Cancer*. 2008;8:167–79. doi:10.1038/nrc2275

- [3] Longhese MP. DNA damage response at functional and dysfunctional telomeres. *Genes and Development*. 2008;22:125–40. doi:10.1101/gad.1626908
- [4] Mu J, Wei LX. Telomere and telomerase in oncology. *Cell Research*. 2002;12:1–7. doi:10.1038/sj.cr.7290104
- [5] Griffith JD, Comeau L, Rosenfield S, et al. Mammalian telomeres end in a large duplex loop. *Cell*. 1999;97:503–14.
- [6] O'Sullivan RJ, Karlseder J. Telomeres: protecting chromosomes against genome instability. *Nature Reviews Molecular Cell Biology*. 2010;11:171–81. doi:10.1038/nrm2848
- [7] Parkinson GN, Lee MP, Neidle S. Crystal structure of parallel quadruplexes from human telomeric DNA. *Nature*. 2002;417:876–80. doi:10.1038/nature755
- [8] Oganessian L, Bryan TM. Physiological relevance of telomeric G-quadruplex formation: a potential drug target. *BioEssays: News and Reviews in Molecular, Cellular and Developmental Biology*. 2007;29:155–65. doi:10.1002/bies.20523
- [9] Rhodes D, Giraldo R. Telomere structure and function. *Current Opinion in Structural Biology*. 1995;5:311–22.
- [10] Palm W, de Lange T. How shelterin protects mammalian telomeres. *Annual Review of Genetics*. 2008;42:301–34. doi:10.1146/annurev.genet.41.110306.130350
- [11] Billaud T, Brun C, Ancelin K, Koering CE, Laroche T, Gilson E. Telomeric localization of TRF2, a novel human telobox protein. *Nature Genetics*. 1997;17:236–9. doi:10.1038/ng1097-236
- [12] Chong L, van Steensel B, Broccoli D, et al. A human telomeric protein. *Science (New York, NY)*. 1995;270:1663–7.
- [13] Broccoli D, Smogorzewska A, Chong L, de Lange T. Human telomeres contain two distinct Myb-related proteins, TRF1 and TRF2. *Nature Genetics*. 1997;17:231–5. doi:10.1038/ng1097-231
- [14] Court R, Chapman L, Fairall L, Rhodes D. How the human telomeric proteins TRF1 and TRF2 recognize telomeric DNA: a view from high-resolution crystal structures. *EMBO Reports*. 2005;6:39–45. doi:10.1038/sj.embor.7400314
- [15] Kim NW, Piatyszek MA, Prowse KR, et al. Specific association of human telomerase activity with immortal cells and cancer. *Science (New York, NY)*. 1994;266:2011–5.
- [16] Baumann P, Cech TR. POT1, the putative telomere end-binding protein in fission yeast and humans. *Science (New York, NY)*. 2001;292:1171–5. doi:10.1126/science.1060036

- [17] Houghtaling BR, Cuttonaro L, Chang W, Smith S. A dynamic molecular link between the telomere length regulator TRF1 and the chromosome end protector TRF2. *Current Biology*. 2004;14:1621–31. doi:10.1016/j.cub.2004.08.052
- [18] Liu D, O'Connor MS, Qin J, Songyang Z. Telosome, a mammalian telomere-associated complex formed by multiple telomeric proteins. *The Journal of Biological Chemistry*. 2004;279:51338–42. doi:10.1074/jbc.M409293200
- [19] Ye JZ, Hockemeyer D, Krutchinsky AN, et al. POT1-interacting protein PIP1: a telomere length regulator that recruits POT1 to the TIN2/TRF1 complex. *Genes and Development*. 2004;18:1649–54. doi:10.1101/gad.1215404
- [20] Nandakumar J, Cech TR. Finding the end: recruitment of telomerase to telomeres. *Nature Reviews Molecular Cell Biology*. 2013;14:69–82. doi:10.1038/nrm3505
- [21] Verdun RE, Karlseder J. Replication and protection of telomeres. *Nature*. 2007;447:924–31. doi:10.1038/nature05976
- [22] Bianchi A, Smith S, Chong L, Elias P, de Lange T. TRF1 is a dimer and bends telomeric DNA. *EMBO Journal*. 1997;16:1785–94. doi:10.1093/emboj/16.7.1785
- [23] Martinez P, Thanasoula M, Munoz P, et al. Increased telomere fragility and fusions resulting from TRF1 deficiency lead to degenerative pathologies and increased cancer in mice. *Genes and Development*. 2009;23:2060–75. doi:10.1101/gad.543509
- [24] Sfeir A, Kosiyatrakul ST, Hockemeyer D, et al. Mammalian telomeres resemble fragile sites and require TRF1 for efficient replication. *Cell*. 2009;138:90–103. doi:10.1016/j.cell.2009.06.021
- [25] Amiard S, Doudeau M, Pinte S, et al. A topological mechanism for TRF2-enhanced strand invasion. *Nature Structural and Molecular Biology*. 2007;14:147–54. doi:10.1038/nsmb1192
- [26] De Lange T. Protection of mammalian telomeres. *Oncogene*. 2002;21:532–40. doi:10.1038/sj.onc.1205080
- [27] Stansel RM, de Lange T, Griffith JD. T-loop assembly in vitro involves binding of TRF2 near the 3' telomeric overhang. *EMBO Journal*. 2001;20:5532–40. doi:10.1093/emboj/20.19.5532
- [28] Denchi EL, de Lange T. Protection of telomeres through independent control of ATM and ATR by TRF2 and POT1. *Nature*. 2007;448:1068–71. doi:10.1038/nature06065
- [29] Smogorzewska A, Karlseder J, Holtgreve-Grez H, Jauch A, de Lange T. DNA ligase IV-dependent NHEJ of deprotected mammalian telomeres in G1 and G2. *Current Biology*. 2002;12:1635–44.
- [30] Benetti R, Schoeftner S, Munoz P, Blasco MA. Role of TRF2 in the assembly of telomeric chromatin. *Cell Cycle (Georgetown, TX)*. 2008;7:3461–8. doi:10.4161/cc.7.21.7013

- [31] Celli GB, de Lange T. DNA processing is not required for ATM-mediated telomere damage response after TRF2 deletion. *Nature Cell Biology*. 2005;7:712–8. doi:10.1038/ncb1275
- [32] Karlseder J, Broccoli D, Dai Y, Hardy S, de Lange T. p53-and ATM-dependent apoptosis induced by telomeres lacking TRF2. *Science (New York, NY)*. 1999;283:1321–5.
- [33] Loayza D, De Lange T. POT1 as a terminal transducer of TRF1 telomere length control. *Nature*. 2003;423:1013–8. doi:10.1038/nature01688
- [34] Wang F, Podell ER, Zaug AJ, et al. The POT1-TPP1 telomere complex is a telomerase processivity factor. *Nature*. 2007;445:506–10. doi:10.1038/nature05454
- [35] Xin H, Liu D, Wan M, et al. TPP1 is a homologue of ciliate TEBP-beta and interacts with POT1 to recruit telomerase. *Nature*. 2007;445:559–62. doi:10.1038/nature05469
- [36] Benarroch-Popivker D, Pisano S, Mendez-Bermudez A, et al. TRF2-mediated control of telomere DNA topology as a mechanism for chromosome-end protection. *Molecular Cell*. 2016;61:274–86. doi:10.1016/j.molcel.2015.12.009
- [37] Bae NS, Baumann P. A RAP1/TRF2 complex inhibits nonhomologous end-joining at human telomeric DNA ends. *Molecular Cell*. 2007;26:323–34. doi:10.1016/j.molcel.2007.03.023
- [38] Sarthy J, Bae NS, Scraftford J, Baumann P. Human RAP1 inhibits non-homologous end joining at telomeres. *EMBO Journal* 2009;28:3390–9. doi:10.1038/emboj.2009.275
- [39] Huffman KE, Levene SD, Tesmer VM, Shay JW, Wright WE. Telomere shortening is proportional to the size of the G-rich telomeric 3'-overhang. *The Journal of Biological Chemistry*. 2000;275:19719–22. doi:10.1074/jbc.M002843200
- [40] Olovnikov AM. A theory of marginotomy: the incomplete copying of template margin in enzymic synthesis of polynucleotides and biological significance of the phenomenon. *Journal of Theoretical Biology*. 1973;41:181–90.
- [41] Watson JD. Origin of concatemeric T7 DNA. *Nature: New Biology*. 1972;239:197–201.
- [42] Harley CB, Futcher AB, Greider CW. Telomeres shorten during ageing of human fibroblasts. *Nature*. 1990;345:458–60. doi:10.1038/345458a0
- [43] Vaziri H, Schachter F, Uchida I, et al. Loss of telomeric DNA during aging of normal and trisomy 21 human lymphocytes. *American Journal of Human Genetics*. 1993;52:661–7.
- [44] Greider CW, Blackburn EH. Identification of a specific telomere terminal transferase activity in *Tetrahymena* extracts. *Cell*. 1985;43:405–13.
- [45] Liu T, Chung MJ, Ullenbruch M, et al. Telomerase activity is required for bleomycin-induced pulmonary fibrosis in mice. *Journal of Clinical Investigation*. 2007;117:3800–9. doi:10.1172/jci32369

- [46] Hug N, Lingner J. Telomere length homeostasis. *Chromosoma*. 2006;115:413–25. doi:10.1007/s00412-006-0067-3
- [47] Bechter OE, Eisterer W, Dlaska M, Kuhr T, Thaler J. CpG island methylation of the hTERT promoter is associated with lower telomerase activity in B-cell lymphocytic leukemia. *Experimental Hematology*. 2002;30:26–33.
- [48] Collins K. Ciliate telomerase biochemistry. *Annual Review of Biochemistry*. 1999;68:187–218. doi:10.1146/annurev.biochem.68.1.187
- [49] Greider CW. Telomerase activity, cell proliferation, and cancer. *Proceedings of the National Academy of Sciences of the United States of America*. 1998;95:90–2.
- [50] Greider CW. Telomeres do D-loop-T-loop. *Cell*. 1999;97:419–22.
- [51] Cong YS, Wen J, Bacchetti S. The human telomerase catalytic subunit hTERT: organization of the gene and characterization of the promoter. *Human Molecular Genetics*. 1999;8:137–42.
- [52] Gu J, Andreeff M, Roth JA, Fang B. hTERT promoter induces tumor-specific Bax gene expression and cell killing in syngenic mouse tumor model and prevents systemic toxicity. *Gene Therapy*. 2002;9:30–7. doi:10.1038/sj.gt.3301619
- [53] Avilion AA, Piatyszek MA, Gupta J, Shay JW, Bacchetti S, Greider CW. Human telomerase RNA and telomerase activity in immortal cell lines and tumor tissues. *Cancer Research*. 1996;56:645–50.
- [54] Yi X, Tesmer VM, Savre-Train I, Shay JW, Wright WE. Both transcriptional and posttranscriptional mechanisms regulate human telomerase template RNA levels. *Molecular and Cellular Biology*. 1999;19:3989–97.
- [55] Chan SW, Blackburn EH. New ways not to make ends meet: telomerase, DNA damage proteins and heterochromatin. *Oncogene*. 2002;21:553–63. doi:10.1038/sj.onc.1205082
- [56] Bagheri S, Nosrati M, Li S, et al. Genes and pathways downstream of telomerase in melanoma metastasis. *Proceedings of the National Academy of Sciences of the United States of America*. 2006;103:11306–11. doi:10.1073/pnas.0510085103
- [57] Hoffmeyer K, Raggioli A, Rudloff S, et al. Wnt/beta-catenin signaling regulates telomerase in stem cells and cancer cells. *Science (New York, NY)*. 2012;336:1549–54. doi:10.1126/science.1218370
- [58] Li S, Crothers J, Haqq CM, Blackburn EH. Cellular and gene expression responses involved in the rapid growth inhibition of human cancer cells by RNA interference-mediated depletion of telomerase RNA. *Journal of Biological Chemistry*. 2005;280:23709–17. doi:10.1074/jbc.M502782200
- [59] Maida Y, Yasukawa M, Furuuchi M, et al. An RNA-dependent RNA polymerase formed by TERT and the RMRP RNA. *Nature*. 2009;461:230–5. doi:10.1038/nature08283

- [60] Park JI, Venteicher AS, Hong JY, et al. Telomerase modulates Wnt signalling by association with target gene chromatin. *Nature*. 2009;460:66–72. doi:10.1038/nature08137
- [61] Smith LL, Collier HA, Roberts JM. Telomerase modulates expression of growth-controlling genes and enhances cell proliferation. *Nature Cell Biology*. 2003;5:474–9. doi:10.1038/ncb985
- [62] Ahmed S, Passos JF, Birket MJ, et al. Telomerase does not counteract telomere shortening but protects mitochondrial function under oxidative stress. *Journal of Cell Science*. 2008;121:1046–53. doi:10.1242/jcs.019372
- [63] Haendeler J, Hoffmann J, Brandes RP, Zeiher AM, Dimmeler S. Hydrogen peroxide triggers nuclear export of telomerase reverse transcriptase via Src kinase family-dependent phosphorylation of tyrosine 707. *Molecular and Cellular Biology*. 2003;23:4598–610.
- [64] Haendeler J, Droese S, Buchner N, et al. Mitochondrial telomerase reverse transcriptase binds to and protects mitochondrial DNA and function from damage. *Arteriosclerosis, Thrombosis, and Vascular Biology*. 2009;29:929–35. doi:10.1161/atvbaha.109.185546
- [65] Indran IR, Hande MP, Pervaiz S. hTERT overexpression alleviates intracellular ROS production, improves mitochondrial function, and inhibits ROS-mediated apoptosis in cancer cells. *Cancer Research*. 2011;71:266–76. doi:10.1158/0008-5472.can-10-1588
- [66] Santos JH, Meyer JN, Skorvaga M, Annab LA, Van Houten B. Mitochondrial hTERT exacerbates free-radical-mediated mtDNA damage. *Aging Cell*. 2004;3:399–411. doi:10.1111/j.1474-9728.2004.00124.x
- [67] Santos JH, Meyer JN, Van Houten B. Mitochondrial localization of telomerase as a determinant for hydrogen peroxide-induced mitochondrial DNA damage and apoptosis. *Human Molecular Genetics*. 2006;15:1757–68. doi:10.1093/hmg/ddl098
- [68] Singhapol C, Pal D, Czapiewski R, Porika M, Nelson G, Saretzki GC. Mitochondrial telomerase protects cancer cells from nuclear DNA damage and apoptosis. *PloS one*. 2013;8:e52989. 1-11 doi:10.1371/journal.pone.0052989
- [69] Wright WE, Piatyszek MA, Rainey WE, Byrd W, Shay JW. Telomerase activity in human germline and embryonic tissues and cells. *Developmental Genetics*. 1996;18:173–9. doi:10.1002/(sici)1520-6408(1996)18:2<173::aid-dvg10>3.0.co;2-3
- [70] Shay JW, Bacchetti S. A survey of telomerase activity in human cancer. *European Journal of Cancer (Oxford, England: 1990)*. 1997;33:787–91. doi:10.1016/s0959-8049(97)00062-2
- [71] Stewart SA, Hahn WC, O'Connor BF, et al. Telomerase contributes to tumorigenesis by a telomere length-independent mechanism. *Proceedings of the National Academy of Sciences of the United States of America*. 2002;99:12606–11. doi:10.1073/pnas.182407599

- [72] Hiyama E, Hiyama K, Yokoyama T, et al. Rapid detection of MYCN gene amplification and telomerase expression in neuroblastoma. *Clinical Cancer Research: An Official Journal of the American Association for Cancer Research*. 1999;5:601–9.
- [73] Poremba C, Scheel C, Hero B, et al. Telomerase activity and telomerase subunits gene expression patterns in neuroblastoma: a molecular and immunohistochemical study establishing prognostic tools for fresh-frozen and paraffin-embedded tissues. *Journal of Clinical Oncology: Official Journal of the American Society of Clinical Oncology*. 2000;18:2582–92.
- [74] Yasui W, Tahara H, Tahara E, et al. Expression of telomerase catalytic component, telomerase reverse transcriptase, in human gastric carcinomas. *Japanese Journal of Cancer Research: Gann*. 1998;89:1099–103.
- [75] Gupta J, Han LP, Wang P, Gallie BL, Bacchetti S. Development of retinoblastoma in the absence of telomerase activity. *Journal of the National Cancer Institute*. 1996;88:1152–7.
- [76] Langford LA, Piatyszek MA, Xu R, Schold SC, Jr., Shay JW. Telomerase activity in human brain tumours. *Lancet (London, England)*. 1995;346:1267–8.
- [77] Klingelhutz AJ, Foster SA, McDougall JK. Telomerase activation by the E6 gene product of human papillomavirus type 16. *Nature*. 1996;380:79–82. doi:10.1038/380079a0
- [78] Kyo S, Takakura M, Kohama T, Inoue M. Telomerase activity in human endometrium. *Cancer Research*. 1997;57:610–4.
- [79] Tanaka M, Kyo S, Takakura M, et al. Expression of telomerase activity in human endometrium is localized to epithelial glandular cells and regulated in a menstrual phase-dependent manner correlated with cell proliferation. *The American Journal of Pathology*. 1998;153:1985–91. doi:10.1016/s0002-9440(10)65712-4
- [80] Cong Y-S, Wright WE, Shay JW. Human telomerase and its regulation. *Microbiology and Molecular Biology Reviews*. 2002;66:407–25. doi:10.1128/MMBR.66.3.407-425.2002
- [81] Fuxe J, Akusjarvi G, Goike HM, Roos G, Collins VP, Pettersson RF. Adenovirus-mediated overexpression of p15INK4B inhibits human glioma cell growth, induces replicative senescence, and inhibits telomerase activity similarly to p16INK4A. *Cell Growth and Differentiation: The Molecular Biology Journal of the American Association for Cancer Research*. 2000;11:373–84.
- [82] Henderson YC, Breau RL, Liu TJ, Clayman GL. Telomerase activity in head and neck tumors after introduction of wild-type p53, p21, p16, and E2F-1 genes by means of recombinant adenovirus. *Head & Neck*. 2000;22:347–54.
- [83] Xu D, Erickson S, Szeps M, et al. Interferon alpha down-regulates telomerase reverse transcriptase and telomerase activity in human malignant and nonmalignant hematopoietic cells. *Blood*. 2000;96:4313–8.

- [84] Hande MP, Balajee AS, Natarajan AT. Induction of telomerase activity by UV-irradiation in Chinese hamster cells. *Oncogene*. 1997;15:1747–52. doi:10.1038/sj.onc.1201327
- [85] Xu Y, He K, Goldkorn A. Telomerase targeted therapy in cancer and cancer stem cells. *Clinical Advances in Hematology and Oncology*. 2011;9:442–55.
- [86] Blasco MA, Rizen M, Greider CW, Hanahan D. Differential regulation of telomerase activity and telomerase RNA during multi-stage tumorigenesis. *Nature Genetics*. 1996;12:200–4. doi:10.1038/ng0296-200
- [87] Horn S, Figl A, Rachakonda PS, et al. TERT promoter mutations in familial and sporadic melanoma. *Science (New York, NY)*. 2013;339:959–61. doi:10.1126/science.1230062
- [88] Fajkus J. Detection of telomerase activity by the TRAP assay and its variants and alternatives. *Clinica Chimica Acta; International Journal of Clinical Chemistry*. 2006;371:25–31. doi:10.1016/j.cca.2006.02.039
- [89] Hiyama K, Ishioka S, Shay JW, et al. Telomerase activity as a novel marker of lung cancer and immune-associated lung diseases. *International Journal of Molecular Medicine*. 1998;1:545–9.
- [90] Sugino T, Yoshida K, Bolodeoku J, et al. Telomerase activity in human breast cancer and benign breast lesions: diagnostic applications in clinical specimens, including fine needle aspirates. *International Journal of Cancer*. 1996;69:301–6. doi:10.1002/(sici)1097-0215(19960822)69:4<301::aid-ijc11>3.0.co;2-8
- [91] Ludlow AT, Robin JD, Sayed M, et al. Quantitative telomerase enzyme activity determination using droplet digital PCR with single cell resolution. *Nucleic Acids Research*. 2014;42:e104-e. 1-12 doi:10.1093/nar/gku439
- [92] Sato S, Kondo H, Nojima T, Takenaka S. Electrochemical telomerase assay with ferrocenylnaphthalene diimide as a tetraplex DNA-specific binder. *Analytical Chemistry*. 2005;77:7304–9. doi:10.1021/ac0510235
- [93] Hayakawa M, Kodama M, Sato S, et al. Electrochemical telomerase assay for screening for oral cancer. *British Journal of Oral and Maxillofacial Surgery*. 2016;54:301–5. doi:10.1016/j.bjoms.2016.01.003
- [94] Hartmann U, Brummendorf TH, Balabanov S, Thiede C, Illme T, Schaich M. Telomere length and hTERT expression in patients with acute myeloid leukemia correlates with chromosomal abnormalities. *Haematologica*. 2005;90:307–16.
- [95] Boultonwood J, Fidler C, Shepherd P, et al. Telomere length shortening is associated with disease evolution in chronic myelogenous leukemia. *American Journal of Hematology*. 1999;61:5–9.

- [96] Ohyashiki JH, Ohyashiki K, Fujimura T, et al. Telomere shortening associated with disease evolution patterns in myelodysplastic syndromes. *Cancer Research*. 1994;54:3557–60.
- [97] Donaldson L, Fordyce C, Gilliland F, et al. Association between outcome and telomere DNA content in prostate cancer. *Journal of Urology*. 1999;162:1788–92.
- [98] Svenson U, Roos G. Telomere length as a biological marker in malignancy. *Biochimica et Biophysica Acta*. 2009;1792:317–23. doi:10.1016/j.bbadis.2009.01.017
- [99] Nussey DH, Baird D, Barrett E, et al. Measuring telomere length and telomere dynamics in evolutionary biology and ecology. *Methods in Ecology and Evolution/British Ecological Society*. 2014;5:299–310. doi:10.1111/2041-210x.12161
- [100] Zhang X, Mar V, Zhou W, Harrington L, Robinson MO. Telomere shortening and apoptosis in telomerase-inhibited human tumor cells. *Genes and Development*. 1999;13:2388–99.
- [101] Martinez P, Blasco MA. Telomeric and extra-telomeric roles for telomerase and the telomere-binding proteins. *Nature Reviews Cancer*. 2011;11:161–76. doi:10.1038/nrc3025