We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Phenolic Compounds from the Natural Sources and Their Cytotoxicity

Shagufta Perveen and Areej Mohammad Al-Taweel

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/66898>

Abstract

Natural phenolic compounds are considered as one of the important secondary metabolites for their chemopreventive and chemotherapeutic effects in cancer. These compounds show potent activities for cancer prevention and its treatment. There are many phenolic compounds present in medicinal and edible plants such as, flavonoids, bioflavonoids, stilbenes, chalconoids, chromones, phenylpropanoids, curcuminoids, coumarins, tannins, lignans, neolignans, anthraquinones, quinones, xanthones, phenolic acids and their glycosides and many more. The antioxidant potential of phenolic compounds is almost bolded in the treatment and prevention of cancer. Mono phenolic, polyphenolic and phenolic acids compounds from a large variety of plants, foods, spices, insects, fungus, beverages, lichens, algae and mammals have been shown to inhibit or attenuate the initiation, progression and spread of cancers in cells in vitro and in animals in vivo. In this chapter, we try to cover general view and the recent literature to summarize structural information and cytotoxic effects of phenolic compounds on different cancer cell lines from medicinal herbs and plants.

Keywords: natural phenolic constituents, cytotoxic activities, cancer cell lines

1. Introduction

Natural products offer opportunities for innovation in drug discovery and play a major role for cancer cure. A considerable number of antitumor agents currently used in the clinic are of natural origin. For instance, over half of all anticancer prescription drugs approved internationally between the 1940s and 2006 were natural products or their derivatives. Among

them, plants have been the chief source of natural compounds used for medicine [1]. Many traditional and folk medicinal plants have been use for cancer therapy throughout the world and they work very well for the prevention of cancers. Phenolic compounds play very important role for the treatment and prevention on different types of cancers which are a major health problem around the world. These secondary metabolites associated with the health benefits of human derived from consuming high levels of fresh and dried fruits and vegetables. These compounds with hydroxyl bearing aromatic ring skeleton exhibit wide range of different biological activities including: anti-inflammatory, antioxidant, cytotoxicity, anti-microbial, anti-allergic. According to WHO, 8.2 million people die each year from cancer and it is estimated 13% of all deaths worldwide [2]. In this chapter, we discuss the most recent literature on phenolic constituents from natural sources and their anticancer activities on different cancer cell lines (**Table 1**).

Source	Compound name	Class of compounds	Cytotoxic activity	References
<i>Artocarpus heterophyllus</i>	Artocarpin (1), cudraflavone C (2), 6-prenylapigenin (3), kuwanon C (4), norartocarpin (5), albanin A (6), cudraflavone B (7)	Isoprenoid-substituted flavonoids	B16 melanoma cells	[3]
Apple, grapes, tomato green tea, pine and many other	Kaempferol (8)	Flavonoid	p53 or PLK-1, in MCF-7 breast cancer and HeLa cervical cancer cells, U-2 OS human osteosarcoma cell, A549 lung cancer cell, Miapaca-2, Panc-1 and SNU-213 human pancreatic cancer cells	[4]
<i>Dorstenia mannii</i>	Dorsmanin F (9)	Flavanone	LeukaemiaCCRF-CEM, MDA-MB-231-BCRP, CEM/ADR5000 cells	[5]
<i>Morus mesozygia</i>	Artocarpesin (10), cycloartocarpesin (11)	Flavonoid	Induced apoptosis in CCRF-CEM leukemia cells	[6]
Oriental tobacco, <i>Nicotiana tabacum</i>	6,7-dimethoxy-4'hydroxy-8-formylflavon (12), 4',7-dihydroxy-8-formyl-6-methoxyflavon (13)	Flavonoid	Human tumor (NB4, A549, SHSY5Y, PC3 and MCF7) cell lines	[7]
<i>Sophora flavescens</i>	Isoxanthohumol (14)	Flavonoid	Breast cancer (MCF-7), ovarian cancer (A-2780), prostate cancer (DU145 and PC-3) and colon cancer (HT-29 and SW620) cells, human cytochrome P450 (CYP1A2).	[8]
Kushen <i>Sophora flavescens</i>	Kushecarpin D (15)	Novel flavonoid	Human umbilical vein endothelial cell line (ECV304), antiangiogenic activity, together with its antiproliferative effect on endothelial cells without causing apoptosis	[9]
<i>Scutellaria barbata</i>	Luteolin (16)	Flavonoid	HepG2 and Bel7402 cells, human hepatocellular carcinoma cells, prostate cancer (PCa), assessing the PC3 and LNCaP cells, MCF-7 human breast cancer cells, PC12 cell line	[10–12]

Source	Compound name	Class of compounds	Cytotoxic activity	References
Propolis and in honey	Chrysin (17)	Flavonoid	Oropharyngeal KB, mammary LM3, anaplastic thyroid KAT18, anaplastic thyroid HTh7, pancreatic PANC-1, liver H22, gastric SGC-7901, colon HT-29, cervical Hela, melanoma A375, oesophageal OE33, lung A549, colon DLD-1, rectal SW837, breast MDA-MB-231, glioma U87-MG, esophageal squamous KYSE-510, leukemia U937, prostate PC-3, hepatocellular HepG2, acute T-lymphoblastic leukemia CEM, neuroblastoma SH-SY5Y, squamous cell carcinoma FaDu, breast MCF-7, oral SCC-9, prostate DU-145, leukemia K562, cell line PC12	[13]
<i>Macrothelypteris torresiana</i>	DICO (18)	Novel nonaromatic B-ring flavonoid	Human hepatoma HepG2 cells	[14]
Mulberry tree <i>Morus</i> species	Morusin (19)	Prenylated flavonoid	Human hepatoma SK-Hep1 cells	[15]
<i>Artocarpus communis</i>	Artocarpin (20)	Prenylated flavonoid	HepG2 and PLC/PRF/5 hepatoma cells, human cutaneous SCC cell line HSC-1, human T47D breast cancer cells	[16, 17]
<i>Daphne genkwa</i> Sieb	Genkwanin (21)	Flavonoid	HT-29 and SW-480 human colorectal cancer cells	[18]
Strawberries, apples, persimmons, grapes, onions and cucumbers	Fisetin (22)	Flavonoid	Human malignant melanoma cells A375 (CLR- 1619) and RPMI-7951 (HTB-66), SK-MEL-28 cells, A375 and SK-MEL-28 cells, RPMI-7951 cells	[19]
<i>Clerodendrum inerme</i> and various medicinal herbs	Hispidulin (23)	Flavonoid	Human HepG2 hepatocarcinoma cell line and the mouse L929 fibroblast cell line, human HEK293 fibroblast cell lines, human renal cancer cell (HRCC) lines 786-0 and Caki-1, PANC-1 cancer cells from human pancreas	[20, 21]
Fruits, vegetables, leaves and grains	Quercetin (24)	Flavonoid	HepG2 cells, Human GL-15 glioblastoma cells, breast cancer cells MCF-7 and MCF-7/dox	[22, 23]
All citrus fruits and <i>Cordia obliqua</i>	Hesperetin (25)	Flavonoid	Human esophageal squamous cell carcinoma Eca109 cells, human breast cancer MCF-7 cells, prostate cancer PC-3 cells, HT-29 human colon adenocarcinoma cell, human cervical cancer cell line HeLa cells	[24]
<i>Hypericum perforatum</i>	Apigenin (26)	Flavonoid	MCF-7 breast carcinoma cells, breast tumor cell line MDA-MB231, human cervical carcinoma HeLa cells, human colon carcinoma cell, human leukemia cells, A549 lung cancer cell, human ovarian carcinoma HO-8910PM cells, human prostate cancer PC-3 cell, thyroid carcinoma cell lines, UCLA NPA-87-1 (NPA), cell line PC12	[25, 26]

Source	Compound name	Class of compounds	Cytotoxic activity	References
<i>Azadirachta indica</i>	Quercetin-3-O- β -D-glucopyranoside (27)	Flavonoid	Protect cells from H ₂ O ₂ -induced cytotoxicity	[27]
<i>Polygonum amplexicaule</i>	Amplexicaule A (28)	Flavonoid glycoside	Human breast cancer cell lines MCF-7 and MDA-MB-435	[28]
Wheat bran	Triticuside A (29)	Flavonoid glycoside	Human breast cancer cells (MCF-7 and MDA-MB-231)	[29]
Citrus fruits	Naringin (30)	Flavonoid diglycoside	Breast cancer (TNBC), MDA-MB-231, MDA-MB-468 and BT-549 cells, K562 (human leukemia cell line), Raji (human Burkitt's lymphoma cell line) and NK-92MI	[30, 31]
Filamentous bacterium streptomyces	Quercetin-3-O- α -L-rhamnopyranosyl-(1 $\rightarrow$ 6)- β -D-glucopyranoside (31)	Flavonoid diglycoside	Human lung cancer A549 cells through p53 and cytochrome c	[32]
<i>Cirsium setidens</i> , <i>Aster scaber</i>	Pectolinarin (32) and astragalin (33)	Flavonoid glycoside	Human brain neuroblastoma SK-N-SH cells	[33]
<i>Sophora japonica</i>	Troxeutin (34)	Flavonoid diglycoside	Human prostate cancer radioresistant (DU145) and radiosensitive (PC3) cells	[34]
<i>Bupleurum flavum</i> , <i>Artemisia capillaris</i>	Vicenin (35)	Flavonoid glycoside	Hepatocellular carcinoma HEP-G2 cells, human prostate carcinoma LNCaP, PC-3 & DU-145	[35, 36]
Soybean	Genistein (36)	Isoflavone	CIP2A in MCF-7-C3 and T47D breast cancer cells, human prostate cancer cell PC3-M, PC3 and DU-145 human PCa cell lines, sarcomatoid mammary carcinoma cell line F3II, B16F0 melanoma cell line	[37, 38]
Soybean	Daidzein (37)	Isoflavone	LnCaP, DU145 and PC3 human prostate cancer cell lines, MCF-7 breast cancer cell	[39]
<i>Ateleia glazioviana</i>	Biochanin A (38)	Isoflavone	Pancreatic cancer cells (Panc1 and AsPC-1)	[40]
<i>Erythrina stricta</i> , <i>E. variegata</i>	Alpinum isoflavone (39)	Isoflavon	Leukemia CEM/ADR5000 cells, drug-resistant breast adenocarcinoma MDA-MB-231-BCRP cells, CEM/ADR5000 cells, colon carcinoma HCT116	[41]
<i>Pueraria lobata</i>	Puerarin (40)	Isoflavone-C-glucoside	Myeloid leukemia cell lines, U937, Kasumi-1, HL-60 and NB4 cells, breast cancer MCF-7/adriamycin (MCF-7/adr) cells, colon cancer HT-29 cell	[42]
<i>Toxicodendron vernicifluum</i>	Butein (41)	Chalcone	Human colon adenocarcinoma cell line 220.1, human leukemia cells HL-60, HER2 ⁺ HCC-1419, HCC-2218 and SKBR-3 breast cancer cells, human PCa (LNCaP, CWR22Rv1 and PC-3), human uveal melanoma cell lines (M17, SP6.5 and C918), HeLa human cervical cancer cell line	[43–47]

Source	Compound name	Class of compounds	Cytotoxic activity	References
<i>Spatholobus suberectus</i>	Isoliquiritigenin (42)	Chalcone	Human leukemia cells HL-60 cell, human U373 glioblastoma cells, human hepatoma HepG2) cells, DU145 human prostate cancer cells	[48]
<i>Eugenia aquea</i>	2',4'-dihydroxy-6-methoxy-3,5-dimethylchalcone (43)	Chalcone	Human cell lines liver cancer SMMC-7721 cells, pancreas cancer 8898 cells, tumor of cervix uteri HeLa cells, lung cancer SPC-A-1 cells, high metastatic lung carcinoma 95-D cells and gall bladder carcinoma GBC-SD cells	[49, 50]
<i>Polygonum limbatum</i>	4'-hydroxy-2',6'-dimethoxychalcone (44)	Chalcone	MDA-MB-231-pcDNA3 breast cancer, HCT116 (p53+/+) colon cancer cells(-), the U87MG glioblastoma cells	[51]
<i>Humulus lupulus</i>	Xanthohumol (45)	Prenylated Chalcone	Human breast cancer (MCF-7), colon cancer (HT-29) and ovarian cancer (A-2780) cells	[52, 53]
<i>Piper methysticum</i>	Flavokawain B (46)	Chalcone	Breast cancer cell lines, MCF-7 and MDA-MB231, HCT116 human colon cancer	[54]
<i>Artocarpus communis</i>	Isolespeol (47)	Chalcone	Liver cell line Hep3B, PLC5, Huh7, human colon cancer HT-29, COLO205 cell, SW 872 human liposarcoma cell	[55]
<i>Ashitaba (Angelica keiskei)</i>	Xanthoangelol (48), 4-hydroxyderricin (49)	Chalcone	Human neuroblastoma (IMR-32) and leukemia (Jurkat) cells, SW 872, HT-29, COLO205, Hep3B, PLC5, Huh7, HepG2, KATO III, human tumor cell lines, HL60 (leukemia), CRL1579 (melanoma), A549 (lung) and AZ521 (stomach)	[56, 57]
<i>Dorstenia barteri, Psoralea corylifolia</i>	Isobavachalcone (50)	Chalcone	CCRF-CEM leukemia cells, human neuroblastoma cell lines (IMR-32 and NB-39), Induced apoptosis in CCRF-CEM leukemia cells	[58]
<i>Dorstenia poinsettifolia</i>	Poinsettifolin B (51)	Chalcone	Leukaemia CCRF-CEM, MDA-MB-231-BCRP, M/ADR5000 cells	[59]
<i>Citrus kinokuni</i>	2'-hydroxy-3,4,4', 5,6'-pentamethoxychalcone (52)	Chalcone	Human breast adenocarcinoma (MCF-7), non-small cell lung cancer (NCI-H460) and melanoma (A375-C5)	[60]
<i>Rhuspyroides</i>	Rhuschalcones II-VI (53, 54) (53-57)	Bichalcone	HT29 and HCT-116 colon tumor cell lines	[61]
<i>Hibiscus syriacus</i>	Syriacusin A (58)	Naphthalene	Human skin fibroblast cells (CRL-2076), human tumor cell lines UACC62, ACHN, SW620 and SF539	[62]
<i>Rumex nepalensis</i>	Rumexneposide A (59)	Naphthalene acylglucoside	A549 (human non-small cell lung cancer), SKBR3, MCF-10A, MCF-7 and H522 cells (nonsmall-cell lung carcinoma)	[63]
<i>Pentas parvifolia and Pentas bussei</i>	Parvinaphthol B (60), parvinaphthols C (61)	Naphthalene	MDA-MB-231 human triple-negative breast cancer cell line	[64]

Source	Compound name	Class of compounds	Cytotoxic activity	References
<i>Rumex dentatus</i>	Chrysophanol (62)	Anthraquinone	MCF-7 breast cancer cell line, gastric cancer 7901 cells, Melanoma A375 cells and oophoroma SKOV-3 cells	[65]
<i>Streptomyces</i> sp. ERINLG-26	9,10-anthraquinone (63), 2-hydroxy-9,10-anthraquinone (64), 2,3-dihydroxy-9,10-anthraquinone (65)	Anthraquinone	A549 lung adenocarcinoma and COLO320 cancer cell line	[66–68]
<i>Rheum palmatum</i>	Chrysophanol 8-O-beta-(6'-acetyl) glucopyranoside (66)	Anthraquinone glucoside	Human oral squamous cell carcinoma (HSC-2) and salivary gland tumor (HSG) cell lines than against normal human gingival fibroblasts (HGF)	[69]
<i>Cratoxylum maingayi</i> and <i>C. cochinchinense</i>	Formoxanthone C (67)	Xanthone	Human lung cancer (NCI-H187), MCF-7 (breast adenocarcinoma), KB (human oral cancer), HeLa (human cervical cancer) and HT-29 (colon cancer)	[70, 71]
Mangrove fungus <i>Phomopsis longicolla</i> , Fungus <i>Phomopsis longicolla</i> , <i>Rhizhopora mucronata</i>	Phomoxanthone A (68)	Xanthone	Tumour cell lines or of blood cancer cell lines	[72]
<i>Garcinia mangostana</i>	Garcinone C (69)	Xanthone	MCF-7, A549, Hep-G2 and CNE cell lines	[73]
<i>Garcinia nobilis</i>	Morusignin I (70), 8-hydroxycudraxanthone G (71), cudraxanthone I (72)	Xanthone	Breast cancer cells transduced with control vector (MDA-MB-231-pcDNA3), Human wild-type HCT116 (p53+/+) colon cancer cells, Human glioblastoma multiforme U87MG cells, Human HepG2 hepatocellular carcinoma cells and normal AML12	[74]
<i>Securidaca longepedunculata</i>	1,6,8-trihydroxy-2,3,4,5-tetramethoxyxanthone (73), 1,6-dihydroxy-2,3,4,5,8-pentamethoxyxanthone (74)	Xanthone	Human pancreatic cancer cell line, PANC-1	[75]
<i>Garcinia hunburyi</i>	Desoxymorellin (75)	Xanthone	HEL (human embryonic lung fibroblasts) and HeLa (Henrietta Lacks cervical cancer)	[76]
<i>Garcinia cantleyana</i>	Cantleyanone (76)	Xanthone	Breast cancer (MDA-MB-231 and MCF-7), ovarian cancer (CaOV-3) and HeLa cells	[77]
<i>Garcinia lateriflora</i>	Lateriflorone (77)	Xanthone	P388 cancer cell line	[78]
<i>Garcinia morella</i>	Gambogic acid (78)	Xanthone	T47D and DLD-1 breast cancer cells	[79]
<i>Garcinia gaudichaudii</i>	Gaudichaudione (79)	Xanthone	Parental murine leukemia P388 and P388/doxorubicin-resistant cell lines	[80]
<i>Garcinia cowa</i>	Cowaxanthones G (80)	Xanthone	Human cancer cell lines (HeLa, PANC-1 and A549)	[81]

Source	Compound name	Class of compounds	Cytotoxic activity	References
<i>Oncidium baueri</i>	Batatasin III (81)	Stilbenoid	U251 (glioma, CNS), MCF-7 (breast), NCI-ADR/RES (ovarian expressing phenotype multiple drugs resistance), 786-0 (renal), NCI-H460 (lung, non-small cell), HT29 (colon), HaCat (human keratinocytes, immortalized non-tumoral cell)	[82]
<i>Scirpus yagara</i>	Sciryagarol I (82) and II (83)	Cis-stilbenoid	HeLa Cell Line human epitheloid cervix carcinoma	[83]
<i>Gnetum macrostachyum</i>	Macrostachyols D (84)	Oligostilbenoid	Human cervical carcinoma (HeLa) and human mouth epidermal carcinoma (KB) cell lines	
<i>Cajanus cajan</i>	Cajanotone (85), cajaninstilbene acid (86), pinosylvin monomethyl ether (87), longistylin A (88), longistylin C(89)	Stilbenoid	Human hepatoma cell line HepG2, human breast adenocarcinoma MCF-7 and human lung cancer cell line A549	[84]
Peanuts	Arachidin-1 (90)	Peanut Stilbenoid	Human leukemia HL-60 cells	[85]
<i>Combretum cafferum</i>	Combretastatin A-4 (91)	Stilbenoid	Human hepatocellular carcinoma HepG2, SMMC-7721, gastric carcinoma BGC-803 cells, breast carcinoma MDA-MB-435	[86]
<i>Macaranga siamensis</i>	Macasiamenenes A (92), macasiamenene K (93), Macasiamenene L (94), acasiamenenes M (95)	Prenylated stilbenoid	MOLT-3 (acute lymphoblastic leukemia)	[87]
<i>Monomeria barbata</i>	1,4,7-trihydroxy-2-methoxy-9,10-dihydrophenanthrene (96), 1,3,8-tri (phydroxybenzyl)-4-methoxy-phenanthrene-2,7-diol (97)	Phenanthrene	HepG-2, promyelocytic leukaemia HL-60, ovarian carcinoma Skov-3	[88]
<i>Bulbophyllum odoratissimum</i>	Bulbophythrins A (98) and B (99)	Dimeric phenanthrene	Human leukemia cell lines K562 and HL-60, human lung adenocarcinoma A549, human hepatoma BEL-7402 and human stomach cancer SGC-7901	[89]
<i>Helicteres hirsuta</i>	Pinoresinol (100), boehmenan (101) and boehmenan H (102)	Lignan	LNCaP, Lu1 (human lung cancer), MCF-7 (human breast cancer) and HUVEC (human umbilical vein endothelial) cell lines,	[90]
<i>Sambucus williamsii</i>	Sambucasinol A (103), B (104), C (105)	Lignan	Human cell lines A549, SK-OV-3, SK-MEL-2 and XF498	[91]
<i>Pycnanthus angolensis</i>	Pycnanthulignene A (106)	Lignan	CCRF-CEM leukemia cell line, CEM/ADR50 0 0 cells	[41]
<i>Phyllanthus glaucus</i>	Phyllanthusmin C (107)	Lignan glycoside	HL-60, MCF-7 and SW480 cells	[92]

Table 1. Cytotoxic phenolics from different natural source.

2. Natural phenolics with anticancer properties

2.1. Flavonoids and their cytotoxic activities

Flavonoids and their glycosides are the main phenolic constituents isolate from many natural sources in a reasonable amount. These phenolic compounds are very well-known for their cytotoxic activity and gave very promising results against different types of tumors. The basic structures of flavonoids are based on a C₆-C₃-C₆ skeleton derived from 1,3-diphenylpropane. Flavonoids are further divided into three subclasses on the basis of their little structural changes, substitutions and degree of hydroxylation and polymerization, such as, flavonoids (2-phenylchromen-4-one), isoflavonoids (3-phenylchromen-4-one) and neoflavonoids (4-phenylchromen with no hydroxyl group substitution at position 2). More than ten thousand flavonoids have been identified from natural sources and it exhibit a number of human health benefits due to their interactions with different cellular targets including antioxidant, anti-inflammatory, antiviral and anticancer properties. Isoprenylated flavonoids isolated from Moraceae family has many cytotoxic activity in different cancer cell lines including MCF-7 human breast cancer, TK-10 human renal cancer and UACC-62 human melanoma cells, A549 human lung cancer, Hep3B human hepatocellular cancer, HT-29 human colorectal cancer cells.

The cell viability of B16 melanoma cells after 3 days culture with compounds Artocarpin, cudraflavone C, prenylapigenin, kuwanon C, norartocarpin, albanin A and cudraflavone B at different concentrations was determined using 3-(4,5-dimethylthiazole-2-yl)-2,5-diphenyltetrazolium bromide MTT colorimetric assays. These compounds 1–7 exhibit potent cytotoxicity in a concentration-dependent manner. The IC₅₀ values are 10.3, 9.2, 32.5, 14.2, 7.8, 84.7 and 12.5 μM [3].

Kaempferol is a dietary flavonoid and it has many protective effects for human health, was treated with human breast cancer cell line, MCF-7 and it regulated down the expression of polo-like kinase PLK-1, which has reported to regulate mitotic progression and to be upregulated in several human tumors. This polyphenolic compound significantly reduces cell viabilities of U-2 OS, HOB and 143B cells, but exerts low cytotoxicity on human fetal osteoblast progenitor hFOB cells. It has anticancer effects on Miapaca-2, Panc-1 and SNU-213 human pancreatic cancer cells. In a dose-dependent manner, its decreased viability of pancreatic cancer cells by increasing apoptosis. The anticancer effect of kaempferol mediated by inhibition of EGFR-related Src, ERK1/2 and AKT pathways and it could act as potent pancreatic cancer cells inhibitor [4]. The IC₅₀ value of Dorsmanin F is a prenylated flavonoid and it ranged from 5.34 to 1.94 μM towards leukemia CCRF-CEM cells to 33.30–28.92 μM towards MDA-MB-231-BCRP cells, respectively and from 0.20 to 195.12 μM against CCRF-CEM. This compound induced apoptosis in CCRF-CEM leukemia cells, mediated by MMP disruption and increased reactive oxygen spp (ROS) production [5]. Artocarpesin, cycloartocarpesin (unique flavonoid structures) displayed cytotoxic effect on four cell lines with IC₅₀ values, respectively, below 106, 50 and 25 μM. The IC₅₀ values of these compounds ranged from 23.95 μM for hepatocarcinoma HepG2 cell, 105 μM towards colon carcinoma HCT116 (p53-/-) cells for artocarpesin, from 15.51 μM for leukemia CCRF-CEM cells and 49.83 μM for glioblastoma U87MG. EGFR cells for cycloartocarpesin [6]. 6,7-dimethoxy-4'-hydroxy-8-formylflavon and 4',7-dihydroxy-

8-formyl-6-methoxyflavon are formylated flavonoid and showed significant cytotoxicity against PC3 and A549 cell lines with IC_{50} values of 2.6 and 1.6 μ M, respectively [7]. Isoxanthohumol is known as hops and wines flavonoid and it exhibits antiproliferative activity against human breast cancer MCF-7 cell lines, A-2780 ovarian cancer cell lines, prostate cancer PC-3 and DU145 and colon cancer SW620 and HT-29 cells lines. It inhibits the activation of carcinogens: 2-amino-3-methylimidazo[4, 5-f]quinoline and aflatoxin B1, AFB1 via human cytochrome P450 CYP1A2 [8]. Kushecarpin D is a Chinese traditional herbal medicine and exhibits antiangiogenic activity demonstrated by its effects on migration, adhesion, tube formation of an endothelial cell line. The antiangiogenic activity, together with its antiproliferative effects on endothelial cells, indicates that kushecarpin D is an excellent candidate for development as chemo preventive molecule to be used in combating tumor development [9]. Luteolin is one of the very important flavonoid found in many foods with lot of health benefits and it prevent cancers strongly, it induce cell cycle arrest and apoptosis in various human cancer cells and it synergize antitumor effects of 5-FU on Bel7402 and HepG2 cells, which can related with apoptosis and regulation of 5-FU metabolism. The IC_{50} value of luteolin on PC₃ and LNCaP cells was 31.44 and 32.05 μ M, respectively. It resulted in a marked reduction in cell proliferation in a dose-dependent manner and enhanced the paclitaxel-induced apoptosis in human breast cancer MDA-MB-231 cells by blocking STAT3 [10–12]. Chrysin is found in many flowers and honey and it has preventive effect on cancer induced chemically on xenograft tumor by inducing activity of antioxidant and detoxification enzyme, reducing the activities of cytochrome P450 (CytP450)-dependent monooxygenases, inhibiting cellular proliferation and inducing apoptosis. Chrysin has induced breast cancer resistance protein (BCRP) in Caco-2 cells. Chrysin and many other flavonoids such as fisetin, kaempferol, galangin, myricetin and apigenin reported as potent inhibitors of P-form phenol sulfo transferase mediated sulfation induced carcinogenesis in human hepatoma cell line HepG2. Chrysin kill several histotype cancer cells, including hematological, cervical, liver, colon, lung, breast, nasopharyngeal, glioblastoma, prostate, thyroid and pancreatic cancer [13]. DICO is a nonaromatic B ring flavonoid having potent antitumor activity and inhibited growth of HepG2 cells in different dose/time dependent manners. It induced G2/M cell cycle arrest and apoptosis via a ROS-mediated mitochondrial pathway. It has significant antitumor effect through G2/M cell cycle arrest and apoptosis induction, which suggested DICO has therapeutic potential against tumors [14]. Morusin belongs to the prenylated class of flavonoids and it suppressed signal transducer and activator of transcription 3 STAT3 and nuclear factor-kB/NFkB signaling pathway, which modulate protein expression involved in invasion process. Morusin decreased lung colonization of SK-Hep1 cells in mice. It indicates that morusin possesses antitumor progression potential by suppressing STAT3 and NFkB [15]. Artocarpin showed significant anticancer activities on breast cancer cells (T47D cells) with IC_{50} value on T47D cells was 12.6 μ M on concentration-dependent manner. Anticancer effect of artocarpin in HepG2 and PLC/PRF/5 hepatoma cells is mediated through autophagic cell death mechanism and it showed dose-dependent reduction on cancer cell viability after 24 h of treatment and IC_{50} value was calculated to be approximately 15 μ M in both cell lines [16, 17]. Genkwanin a methoxy flavonoid, significantly inhibited HT-29 and SW-480 human colorectal cancer cells proliferation and inflammatory cytokine IL-8 secretion. It has a better antitumor activity via enhancing host immunity and decreasing inflammatory cytokine levels [18]. Fisetin is a

tetrahydroxy flavonoid having many health benefits and its effects on short- and long-term growth of BRAF-mutated A375, SK-MEL-28 and RPMI-7951 melanoma cancer cells. Results of MTT assay demonstrated that fisetin (10–60 μ M) treatment significantly decreased the growth of A375 8.64–61.75%, SK-MEL-28, 6.94–59.79% and RPMI-7951, 11.60–64.11% cells in a concentration-dependent manner [19]. Hispidulin is found in many herbs and significantly inhibited HepG2 cell growth in a time and dose-dependent manner. Hispidulin (at 200 μ M) inhibited the growth of HepG2 cells by nearly 50, 70 and 90% after 24, 48 and 72 h of treatment, respectively, which suggests that hispidulin promotes HepG2 cell death through apoptosis. It inhibited cell growth in a dose-dependent manner in (HRCC) lines 786-0 and Caki-1 cell lines, whereas Caki-1 cells were found to be more resistant to hispidulin treatment [20, 21]. Quercetin is a dietary flavonoid (berries flavonoid) showed strong cytotoxicity as 1.49-fold in MCF-7 cells and 1.98-fold in MCF-7/dox cells. It suppressed proliferation and survival of HepG2 cancer cells and induced apoptosis by enhancing the expression of p53 and BAX through downregulation of ROS, PKC, PI3K and COX-2 [22, 23]. Hesperetin a citrus flavonoid exhibited inhibition of cell growth in a concentration and time-dependent manner with IC_{50} at 72 h 200 μ M. It could significantly promote apoptosis of Eca109 cells in a dose-/time-dependent manner [24]. Apigenin a trihydroxy flavonoid with potential health benefits exhibited strong growth inhibitory activity in HER2/neu breast cancer cells but was much less effective in inhibiting growth of cells expressing basal levels of HER2/neu. It induces apoptosis in MDA-MB-453 breast cancer cells with involving intrinsic and extrinsic apoptotic pathways. It has shown to downregulate levels of cyclin D1, D3 and cdk4 and increases p27 protein levels in breast cancer cells. Apigenin inhibited human cervical carcinoma HeLa cells growth through an apoptotic pathway. In various human colon carcinoma cells, it resulted cell growth inhibition and G2/M cell cycle arrest. The effects of apigenin on lung cancer cells were evaluated and it inhibited A549 lung cancer cell. It has capability to significantly reduce cell number and induce apoptosis in PWR-1E, LNCaP, PC-3 and DU145 cells [25, 27]. Quercetin-3-O- β -D-glucopyranoside also known as isoquercetin and found in mangoes in high amount and it is an interesting dietary compound worth further investigation as a cytoprotective agent. Pretreatment of PC12 cells with nontoxic concentrations of this compound protect cells from H_2O_2 induced cytotoxicity with decrease in generation of reactive oxygen species (ROS). These observations qualify it as cytoprotective dietary compound [27]. Amplexicaule A is found in many herbs and it increased levels of cleaved caspase-3, -8, -9 and PARP, resulted from suppression of MCL-1 and BCL-2 expression in cells. This compound inactivated the Akt/mTOR pathway of breast cancer cells. It influenced strongly on breast cancer cells, most likely by induction of apoptosis [28]. Triticuside A a dietary flavonoid found in wheat and induced apoptosis accompanied by a significant decrease in Mcl-1 and Bcl-2 proteins and by increase in cleavage of caspases-3, -7, -9 and PARP. It suppressed the level of phospho-Akt and its downstream targets, mTOR and P70 S6 kinase. LY294002, a specific inhibitor of PI3K, significantly enhanced the triticuside A induced apoptosis. It may be a potentially useful wheat bran component and can be used for treatment of breast cancer [29].

Naringin is a diglycoside flavonoid and found in many citrus fruits and it inhibited cell proliferation and promoted cell apoptosis and G1 cycle arrest, accompanied by increased p21 and decreased surviving. Significant inhibitory effects of naringin on the cell proliferation of TNBC cells were observed. MDA-MB-231 and BT-549 cells treatment with naringin (50, 100

and 200 M) for 48 h, significantly increased apoptosis in breast cancer cells. It exerts significant effects on inhibition of breast cancer cells growth in through mediating-catenin pathway. Treatment of Raji with naringin showed maximum sensitivity towards NK cell lysis and the activity was 2.5-fold with naringin treatment [30–32]. Quercetin-3-*O*- α -L-rhamnopyranosyl-(1 $\rightarrow$ 6)- β -D-glucopyranoside induces apoptosis in A549 cancer cells via caspase activation through cytochrome release from mitochondria. It tested against A549 lung cancer cell line, COLO320DM cancer cell line and Vero cell line and it showed prominent cytotoxic activity against A549 lung cancer cell. This compound showed 87.41% activity at dose of 164 μ M with IC_{50} value of 82 μ M. COLO320DM cancer cell line was maintained in complete tissue culture medium [32]. Pectolinarin and astragalin isolated from edible plants against H_2O_2 -induced cell death of human brain neuroblastoma SK-N-SH cells. These compounds showed protective effects against H_2O_2 -induced cell death and inhibited ROS generation by oxidative stress [33]. Troxerutin showed strong binding with calf thymus DNA in vitro and DNA interaction was confirmed by CD spectropolarimetry. The mode of binding of troxerutin to DNA was assessed by competing with EtBr or DAPI, known DNA intercalator and minor groove binder, respectively. It induced cytotoxicity in radioresistant DU145 and sensitive PC3 prostate cancer cells. When troxerutin was pretreated with DU145 and PC3 cells and it were exposed to gamma-radiation, the cytotoxicity induced in PC3 and DU145 prostate cancer cells and it was monitored by MTT assay. The toxicity induced by troxerutin 5 μ M was more than 40% over control at 24 h, but more than 50% increase was observed at 48 h. Radiation per se induced around 18% cell death in PC3 cells [34]. Vicenin gave IC_{50} values 141.7 μ M, 195.7 μ M and 369.3 μ g mL⁻¹ on treatment with PC-3, DU-145 and LNCaP cells. Curcumin was used as positive control with IC_{50} 17.90 μ M. It is an active constituent of the medicinal herbs (Tulsi) and it effectively induced antiproliferative, antiangiogenic and proapoptotic effect in CaP cells (PC-3, DU-145 and LNCaP) [35, 36] (**Figure 1**).

2.2. Isoflavonoids and their cytotoxic activities

Isoflavonoids play very important role in human health-promoting natural chemicals. They belong to plants secondary metabolites, mediate diverse biological functions through numerous pathways. Isoflavonoids are phenolic compounds and possess a 3-phenylchroman skeleton that biogenetically derived from 2-phenyl chroman skeleton of flavonoids. Some studies reported that anticarcinogenic activities of dietary soy isoflavonoids play important role for preventing colorectal cancer. Isoflavonoids have shown to possess many biological properties that can account for cancer prevention. Isoflavones exert their effects through numerous pathways with respect to the cancer prevention; and it use mechanisms of action which appear to be various, complementary and overlapping.

Genistein is a soybeans isoflavones and it decrease the risk of breast cancer and induced downregulation of CIP2A in breast cancer cells MCF-7-C3 and T47D, which correlates with its growth inhibition and apoptotic activities. In a model of human prostate carcinoma, treatment with genistein, decreased the metastatic burden, without changing the size of primary tumor and cell detachment were also decreased. According to studies, patients with PCa effectively tolerate genistein and therapeutic modulation of metastasis, specifically MMP-2 (matrix metalloproteinase 2) is possible [39, 40]. Inhibitory effect of daidzein investigated by

Figure 1. Structures of flavonoids 1–35.

bacterial flora in intestines, on DU145, LnCaP and PC3 human prostate cancer cell. The results provide a better understanding of the biomolecular mechanisms of this compound as natural anticancer agent and provide base for development of daidzein and its analogs as potent anticancer molecules [39]. Biochanin A is found in nuts and beers and it effects on pancreatic cancer progression and it induced dose-dependent toxicity on pancreatic cancer cells Panc1 and AsPC-1. It reduced colony formation ability of Panc1 cells and induced dose-dependent apoptosis [40]. The IC_{50} value of Alpinum isoflavone ranged from 5.91 μ M towards leukemia CEM/ADR5000 cell and to 65.65 μ M towards drug-resistant breast adenocarcinoma MDA-MB-231-BCRP cells. This sioflavone induced apoptosis of CCRF-CEM cells, mediated by the loss of MMP and increase the ROS production [41]. Puerarin prevents the proliferation of breast cancer cells (MDA-MB-231, HS578T and MCF-7) at 50% of cell growth inhibition with concentration of 46, 71 and 69 μ M, respectively. Puerarin was further preventing three different types of breast cancer in the G0/G1 phase of the cell cycle and stimulated apoptosis in these cells [42] (**Figure 2**).

Figure 2. Structure of isoflavonoids 36–40.

2.3. Chalcone and their cytotoxic activities

Chalcones mainly belongs to flavonoids family and known as open chain flavonoid in which two aromatic rings A and B structurally joined by α,β -unsaturated carbonyl system. Plants containing chalcones have been used traditionally as anti-inflammatory, antioxidant, antimalarial, antimicrobial, antifungal, antitubercular, cytotoxic, antiviral, antitumor and chemopreventive agent. These are very common phenolics specially founds in Leguminosae, Moraceae and Asteraceae families.

Butein is a natural dietary chalcone has many traditional uses and a herbal medicine. It reduced the cell viability of cultured human uveal melanoma cells in a dose-dependent manner with IC_{50} at 13.3 and 15.8 μ M in SP6.5 and M17 cell lines, respectively. Similar effects were also found in a highly aggressive and metastatic C918 cell line (IC_{50} 16.7 μ M). At 2 μ M concentration, it inhibited the incorporation of ^{14}C -labelled thymidine, uridine and leucine into the colon cancer cells whilst 5-fluorouracil (5-FU, a chemotherapeutic drug) at 50 μ M concentration. The cytotoxic action of butein was different from 5-FU but may be similar to colchicine, a known HeLa cell inhibitor. Butein inhibit telomerase activity by downregulating hTERT gene expression in human leukemia cells and it causes apoptosis of breast cancer cells, while luminal HER2⁺ HCC-1419, HCC-2218 and SKBR-3 breast cancer cells. Treatment with butein (10–30 μ M) decreased cell viability in LNCaP (29, 42 and 52%), CWR22Rv1 (20, 31 and 42%) and PC-3 (11, 22 and 35%) cells. It inhibited colony formation, cell viability, migration, invasion, induced cell cycle at G2/M stage, cell apoptosis and enhanced caspase-3, -8 and -9 activity in HeLa cells in a dose-dependent manner [43–47]. Isoliquiritigenin is a natural chalcone and used for the treatment of cancer, it is cytostatic and able to overcome the intrinsic resistance of U373 cancer cells to proapoptotic stimuli. After 72 h treatment with 10 μ g mL⁻¹ of isoliquiritigenin, a typical differentiated morphology observed in HL-60 cancer cells, including the decrease in karyoplasmic ratio and the increase in kidney-shape nuclear cells. This compound is able to induce the monocytic differentiation in leukemia cells. It has potential as a drug for leukemia. HepG2 cells are significantly more resistant to isoliquiritigenin when the activity of p53 was blocked. Isoliquiritigenin inducible p53 plays a key apoptotic role and may do so by regulating the expression of specific target molecules that promotes efficient apoptotic cell death following G2/M-cell cycle arrest. It inhibits the proliferation of prostate cancer cells, via inhibition of ErbB3 signaling and PI3K/Akt pathway [48]. 2',4'-dihydroxy-6-methoxy-3,5-dimethylchalcone tested on human cell lines including liver cancer SMMC-7721 cells, pancreas cancer 8898 cells, tumor of cervix uteri HeLa cells, lung cancer SPC-A-1 cells, high metastatic lung carcinoma 95-D cells and gall bladder carcinoma GBC-SD cell lines was dose-dependent. It observed that different cells had a different sensitivity for inhibition effect of this chalcone. The IC_{50} values on cytotoxicity were 32.3, 37.2, 37.7, 81.3, 84.6 and 84.8 μ M for SMMC-7721, 8898, HeLa, GBC-SD SPC-A-1 and 95-D cells, respectively [49, 50]. 4'-hydroxy-2',6'-dimethoxychalcone showed IC_{50} values in a range of 2.54 μ M against CEM/ADR5000 leukemia cells to 58.63 μ M towards hepatocarcinoma HepG2 cells. This compound arrested cell cycle between Go/G1 phase and induced apoptosis via disrupted mitochondrial membrane potential MMP and increased production of reactive oxygen species (ROS) in the studied leukemia cell line [51]. Xanthohumol is a prenylated chalcone found in beers; it caused dose dependent (0.1–100 μ M) decrease in growth of MCF-7, HT-29 and A-2780 cancer

cell lines. After two-day treatment, the concentrations at which growth of MCF-7 cells was inhibited by 50% (IC_{50}) were 13.3 μ M for xanthohumol. After four-day treatment, IC_{50} for xanthohumol was 3.47 μ M [52, 53]. Flavokawain B is known as kava chalcone and it induced caspase and mitochondria-dependent apoptosis which characterized by cytochrome c release and Bak translocation to mitochondria. It induces G2/M accumulation, autophagy and apoptosis, leading to HCT116 colon cancer cell growth inhibition. It further induced both MCF-7 and MDA-MB231 and significant G2/M arrest was seen in MDA-MB231 cells [54]. Isolespeol is a geranyl chalcone and it showed inhibitory activity against human liposarcoma cells SW 872 with IC_{50} values of 3.8 μ M. Treatment of SW 872 human liposarcoma cells with this compound stimulated increase protein expression of Fas, FasL and p53 [55]. Xanthoangelol and 4-hydroxyderricin inhibit adipocytes differentiation through AMPK and mitogen-activated protein kinase pathways, resulting in the down-expression of adipocyte-specific transcription factors. Xanthoangelol induce apoptotic cell death by activation of caspase-3 in neuroblastoma and leukemia cells through a mechanism that does not involve Bax/Bcl-2 signal transduction. Therefore, this compound may effective drug for the treatment of neuroblastoma and leukemia. 4-Hydroxyderricin showed significant cytotoxicity in four human tumor cell with IC_{50} values 4.8 μ M (CRL1579), 5.5 μ M (HL60), 4.2 μ M (AZ521) and 10.2 μ M (A549). 4-Hydroxyderricin induced further the early apoptosis of HL60 cells and it observed as membrane lipid exposure in the flow cytometry [56, 57]. The IC_{50} values for Isobavachalcone is a prenylated chalcone and having wide range of biological activities including anticancer, it ranged from 0.20 μ M (towards CCRF-CEM cells) to 195.12 μ M (towards leukemia CEM/ADR 5000 cells) for doxorubicin. It induces apoptosis in CCRF-CEM leukemia cells, mediated by caspase activation and the disruption of MMP [58]. Cytotoxicity of Poinsettifolin B assessed against different sensitive and multidrug-resistant cancer cell lines. The IC_{50} values for this compound ranged from 5.34 to 1.94 μ M towards CCRF-CEM leukemia cells, to 33.30 and 28.92 μ M towards human breast cells MDA-MB-231-BCRP, respectively and from 0.20 μ M against human leukemia cell CCRF-CEM, to 195.12 μ M against CEM/ADR5000 cells [59]. *2'-hydroxy-3,4,4',5,6'-pentamethoxychalcone* is very potent in inhibiting breast adenocarcinoma (MCF-7), lung cancer (NCI-H460) and melanoma (A375-C5) cell lines [60]. Rhuschalcones II–VI showed potent cytotoxic activities against HCT-116 and HT29 human colon cancer cells [61] (Figure 3).

2.4. Naphthalenes and their cytotoxic activities

Naphthalenes are simplest and most important member of arenas, in which two benzene rings are fused with each other and it based on a C_6-C_4 skeleton. Many naphthalene-based natural products have been isolated from different sources and showed significant biological activities.

Syriacusin A inhibit activity of human neutrophil elastase HNE, a serine protease to degrade extracellular matrix ECM proteins including elastin, with IC_{50} values 8.0, 5.2 and 6.1 μ M, respectively [62]. Rumexneposide A showed against broad spectrum activity against human cancer cells lines, including lung and breast cancer cells (A549, H522, MCF-7, MCF-10A, SKBR3) with IC_{50} value 31.0, 15.7, 21.8, 22.8 and 20.7 μ M [63]. Parvinaphthol B, parvinaphthols C showed marginal cytotoxicity against the MDA-MB-231 human triple-negative breast cancer cell line with IC_{50} values ranging from 62.3 to 129.6 μ M [64].

Figure 3. Structure of chalcones 41–57.

2.5. Anthraquinone and their cytotoxic activities

Anthraquinones are the active constituents of many herbs with a tricyclic C6-C2-C6-based skeleton. They are widely distributed in the *Fabaceae*, *Liliaceae*, *Labiata*, *Polygonaceae* and *Rhamnaceae* families. Anthraquinone found in herbs (*Polygonum*) in vegetable (cabbage, lettuce and beans).

Chrysophanol was active in the oophoroma SKOV-3 cell line, where the IC_{50} value was $5.62 \mu M$ and with IC_{50} value $20.4 \mu M$ for MCF-7 breast cancer cell line [65]. 9,10-anthraquinone showed strong cytotoxic activity against COLO320 human colon carcinoma cell line. It gave 79.7% cytotoxicity at $300 \mu g mL^{-1}$ concentration of compound with $75 \mu g mL^{-1}$ IC_{50} value. The treatment of human colon carcinoma cell line COLO320 with 9,10-anthraquinone significantly decreased the proliferation of cells and enhanced the formation of apoptotic bodies and fragmented DNA. The expressions of p53 and caspase-3 was upregulated from 9,10-anthraquinone in colon adenocarcinoma cells [66]. 2-hydroxy-9,10-anthraquinone showed cytotoxic activity against A549 lung and COLO320 cells lines and it showed 62.7% activity at the dose of $500 \mu g mL^{-1}$ with IC_{50} value of $400 \mu g mL^{-1}$ against COLO320 cells [67]. 2,3-dihydroxy-9,10-anthraquinone inhibits PI3K/AKT activity after treatment. Also, COX-2 enzyme plays a major role in colorectal cancer and significantly reduced COX-2 enzyme in COLO320 cells.

It involves in apoptotic pathway, mitochondrial function, cell cycle checkpoint and control the over expression gene during the colorectal cancer [66]. Chrysophanol 8-O-beta-(6'-acetyl) glucopyranoside exhibited relatively higher cytotoxic activities against human oral squamous cell carcinoma HSC-2 and salivary gland tumor HSG cell lines than against normal human gingival fibroblasts HGF cells lines [69] (**Figure 4**).

2.6. Xanthone and their cytotoxic activities

Xanthones are three-membered ring compounds with (C6-C1-C6) skeleton and are mainly found in higher plants and microorganisms. Now days, they gained great importance due to their significant pharmacological and biological properties. These types of natural compounds have broad biological profile, such as, antihypertensive, anti-inflammatory, antioxidant antithrombiol, anticancer and antiviral activities. Xanthones are commonly found in Gentianaceae, Moraceae, Guttiferae, Clusiaceae, Polygalaceae in citrus fruits and mangosteen.

Formoxanthone C has α,α,β -trimethylfuran ring and it showed potent cytotoxic activity against NCI-H187 with IC_{50} $0.22 \mu\text{g mL}^{-1}$, which is stronger than elliptecine, a standard drug with IC_{50} $0.45 \mu\text{g mL}^{-1}$ [70, 71]. Phomoxanthone A displayed strong anticancer activity, on the treatment against cisplatin resistant (CisR) cancer cell lines or blood cancer cell lines with an IC_{50} values in sub-micromolar concentration and it was up to 100-folds less active against PBMC peripheral blood cells from a healthy donors [72]. Garcinone C was tested for cytotoxicity against MCF-7, A549, Hep-G2 and CNE cell lines by 3-(4,5-dimethylthiazol2-yl)-2,5-diphenyltetrazolium bromide (MTT) colorimetric assay, while doxorubicin used as positive control. Garcinone C showed strong cytotoxicity against all cell lines with IC_{50} values from 4.3 ± 0.1 to $7.1 \pm 0.5 \mu\text{M}$ [73]. Morusignin I 70 and cudraxanthone I inhibited proliferation of all cancer cell lines including sensitive and drug-resistant phenotype. 8-hydroxycudraxanthone G showed activity on many cell lines with IC_{50} values ranged from $16.65 \mu\text{M}$ against CCRF-CEM leukemia cells to $70.38 \mu\text{M}$ against HepG2 hepatocarcinoma cells. The IC_{50} values of different cells ranged from $7.15 \mu\text{M}$ against CCRF-CEM human leukemic cells to $53.85 \mu\text{M}$ against U87MG. Δ EGFR human glioblastoma cells for morusignin I and $2.78 \mu\text{M}$ against MDA-MB231 BCRP breast cancer cells to $22.49 \mu\text{M}$ against U87MG human glioblastoma cell line for cudraxanthone I [74]. 1,6,8-trihydroxy-2,3,4,5-tetramethoxyxanthone, 1,6-dihydroxy-2,3,4,5,8-pentamethoxyxanthone were tested for preferential cytotoxic activity against human pancreatic cancer cells PANC-1 under nutrient deprived condition. These compound displayed potent cytotoxicity with PC_{50} of 22.8 and $17.4 \mu\text{M}$, respectively. They triggered apoptosis-like PANC-1 cell death in NDM with glucose sensitive mode [75]. Desoxymorellin inhibited the growth of HEL human embryonic lung fibroblasts and HeLa Henrietta Lacks cervical cancer cell with a minimum inhibitory concentration MIC of 0.39 mg mL^{-1} [76]. Cantleyanone displayed significant cytotoxicity against breast cancer MDA-MB-231 and MCF-7, ovarian cancer CaOV-3 and HeLa cells with EC_{50} values ranging from 0.22 to 17.17 mg mL^{-1} [77]. Lateriflorone was cytotoxic against P388 cancer cell line with an ED_{50} value 5.4 mg mL^{-1} [78]. Gambogic acid inhibited proliferation of T47D and DLD-1 breast cancer cells with GI_{50} values of 0.04 and 0.03 mm , respectively [79]. Gaudichaudione displayed strong growth inhibitory activity against parental murine leukemia P388 and P388/doxorubicin resistant cell lines at low μM concentrations [80]. Cowaxanthones G showed potent inhibition on cell viability $IC_{50} < 10 \mu\text{M}$, while etoposide was used as positive control [81] (**Figure 5**).

Figure 4. Structure of naphthalenes 58–61 and Anthraquinones 62–66.

Figure 5. Structure of structure of xanthenes 67–80.

2.7. Stilbene and their cytotoxic activities

Stilbenoids are formed by flavonoid biosynthesis pathway and it consists on C6-C2-C6-based skeleton and usually found in peanuts, grapes and wines. Stilbenoid have a great interest on account due to their promising pharmacological activities including anticancer, antimicrobial, antioxidant and anti-inflammatory. Stilbenoid play important role of phytoalexin in plants. They play very important role in the defense of different pathogen.

Batatasin III inhibited growth of all cell lines including U251 glioma, CNS, MCF-7 breast, NCI-ADR/RES (ovarian expressing phenotype multiple drugs resistance), 786-0 renal NCI-H460 lung, non-small cell, HT29 colon, HaCat human keratinocytes, immortalized non-tumoral cell with GI_{50} values close to 30 mg mL^{-1} [82]. The cytotoxicity of Sciryagarol I and II was evaluate by MTT assay against human tumor cell lines MGC803, SMMC7721 and Hela and it showed significant cytotoxicity against Hela cell lines with IC_{50} values 61.21 and $7.21 \mu\text{M}$, respectively [83]. Oligostilbene macrostachyol D showed significant cytotoxicity to HeLa cells with IC_{50} value of $4.13 \mu\text{M}$. Cajanotone, cajaninstilbene acid, pinosylvin monomethyl ether, longistylin A and longistylin C showed strong cytotoxicities against human hepatoma HepG2, human breast adenocarcinoma MCF-7 and human lung cancer A549 cell lines with IC_{50} values in the range of $3.5\text{--}15.5 \mu\text{M}$, while doxorubicin used as a positive control [84]. Arachidin-1 induced human leukemia HL-60 cells death with EC_{50} value $4.2 \mu\text{M}$ and it was more potent than resveratrol and induces cell death in HL-60 cells through only the intrinsic apoptotic pathway [85]. Combretastatin A-4 showed strong cytotoxicity against human cancer cell lines: HepG2 with IC_{50} value 9.2, SMMC-7721 with IC_{50} value 12.8, BGC-803 with IC_{50} value 12.2, MDA-MB-231 with IC_{50} value $17.6 \mu\text{M}$ [86]. Macasiamenenes A2, macasiamenene K, Macasiamenene L, acasiamenenes M exhibited significant cytotoxicity against MOLT-3 cancer cell line with the IC_{50} values in the range of $0.66\text{--}9.78 \mu\text{M}$, etoposide used as Positive control standard [87] (Figure 6).

2.8. Phenanthrene and their cytotoxic activities

Phenanthrenes consist on three fused aromatic rings and found in *Dendrobium* and *Dioscorea* spp.

Cytotoxicities of 1,4,7-trihydroxy-2-methoxy-9,10-dihydrophenanthrene, 1,3,8-tri (phydroxybenzyl)-4-methoxy-phenanthrene-2,7-diol towards liver carcinoma HepG-2, promyelocytic leukaemia HL-60, ovarian carcinoma Skov-3 and epidermoid carcinoma A431 cancer cell lines were determined by MTT method. 1,4,7-trihydroxy-2-methoxy-9,10-dihydrophenanthrene is the strongest one to HepG-2 and Skov-3 with IC_{50} of 15.9 and $124.0 \text{ mmol L}^{-1}$, respectively; 1,3,8-tri (phydroxybenzyl)-4-methoxy-phenanthrene-2,7-diol is the strongest one to HL-60 and A431 with IC_{50} of 34.9 and 46.4 mmol L^{-1} , respectively [88]. Biphenanthrenes bulbophythrins A and B were evaluated in vitro for their inhibitory ability against human leukemia cell lines K562 and HL-60, human lung adenocarcinoma A549, human hepatoma BEL-7402 and human stomach cancer SGC-7901, using cisplatin as a positive control. Bulbophythrins A exhibited some selectivity against HL-60 and BEL-7402 with IC_{50} values of 1.27×10^{-3} and $1.22 \times 10^{-3} \mu\text{mol mL}^{-1}$, respectively, whereas bulbophythrins B was most active against A549 with IC_{50} value of $1.18 \times 10^{-3} \mu\text{mol mL}^{-1}$ [89] (Figure 7).

Figure 6. Structure of structure of stilbenoids 81–95.

2.9. Lignan and their cytotoxic activities

Lignans based on (C6-C3)₂ biphenolic skeleton and they consists on a large group of plant phenolic produced by the oxidative dimerization of two phenyl propanoid moieties. They are found in black berries, strawbwerries, raspberries, blue berries, broccoli, apricot, cabbage and many seeds. The antiestrogenic effects of lignans could help to reduce the risk of hormone-associated breast, uterine, ovarian and prostate cancers.

Pinoresinol, boehmenan and boehmenan H were found the most cytotoxic against human cancer cell lines Lu1, LNCaP and MCF-7 and a normal cell line HUVEC with ED₅₀ values for Lu1 cell line was 0.8, 10.4 and 5.3 g mL⁻¹, for LNCaP cell line was 0.5, 9.5 and 7.7 g mL⁻¹, while the ED₅₀ values for MCF-7 cell line was 1.7, 10.0 and 10.2 g mL⁻¹, ED₅₀ values for HUVEC 1.1, 9.0 and 6.2 g mL⁻¹ were observed for these lignans. The cytotoxic activity of pinoresinol against various cancer cell lines has been mentioned in the literature. It exhibited cytotoxicity against the KB cell line with an IC₅₀ value of 2.2 µg mL⁻¹ [90]. The cytotoxic activity of Sambucasinol A, B and C was evaluated by determining their inhibitory effects on human tumor cell lines A549, SK-OV-3, SK-MEL-2 and XF498 using the SRB bioassay. These compounds showed cytotoxicity against all cell lines, with IC₅₀ values in the range of 11.07–19.62 µM [89]. The IC₅₀ values for the pycnanthulignene A ranged from 0.20 µM (towards CCRF-CEM cells) for pycnanthulignene A to 195.12 µM (towards CEM/ADR5000 cells) for doxorubicin [41]. Phyllanthusmin C showed significant cytotoxicity against human leukemia HL-60 cells, human breast MCF-7 cells and human colon SW480 cell lines with IC₅₀ values of 9.2, 19.2 and 20.5 µM, compared with the cisplatin with IC₅₀ values of 1.7, 10.9 and 9.9 µM, respectively [92] (**Figure 8**).

Figure 7. Structure of phenanthrenes 96–99.

Figure 8. Structure of lignans 100–107.

3. Conclusion

Phenolic plays an important role in human nutrition and health benefit and chemoprevention is one of the most realistic and promising approaches for the prevention of malignant disorders. The results of this chapter may help to identify the phenolic compounds from natural sources with optimized cytotoxic activity to be tested for the treatment of different types of tumors. Phenolic compounds are incorporated into them and the order of their incorporation efficiency is similar to its cytotoxic activity. The cytotoxic data of phenolic compounds compiled in this chapter and relationships presented here cannot only be useful in chemoprevention to choose the correct source of these compounds containing most active natural polyphenols, considering the individual genetic cancer risks and familial anamnesis but also in the selection of parent compounds to design and synthesize novel chemotherapy drugs starting from the valuable material given to us by the nature.

Acknowledgements

This book chapter was supported by a grant from the “Research Center of the Female Scientific and Medical Colleges”, Deanship of Scientific Research, King Saud University.

Author details

Shagufta Perveen* and Areej Mohammad Al-Taweel

*Address all correspondence to: shagufta792000@yahoo.com

Department of Pharmacognosy, College of Pharmacy, King Saud University, Riyadh, Kingdom of Saudi Arabia

References

- [1] Huang WY, Cai YZ. Natural phenolic compounds from medicinal herbs and dietary plants: potential use for cancer prevention. *Nutrition and Cancer*. 2010;**62**:1–20. doi:10.1080/01635580903191585
- [2] World Health Organization, Cancer [Internet]. 2016. Available from: <http://www.who.int/cancer/en/>
- [3] Enos TA, Keisuke Y, Kuniyoshi S, Ryuichiro K. Isoprenoid-substituted flavonoids from wood of *Artocarpus heterophyllus* on B16 melanoma cells: cytotoxicity and structural criteria. *Fitoterapia*. 2010;**81**:120–123. doi:10.1016/j.fitote.2009.08.001
- [4] Jungwhoi L, Jae HK. Kaempferol inhibits pancreatic cancer cell growth and migration through the blockade of EGFR-related pathway in vitro. *PLoS One*. 2016. doi:10.1371/journal.pone.0155264.2016

- [5] Victor K, Armelle TM, Maen Z, Bathelemy N, Gilbert DWFK, Simeon FK, Bonaventure TN, Thomas E. Cytotoxicity of two naturally occurring flavonoids (dorsmanin F and poinsettifolin B) towards multi-factorial drug-resistant cancer cells. *Phytomedicine*. 2015;**22**:737–743. doi:10.1016/j.phymed.2015.04.007
- [6] Victor K, Armelle TM, Maen Z, Bathelemy N, Gilbert DWFK, Simeon FK, Bonaventure TN, Thomas E. Cytotoxicity of three naturally occurring flavonoid derived compounds (artocarpesin, cycloartocarpesin and isobavachalcone) towards multi-factorial drug-resistant cancer cells. *Phytomedicine*. 2015;**22**:1096–1102. doi:10.1016/j.phymed.2015.07.006
- [7] Jinxiong C, Hongqiong L, Yuanxing D, Wei Z, Guangyu Y, Yadong G, Yongkuan C, Qiufen H. Three new flavonoids from the leaves of oriental tobacco and their cytotoxicity. *Phytochemistry Letters*. 2013;**6**:144–147. doi:10.1016/j.phytol.2012.12.001
- [8] Anna KZ, Wanda KM, Malgorzata G, Katarzyna W, Edyta W, Mirosław A. Isoxanthohumol- biologically active hop flavonoid. *Fitoterapia*. 2015;**103**:71–82. doi:10.1016/j.fitote.2015.03.007
- [9] Li-Ping P, He-Ping C, Mei-Ai C, Xiu-Li Z, Qing-Xiang G, Cheng-Shan Y, Chun-Ming W. The antiangiogenic activity of Kushecarpin D, a novel flavonoid isolated from *Sophora flavescens* Ait. *Life Sciences*. 2013;**93**:791–797. doi:10.1016/j.lfs.2013.09.025
- [10] Huanli X, Tao Y, Xiaohui L, Ye T, Xiaoliang C, Ru Y, Shuonan S, Xiukun L, Guanhua D. Luteolin synergizes the antitumor effects of 5-fluorouracil against human hepatocellular carcinoma cells through apoptosis induction and metabolism. *Life Sciences*. 2016;**144**:138–147. doi:10.1016/j.lfs.2015.12.002
- [11] Han K, Meng W, Zhang JJ, Zhou Y, Wang YL, Su Y, Lin SC, Gan ZH, Sun YN, Min DL. Luteolin inhibited proliferation and induced apoptosis of prostate cancer cells through miR-301. *Oncotargets and Therapy*. 2016;**9**:3085–3094. doi:10.2147/OTT.S102862
- [12] Sato Y, Sasaki N, Saito M, Endo N, Kugawa F, Ueno A. Luteolin attenuates doxorubicin-induced cytotoxicity to MCF-7 human breast cancer cells. *Biological & Pharmaceutical Bulletin*. 2015;**38**:703–709. doi:10.1248/bpb.b14-00780
- [13] Eshvendar RK, Lakshmi NB, Rajaram MM, Athira KV, Ranadeep G, Chandana CB. Chemopreventive and therapeutic potential of chrysin in cancer: mechanistic perspectives. *Toxicology Letters*. 2015;**233**:214–225. doi:10.1016/j.toxlet.2015.01.008
- [14] Daonian Z, Anhua W, Cheng C, Jinlan R. DICO, a novel nonaromatic B-ring flavonoid, induces G2/M cell cycle arrest and apoptosis in human hepatoma cells. *Food and Chemical Toxicology*. 2013;**57**:322–329. doi:10.1016/j.fct.2013.03.032
- [15] Wea-Lung L, Deng-Yu L, Yean-Jang L, Nai-Fang C, Tsui-Hwa T. Antitumor progression potential of morusin suppressing STAT3 and NFκB in human hepatoma SK-Hep1 cells. *Toxicology Letters*. 2015;**232**:490–498. doi:10.1016/j.toxlet.2014.11.031
- [16] Arung ET, Wicaksono BD, Handoko YA, Kusuma IW, Shimizu K, Yulia D, Sandra F. Cytotoxic effect of artocarpin on T47D cells. *Journal of Natural Medicines*. 2010;**64**:423–429. doi:10.1007/s11418-010-0425-6

- [17] Cheng-Wei T, Wen-Sheng T, Liang-Tzung L, Chiang-Wen L, Feng-Lin Y, Chun-Ching L. Enhanced autophagic activity of artocarpin in human hepatocellular carcinoma cells through improving its solubility by a nanoparticle system. *Phytomedicine*. 2016;**23**:528–540. doi:10.1016/j.phymed.2016.02.010
- [18] XueW, Zi-Jing S, Xin H, Run-Qi Z, Chun-Feng Z, Fei L, Chong-Zhi W, Chun-Su Y. Antitumor and immunomodulatory activity of genkwanin on colorectal cancer in the APCMin/+ mice. *International Immunopharmacology*. 2015;**29**:701–707. doi:10.1016/j.intimp.2015.09.006
- [19] Harish CP, Ronald DB, Katherine MH, Jyoti A, Craig AE, Mohammad A, Farrukh A. Fisetin, a phytochemical, potentiates sorafenib-induced apoptosis and abrogates tumor growth in athymic nude mice implanted with BRAF-mutated melanoma cells. *Oncotarget*. 2015;**6**:28296–28311. doi:10.18632/oncotarget.5064
- [20] Scoparo CT, Valdameri G, Worfel PR, Guterres FA, Martinez GR, Winnischofer SM, Di Pietro A, Rocha ME. Dual properties of hispidulin: antiproliferative effects on HepG2 cancer cells and selective inhibition of ABCG2 transport activity. *Molecular and Cellular Biochemistry*. 2015;**409**:123–133. doi:10.1007/s11010-015-2518-8
- [21] Hui G, Qixiao J, Yantao H, Jianjun P, Chunbo W. Hispidulin potentiates the antitumor effect of sunitinib against human renal cell carcinoma in laboratory models. *Cell Biochemistry and Biophysics*. 2015;**71**:757–764. doi:10.1007/s12013-014-0260-6
- [22] Li SZ, Li K, Zhang JH, Dong Z. The effect of quercetin on doxorubicin cytotoxicity in human breast cancer cells. *Anticancer Agents in Medicinal Chemistry*. 2013;**13**:352–355. doi:10.2174/1871520611313020020
- [23] Akhilendra KM, Manjula V. Anticarcinogenic action of quercetin by downregulation of phosphatidylinositol 3-kinase (PI3K) and protein kinase C (PKC) via induction of p53 in hepatocellular carcinoma (HepG2) cell line. *Molecular Biology Reports*. 2015;**42**:1419–1429. doi:10.1007/s11033-015-3921-7
- [24] Dandan W, Jixiang Z, Jing W, Jiao L, Fei L, Weiguo D. Hesperetin induces apoptosis of esophageal cancer cells via mitochondrial pathway mediated by the increased intracellular reactive oxygen species. *Tumor Biology*. 2016;**37**:3451–3459. doi:10.1007/s13277-015-4176-6
- [25] Sanjeev S, Sanjay G. Apigenin: a promising molecule for cancer prevention. *Pharmaceutical Research*. 2010;**27**:962–978. doi:10.1007/s11095-010-0089-7
- [26] Balbino LS, Mona NO, Paulo LCC, Bruno PSP, Alessandra BS, Tais A, Victor DAS, Maria FDC, Ramon SE, Marcienne T, Herve C, Marie-Pierre J, Vivaldo M, Silvia LC, Flavonoids suppress human glioblastoma cell growth by inhibiting cell metabolism, migration and by regulating extracellular matrix proteins and metalloproteinases expression. *Chemico-Biological Interactions*. 2015;**242**:123–138. doi:10.1016/j.cbi.2015.07.014.
- [27] Yalda S, Maryam R, Leila H, Zohreh J, Quercetin-3-O-β-D-glucopyranoside, a dietary flavonoid, protects PC12 cells from H₂O₂-induced cytotoxicity through inhibition of reactive oxygen species, *Food Chemistry*. 2015;**167**:162–167. doi:10.1016/j.foodchem.2014.06.079

- [28] Meixian X, Hanwen S, Guangwen S, Dingrong W, Feng H, Morgann L, Yali D, Jun L, Sinisa D, Gaungzhong Y, Chunhua S. Amplexicaule A exerts anti-tumor effects by inducing apoptosis in human breast cancer. *Oncotarget*. 2016;**7**:18521–18530. doi:10.18632/oncotarget.7848
- [29] Yu S, Yan C, Yi Z, Fu-Qin G, Hao S, Xing-cong R, Yu C, Xu Feng, Jin-Ming Y. Triticuside A, a dietary flavonoid, inhibits proliferation of human breast cancer cells via inducing apoptosis. *Nutrition and Cancer*. 2013;**65**:891–899. doi:10.1080/01635581.2013.802001.
- [30] Hongzhong L, Bing Y, Jing H, Tingxiu X, Xuedong Y, Jingyuan W, Fuling L, Li Z, Hongyuan L, Guosheng R. Naringin inhibits growth potential of human triple-negative breast cancer cells by targeting β -catenin signaling pathway. *Toxicology Letters*. 2013;**220**:219–228. doi:10.1016/j.toxlet.2013.05.006
- [31] Jeong HK, Jae KL. Naringenin enhances NK cell lysis activity by increasing the expression of NKG2D ligands on Burkitt's lymphoma cells. *Archives of Pharmacal Research*. 2015;**38**:2042–2048. doi:10.1007/s12272-015-0624-5
- [32] Balachandran C, Sangeetha B, Duraipandiyan V, Raj MK, Ignacimuthu S, Al-Dhabi NA, Balakrishna K, Parthasarathy K, Arulmozhi NM, Arasu MV. A flavonoid isolated from *Streptomyces* sp. (ERINLG-4) induces apoptosis in human lung cancer A549 cells through p53 and cytochrome c release caspase dependant pathway. *Chemico-Biological Interactions*. 2014;**224**:24–35. doi:10.1016/j.cbi.2014.09.019
- [33] Mi JC, Sanghyun L, Yong P, Jisun L, Ki HK. Neuroprotective effects of phytosterols and flavonoids from *Cirsium setidens* and *Aster scaber* in human brain neuroblastoma SK-N-SH cells. *Life Sciences*. 2016;**148**:173–182. doi:10.1016/j.lfs.2016.02.035
- [34] Panat NA, Singh BG, Maurya DK, Sandur SK, Ghaskadbi SS. Troxerutin, a natural flavonoid binds to DNA minor groove and enhances cancer cell killing in response to radiation, *Chemico-Biological Interactions*. 2016;**251**:34–44. doi:10.1016/j.cbi.2016.03.024
- [35] Reneta G, Dimitrina Z, Silviya R, Nikolay D, Spiro K, Valentin L, Vanio M. Cytotoxic and hepatoprotective effects of *Bupleurum flavum* flavonoids on hepatocellular carcinoma HEP-G2 cells. *British Journal of Pharmaceutical Research*. 2016;**11**:1–8. doi:10.9734/BJPR/2016/25785
- [36] Lokesh DN, Rit V, Jyotsana S, Spence F, Rhonda R, Sanjay A, Sharad SS. Anti-cancer effects of novel flavonoid vicenin-2 as a single agent and in synergistic combination with docetaxel in prostate cancer. *Biochemical Pharmacology*. 2011;**82**:1100–1109. doi:10.1016/j.bcp.2011.07.078
- [37] Zhao Q, Zhao M, Parris AB, Xing Y, Yang X. Genistein targets the cancerous inhibitor of PP2A to induce growth inhibition and apoptosis in breast cancer cells. *International Journal of Oncology*, 2016;**49**:1203–1210. doi:10.3892/ijo.2016.3588
- [38] Janet MP, Sankar NK, Raymond CB. Genistein inhibits human prostate cancer cell detachment, invasion and metastasis. *The American Journal of Clinical Nutrition*. 2014;**100**:431S–436S. doi:10.3945/ajcn.113.071290

- [39] Lu Z, Zhou R, Kong Y, Wang J, Xia W, Guo J, Liu J, Sun H, Liu K, Yang J, Mi M, Xu H. S-equol, secondary metabolite of natural anticancer isoflavone daidzein, inhibits prostate cancer growth in vitro and in vivo, though activating the Akt/FOXO3a pathway. *Current Cancer Drug Targets*. 2016;**16**:455–465. doi:10.2174/1568009616666151207105720
- [40] Bhardwaj V, Tadinada SM, Jain A, Sehdev V, Daniels CK, Lai JC, Bhushan A. Biochanin A reduces pancreatic cancer survival and progression. *Anticancer Drugs*. 2014;**25**:296–302. doi:10.1097/CAD.0000000000000044.
- [41] Victor K, Armelle TM, Eric CNN, Christophe CS, Maen Z, Augustin EN, Thomas E. Cytotoxicity of seven naturally occurring phenolic compounds towards multi-factorial drug-resistant cancer cells . *Phytomedicine*. 2016;**23**:856–863. doi:10.1016/j.phymed.2016.04.007
- [42] Yan-Xi Z, Hong Z, Cheng P. Puerarin: a review of pharmacological effects. *Phytotherapy Research*. 2014;**28**:961–975. doi:10.1002/ptr.5083
- [43] Sung-Gook C, Sang-Mi W, Seong-Gyu K. Butein suppresses breast cancer growth by reducing a production of intracellular reactive oxygen species. *Journal of Experimental & Clinical Cancer Research*. 2014;**33**:1–11. doi:10.1186/1756-9966-33-51
- [44] Moon DO, Kim MO, Lee JD, Choi YH, Kim GY. Butein suppresses c-Myc-dependent transcription and Akt-dependent phosphorylation of hTERT in human leukemia cells. *Cancer Letters*. 2009;**286**:172–179. doi:10.1016/j.canlet.2009.05.028
- [45] Yit CC, Das NP, Cytotoxic effect of butein on human colon adenocarcinoma cell proliferation, *Cancer Letters*. 1994;**82**:65–72. doi:10.1016/0304-3835(94)90147-3
- [46] Cui Z, Song E, Hu DN, Chen M, Rosen R, McCormick SA. Butein induces apoptosis in human uveal melanoma cells through mitochondrial apoptosis pathway. *Current Eye Research*. 2012;**37**:730–9. doi:10.3109/02713683.2012.671436
- [47] Xue B, Yaxin M, Guobin Z. Butein suppresses cervical cancer growth through the PI3K/AKT/mTOR pathway. *Oncology Reports*. 2015;**33**:3085–3092. doi:10.3892/or.2015.3922
- [48] Fu P, Chun-Wang M, Qin-Mei Z, Jian-Ping C, Liang X. Cytotoxic evaluation against breast cancer cells of isoliquiritigenin analogues from *Spatholobus suberectus* and their synthetic derivatives. *Journal of Natural Products*. 2016;**79**:248–251. doi:10.1021/acs.jnatprod.5b00774
- [49] Chun-Lin Y, Jian-Wen L, Dong-Zhi W, Yan-Hua L. Feng Qian. In vitro anti-tumor activity of 2',4'-dihydroxy-6'-methoxy-3',5'-dimethylchalcone against six established human cancer cell lines. *Pharmacological Research*. 2004;**50**:505–10. doi:10.1016/j.phrs.2004.05.004
- [50] Huang HY, Niu JL, Zhao LM, Lu YH. Reversal effect of 2',4'-dihydroxy-6'-methoxy-3',5'-dimethylchalcone on multi-drug resistance in resistant human hepatocellular carcinoma cell line BEL-7402/5-FU. *Phytomedicine*. 2011;**18**:1086–1092. doi:10.1016/j.phymed.2011.04.001

- [51] Victor K, Antoine HLN, Armelle TM, Benjamin W, Hippolyte KW, Pierre T, Thomas E. Cytotoxicity and modes of action of 4'-hydroxy-2',6'-dimethoxychalcone and other flavonoids toward drug-sensitive and multidrug-resistant cancer cell lines. *Phytomedicine*. 2014;**21**:1651–57. doi:10.1016/j.phymed.2014.08.001
- [52] Miranda CL, Stevens JF, Helmrich A, Henderson MC, Rodriguez RJ, Yang YH, Deinzer ML, Barnes DW, Buhler DR. Antiproliferative and cytotoxic effects of prenylated flavonoids from hops (*Humulus lupulus*) in human cancer cell lines. *Food and Chemical Toxicology*. 1999;**37**:271–285. doi:10.1016/S0278-6915(99)00019-8
- [53] Tetsuro Y, Masato K, Etsu T, Masaya I, Hiroyuki O, Yasuo M, Junko H, Mitsunori S, Toyomasa K. Xanthohumol suppresses oestrogen-signalling in breast cancer through the inhibition of BIG3-PHB2 interactions. *Scientific Reports*. 2014;**7355**:1–9. doi:10.1038/srep07355
- [54] Nadiah A, Muhammad NA, Swee KY, Kian LL, Wan YH, Mohd PA, Chai LH, Abdul RO, Jamil I, Noorjahan B. Flavokawain B induced cytotoxicity in two breast cancer cell lines, MCF-7 and MDA-MB231 and inhibited the metastatic potential of MDA-MB231 via the regulation of several tyrosine kinases *In vitro*. *BMC Complementary and Alternative Medicine*. 2016;**16**:1–14. doi:10.1186/s12906-016-1046-8
- [55] Fang SC, Hsu CL, Yu YS, Yen GC. Cytotoxic effects of new geranyl chalcone derivatives isolated from the leaves of *Artocarpus communis* in SW 872 human liposarcoma cells. *Journal of Agricultural and Food*. 2008;**56**:8859–8868. doi:10.1021/jf8017436
- [56] Sumiyoshi M, Taniguchi M, Baba K, Kimura Y. Antitumor and antimetastatic actions of xanthoangelol and 4-hydroxyderricin isolated from *Angelica keiskei* roots through the inhibited activation and differentiation of M2 macrophages. *Phytomedicine*. 2015;**22**:759–767. doi:10.1016/j.phymed.2015.05.005
- [57] Keiichi T, Kou M, Noriya T, Reiko N, Satoru A, Yumiko K, Motohiko U, Daisuke H, Toshihiro A, Takashi S. Xanthoangelol, a major chalcone constituent of *Angelica keiskei*, induces apoptosis in neuroblastoma and leukemia cells. *Biological and Pharmaceutical Bulletin*. 2005;**28**:1404–1407. doi:10.1248/bpb.28.1404
- [58] Victor K, Armelle TM, Bonaventure TN, Thomas E. Cytotoxicity of three naturally occurring flavonoid derived compounds (artocarpesin, cycloartocarpesin and isobavachalcone) towards multi-factorial drug-resistant cancer cells. *Phytomedicine*. 2015;**22**:1096–1102. doi:10.1016/j.phymed.2015.07.006
- [59] Kuete V, Mbaveng AT, Zeino M, Ngameni B, Kapche GD, Kouam SF, Ngadjui BT, Efferth T. Cytotoxicity of two naturally occurring flavonoids (dorsmanin F and poinsettifolin B) towards multi-factorial drug-resistant cancer cells. *Phytomedicine*. 2015;**22**:737–743. doi:10.1016/j.phymed.2015.04.007
- [60] Cazal CM, Choosang K, Severino VG, Fernandes JB, da Silva MF, Vieira PC, Nascimento MS, Almeida GM, Vasconcelos MH, Pakkong P, Pinto MM. Natural compounds with cell growth inhibitory activity in human tumor cell lines. *Anticancer Agents in Medicinal Chemistry*. 2013;**13**:1582–1589. doi:10.2174/1871520613666131125122857

- [61] Ladislaus K. Mdee, Samuel O. Yeboah, Berhanu M. Abegaz, rhuschalcones II–VI, five new bichalcones from the root bark of *Rhuspyroides*. Journal of Natural Product. 2003;**66**:599–604. doi:10.1021/np020138q
- [62] Yoo I, Yun B, Lee I, Ryoo I, Choung D, Han K. Three naphthalenes from root bark of *Hibiscus syriacus*. Phytochemistry. 1998;**47**:799–802. doi:10.1016/S0031-9422(97)00674-2
- [63] Heng-Xing L, Huan-Qin D, Hai-An F, Xiao-Ping D, Abiodun HA, Li-Xin Z, Yong-Xian C. Bioactive compounds from *Rumex* plants. Phytochemistry Letters. 2010;**3**:181–184. doi:10.1016/j.phytol.2010.05.005
- [64] Negera A, Fangfang P, Amra G, Jurgen G, Paul AF, Goran L, Kari R, Abiy Y, Mate E. Naphthalene derivatives from the roots of *Pentas parvifolia* and *Pentas bussei*. Journal of Natural Products. 2016;**79**:2181–2187. doi:10.1021/acs.jnatprod.6b00178
- [65] Hui Z, Zengjun G, Nan W, Wenming X, Ling H, Nan L, Yanxia H. Two novel naphthalene glucosides and an anthraquinone isolated from *Rumex dentatus* and their antiproliferation activities in four cell lines. Molecules. 2012;**17**:843–850. doi:10.3390/molecules17010843
- [66] Chandrasekar B, Veeramuthu D, Al-Dhabi NA, Stalin A, Balakrishna K, Ignacimuthu S, Tilton F. Isolation and characterization of Anthraquinone from *Streptomyces* sp. ERINLG-26 with anticancer activity against adenocarcinoma cell line COLO320. Applied Biochemistry and Microbiology. 2015;**51**:522–529. doi:10.1134/S0003683815050038
- [67] Chandrasekar B, Veeramuthu D, Yuvaraj A, Balachandran S, Nobuhiko E, Naif AA, Savarimuthu I, Yoko I, Akinao O, Paramasivan TP. Isolation and characterization of 2-hydroxy-9,10-anthraquinone from *Streptomyces olivochromogenes* (ERINLG-261) with antimicrobial and antiproliferative properties. Brazilian Journal of Pharmacognosy. 2016;**26**:285–295. doi:10.1016/j.bjp.2015.12.003
- [68] Balachandran C, Emi N, Arun Y, Yamamoto N, Duraipandiyan V, Inaguma Y, Okamoto A, Ignacimuthu S, Al-Dhabi NA, Perumal PT. In vitro antiproliferative activity of 2,3-dihydroxy-9,10-anthraquinone induced apoptosis against COLO320 cells through cytochrome c release caspase mediated pathway with PI3K/AKT and COX-2 inhibition. Chemico-Biological Interactions. 2016;**249**:23–35. doi:10.1016/j.cbi.2016.02.016
- [69] Shi YQ, Fukai T, Sakagami H, Kuroda J, Miyaoka R, Tamura M, Yoshida N, Nomura T. Cytotoxic and DNA damage-inducing activities of low molecular weight phenols from rhubarb. Anticancer Research. 2001;**21**:2847–2853.
- [70] Surat L, Wisanu M, Sorwaporn K. Antimalarial and cytotoxic phenolic compounds from *Cratoxylum maingayi* and *Cratoxylum cochinchinense*. Molecules. 2009;**14**:1389–1395. doi:10.3390/molecules14041389
- [71] Sompong B, Chatchanok K, Chanita P, Akkharawit K, Kan C. Antibacterial and cytotoxic xanthenes from the roots of *Cratoxylum formosum*. Phytochemistry. 2006;**67**:723–727. doi:10.1016/j.phytochem.2006.01.007

- [72] Frank M, Niemann H, Bohler P, Stork B, Wesselborg S, Lin W, Proksch P. Phomoxanthone A-from mangrove forests to anticancer therapy. *Current Medicinal Chemistry*. 2015;**22**:3523–3532. doi:10.2174/0929867322666150716115300
- [73] Manqin F, Samuel XQ, Yujuan X, Jijun W, Yulong C, Yuanshan Y, Gengsheng X. A new xanthone from the pericarp of *Garcinia mangostana*. *Natural Product Communications*. 2013;**8**:1733–1734.
- [74] Victor K, Louis PS, Judith LNO, Hugues F, Benjamin W, Thomas E. Cytotoxicity and modes of action of three naturally occurring xanthones (8-hydroxycudraxanthone G, morusignin I and cudraxanthone I) against sensitive and multidrug-resistant cancer cell lines. *Phytomedicine*. 2014;**21**:315–322. doi:10.1016/j.phymed.2013.08.018
- [75] Dya FD, Suresh A, Shigetoshi K, Hiroyuki M, Yasuhiro T. Hepta-oxygenated xanthones as anti-austerity agents from *Securidaca Longepedunculata*. *Bioorganic & Medicinal Chemistry*. 2013;**21**:7663–7668. doi:10.1016/j.bmc.2013.10.027
- [76] Asano J, Chiba K, Tada M, Yoshii T. Cytotoxic xanthones from *Garcinia hanburyi*. *Phytochemistry*. 1996;**41**:815–820. doi:10.1016/0031-9422(95)00682-6
- [77] Shadid KA, Shaari K, Abas F, Israf DA, Hamzah AS, Syakroni N, Lajis NH. Cytotoxic caged-polyprenylated xanthonoids and a xanthone from *Garcinia cantleyana*. *Phytochemistry*. 2007;**68**:2537–2544. doi:10.1016/j.phytochem.2007.05.024
- [78] Soleh K, Shu-Geng C, Xiao-Hua W, Jagadese JV, Tony S, Masdianto SH, Goh, Keng-Yeow Sim. Lateriflorone, a cytotoxic spiroxalactone with a novel skeleton, from *Garcinia lateriflora* BL. *Tetrahedron Letters*. 1999;**40**:157–160. doi:10.1016/S0040-4039(98)80046-4
- [79] Zhang HZ, Kasibhatla S, Wang Y, Herich J, Guastella J, Tseng B, Drewe J, Cai SX. Discovery, characterization and SAR of gambogic acid as a potent apoptosis inducer by a HTS assay. *Bioorganic and Medicinal Chemistry*. 2004;**12**:309–317. doi:10.1016/j.bmc.2003.11.013
- [80] Xiaohua W, Shugeng C, Sweehock G, Anne H, Benny KHT. Mitochondrial destabilisation and Caspase-3 activation are involved in the apoptosis of jurkat cells induced by gaudichaudione A, a cytotoxic xanthone. *Planta Medica*. 2002;**68**:198–203. doi:10.1055/s-2002-23142
- [81] Zhengxiang X, Hong Z, Danqing X, Yuanzhi L, Wenwei F, Hongsheng T, Peng C, Ling Y, Hongxi X. Xanthones from the leaves of *Garcinia cowa* induce cell cycle arrest, apoptosis and autophagy in cancer cells. *Molecules*. 2015;**20**:11387–11399. doi:10.3390/molecules200611387
- [82] Josiane AM, Ivania TAS, Thiago LA, Silvana MOS, Cleuza CS, Lucas URC, Ana LTGR, Joao EC, Debora BV, Celso VN, Debora BS, Vanessa K, Ricardo TF, Armando MP. Oncibauerins A and B, new flavanones from *Oncidium baueri* (Orchidaceae). *Phytochemistry Letters*. 2014;**9**:141–148. doi:10.1016/j.phytol.2014.06.004

- [83] Qiao-Li L, Ling-Ling L, Xiaodong C, Nuo-Shu Z, Jin-Ao D. Bioactive cis-stilbenoids from the tubers of *Scirpus yagara*. *Fitoterapia*. 2013;**84**:170–73. doi:10.1016/j.fitote.2012.11.012
- [84] Nen-Ling Z, Yue-Hui Z, Ri-Ming H, Man-Qin F, Zhi-Wei S, Jia-Zhong C, Ying-Jie H, Sheng-Xiang Q. Two new stilbenoids from *Cajanus cajan*. *Zeitschrift fur Naturforschung B*. 2012;**67b**:1314–18. doi:10.5560/ZNB.2012-0184
- [85] Huang CP, Au LC, Chiou RY, Chung PC, Chen SY, Tang WC, Chang CL, Fang WH, Lin SB. Arachidin-1, a peanut stilbenoid, induces programmed cell death in human leukemia HL-60 cells. *Journal of Agricultural and Food Chemistry*. 2010;**58**:12123–29. doi:10.1021/jf102993j
- [86] Jie M, Duowei W, Zhuo W, Wei T, Xianjing L, Jingjing D, YunW, Hongbao Y, Linjun Y, Yan C, Jinsong B, Zhen C, Yong Y. Combretastatin A-1 phosphate, a microtubule inhibitor, acts on both hepatocellular carcinoma cells and tumor-associated macrophages by inhibiting the Wnt/ β -catenin pathway. *Cancer Letters*. 2016;**80**:134–143. doi:10.1016/j.canlet.2016.06.020
- [87] Phanruethai P, Suwannee S, Chulabhorn M, Somsak R, Vilailak P. Cytotoxic and cancer chemopreventive properties of prenylated stilbenoids from *Macaranga siamensis*. *Tetrahedron*. 2015;**71**:5562–5571. doi:10.1016/j.tet.2015.06.058
- [88] Ming HY, Le C, Mei HL, Xiang HZ, Ya BY, Zhong TD. Three new phenanthrenes from *Monomeria barbata* Lindl. *Chinese Chemical Letters*. 2010;**21**:325–328. doi:10.1016/j.cclet.2009.11.006
- [89] Junju X, Hong Y, Chen Q, Yanli Z, Ying L, Yegao C. Two new biphenanthrenes with cytotoxic activity from *Bulbophyllum odoratissimum*. *Fitoterapia*. 2009;**80**:381–84. doi:10.1016/j.fitote.2009.05.007
- [90] Young-Won C, William PJ, Ismail R, Soedarsono R, Leonardus BSK, Hee-Byung C, Norman RF, Geoffrey AC, Steven MS, John MC, Kinghorn AD. Cytotoxic lignans from the stems of *Helicteres hirsuta* collected in Indonesia. *Phytotherapy Research*. 2006;**20**:62–65. doi:10.1002/ptr.1806
- [91] Won SS, Lalita S, Sun YK, Sang UC, Kang RL. Bioactive lignan constituents from the twigs of *Sambucus williamsii*. *Bioorganic & Medicinal Chemistry Letters*. 2016;**26**:1877–1880. doi:10.1016/j.bmcl.2016.03.023
- [92] Shan Y, Jun-Jiang L, Jian-Qiang Z, Yan L, Dong W, Chong-Ren Y, Min X, Ying-Jun Z. New cytotoxic lignan glycosides from *Phyllanthus glaucus*. *Natural Product Research*. 2016;**30**:419–425. doi:10.1080/14786419.2015.1023198