We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Foods with Functional Properties and Their Potential Uses in Human Health

Ricardo Tighe-Neira, Miren Alberdi,
Patricio Arce-Johnson, Jesús L. Romero-Romero,
Marjorie Reyes-Díaz and
Claudio Inostroza-Blancheteau

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/67077>

Abstract

Vegetables and fruits have been a part of human diet since ancient times; nevertheless, as countries develop, its population's feeding habits change and tend to have a diet poor in vegetables and fruits, with well-known consequences. Several food plant products with massive consumption and within the reach of the population are products such as artichoke, leek, hot chili pepper, coriander, kiwifruit, sweet orange, highbush blueberry, and maracuyá to name a few. They have many beneficial properties principally by its content of phytochemicals with high impact on human health, beyond nutritional support. The phytochemicals are bioactive compounds such as vitamins, carotenoids, phenolic acid, and flavonoids, which contribute to antioxidant capacity and as a whole prevent chronic nontransmissible diseases such as: diabetes, high blood pressure, high cholesterol in blood, cardiovascular risks, among others. This relationship between food plant for human consumption and its impacts on human health is discussed in this chapter, highlighting coriander and kiwifruit by its wide range of benefits.

Keywords: phytochemicals, antioxidant capacity, polyphenols, chronic diseases, healthy feeding habits, life style

1. Introduction

Foods are neither inherently good nor bad. Rather, good or bad eating habits, as well as other factors such as smoking and physical activity, all influence human health. If we desire a healthy lifestyle and wish to avoid chronic nontransmissible disorders such as diabetes, high

levels of cholesterol, cardiovascular diseases, etc., foods, especially those that are functional, are only a part of the solution [1].

Despite a wealth of information, there is no universal definition about what constitutes a functional food. However, there is consensus concerning central concepts, which are associated with their benefit for human health beyond their traditional nutrients [2]. Along the same lines, the importance of phytochemicals as a class of biologically active metabolites in plants is accepted [3]. When discussing “potential use for human health” to refer to a particular plant, preliminary evidence on its outstanding phytochemical content must already exist, which means it can be used in the future as a source to investigate more profoundly its beneficial implications in human health.

Some processes such as cooking alter the content and composition of phytochemicals present in vegetables, reducing their concentrations by thermal degradation or augmenting their concentration with respect to the raw material. However, these effects are varying with the cooking method and type of phytochemical [4]. These, together with the growing consumption of fiber, are the principal reasons to recommend the regular intake of fresh vegetables [5].

Functional foods may be plant or animal products, that are fresh, semi-processed or processed, but in this chapter we will refer mainly to fresh plants and their properties beyond their nutritional characteristics. In addition, we will also discuss the existence of several common horticultural and fruit plants that are widely available and consumed by the human population, whose functional properties have yet to be systematized and categorized. Vegetables with high functional interest such as artichoke, leek, hot chili pepper, and coriander, as well as fruit plants such as kiwifruit, sweet orange, and highbush blueberry are considered [6–9]. Within the species discussed in this chapter, only artichokes must be consumed cooked, whilst the others may be eaten fresh.

We believe that the updated information about plants with characteristics as functional foods responds to a need of the population and scientists to learn more about healthy habits and how consumption of natural foods can improve their quality of life.

2. Horticultural species with functional properties and their potential use in human health

The properties and features of some horticultural species that have beneficial effects on human health are mentioned here. It is important to note that the artichoke, leek, hot chili pepper, and coriander are considered in this section, because these species are widely consumed by the human population.

2.1. Artichoke (*Cynara cardunculus* var. *scolymus* and *Cynara scolymus*)

These species belong to the *Asteraceae* family (**Figure 1**) are native to the Mediterranean region [10] and grow in many parts of the world [11]. *Cynara scolymus* is cultivated due to its large immature flower heads, which have special functional and nutraceutical characteristics, and a

high antioxidant capacity, with more than 9000 μmol trolox equivalents (TE) 100 g^{-1} FW [12]. This antioxidant capacity is due to its high content of total polyphenols like caffeoylquinic acids and flavonoids [7]. These polyphenols are present in flower heads, with values ranging from 4.8 to 29.8 mg g^{-1} FW in different Italian varieties [7, 13, 14]. It is important to mention that these values are not only dependent on the genetic background, as the interaction genotype-environment also has an influence [7]. Therefore, antioxidant content may be affected by agricultural practices, because in different locations, the same variety may have different antioxidant capacities.

Figure 1. Immature flower head of artichoke.

Choleretic and hypocholesterolemic activities, due to the presence of chlorogenic acid, cynarin, and lutein, have been reported in clinical studies, which demonstrated the effect of leaf extracts on the inhibition of the biosynthesis of cholesterol in rat hepatocytes [15]. Furthermore, these extracts prevented necrosis in rat hepatocytes provoked by hydroperoxides indicated for treatment of dyspepsia, or dyskinesia of the bile ducts, as well as disorders in the assimilation of fats in humans [16–18]. The nutraceutical and therapeutic actions of several metabolites are summarized in **Table 1**. The positive effects of ingesting *C. cardunculus* flower heads have been widely demonstrated; nevertheless, artichoke leaves and external parts of bracts may be used in industrial processes to obtain functional metabolites for use in human health.

Species	Molecule	Part of plant	Specific function for human health	References
<i>Cynara cardunculus</i> L. var. <i>scolymus</i> L. / <i>Cynara scolymus</i>	Caffeoylquinic acids, flavonoids, and cynarin	Flower head and leaves	Antioxidant capacity	[7, 12–14]
	Chlorogenic acid and cynarin	Flower head and leaves	Choleretic, hepatoprotective	[168, 169, 170]
	Luteolin	Flower head and leaves	Hypocholesterolemic	[171]
	Inulin	Flower head, stem and leaves	Stimulates the intestinal flora, hypocholesterolemic	[170]
<i>Allium ampeloprasum</i> Var. <i>porrum</i> / <i>Allium porrum</i>	Polyphenols	White part and green leaves	Antioxidant capacity	[28]
	Ascorbate	White part and green leaves	Growth and tissue repair, antioxidant	[28, 30]
	Steroidal saponines	White part	Antiinflammatory, gastroprotective and antiulcerogenesis	[31, 32]
	Methanolic extracts	White part and green leaves	Antimicrobial	[35]
	Steroidal saponines/polyphenols	White part	Cytotoxic activity: a potential anticancer agent	[33, 35]
	Hydroalcoholic extracts	White part and green leaves	Hypolipidemic/hypocholesterolemic	[172, 173]
<i>Capsicum annuum</i> L.	Polyphenols	Fruit	Antioxidant capacity	[37, 174]
	Dietary fiber	Fruit	Anticarcinogenic, prevents diabetes, hypocholesterolemic	[9, 175]
<i>Coriandrum sativum</i> L.	Essential oil, aqueous, methanolic and ethanolic extracts	Fruit and leaves	Prevention of several chronic degenerative disorders	[8, 44, 48]
	Essential oils	Fruit and leaves	Anticancer activity	[176, 177]
	Extracts	Seeds	Hypocholesterolemic	[51]
	Extracts	Seeds	Antidiabetic	[49]
	Diethyl ether and aqueous extracts	Fruit, seeds and leaves	Anxiolytic, sedatives, antidepressant	[51, 54, 56]
	Aqueous, ethanolic and chloroformic extracts	Aerial parts	Analgesic	[55]
	Ethanolic extract	Root	Antimicrobial	[48]
	Extracts	Aerial parts	Antioxidant activity	[47]
	Hydroalcoholic extract	Aerial parts	Anticonvulsant	[15]
	Ethanolic extract	Seeds	Cognitive effects (improves learning in the long term)	[53]

Table 1. Selected plant species and their compounds that are beneficial for human health.

Inositol and inulin are soluble carbohydrates, which are present in external bracts of artichokes. In the case of inositol (chiro-scyllo- and myo-inositol), values fluctuate from 6.7 to 9.3 mg g⁻¹ DW while for inulin they fluctuate from 69.8 to 114.6 mg g⁻¹ DW [16]. These values are higher than the ones reported by Hernández-Hernández et al. [17] in edible bracts of artichoke. Regarding the beneficial properties, inositols have been used in treatments against diabetes mellitus [18]. In this sense, Crawford et al. [19] reported that inositol prevents diabetes mellitus in pregnant women and concluded that myo-inositol shows promising results by preventing the onset of the disease. Furthermore, inulin has been associated with some beneficial functional properties, as it can be a good source of carbohydrates and fiber, associated with positive effects in the prevention of colon cancer [20, 21]. In addition, prebiotic properties and effects on the absorption of calcium have been reported [20, 21]. In this sense, research on mineral absorption of calcium and magnesium concluded that inulin can reduce risk for osteoporosis by increasing their absorption [22].

2.2. Leek (*Allium ampeloprasum* var. *porrum* or *Allium porrum*)

It has been reported that this species has a substantial nutraceutical and functional properties (see **Table 1**) and is cultivated in Asia, America, and Europe, especially in the Mediterranean region [23, 24]. *A. ampeloprasum*, belongs to subgenus *Allium* section *Allium* (**Figure 2**), is considered a "complex species" due to different ploidy levels and genome constitution, and diploid, tetraploid, and octoploid accessions have been described [25, 26]. In fact, Hirschegger et al. [27] suggested that molecular evidence could be used to consider this species as a tetraploid horticultural group together with other *Allium* genus species. This condition is particularly important in accounting for its nutritional, functional, and nutraceutical characteristics [28].

Figure 2. Green leaves and white part of leek.

The principal beneficial properties of the *Allium* genus are mainly due to the presence of many sulfur compounds containing bioactive constituents, including: dimethyl disulfide, methyl propenyl disulfide, propyl propenyl disulfide, dimethyl trisulfide, methyl propyl trisulfide, methyl propenyltrisulfide, S-methyl cysteine sulfoxide, S-propyl cysteine sulfoxide, S-propenyl cysteine sulfoxide, and N-(γ -glutamyl)-S-(E-1-propenyl) cysteine [6].

In the evaluation of 30 leek cultivars, the content of total phenols varied from 5 to 15 mg gallic acid equivalents (GAE) g^{-1} DW for whole plants [28]. Other studies reported values of 5.5–6.0 mg GAE g^{-1} DW in whole leeks [29]. These differences in the total phenolic content could be attributed to the genetic variability of this species and agricultural systems [28]. Moreover, in the same study, the oxygen radical absorbance capacity (ORAC) was evaluated, where the green leaves possessed 82–135 $\mu\text{mol TE g}^{-1}$ DW, whereas the white part contained just 27–88 $\mu\text{mol TE g}^{-1}$ DW. Additionally, Vandekinderen et al. [30] determined the total vitamin C content (ascorbic acid + dehydroascorbic acid) whose values reached 9.65 mg 100 g^{-1} FW in whole leeks. Bernaert et al. [28] reported 5.54 mg ascorbic acid (AA) g^{-1} DW in whole leeks and higher values in green leaves than in white parts (2.77–8.52 mg AA g^{-1} DW and 0.89–3.55 mg AA g^{-1} DW, respectively). Values of polyphenols, AA, and antioxidant activity may be influenced by the season of year, genetic characteristics, and biotic and abiotic factors during vegetative growth, as well as agricultural practices.

The antiinflammatory, gastroprotective, and cytotoxic activities of organosulfate compounds, saponins, particularly steroidal saponines, have been well documented in *A. ampeloprasum* var. *porrum* [31–33].

Another property that is exclusive to the *Allium* genus is their antimicrobial activity (see **Table 1**). This has been reported since ancient times, and leeks have been used to treat wounds and respiratory diseases, as well as acting as an antibacterial agent due to the presence of alliin-containing structures [34, 35]. Polyphenols of methanolic extracts of green leaves and white parts of *A. porrum* are potent against Gram-positive (*Staphylococcus aureus* ATCC 25923 and *Bacillus subtilis* ATCC 6633) and Gram-negative (*Klebsiella pneumoniae* ATCC 13883, *E. coli* ATCC 25922, *Proteus vulgaris* ATCC 13315 and *Proteus mirabilis* ATCC 14153) bacteria, as well as fungal species (*Candida albicans* ATCC 10231 and *Aspergillus niger* ATCC 16404) [28]. These authors affirm a negative effect of methanolic extracts of *A. porrum* on Hep2c, L2OB, and RD cell cultures. This cytotoxic activity could indicate a future use of these natural biological compounds in human health.

2.3. Hot chili pepper (*Capsicum annuum* L. var. *longum*)

The fruit of this species (**Figure 3**), immature or mature and leaves, contains at least two groups of bioactive compounds of significance for human health, polyphenols and carotenoids. The polyphenol content is variable but reaches over 20 mg GAE g^{-1} DW in mature and dried fruits [36], and 40 mg GAE g^{-1} DW in leaves [37]. The polyphenols of fruits have a total antioxidant capacity of 26.6–44.4 $\mu\text{mol TE g}^{-1}$ DW, depending on the variety [36].

According to Serrano et al. [38], the small intestine has around 25% of bioavailability of total polyphenols.

Figure 3. Immature fruit of red hot chili pepper.

Regarding the total carotenoids present in different varieties of red hot chili peppers, Hervert-Hernández et al. [36] indicated values from 87.6 to 373.3 mg 100 g⁻¹ DW. In addition, the same authors determined that the bioavailability of chili carotenoids in the small intestine ranges from 20 to 50% of the total content, depending on the variety [36]. In addition, *C. annuum* is also a good source of vitamin C (up to 26.5 mg g⁻¹ DW) [39]. Minerals, mainly potassium and magnesium, as well as dietary fiber, reducing sugars (around 20% DW), and antimicrobial activity useful for functional food production are also characteristic of this species [9]. Regarding the antimicrobial activity, De et al. [40] identified three pathogens that are susceptible to ethyl alcohol extracts of chili: *Bacillus subtilis* ATCC6633 (minimum inhibitory concentration (MIC) 10–25 mg mL⁻¹), *Escherichia coli* ATCC10536 (MIC 25–50% mg mL⁻¹), and *Saccharomyces cerevisiae* ATCC 9763 (MIC 2–5% mg mL⁻¹). In addition, an antimicrobial activity in *Staphylococcus aureus* ATCC14154, *Escherichia coli* ATCC-1698, *Pseudomonas aeruginosa* ATCC-23993, *Candida albicans* ATCC-14053, and *Sarcina lutea* (Collection of Microbiology Laboratory of Chemical Engineering Department, Institut Teknologi Bandung) has also been reported using ethanolic extracts of chili [41]. These authors indicate that capsaicin may be one of the main responsables of microorganism inhibition. Likewise, Huang et al. [42] affirm that developments of *Staphylococcus aureus*, *Saccharomyces cerevisiae*, and *Aspergillus niger* were negatively affected by capsaicin.

2.4. Coriander (*Coriandrum sativum* L.)

This species of the *Apiaceae* family (**Figure 4**) is usually cultivated throughout the year in diverse edafoclimatic areas [43]. It is native to Italy and is currently propagated in several Mediterranean regions of Europe, as well as in America and Asia [44]. It is consumed principally fresh, either alone or in salads. This species has multiple human health benefits and has considerable potential as a functional horticulture species (see **Table 1**). *C. sativum* contains essential oils in seeds and in the pericarp whose content and composition appear to be dependent on biological and geographical traits [35]. The oil content is approximately 1% of seed weight of which linalool is the major component (73%) [44–46]. In the stem and immature leaves, the most important compounds are essential oils, flavonoids (quercetin, kaempferol, and acacetin), phenolic acids (vanillic acid, p-coumaric acid, syringic acid, p-OH benzoic acid, cis-ferulic acid, and trans-ferulic acid), and polyphenols [8]. The principal phytochemicals are β -carotene ($5.1 \text{ mg g}^{-1} \text{ DW}$), AA ($1.16 \text{ mg } 100 \text{ g}^{-1} \text{ FW}$), total phenolics ($2.05 \text{ mg GAE } 100 \text{ g}^{-1} \text{ DW}$), and antioxidants ($1.12 \text{ mg GAE } 100 \text{ g}^{-1} \text{ DW}$) [47].

Figure 4. Stem and immature leaves of coriander.

According to Kumar et al. [48], ethanolic extracts of fresh coriander roots contain alkaloids, flavonoids, terpenoids, sterols, carbohydrates, saponins, and phenolic compounds. This extract and its fractions possess significant antibiotic activity against *Staphylococcus aureus*, *Bacillus cereus*, *Salmonella typhi*, and *Klebsiella*.

The antihyperglycemic (antidiabetic) activity of coriander has been studied by several authors. Deepa and Anuradha [49] analyzed the effects of coriander seed extracts in rats, which showed decreases by 44% in blood glucose and by 58% in glycosylated hemoglobin levels with respect to untreated rats. At the same time, the insulin level in plasma increased to 40%. They also reported beneficial effects in kidney and pancreas. Moreover, *C. sativum* seeds have hypocholesterolemic properties in rats, increasing hydroxymethyl glutaryl CoA (HMG-CoA) reductase and plasma lecithin cholesterol acyl transferase activities [50, 51].

Additional properties such as analgesic, anticonvulsive, anxiolytic, sedative, antidepressant, and cognitive effects of coriander have been tested *in vivo* in mice. In many cases, its effect was comparable with equivalent standardized doses of the typical drugs used to treat these diseases, elegantly demonstrating the beneficial properties of coriander in human health [15, 52–56].

3. Fruit species with functional properties and their potential use in human health

Fruits, in addition to horticultural species, constitute a group of foods for humans with important functional characteristics. In this chapter, we consider kiwifruit, sweet orange, and high-bush blueberry given their extensive geographical distribution, consumption, and richness in biocompounds with nutraceutical properties.

3.1. Kiwifruit (*Actinidia deliciosa* [A. Chev] C.F Liang et A.R Ferguson/*Actinidia chinensis* [Planch])

This species originated in Asia [56] and belongs to the *Actinidiaceae* family (**Figure 5**). The *Actinidia* genus includes 66 species, but only four are cultivated for fruit production. Of these, *A. deliciosa* and *A. chinensis* are the most accepted by consumers worldwide [58, 59]. Here, we will refer to both species indistinctly. Kiwifruits have multiple sensory, nutritional, and phytochemical properties and are rich in dietary fiber, acids, phenols, and vitamins [57, 60] (see **Table 2**). These contribute to antioxidant activity, which varies with the variety and the part of fruit consumed. For example, Soquetta et al. [57] found higher values of antioxidant activity measured by the ferric reducing ability of the plasma (FRAP) method as well as carotenoids, flavonoids, and vitamin C in flour of the Monty variety compared to the Bruno variety. Moreover, the same authors found the highest content of these compounds in flours from kiwifruit skin compared to flour from kiwi fruit bagasse, reporting values from 59 to 189 mg AA in 100 g of kiwifruit flour, almost double that found in oranges and strawberries [61], and 200–1200 mg GAE 100 g⁻¹ for phenolic compounds. D'Evoli et al. [60] indicated that in the total fresh kiwifruit, the content of oxalic acid was 8 mg 100 g⁻¹ FW, while citric and malic acid contents were 1.2 and 0.24 g 100 g⁻¹ FW, respectively. Furthermore, the same authors indicated that kiwifruits contain 90 mg GAE 100 g⁻¹ FW of the total polyphenols, 0.2 mg 100 g⁻¹ FW of lutein, and 0.06 mg 100 g⁻¹ FW of β -carotene, in addition to α -tocopherol, γ -tocopherol, and γ -tocotrienol which represent 0.9, 0.04, and 0.12 mg 100 g⁻¹ FW, respectively. All these compounds give kiwifruit

strong antioxidant properties that contribute to protect cells against oxidative damage [62]. Its antioxidant capacity (ORAC) varies from 0.06 to 1.4 $\mu\text{mol TE } 100 \text{ g}^{-1} \text{ FW}$ of fruit, depending on the hydrophilic or lipophilic fraction used in the analysis [60]. Lee et al. [63], using the same method but expressed as vitamin C equivalents (VCE), reported values from 595.7 to 2662.7 VCE $100 \text{ g}^{-1} \text{ FW}$. Clinical studies indicate that the uptake of vitamin C derived from kiwifruit reaches 40% in humans, similar to the rate of synthetic vitamin C uptake [64, 65]. This was tested by Vissers et al. [66] in a mouse model, where they found highly effective delivery to tissues. Together with vitamin C uptake, several other nutrients and beneficial phytochemicals are consumed, with synergistic effects, among them, of iron (Fe). A clinical study with women [67, 68] indicated that a breakfast fortified with Fe, when consumed with kiwifruit, can improve Fe content in women with low Fe stores; this may be related with the high values of AA, lutein, and zeaxanthin of kiwifruit.

Figure 5. Fruits of *Actinidia deliciosa* commonly name kiwifruit.

The dietary fiber content of kiwifruit is 2–3.39 g $100 \text{ g}^{-1} \text{ FW}$ [60, 69] of which 25–30% is found in the flour skin and fruit bagasse [57]. These values are higher than several of the widely consumed fruits such as orange, apple, banana, strawberries, and blueberries [61]. Thus, kiwifruits contain sufficient fiber thus improving digestive performance, ameliorating digestive transit, alleviating constipation, and irritable bowel syndrome [60, 70–72]. Studies performed in rat have reported that kiwifruits improve digestion of the principal proteins of beef muscle, soy protein, gelatin, and gluten [73]. This is related with its content of actinidin, a very active proteolytic enzyme, which acts in concert with the gastric and intestinal proteases, pepsin, and pancreatin, generating an increment in protein digestion in the gastric and intestinal tracts [74, 75].

Species	Molecule	Part of plant	Specific function for human health	References
<i>Actinidia deliciosa</i> (A. Chev) C.F Liang et A.R Ferguson/ <i>actidinia chinensis</i> Planch.	Dietary fiber	Fruit and flour skin and bagasse of fruit	Alleviate constipation/laxative	[57, 60, 71]
	Actinidin	Whole fruit	Increment on protein digestion	[73 – 75]
	Polyphenols, flavonoids Vitamin C, E, lutein, zeaxanthin and other phytochemicals	Fruit and flour skin and bagasse of fruit	Antioxidant activity/ reduces cellular damage from oxidative stress	[57, 62, 71, 86]
	Vitamin C	Fruit	Positive synergic effect with nutrients and phytochemicals	[61]
	Essential amino acids, linolenic acid, folic acid, vitamins A, B6, B12, C, and E, minerals like Zn, Cu, Fe, and Se.	Fruit	Strengthen the immune system	[82, 84]
	Phytosterols and ursolic acid/independent or synergic actions of polyphenols, vitamin C, and E	Fruit	Improve lipid profile in men/reduction of risk of cardiovascular diseases	[61, 76, 77, 79, 86]
	Unidentified	Fruit	Antihypertensive	[79, 80]
	Unidentified	Fruit	Antithrombotic	[78]
	Vitamin C and inositol and other unidentified compounds	Fruit	Antiinflammatory activities	[71, 178]
	Phytosterols and ursolic acid	Fruit	Inhibit carcinogenesis processes	[86]
<i>Citrus sinensis</i> (L.) Osbeck	Fiber	Fruit	Antiirritable bowel syndrome	[72]
	Unidentified	Juice and ethanolic peel extract	Antihyperglycemic activity	[89, 104, 105]
	Flavonoids and other polyphenols	Juice	Antihypercholesterolemic activity	[107, 110, 111]
	Flavonoids other than anthocyanins	Juice	Antithrombotic	[112]
	Flavonoids and other polyphenols	Juice	Cardioprotective	[108, 111]
	Flavones	Juice	Enhanced antioxidant defense system	[114]

Species	Molecule	Part of plant	Specific function for human health	References
<i>Vaccinium corymbosum</i> L.	Unidentified	Juice	Antiinflammatory activity	[119, 120]
	Flavones, flavonoids and flavonols	Juice	Microbial activity	[101, 122]
	Unidentified	Juice	Hypoglycemic activity	[135]
	Unidentified	Juice	Antiinflammatory	[140]
	Interaction phenolic compounds	Fruit and leaf aqueous extract	Antimicrobial	[138]
	Anthocyanins and other phytochemicals	Fruit	Modulation of vascular function	[178, 179]
	Anthocyanins and other polyphenols	Extract hydroalcoholic of fruit	Cytotoxic activity	[142]
	Antioxidant action	Fruit	Antiatherogenic effect/hypocholesterolemic	[144]

Table 2. Selected fruit species and their compounds that are beneficial for human health.

Kiwifruits possess hypocholesterolaemic activity in hypercholesterolemic men [61]. This property may be related with the expression of the *Taq1B* gene in response to the consumption of kiwifruits, which modulates the content of lipids in blood plasma and has been associated with a reduced risk of cardiovascular diseases [76, 77]. However, other researchers did not find the same effect on cholesterol levels, but concluded that the consumption of two or three fruits per day can reduce levels of blood triglycerides by 15%, compared with the control [78]. Similar clinical studies have demonstrated that kiwifruits (two or three per day) can reduce blood pressure in male smokers [79], possibly related with 11% reduction in angiotensin-converting enzyme (ACE) activity. This finding is considered relevant because it is very difficult to modulate hypertension by diet [79, 80]. In addition, a clinical study revealed that daily consumption of kiwifruit produces a reduction of 15% in platelet aggregation, which can be understood as antithrombotic activity [79, 81]. Nonetheless, Brevik et al. [81] discussed this effect because it may be influenced by the rate of kiwifruit consumption.

Additionally, Hunter et al. [82] and Skinner [83] affirm that kiwifruits have an important function in the modulation of the immune system. In this context, Hunter et al. [62] indicate that kiwifruit contributes significantly to lessening upper respiratory tract infections, head congestion, and sore throats in older individuals. Even though, there is a large source of variation in immune function, the nutrient status of this fruit is crucial. The most important phytochemicals present in kiwifruit include essential amino acids, linolenic acid, folic acid, vitamins A, B6, B12, C, and E, and minerals such as zinc, copper, iron, and selenium [84]. Given the type and content of phytochemicals, beneficial immune effects are not unexpected [82], although the mechanisms and the specific molecules underlying these effects are unknown. Moreover,

preliminary studies under *in vitro* and *ex vivo* conditions found a protective effect of kiwifruit over oxidative damage of DNA, which may be interpreted as inhibition of the carcinogenesis process [85, 86]. Subsequently, Collins et al. [87] determined that kiwifruit consumption could protect against oxidative DNA damage protection in humans and *ex vivo* by both increasing the antioxidant status in the plasma, and stimulating DNA repair.

3.2. Sweet orange (*Citrus sinensis* (L.) Osbeck.)

The sweet orange is one of most economically important fruits in worldwide [88, 89]. It is believed that the *Citrus* genus is native to Asia, specifically from Southern China (Yunnan), which may be the origin and point of distribution of several contemporaneous *Citrus* species [90]. *Citrus sinensis* (L.) Osbeck belongs to the *Rutaceae* family and is believed to be a backcross hybrid between pummelo and mandarin (**Figure 6**) [91, 92]. The sweet orange species have several cultivated varieties, some of which are mentioned by Grosso et al. [93], but in this chapter we will discuss this species without distinguishing between varieties.

Figure 6. Fruits of *Citrus sinensis* commonly name sweet orange.

The sweet orange harbors several interesting phytochemical compounds that play an important role in human health (see **Table 2**). These include vitamins and polyphenols such as hesperidin, gallic acid, sinapic acid, caffeic acid, p-hydroxybenzoic acid, vanillic acid, narirutin,

naringin, p-cumaric, and ferulic acid [93–96]. Hesperidin is the major polyphenol of sweet oranges, accounting for over 77% of the flavonol content [98–100]. These compounds are present in the edible fruit, juice, and/or peel, and here we concentrate on the juice and the edible fresh fruit, due to their direct implications in human health. As a functional food, Letaief et al. [97] determined that the AA content in orange juice fluctuates from 551 to 614 mg L⁻¹, total phenolics range from 413 to 417 mg GAE L⁻¹, and flavonoids from 25 to 60 mg catechin equivalents (CE) g⁻¹ DW. Roussos [94] measured total phenols (964–1215 mg TAE L⁻¹). The percentage of antioxidant activity of the juice, evaluated by the DPPH method, fluctuated from 36.4% to 56.6% [94, 97]. The antioxidant activity of sweet orange juice is dependent on the state of maturity of the fruit. Indeed, Adu et al. [101] noted higher levels of antioxidant activity in fruits of 3–6 months (over 80%) than in fruits of 10–12 months (around 70%). Fiber and amino acids are also important in juice. In this sense, Aschoff et al. [96] informed 1.4 g 100 g⁻¹ of dietary fiber, and Roussos [94] mentioned that juice contains 18 amino acids (included the essentials amino acids), especially proline, arginine, asparagine, glycine, serine, and γ -aminobutyric acid. Other authors have determined some of these and other phytochemicals in homogenate orange segments without peel, where Aschoff et al. [96] reported 36.2 mg 100 g⁻¹ FW of AA, 271.5 μ g 100 g⁻¹ FW of carotenoids, and 13.6 g 100 g⁻¹ FW of dietary fiber. Recently, Molan et al. [102] evaluated some compounds and properties of sweet orange seeds, such as total polyphenols (10.9–39.4 mg GAE g⁻¹ DW) and antioxidant activity (around 50%) by the DPPH method.

It has been reported that to maintain sufficient antioxidant protection, an estimated average consumption of 60 and 75 mg d⁻¹ of vitamin C is required for young women and men, respectively; however, it is suggested an increase of 35 mg d⁻¹ for smokers [103]. This is important because orange consumption provides other phytochemicals with multiple benefits to human health. Several clinical studies confirm this assertion. For example, sweet orange juice also harbors antidiabetic activity, as determined in rats by metabolome analysis [104]. This agrees with research performed by Kumar and Bhaskar [105] in rats, using ethanolic orange peel extract, where blood glucose decreased around 60% with respect to the control after 3 weeks of treatment, similar to the drug, glibenclamide. Furthermore, Mallick and Khan [89] suggest a combination of juice of *C. sinensis* and *C. paradisi* in order to reduce the level of glucose and improve the insulin level in the plasma of diabetic rats.

Hypocholesterolemic activity was demonstrated in women with aerobic exercise and a consumption of 500 mL of sweet orange juice daily [106]. These authors found a 15% decrease of low-density lipoprotein (LDL-C) in serum and an 18% increase of high-density lipoprotein (HDL-L), whereas the ratio LDL/HDL-cholesterol decreased by 27%. Furthermore, they also noted an improved performance during physical activity, by a reduction of blood lactate. Moreover, a long-term study (twelve months) showed that consumption of orange juice (480 mL daily) triggered reductions of 11% in total cholesterol, 18% in LDL-cholesterol, 12% in apolipoprotein B, and 12% in the LDL/HDL ratio in comparison to nonconsumers [107]. In addition, an increase in antiatherogenic activity levels with the consumption of sweet orange juice was found [108–110]. Recently, it was informed that in rats, antihyperlipidemic activity is due to phytochemical compounds like flavonoids and other polyphenols with antioxidant capacity present in the juice of sweet oranges [111]. Therefore, the juice of sweet oranges may play an important cardioprotective role by preventing thrombosis [111]. In humans, orange

juice intake also decreases procoagulant activity, possibly due to flavonoids, like anthocyanins, or other juice components [112].

Another feature of sweet orange juice that supports its cardioprotective role is its effect on diastolic blood pressure, which was significantly lower in men after the daily consumption of 500 mL orange juice for 4 weeks, and an enhancement of endothelium-dependent microvascular reactivity [113]. These authors also suggest that hesperidin could be related to the beneficial effect of orange juice in cardioprotection. Likewise, Rangel-Huerta et al. [114] related the reduction of blood pressure in obese adults with the consumption of at least 300 mg flavanones over 12 weeks. On the contrary, Schär et al. [115] found a relatively high flavanone and phenolic metabolite content in plasma, but no effects were observed on blood pressure and cardiovascular risk biomarkers. Additionally, Giordano et al. [116] reported that a daily intake of 1 L of orange juice for 4 weeks was not effective in reducing cellular markers associated with cardiovascular risks. Nevertheless, in general, more evidence of positive rather than neutral or negative effects on cardiovascular risk of sweet orange juice consumption exists. In fact, risk factors are mainly associated with metabolic syndromes such as cholesterol, blood pressure, and blood coagulation, and frequent intake of orange juice may be a useful delaying strategy [117].

The antiinflammatory activity of sweet orange juice has been reported by Mohanty et al. [118] where glucose induced an acute increase in ROS and inflammation, and orange juice intake prevented meal-induced oxidative and inflammatory stress [119]. Recent studies in rats revealed the positive effect of orange juice over histological and biochemical changes related with a progress in colonic oxidative status [120]. Besides, the antimicrobial activity of sweet orange juice has been reported by several authors. Recently, Adu et al. [101] indicated an inhibitory effect of orange juice from fruits at different stages of development against Gram-positive and Gram-negative bacteria and fungi, like *B. subtilis* NCTC 10073, *C. albicans* ATCC 10231, *E. coli* ATCC 25922, *P. vulgaris* NCTC 4175, *Pseudomonas aeruginosa* ATCC 27853, and *S. aureus* ATCC. Similar results on bacteria and fungi were found by Javed et al. [121] using essential oils of orange peel. This positive effect appears to be related with flavones, flavonoids, and flavonols [122].

3.3. Highbush blueberry (*Vaccinium corymbosum* L.)

The highbush blueberry is a species that belongs to the *Ericaceae* family (**Figure 7**) [123] exhibiting a high level of morphological diversity [124]. It is native to eastern United States and was domesticated during the twentieth century [125, 126]. Its distribution and consumption is extensive due to the human health benefits (antioxidant and mineral characteristics) of fruits and leaves [123] (see **Table 2**). This fruit has a wide range of phenolic compounds, especially flavonols, such as quercetin, as well as anthocyanins [127]. Some values of the main phytochemicals that contribute to antioxidant capacity are: phenolic compounds (261–585 mg g⁻¹ FW), flavonoids (50 mg g⁻¹ FW), and anthocyanins (25–495 mg g⁻¹ FW) [128–130].

The antioxidant activity is higher in wild blueberry species, and part of this activity is conserved in cultivated varieties [131]. The total antioxidant activity of blueberry species ranges from 15.88 to 18.41 μmol Fe²⁺ kg⁻¹ FW, using the FRAP reagent [130]. Contreras et al. [132] showed values near to 80% of antioxidant capacity measured by the DPPH method under *in vitro* conditions. The same authors affirmed that antioxidant capacity is related with the content of chlorogenic acid, myricetin, syringic acid, and rutin.

Figure 7. Fruits of *Vaccinium corymbosum* commonly name highbush blueberry.

Plasma antioxidant capacity (PAC) is considered a biomarker for antioxidant status of humans. In this context, Fernández-Panchon et al. [129] indicated that PAC increased following consumption of some foods rich in phenols, which could be related with *in vivo* bioactivity, and its consequent positive effects for human health. More specific biological properties of blueberry have been described, such as anticarcinogenic, antidiabetic, antiinflammatory, antimicrobial, and reducing cholesterol, among other activities [133–135]. The hypoglycemic activity of blueberry is mentioned by several authors. Aktan et al. [135] reported a severe case of hypoglycemia in a patient of 75 years old, who had diagnosed but untreated prediabetes. Just before the episode, this patient consumed about 500 mL juice of blueberry and *Laurocerasus officinalis*, which is also considered hypoglycemic. Similarly, Cheplick et al. [136] affirmed from *in vitro* studies that blueberry fruit has potential for diet-based management of hyperglycemia, especially in the early stages of disease.

Blueberry extracts also have antimicrobial activity, which have interest considering that many microorganisms are pathogenic to humans. In this line, a significant effect of extracts on *Listeria monocytogenes* and *Salmonella enteritidis* was found under laboratory conditions by Shen et al. [137]. In the same conditions (laboratory), other extracts from dried fruits and leaves were tested on contaminant/pathogenic microorganisms. The findings indicate good results in the inhibition of development of *S. aureus* ATCC 29213, *Enterococcus faecalis* ATCC 29212, *E. coli* ATCC 27853, *K. pneumoniae* ATCC10031, *Acetobacter baumannii* ATCC 19609, *S. enteritidis* ATCC 3076, *Salmonella typhimurium* KCCM 11862, *Enterococcus faecium* LGM 11423, *Listeria innocua* NCTC 11286, *Bacillus cereu* ATCC 11778, and *P. aeruginosa* ATCC 27853 [134, 138, 139].

Zhong et al. [140] tested homogenized fresh blueberry juice as a therapy of juvenile idiopathic arthritis. The combined therapy of blueberry juice and etanercept (the typical drug used to treat this condition), improved the therapeutic effect of etanercept in patients with this pathology. Samad et al. [141] confirmed the antiinflammatory activity of extracts of blueberry in an *in vitro* study.

Yi et al. [133] studied the effect of phenolic compounds over colon cancer cell proliferation. Results indicated that these phytochemicals could inhibit the carcinogenic cells. Massarotto

et al. [142] demonstrated that anthocyanins and other phenolic compounds have cytotoxic activity, as tested in tumoral cell lines under *in vitro* conditions; thus, blueberry extracts could be useful for future treatment of cancer, as a natural cytotoxic agent. In this context, Tsuda et al. [143] obtained similar results with human leukemia cells and ethanolic extracts of several berries that include blueberry fruits. In all cases, the induction of apoptosis in the cancerous cells may be the mechanism triggered by blueberry.

The antilipidemic and antiatherogenic actions of blueberry have been reported by several authors. Coban et al. [144] indicated that the fresh fruit is food supplements that generate a positive effect over aorta and liver of hypercholesterolemic Guinea pigs [144]. In this respect, Cutler et al. [145] confirmed that berries are a special source of phytochemicals (anthocyanins and other phenolic compounds) and can be exploited as natural phytochemicals to contribute toward the amelioration of several chronic diseases, including those derived from alterations in the lipid profile in vascular systems.

3.4. Maracuyá (*Passiflora edulis* Sims)

Maracuyá (*Passiflora edulis* Sims) is a species that belongs to the *Passifloraceae* family (**Figure 8**) [146, 147] which is native to Brazil, South America. Nevertheless, some authors report that its real origin is Australia and is called *Passiflora edulis* forma *flavicarpa* [148, 149]. Variability studies have been carried out in South America, mainly in Colombia, as this region is particularly rich in this genus, although a low variability has been reported [150, 151]. Both scientific names *Passiflora edulis* Sims and *Passiflora edulis* forma are indistinctly considered in this chapter. Talcott et al. [152] identified several phenolic acids such as galacturonic acid, p-hydroxybenzoic acid, syringic acid, caffeic acid, p-coumaric acid, tryptophan, flavonoid glycoside, sinapic acid, ferulic acid, o-coumaric acid, and syringic acid in this species. Some compounds such as tryptophan, sinapic acid, and p-coumaric acid are in higher quantity with 733, 626, and 623 $\mu\text{g L}^{-1}$ DW, respectively. Within the latter, total phenolic compounds fluctuated from 342.8 to 382 mg GAE L^{-1} FW [153]; total carotenoids varied between 22.4 and 29.1 mg L^{-1} DW, and the ascorbic acid content from 0.22 to 0.33 g kg^{-1} FW [152, 153]. The anthocyanin concentration in pulps and by product, on the other hand, were 3.48 and 3.7 mg 100 g^{-1} DW, respectively, while the flavonoids in pulps and by product were 60.3 and 40 mg 100 g^{-1} DW, respectively [154]. Regarding the above, Zucolotto et al. [155] indicated that C-glycosyl flavonoids are present in several species of *Passiflora* in South America. Furthermore, Da Silva et al. [154] informed that values for β -carotene fluctuated from 57.93 to 1362.07 $\mu\text{g } 100 \text{ g}^{-1}$ DW for both pulp and by product. It is worth noting, that piceatannol, a compound with an important antioxidant characteristic, is present in peel and seeds of the maracuyá fruit [156]. Moreover, the total antioxidant activity was found to reach values from 409.13 to 805.5 $\mu\text{M TE L}^{-1}$ FW in this fruit [153], while Marcoris et al. [157] indicate values ranging from 1279 to 1460 $\mu\text{M TE L}^{-1}$ FW. Total dietary and soluble fiber is another important characteristic attributed to this species, with values fluctuating from 35.5 to 81.5 g 100 g^{-1} DM. These values are higher in comparison to other tropical fruits such as *Mangifera indica*, *Ananas comosus*, and *Psidium guajava* [158]. The maracuyá fruit has several special characteristics that are beneficial for human health, described in **Table 2**. Within the latter, the most important properties are the sedative

and anxiolytic activities, which are common to several other species of the *Passiflora* genus [159]. Evaluation of aqueous extracts of pericarp fruit on rats concluded that a sedative effect was obtained with an oral administration of 300 mg kg^{-1} [160]. This effect was corroborated by a dose-dependent decrease of the locomotor-activity. Similar studies using ethanolic extracts of maracuyá leaves in rats exhibited sedative effects at 400 mg kg^{-1} [161]. Figueiredo et al. [162] reported that 130 mg kg^{-1} of bark flour of maracuyá fruits showed a sedative effect in rats. On the other hand, antiproliferative properties on cancer cells evaluated in SW480 and SW620 cells lines showed that cell growth in both lines was inhibited with $50\text{--}500 \mu\text{g mL}^{-1}$ of leaf ethanolic extracts and maracuyá fruit juice [163]. The polysaccharide peel of maracuyá fruits was also evaluated for its antiinflammatory effects and antidiabetic properties. In this context, the reduction of the inflammation was associated with the liberation or synthesis of histamine and serotonin, in response to a polysaccharide fraction of the maracuyá fruit [164]. Moreover, flour peel of maracuyá fruits in diabetic rats showed a decline in glucose content in the blood [165], probably associated with the high fiber level in this fruit tissue, which could prevent absorption of carbohydrates [166, 167]. Furthermore, triglycerides levels significantly decreased with 25 mg kg^{-1} of flour peel, however, no changes in total cholesterol levels were observed [165]. Still, Barbalho et al. [167] concluded that maracuyá fruit juice could improve the lipid profile, including the triglycerides, cholesterol, LDL-cholesterol, and HDL-cholesterol levels.

Figure 8. Fruits of *Passiflora edulis* commonly name maracuyá.

4. Conclusion and perspectives

A wealth of information in the field of phytochemical compounds and their impact on human health has been generated. Nowadays, it is possible to affirm that fruits and vegetables must be a part of daily diet. This is not simply a recommendation, but must be treated as an urgent

requirement to ameliorate human health, especially in decreasing chronic nontransmissible diseases. We believe that additional efforts of governments and diverse organisms related with human health are necessary in order to highlight the benefits of these food types. Coriander and kiwifruit have remarkable characteristics and are excellent functional foods. We highlight these species for their wide range of benefits in different human diseases and their worldwide distribution. Likewise, further investigation is required to understand the mechanisms associated with several biochemical and physiological processes induced by fruit and vegetable intake in humans. Furthermore, we consider that leeks and artichokes have special potential as functional foods. Although there is a lot of information about the beneficial effects of fruits, we believe it is possible to extend studies to other organs like leaves and stems in artichokes, and roots in leeks, because they can offer additional benefits to human health.

Acknowledgments

We are grateful to FONDECYT 1110726, 1120917, and 1150220 projects and FONDECYT-POSDOCTORAL 3120248, Doctoral Fellowship CONICYT-PCHA/Doctorado Nacional/2016-21160984, as well as the Millennium Nucleus for Plant Synthetic Biology and Systems Biology NC130030. We would like to thank Dr. Michael Handford and Dr. Franko Restovic for language support.

Author details

Ricardo Tighe-Neira¹, Miren Alberdi^{2,3}, Patricio Arce-Johnson⁴, Jesús L. Romero-Romero^{4,5,6}, Marjorie Reyes-Díaz^{2,3} and Claudio Inostroza-Blancheteau^{1,7*}

*Address all correspondence to: claudio.inostroza@uct.cl

1 School of Agronomy, Faculty of Natural Resources, Universidad Católica de Temuco, Temuco, Chile

2 Department of Chemical Sciences and Natural Resources, Faculty of Engineering and Sciences, Universidad de La Frontera, Temuco, Chile

3 Center of Plant, Soil Interaction and Natural Resources Biotechnology, Scientific and Technological Bioresource Nucleus (BIOREN), Universidad de La Frontera, Temuco, Chile

4 Department of Molecular Genetics and Microbiology, Faculty of Biological Sciences, Pontificia Universidad Católica de Chile, Santiago, Chile

5 Faculty of Agronomy and Forest Engineering, Pontificia Universidad Católica de Chile, Santiago, Chile

6 Department of Agricultural Biotechnology, Instituto Politécnico Nacional, CIIDIR Sinaloa Unid, Guasave, Mexico

7 Nucleus of Research in Food Production, Faculty of Natural Resources, Universidad Católica de Temuco, Temuco, Chile

References

- [1] Hasler CM. Functional foods: their role in disease prevention and health promotion. *Food Technology*. 1998; 52: 63–147.
- [2] Institute of Medicine. Opportunities in the Nutrition and Food Sciences: Research Challenges and the Next Generation of Investigators. Washington, DC: The National Academies Press, 1994. p. 328. DOI:10.17226/2133
- [3] Steinmetz KA, Potter JD. Vegetables, fruit, and cancer. II. Mechanisms. *Cancer Causes & Control*. 1991; 2: 427–442. DOI:10.1007/BF00054304
- [4] Palermo M, Pellegrini N, Fogliano V. The effect of cooking on the phytochemical content of vegetables: Effect of cooking on vegetable phytochemicals. *Journal of the Science of Food and Agriculture*. 2014; 94: 1057–1070. DOI:10.1002/jsfa.6478
- [5] Slavin JL, Lloyd B. Health benefits of fruits and vegetables. *Advances in Nutrition*. 2012; 3: 506–516. DOI:10.3945/an.112.002154
- [6] Uchida A, Ogihara J, Kumagai H, Akao M, Matsufuji H, Sakurai H, Sekiguchi I. Mechanisms of sulfide components expression and structural determination of substrate precursor in jumbo leek (*Allium ampeloprasum* L.). *Nippon Shokuhin Kagaku Kogaku Kaishi*. 2009; 56: 280–285. DOI:10.3136/nskkk.56.280
- [7] Ceccarelli N, Curadi M, Picciarelli P, Martelloni L, Sbrana C, Giovannetti M. Globe artichoke as a functional food. *Mediterranean Journal of Nutrition and Metabolism*. 2010; 3: 197–201. DOI:10.1007/s12349-010-0021-z.
- [8] Nambiar VS, Daniel M, Guin P. Characterization of polyphenols from coriander leaves (*Coriandrum sativum*), Red amaranthus (*A. paniculatus*) and green amaranthus (*A. frumentaceus*) using paper chromatography and their health implications. *Journal of Herbal Medicine and Toxicology*. 2010; 4: 173–177.
- [9] Simonovska J, Rafajlovska V, Kavrakovski Z, Srbinoska M. Nutritional and bioactive compounds in hot fruits of *Capsicum annuum* L. from Macedonia. *Macedonian Journal of Chemistry and Chemical Engineering*. 2014; 33: 95–102.
- [10] Bratsch A, Arancibia A. Specialty Crop Profile: Globe Artichoke. College of Agriculture and Life Sciences, Virginia Polytechnic Institute and State University. 2014; Publication 438–108, p. 2–9.
- [11] Bianco VV. Present situation and future potential of artichoke in the mediterranean basin. *Acta Horticulturae*. 2005; 681: 39–58. DOI:10.17660/ActaHortic.2005.681.2
- [12] Pennington JAT, Fisher RA. Classification of fruits and vegetables. *Journal of Food Composition and Analysis*. 2009; 22: S23–S31. DOI:10.1016/j.jfca.2008.11.012
- [13] Alamanni MC, Cossu M. Antioxidant activity of the extracts of the edible part of artichoke (*Cynaras colymus* L.) var. spinososardo. *Italian Journal of Food Science*. 2003; 15: 187–195.

- [14] Romani A, Pinelli P, Cantini C, Cimato A, Heimler D. Characterization of Violetto di Toscana, a typical Italian variety of artichoke (*Cynaras colymus* L.). Food Chemistry. 2006; 95: 221–225. DOI:10.1016/j.foodchem.2005.01.013
- [15] Karami R, Hosseini M, Mohammadpour T, Ghorbani A, Sadeghnia HR, Rakhshandeh H, Vafae F, Esmailizadeh M. Effects of hydroalcoholic extract of *Coriandrum sativum* on oxidative damage in pentylenetetrazole-induced seizures in rats. Iranian Journal of Neurology. 2015; 14: 59–66.
- [16] Ruiz-Aceituno L, García-Sarrió M J, Alonso-Rodriguez B, Ramos L, Sanz L. Extraction of bioactive carbohydrates from artichoke (*Cynara scolymus* L.) external bracts using microwave assisted extraction and pressurized liquid extraction. Food Chemistry. 2016; 196: 1156–1162.
- [17] Hernández-Hernández O, Ruiz-Aceituno L, Sanz M, Martínez-Castro I. Determination of free inositols and other low molecular weight carbohydrates in vegetables. Journal of Agricultural and Food Chemistry. 2011; 6: 2451–2455.
- [18] Ostlund RE, Sherman WR. Pinitol and derivatives thereof for the treatment of metabolic disorders (US Patent 5827896). 1998.
- [19] Crawford TJ, Crowther CA, Alsweiler J, Brown J. Antenatal dietary supplementation with myo-inositol in women during pregnancy for preventing gestational diabetes. The Cochrane Library. 2015.
- [20] Roberfroid, M. Dietary fiber, inulin, and oligofructose: a review comparing their physiological effects. Critical Reviews in Food Science & Nutrition. 1993; 33, 103–148.
- [21] Shoaib M, Shehzad A, Omar M, Rakha A, Raza H, Sharif HR, Shakeel A, Ansari A, Niazi S. Inulin: Properties, health benefits and food applications. Carbohydrate Polymers. 2016; 147: 444–454.
- [22] Bosscher D, Van Loo J, Franck A. Inulin and oligofructose in lowering the risk of osteoporosis: a major concern for women's health. Agro Food Industry HI-TECH. 2006; 1: 10–12.
- [23] Jones HA, Mann LK. Onions and their allies: botany, cultivation, and utilization. Hill L, editor. Cornell University, 1963. p. 286.
- [24] Fritsch RM, Friesen N. Evolution, Domestication and Taxonomy, Chapter 1. In: Rabinowitch HD, Currah L, editors. Allium Crop Science: Recent Advances. Wallingford, UK, CABI Publishing; 2002. p. 5–30. DOI:10.1079/9780851995106.0005
- [25] Hirschegger P, Galmarini C, Bohanec B. Characterization of a novel form of fertile great headed garlic (*Allium* sp.). Plant Breeding. 2006; 125: 635–637. DOI:10.1111/j.1439–0523.2006.01279.x
- [26] Mathew BA. Review of Allium Section Allium. Royal Botanic Gardens. Kew, editors. Chicago USA. Kew Publishing. 1996. p. 182.

- [27] Hirschegger P, Jakše J, Trontelj P, Bohanec B. Origins of *Allium ampeloprasum* horticultural groups and a molecular phylogeny of the section *Allium* (*Allium*: Alliaceae). *Molecular Phylogenetics and Evolution*. 2010; 54: 488–497. DOI:10.1016/j.ympev.2009.08.030
- [28] Bernaert N, De Paepe D, Bouten C, De Clercq H, Stewart D, Van Bockstaele E, Van Droogenbroeck B. Antioxidant capacity, total phenolic and ascorbate content as a function of the genetic diversity of leek (*Allium ampeloprasum* var. *porrum*). *Food Chemistry*. 2012; 134: 669–677. DOI:10.1016/j.foodchem.2012.02.159
- [29] Bernaert N, De Clercq H, Van Bockstaele E, De Loose M, Van Droogenbroeck B. Antioxidant changes during postharvest processing and storage of leek (*Allium ampeloprasum* var. *porrum*). *Postharvest Biology and Technology*. 2013; 86: 8–16.
- [30] Vandekinderen I, Van Camp J, Devlieghere F, Ragaert P, Veramme K, Bernaert N, Denon Q, De Meulenaer B. Evaluation of the use of decontamination agents during fresh-cut leek processing and quantification of their effect on its total quality by means of a multidisciplinary approach. *Innovative Food Science & Emerging Technologies*. 2009; 10: 363–373. DOI:10.1016/j.ifset.2009.02.002
- [31] Adao CR, da Silva BP, Parente JP. A new steroidal saponin from *Allium ampeloprasum* var. *porrum* with antiinflammatory and gastroprotective effects. *Phytochemistry Letters*. 2011a; 4: 306–310. DOI:10.1016/j.phytol.2011.05.009
- [32] Adao CR, da Silva BP, Parente JP. A new steroidal saponin with antiinflammatory and antiulcerogenic properties from the bulbs of *Allium ampeloprasum* var. *porrum*. *Fitoterapia*. 2011b; 82: 1175–1180. DOI:10.1016/j.fitote.2011.08.003
- [33] Sobolewska D, Michalska K, Podolak I, Grabowska K. Steroidal saponins from the genus *Allium*. *Phytochemistry Reviews* 2014; 15: 1–35. DOI:10.1007/s11101-014-9381-1
- [34] Maidment DCJ, Dembny Z, Watts DI. The anti-bacterial activity of 12 *Alliums* against *Escherichia coli*. *Nutrition & Food Science*. 2001; 31: 238–241. DOI:10.1108/EUM000000000005614
- [35] Radovanović B, Mladenović J, Radovanović A, Pavlović R, Nikolić V. Phenolic composition, antioxidant, antimicrobial and cytotoxic activities of *Allium porrum* L. (Serbia) extracts. *Journal of Food and Nutrition Research*. 2015; 3: 564–569. DOI:10.12691/jfnr-3-9-1
- [36] Hervet-Hernández D, Sáyo-Ayerdi SG, Goñi I. Bioactive compounds of four hot pepper varieties (*Capsicum annuum* L.), antioxidant capacity, and intestinal bioaccessibility. *Journal of Agricultural and Food Chemistry*. 2010; 58: 3399–3406. DOI:10.1021/jf904220w
- [37] Tighe-Neira R, Navarro RM, Cantergiani GL, Novoa AC. Efecto de dos extractos botánicos en el desarrollo y contenido de polifenoles de ají (*Capsicum annuum* L.). *Revista Mexicana de Ciencias Agrícolas*. 2013; 5: 115–127

- [38] Serrano J; Goñi I, Saura-Calixto F. Determination of β -carotene and lutein available from green leafy vegetables by an in vitro digestion and colonic fermentation method. J. Agric. Food Chem. 2005; 53: 2936–2940. DOI:10.1021/jf0480142
- [39] Golubkina NA, Dzhos EA, Pishnaya ON, Mamedov MI, Nadezhkin SM. Biochemical characteristics of hot pepper fruit grown in Moscow region. Russian Agricultural Sciences. 2014; 40: 27–31. DOI:10.3103/S1068367414010042
- [40] De M, De Ak, Banerjee AB. Screening of spices for antimicrobial activity. Journal of Spices and Aromatic Crops. 1999; 2.
- [41] Soetarno S, Sukrasno S, Yulinah E, Sylvia S. Antimicrobial activities of the ethanol extracts of capsicum fruits with different levels of pungency. Jurnal Matematika & Sains. 1997; 2: 57–63.
- [42] Huang X, Huadan TAN, Zheng Y, Zhang J, Yijun ZHOU. Study on antimicrobial activities of ethanol extracts from different capsicum cultivars. 农业科学与技术 (英文版). 2012; 13: 2514–2517.
- [43] Mhemdi H, Rodier E, Kechaou N, Fages J. A supercritical tuneable process for the selective extraction of fats and essential oil from coriander seeds. Journal of Food Engineering. 2011; 105: 609–616. DOI:10.1016/j.jfoodeng.2011.03.030
- [44] Sahib NG, Anwar F, Gilani AH, Hamid AA, Saari N, Alkharfy KM. Coriander (*Coriandrum sativum* L.): A potential source of high-value components for functional foods and nutraceuticals - A review. Phytotherapy Research 2012; 27: 1439–1456. DOI:10.1002/ptr.4897
- [45] Bhuiyan MNI, Begum J, Sultana M. Chemical composition of leaf and seed essential oil of *Coriandrum sativum* L. from Bangladesh. Bangladesh Journal of Pharmacology. 2009; 4: 150–153. DOI:10.3329/bjp.v4i2.2800
- [46] Zoubiri S, Baaliouamer A. Essential oil composition of *Coriandrum sativum* seed cultivated in Algeria as food grains protectant. Food Chemistry. 2010; 122: 1226–1228. DOI:10.1016/j.foodchem.2010.03.119
- [47] Buthlezi MND, Soundy P, Jifon J, Sivakumar D. Spectral quality of photo-selective nets improves phytochemicals and aroma volatiles in coriander leaves (*Coriandrum sativum* L.) after postharvest storage. Journal of Photochemistry and Photobiology B: Biology. 2016; 161: 328–34. DOI:10.1016/j.jphotobiol.2016.05.032
- [48] Kumar RS, Balasubramanian P, Govindaraj P, Krishnaveni T. Preliminary studies on phytochemicals and antimicrobial activity of solvent extracts of *Coriandrum sativum* L. roots (Coriander). Journal of Pharmacognosy and Phytochemistry. 2014; 2: 74–78.
- [49] Deepa B, Anuradha CV. Antioxidant potential of *Coriandrum sativum* L. Indian Journal Experimental Biology. 2011; 49: 30–8.

- [50] Dhanapakiam P, Joseph JM, Ramaswamy VK, Moorthi M, Kumar AS. The cholesterol lowering property of coriander seeds (*Coriandrum sativum*): mechanism of action. *Journal of Environmental Biology*. 2008; 29: 53–56.
- [51] Saravanan M, Ignacimuthu S. Hypocholesterolemic effect of indian medicinal plants - A review. *Medicinal Chemistry*. 2015; 5: 040–049. DOI:10.4172/2161-0444.10002
- [52] Kharade SM, Gumate DS, Patil VM, Kokane SP, Naikwade NS. Behavioral and biochemical studies of seeds of *Coriandrum sativum* in various stress models of depression. *International Journal of Current Research and Review*. 2011; 1003: 4–8.
- [53] Zargar-Nattaj SS, Tayyebi P, Zangoori V, Moghadamnia Y, Roodgari H, Jorsaraei SG, Moghadamnia AA. The effect of *Coriandrum sativum* seed extract on the learning of new born mice by electric shock: interaction with caffeine and diazepam. *Psychology Research and Behavior Management*. 2011; 4: 13–19. DOI:10.2147/PRBM.S15905
- [54] Mahendra P, Bisht S. Anti-anxiety activity of *Coriandrum sativum* assessed using different experimental anxiety models. *Indian Journal of Pharmacology*. 2011; 43: 574–577. DOI:10.4103/0253-7613.84975
- [55] Fatemeh KS, Vafadar LS, Hosseini M, Naser SM, Ghorbani A, Pourganji M. The analgesic effects of different extracts of aerial parts of *Coriandrum Sativum* in mice. *International Journal of Biomedical Science*. 2015; 11: 23–28.
- [56] Latha K, Rammohan B, Sunanda BPV, Maheswari MSU, Mohan SK. Evaluation of anxiolytic activity of aqueous extract of *Coriandrum sativum* Linn. in mice: A preliminary experimental study. *Pharmacognosy Research*. 2015; 7: (Suppl 1): S47–S51. DOI:10.4103/0974-8490.157996
- [57] Soquetta MB, Stefanello FS, Da Mota Huerta K, Monteiro SS, Da Rosa CS, Terra NN. Characterization of physiochemical and microbiological properties, and bioactive compounds, of flour made from the skin and bagasse of kiwi fruit (*Actinidia deliciosa*). *Food chemistry*. 2016. 199: 471–478. DOI:10.1016/j.foodchem.2015.12.022
- [58] Liang Chou-Fen, Ferguson AR. The botanical nomenclature of the kiwifruit and related taxa. *New Zealand Journal of Botany*. 1986; 24: 183–184. DOI:10.1080/0028825X.1986.10409728
- [59] Drzewiecki J, Latocha P, Leontowicz H, Leontowicz M, Park YS, Najman K, Weisz M, Ezra A, Gorinstein S. Analytical methods applied to characterization of *Actinidia arguta*, *Actinidia deliciosa*, and *Actinidia eriantha* kiwi fruit cultivars. *Food Analytical Methods*. 2016; 9: 1353–1366. DOI:10.1007/s12161-015-0309-1
- [60] D'Evoli L, Moscatello S, Lucarini M, Aguzzi A, Gabrielli P, Proietti S, Battistelli A, Famianic F, Böhm V, Lombardi-Boccia G. Nutritional traits and antioxidant capacity of kiwifruit (*Actinidia deliciosa* Planch., cv. Hayward) grown in Italy. *Journal of Food Composition and Analysis*. 2015; 37: 25–29. DOI:10.1016/j.jfca.2014.06.012
- [61] Gammon CS, Kruger R, Minihaue AM, Conlon CA, von Hurst PR, Stonehouse W. Kiwifruit consumption favourably affects plasma lipids in a randomised controlled

- trial in hypercholesterolaemic men. *British Journal of Nutrition*. 2013; 109: 2208–2218. DOI:10.1017/S0007114512004400
- [62] Hunter DC, Skinner MA, Wolber FM, Booth CL, Loh JM, Wohlers M, Stevenson LM, Kruger MC. Consumption of gold kiwifruit reduces severity and duration of selected upper respiratory tract infection symptoms and increases plasma vitamin C concentration in healthy older adults. *British Journal of Nutrition*. 2012; 108: 1235–1245. DOI:10.1017/S0007114511006659
- [63] Lee I, Im S, Jin CR, Heo HJ, Cho YS, Baik MY, Kim DO. Effect of maturity stage at harvest on antioxidant capacity and total phenolics in kiwifruits (*Actinidia* spp.) grown in Korea. *Horticulture, Environment, and Biotechnology*. 2015; 56: 841–848. DOI:10.1007/s13580-015-1085-y.
- [64] Carr AC, Bozonet SM, Vissers MCM. A randomised cross-over pharmacokinetic bioavailability study of synthetic versus kiwifruit-derived vitamin C. *Nutrients*. 2013a; 5: 4451–4461. DOI:10.3390/nu5114451
- [65] Carr AC, Bozonet SM, Pullar JM, Simcock JW, Vissers M. A randomized steady-state bioavailability study of synthetic versus natural (Kiwifruit-Derived) vitamin C. *Nutrients*. 2013b; 5: 3684–3695. DOI:10.3390/nu5093684
- [66] Vissers MC, Bozonet SM, Pearson JF, Braithwaite LJ. Dietary ascorbate intake affects steady state tissue concentrations in vitamin C-deficient mice: tissue deficiency after suboptimal intake and superior bioavailability from a food source (kiwifruit). *The American journal of clinical nutrition*. 2011; 93: 292–301. DOI:10.3945/ajcn.110.004853
- [67] Beck K, Conlon C, Kruger R, Coad J, Stonehouse W. The effect of gold kiwifruit consumed with an iron fortified breakfast cereal meal on iron status in women with low iron stores: A 16 week randomised controlled intervention study. *BMC Public Health*. 2010; 10: 1. DOI:10.1186/1471-2458-10-36
- [68] Beck K, Conlon CA, Kruger R, Coad J, Stonehouse W. Gold kiwifruit consumed with an iron-fortified breakfast cereal meal improves iron status in women with low iron stores: a 16-week randomised controlled trial. *British Journal of Nutrition*. 2011; 105: 101–109. DOI:10.1017/S0007114510003144
- [69] Schakel S, Sievert YA, Buzzard IM. In: *Dietary fiber values for common foods*. The CRC handbook of dietary fiber in human nutrition 3rd ed. Spiller G. Editor. London: CRC Press. 2001.
- [70] Chan AO, Leung G, Tong T, Wong NY. Increasing dietary fiber intake in terms of kiwifruit improves constipation in Chinese patients. *World journal of gastroenterology*. 2007; 13: 4771–4775.
- [71] Singletary K. Kiwifruit: overview of potential health benefits. *Nutrition Today*. 2012; 47: 133–147. DOI:10.1097/NT.0b013e31825744bc
- [72] Chang CC, Lin YT, Lu YT, Liu YS, Liu JF. Kiwifruit improves bowel function in patients with irritable bowel syndrome with constipation. *Asia Pacific Journal of Clinical Nutrition*. 2010; 19: 451–457. DOI: 10.6133/apjcn.2010.19.4.01.

- [73] Rutherford SM, Montoya CA, Zou ML, Moughan PJ, Drummond LN, Boland MJ. Effect of actinidin from kiwifruit (*Actinidiadeliciosa* cv. Hayward) on the digestion of food proteins determined in the growing rat. *Food Chemistry*. 2011; 129: 1681–1689. DOI:10.1016/j.foodchem.2011.06.031
- [74] Kaur L, Rutherford SM, Moughan PJ, Drummond L, Boland MJ. Actinidin enhances gastric protein digestion as assessed using an in vitro gastric digestion model. *Journal of agricultural and food chemistry*. 2010a; 58: 5068–5073. DOI:10.1021/jf903332a
- [75] Kaur L, Rutherford SM, Moughan PJ, Drummond L, Boland MJ. Actinidin enhances protein digestion in the small intestine as assessed using an in vitro digestion model. *Journal of Agricultural and Food Chemistry*. 2010b; 58: 5074–5080. DOI:10.1021/jf903835g
- [76] Gammon CS, Minihaue AM, Kruger R, Conlon CA, von Hurst PR, Jones B, Stonehouse W. TaqIB polymorphism in the cholesteryl ester transfer protein (CETP) gene influences lipid responses to the consumption of kiwifruit in hypercholesterolaemic men. *British Journal of Nutrition*. 2014a; 111: 1077–1084. DOI:10.1017/S0007114513003437
- [77] Gammon CS, Kruger R, Conlon CA, von Hurst PR, Jones B, Stonehouse W. Inflammatory status modulates plasma lipid and inflammatory marker responses to kiwifruit consumption in hypercholesterolaemic men. *Nutrition, Metabolism and Cardiovascular Diseases*. 2014b; 24: 91–99. DOI:10.1016/j.numecd.2013.06.001
- [78] Duttaroy AK, Jørgensen A. Effects of kiwi fruit consumption on platelet aggregation and plasma lipids in healthy human volunteers. *Platelets*. 2004; 15: 287–292. DOI:10.1080/09537100410001710290
- [79] Karlsen A, Svendsen M, Seljeflot I, Laake P, Duttaroy AK, Drevon CA, Arnesen H, Tonstad S, Blomhoff R. Kiwifruit decreases blood pressure and whole-blood platelet aggregation in male smokers. *Journal of human hypertension*. 2013; 27: 126–130. DOI:10.1038/jhh.2011.116
- [80] Svendsen M, Tonstad S, Heggen E, Pedersen TR, Seljeflot I, Bøhn SK, Bastani NE, Blomhoff R, Holme IM, Klemsdal TO. The effect of kiwifruit consumption on blood pressure in subjects with moderately elevated blood pressure: A randomized, controlled study. *Blood Pressure*. 2014; 24: 48–54. DOI:10.3109/08037051.2014.976979
- [81] Brevik A, Gaivão I, Medin T, Jørgensen A, Piasek A, Elilasson J, Karlsen A, Blomhoff R, Veggan T, Duttaroy AK, Collins AR. Supplementation of a western diet with golden kiwifruits (*Actinidia chinensis* var. 'Hort 16A') effects on biomarkers of oxidation damage and antioxidant protection. *Nutrition Journal*. 2011; 10: 54. DOI:10.1186/1475-2891-10-54
- [82] Hunter DC, Greenwood J, Zhang J, Skinner MA. Antioxidant and “natural protective” properties of kiwifruit. *Current Topics in Medicinal Chemistry*. 2011; 11: 1811–1820. DOI:10.2174/156802611796235134
- [83] Skinner MA. Wellness foods based on the health benefits of fruit: gold kiwifruit for immune support and reducing symptoms of colds and influenza. *J. Food Drug Anal.* 2012; 20 (Suppl 1): 261–264.

- [84] Calder PC, Kew S. The immune system: a target for functional British Journal of Nutrition. 2002; 88: S165-S176. DOI:10.1079/BJN2002682
- [85] Collins BH, Horská A, Hotten PM, Riddoch C, Collins AR. Kiwifruit protects against oxidative DNA damage in human cells and in vitro. Nutrition and Cancer. 2001; 39: 148–153. DOI:10.1207/S15327914nc39120
- [86] Fiorentino A, D'Abrosca B, Pacifico S, Mastellone C, Scognamiglio M, Monaco P. Identification and assessment of antioxidant capacity of phytochemicals from kiwi fruits. Journal of Agricultural and Food Chemistry. 2009; 57: 4148–4155. DOI:10.1021/jf900210z
- [87] Collins AR, Harrington V, Drew J, Melvin R. Nutritional modulation of DNA repair in a human intervention study. Carcinogenesis. 2003; 24: 511–515. DOI:10.1093/carcin/24.3.511
- [88] Moore GA. Oranges and lemons: clues to the taxonomy of Citrus from molecular markers. Trends in Genetics. 2001; 17: 536–540. DOI:10.1016/S0168-9525(01)02442-8
- [89] Mallick N, Khan RA. Effect of *Citrus paradisi* and *Citrus sinensis* on glycemic control in rats. African Journal of Pharmacy and Pharmacology. 2015; 9: 60–64. DOI:10.5897/AJPP2014.4185
- [90] Gmitter FG, Hu X. The possible role of Yunnan, China, in the origin of contemporary Citrus species (Rutaceae). Economic Botany. 1990; 44: 267–277. DOI:10.1007/BF02860491
- [91] Bayer RJ, Mabberley DJ, Morton C, Miller CH, Sharma IK, Pfeil B E, Rich S, Hitchcock R, Sykes S. A molecular phylogeny of the orange subfamily (Rutaceae: Aurantioideae) using nine cpDNA sequences. American Journal of Botany. 2009; 96: 668–685. DOI:10.3732/ajb.0800341
- [92] Xu Q, Chen LL, Ruan X, Chen D, Zhu A, Chen C, Bertrand D, Jiao Wen-Biao, Hao Bao-Hai, Lyon MP, Chen J, Gao S, Xing F, Lan H, Chang Ji-Wei, Ge X, Lei Y, Hu Q, Miao Y, Wang L, Xiao S, Biswas MK, Zeng W, Guo F, Cao H, Yang X, Xu Xi-Wen, Cheng Yun-Jiang, Xu J, Liu Ji-Hong, Luo OJ, Tang Z, Guo Wen-Wu, Kuang H, Zhang Hong-Yu, Roose ML, Nagarajan N, Deng Xiu-Xin, Ruan Y. The draft genome of sweet orange (*Citrus sinensis*). Nature genetics. 2013; 45: 59–66. DOI:10.1038/ng.2472
- [93] Grosso G, Galvano F, Mistretta A, Marventano S, Nolfo F, Calabrese G, Buscemi S, Drago F, Veronesi U, Scuderi A. Red orange: Experimental models and epidemiological evidence of its benefits on human health. Oxidative Medicine and Cellular Longevity. 2013; Article ID 157240: 11. DOI:10.1155/2013/157240
- [94] Roussos PA. Phytochemicals and antioxidant capacity of orange (*Citrus sinensis* (L.) Osbeck cv. Salustiana) juice produced under organic and integrated farming system in Greece. Scientia horticulturae. 2011; 129: 253–258. DOI:10.1016/j.scienta.2011.03.040
- [95] Goulas V, Manganaris GA. Exploring the phytochemical content and the antioxidant potential of Citrus fruits grown in Cyprus. Food Chemistry. 2012; 131: 39–47. DOI:10.1016/j.foodchem.2011.08.007

- [96] Aschoff JK, Kaufmann S, Kalkan O, Neidhart S, Carle R, Schweiggert RM. In vitro bioaccessibility of carotenoids, flavonoids, and vitamin C from differently processed oranges and orange juices [*Citrus sinensis* (L.) Osbeck]. *Journal of Agricultural and Food Chemistry*. 2015; 63: 578–587. DOI:10.1021/jf505297t
- [97] Letaief H, Zemni H, Mliki A, Chebil S. Composition of *Citrus sinensis* (L.) Osbeck cv «Maltaise demi-sanguine» juice. A comparison between organic and conventional farming. *Food Chemistry*. 2016; 194: 290–295. DOI:10.1016/j.foodchem.2015.08.025
- [98] Mattila P, Astola J, Kumpulainen J. Determination of flavonoids in plant material by HPLC with diode-array and electro-array detections. *Journal of Agricultural and Food Chemistry*. 2000; 48: 5834–5841. DOI:10.1021/jf000661f
- [99] Neveu V, Perez-Jimenez J, Vos F, Crespy V, du Chaffaut L, Mennen L., Knox C, Eisner R, Cruz J, Wishart D, Scalbert A. Phenol-Explorer: an online comprehensive database on polyphenol contents in foods. *Database*. 2010. DOI:10.1093/database/bap024
- [100] Perche O, Vergnaud-Gauduchon J, Morand C, Dubray C, Mazur A, Vasson, MP. Orange juice and its major polyphenol hesperidin consumption do not induce immunomodulation in healthy well-nourished humans. *Clinical Nutrition*. 2014; 33: 130–135. DOI:10.1016/j.clnu.2013.03.012
- [101] Adu F, Apenteng JA, Kuntworbe N, Akanwariwiak WG, Appiah T, Mintah DN. Antimicrobial and antioxidant activities of *Citrus sinensis* var. late Valencia fruits at various stages of development. *African Journal of Microbiology Research*. 2016; 10: 73–78. DOI:10.5897/AJMR2015.7725
- [102] Molan AL, Ismail MH, Nsaif RH. Phenolic contents and antioxidant activity of peels and seeds of orange (*Citrus sinensis*) cultivated in Iraq. 2016. DOI:10.20959/wjpps20164-6478
- [103] Fallico B, Ballistreri G, Arena E, Brighina S, Rapisarda P. Bioactive compounds in blood oranges (*Citrus sinensis* (L.) Osbeck): Level and intake. *Food Chemistry*. 2017; 215: 67–75. DOI:10.1016/j.foodchem.2016.07.142
- [104] Ahamad SR. Phytochemical, heavy metal and metabolomic anti-diabetic analysis of *Citrus sinensis*. *Bulletin of Environment, Pharmacology and Life Sciences*. 2015; 5: 22–27.
- [105] Kumar PRZA, Bhaskar A. Evaluation of antihyperglycaemic and antihyperlipidemic activity of *Citrus sinensis* peel extract on streptozotocin-induced diabetic rats. *International Journal of Diabetes in Developing Countries*. 2015; 35: 448–453. DOI:10.1007/s13410-015-0310-7
- [106] Aptekmann NP, Cesar TB. Orange juice improved lipid profile and blood lactate of overweight middle-aged women subjected to aerobic training. *Maturitas*. 2010; 67: 343–347. DOI:10.1016/j.maturitas.2010.07.009
- [107] Aptekmann NP, Cesar TB. Long-term orange juice consumption is associated with low LDL-cholesterol and apolipoprotein B in normal and moderately hypercholesterolemic subjects. *Lipids in health and disease*. 2013; 12: 1. DOI:10.1186/1476-511X-12-119

- [108] Choi JS, Yokozawa T, Oura H. Antihyperlipidemic effect of flavonoids from *Prunus davidiana*. *Journal of Natural Products*. 1991; 54: 218–224. DOI:10.1021/np50073a022
- [109] Cesar TB, Aptekmann NP, Araujo MP, Vinagre CC, Maranhão R C. Orange juice decreases low-density lipoprotein cholesterol in hypercholesterolemic subjects and improves lipid transfer to high-density lipoprotein in normal and hypercholesterolemic subjects. *Nutrition Research*. 2010; 30: 689–694. DOI:10.1016/j.nutres.2010.09.006
- [110] Kurowska EM, Spence JD, Jordan J, Wetmore S, Freeman DJ, Piché LA, Serratore P. HDL-cholesterol-raising effect of orange juice in subjects with hypercholesterolemia. *The American Journal of Clinical Nutrition*. 2000; 72: 1095–1100.
- [111] Mallick N, Khan RA. Antihyperlipidemic effects of *Citrus sinensis*, *Citrus paradisi*, and their combinations. *Journal of Pharmacy & Bioallied Sciences*. 2016; 8: 112.
- [112] Napoleone E, Cutrone A, Zurlo F, Di Castelnuovo A, D’Imperio M, Giordano L., De Curtis A, Iacoviello L, Rotilio D, Cerletti C, de Gaetano G, Donati MB, Lorenzet R. Both red and blond orange juice intake decreases the procoagulant activity of whole blood in healthy volunteers. *Thrombosis Research*. 2013; 132: 288–292. DOI:10.1016/j.thromres.2013.06.022
- [113] Morand C, Dubray C, Milenkovic D, Lioger D, Martin JF, Scalbert A, Mazur A. Hesperidin contributes to the vascular protective effects of orange juice: a randomized crossover study in healthy volunteers. *The American Journal of Clinical Nutrition*. 2011; 93: 73–80. DOI:10.3945/ajcn.110.0047945
- [114] Rangel-Huerta OD, Aguilera CM, Martin MV, Soto MJ, Rico MC, Vallejo F, Tomas-Barberan F, Perez-de-la-Cruz JA, Gil A, Mesa MD. Normal or high polyphenol concentration in orange juice affects antioxidant activity, blood pressure, and body weight in obese or overweight adults. *The Journal of Nutrition* 2015; 145: 1808–1816. DOI:10.3945/jn.115.213660
- [115] Schär MY, Curtis PJ, Hazim S, Ostertag LM, Kay CD, Potter J Cassidy A. Orange juice-derived flavanone and phenolic metabolites do not acutely affect cardiovascular risk biomarkers: a randomized, placebo-controlled, crossover trial in men at moderate risk of cardiovascular disease. *The American Journal of Clinical Nutrition*. 2015; 101: 931–938. DOI:10.3945/ajcn.114.104364
- [116] Giordano L, Coletta W, Tamburrelli C, D’Imperio M, Crescente M, Silvestri C, Rapisarda P, Reforgiato Recupero G, De Curtis A, Iacoviello L, de Gaetano G, Rotilio D, Cerletti C, Donati MB. Four-week ingestion of blood orange juice results in measurable anthocyanin urinary levels but does not affect cellular markers related to cardiovascular risk: a randomized cross-over study in healthy volunteers. *European Journal of Nutrition*. 2012; 51: 541–548. DOI:10.1007/s00394-011-0237-9
- [117] Silveira JQ, Dourado GK, Cesar TB. Red-fleshed sweet orange juice improves the risk factors for metabolic syndrome. *International Journal of Food Sciences and Nutrition*. 2015; 66: 830–836. DOI:10.3109/09637486.2015.1093610

- [118] Mohanty P, Hamouda W, Garg R, Aljada A, Ghanim H, Dandona P. Glucose challenge stimulates reactive oxygen species (ROS) generation by leucocytes. *The Journal of Clinical Endocrinology & Metabolism*. 2000; 85: 2970–2973.
- [119] Ghanim H, Sia CL, Upadhyay M, Korzeniewski K, Viswanathan P, Abuaysheh S, Mohanty P, Dandona P. Orange juice neutralizes the proinflammatory effect of a high-fat, high-carbohydrate meal and prevents endotoxin increase and Toll-like receptor expression. *The American Journal of Clinical Nutrition*. 2010; 91: 940–949. DOI:10.3945/ajcn.2009.28584
- [120] Khan RA, Mallick N, Feroz Z. Anti-inflammatory effects of *Citrus sinensis* L., *Citrus paradisi* L. and their combinations. *Pakistan Journal of Pharmaceutical Sciences*. 2016; 29: 843–852.
- [121] Javed S, Javaid A, Nawaz S, Saeed MK, Mahmood Z, Siddiqui SZ, Ahmad R. Phytochemistry, GC-MS analysis, antioxidant and antimicrobial potential of essential oil from five Citrus species. *Journal of Agricultural Science*. 2014; 6: 201–208. DOI:10.5539/jas.v6n3p201
- [122] Dixon, R. A., Dey, P. M. and Lamb, C. J. Phytoalexins: Enzymology and Molecular Biology, in *Advances in Enzymology and Related Areas of Molecular Biology*. 1983; 55: (ed A. Meister), John Wiley & Sons, Inc., Hoboken, NJ, USA. DOI:10.1002/9780470123010
- [123] Carvalho M, Matos M, Carnide V. Fingerprinting of *Vaccinium corymbosum* cultivars using DNA of fruits. *Horticultural Science*. 2014; 41: 175–184.
- [124] Luby JJ, Ballington JR, Draper AD, Pliszka K, Austin ME. Blueberries and cranberries (*Vaccinium*). *Genetic Resources of Temperate Fruit and Nut Crops*. 1991; 290: 393–458. DOI:10.17660/ActaHortic.1991.290.9
- [125] Galleta, GJ, Ballington JR. Blueberries, cranberries and lingonberries In: Janick J, Moore JN, Editors. *Fruit breeding*. Vol 2. New York: Wiley; 1996. p. 1–107.
- [126] Levi A, Rowland LJ. Identifying blueberry cultivars and evaluating their genetic relationships using randomly amplified polymorphic DNA (RAPD) and simple sequence repeat-(SSR-) anchored primers. *Journal of the American Society for Horticultural Science*. 1997; 122: 74–78.
- [127] Figueira ME, Oliveira M, Direito R, Rocha J, Alves P, Serra AT, Duarte C, Bronze R, Fernandes A, Brites D, Freitas M, Fernandes E, Freitas M. Protective effects of a blueberry extract in acute inflammation and collagen-induced arthritis in the rat. *Biomedicine & Pharmacotherapy*. 2016; 83: 1191–1202. DOI:10.1016/j.biopha.2016.08.040
- [128] Skupien K, Oszmiański J. Comparison of six cultivars of strawberries (*Fragaria x ananassa* Duch.) grown in northwest Poland. *European Food Research and Technology*. 2004; 219: 66–70. DOI:10.1007/s00217-004-0918-1
- [129] Fernández-Pancho MS, Villano D, Troncoso AM, Garcia-Parrilla MC. Antioxidant activity of phenolic compounds: from in vitro results to in vivo evidence. *Critical Reviews in Food Science and Nutrition*. 2008; 48: 649–671. DOI:10.1080/10408390701761845

- [130] Chiabrando V, Giacalone G. Anthocyanins, phenolics and antioxidant capacity after fresh storage of blueberry treated with edible coatings. *International Journal of Food Sciences and Nutrition*. 2015; 66: 248–253. DOI:10.3109/09637486.2014.986075
- [131] Braga PC, Antonacci R, Wang YY, Lattuada N, Dal Sasso M, Marabini L, Fibiani M, Lo Scalzo R. Comparative antioxidant activity of cultivated and wild *Vaccinium* species investigated by EPR, human neutrophil burst and COMET assay. *European Review for Medical and Pharmacological Sciences*. 2013; 17: 1987–1999.
- [132] Contreras RA, Köhler H, Pizarro M, Zúiga GE. In vitro cultivars of *Vaccinium corymbosum* L. (Ericaceae) are a source of antioxidant phenolics. *Antioxidants*. 2015; 4:281–292. DOI:10.3390/antiox4020281
- [133] Yi W, Fischer J, Krewer G, Akoh CC. Phenolic compounds from blueberries can inhibit colon cancer cell proliferation and induce apoptosis. *Journal of agricultural and food chemistry* 2005; 53: 7320–7329. DOI:10.1021/jf051333o
- [134] Pervin M, Hasnat MA, Lim BO. Antibacterial and antioxidant activities of *Vaccinium corymbosum* L. leaf extract. *Asian Pacific Journal of Tropical Disease*. 2013; 3: 444–453. DOI:10.1016/S2222-1808(13)60099-7
- [135] Aktan AH, Ozcelik A, Cure E, Cure MC, Yuce S. Profound hypoglycemia-induced by *Vaccinium corymbosum* juice and laurocerasus fruit. *Indian Journal of Pharmacology*. 2014; 46: 446–447. DOI:10.4103/0253-7613.135963
- [136] Cheplick S, Sarkar D, Bhowmik P, Shetty K. Phenolic bioactives from developmental stages of highbush blueberry (*Vaccinium corymbosum*) for hyperglycemia management using in vitro models. *Canadian Journal of Plant Science*. 2015; 95: 653–662. DOI:10.4141/CJPS-2014-352
- [137] Shen X, Sun X, Xie Q, Liu H, Zhao Y, Pan Y, Cheng-An H, Wu VC. Antimicrobial effect of blueberry (*Vaccinium corymbosum* L.) extracts against the growth of *Listeria monocytogenes* and *Salmonella enteritidis*. *Food Control*. 2014; 35: 159–165. DOI:10.1016/j.foodcont.2013.06.040
- [138] Silva S, Costa EM, Pereira MF, Costa MR, Pintado ME. Evaluation of the antimicrobial activity of aqueous extracts from dry *Vaccinium corymbosum* extracts upon food micro-organism. *Food Control*. 2013; 34: 645–650. DOI:10.1016/j.foodcont.2013.06.012
- [139] Silva S, Costa EM, Costa MR, Pereira MF, Pereira JO, Soares, JC, Pintado MM. Aqueous extracts of *Vaccinium corymbosum* as inhibitors of *Staphylococcus aureus*. *Food Control*. 2015; 51: 314–320. DOI:10.1016/j.foodcont.2014.11.040
- [140] Zhong Y, Wang Y, Guo J, Chu H, Gao Y, Pang L. Blueberry improves the therapeutic effect of etanercept on patients with juvenile idiopathic arthritis: Phase III Study. *The Tohoku Journal of Experimental Medicine*. 2015; 237:183–191. DOI:10.1620/tjem.237.183
- [141] Samad NB, Debnath T, Ye M, Hasnat MA, Lim BO. In vitro antioxidant and anti-inflammatory activities of Korean blueberry (*Vaccinium corymbosum* L.) extracts. *Asian Pacific Journal of Tropical Biomedicine*. 2014; 4: 807–815. DOI:10.12980/APJTB.4.2014C1008

- [142] Massarotto G, Barcellos T, Garcia CSC, Brandalize APC, Moura S, Schwambach J, Henriques JAP, Roesch-Ely M. Chemical characterization and cytotoxic activity of blueberry extracts (cv. Misty) cultivated in Brazil. *Journal of Food Science*. 2016; 81. DOI:10.1111/1750-3841.13385
- [143] Tsuda H, Kunitake H, Kawasaki-Takaki R, Nishiyama K, Yamasaki M, Komatsu H, Yukizaki C. Antioxidant activities and anti-cancer cell proliferation properties of natuhaze (*Vaccinium oldhamii* Miq.), shashanbo (*V. bracteatum* Thunb.) and blueberry cultivars. *Plants*. 2013; 2: 57–71. DOI:10.3390/plants2010057
- [144] Coban J, Evran B, Özkan F, Çevik A, Doğru-Abbasoğlu S, Uysal M. Effect of blueberry feeding on lipids and oxidative stress in the serum, liver and aorta of guinea pigs fed on a high-cholesterol diet. *Bioscience, Biotechnology, and Biochemistry*. 2013; 77: 389–391. DOI:10.1271/bbb.120722
- [145] Cutler BR, Petersen C, AnandhBabu PV. Mechanistic insights into the vascular effects of blueberries: Evidence from recent studies. *Molecular Nutrition and Food Research*. 2016. DOI:10.1002/mnfr.201600271
- [146] Oliveira GAF, Pádua JG, Costa JL, Jesus OND, Carvalho FMD, Oliveira EJD. Cross-species amplification of microsatellite loci developed for *Passiflora edulis* Sims. in related *Passiflora* Species. *Brazilian Archives of Biology and Technology*. 2013; 5: 785–792.
- [147] Vanderplank J, *Passion flowers*. 2nd ed. MIT Press, Cambridge, Mass; 1996.
- [148] Vitta FA, Bernacci LC A new species and two overlooked species of *Passiflora* (*Passifloraceae*) from Brazil. *Brittonia*. 2004; 56: 89–95.
- [149] Bernacci LC, Soares-Scott MD, Junqueira NTV, Passos IRDS, Meletti LMM. *Passiflora edulis* Sims: the correct taxonomic way to cite the yellow passion fruit (and of others colors). *Revista Brasileira de Fruticultura*. 2008; 2: 566–576.
- [150] Ocampo J, Coppens D’G, Jarvi A. Distribution of the genus *Passiflora* L. diversity in Colombia and its potential as an indicator for biodiversity management in the coffee growing zone. *Diversity*. 2010; 11: 1158–1180.
- [151] Ortiz DC, Bohórquez A, Duque MC, Tohme J, Cuéllar D, Vásquez TM. Evaluating purple passion fruit (*Passiflora edulis* Sims f. *edulis*) genetic variability in individuals from commercial plantations in Colombia. *Genetic Resources and Crop Evolution*. 2012; 6: 1089–1099.
- [152] Talcott ST, Percival SS, Pittet-Moore J, Celoria C. Phytochemical composition and antioxidant stability of fortified yellow passion fruit (*Passiflora edulis*). *Journal of Agricultural and Food Chemistry*. 2003; 4: 935–941.
- [153] Ramaiya S, Bujang JS, Zakaria MH, King WS, Sahrir S, Arif M. Sugars, ascorbic acid, total phenolic content and total antioxidant activity in passion fruit (*Passiflora*) cultivars. *Journal of the Science of Food and Agriculture*. 2012; 5: 1198–1205.

- [154] Da Silva LMR, De Figueiredo EAT, Ricardo NMPS, Vieira IGP, De Figueiredo RW, Brasil IM, Gomes CL. Quantification of bioactive compounds in pulps and by-products of tropical fruits from Brazil. *Food Chemistry*. 2014; 143: 398–404.
- [155] Zucolotto SM, Fagunde C, Reginatto FH, Ramos FA, Castellanos L, Duque C, Schenkel EP. Analysis of C-glycosyl Flavonoids from South American *Passiflora* Species by HPLC-DAD and HPLC-MS. *Phytochemical Analysis*. 2012, 3; 232–239.
- [156] Maruki-Uchida H, Kuritak, Sugiyam, K, Sai M, Maeda, K., Ito, T. (2013). The protective effects of piceatannol from passion fruit (*Passiflora edulis*) seeds in UVB-irradiated keratinocytes. *Biological and Pharmaceutical Bulletin*, 36(5), 845–849.
- [157] Marcoris MS, De Marchi R, Janzantti N, Monteiro M. The influence of ripening stage and cultivation system on the total antioxidant activity and total phenolic compounds of yellow passion fruit pulp. *Journal of the Science of Food and Agriculture*. 2012; 9: 1886–1891.
- [158] Martínez R, Torres P, Meneses MA, Figueroa JG, Pérez-Álvarez JA, Viuda-Martos M. Chemical, technological and *in vitro* antioxidant properties of mango, guava, pineapple and passion fruit dietary fibre concentrate. *Food Chemistry*. 2012; 3: 1520–1526.
- [159] Sato CA, Andrade AS, Brito VM, Sampaio MAUM, Maffei H de AF, Oliva LVM. Effects of compounds from *Passiflora edulis* Sims f. flavicarpa juice on blood coagulation and on proteolytic enzymes. *Protein and Peptide Letters*. 2012; 5: 501–508.
- [160] Klein N, Gazola AC, Lima TCM, Schenkel E, Nieber K, Butterweck V. Assessment of sedative effects of *Passiflora edulis* f. flavicarpa and *Passiflora alata* extracts in mice, measured by telemetry. *Phytotherapy Research*. 2014; 5: 706–713.
- [161] Li H, Zhou P, Yang Q, Shen Y, Deng J, Li L, Zhao D. Comparative studies on anxiolytic activities and flavonoid compositions of *Passiflora edulis* 'edulis' and *Passiflora edulis* 'flavicarpa'. *Journal of Ethnopharmacology*. 2011; 3: 1085–1090.
- [162] Figueiredo DA, Pordeus L, Paulo LL, Braga RM, Fonsêca DV, Sousa BS, Costa MJC, Goncalves MCR, Oliveir KH. Effects of bark flour of *Passiflora edulis* on food intake, body weight and behavioral response of rats. *Revista Brasileira de Farmacognosia*. 2016; 5: 595–600.
- [163] Aguillón OJ, Maldonado ME, Loango CN, Arango VSS, Landázuri P. (2013). Antioxidant and antiproliferative activity of ethanolic and aqueous extracts from leaves and fruits juice *Passiflora edulis*. *Perspectivas en Nutrición Humana*. 2013; 1: 13–25.
- [164] Silva RO, Damasceno SR, Brito TV, Dias JM, Fontenele AM, Braúna IS, Junior JSC, Miaciél JS, Paulac RCM, Ribeiroa RA, Souza HLP, Freitas ALP, Medeiros Jand-Venes, Silva DC, Barbosa ALR, Polysaccharide fraction isolated from *Passiflora edulis* inhibits the inflammatory response and the oxidative stress in mice. *Journal of Pharmacy and Pharmacology*. 2015; 7: 1017–1027.

- [165] Silva DC, Freitas AL, Pessoa CD, Paula RC, Mesquita JX, Leal L K, Brito GAC, Goncalves OD, Viana GS. Pectin from *Passiflora edulis* shows anti-inflammatory action as well as hypoglycemic and hypotriglyceridemic properties in diabetic rats. *Journal of Medicinal Food*. 2011; 10: 1118–1126.
- [166] Salgado JM, Bombarde TAD, Mansi DN, Piedade SMDS, Meletti LMM. Effects of different concentrations of passion fruit peel (*Passiflora edulis*) on the glicemic control in diabetic rat. *Food Science and Technology (Campinas)*, 2010; 3: 784–789.
- [167] Barbalho SM, Damasceno DC, Spada APM, Lima IEDR, Araújo AC, Guiguer EL, Martuchi KA, Oshiiwa M, Mendes CG. Effects of *Passiflora edulis* on the metabolic profile of diabetic wistar rat offspring. *Journal of Medicinal Food*. 2011; 12: 1490–1495.
- [168] Kirchhoff R, Beckers C, Kirchhoff GM, Trinczek-Gärtner H, Petrowicz O, Reimann HJ. Increase in choleresis by means of artichoke extract. *Phytomedicine*. 1994; 1: 107–115. DOI:10.1016/S0944-7113(11)80027-9
- [169] Gebhardt R. Antioxidative and protective properties of extracts from leaves of the artichoke (*Cynara scolymus* L.) against hydroperoxide-induced oxidative stress in cultured rat hepatocytes. *Toxicology and Applied Pharmacology*. 1997; 144: 279–286. DOI:10.1006/taap.1997.8130
- [170] Lattanzio V, Kroon PA, Linsalata V, Cardinali A. Globe artichoke: a functional food and source of nutraceutical ingredients. *Journal of Functional Foods*. 2009; 1: 131–144. DOI:10.1016/j.jff.2009.01.002
- [171] Gebhardt R. Inhibition of cholesterol biosynthesis in HepG2 cells by artichoke extracts is reinforced by glucosidase pretreatment. *Phytotherapy Research*. 2002; 16: 368–372. DOI:10.1002/ptr.960
- [172] Movahedian A, Sadeghi H, Ghannadi A, Gharavi M, Azarpajoo S. Hypolipidemic activity of *Allium porrum* L. in cholesterol-fed rabbits. *Journal of Medicinal Food*. 2006; 9: 98–101. DOI:10.1089/jmf.2006.9.98
- [173] Roghani M, Aghaei M. The effect *Allium ampeloprasum* feeding on serum level of glucose, triglyceride, and total cholesterol of diabetic rats. *Journal of Semnan University of Medical Science*. 2007; 8: 73–78.
- [174] Del Rio D, Rodriguez-Mateos A, Spencer JP, Tognolini M, Borges G, Crozier A. Dietary (poly) phenolics in human health: structures, bioavailability, and evidence of protective effects against chronic diseases. *Antioxidants & Redox Signaling*. 2013; 18: 1818–1892. DOI:10.1089/ars.2012.4581
- [175] Pak N. La fibra dietética en la alimentación humana, importancia en la salud. *Anales de la Universidad de Chile*. 2000. DOI:10.5354/0717-8883.2000.2504
- [176] Bhat S, Kaushal P, Kaur M, Sharma HK. Coriander (*Coriandrum sativum* L.): Processing, nutritional and functional aspects. *African Journal of Plant Science*. 2014; 8: 25–33. DOI:10.5897/AJPS2013.1118

- [177] Forastiere F, Pistelli R, Sestini P, Fortes C, Renzoni E, Rusconi F, Dell' Orco V, Ciccone G, Bisanti L. Consumption of fresh fruit rich in vitamin C and wheezing symptoms in children. *Thorax*. 2000; 55: 283–288. DOI:10.1136/thorax.55.4.283
- [178] Del Bo C, Riso P, Campolo J, Møller P, Loft S, Klimis-Zacas D, Brambilla A, Rizzolo A, Porrini M. A single portion of blueberry (*Vaccinium corymbosum* L) improves protection against DNA damage but not vascular function in healthy male volunteers. *Nutrition Research*. 2013; 33: 220–227. DOI:10.1016/j.nutres.2012.12.009
- [179] Del Bo C, Porrini M, Fracassetti D, Campolo J, Klimis-Zacas D, Riso P. A single serving of blueberry (*V. corymbosum*) modulates peripheral arterial dysfunction induced by acute cigarette smoking in young volunteers: a randomized-controlled trial. *Food and Function*. 2014; 5: 3107–3116. DOI:10.1039/c4fo00570h