We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Near-Infrared Spectroscopy Combined with Multivariate Tools for Analysis of Trace Metals in Environmental Matrices

Philiswa N. Nomngongo,
Tshimangadzo S. Munonde, Anele Mpupa and
Nkositatile Raphael Biata

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/67199>

Abstract

Environmental contamination by trace elements is becoming increasingly important problem worldwide. Trace metals such as cadmium, copper, lead, chromium, and mercury are major environmental pollutants that are predominantly found in areas with high anthropogenic activities. Therefore, there is a need for rapid and reliable tools to assess and monitor the concentration of heavy metal in environmental matrices. A nondestructive, cost-effective, and environmentally friendly procedure based on near-infrared reflectance spectroscopy (NIRS) and chemometric tools has been used as alternative technique for the simultaneous estimation of various heavy metal concentrations in environmental sample. The metal content is estimated by assigning the absorption features of metals associated with molecular vibrations of organic and inorganic functional groups in organic matter, silicates, carbonates, and water at 780–2500 nm in the near-infrared region. This chapter, reviewed the application of NIRS combined with chemometric tools such as multiple linear regression (MLR), principal component regression (PCR), and partial least squares (PLS) regression. The disadvantages and advantages of each chemometric tool are discussed briefly.

Keywords: near-infrared spectroscopy, principal component regression, partial least squares, multiple linear regression, trace metals

1. Introduction

Due to fast industrial development and growth that have happened in most areas of the world during the recent years, water and soil are getting a large amount of pollutants such as trace

elements from different sources [1]. This can, however, lead to environmental contamination, thus affecting the ecosystem. Contamination refers to the condition of the land or water where any chemical substance or waste has been added at the above background level. The signifies of water, air, and land pollution include an adverse health or environmental impact [2–4] runoff, aerial deposition of chemicals used for agriculture or industry, materials stored or dumped on the site, and contaminants in imported fill, and building demolition can also result in contamination of the soil and water that are close to residential communities [5]. Contaminants such as trace metals may be introduced into drinking water via the aforementioned activities or leached from the soil into groundwater [6]. Additionally, trace metals occur naturally in the earth's crust [7]. For this reason, they can be present in soils at a background level. Trace metals persist for a long time in the environment because they are not degradable. In addition, they are translocated to different components, thus affecting the biota [2–4]. The persistence of trace metals can result in bioaccumulation and biomagnifications causing heavier exposure for some living organisms that are present in the environment [8].

Trace metal contaminations threaten agriculture and other food sources for human population as well as poor vegetation growth and that lower plant resistance against pests [9]. This situation poses different kinds of challenges for remediation. Furthermore, people can be exposed to contaminants in the soil through different ways. These include dermal exposure or inhalation and penetration via the skin or eyes (includes exposure to dust) [5]. Trace metal exposure is normally chronic (exposure over a longer period of time), due to food chain transfer [2]. But the case of acute (immediate) poisoning is rare through ingestion or dermal contact but is possible [2]. The toxicity of trace metals is one of the major environmental health concerns and potentially dangerous due to bioaccumulation through the food chain [4].

In view of the abovementioned challenges, the development of sensitive and selective analytical procedure for the determination of trace metals is of great importance. Flame atomic absorption spectrometry (FAAS) [10], graphite furnace atomic absorption spectrometry (GFAAS) [11], cathodic and anodic stripping voltammetry [12, 13], inductively coupled plasma optical emission spectrometry (ICP-OES) [14], and inductively coupled plasma mass spectrometry (ICP-MS) [14] are the most widely used analytical techniques for determination of trace metals in different matrices. However, these techniques are expensive, tedious, complex, and highly time-consuming [15–17]. In addition, investigation of trace metal concentration distributions in environmental matrices is based on numerous samples and laboratory analysis. Therefore, a rapid, reliable, and environmentally friendly method is required to detect and survey the distribution of trace metals in environmental matrices. This is done in order to diagnose suspected contaminated areas as well as control the rehabilitation processes [18]. Reflectance spectroscopy is the study of the absorption and emission of light and other radiations by matter as related to the dependence of these processes on the wavelength of the radiation [19]. It is based on the distinct vibrations and electronic processes of chemical bonds in molecules [20]. These vibrations can be observed in three regions, namely, far IR ($25 \times 10^3 \text{ nm} - 1 \times 10^6 \text{ nm}$), mid-IR ($25 \times 10^2 \text{ nm} - 25 \times 10^3 \text{ nm}$), and near IR ($8 \times 10^2 \text{ nm} - 25 \times 10^2 \text{ nm}$), with mid-IR and near IR being the most useful for qualitative and quantitative analysis [21]. The MIRS is known to have a greater predictive ability for soil geochemistry compared with the visible-near infrared (vis-NIR) [22, 23]. However, more

interest is on NIRs because it is generally cheaper [23]. In NIR, many trace elements are spectrally featureless and only exhibit characteristic spectral features at high concentrations ($>4000 \text{ mg kg}^{-1}$) [23, 24]. Therefore, due to high detection, low levels of heavy metals can be indirectly determined from spectra due to their association with Fe oxides, clays, and organic matter [24]. For this reason, cost-effective and nondestructive analytical techniques based on near-infrared (NIR) spectroscopy coupled with chemometrics have been developed to overcome the problems encountered when using traditional methods [17, 25].

The aim of this chapter is to review the application of near-infrared spectroscopy (NIRS) combined with chemometrics for the estimation of the concentration and distribution of trace elements in environmental matrices. The disadvantages of NIRS without chemometrics for analysis of trace elements are discussed. In addition, this chapter aims at promoting the application of simpler and greener methods such as the combination of NIRS and multivariate tools for monitoring of trace metal contaminations in different matrices.

2. Application of near-infrared spectroscopy (NIRS) for analysis of trace metals

Near-infrared spectroscopy (NIRS) is a fast and nondestructive analytical technique that is used to provide multi-constituent analysis of almost all types of sample matrices [15–17, 25]. This technique covers wavelength range closer to the mid-infrared and broadens up to the visible region [26, 27]. Typically, NIRS is primarily based on absorbance characteristics caused by vibrations of covalent bonds between H, C, O, S, and N, which are the main components of the organic matter [28]. Pure metals do not absorb in the NIR region [29]. However, their indirect detection is possible via their complexation with organic molecules containing C–H, N–H, and O–H bonds, which are detectable [30]. This concept was termed “aquaphotomics” which is based on fact that the characteristic absorbance pattern of water (O–H overtones) can change as a consequence of the binding reaction with the metal [31, 32]. This effect is described and demonstrated in the study carried out by Putra et al.; although some complexes might be similar in different samples, slight differences in spectral features such as shifts in peak wavelength may still be seen depending on the nature of the cation [30]. In addition, the electromagnetic radiation spectrum in the near-infrared region contains useful information about environmental sample constituents such as soil that can be used for prediction of metal concentrations [25, 28, 33]. For instance, the absorption features associated with electronic transitions of Fe^{3+} and Fe^{2+} ions in Fe-bearing minerals can be found in the near-infrared region at 780–1200 nm [33–35]. In addition, the absorption features of metals associated with molecular vibrations of organic and inorganic functional groups in organic matter, silicates, carbonates, and water can be found in near infrared at 780–2500 nm region [24, 33, 34, 36].

Wu et al. [24, 36] reported the feasibility of using NIRS for monitoring and predicting trace metals in suspended solids, sediment, and soil. This was achieved by quantitative evaluations of the spectral activity of sediment and soil properties [23–37]. However, due to challenges such as the collinearity, band overlaps, and interactions for some soil properties, the spectra of soil, sediment, or suspended solids are often broad and nonspecific [25]. To overcome these

challenges, some chemometric tools have been used to be applied to the quantitative analysis of the spectroscopic data [38]. These chemometric tools include multiple linear regression (MLR) [39], principal component regression (PCR) [40, 41], and partial least squares (PLS) regression [42]. These chemometric tools have been used to characterize soil spectra and build models for estimating the trace metal concentrations in soil or sediments and other matrices [25].

2.1. Applications of NIRS-multi-linear regression determination trace metals in environmental samples

Multiple linear regression (MLR) is a conventional chemometric method that commonly correlates a linear combination of several selected spectral bands/indices which have high correlations with the heavy metal concentrations [18]. The disadvantage of using MLR is that it does not perform well with hyperspectral measurements. This is because the NIR spectral data usually exhibit high collinearity [18, 39]. This challenge has been solved by applying the enter and stepwise MLR approaches [18, 39]. In enter-MLR approach, a procedure for variable selection is adopted, and the selected variables are then used to calibrate the MLR model [18]. In stepwise-MLR approach, on the other hand, a forward or backward method is applied to progressively select the independent variables according to a tolerance significance level, which is generally set to 0.05 [18, 39]. Due to the challenges associated with MLS, there are very few reports on its application together with NIRS for determination of trace elements in environmental samples.

Kemper and Sommer [39] explored the possibility to adapt chemometrics approaches for the quantitative estimation of As, Cd, Cu, Fe, Hg, Pb, S, Sb, and Zn in polluted soils using stepwise multiple linear regression (MLR) analysis and an artificial neural network (ANN) approach. The authors reported that the models predicted six out of nine elements with high accuracy. In addition, it was discovered that most wavelengths important for prediction were attributed to absorption features of iron and iron oxides. Furthermore, their results revealed the feasibility to predict heavy metals in contaminated soils using the rapid and cost-effective NIRS. Other applications are reported by Malley et al. [40] and Choe et al. [41].

2.2. Applications of NIRS-PCR for determination trace metals in environmental samples

Principal component regression (PCR) is a chemometric tool that combines principal component analysis and MLR [18, 42]. In this method, the independent variables are first decomposed into orthogonal principal components using the nonlinear iterative partial least squares algorithm and full cross validation of the calibration set [18]. The maximum number of principal components is then defined according to the minimum value of the root-mean-square error of the cross validation [18]. In the final step, the chosen principal components are used to calibrate the MLR models [43]. The advantage of PCR over the normal MLR is that the principal components are uncorrelated and the noise is filtered [18]. There is very limited information on the application of NIRS-PCR on the analysis of trace metals in different matrices. However, some of the reports are available in the literature [42, 43].

Wu et al. [36] reported the practicality of using NIRS for the determination of Hg concentration agricultural soil samples. The accuracy of the prediction models was optimized by applying several spectral pretreatments to the reflectance spectra. The univariate regression and principal component regression were used for the prediction of Hg concentration. According to their results, the optimal model was achieved using the PCR combined with Kubelka-Munk transformation. In addition, the results obtained from the correlation analysis revealed that Hg concentration correlated negatively with soil reflectance, while positively with the absorption depths of goethite at 0.496 μm and clay minerals at 2.21 μm [36]. The findings suggest that the adsorption of Hg by clay-size mineral accumulations in soils was the mechanism that can be used to predict the spectral absorption band of Hg.

2.3. Applications of NIRS-partial least squares regression environmental samples

Partial least squares regression (PLSR) is a chemometric method that is widely used to quantitatively derive information from NIR spectra [18, 44]. The PLSR allows a refined statistical approach using the full spectral region rather than unique and isolated analytical bands [44]. The principle of PLSR is based on incorporation of the dependent variables in the calculation of the principal components [45, 46]. For this reason, the PLSR is able to handle data with strong collinearity and noise [18, 44]. In addition, PLSR provides the possibility of handling cases where the number of variables significantly exceeds the number of available samples [47]. The applications of NIRS combined PLSR for the determination of metals in different environmental matrices have been widely reported in the literature (see **Table 1**).

Moros et al. [15] evaluated the potential of near-infrared (NIR) diffuse reflectance infrared Fourier transform spectroscopy (DRIFTS) combined with PLSR for nondestructive determinations of trace elements in foods. This analysis was achieved without physical or chemical sample pretreatment. The authors compared two spectral pretreatments that are multiplicative signal correction (MSC) and standard normal variate (SNV). Their results revealed that the PLS models built after using SNV provided the best prediction results for the determination of arsenic and lead in powdered red paprika samples. The concluded results showed that NIR diffuse reflectance spectroscopy combined with the PLS could be used to estimate the concentration of As and Pb at 100 $\mu\text{g kg}^{-1}$ level with a standard error of prediction of 39 and 50 $\mu\text{g kg}^{-1}$ for As and Pb, respectively. Furthermore, the estimated percentage errors were lower than 25% without the need of using sophisticated and high-cost instrumentation (such as ICP-MS and GFAAS) together with tedious and expensive digestion procedures for sample preparation. Moreover, the suggested NIRS-PLSR methodology was found to be an important tool for screening of trace elements in foods in the laboratories [15].

In another study, Li et al. [48] reported a method for simultaneous determination of mercury, lead, and cadmium ions in water samples using solid-phase extraction and near-infrared diffuse reflectance spectroscopy (NIRDRS). In order to analyze trace metal content in water samples using NIRS, thiol-functionalized magnesium phyllosilicate (Mg-MTMS) was used as an adsorbent for extraction of target analytes from aqueous solution. The adsorbed metals were measured using NIRDRS combined with PLS models. This combination resulted in fast

Analytes	Matrix	Concentra ions	Ref.
Cr, Co, Ni, Cu, Zn, As, Se, Cd, and Tl	Soil	0.32–110 $\mu\text{g g}^{-1}$	[16]
Zn, Pb, and Cd	Soil	0.17–6530 mg kg^{-1}	[41]
Pb^{2+} , Zn^{2+} , Cu^{2+} , Cd^{2+} , and Cr^{3+}	Water		[42]
Cd and Zn	Soil	2.25–51.48 mg L^{-1}	[43]
K, Ca, Mg, Fe, and Zn	Manure compost	0.676–80.97 mg kg^{-1}	[44]
Cd, Cu, Zn, Pb, Ni, Mn, and Fe	Freshwater sediments	7.63–198.20 g kg^{-1}	[45]
Zn and Pb	Soil	2–425 mg g^{-1}	[46]
Al, Ag, As, Ba, Be, Bi, Ca, Cd, Ce, Cs, Co, Cr, Cu, Fe, Ga, In, K, La, Li, Mg, Mn, Mo, Na, Nb, Ni, P, Pb, Rb, S, Sb, Sc, Sn, Sr, Te, Th, Ti, Tl, U, V, W, Y, and Zn	Soil	1672–4601 mg kg^{-1}	[47]
Fe, Zn, Mn, and Cu	Agricultural soils	50–100 mg kg^{-1}	[48]
Cu, Mn, Zn, and Fe	Water and HNO_3	–	[49]
		1–10,000 $\mu\text{g L}^{-1}$	

Table 1. Applications of NIRS-PLS regression for determination of trace metals in environmental samples.

and simultaneous quantitative prediction. The metal ions interacted with a functional group of the adsorbent and their absorption bands were observed in the spectra, thus leading to an efficient and precise prediction models. The concentration of the three metals that can be correctly determined was found to range between 4 and 6 mg L^{-1} . This proved that the adsorbent used (thiol-functionalized magnesium phyllosilicate) had a high efficiency for the enrichment of Hg, Pb, and Cd in dilute solution. Furthermore, the results obtained revealed the feasibility of NIRDRS-PLSR for quantitative analysis metal ions in river water. Other applications of NIRS combined with PLSR are presented in **Table 1**.

2.4. Nonlinear calibration models for near-infrared spectroscopy

The abovementioned linear calibration models (especially PLS and PCR) have been extensively due to their ease of use, fast computation, good predictive performance, and easy interpretable representations [49]. However, the linearity assumption is not always valid, and when the spectra exhibit nonlinearities, they tend to give nonoptimal results [49]. Therefore, in such cases, it is of greater significance to develop the robust model system based on different nonlinear calibrations [49, 50]. These models include kernel PLS (KPLS), support vector machines (SVM), least squares SVM (LS-SVM), and among other artificial neural networks (ANN). Brief descriptions of these models are given in the subsequent paragraphs.

Kernel PLS is a nonlinear extension of linear PLS in which input data are transformed into a high-dimensional feature space via nonlinear mapping [51]. Briefly, the KPLS includes two steps. The first step includes embedding data in an input space via nonlinear mapping. The second step is that a linear algorithm is designed to discover the linear relationship [52]. The ANN on the other hand is a flexible mathematical structure capable of identifying complex

nonlinear interactions between input and output data sets. This method is reported to be useful and efficient, especially in problems for which the characteristics are difficult to describe using physical equations [53]. The ANN model has been shown to perform better than linear models [54].

Support vector machines are the models that involve a solution of a quadratic programming problem leading to global models that are often unique [55]. The application of this type of model is further discussed and investigated by the authors in Refs. [56–59]. Finally, LS-VSM model is a simplification of the computational calculations of SVM by implementation of a least squares version for SVM [51]. Least squares SVM is capable of dealing with both linear and nonlinear multivariate calibration problems relatively fast [60]. In LS-VSM, a linear estimation is done in a kernel-induced feature space; the use of LS-SVM and NIR has been investigated by Borin et al. [55].

There are few studies on application of the nonlinear multivariate calibration in NIR spectroscopy that have been reported for analysis of trace metals. For instance, Shao and He [61] investigated the two sensitive wavelength (SW) selection methods combined with visible-near-infrared (vis/NIR) spectroscopy to determine the levels of some trace elements (Fe, Zn) in rice leaf. Calibration models using SWs selected by latent variables analysis (LVA) and independent component analysis (ICA) and nonlinear regression of a least squares support vector machine (LS-SVM) were developed. In the nonlinear models, six SWs selected by ICA provide the optimal ICA-LS-SVM model when compared with LV-LS-SVM. The coefficients of determination (R^2), root-mean-square error of prediction (RMSEP), and bias by ICA-LS-SVM were 0.6189, 20.6510, and -12.1549 ppm, respectively, for Fe, and 0.6731, 5.5919, and 1.5232 ppm, respectively, for Zn [62]. The overall results indicated vis-NIR spectroscopy combined with ICA-LS-SVM provided accurate determination of trace elements in rice leaf. Other methods are reported by Xu et al. [62] and Barbosa et al. [63], among others.

3. Conclusions

This chapter revealed that NIRS combined with multivariate tools has a great potential tool to improve the understanding of trace metal concentration in environmental matrices. It was also concluded that the use of chemometrics offered a rapid and cost-effective alternative to measure multielement particularly in soils and sediments. However, according to literature chemometric models such as MLS and PCR used in the NIRS are not reliable as compared to PLSR. Therefore, it is important for researchers to select proper chemometrics for their application. Due to the limitations encountered when using some of the chemometric tools, it is necessary to develop new chemometric methods or modify the conventional ones so as to improve their reliability and accuracy. It is reported in the literature that the limitation of NIRS is that some trace metals and other mineral compounds such as phosphorus do not absorb radiation in the NIR region. However, in most cases, this problem is solved by using the absorption features of metals associated with molecular vibrations of organic and inorganic functional groups in organic matter, silicates, carbonates, and water in near infrared (780–2500 nm region). Alternatively, some researchers combine different detection techniques such as UV-visible with NIRS [17, 64].

Author details

Philiswa N. Nomngongo*, Tshimangadzo S. Munonde, Anele Mpupa and Nkositile Raphael Biata

*Address all correspondence to: pnnomngongo@uj.ac.za

Department of Applied Chemistry, University of Johannesburg, Johannesburg, South Africa

References

- [1] Wei, B., & Yang, L. (2010). A review of heavy metal contaminations in urban soils, urban road dusts and agricultural soils from China. *Microchemical Journal*, 94(2), 99–107.
- [2] Kumar, P., & Singh, A. (2010). Cadmium toxicity in fish: an overview. *GERF Bulletin of Biosciences*, 1(1), 41–47.
- [3] Obodai, E. A., Boamponsem, L. K., Adokoh, C. K., Essumang, D. K., Villawoe, B. O., Aheto, D. W., & Debrah, J. S. (2011). Concentrations of heavy metals in two Ghanaian Lagoons. *Archives of Applied Science Research*, 3, 177–187.
- [4] Rajaganapathy, V., Xavier, F., Sreekumar, D., & Mandal, P. K. (2011). Heavy metal contamination in soil, water and fodder and their presence in livestock and products: a review. *Journal of Environmental Science and Technology*, 4(3), 234–249.
- [5] Li, Z., Ma, Z., van der Kuijp, T. J., Yuan, Z., & Huang, L. (2014). A review of soil heavy metal pollution from mines in China: pollution and health risk assessment. *Science of the Total Environment*, 468, 843–853.
- [6] El-Bouraie, M. M., El-Barbary, A. A., Yehia, M. M., & Motawea, E. A. (2010). Heavy metal concentrations in surface river water and bed sediments at Nile Delta in Egypt. *Suo*, 61(1), 1–12.
- [7] Aderinola, O. J., Clarke, E. O., Olarinmoye, O. M., Kusemiju, V., & Anatekhai, M. A. (2009). Heavy metals in surface water, sediments, fish and periwinkles of Lagos Lagoon. *American-Eurasian Journal of Agricultural & Environmental Sciences*, 5(5), 609–617.
- [8] Adelekan, B. A., & Abegunde, K. D. (2011). Heavy metals contamination of soil and groundwater at automobile mechanic villages in Ibadan, Nigeria. *International Journal of Physical Sciences*, 6(5), 1045–1058.
- [9] Shaylor, H., McBride, M., Harrison, E. (2009). Sources and Impacts of Contaminants in Soil. Cornell Waste Management Institute, New York (USA), <http://cwmi.css.cornell.edu> accessed 15 May 2016.
- [10] Dimpe, K. M., Ngila, J. C., Mabuba, N., & Nomngongo, P. N. (2014). Evaluation of sample preparation methods for the detection of total metal content using inductively coupled plasma optical emission spectrometry (ICP-OES) in wastewater and sludge. *Physics and Chemistry of the Earth, Parts A/B/C*, 76, 42–48.

- [11] Guo, W., Zhang, P., & Jin, L. (2015). Heat-assisted slurry sampling GFAAS method for determination of lead in food standard reference materials. *Journal of Food Composition and Analysis*, 42, 78–83.
- [12] Armstrong, K. C., Tatum, C. E., Dansby-Sparks, R. N., Chambers, J. Q., & Xue, Z. L. (2010). Individual and simultaneous determination of lead, cadmium, and zinc by anodic stripping voltammetry at a bismuth bulk electrode. *Talanta*, 82(2), 675–680.
- [13] Raoof, J. B., Ojani, R., Alinezhad, A., & Rezaie, S. Z. (2010). Differential pulse anodic stripping voltammetry of silver(I) using p-isopropylcalix[6]arene modified carbon paste electrode. *Monatshefte für Chemie-Chemical Monthly*, 141(3), 279–284.
- [14] Vanini, G., Souza, M. O., Carneiro, M. T., Filgueiras, P. R., Bruns, R. E., & Romão, W. (2015). Multivariate optimisation of ICP OES instrumental parameters for Pb/Ba/Sb measurement in gunshot residues. *Microchemical Journal*, 120, 58–63.
- [15] Moros, J., Llorca, I., Cervera, M. L., Pastor, A., Garrigues, S., & De la Guardia, M. (2008). Chemometric determination of arsenic and lead in untreated powdered red paprika by diffuse reflectance near-infrared spectroscopy. *Analytica Chimica Acta*, 613(2), 196–206.
- [16] Moros, J., Martínez-Sánchez, M. J., Pérez-Sirvent, C., Garrigues, S., & de la Guardia, M. (2009). Testing of the Region of Murcia soils by near infrared diffuse reflectance spectroscopy and chemometrics. *Talanta*, 78(2), 388–398.
- [17] Wang, J., Cui, L., Gao, W., Shi, T., Chen, Y., & Gao, Y. (2014). Prediction of low heavy metal concentrations in agricultural soils using visible and near-infrared reflectance spectroscopy. *Geoderma*, 216, 1–9.
- [18] Shi, T., Chen, Y., Liu, Y., & Wu, G. (2014). Visible and near-infrared reflectance spectroscopy—an alternative for monitoring soil contamination by heavy metals. *Journal of Hazardous Materials*, 265, 166–176.
- [19] Tripathy, S., Middha, A., Swain, S. R. (2016). Assessment of phytoconstituent of five faced *Elaeocarpus ganitrus* beads by FTIR and UV-VIS spectroscopic analysis. *Imperial Journal of Interdisciplinary Research*, 2(10).
- [20] Yuan, X., Luo, K., Zhang, K., He, J., Zhao, Y., Yu, D. (2016). Combinatorial vibration-mode assignment for the FTIR spectrum of crystalline melamine: a strategic approach toward theoretical IR vibrational calculations of triazine-based compounds. *The Journal of Physical Chemistry A*, 120(38), 7427–7433.
- [21] Reichert, A., Sussmann, R. (2016). The Zugspitze radiative closure experiment for quantifying water vapor absorption over the terrestrial and solar infrared—part 3: quantification of the mid-and near-infrared water vapor continuum in the 2500 to 7800 cm^{-1} spectral range under atmospheric conditions. *Atmospheric Chemistry and Physics*, 16(18), 11671–11686.
- [22] Reeves, J. B., & Smith, D. B. (2009). The potential of mid-and near-infrared diffuse reflectance spectroscopy for determining major-and trace-element concentrations in soils from a geochemical survey of North America. *Applied Geochemistry*, 24(8), 1472–1481.

- [23] O'Rourke, S. M., Minasny, B., Holden, N. M., & McBratney, A. B. (2016). Synergistic use of Vis-NIR, MIR, and XRF spectroscopy for the determination of soil geochemistry. *Soil Science Society of America Journal*, 80(4), 888–899.
- [24] Wu, D., He, Y., Shi, J., & Feng, S. (2009). Exploring near and midinfrared spectroscopy to predict trace iron and zinc contents in powdered milk. *Journal of Agricultural and Food Chemistry*, 57(5), 1697–1704.
- [25] Song, S., Yuan, L., Zhang, X., Hayat, K., Chen, H., Liu, F., ... & Niu, Y. (2013). Rapid measuring and modelling flavour quality changes of oxidised chicken fat by electronic nose profiles through the partial least squares regression analysis. *Food Chemistry*, 141(4), 4278–4288.
- [26] Reich, G. (2005). Near-infrared spectroscopy and imaging: basic principles and pharmaceutical applications. *Advanced Drug Delivery Reviews*, 57(8), 1109–1143.
- [27] Chavez, P. F., Sacre, P. Y., De Bleye, C., Netchacovitch, L., Mantanus, J., Motte, H., Schubert, M., Hubert, P., & Ziemons, E. (2015). Active content determination of pharmaceutical tablets using near infrared spectroscopy as Process Analytical Technology tool. *Talanta*, 144, 1352–1359.
- [28] Kleinebecker, T., Poelen, M. D., Smolders, A. J., Lamers, L. P., & Hölzel, N. (2013). Fast and inexpensive detection of total and extractable element concentrations in aquatic sediments using near-infrared reflectance spectroscopy (NIRS). *PLoS One*, 8(7), e70517.
- [29] Shamsoddini, A., Raval, S., & Taplin, R. (2014). Spectroscopic analysis of soil metal contamination around a derelict mine site in the Blue Mountains, Australia. *ISPRS Annals of the Photogrammetry, Remote Sensing and Spatial Information Sciences*, 2(7), 75.
- [30] Putra, A., Meilina, H., & Tsenkova, R. (2012). Use of near-infrared spectroscopy for determining the characterization metal ion in aqueous solution. In *Proceedings of the Annual International Conference, Syiah Kuala University-Life Sciences & Engineering Chapter* (Vol. 2, No. 2), Banda Aceh.
- [31] Tsenkova, R., Kovacs, Z., & Kubota, Y. (2015). Aquaphotomics: near infrared spectroscopy and water states in biological systems. In *Membrane Hydration* (pp. 189–211). Springer International Publishing, Switzerland.
- [32] Schwartz, G., Ben-Dor, E., & Eshel, G. (2012). Quantitative analysis of total petroleum hydrocarbons in soils: comparison between reflectance spectroscopy and solvent extraction by 3 certified laboratories. *Applied and Environmental Soil Science*, 2012, 1–11.
- [33] Xian-Li, X. I. E., Xian-Zhang, P. A. N., & Bo, S. U. N. (2012). Visible and near-infrared diffuse reflectance spectroscopy for prediction of soil properties near a copper smelter. *Pedosphere*, 22(3), 351–366.
- [34] Horta, A., Malone, B., Stockmann, U., Minasny, B., Bishop, T. F. A., McBratney, A. B., Pallasser, P., & Pozza, L. (2015). Potential of integrated field spectroscopy and spatial analysis for enhanced assessment of soil contamination: a prospective review. *Geoderma*, 241, 180–209.

- [35] Huang, Z., Tao, W., Fang, J., Wei, X., & Du, Y. (2009). Multivariate calibration of on-line enrichment near-infrared (NIR) spectra and determination of trace lead in water. *Chemometrics and Intelligent Laboratory Systems*, 98(2), 195–200.
- [36] Wu, Y. Z., Chen, J., Ji, J. F., Tian, Q. J., Wu, X. M. (2005). Feasibility of reflectance spectroscopy for the assessment of soil mercury contamination. *Environmental Science & Technology*, 39(3), 873–878.
- [37] Soriano-Disla, J. M., Janik, L., McLaughlin, M. J., Forrester, S., Kirby, J., Reimann, C., & Team, T. E. G. P. (2013). The use of diffuse reflectance mid-infrared spectroscopy for the prediction of the concentration of chemical elements estimated by X-ray fluorescence in agricultural and grazing European soils. *Applied Geochemistry*, 29, 135–143.
- [38] Hemmateenejad, B., Akhond, M., & Samari, F. (2007). A comparative study between PCR and PLS in simultaneous spectrophotometric determination of diphenylamine, aniline, and phenol: effect of wavelength selection. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, 67(3), 958–965.
- [39] Kemper, T., & Sommer, S. (2002). Estimate of heavy metal contamination in soils after a mining accident using reflectance spectroscopy. *Environmental Science & Technology*, 36(12), 2742–2747.
- [40] Malley, D. F., Yesmin, L., Wray, D., & Edwards, S. (1999). Application of near-infrared spectroscopy in analysis of soil mineral nutrients. *Communications in Soil Science & Plant Analysis*, 30(7–8), 999–1012.
- [41] Choe, E., Kim, K. W., Bang, S., Yoon, I. H., & Lee, K. Y. (2009). Qualitative analysis and mapping of heavy metals in an abandoned Au–Ag mine area using NIR spectroscopy. *Environmental Geology*, 58(3), 477–482.
- [42] Chang, C. W., Laird, D. A., Mausbach, M. J., & Hurburgh, C. R. (2001). Near-infrared reflectance spectroscopy–principal components regression analyses of soil properties. *Soil Science Society of America Journal*, 65(2), 480–490.
- [43] Ko, H. J., Choi, H. L., Park, H. S., & Lee, H. W. (2004). Prediction of heavy metal content in compost using near-infrared reflectance spectroscopy. *Asian Australasian Journal of Animal Sciences*, 17(12), 1736–1740.
- [44] Dupuy, N., & Douay, F. (2001). Infrared and chemometrics study of the interaction between heavy metals and organic matter in soils. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, 57(5), 1037–1047.
- [45] Shi, T., Wang, J., Chen, Y., & Wu, G. (2016). Improving the prediction of arsenic contents in agricultural soils by combining the reflectance spectroscopy of soils and rice plants. *International Journal of Applied Earth Observation and Geoinformation*, 52, 95–103.
- [46] Shi, T., Liu, H., Wang, J., Chen, Y., Fei, T., & Wu, G. (2014). Monitoring arsenic contamination in agricultural soils with reflectance spectroscopy of rice plants. *Environmental Science & Technology*, 48(11), 6264–6272.
- [47] Shao, X., Bian, X., Liu, J., Zhang, M., & Cai, W. (2010). Multivariate calibration methods in near infrared spectroscopic analysis. *Analytical Methods*, 2(11), 1662–1666.

- [48] Li, J., Zhang, Y., Cai, W., & Shao, X. (2011). Simultaneous determination of mercury, lead and cadmium ions in water using near-infrared spectroscopy with preconcentration by thiol-functionalized magnesium phyllosilicate clay. *Talanta*, 84(3), 679–683.
- [49] Ni, W., Nørgaard, L., & Mørup, M. (2014). Non-linear calibration models for near infrared spectroscopy. *Analytica Chimica Acta*, 813, 1–14.
- [50] Chen, Q., Cai, J., Wan, X., & Zhao, J. (2011). Application of linear/non-linear classification algorithms in discrimination of pork storage time using Fourier transform near infrared (FT-NIR) spectroscopy. *LWT-Food Science and Technology*, 44(10), 2053–2058.
- [51] Gu, C., Xiang, B., Su, Y., & Xu, J. (2013). Near-infrared spectroscopy coupled with kernel partial least squares-discriminant analysis for rapid screening water containing malathion. *American Journal of Analytical Chemistry*, 4(3), 111–116. doi: 10.4236/ajac.2013.43015.
- [52] Rosipal, R., & Trejo, L. J. (2001). Kernel partial least squares regression in reproducing kernel Hilbert space. *Journal of Machine Learning Research*, 2(Dec), 97–123.
- [53] Wang, K., Chi, G., Lau, R., & Chen, T. (2011). Multivariate calibration of near infrared spectroscopy in the presence of light scattering effect: a comparative study. *Analytical Letters*, 44(5), 824–836.
- [54] Hsu, K. L., Gupta, H. V., & Sorooshian, S. (1995). Artificial neural network modeling of the rainfall-runoff process. *Water Resources Research*, 31(10), 2517–2530.
- [55] Borin, A., Ferrao, M. F., Mello, C., Maretto, D. A., & Poppi, R. J. (2006). Least-squares support vector machines and near infrared spectroscopy for quantification of common adulterants in powdered milk. *Analytica Chimica Acta*, 579(1), 25–32.
- [56] Zomer, S., Brereton, R. G., Carter, J. F., & Eckers, C. (2004). Support vector machines for the discrimination of analytical chemical data: application to the determination of tablet production by pyrolysis-gas chromatography-mass spectrometry. *Analyst*, 129(2), 175–181.
- [57] Brudzewski, K., Osowski, S., & Markiewicz, T. (2004). Classification of milk by means of an electronic nose and SVM neural network. *Sensors and Actuators B: Chemical*, 98(2), 291–298.
- [58] Fernandez Pierna, J. A., Volery, P., Besson, R., Baeten, V., & Dardenne, P. (2005). Classification of modified starches by Fourier transform infrared spectroscopy using support vector machines. *Journal of Agricultural and Food Chemistry*, 53(17), 6581–6585.
- [59] Van Gestel, T., De Brabanter, J., De Moor, B., Vandewalle, J., Suykens, J. A. K., & Van Gestel, T. (2002). *Least Squares Support Vector Machines* World Scientific, Singapore, 2002.
- [60] Suykens, J. A., & Vandewalle, J. (1999). Least squares support vector machine classifiers. *Neural Processing Letters*, 9(3), 293–300.
- [61] Shao, Y., & He, Y. (2013). Visible/Near infrared spectroscopy and chemometrics for the prediction of trace element (Fe and Zn) levels in rice leaf. *Sensors*, 13(2), 1872–1883.

- [62] Xu, L., Yan, S. M., Cai, C. B., & Yu, X. P. (2014). Nondestructive discrimination of lead (Pb) in preserved eggs (pidan) by near-infrared spectroscopy and chemometrics. *Journal of Spectroscopy*, 2014.
- [63] Barbosa, R. M., Batista, B. L., Varrique, R. M., Coelho, V. A., Campiglia, A. D., & Barbosa, F. (2014). The use of advanced chemometric techniques and trace element levels for controlling the authenticity of organic coffee. *Food Research International*, 61, 246–251.
- [64] Huang, G., Han, L., Yang, Z., & Wang, X. (2008). Evaluation of the nutrient metal content in Chinese animal manure compost using near infrared spectroscopy (NIRS). *Bioresource Technology*, 99(17), 8164–8169.

IntechOpen