We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Lipoproteins and Diseases of the Brain

Kwang-Min Kim and G. Tayhas R. Palmore

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/67053>

Abstract

Apolipoprotein E4 (apoE4) and outer surface protein A (ospA) are pathogenic lipoproteins involved in the progression of Alzheimer's disease and Lyme neuroborreliosis, respectively. Results from previous studies indicate that apoE4 exhibits neurotoxicity by activating amyloid beta pathways, and ospA causes damage to the brain by stimulating immune activity of microglia and astrocytes. These results, however, lack information about the specific interactions that develop between neurons and these two lipoproteins. It is essential to investigate the effect of these lipoproteins on neuronal morphology and function to better understand the mechanism of damage and disease of the brain. This chapter summarizes previous studies on the role of apoE4 and ospA in diseases of the brain and discusses experimental results from our own work that suggests new roles for apoE4 and ospA in neuronal outgrowth and synaptic loss.

Keywords: apolipoprotein E4, bacterial outer surface protein A, neurodegeneration, neuroinflammation, nerve regeneration, synaptic loss

1. Introduction

Lipoproteins in the brain are involved in the onset and progression of neurodegenerative diseases (e.g., Alzheimer's disease) [1, 2] and neuroinflammatory disorders (e.g., neuroborreliosis) [3, 4]. These lipoproteins are either endogenously expressed by astrocytes [5] and microglia [6, 7] or exogenously produced by bacterial pathogens (e.g., *Borrelia burgdorferi*, *Streptococcus pneumoniae*) [8].

The most abundant endogenous lipoproteins in the brain include apolipoprotein E (apoE) and apoJ [2]. These endogenous lipoproteins mediate transport of lipids between various cells in the brain to maintain and regulate the brain structure and function [9, 10]. The apoE isoform, apoE4, has been investigated intensively because previous studies showed that lipidation of

apoE4 (i.e., apoE4 carrying cholesterol and phospholipids) is the major risk factor indicative of the onset of Alzheimer's disease (AD) [11].

The exogenous lipoprotein most studied in the brain is the bacterial outer surface protein A (ospA), which is produced by *B. burgdorferi* [12, 13]. *B. burgdorferi* causes Lyme disease, which is the most common tick-borne infection in Europe and in North America [14]. A recent study using rats infected with *B. burgdorferi* demonstrated that *B. burgdorferi* was observed across the blood-brain barrier (BBB) and that the expression level of ospA was augmented significantly in the brain [4].

Thus, apoE and ospA have been of interest to both scientists and clinicians who seek to develop new strategies for treatment of brain injuries and brain disorders induced by these pathogenic lipoproteins. It still remains unclear however, if apoE and ospA interact directly with neurons to disrupt the structure and function of the brain, whereas it is documented extensively that these lipoproteins induce pathological states via amyloid beta (A β) aggregation [15, 16] and immune activation of microglia and astrocytes [17, 18]. To address the absence of direct evidence of interaction between lipoproteins and neurons, we have studied the effect of apoE4 and ospA on neurons in terms of axonal outgrowth and synaptic loss. This chapter discusses these findings and the potential new roles of apoE4 and ospA in the context of previous studies on these lipoproteins in neurodegeneration and neuroinflammation.

2. ApoE4 and neuronal outgrowth

2.1. Lipidation of apoE isoforms

ApoE transports and clears lipids from one cell to another to maintain lipid homeostasis of the brain [9, 10]. To carry lipids (e.g., cholesterol, phospholipids, and lipoproteins), apoE is lipidated (i.e., lipid-bound apoE) by adenosine triphosphate (ATP)-binding cassette A1 (ABCA1) transporters on astrocytes [19] (**Figure 1a**). The lipidation status of apoE depends on its three isoforms (i.e., apoE2, apoE3, and apoE4) coded by three alleles (ϵ 2, ϵ 3, and ϵ 4 of *APOE* gene) on chromosome 19 [20]. The three isoforms of apoE differ from one another by amino acid interchanges at two residue sites (**Table 1**).

These minor variations cause a change in the structure and function of apoE, which eventually leads to distinct disease mechanisms in AD [21]. ApoE4 has an arginine at residue 112 that connects the N terminus (Arg 61) to the C terminus (Glu 255) to form a folded structure of apoE called *domain interaction* [22]. ApoE2 and apoE3 have a cysteine at residue 112, which is less likely to create the folded structure of domain interaction. The presence of domain interaction results in distinct lipid-bound forms among apoE isoforms. ApoE4 binds preferentially to larger lipid particles due to its folded structure, which interferes with internalization of lipids into neurons [11]. In contrast, apoE2 and E3 bind to various sizes of lipids in more ways that are efficient and thus facilitate lipid transport between cells in the brain. The lipidated apoE can be internalized into cells in the brain (i.e., astrocytes, microglia, and neurons) through the family of low-density lipoprotein receptors (LDLR), low-density lipoprotein receptor-related protein 1 (LRP1), or heparan sulfate proteoglycans (HSPGs) [11, 16].

Figure 1. The interaction between apoE and neurons illustrated. (a) ApoE transports lipids to neurons by forming lipopeptide particles (i.e., lipidation of apoE). ApoE is lipidated by ATP-binding cassette A1 (ABCA1) transporters of astrocytes. The lipid-binding affinity of apoE4 is different from that of apoE2 and apoE3 because of structural differences in its domain interaction. Both lipid and apoE can bind to Aβ monomers and oligomers. The Aβ-lipidated apoE2/3 complex can be internalized by LDLR, LRP1, or HSPG, which clears Aβ. The efficiency of internalizing large lipid-bound apoE4 into cells is low, which increases the probability of Aβ plaque formation because of poor Aβ clearance. (b) ApoE alone can bind to Aβ monomers and oligomers regardless of its isoforms. The Aβ-nonlipidated apoE complex increases the probability of forming Aβ plaques because nonlipidated apoE cannot be internalized via LDLR or LRP1. (c) Nonlipidated apoE4 enhances neuronal adhesion, axon outgrowth, and neurite branching. The receptor in neurons that regulates growth-enhancing effects of nonlipidated apoE4 remains unknown. Abbreviations: ABCA1, ATP-binding cassette A1 transporter; LDLR, low-density lipoprotein receptor; LRP1, low-density lipoprotein receptor-related protein 1; HSPG, heparan sulfate proteoglycans.

ApoE isoforms	ApoE amino acid residue	
	112	158
ApoE2	Cys	Cys
ApoE3	Cys	Arg
ApoE4	Arg	Arg

Table 1. Differences of apoE isoforms in amino acid residues.

When the lipidated apoE is internalized into cells, Aβ monomers and oligomers are also cleared because they bind to both lipids and apoE at residues 12–28 [23]. Thus, Aβ clearance is dependent on the structural difference of apoE isoforms, and this mechanism helps to prevent

the A β aggregation that is associated with the progression of AD. The A β -bound apoE, however, forms aggregates regardless of the isoform of apoE when they are not lipidated and thus, are not internalized [16, 23] (**Figure 1b**).

2.2. Nonlipidated apo E4 and neuronal outgrowth

When lipidated, apoE4 is known to be toxic to neurons through various pathogenic pathways such as A β aggregation and apoE fragment formation [21]. The effect of apoE4 on neurons when it is not lipidated, however, remains unclear. To address this knowledge gap, the effect of apoE4 on neuronal outgrowth was studied *in vitro* without lipids in the medium [24]. This study compared neuronal responses to various culture substrates including glass, laminin-coated glass, and apolipoprotein E4-coated glass by quantifying key neuronal outgrowth parameters in terms of cell adhesion, axon length, number of neurites, and number of branches on axons. The results of this study demonstrated that apoE4 not only enhances neuronal adhesion but also significantly increases axon outgrowth and branching when compared to laminin, a protein that is recognized as one of the best extracellular matrix (ECM) proteins for enhancing neuronal growth [25]. As such, results from this study contradict the prevailing view that apoE4 has only a degenerative effect on cells in the brain. Although apoE4 when lipidated predominantly exhibits neurotoxicity when studied *in vivo* and in clinical models, it should be considered that both lipidated and nonlipidated apoE in these models constantly interact with neurons to mediate brain activity. Thus, the results from this study provide a complementary mechanism of action of apoE. In addition, the neuron-growing potential of apoE4 can be applied to transplantable therapeutic systems using stem cells or microstructure devices prior to interaction of lipids *in vivo*.

The mechanism by which nonlipidated apoE mediates axon outgrowth and branching remains elusive, whereas lipidated apoE is known to interact with cells via LRP1, LDLR, or HSPG [16]. It has been reported that apoE does not bind to LDLR or LRP1 without lipidation [26, 27]. Integrin and HSPGs also were tested for their involvement in apoE4-induced axon outgrowth by inhibiting these receptors. Neither of these receptors was found to be responsible for apoE4-induced neuronal outgrowth (**Figure 1c**). The mechanism of interaction between neurons and nonlipidated apoE4 is the subject of ongoing studies.

3. Bacterial lipoprotein and synaptic loss

3.1. Bacterial lipoproteins and neuroinflammation

Bacterial surface components including lipoproteins and lipopolysaccharide (LPS) have been reported to be elevated in the cerebrospinal fluid (CSF) of patients suffering from a bacterial infection such as bacterial meningitis [28]. These components can cause neuropsychiatric manifestations such as lymphocytic meningitis, cranial and peripheral neuropathy, and cerebral infarcts [29, 30]. When compared to LPS, bacterial lipoproteins activate inflammatory pathways more vigorously [31], leading to more severe damage to tissue [32]. Bacterial lipoproteins still remain in the tissue even after the degradation of bacteria by antibiotic

therapies [33, 34]. As a result, many studies suggest that minimizing the production of bacterial proteins or inhibiting bacterial protein synthesis is more effective at preventing neural injury from bacterial infections in animal models or patients [35, 36] than simply using antibiotics to kill bacteria. Bacterial lipoproteins in the brain trigger microglia activation via the toll-like receptors (TLRs) to produce inflammatory mediators (e.g., cytokines and reactive oxygen species) [37–39] and induce migration of immune cells across the BBB [40, 41]. The result is damaged brain tissue including cell death of astrocytes, oligodendrocytes, and neurons [42, 43].

The outer surface protein (osp) is the most studied bacterial lipoprotein that includes ospA, ospB, and ospC from *B. burgdorferi* [12, 44, 45]. Three palmitoyl groups (i.e., the lipid portion) at the N-terminus of the peptide is responsible for immune activation and tissue injury [46, 47], whereas the peptide portion of ospA is not effective at activating immune pathways [32]. Thus, tripalmitoyl-S-glyceryl-cysteine (Pam3-Cys), a synthetic lipopeptide mimicking the N-termini of osp, is often used for studying bacterial infection in a wide range of research fields involving immunology and neuroscience [48, 49]. Although all of ospA, ospB, and ospC share common immune pathways (e.g., NF- κ B activation) via TLR2, ospA shows higher toxicity to tissues when compared with ospB and ospC [32]. The reason for distinct toxic effects among these different lipoproteins continues to remain elusive.

3.2. OspA and presynaptic loss

OspA from *B. burgdorferi* is able to cross the BBB by binding to CD40 of brain-microvascular endothelial cells [4]. OspA in the brain activates TLR2 on microglia and astrocytes, which initiates immune activity and causes damage to brain tissue [14, 50]. However, information regarding the interaction between ospA and neurons is lacking because the expression level of TLR2 in neurons is extremely low when compared with that of microglia or astrocytes. Thus, the interaction between ospA and neurons has been overlooked [51, 52]. To address this question, the effect of ospA on neurons has been investigated with a specific focus on synaptic loss. The density and transmission of synapses are considered to be the key parameters in determining the functional state of brain tissue (e.g., information processing and storage) because neurons transmit electrical and biochemical signals to adjacent neurons through the synapse. The signal-sending synapse (i.e., presynapse) is located on the axon and the signal-receiving synapse (i.e., postsynapse) is located on the dendrite of a neuron. If neurons lose one of these synapses or have misaligned synapses, the brain cannot function properly even when neurons survive from brain injuries or diseases. Thus, the change in pre- and postsynaptic density was quantified following treatment of cultures of rat E18 hippocampal neurons with ospA (2 μ M) for 24 h (**Figure 2**). The quantification of synaptic density was determined by counting the number of synaptic sites (i.e., synapsin or postsynaptic density protein 95 (PSD-95)) in a randomly selected secondary dendrite. OspA expressed from *Escherichia coli* (prepared by the Biomaterials and Advanced Drug Delivery Laboratory at Stanford University) showed that ospA significantly decreased the number of presynaptic sites (i.e., synapsin) ($p = 0.04$), whereas it did not affect the number of postsynaptic sites (i.e., PSD-95) ($p > 0.05$) (**Figure 2**). This result suggests that ospA directly disrupts neuronal function by damaging presynapses exclusively.

Figure 2. OspA and synaptic density. (a) Fluorescent images showing rat primary hippocampal neurons treated with vehicle (endotoxin-free water). (b) Fluorescent images showing rat primary hippocampal neurons treated with ospA (2 μ M). (a) and (b) Neurons stained with anti-PSD-95 (postsynaptic protein), anti-synapsin (presynaptic protein), merge of PSD-95 and synapsin, merge of PSD-95, synapsin, and MAP2 (dendrite) from left to right are shown. (c) The postsynaptic density was measured by the number of postsynaptic sites (puncta) per length of selected dendrite. The postsynaptic density was not affected by ospA ($P > 0.05$). (d) The presynaptic density was measured by the number of presynaptic sites (puncta) per length of selected dendrite. The presynaptic density decreased significantly by ospA ($P = 0.041$).

3.3. Mechanism of synaptic dysfunction induced by bacterial lipoproteins

A recent study demonstrated that viral infection leads to cognitive dysfunction by microglial engulfment of presynapses via the complement C3 pathway [53]. Another recent study showed that viral infection impairs synaptic function via glycogen synthase kinase 3 (GSK-3) activation and intracellular accumulation of A β [54]. Thus, an increasing number of studies are being reported that elucidate the mechanism underlying synaptic dysfunction induced by viral infection. Although there is evidence that bacterial lipoprotein ospA also damages presynapses (**Figure 2**), information as to how bacterial infection impairs synaptic function is lacking. Three possible mechanisms may account for synaptic dysfunction during bacterial infection. First, bacterial lipoproteins damage synapses via activation of inflammatory pathways (e.g., TLR2 and TLR4) as discussed in Section 3.1. Second, bacterial lipoproteins damage synapses through neurotransmitter-mediated excitotoxicity. It has been demonstrated that the level of quinolinic acid, the N-methyl-D-aspartate (NMDA) receptor agonist, was elevated significantly in the CSF of Lyme neuroborreliosis patients [55]. The

NMDA receptor mediates synaptic transmission, plasticity, and excitotoxicity in the central nervous system (CNS) and it exhibits excitotoxic effects when an excessive flux of calcium occurs by the increase of a neurotransmitter such as glutamate [56]. However, it is yet to be determined whether the presence of bacterial lipoproteins directly mediates the elevation of quinolinic acid. Third, bacterial lipoproteins damage synapses through physical interaction with synapses independent of biochemical pathways (i.e., inflammation and receptor activation). It has been suggested that the physical properties of proteins (e.g., aggregate pattern and size) is a crucial determinant in mediating pathogenic toxicity [57, 58]. This toxicity occurs independent of their sequences or lengths [59] in a manner that is similar to the aggregation of A β in Alzheimer's disease [60] or α -synuclein in Parkinson's disease [61]. Previous studies showed that Pam3-Cys, the synthetic N-terminus of ospA, self-assembled and showed aggregating potential *in vitro* assays [58, 62], which can be related to brain tissue damage including the disruption of synaptic function.

4. Conclusions

This chapter describes the new roles of apoE4 and ospA as major pathogenic endogenous and exogenous lipoproteins, respectively, in neuronal outgrowth and function by discussing recent experimental data in the context of previous reports. Recent studies show that apoE4 enhances neuronal adhesion and axonal outgrowth *in vitro* when it acts alone without lipids. New studies also demonstrate the possibility that ospA can induce synaptic dysfunction by damaging exclusively presynaptic sites. These results contribute to a new understanding of how lipoproteins are involved in developing neuropathology by interacting with neurons. Future studies should focus on the specific mechanism of interaction between apoE4 and neurons and the effect of ospA on synaptic function using *in vivo* models. Along with many pathogenic pathways governed by various cell types in the brain (e.g., microglia, astrocytes, and oligodendrocytes), the effect of pathogenic factors on neuronal activity provides a deeper understanding of structural and functional abnormality in neurodegeneration and neuroinflammation [63]. Understanding the interaction between lipoproteins and neurons in the brain should yield new approaches to the treatment of brain injuries and brain disorders.

Author details

Kwang-Min Kim¹ and G. Tayhas R. Palmore^{2*}

*Address all correspondence to: tayhas_palmore@brown.edu

¹ Department of Neurosurgery, Stanford University, Stanford, CA, USA

² School of Engineering, Brown University, Providence, RI, USA

References

- [1] Urosevic N, Martins RN. Infection and Alzheimer's disease: the APOE epsilon 4 connection and lipid metabolism. *J Alzheimers Dis.* 2008 May;13(4):421–35. PubMed PMID: WOS:000256230100007. English.
- [2] Wang H, Eckel RH. What are lipoproteins doing in the brain?. *Trends Endocrin Met.* 2014 Jan;25(1):8–14. PubMed PMID: WOS:000330606100005. English. DOI:10.1016/j.tem.2013.10.003
- [3] Latov N, Wu AT, Chin RL, Sander HW, Alaedini A, Brannagan TH. Neuropathy and cognitive impairment following vaccination with the OspA protein of *Borrelia burgdorferi*. *J Peripher Nerv Syst.* 2004 Sep;9(3):165–7. PubMed PMID: WOS:000224002700007. English. DOI:10.1111/j.1085-9489.2004.09306.x
- [4] Pulzova L, Kovac A, Mucha R, Mlynarcik P, Bencurova E, Madar M, et al. OspA-CD40 dyad: ligand-receptor interaction in the translocation of neuroinvasive *Borrelia* across the blood-brain barrier. *Sci Rep.* 2011 Sep 8; 1: 86. PubMed PMID: WOS:000296053800002. English. DOI:10.1038/Srep00086
- [5] Mahley RW. Apolipoprotein-E - cholesterol transport protein with expanding role in cell biology. *Science.* 1988 Apr 29;240(4852):622–30. PubMed PMID: WOS:A1988N126500025. English. DOI:10.1126/science.3283935
- [6] Uchihara T, Duyckaerts C, He Y, Kobayashi K, Seilhean D, Amouyel P, et al. Apoe immunoreactivity and microglial cells in Alzheimers-disease brain. *Neurosci Lett.* 1995 Jul 28;195(1):5–8. PubMed PMID: WOS:A1995RM14300002. English. DOI:10.1016/0304-3940(95)11763-M
- [7] Hoffmann O, Braun JS, Becker D, Halle A, Freyer D, Dagand E, et al. TLR2 mediates neuroinflammation and neuronal damage. *J Immunol.* 2007 May 15;178(10):6476–81. PubMed PMID: WOS:000246286200055. English.
- [8] Kovacs-Simon A, Titball RW, Michell SL. Lipoproteins of bacterial pathogens. *Infect Immun.* 2011 Feb;79(2):548–61. PubMed PMID: WOS:000286462000001. English. DOI:10.1128/iai.00682-10
- [9] Mahley RW, Rall SC. Apolipoprotein E: far more than a lipid transport protein. *Annu Rev Genom Hum G.* 2000;1:507–37. PubMed PMID: WOS:000165768900019. English. DOI:10.1146/annurev.genom.1.1.507
- [10] Raber J, Wong D, Yu GQ, Buttini M, Mahley RW, Pitas RE, et al. Alzheimer's disease—apolipoprotein E and cognitive performance. *Nature.* 2000 Mar 23;404(6776):352–4. PubMed PMID: WOS:000086119000038. English. DOI:10.1038/35006165
- [11] Bu GJ. Apolipoprotein E and its receptors in Alzheimer's disease: pathways, pathogenesis and therapy. *Nat Rev Neurosci.* 2009 May;10(5):333–44. PubMed PMID: WOS:000265365300010. English. DOI:10.1038/nrn2620

- [12] Barbour AG, Tessier SL, Todd WJ. Lyme-disease spirochetes and ixodid tick spirochetes share a common surface antigenic determinant defined by a monoclonal-antibody. *Infect Immun.* 1983;41(2):795–804. PubMed PMID: WOS:A1983RA43900050. English.
- [13] Burgdorfer W, Barbour AG, Hayes SF, Benach JL, Grunwaldt E, Davis JP. Lyme-disease—a tick-borne spirochetosis. *Science.* 1982;216(4552):1317–9. PubMed PMID: WOS:A1982NS80100029. English. DOI:10.1126/science.7043737
- [14] Pfister HW, Wilske B, Weber K. Lyme borreliosis—basic science and clinical aspects. *Lancet.* 1994 Apr 23;343(8904):1013–6. PubMed PMID: WOS:A1994NG68200013. English. DOI:10.1016/S0140-6736(94)90130-9
- [15] Ittner LM, Gotz J. Amyloid-beta and tau—a toxic pas de deux in Alzheimer’s disease. *Nat Rev Neurosci.* 2011 Feb;12(2):67–72. PubMed PMID: WOS:000286420100001. English. DOI:10.1038/nrn2967
- [16] Liu CC, Kanekiyo T, Xu HX, Bu GJ. Apolipoprotein E and Alzheimer disease: risk, mechanisms and therapy. *Nat Rev Neurol.* 2013 Feb;9(2):106–18. PubMed PMID: WOS:000315250400010. English. DOI:10.1038/nrneurol.2012.263
- [17] Lien E, Sellati TJ, Yoshimura A, Flo TH, Rawadi G, Finberg RW, et al. Toll-like receptor 2 functions as a pattern recognition receptor for diverse bacterial products. *J Biol Chem.* 1999 Nov 19;274(47):33419–25. PubMed PMID: WOS:000083745200039. English. DOI:10.1074/jbc.274.47.33419
- [18] Koch S, Donarski N, Goetze K, Kreckel M, Stuerenburg HJ, Buhmann C, et al. Characterization of four lipoprotein classes in human cerebrospinal fluid. *J Lipid Res.* 2001 Jul;42(7):1143–51. PubMed PMID: WOS:000169784300018. English.
- [19] Fitz NF, Cronican AA, Saleem M, Fauq AH, Chapman R, Lefterov I, et al. Abca1 deficiency affects Alzheimer’s disease-like phenotype in human ApoE4 but not in ApoE3-targeted replacement mice. *J Neurosci.* 2012 Sep 19;32(38):13125–36. PubMed PMID: WOS:000309258100016. English. DOI:10.1523/Jneurosci.1937-12.2012
- [20] Farrer LA, Cupples LA, Haines JL, Hyman B, Kukull WA, Mayeux R, et al. Effects of age, sex, and ethnicity on the association between apolipoprotein E genotype and Alzheimer disease—a meta-analysis. *Jama-J Am Med Assoc.* 1997 Oct 22;278(16):1349–56. PubMed PMID: WOS:A1997YA90500037. English. DOI:10.1001/jama.278.16.1349
- [21] Mahley RW, Huang YD. Apolipoprotein E sets the stage: response to injury triggers neuropathology. *Neuron.* 2012 Dec 6;76(5):871–85. PubMed PMID: WOS:000312283400005. English. DOI:10.1016/j.neuron.2012.11.020 (2012).
- [22] Dong LM, Wilson C, Wardell MR, Simmons T, Mahley RW, Weisgraber KH, et al. Human apolipoprotein E—role of arginine-61 in mediating the lipoprotein preferences of the E3-isoform and E4-isoform. *J Biol Chem.* 1994 Sep 2;269(35):22358–65. PubMed PMID: WOS:A1994PE09800065. English.
- [23] Sadowski MJ, Pankiewicz J, Scholtzova H, Mehta PD, Prelli F, Quartermain D, et al. Blocking the apolipoprotein E/amyloid-beta interaction as a potential therapeutic

- approach for Alzheimer's disease. *Proc Natl Acad Sci USA*. 2006 Dec 5;103(49):18787–92. PubMed PMID: WOS:000242689800072. English. DOI:10.1073/pnas.0604011103
- [24] Kim KM, Vicenty J, Palmore GTR. The potential of apolipoprotein E4 to act as a substrate for primary cultures of hippocampal neurons. *Biomaterials*. 2013 Apr;34(11):2694–700. PubMed PMID: WOS:000315748200012. English. DOI:10.1016/j.biomaterials.2013.01.012
- [25] LuckenbillEdds L. Laminin and the mechanism of neuronal outgrowth. *Brain Res Rev*. 1997 Feb;23(1–2):1–27. PubMed PMID: WOS:A1997WN18500001. English. DOI:10.1016/S0165-0173(96)00013-6
- [26] Wilson C, Wardell MR, Weisgraber KH, Mahley RW, Agard DA. Three-dimensional structure of the LDL receptor-binding domain of human apolipoprotein E. *Science*. 1991 Jun 28;252(5014):1817–22. PubMed PMID: WOS:A1991FU06100031. English. DOI:10.1126/science.2063194
- [27] Blacklow SC. Versatility in ligand recognition by LDL receptor family proteins: advances and frontiers. *Curr Opin Struc Biol*. 2007 Aug;17(4):419–26. PubMed PMID: WOS:000250423800006. English. DOI:10.1016/j.sbi.2007.08.017
- [28] Goos M, Lange P, Hanisch UK, Prinz M, Scheffel J, Bergmann R, et al. Fibronectin is elevated in the cerebrospinal fluid of patients suffering from bacterial meningitis and enhances inflammation caused by bacterial products in primary mouse microglial cell cultures. *J Neurochem*. 2007 Sep;102(6):2049–60. PubMed PMID: WOS:000249452000031. English. DOI:10.1111/j.1471-4159.2007.04683.x
- [29] Veenendaalhilbers JA, Perquin WVM, Hoogland PH, Doornbos L. Basal meningo-vasculitis and occlusion of the basilar artery in two cases of *Borrelia burgdorferi* infection. *Neurology*. 1988 Aug;38(8):1317–9. PubMed PMID: WOS:A1988P586300030. English.
- [30] Pfister HW, Rupprecht TA. Clinical aspects of neuroborreliosis and post-Lyme disease syndrome in adult patients. *Int J Med Microbiol*. 2006 May;296:11–6. PubMed PMID: WOS:000238550200003. English. DOI:10.1016/j.ijmm.2005.12.003
- [31] Hirschfeld M, Kirschning CJ, Schwandner R, Wesche H, Weis JH, Wooten RM, et al. Cutting edge: inflammatory signaling by *Borrelia burgdorferi* lipoproteins is mediated by toll-like receptor 2. *J Immunol*. 1999 Sep 1;163(5):2382–6. PubMed PMID: WOS:000082125400007. English.
- [32] Batsford S, Dunn J, Mihatsch M. Outer surface lipoproteins of *Borrelia burgdorferi* vary in their ability to induce experimental joint injury. *Arthritis Rheum*. 2004 Jul;50(7):2360–9. PubMed PMID: WOS:000222820300038. English. DOI:10.1002/art.20337
- [33] Bockenstedt LK, Gonzalez DG, Haberman AM, Belperron AA. Spirochete antigens persist near cartilage after murine Lyme borreliosis therapy. *J Clin Invest*. 2012 Jul;122(7):2652–60. PubMed PMID: WOS:000306044600035. English. DOI:10.1172/JCI58813

- [34] Evans ME, Pollack M. Effect of antibiotic class and concentration on the release of lipopolysaccharide from *Escherichia coli*. J Infect Dis. 1993 Jun;167(6):1336–43. PubMed PMID: WOS:A1993LD69400011. English.
- [35] Nau R, Wellmer A, Soto A, Koch K, Schneider O, Schmidt H, et al. Rifampin reduces early mortality in experimental *Streptococcus pneumoniae* meningitis. J Infect Dis. 1999 Jun;179(6):1557–60. PubMed PMID: WOS:000080561100034. English. DOI:10.1086/314760
- [36] Bottcher T, Gerber J, Wellmer A, Smirnov AV, Fakhrijanali F, Mix E, et al. Rifampin reduces production of reactive oxygen species of cerebrospinal fluid phagocytes and hippocampal neuronal apoptosis in experimental *Streptococcus pneumoniae* meningitis. J Infect Dis. 2000 Jun;181(6):2095–8. PubMed PMID: WOS:000087923900038. English. DOI:10.1086/315518
- [37] Akira S, Takeda K. Toll-like receptor signalling. Nat Rev Immunol. 2004 Jul;4(7):499–511. PubMed PMID: WOS:000222435700012. English. DOI:10.1038/nri1391
- [38] Moresco EMY, LaVine D, Beutler B. Toll-like receptors. Curr Biol. 2011 Jul 12;21(13):R488–93. PubMed PMID: WOS:000292805400004. English.
- [39] O'Neill LAJ, Golenbock D, Bowie AG. The history of Toll-like receptors—redefining innate immunity. Nat Rev Immunol. 2013 Jun;13(6):453–60. PubMed PMID: WOS:000319445300012. English. DOI:10.1038/nri3446
- [40] Ek M, Engblom D, Saha S, Blomqvist A, Jakobsson PJ, Ericsson-Dahlstrand A. Inflammatory response—pathway across the blood-brain barrier. Nature. 2001 Mar 22;410(6827):430–1. PubMed PMID: WOS:000167583800030. English. DOI: 10.1038/35068632
- [41] Ramesh G, MacLean AG, Philipp MT. Cytokines and chemokines at the crossroads of neuroinflammation, neurodegeneration, and neuropathic pain. Mediat Inflamm. 2013; 2013:480739. PubMed PMID: WOS:000323444900001. English. DOI: 10.1155/2013/480739
- [42] Parthasarathy G, Fevrier HB, Philipp MT. Non-viable *Borrelia burgdorferi* induce inflammatory mediators and apoptosis in human oligodendrocytes. Neurosci Lett. 2013 Nov 27;556:200–3. PubMed PMID: WOS:000328520900040. English. Doi:10.1016/j.neulet.2013.10.032
- [43] Ramesh G, Alvarez AL, Roberts ED, Dennis VA, Lasater BL, Alvarez X, et al. Pathogenesis of Lyme neuroborreliosis: *Borrelia burgdorferi* lipoproteins induce both proliferation and apoptosis in rhesus monkey astrocytes. Eur J Immunol. 2003 Sep;33(9):2539–50. PubMed PMID: WOS:000185323200020. English. DOI:10.1002/eji.200323872
- [44] Barbour AG, Tessier SL, Hayes SF. Variation in a major surface protein of Lyme-disease spirochetes. Infect Immun. 1984;45(1):94–100. PubMed PMID: WOS:A1984SX63000017. English.
- [45] Livey I, Gibbs CP, Schuster R, Dorner F. Evidence for lateral transfer and recombination in *OspC* variation in Lyme-disease *Borrelia*. Mol Microbiol. 1995 Oct;18(2):257–

69. PubMed PMID: WOS:A1995TG43300008. English. DOI:10.1111/j.1365-2958.1995.mmi_18020257.x
- [46] Erdile LF, Brandt MA, Warakowski DJ, Westrack GJ, Sadziene A, Barbour AG, et al. Role of attached lipid in immunogenicity of *Borrelia burgdorferi* OspA. *Infect Immun*. 1993 Jan;61(1):81–90. PubMed PMID: WOS:A1993KF63600011. English.
- [47] Weis JJ, Ma Y, Erdile LF. Biological-activities of native and recombinant *Borrelia burgdorferi* outer surface protein A: dependence on lipid modification. *Infect Immun*. 1994 Oct;62(10):4632–6. PubMed PMID: WOS:A1994PH29800073. English.
- [48] Aliprantis AO, Yang RB, Mark MR, Suggett S, Devaux B, Radolf JD, et al. Cell activation and apoptosis by bacterial lipoproteins through toll-like receptor-2. *Science*. 1999 Jul 30;285(5428):736–9. PubMed PMID: WOS:000081765100055. English. DOI:10.1126/science.285.5428.736
- [49] Du XN, Fleiss B, Li HF, D'angelo B, Sun YY, Zhu CL, et al. Systemic stimulation of TLR2 impairs neonatal mouse brain development. *PLoS One*. 2011 May 6;6(5): e19583. PubMed PMID: WOS:000290305600031. English. DOI:10.1371/journal.pone.0019583
- [50] Saijo K, Glass CK. Microglial cell origin and phenotypes in health and disease. *Nat Rev Immunol*. 2011 Nov;11(11):775–87. PubMed PMID: WOS:000296584700015. English. DOI:10.1038/nri3086
- [51] Tang SC, Arumugam TV, Xu XR, Cheng AW, Mughal MR, Jo DG, et al. Pivotal role for neuronal Toll-like receptors in ischemic brain injury and functional deficits. *Proc Natl Acad Sci USA*. 2007 Aug 21;104(34):13798–803. PubMed PMID: WOS:000249064700050. English. DOI:10.1073/pnas.0702553104
- [52] Downer EJ, Johnston DGW, Lynch MA. Differential role of Dok1 and Dok2 in TLR2-induced inflammatory signaling in glia. *Mol Cell Neurosci*. 2013 Sep;56:148–58. PubMed PMID: WOS:000325834100015. English. DOI:10.1016/j.mcn.2013.04.007
- [53] Vasek MJ, Garber C, Dorsey D, Durrant DM, Bollman B, Soung A, et al. A complement-microglial axis drives synapse loss during virus-induced memory impairment. *Nature*. 2016 Jun 23;534(7608):538–43. PubMed PMID: WOS:000378270300051. English. DOI:10.1038/nature18283
- [54] Piacentini R, Li Puma DD, Ripoli C, Marrocci ME, De Chiara G, Garaci E, et al. Herpes Simplex Virus type-1 infection induces synaptic dysfunction in cultured cortical neurons via GSK-3 activation and intraneuronal amyloid-beta protein accumulation. *Sci Rep*. 2015 Oct 21;5:15444. PubMed PMID: WOS:000363122200004. English. DOI:10.1038/Srep15444
- [55] Halperin JJ, Heyes MP. Neuroactive kynurenines in Lyme borreliosis. *Neurology*. 1992 Jan;42(1):43–50. PubMed PMID: WOS:A1992HA89500007. English.
- [56] Cull-Candy S, Brickley S, Farrant M. NMDA receptor subunits: diversity, development and disease. *Curr Opin Neurobiol*. 2001 Jun;11(3):327–35. PubMed PMID: WOS:000169286200009. English. DOI:10.1016/S0959-4388(00)00215-4

- [57] Bucciantini M, Giannoni E, Chiti F, Baroni F, Formigli L, Zurdo JS, et al. Inherent toxicity of aggregates implies a common mechanism for protein misfolding diseases. *Nature*. 2002 Apr 4;416(6880):507–11. PubMed PMID: WOS:000174756500035. English. DOI:10.1038/416507a
- [58] Schromm AB, Howe J, Ulmer AJ, Wiesmuller KH, Seyberth T, Jung G, et al. Physicochemical and biological analysis of synthetic bacterial lipopeptides—validity of the concept of endotoxic conformation. *J Biol Chem*. 2007 Apr 13;282(15):11030–7. PubMed PMID: WOS:000245941500023. English. DOI:10.1074/jbc.M700287200
- [59] Pastor MT, Kummerer N, Schubert V, Esteras-Chopo A, Dotti CG, de la Paz ML, et al. Amyloid toxicity is independent of polypeptide sequence, length and chirality. *J Mol Biol*. 2008 Jan 18;375(3):695–707. PubMed PMID: WOS:000253124600010. English. DOI:10.1016/j.jmb.2007.08.012
- [60] Ono K, Condrón MM, Teplow DB. Structure-neurotoxicity relationships of amyloid beta-protein oligomers. *Proc Natl Acad Sci USA*. 2009 Sep 1;106(35):14745–50. PubMed PMID: WOS:000269481000005. English. DOI:10.1073/pnas.0905127106
- [61] Conway KA, Lee SJ, Rochet JC, Ding TT, Williamson RE, Lansbury PT. Acceleration of oligomerization, not fibrillization, is a shared property of both alpha-synuclein mutations linked to early-onset Parkinson's disease: implications for pathogenesis and therapy. *Proc Natl Acad Sci USA*. 2000 Jan 18;97(2):571–6. PubMed PMID: WOS:000084873100012. English. DOI:10.1073/pnas.97.2.571
- [62] Hamley IW, Kirkham S, Dehsorkhi A, Castelletto V, Reza M, Ruokolainen J. Toll-like receptor agonist lipopeptides self-assemble into distinct nanostructures. *Chem Commun*. 2014;50(100):15948–51. PubMed PMID: WOS:000345452400028. English. DOI:10.1039/c4cc07511k
- [63] Xanthos DN, Sandkuhler J. Neurogenic neuroinflammation: inflammatory CNS reactions in response to neuronal activity. *Nat Rev Neurosci*. 2014 Jan;15(1):43–53. PubMed PMID: WOS:000329022000010. English. DOI:10.1038/nrn3617

IntechOpen