We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Promoting Patient-Centered Care in Chronic Disease

Ana Monteiro Grilo,
Margarida Custódio dos Santos,
Ana Isabel Gomes and Joana Santos Rita

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/67380>

Abstract

Nowadays, many people around the world are seeing their lives being shattered and even shortened due to one or more chronic conditions. Chronic illness is a dynamic ongoing process that is determined by a complexity of factors. Patient literacy, motivation, emotional well-being, and resources play an important role on patient adaption and are important challenges for healthcare providers. A systematic patient-centered approach that enables chronic patients to play an active role in their condition management and in the decision-making process on a day-to-day basis is required. However, some studies show that health professionals do not always guide their actions by Patient-centered orientation, either by personal issues or by professional and/or institutional barriers. The present chapter aimed to provide a comprehensive approach to patient-centered care in chronic disease and offer a structured guideline as a tool for formal academic education in chronic patient-centered care. This chapter is structured in five sections: (1) Chronic disease: the challenge of the twenty-first century, (2) The patient with a chronic disease, (3) Patient-centered care in chronic diseases, (4) Issues and barriers to achieve patient-centered care, and (5) Guide tool for health professionals' training and education in patient-centered care.

Keywords: patient care, chronic disease, patient empowerment, medical students, health professionals

1. Introduction

Patient-centered care is defined as the professional's attitude during healthcare that is closely congruent with a responsive to patient's wants, needs, and preferences [1]. Although patients differ in their preferences and interaction styles, the patient-centered style has been associated with higher rates of satisfaction, better adherence, and better treatment,

particularly in the management of chronic illness [2]. Effective chronic disease management involves continued and coordinated action by the patient and the physician: dynamic partnership with mutual responsibilities and accountabilities.

Nevertheless, some studies conclude that health professionals do not always guide their actions by patient-centered orientation, either by personal issues (e.g., personality characteristics, development of specific communication skills) or by professional and/or institutional barriers [3]. Training health professionals, either during the academic years or in postgraduate courses, can promote the change that is required to accomplish patient centeredness.

The present chapter aimed to provide a comprehensive approach to patient-centered care in chronic disease and offer a structured guideline as a tool for health professionals' training and education in patient-centered care. It will be structured in five sections: (1) Chronic disease: the challenge of the twenty-first century, (2) The patient with a chronic disease, (3) Patient-centered care in chronic diseases, (4) Issues and barriers to achieve patient-centered care, and (5) Guide tool for health professionals' training and education in patient-centered care.

2. Chronic diseases: the challenge of the twenty-first century

The World Health Organization (WHO) defines chronic illnesses as diseases of long duration and generally slow progression. These diseases can usually be controlled but not cured and include persistent and repeated health problems, gradual changes over time, and an asynchronous evolution [4]. Many people around the world are seeing their lives being shattered and even shortened due to one or more chronic conditions [5]. Noncommunicable diseases such as diabetes mellitus, heart disease, obesity, chronic respiratory diseases, and cancer are among the most common [6, 7]. These diseases are the leading cause of death among adults, as well as mortality rates due to them in middle-aged people are higher in some high-income countries [8]. In 2012, about half of adult population had one or more chronic disease, and one of four adults had two or more chronic conditions [9]. Chronic diseases are major causes for disability, including chronic pain, motor and sensory dysfunction, blindness, lower limb amputation, and impaired functioning. In addition, chronic conditions are not only the principal cause for the increased costs in the healthcare system but also for the social costs, like absenteeism at work and decreased productivity [7]. Because of an aging population and the adoption of unhealthy lifestyles, chronic diseases are estimated to grow even further in the next decades. Therefore, it is not surprising that WHO considered chronic conditions the healthcare challenge of this century.

The principal causes of chronic diseases have been identified and well established. Age and hereditary are the only non-modifiable risk factors; in almost all developed countries, most common causes are modifiable, as unhealthy diet and excessive energy intake, tobacco use, physical inactivity, and alcohol abuse [8].

Chronic disease can impact quality of life and daily activities and should require ongoing actions on a long-term basis [10] (**Table 1**). Often, people living with chronic disease face significant life

Dimensions	Examples
Personal	Age, gender
Sensorial	Pain, fatigue, incontinence, joint swelling
Emotional	Anxiety, depression, anger, distress
Cognitive	Negative thoughts, dealing with uncertainty about future, request for information on disease and treatments
Behavioral	Difficulties in activities of daily life, problems with mobility and balance, sleep problems, functional restrictions, medication management
Social	Discrimination, adjustment or end of career, changes in personal relationships
Self/identity	Changes in self-image and self-esteem
Healthcare	More contact with health professionals

Table 1. Experience of chronic disease (adapted from Ref. [10]).

changes and deal with constant threats to their personal health [11]. Most of them must manage daily symptoms, special diets, physical exercise plans, adherence to medication regimens, and systematic monitoring to identify new problems. Each person is affected in a different way by chronic disease; experiences diverge according to personal (e.g., coping strategies), contextual (e.g., access to health systems), and physical (e.g., symptoms) contexts [10].

As indicated in **Table 1**, patients with chronic disease face significant stresses in managing their illness [11] that go far beyond the functional impairments. Therefore, the major goal for healthcare system, and particularly for health professionals, is to address the needs of chronic patients and their care [12].

Nowadays, with effective behavior change efforts and adherence to medication regimens, chronic diseases and their consequences can often be better managed [8, 13]. However, the traditional paternalistic model of care focused on the control of acute symptoms shows itself inadequate to meet the challenge of chronic disease and particularly the needs of chronic patients [14]. A systematic patient-centered approach that enables chronic patients to play an active role in their condition management and in the decision-making process on a day-to-day basis is required [8].

3. The patient with a chronic disease

Patient adaptation to illness has been found to be associated with patient engagement to treatment [15]. Despite differences among patients, studies revealed that the majority of patients and families wish to be more active and involved in treatment decisions and procedures [16]. This involvement, the amount of information, and the development of competencies that usually came along with it may indeed help patients to succeed dealing with disease and treatment’s challenges.

Currently, chronic patients (and their families) are, mostly, their own primary healthcare provider. Although they receive support and counselling from health professionals, most patients live the major part of their lives outside of health institutions, they work and have their families. Regarding their illness, patients are expected to motorize their symptoms, to make decisions about adherence behaviors, often to follow complex treatment regimens, to make decisions about when they need to seek professional care, and to make necessary life changes to reach or maintain better health outcomes. Effective self-management is, then, considered critical to meet the needs of people living with long-term conditions. Self-management has been defined as “the care taken by individuals towards their own health and well-being; it comprises the actions they take to lead a healthy lifestyle; to meet their social, emotional and psychological needs; to care for their long-term condition; and to prevent further illness or accidents” [17].

Some factors related to the patient have been repeatedly highlighted to influence the way patient's cope with this challenges.

3.1. Patient literacy

Health literacy respects to the ability to read, understand, and act upon health information and has been associated with the skills for making appropriate health decisions [18]. Receiving clear information and education increases patient understanding of the disease implications. On the other hand, information about treatment procedures enhances patient's auto-efficacy and self-confidence.

Three distinct levels or purposes of health literacy have been identified: *functional*, basic skills in reading and writing necessary for effective functioning in a health context; *interactive*, more advanced cognitive literacy and social skills that enable active participation in healthcare; and *critical*, the ability to critically analyze and use information to participate in actions that overcome structural barriers to health [16].

A systematic review by Berkman and colleagues to determine whether low health literacy is related to the poorer use of healthcare, outcomes, costs, and disparities in health outcomes among persons of all ages found that low literacy was associated with poorer health outcomes and the poorer use of healthcare services [19]. Association between good level of health literacy and treatment outcomes, patient satisfaction, and better quality of care was also found in recent studies [20, 21].

3.2. Patient willingness and motivation

Studies show that, in general, patients wish some degree of participation on their care [22, 23]. Nevertheless, despite the evident benefits, not all patients feel motivated to participate in their care, to adhere to medical recommendations, or to change lifestyles or health behaviors that are key stones for their treatment. Hibbard and colleagues, in a study with adults aged 45–97 years old with chronic diseases, found that only 22% adopted more involved and active behaviors to participate in their treatment, 12% do not think that they must play an active role in their own health believing that they are just recipients of healthcare, 29% do not have enough information or understanding about their clinical situation and their treatment, and 37% have the understanding but not the confidence to be more involved [24].

In a qualitative study that aimed to explore patients' and practitioners' views on factors influencing engagement in self-management in the context of multimorbidity, with a sample drawn from four general practices in Greater Manchester, motivation emerged as one of the three themes that captured patients' and professionals' views. In this study, motivation drew on understandings that successful self-management was partly contingent on patients' belief and expectation that self-management would improve their health and how low mood can negatively influence patients' capacity and sense of responsibility for self-management [25].

Regarding adherence, in a study focused in the relation between physiotherapist's attitudes and patient adherence, Chan and colleagues found that patient's treatment motivation mediated the relationship between physiotherapist's autonomy supportive behaviors and rehabilitation adherence [26]. Increasing patient motivation through motivational strategies and motivational interviewing can be used to reduce resistance and improve the odds of achieving positive clinical outcomes among noncompliant/resistant patients [27]. Highlighting the importance of motivation on patient adherence [28], a meta-analysis of systematic reviews about the effectiveness of adherence interventions published between 1990 and 2005 found that interventions focused on behavioral interventions that used incentives and patient motivation were among the most effective to promote adherence.

3.3. Patient emotional disorders

Depression is frequently associated with chronic conditions and, if untreated, can adversely affect the course of the disease and limit effective treatment for the chronic condition [29]. For instance, depression was found in 27% of diabetic population compared with 17% in the non-diabetic population. Those patients with diabetes and depression experienced an impact with a large effect size on quality of life, as compared with those who suffered diabetes and who were not depressed [30]. In another study, depression coexisting with diabetes was associated with poorer glycemic control, increased risk of complications, increased healthcare utilization and costs, increased functional disability, and lost work productivity [31]. There is also some evidence that depression affects self-management. Patients with depressive symptoms (including subclinical depression) were found much less likely to gain in activation and in their self-management behaviors [32].

3.4. Patient resources

Social support refers to the degree to which interpersonal relationships correspond to certain functions of material support, affective, emotional, informative, and positive social interaction [33] and is a multifaceted experience that involves voluntary associations and formal and informal relationships with others [34]. Social support can buffer the negative impact of life events on health and positively influence psychosocial adjustment and self-management of chronic illness [35]. A large amount of studies found associations between social support and clinical and self-management/self-care behaviors [36], health-related quality of life [37, 38], and patient adherence to treatment [39]. Patient's network for social support includes family and friends, social environment, and all health providers [40].

In conclusion, adaptation to chronic illness is a dynamic ongoing process that is determined by a complexity of factors. Evidence has shown that patients are decision-makers that may actively contribute to their own care. Patient literacy, motivation, emotional well-being, and resources play an important role on patient adaption and are important challenges for healthcare providers. These challenges are better attended within the framework of the patient-centered model of care.

4. Patient-centered care in chronic diseases

Chronic disease management demands the adaptation of the patient-centered care model to the changing patients' needs. Hudon and colleagues developed a thematic analysis based on Stewart's¹ model of patient-centered care, based on articles published between 1980 and 2009. They found six major themes: (a) *Starting from the patient's situation*, (b) *Legitimizing the illness experience*, (c) *Acknowledging the patient's expertise*, (d) *Developing an ongoing partnership*, (e) *Offering realistic hope*, and (f) *Providing patients' advocacy in the healthcare system* [2].

4.1. Starting from the patient situation

Chronic patient-centered care requires an extended comprehension of the patient's situation. This theme included two subthemes corresponding to Stewart et al. dimensions *exploring both the disease and the illness* and *understanding the whole person*. *Exploring both the disease and the illness* means to discover the patient's personal and subjective experience of sickness: worries, feelings, expectations, previous experience of care, health behaviors, and confidence with chronic conditions management, with the same interested as the biological dimensions like physical evaluation and laboratory results.

Understanding the whole person, as the act of being aware of the surrounding circumstances of patient's life, implies the acknowledgment that chronic health meanings could vary with adjacent environments. Stewart and colleagues (2003) recognize that take into account various aspects of the patient's personal and life context allowed health professional's comprehension to the several factors that influence patient's complex dynamic responses to chronic condition [41]. Contextual factors could be proximal and distal to the patient [41]. Proximal factors refer to immediate and specific categories and comprise family, financial security, education, employment, leisure, and social support. Distal factors are related with general and meta-context categories and include community, culture, healthcare system, sociohistorical period, geography, and the media.

4.2. Legitimizing the illness experience

Chronic patient-centered care includes attending to patient's needs, worries, and emotions as grief over their previous skills and allows them to express their concerns related to their

¹Stewart et al. (2006) model included six components: exploring both the disease and the illness experience, understanding the whole person, finding common ground, incorporating prevention and health promotion, enhancing the patient-doctor relationship, and being realistic.

chronic conditions. By recognizing subjective perceptions and sensations of illness, health professionals formally acknowledge the reality of the patient's experiences and allow the patient to feel that his definition of reality is confirmed [42]. Legitimation helped patient in his/her illness adaptation process and improve patient-health professional relationship. Lehman and colleagues show that patients with chronic fatigue syndrome report more psychological distress and dissatisfaction when physician failed to legitimize their illness [43]. This theme is one of the most important, when improving centered care is a priority.

4.3. Acknowledging patient expertise on his/her own life

One of the central premises of Carl Rogers' patient-centered model is the idea that each patient is the world best expert on himself [44]. Patient's expertise should be acknowledged and respected by the caregiver. In chronic disease, great management of the conditions depends on patient's actions; therefore, all the interactions with health professionals should centered on patients' strengths and challenges on running their disease condition. The development of self-management plans should integrate not only the health professional expertise but also the patient's concerns, priorities, and resources. Usually, patients appreciate being treated as a full member of the health team, as this promotes their sense of control and empowerment [10]. This requires a new empowerment paradigm, with redefinition of patient and health professional roles [45].

4.4. Developing an ongoing partnership

Effective chronic disease management entails an ongoing partnership between patient and health professionals. Partnership refers to both a relationship and a process [10, 35] and promotes patient empowering to take control over his/her situation. Therefore, patients can be involved in their care as an active partner, helping to clarify symptoms and diagnosis, sharing decisions and treatments responsibilities, discussing treatment outcomes and alternatives, and agreeing on a treatment plan. This theme corresponds to two subthemes on Stewart and colleagues model: enhancement of the physician-patient relationship and finding common ground [41].

Mutual respect, trust, and collaborative action are essential ingredients to build and *enhance the patient-physician relationship*. Both patient and health professional need to feel committed to sharing power and responsibility [10]. In chronic disease, physical conditions and patients' preferences and capacities change over time and determine the nature of partnership in which moment. On the other hand, continuity of care is an opportunity to improve the patient-health professional relationship, to empower the patient and to support during the most difficult phases of chronic condition. According to this author, an effective partnership depends on four factors: patient (e.g., stage of adjustment to chronic illness, self-esteem), adjunctive factors (e.g., family support, health service resources), communication process (patient and health professional verbal and no verbal skills), and health professional (e.g., attitudes related to own role, self-awareness).

Finding a common ground is the process by which health professional and patient range a mutual understanding and agreement related to chronic disease (the problem), goals and priorities in management and/or treatment of chronic disease, and roles to be assumed both by patient and health professional [41]. Patient needs, preferences, and beliefs should be respected always.

4.5. Offering realistic hope

This theme emphasizes on hope and support in the context of uncertainty or the inevitable deterioration of chronic conditions and contrasts from Stewart and colleagues (2006) *being realistic* issue. Hope has been identified as a key attribute required for patient adaption to illness and disability [46]. Giving hope tempered with realism allows patients to deal with uncertainty and promotes patient well-being and adaptation. Hope does not suppress negative aspects of the illness but includes and focuses on the positive, less negative, and possible developments that may be considered better outcomes or better patient quality of life [47].

4.6. Providing patients' advocacy in the healthcare system

This theme is new considering Stewart's model and relates to the health professional's role in guiding the patient through the healthcare system. That means defending their rights, interests and safety. Often, chronic conditions are complex and patients need different levels of care (e.g., physiotherapy), support groups, and community services.

4.7. Over time

The six themes previously described comprise a longitudinal dimension. Patient's situation could change depending on the disease course, life context, and illness experience. On the other hand, patients' expectations, knowledge, and expertise in self-management of his/her chronic conditions could improve (or not) over time and influence patient-health professional relationship. As the disease progresses, values, goals, and preferences must be reassessed and discussed. Even health professional advocacy role should be adjusted to patient desires at a particular time.

5. Issues and barriers to achieve patient-centered care

The WHO (2005) recognizes the potential benefits of chronic patient-centered care but also identifies many challenges when applying this approach [16]. In fact, even when health professionals recognize patient-centered approach as an essential feature in clinical practice, they report difficulties in maintaining these behaviors during care. Personal and professional barriers need to be considered to carry on this challenge.

5.1. Organizational barriers

Time constraints, staff shortage, and work overload are frequently mentioned as major difficulties for a patient-centered care by health professionals [48]. In a study developed by West and colleagues, nurses acknowledge that lack of time frequently compromises their ability to provide clear and complete information to patients and their families, to give effective emotional support, and to offer the appropriate treatment according to patient's clinical needs and safety [49]. The limited time available for medical appointment is also referred by doctors [50, 51] and patients [52] as a barrier to provide information about diagnosis and to discuss

treatment options. The impossibility to control physical environment features like temperature or noise can reduce the possibility of health professionals to comply with patient's specific needs during recovery [49]. In addition, a private space that guarantees patient's confidentiality and dignity during medical procedures is a condition difficult to offer in health-care institutions [49].

5.2. Interpersonal barriers

The lack of a common acknowledgment about the relevance of patient-centered practices in healthcare among professionals' team is mentioned as an important barrier [53]. The health team relationships and the conditions given to the staff to effectively consider patients' perspectives appear to facilitate a more generalized patient-centered attitude. When clinical and personal information about the patient is shared by the health team, the health professionals have major opportunities to make stronger and more consistent decisions about the best interventions toward the patient [54]. A prior and clear definition of the clinical practice role of each health professional regarding patients' care and communication seems to be very useful to overcome difficulties in applying correctly a patient-centered attitude [55].

5.3. Individual barriers

Several studies have highlighted specific health professionals' beliefs and feelings as possible barriers to a more patient-centered attitude. The fear of leading with patient's emotional distress has been identified as an important obstacle for the adoption of relational models that approach nurses from their patients [49, 56]. Nurses feel frequently that the emotional detachment from their work and their patients is essential to keep going emotionally and mentally healthy and to prevent stress and burnout [57].

The professionals' perception of patient roles and responsibilities can also affect their intervention during the medical procedures. In a study presented by Jallinoja and colleagues, nurses and physicians identify patients' willingness to make changes in their lifestyles in a chronic disease context as a barrier to treatment [58]. Pessimistic judgments or unrealistic expectations about patients' abilities to make effective lifestyle changes can compromise medical actions based in promoting empowerment. In a systematic review about barriers to the adoption of a shared decision-making process by physicians, doctors perceived inability to balance patients' preferences with the clinical guidelines available, and they are afraid that the implementation of a shared decision-making process leads to patients' discomfort or threatens health professionals' autonomy. Moreover, the perception that shared decision-making practices does not contribute significantly to better patient's outcomes is identified as a possible difficulty [48]. Lack of motivation and difficulties in following a holistic approach are another two important problems referred by nurses

The gap between the "ideal" academically taught and the "real" clinical practice can lead to the withdrawal of patient-centered attitudes and to the adoption of more traditional paternalistic models [59, 60]. During the academic internship and among recently graduated health professionals is evident some inconstancy in orientation assumed toward patients.

That is students and professionals range from focusing exclusively in accomplish procedures and tasks correctly, or accent primary in patients' needs and worries [51, 61, 62]. The newness of the experience endorses lack of confidence and uncertainly, when students have to decide about the right action to implement in specific situations during healthcare [63], and gives more strength to students' preconceptions and stereotypes about patients [62]. In this period, senior colleagues who have responsibilities in tutoring students in their first interactions with patients play an important role as models [64]. However, not always those first clinical experiences are properly supervised [65] nor patient-centered care orientation is defended and practiced by tutors [51, 60, 66]. In this context, students face problems in really translating patient-centered approach into nursing care, because poor or ineffective teaching strategies were used to training patient-centered care skills [67, 68].

5.4. Providing healthcare to specific chronic patient groups

Difficulties in effective communication and patient-centered attitudes seem to exacerbate when healthcare is provided to certain patient groups, namely, patients with intellectual disabilities [69], critically ill patients [70], patients with severe communication impairment [71], elderly [63, 72] and pediatrics patients [73], and cancer patients [55].

6. Guide tool for health professionals' training and education in patient-centered care

Patient-centered care requires competences in communication skills. Nowadays, many medical schools recognize the impact of communication skills and use guidelines to teach and assess these competences during undergraduate medical training [74, 75]. The recent European consensus on learning objectives for a core communication curriculum in healthcare professions [47] states as key tasks: (a) build relationship and develop trust and rapport; (b) elicit information from the patient/problems and concerns; (c) consider the patient's perspective of the illness; (d) give relevant information and explanations; (e) develop shared plan of care; and (f) close the interview and set up next meeting.

Considering the effectiveness of several earlier guidelines about communication skills training in healthcare professional's education, namely, the Calgary Cambridge [76], the Four Habits Model [77], Kalamazoo Consensus Framework [75], REDE Model Skills Checklist [74], and the Health Professionals Core Communication Curriculum [47], we propose a tool aimed to systematize major dimensions of patient-centered care in chronic illness as defined by Hudon and colleagues into their specific core conditions and respective communication skills, with examples of interactions with the patient (**Table 2**).

Regarding methods for effective training and education of health professionals' students, active learning and reflective learning seem to be the most effective teaching strategies [75]. At the start, core conditions and communication skills have to be explained [76], and evidence of their utility in chronic patient-centered care should be provided. In formal

1. Starting from the patient situation

Core conditions	Gather information: (a) encourage patients to tell story of problems in own words; (b) explore both the disease and the illness means (scan how each problem affects the patient's life); examine patient beliefs, concerns, and expectations. Enhance patient to talk and express feelings.
Skills	Use open-ended questions to initiate patient narrative. Identify patient's problems or subjects that patients desire to discuss with appropriate open questions . Use VIEW questions [74]: Vital activities: <i>How does chemotherapy treatment impact on daily routines?</i> Ideas: <i>What do you think of talking to our psychologist about the emotional impact of chemotherapy treatment?</i> Expectations: <i>You are going to finish your chemotherapy treatment today. What are your expectations about it?</i> Worries: <i>What worries you most about chemotherapy treatment side effects?</i> Listen attentively without interruption, facilitating patient's responses verbally (e.g., <i>yes; ah, ah; go on; mm-hmm; I see</i>) and nonverbal (e.g., eye contact, facial expression) facilitation. Attend to nonverbal cues , checking out body language and facial expression (e.g., <i>This seems to be a difficult problem of you...</i>). Clarify statements that are unclear or need amplification, with more specify or yes/no questions (e.g., <i>Could you explain what you mean by having two bad days?</i>). Summarize patient narrative to verify understanding of what the patient has said (e.g., <i>So, you feel very tired and you think that your tiredness is not related with the intense work period you have been involved</i>), and give patient opportunity to correct or add information.

2. Legitimizing the illness experience

Core conditions	Recognize emotional cues and clarify the emotion (if necessary). Validate patient's views and feelings, avoiding judgments.
Skills	Demonstrate empathy , showing interest and care in the patient experience. Communicate understanding and appreciation of patient's feeling or situation, using SAVE [74]. Support: <i>We are going to work together to find better solution to control your pain.</i> Acknowledge: <i>The experience you describe during your back pain crisis sound overwhelming.</i> Validate: <i>It's very common for people with persistent pain feel exasperated and discouraged like you do.</i> Emotion naming: <i>You seem more discouraged today than our last appointment.</i> Use no verbal cues (tone, eye contact, and posture) that show care and concern.

3. Acknowledging patient expertise on their own lives

Core conditions	Recognize patient's attributes and expertise. Reinforce patient's skills and strengths. Determine how much the patient desire to be included in choices and decisions and clarify your own role as a supporter and a trustful information provider. Negotiate a mutually acceptable plan.
Skills	Use open-ended questions to encourage patients to talk about illness, their experience, and their sources of information (e.g., <i>What do you think are the most difficult task of your treatment? Where did you look for information on your illness?</i>). Listen attentively , allowing patients to complete what they have to say without interruptions. Use reflection , to acknowledge what has worked with the patient (e.g., <i>I see you already exercised three times a day. It is very important for your diabetes to exercise</i>) and/or his/her resources and strengths (<i>I realize you are very motivated to change your diet and start eating more health food</i>). Support patients, using clarification questions that help them to rephrase information that is unclear or needs amplification (e.g., <i>You are saying that having to take that pill worries you. Can you explain a little more?</i>). Express the importance of patient expertise by incorporating patient's opinions, values, and needs in treatment plans (e.g., <i>Based on your concerns, I think that we should change our initial plan. What is your opinion?</i>).

Use **negotiation**, showing respect about patient’s needs, expectations, and objectives (*What do you expect from this treatment?*), asking questions and promoting answers that may clarify the object of negotiation (*Can you explain what you meant by having difficulties in think about this diet?*), providing information that allows the discussion of different alternatives, understanding patient limitations (e.g., the resources that are need to implement a specific treatment), highlighting shared points of view and showing pleasure in agreement (*I am very glad that we agree on several issues about your diet plan*), and arranging plans that recurrently reassess decisions and help to adjust them if needed.

4. Developing an ongoing partnership

Core conditions	Enhancing the patient-physician relationship	<p>Reveal interest in the patient as a person.</p> <p>Develop trust and rapport, accepting legitimacy of patient’s views and feeling.</p> <p>Provide support adjusted to the stage of chronic illness.</p>
	Finding common ground	<p>Focus on working collaboratively to meet patient goals.</p> <p>Provide reliable, clear, evidence-based, and individualized information that promotes sharing decision and responsibilities.</p> <p>Use adequate strategies to clarify patient’s doubts and to support patient’s decisions.</p> <p>Promote the sense of “joining together” with patients in supportive partnerships collaboration to meet patient’s goals.</p> <p>Promote patient’s sharing information and educate patients on how to reach information and how to motorize treatment outcomes.</p> <p>Develop and use systems to record, communicate, and monitor the implementation of patient’s preferences.</p>
Skills	Enhancing the patient-physician relationship	<p>Attend to patient’s privacy, confidentiality, and autonomy.</p> <p>Acknowledge difficult situations and challenges during the chronic illness stages, providing continued, willing, helpful, gentle, supportive care.</p> <p>Deal kindly with embarrassing and disturbing issues.</p> <p>Avoid interrupting and respect the silences.</p> <p>Avoid direct advice (<i>You should...</i>), impositions (<i>You have to...</i>), or over, not realistic tranquilization (<i>I’m sure you are going to be fine.</i>).</p> <p>Consider patient’s degree of understanding and language, avoiding jargons.</p> <p>Use BUSTER [78]. Be prepared: define a plan to manage for specific patient’s emotions and practice self-regulation. Use nonjudgmental listening. Six second rule: avoid escalation of conversation. “Tell me more” statements: <i>Can you tell me more about that worry that you just mentioned now?</i> Empathize and validate: <i>It isn’t easy to talk about your illness, is it?</i> Respond with a wish statement: <i>I wish this treatment had been more effective</i></p>
	Finding common ground	<p>Provide information, tailoring information to what the patient wants and/or is able to know, using words that show that you prioritize patient information needs, thru categories (e.g. <i>Let’s talk about medication side effects</i>) and “7C”: clear, concise, concrete, correct, coherent, complete, and courteous [79].</p> <p>Check if the patients would like more information; if necessary, provide writing or audio-taping information. Be sure that your information promotes patient’s understanding and skills for decision taking.</p> <p>Make suggestions, rather than given directives (e.g., <i>I would like to make a suggestion about your diet</i>) and encourage patient to make choices (e.g., <i>Let’s talk about your alternatives. I think you have three options here...</i>).</p>

Check understanding by asking patients **feedback** or **teach back**, asking to, in their own words, tell what you have just explained. Be careful and try not to sound like you are testing your patient (e.g., *We have been talking about your diet. What do you think is the most important information so far, as for instance? Could you explain it back to me? I really want to know if I was clear enough.*)

Encourage questions, stating, e.g., *Your questions are very important for me and for your process*, and use **open-end questions**, eliciting patient's preference and views in relation to treatment options (e.g., *For you what are the main benefits of this treatment? What about surgery?*).

Reinforce patients for achieving small successes encouraging them to raise expectations about self-efficacy and enhancing involvement in partnership (e.g., *We did so well. Your effort was compensated. I know we can do even more.*).

Provide **education on patient treatment** by **modeling** and/or delivering **written information and instructions** that promote patient feeling of self-efficacy and encourage patients to build their own capacity for self-management.

5. Offering realistic hope

Core conditions	Being responsive to patients' concerns emotions and suffering. Being honest and trustful, providing clear update and realistic information. Having a positive attitude toward the diagnosis, the treatment, and the care. Including patient education and capacitation to decrease uncertainty.
Skills	Promote patients' control toward the treatment and their own quality of life by providing clear and realistic information that allows patients to reduce uncertainty and to be aware of different alternatives and new developments. Promote patients' positive thinking by using positive talk .

Table 2. Guidelines for effective communication with chronic health patients based on patient-centered care principles as defined by Hudon and colleagues [2].

academic education, active patient participation can be achieved either by using actor-simulated patients (sometimes called standardized patients) or by engaging real patient to share their own experiences [80, 81]. Videos can also be used. One good example is the Database of Individual Patient Experience of Illness (DIPEX) [82]. These educational sessions should be an ongoing task in clinical education and must be carefully plan to give space to trainees' reflection. Participants should be able to discuss the skills used and the impact of the trained skills on the patient on health professional and should have the opportunity to obtain feedback on their performance [83].

Moreover, learning and training communication skills should be an ongoing process during academic and professional education and meet the progressively more difficult demands that are placed on students and professionals. Thus, for example, basic communication skills such as welcoming the patient or developing empathy should be the focus of training in the early years. On the other hand, issues such as giving bad news or dealing with emotionally disturbing situations should be integrated into the more advanced years of training.

At the end of the training, participants should be able to use a checklist that helps them to verify if all dimensions of chronic patient-centered care model have been considered during their interaction with patient (Table 3).

	Yes/no	Observation
I understand the patient's perspective of his/her situation		
I validated (with empathy) patient's experience and feelings		
I asked and understand what are the patient's main concerns		
I provided information I want to give and that answered patient's needs		
I check that I have responded adequately to the patient's demands		
I encourage patient's skills and strengths		
We shared decision in the level patient's feel comfortable		
I offer realistic hope considering patient situation		

Table 3. Checklist for health professionals' self-evaluation regarding communication and interaction with patients, based on patient-centered care principles as defined by Hudon and colleagues [2].

7. Conclusions

In this chapter, we have described some of the challenges motivated by chronic disease management demands and outlined some core principles of patient-centered care in chronic disease. Promoting a chronic patient-centered approach is likely depending on giving primacy to the subjective aspects of illness, therefore is not a simple matter and must be included in all levels of medical education. Two important next steps concerning the development of chronic patient-centered model must be considered. First, training skills programs must be integrated in preclinical courses and not only in courses of communication as usually. Second, principles of chronic patient-centered model should be applied to a wide variety of chronic diseases, making them tangibly significant to concrete clinical conditions.

Author details

Ana Monteiro Grilo^{1*}, Margarida Custódio dos Santos^{1,2}, Ana Isabel Gomes³ and Joana Santos Rita⁴

*Address all correspondence to: ana.grilo@estesl.ipl.pt

1 Higher School of Health Technology, Polytechnic Institute of Lisbon, Lisbon, Portugal

2 Faculty of Psychology, University of Lisbon, Lisbon, Portugal

3 Lusíadas Lisbon Hospital, Lisbon, Portugal

4 Lisbon Regional Health Administration, Odivelas Health Care Centers, Lisbon, Portugal

References

- [1] Laine C, Davidoff F. Patient-centered medicine: a professional evolution. *JAMA*. 1996;275(2):152-6.
- [2] Hudon C, Fortin M, Haggerty J, Loignon C, Lambert M, Poitras ME. Patient-centered care in chronic disease management: a thematic analysis of the literature in family medicine. *Patient Education and Counseling*. 2012;88(2):170-6.
- [3] McCabe C. Nurse–patient communication: an exploration of patients’ experiences. *Journal of Clinical Nursing*. 2004;13(1):41-9.
- [4] Bentzen, N, ed. *Wonca Dictionary of General/Family Practice*. Wonca International Classification Committee: Copenhagen, 2003.
- [5] Wu S, Green A. *Projection of chronic illness prevalence and cost inflation*. Santa Monica, CA: RAND Health; 2000.
- [6] Martin C. Chronic disease and illness care. *Canadian Family Physician*. 2007;53:2086-91.
- [7] Mendis S, Armstrong T, Bettcher D, Branca F, Lauer J, Mace C, et al. *GLOBAL STATUS REPORT on noncommunicable diseases 2014*. Geneva: World Health Organization; 2015.
- [8] WHO. *Preventing chronic diseases: a vital investment*. WHO Global Report. Geneva: WHO; 2005.
- [9] Ward B, Schiller J, Goodman R. Multiple chronic conditions among US adults: a 2012 update. *Preventing Chronic Disease*. 2014;11:130389.
- [10] Reynolds F. *Communication and clinical effectiveness in rehabilitation*. Edinburgh: Elsevier Butterworth Heinemann; 2005.
- [11] Boyd CM, Wolff JL, Giovannetti E, Reider L, Weiss C, Xue QL, et al. Health care task difficulty among older adults with multimorbidity. *Medical Care*. 2014;52(3):S118-25.
- [12] Trehearne B, Fishman P, Lin E. Role of the nurse in chronic illness management: making the medical home more effective. *Nursing Economics*. 2014;32(4):178-84.
- [13] McMullen CS, Safford M, Bosworth H, Phansalkar S, Leong A, Fagan M, et al. Patient-centered priorities for improving medication management and adherence. *Patient Education and Counseling*. 2015;98:102-10.
- [14] Brock C, Dost A. *Self-care—a real choice: self-care support—a practical option*. London: Department of Health; 2005.
- [15] Hibbard JH, Greene J. What the evidence shows about patient activation: better health outcomes and care experiences; fewer data on costs. *Health Affairs*. 2013;32(2):207-14.

- [16] Coulter A, Parsons S, Askham J. Policy brief: Where are the patients in decision-making about their own care. WHO 2008 and WHO on behalf of the European Observatory on Health Systems and Policies 2008. Denmark: World Health Organisation; 2008.
- [17] Clark N, Becker M, Janz N, Lorig K, Rawkowski W, Anderson L. Self-management of chronic disease by older adults: a review and questions for research. *Journal of Aging and Health*. 1991;3:3-27.
- [18] Nielsen-Bohlman L, Panzer A, Kindig D. Health literacy: a prescription to end confusion. Institute of Medicine of National Academics. Washington, D.C.: National Academies Press; 2004.
- [19] Berkman ND, Sheridan SL, Donahue KE, Halpern DJ, Crotty K. Low health literacy and health outcomes: an updated systematic review. *Annals of Internal Medicine*. 2011;155(2):97-107.
- [20] Altin SV, Stock S. The impact of health literacy, patient-centered communication and shared decision-making on patients' satisfaction with care received in German primary care practices. *BMC Health Services Research*. 2016;16(1):450-60.
- [21] Fisher CA, Myers M. Patient education: a win-win opportunity. *Nursing Management*. 2016;47(3):18-20.
- [22] Foot C, Gilbert H, Dunn P, Jabbal J, Seale B, Goodrich J, et al. People in control of their own health and care: the state of involvement. London: The Kings Fund; 2014.
- [23] Vahdat S, Hamzehgardeshi L, Hessam S, Hamzehgardeshi Z. Patient involvement in health care decision making: a review. *Iranian Red Crescent Medical Journal*. 2014; 16(1):e12454.
- [24] Hibbard JH, Mahoney ER, Stockard J, Tusler M. Development and testing of a short form of the patient activation measure. *Health services research*. 2005;40(6p1):1918-30.
- [25] Coventry PA, Fisher L, Kenning C, Bee P, Bower P. Capacity, responsibility, and motivation: a critical qualitative evaluation of patient and practitioner views about barriers to self-management in people with multimorbidity. *BMC Health Services Research*. 2014;14(1):536-48.
- [26] Chan DK, Lonsdale C, Ho PY, Yung PS, Chan KM. Patient motivation and adherence to postsurgery rehabilitation exercise recommendations: the influence of physiotherapists' autonomy-supportive behaviors. *Archives of Physical Medicine and Rehabilitation*. 2009;90(12):1977-82.
- [27] Butterworth SW. Influencing patient adherence to treatment guidelines. *Journal of Managed Care & Specialty Pharmacy*. 2008;14(6 Supp B):21-5.
- [28] van Dulmen S, Sluijs E, van Dijk L, de Ridder D, Heerdink R, Bensing J. Patient adherence to medical treatment: a review of reviews. *BMC Health Services Research*. 2007;7(1):1-13.

- [29] Chapman DP, Perry GS, Strine TW. The vital link between chronic disease and depressive disorders. *Preventing Chronic Diseases*. 2005;2(1):1-10.
- [30] Goldney RD, Phillips PJ, Fisher LJ, Wilson DH. Diabetes, depression, and quality of life a population study. *Diabetes Care*. 2004;27(5):1066-70.
- [31] Egede LE. Effect of comorbid chronic diseases on prevalence and odds of depression in adults with diabetes. *Psychosomatic Medicine*. 2005;67(1):46-51.
- [32] Hibbard JH, Mahoney ER, Stock R, Tusler M. Do increases in patient activation result in improved self-management behaviors? *Health Services Research*. 2007;42(4):1443-63.
- [33] Griep RH, Chor D, Faerstein E, Werneck GL, Lopes CS. Social support: scale test-retest reliability in the Pro-Health Study. *Cadernos de Saúde Pública*. 2003;19(2):625-34.
- [34] Bardach SH, Tarasenko YN, Schoenberg NE. The role of social support in multiple morbidity: self-management among rural residents. *Journal of Health Care for the Poor and Underserved*. 2011;22(3):759-71.
- [35] Gallant MP. The influence of social support on chronic illness self-management: a review and directions for research. *Health Education & Behavior*. 2003;30(2):170-95.
- [36] Strom JL, Egede LE. The impact of social support on outcomes in adult patients with type 2 diabetes: a systematic review. *Current Diabetes Reports*. 2012;12(6):769-81.
- [37] Colloca G, Colloca P. The effects of social support on health-related quality of life of patients with metastatic prostate cancer. *Journal of Cancer Education*. 2016;31(2):244-52.
- [38] Degroote S, Vogelaers D, Vandijck DM. What determines health-related quality of life among people living with HIV: an updated review of the literature. *Archives of Public Health*. 2014;72(1):40-50.
- [39] Miller TA, DiMatteo MR. Importance of family/social support and impact on adherence to diabetic therapy. *Diabetes Metabolic Syndrome and Obesity: Targets and Therapy*. 2013;6(6):421-6.
- [40] Neri L, Brancaccio D, Rocca RL, Rossa F, Martini A, Andreucci VE. Social support from health care providers is associated with reduced illness intrusiveness in hemodialysis patients. *Clinical Nephrology*. 2011;75(2):125-34.
- [41] Stewart M, Belle Brown J, Weston W, McWhinney I, McWilliam C, Freeman T. *Patientcentered medicine: transforming the clinical method*. 2nd ed. Oxford UK: Radcliffe Medical Press Ltd; 2003.
- [42] Ware NC. Suffering and the social construction of illness: the delegitimation of illness experience in chronic fatigue syndrome. *Medical Anthropology Quarterly*. 1992;6(4):347-61.

- [43] Lehman AM, Lehman DR, Hemphill KJ, Mandel DR, Cooper LM. Illness experience, depression, and anxiety in chronic fatigue syndrome. *Journal of Psychosomatic Research*. 2002;52(6):461-5.
- [44] Rogers CR. A theory of therapy, personality, and interpersonal relationships, as developed in the client-centered framework. In: Koch S, editor. *Psychology: A Study of a Science*. Vol. 3. New York: McGraw-Hill; 1956. pp. 184-256.
- [45] Anderson R, Funnell M. Patient empowerment: reflections on the challenge of fostering the adoption of a new paradigm. *Patient Education and Counseling*. 2005;57(2):153-7.
- [46] Soundy A, Smith B, Dawes H, Pall H, Gimbrere K, Ramsay J. Patient's expression of hope and illness narratives in three neurological conditions: a meta-ethnography. *Health Psychology Review*. 2013;7(2):177-201.
- [47] Bachmann C, Abramovitch H, Barbu CG, Cavaco AM, Elorza RD, Haak R, et al. A European consensus on learning objectives for a core communication curriculum in health care professions. *Patient Education and Counseling*. 2013;93(1):18-26.
- [48] Légaré F, Ratté S, Gravel K, Graham ID. Barriers and facilitators to implementing shared decision-making in clinical practice: update of a systematic review of health professionals' perceptions. *Patient Education and Counseling*. 2008;73(3):526-35.
- [49] West E, Barron DN, Reeves R. Overcoming the barriers to patient-centred care: time, tools and training. *Journal of Clinical Nursing*. 2005;14:435-43.
- [50] Ford S, Schofield T, Hope T. Barriers to the evidence-based patient choice (EBPC) consultation. *Patient Education and Counseling* 2002;47:179-85.
- [51] Williams C, Cantillon P, Cochrane M. The doctor-patient relationship: from undergraduate assumptions to pre-registration reality. *Medical Education*. 2001;35:743-7.
- [52] Glenton C. Developing patient-centred information for back pain sufferers. *Health Expectations*. 2002;5:319-29.
- [53] Esmaeili M, Cheraghi M, Salsali M. Barriers to patient-centered care: a thematic analysis study. *International Journal of Nursing Knowledge*. 2014;25(1):2-8.
- [54] Lamb B, Taylor C, Lamb J, Strickland S, Vincent C, Green J, et al. Facilitators and barriers to teamworking and patient centeredness in multidisciplinary cancer teams: findings of a national study. *Annals of Surgical Oncology*. 2013;20:1408-16.
- [55] Wittenberg-Lyles E, Goldsmith J, Ferrell B. Oncology nurse communication barriers to patient-centered care. *Clinical Journal of Oncology Nursing*. 2013;17(2):152-8.
- [56] Yam B, Rossiter J. Caring in nursing: perceptions of Hong Kong nurses. *Journal of Clinical Nursing*. 2000;9(2):293-302.
- [57] Kruijver I, Kerkstra A, Bensing J, van de Wiel H. Communication skills of nurses during interactions with simulated cancer patients. *Journal of Advanced Nursing*. 2001;34(6):772-9.

- [58] Jallinoja P, Absetz P, Kuronen R, Nissinen A, Talja M, Uutela A, et al. The dilemma of patient responsibility for lifestyle change: perceptions among primary care physicians and nurses. *Scandinavian Journal of Primary Health Care*. 2007;25(4):244–9.
- [59] Chant S, Jenkinson T, Randle J, Russell G. Communication skills: some problems in nursing education and practice. *Journal of Clinical Nursing*. 2002;11:12-21.
- [60] Haidet P, Dains J, Paterniti D, Hechtel L, Chang T, Tseng E, et al. Medical student attitudes toward the doctor-patient relationship. *Medical Education* 2002;36:568-74.
- [61] Skaalvik M, Normann H, Henriksen N. Student experiences in learning person-centred care of patients with Alzheimer's disease as perceived by nursing students and supervising nurses. *Journal of Clinical Nursing*. 2010;19(17–18):2639-48.
- [62] Suikkala A, Leino-Kilpi H. Nursing student-patient relationship: experiences of students and patients. *Nurse Education Today*. 2005;25(5):344-54.
- [63] Tuohy D. Student nurse-older person communication. *Nurse Education Today*. 2003; 23(1):19-26.
- [64] Kotecki C. Baccalaureate nursing students' communication process in the clinical setting. *Journal of Nursing Education*. 2002;41:61-8.
- [65] Hjorleifsdottir E, Carter D. Communicating with terminally ill cancer patients and their families. *Nurse Education Today*. 2000;20(6):646-53.
- [66] McCarthy B, O'Donovan M, Twomey A. Person-centred communication: design, implementation and evaluation of a communication skills module for undergraduate nursing students—an Irish context. *Contemporary Nurse*. 2008;27(2):207-22.
- [67] Girdley D, Johnsen C, Kwekkeboom K. Facilitating a culture of safety and patient-centered care through use of a clinical assessment tool in undergraduate nursing education. *Journal of Nursing Education*. 2009;48(12):702-5.
- [68] McKeon L, Norris T, Cardell B, Britt T. Developing patient-centered care competencies among prelicensure nursing students using simulation. *Journal of Nursing Education*. 2009;48(12):711-5.
- [69] McKonkey R, Morris I, Purcell M. Communications between staff and adults with intellectual disabilities in naturally occurring settings. *Journal of Intellectual Disability Research*. 1999;43(3):194-205.
- [70] Alasad J, Ahmad M. Communication with critically ill patients. *Journal of Advanced Nursing*. 2005;50(4):356-62.
- [71] Hemsley B, Sigaffos J, Balandin S, Forbes R, Taylor C, Green V, et al. Nursing the patient with severe communication impairment. *Journal of Advanced Nursing*. 2001;35(6):827-35.
- [72] Bastiaens H, Van Royen P, Pavlic D, Raposo V, Baker R. Older people's preferences for involvement in their own care: a qualitative study in primary health care in 11 European countries. *Patient Education and Counseling*. 2007;68(1):33-42.

- [73] Curtis-Tyler K. Levers and barriers to patient-centred care with children: findings from a synthesis of studies of the experiences of children living with type 1 diabetes or asthma. *Child: Care, Health and Development* 2010;37(4):540-50.
- [74] Windover AK, Boissy A, Rice TW, Gilligan T, Velez VJ, Merlino J. The REDE model of healthcare communication: optimizing relationship as a therapeutic agent. *Journal of Patient Experience*. 2014;1(1):8-13.
- [75] Rider EA, Hinrichs MM, Lown BA. A model for communication skills assessment across the undergraduate curriculum. *Medical Teacher*. 2006;28(5):e127-34.
- [76] Kurtz SM, Silverman JD, Draper J. Teaching and learning communication skills in medicine. 2nd ed. Oxford, UK: Radcliffe Medical Press; 2004.
- [77] Frankel RM, Stein T. Getting the most out of the clinical encounter: the four habits model. *Journal of Medical Practice Management*. 2001;16(4):184-91.
- [78] Baile W, Walters R. Challenging emotional conversations with patients and families: a guide to forming a therapeutic alliance with patients and families. www.mdanderson.org/icare: Texas: The University of Texas MD Anderson Cancer Center; 2012.
- [79] Bergin F. Successful presentations. Englewood Cliffs, NJ: Prentice Hall; 1995.
- [80] Thompson BM, Teal CR, Scoot DM, Manning SN, Greenfield E, Shada R, et al. Following the clues: teaching medical students to explore patients' context. *Patient Education and Counseling*. 2010;80(3):345-50.
- [81] Bleiker J, Knapp K, Frampton I. Teaching patient care to students: a blended learning approach in radiography education. *Radiography* 2011; 17(3): 235-40.
- [82] Herxheimer A, McPherson A, Miller R, Shepperd S, Yaphe J, Ziebland S. Database of patients' experiences (DIPEX): a multi-media approach to sharing experiences and information. *The Lancet*. 2000;355(9214):1540-3.
- [83] Maguire P, Pitceathly C. Key communication skills and how to acquire them. *British Medical Journal*. 2002;325(7366):697-700.