We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Optimizing Antimicrobial Agents in Endodontics

Patricia P. Wright and Laurence J. Walsh

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/67711>

Abstract

In endodontic (root canal) treatment, a multispecies bacterial and fungal infection is present in a place that is inaccessible to the host immune system and which offers physical protection from applied topical agents. All current protocols for irrigation suffer various deficits in performance, which is why further research on alternative approaches to using antimicrobial substances is warranted. This chapter examines the technical and clinical factors which influence the performance of antimicrobial biocide-based therapies used in endodontics within dental practice, addressing issues around instability of biocides, the influence of pH, the role of physical agitation and the challenge of penetration into biofilms and into confined spaces. A range of methods to overcome the challenges in performance are described, including novel solvents and vehicles for biocides, stabilizing agents, physical agitation and the use of activation protocols including the use of intense light, ultrasound and laser-generated shockwaves to improve the effect of biocides. While specific examples are given from the dental setting of endodontics, the principles have broader application to medicine and to general industry.

Keywords: biofilms, endodontics, biocides, ultrasonics, lasers, disinfection

1. Introduction

Antimicrobial agents which act as biocides have an important place in modern health care as they overcome many of the limitations of antibiotic and antifungal medicines by attacking not one but many targets, making the development of resistance through spontaneous mutation difficult or impossible. There are a wide range of biocides in common use, and their major mechanisms of attack on bacteria and fungi vary according to the agent chosen (**Table 1**). Considering situations where infection is present and which are very challenging to treat with antibiotics, the situation of the infected root canal is the focus of this chapter.

Oxidation of biomolecules: Sodium hypochlorite (NaOCl), hydrogen peroxide (H ₂ O ₂), ozone (O ₃), photoactivated disinfection (PAD).
Cell membrane damage: Chlorhexidine (CHX), phenolic agents, calcium hydroxide (Ca(OH) ₂), functional antimicrobial peptides, nanoparticles (NPs) (including chitosan, gold (Au) and silver (Ag)).
Protein coagulation: Laser photothermal disinfection.

Table 1. Primary mechanisms of action of broad-spectrum antimicrobial agents.

In this particular clinical situation, there is a polymicrobial infection with multiple species of bacteria as well as occasionally fungi also being present in a dense biofilm that penetrates into the tubules of the dentine of the tooth root. The organisms and their products including endotoxins cause severe inflammatory reactions in the adjacent bone and soft tissues. The location of the biofilm within the root canal system of the tooth makes it inaccessible to the host immune system, while the tubules give physical protection from agents which are applied topically. The goal of endodontic (root canal) treatment is to decontaminate the entire root canal system; however, in many cases, viable bacteria remain at the end of treatment, causing ongoing inflammation [1], with accompanying pain and other complications [2]. Retreatment using either non-surgical or surgical methods is focussed on removing persisting microbial contaminants [3].

2. The rationale for irrigation with disinfectant solutions

The use of physical instrumentation alone, such as files or ultrasonics, does not give adequate debridement of the root canal system because of its complex three-dimensional shape, but creates some space to permit better permeation of irrigation solutions [4, 5]. The mainstay of microbial therapy is the application of multiple chemical agents as partners to physical debridement. Most current protocols involve copious irrigation with 2.5–6% solutions of sodium hypochlorite (NaOCl), supplemented in some cases with further treatments using 2% chlorhexidine (CHX). The antimicrobial actions of such agents are related to their concentration, viscosity and ability to wet the walls of the root canal system [6]. Clinical studies have shown that supplementary steps such as placing a paste of calcium hydroxide (Ca(OH)₂) as an inter-appointment medications can further reduce the microbial bioburden [7].

The properties of an ideal irrigation solution are listed in **Table 2**. None of the current agents used alone fulfil all these requirements. This is why combinations of approaches and improved formulation are such an important goal and the focus of the current discussion.

Since all current irrigation protocols suffer various deficits in performance, further research on alternative approaches to using antimicrobial substances is warranted. The purpose of this chapter is to review the various ways that antimicrobial actions of these materials can be enhanced, using combinations of products, altered vehicles, chemical activation and physical agitation.

Effectiveness: Exerts a rapid antimicrobial action, effective on both Gram-positive and Gram-negative bacteria, effective on fungi, able to penetrate into biofilms, able to penetrate into dentine tubules, easy to dispense and use clinically, easily removed from the root canal, dissolves organic tissue, dissolves smear layer, removes debris from the canal, retains activity in the presence of organic matter.
Safety: Not allergenic, not mutagenic, not toxic to human cells, compatible with periapical tissues.
Tooth considerations: Non-staining, does not react adversely with dental materials, does not weaken tooth structure.
Cost: Low cost of manufacture, easily prepared, good long-term stability and long shelf life

Table 2. Properties of an ideal disinfecting agent for endodontics.

3. Sodium hypochlorite

As a solution ranging in concentration from 0.5 to 6%, sodium hypochlorite (NaOCl) is currently the most popular irrigant used in endodontics [8]. The solutions used in dental practice for root canal treatment are a mixture of NaOCl with sodium chloride (to make the solution isotonic), sodium hydroxide (as a pH modifier) and a surfactant. They differ from domestic preparations used as disinfectants, e.g. for sanitizing baby bottles, by being more alkaline (with a pH from 10 to 12) and having the ability to dissolve vital and non-vital soft tissues [9, 10].

Altering the pH influences both the antibacterial effects and tissue-dissolving capacity of NaOCl preparations. In low pH solutions, hypochlorous acid (HOCl) predominates, which has disinfecting actions. This small neutral molecule enters bacterial cells where it oxidizes lipids and proteins, and also reacts with ferrous ions in a reaction similar to the Fenton reaction, to produce hydroxyl radicals [11]. At high pH, both hydroxyl and hypochlorite anions are present. The hypochlorite anion can destroy protein by the formation of chloramines, while the hydroxyl ion reacts with lipids in a saponification reaction to degrade them, thus increasing tissue-dissolving capabilities [12].

3.1. Advantages

NaOCl solutions have a broad antimicrobial spectrum and are effective on both bacteria and fungi such as *Candida albicans* [13, 14]. A particular challenge in endodontics is the presence of highly resistant organisms, particularly *Enterococcus faecalis*, that survive in extremes of acidic and alkaline pH and can withstand nutritional deprivation [15]. NaOCl can disrupt biofilms of *E. faecalis*, but this requires exposure for up to 5 min [16].

Solutions of NaOCl are able to dissolve necrotic soft tissue remnants [17], because it has strong but non-specific proteolytic activity [18, 19]. A further advantage is that when mixed with hydrogen peroxide (H₂O₂), effervescence is produced which can assist in physical debridement [20].

3.2. Limitations

Bacteria located on the surface are easily accessible and can readily be inactivated. Those lodged deep within the dentinal tubules are protected from contact and thus will persist in a

viable state. In the absence of aggressive agitation, NaOCl has a limited ability to eliminate *E. faecalis* when in the biofilm state and deep within dentine tubules [21].

The potency of NaOCl solutions, expressed as the available chlorine in parts per million, declines over time, because the solutions degrade readily, and even more so when exposed to heat and light. This can occur because the stock solutions are kept too long or have been dispensed into clear or translucent plastic syringes which allow in extraneous light [22].

NaOCl has an unpleasant chlorine odour, and it interacts adversely with certain antibiotic medicaments containing tetracycline, as well as with chlorhexidine, causing inactivation through oxidation. These reactions also produce highly staining end products which can cause tooth discolouration [23].

NaOCl corrodes metallic instruments, which may lead to their premature failure (breakage or separation). The proteolytic actions of NaOCl cause severe irritant reactions when only small volumes of NaOCl are extruded into soft tissues, causing dramatic swelling [24]. These reactions are more severe in formulations which include sodium hydroxide as an alkalinizing agent, because of caustic actions on soft tissues [25].

3.3. Enhancement

Adding calcium hydroxide ($\text{Ca}(\text{OH})_2$) improves the effectiveness of NaOCl in terms of reducing bacterial counts as well as endotoxin levels [26]. Another chemical means of enhancing sodium hypochlorite is the addition of detergents to lower the surface tension and enhance penetration, for example, the quaternary ammonium compounds cetrimide and benzalkonium chloride, which themselves are antibacterial agents [27–29]. Adding in surfactants offers the option of using lower and thus less toxic concentrations of NaOCl for disinfection. The same strategy of adding detergents can be applied to other antimicrobial agents to enhance their effectiveness.

The actions of NaOCl can also be enhanced by physical activation, for example, using ultrasonics or pulsed middle infrared lasers (such as Er:YAG or Er,Cr:YSGG lasers) to agitate the solution. Lasers are better than ultrasonic agitation in this regard [30, 31]. Laser activation of NaOCl also enhances the removal of soft tissue and debris from regions that are difficult to clean [32]. Laser agitation employs low average power settings and short pulse durations; thus, there is no ablation of the root structure [33]. The general principle of using physical agitation to improve the effects of antimicrobial agents also applies to solutions of hydrogen peroxide and $\text{Ca}(\text{OH})_2$.

4. Chlorhexidine

Chlorhexidine (CHX) comes in various forms with the gluconate and the acetate being the most widely used in dentistry as disinfectants. It is typically used as a 2% solution of the gluconate form [10, 14, 34, 35] as a final flush [10, 36, 37]. CHX has a narrow spectrum for Gram-negative bacteria but is effective against most Gram-positive bacteria and also fungi. CHX is the agent of choice when there are Gram-positive resistant enterococci present in the root canal, which may be the case in retreatment situations [38].

4.1. Advantages

The actions of CHX in the root canal are strongly influenced by pH. At pH 5.5–6.0, chlorhexidine exists as a di-cation [39, 40]. This positive charge allows it to bind to negatively charged substances including the hydroxyapatite mineral of tooth structure, bacterial polysaccharides and particularly to Gram-positive bacteria [41] as well as onto the surface of biofilms [42]. Once adsorbed onto tooth structure, CHX can prevent subsequent microbial colonization on the surface [43].

CHX exerts antifungal actions against *C. albicans*, a property it shares with hydrogen peroxide and NaOCl [44]. It also has mild anti-collagenolytic activity, but no ability to dissolve necrotic tissues [42].

4.2. Limitations

While effective against key Gram-negative endodontic pathogens such as *Porphyromonas endodontalis*, *Porphyromonas gingivalis* and *Prevotella intermedia* [38], CHX suffers from problems of inherent resistance with pseudomonads and certain other Gram-negative bacteria. Resistance to CHX occurs in *Enterobacter* spp., *Pseudomonas* spp., *Proteus* spp. and *Providencia* spp. [45].

When used as an irrigant in the root canal in the absence of physical agitation, CHX has only a limited ability to eliminate *E. faecalis* because of poor penetration into the dentinal tubules [46]. Likewise, solutions of 2% CHX penetrate poorly into biofilms and are ineffective for dissolving biofilms [47].

When in contact with tissues, CHX is irritant and can delay healing [48, 49]. As with sodium hypochlorite, care must be taken to prevent accidental extrusion into soft tissues. Using a side-vented needle rather than a conventional needle can reduce the volume of fluid that is extruded during irrigation [50, 51].

Despite reported cases of allergy to CHX in medical settings, few have occurred in dental practice. IgE-mediated allergic responses to CHX manifest as redness, itching (urticaria) and swellings [52] and can progress to anaphylaxis [53]. The high propensity for allergy distinguishes CHX from all other disinfectants discussed in this chapter.

CHX suffers from moderate problems of stability as it can undergo cleavage with oxidation, both during storage and when in contact with oxidants such as hydrogen peroxide, ozone or NaOCl. The key end product of oxidative degradation of CHX is para-chloroaniline, an orange to brown coloured substance that is highly irritant and allergenic [23, 40]. As well as causing staining of tooth structure [40], para-chloroaniline may also interfere with the seal of the final root filling [54].

4.3. Enhancements

To enhance the actions of CHX, it can be combined with low concentrations of hydrogen peroxide, mixed freshly so as to limit the possible oxidation of the molecule [55]. Likewise, adding calcium hydroxide improves the antimicrobial actions of CHX [26].

Adding positively charged detergents such as cetrimide can potentiate the antibacterial actions of CHX [39, 56, 57]; however, because of its cationic nature, it cannot be mixed with anionic detergents [56], since these cause precipitation to occur.

5. Calcium hydroxide

Water-based pastes of calcium hydroxide (Ca(OH)_2) are the most commonly used inter-visit dressing material in endodontics [58]. When placed in the root canal, Ca(OH)_2 elevates the local pH, making this unfavourable for the growth of most bacteria [43, 59]. An in vivo study revealed that after 14 days of exposure to a Ca(OH)_2 paste, for 41 out of 44 endodontic pathogens, both the rate of detection in samples and mean bacterial counts declined [58]. Ca(OH)_2 both inactivates bacteria and reduces endotoxin levels [43], through release of hydroxyl (OH^-) ions over a prolonged period of time [43]. These destroy bacterial cell walls and cytoplasmic membranes by degrading fatty acids, thereby allowing the leakage of cellular components. They also inactivate bacterial enzymes. High pH levels within bacterial cells alter the charge of various organic molecules and so interfere with the transport of bacterial nutrients [60]. OH^- ions also denature proteins and damage DNA [61]. Calcium ions may contribute to antibacterial actions by neutralizing negatively charged molecules [62].

5.1. Advantages

Ca(OH)_2 promotes the formation of dental hard tissues, a characteristic exploited in techniques such as apexification where continued formation of the root is intended. Enhanced mineralization is due to activation of alkaline phosphatase, resulting in the release of phosphate groups which then react with calcium ions [60]. While Ca(OH)_2 is biocompatible, the extremely irritant and caustic nature of sodium and potassium hydroxides makes these both unsuitable for use in the root canal of teeth. Neither sodium nor potassium hydroxides have mineralizing actions.

5.2. Limitations

A saturated solution or paste of Ca(OH)_2 in water has a pH of approximately 12.5–12.8. When placed in a tooth, buffering by dentine proteins and carbonate ions lowers the effective pH that can be achieved by water-based solutions and pastes by 1–2 pH units, making the effective pH achieved within the dentine around 10 [63]. This is a problem because *E. faecalis* can withstand a pH of 10 and thus is resistant to the effects of traditional water-based Ca(OH)_2 pastes [59, 64, 65]. Likewise, many water-based Ca(OH)_2 products exert limited antifungal activity [44].

5.3. Enhancements

While Ca(OH)_2 can be combined with CHX, the combination is not significantly more effective against *E. faecalis* [46]. In contrast, Ca(OH)_2 pastes that are enriched with ibuprofen or diclofenac become more effective [66].

Ca(OH)_2 has low solubility in water (0.17% by weight at room temperature), so only small amounts can be dissolved. While the dissolution characteristics can be improved marginally when nanoparticles are used, there remains an upper ceiling of pH 12.5–12.7 for Ca(OH)_2 in a water-based solvent, which explains why pH values of commercial products are always in this narrow range [43].

Work based on replacing water with non-aqueous solvents has shown that several biocompatible fluids are much better solvents for Ca(OH)_2 than water, including propylene glycol, polyethylene glycol (PEG) and glycerol. In each of these fluids, there is higher measured OH^- release, as measured using special electrodes designed for non-aqueous solvents as well as by titration [67, 68]. A preferred solvent is a mixture of two forms of polyethylene glycol (PEG), one being the foundation (PEG 400) and the other being a thickener (PEG 3350), to generate the preferred creamy consistency required for application into the root canal. This PEG blend is a potent solvent for Ca(OH)_2 which has a high release of OH^- ions into any water-based environment, since PEG 400 is miscible with water [69, 70]. This then translates into greater movement of OH^- ions through the roots of human teeth than water-containing Ca(OH)_2 pastes [70]. This can be explained by the common ion effect, with OH^- being the common ion in water.

6. Hydrogen peroxide

At a final concentration of 3–6%, H_2O_2 is a commonly used disinfectant. It generates oxygen radicals [71], of which the hydroxyl radical is the most important since it is the strongest oxidizer [72]. H_2O_2 generates effervescence which provides physical clearance of microbial deposits [73]. The elevated oxygen concentration it creates is unfavourable for the growth of strict anaerobes.

6.1. Limitations

Elevating the pH using an alkali metal hydroxide such as lithium, sodium or potassium hydroxide will accelerate the decomposition of H_2O_2 and provide an alkaline pH [72]. The limitation in this approach is that such hydroxides are inherently caustic.

6.2. Enhancements

The actions of H_2O_2 as a disinfectant can be enhanced by adding a suitable catalyst such as manganese or ferrous ions [74]. The latter can also be employed in the Fenton reaction, where the ferrous ion reacts with H_2O_2 to form hydroxyl and other radicals [74]. Blue, violet or ultraviolet light can provide photochemical activation for this reaction, enhancing further the generation of reactive oxygen species [75]. The Fenton reaction can also occur within bacterial cells, causing cell death [11]. Likewise, photolysis of H_2O_2 using 405 nm violet light combined with ultrasound activation can potentiate hydroxyl radical formation, with a synergistic antibacterial action between light and ultrasound activation [76].

The temperature of H_2O_2 and thus its rate of breakdown can be increased by using dyes that absorb the appropriate wavelength of intense light, for example, blue light into orange or yellow dyes. Titanium dioxide in forms similar to those used in sunscreens allows broad-spectrum light sources to be used to activate H_2O_2 .

Mixing ozone (O_3) and H_2O_2 gives co-catalysis and enhanced effectiveness [77]. As with H_2O_2 , physical agitation of solutions of O_3 in water using ultrasound improves their effectiveness [78]. When H_2O_2 is activated by ultrasonic agitation, the enhanced production of hydroxyl radicals by sonolysis causes increased bacterial killing [79]. This ultrasonic activation can be augmented with 405 nm violet light, as both activation pathways generate hydroxyl radicals in a synergistic manner [76].

The generation of oxygen gas bubbles from H_2O_2 and the associated disrupting effect of these on microbial deposits can be enhanced using middle infrared lasers which emit in the 2700–3000 nm wavelength range, such as the Er,Cr:YSGG and Er:YAG lasers. The energy from these lasers absorbs strongly into both water and H_2O_2 , creating with each laser pulse bubbles of air, steam and oxygen. Shockwaves generated by bubble implosions create shear stresses on the walls of the root canal. Fluid movements at high speeds (in the order of 100 km/h) cause disruption of microbial deposits and smear layers [80, 81]. Because the streaming movements of such fluids causes most of the fluid movement being directed back towards the point of entry of the fibre into the tooth, the volume of fluids extruded by laser activation of various irrigation fluids is no more than when conventional irrigation is undertaken using syringes [50]. The frequent replacement of fluids also provides a cooling effect, so that heat does not accumulate [82, 83].

When H_2O_2 is added to pure water, the absorption curve of the mixture is left shifted from that of water alone, allowing lasers operating at wavelengths from 900 to 1100 nm to have much stronger absorption [84]. This approach allows handheld near-infrared lasers to be used for cavitation-based removal of bacterial deposits and smear layer from the root canal. There is accompanying photothermal disinfection of bacteria located deep in dentine tubules because such wavelengths have high transmission into dentine [85, 86]. When H_2O_2 is used in this clinical application, the concentration is below 6% by volume, as this is the threshold for soft tissue injury [73].

7. Phenolic agents

The use of phenolic agents such as camphorated monochlorophenol (CMCP) and other related hydrophobic antimicrobial agents such as essential oils has a long history in endodontics. Chlorination of phenols enhances their antibacterial action [61]. The antibacterial effect of CMCP is better at low pH. This agent has a high volatility, which may enhance its penetration into dentinal tubules [43].

7.1. Limitations

Many phenolic agents are hydrophobic and have low water solubility, so they are typically formulated in a hydrophobic solvent [87]. As a result, there are poor wetting of the root canal

walls, limited contact with biofilms in the canal and poor penetration into the water-rich environment of dentine tubules.

7.2. Enhancements

$\text{Ca}(\text{OH})_2$ can be added to phenolic products to make them more effective antimicrobial agents, as this disrupts the biofilm matrix [88]. The contact between hydrophobic phenolic agents and the walls of the root canal can be enhanced by formulating these into a water base containing surfactants or by using water-miscible solvents such as low molecular weight forms of PEG. The proper choice of agents can provide not only disinfectant actions but also anti-inflammatory and analgesic effects. The latter are therapeutically desirable when patients present in pain from endodontic infections [69].

8. Nanoparticles

Nanoparticles (NPs) with diameters in the range of 1–100 nm have special properties, including increased chemical reactivity and a large surface area [89]. They can exist singly, in groups or in clusters. In endodontics, NPs suggested for use as adjuncts to conventional treatments include chitosan, poly(lactic-co-glycolic acid) (PLGA), silver (Ag), gold (Au) and $\text{Ca}(\text{OH})_2$ [90]. Chitosan is derived from the chitin exoskeletons of various arthropods [91]. It is polycationic and attaches strongly to negatively charged bacterial cell walls causing their disruption, with subsequent leakage of cellular contents. Chitosan is also capable of disrupting the extracellular matrix of biofilms [92]. Low molecular weight forms of chitosan are more antimicrobial than those with high molecular weights [93]. Carboxymethyl chitosan has been included in endodontic sealers to provide antibacterial properties [94].

PLGA is a highly biocompatible copolymer of lactic and glycolic acid. Both PLGA and chitosan NPs can be loaded with photosensitizers for use in photoactivated disinfection (PAD). Functionalization of NPs enhances the action of photosensitizers such as rose bengal and methylene blue [95, 96].

NPs of metallic oxides, such as calcium oxide, magnesium oxide, zinc oxide and titanium oxide, can exert antibacterial actions by generating reactive oxygen species [92]. The high chemical reactivity and large surface area of NPs enhance these actions over their normal counterparts, as has been shown for calcium oxide [97]. NPs can also be made from $\text{Ca}(\text{OH})_2$ [69, 98].

Of the various metals that form NPs, silver (Ag) is of particular interest because the antibacterial effects of silver are well known in endodontics. The same principles can be employed with ionic solutions of Ag compounds (such as silver fluoride) although issues of discolouration limit their clinical application. Using NPs of Ag in the 2–12 nm range can reduce problems of tooth discolouration. AgNPs may slowly release silver ions (Ag^+) which then interact with biological systems, attracting electrons away from sulphur and nitrogen atoms in the sulphhydryl and amino groups of biological molecules such as proteins, or with the nitrogenous purines and pyrimidines of DNA and RNA [99].

Various methods are known for producing AgNPs, ranging from chemical reactions to bio-synthesis through to electrochemical methods [100, 101]. Solutions containing AgNPs can be generated by high-voltage ultralow-current electrolysis of water using silver electrodes, followed by irradiation of the solution to violet light (400–430 nm wavelength). This alters the properties of the AgNPs and ensures only clusters remain rather than ions [102]. The solutions of such particles are optically clear, free of Ag ions and quite stable.

8.1. Limitations

While silver, gold, tin and zinc all have potential for use in ionic or NP forms, issues of staining and long-term stability need to be addressed. There are now systems for stabilizing stannous (tin) ions using sodium hexametaphosphate, as well as potassium iodide protocols for reducing staining from ionic silver. These have been employed in preventive and restorative dentistry but are yet to be optimized for endodontics [103, 104].

Toxicity issues with NPs require further investigation. AgNPs are not cytotoxic at 25 µg/mL or lower concentrations; however, this threshold is based on cell culture studies using a mouse fibroblast cell line (L929). Thus, for safe clinical use, animal and human studies are needed to assess the toxicological profile in greater detail [105].

8.2. Enhancements

Displacement of NPs into poorly accessible regions of the root canal, such as dentine tubules, can be achieved by using ultrasonic or pulsed laser agitation of the solution.

The application of NPs in solutions for irrigation is at an early stage of development. AgNPs have been used in combination with traditional $\text{Ca}(\text{OH})_2$ to increase the killing of *E. faecalis* in biofilms [106]. There is also the possibility of combining metal NPs with traditional antibiotics to enhance their effectiveness. The interactions of metals with antibiotics need to be optimized, so that effectiveness is enhanced rather than impaired. Tetracyclines bind metal ions by chelation, and this alters their effectiveness. Tetracyclines with their long history of use in endodontics would be a logical first place to study such interactions. It is also possible that NPs can be functionalized with photosensitizers to increase their antimicrobial effectiveness [89]. There is already evidence that AgNPs can be combined with photosensitizers in a dual-treatment approach [107].

9. Functional peptides

Custom peptides could be designed which can inactivate particular bacterial pathogens such as *E. faecalis*. A typical primary root canal infection is a mixed flora of 20–30 species of micro-organisms, so the first challenge in using antimicrobial peptides is to ensure that the coverage is sufficient. The second challenge is that some common Gram-negative endodontic pathogens such as *P. endodontalis* and *P. gingivalis* produce large quantities of proteases (such as the gingipains) [108], which could readily degrade such peptides and limit their useful life as therapeutic agents. A third challenge with peptides is their higher cost of manufacture and shelf life than other options.

10. Laser-based photothermal disinfection

Penetration of near-infrared light in the wavelength range from 800 to 1100 nm through dentine is high, allowing laser light to reach bacteria embedded deep within dentine tubules [109]. Absorption of laser energy into melanin, water, porphyrins and other molecules then denatures enzymes and so kills bacteria through photothermal actions. As the heat will then be transmitted into adjacent tooth structure and periodontal ligament, the laser energy must be delivered in pulsed mode to allow cooling. Typically, the laser is activated over several passes as the fibre is withdrawn, causing irradiation of the entire root canal system [110].

Laser energy used for photothermal disinfection can also exert biostimulatory effects and so enhance healing and reduce inflammation [111, 112]. A recent study using a 980 nm diode laser showed both bacterial inactivation and biostimulatory effects [113].

11. Photoactivated disinfection

Besides the term photoactivated disinfection (PAD), the literature in this area contains many terms that refer to the same process, such as photodynamic therapy (PDT), antimicrobial PDT (aPDT), light-activated disinfection (LAD), advanced non-invasive LAD (ANILAD) and lethal laser photosensitization (LLP).

Lasers or light-emitting diodes (LEDs) used as sources of intense light cause electronic activation of photosensitizers, which then produce ROS. Examples of light and dye combinations include visible red light with methylene blue or toloum chloride [114], visible blue light with curcumin and green light with rhodamine B dye. The release of ROS is a photodynamic process, i.e. driven by light, with no accompanying thermal effects [115]. Even though the ROS produced are extremely short lived, photosensitizers are able to exert their actions on pathogens effect because they attach directly to microbial cell walls. The porphyrins found within many facultative and strict Gram-negative anaerobes can act as endogenous photosensitizers and absorb blue light, such that external dyes are not needed. Gram-positive bacteria and fungi do not contain large amounts of porphyrins and so are much less sensitive to the effects of blue light.

Maximal activation of photosensitizers requires many parameters to be optimized other than the specific absorption of the dye matching the emission wavelength of the light source. The dye concentration must be optimized, since too little means only a low concentration of ROS will be produced. When the dye solution is too concentrated, the optical density of the solution becomes too great and the penetration of light is limited to being superficial, and the effects become photothermal rather than photodynamic. Concentrated dyes also have the potential to stain tooth structure. Adequate amounts of oxygen need to be present locally to support the reaction. For this reason, some protocols include oxygen carriers or oxidizers such as H_2O_2 to enhance the action of the photosensitizers [116]. The pH of the solvent for the dye must be optimized for the concentration and type of ROS desired, e.g. a higher pH means more perhydroxyl radicals will form with greater antimicrobial actions. Inclusion of a surfactant is important for ensuring penetration of the dye into biofilms and into difficult-to-reach areas such as dentine tubules [117]. The optical fibre delivery systems used to deliver intense

light from lasers or LEDs may include special diffusers to optimize the angular distribution of light into the dye.

PAD should be used after conventional disinfection with NaOCl, but not as a complete replacement to traditional chemical disinfectants [118, 119]. It can also be used as a supplementary form of root canal disinfection in endodontic retreatment cases [120]. An animal study which used PAD as an adjunct to standard disinfection with NaOCl reported a larger reduction in periapical lesions and more periapical regeneration than standard treatment protocols [121]. While some in vivo studies have recommended PAD as a possible alternative to NaOCl [122, 123], there remains a need to standardize the protocol and optimize the light parameters [114, 124].

11.1. Enhancements

To achieve an even distribution of light across the dye solution, a range of optical fibre tips have been developed which have side-firing capabilities, giving even lateral dispersion of light because of surface patterning [125]. Pulsing the light source can enhance the production of ROS by the dye. This has been shown in studies of PAD using thick *E. faecalis* biofilms in the root canal [126]. Pulsing the light source lowers the requirements for cooling the diode laser or LED and also improves the stability of the wavelength emitted by reducing drift to longer wavelengths. Finally, ultrasonic agitation can increase the effectiveness of PAD [127]. Photosensitizer dyes can also be used in combination with NPs as discussed above, as well as with H_2O_2 [128, 129].

12. Fluorescence control of disinfection approaches

Both planktonic microorganisms and biofilms in the root canal system can be detected using fluorescence. The point where the canal is free of microbial contamination can be identified precisely, providing an endpoint to treatment [130, 131]. Fibres with special surface characteristics allowing light delivery for fluorescence excitation and light collection from fluorescence emissions have been designed, and their performance demonstrated in various situations [132–134].

12.1. Limitations

The major limitation to such systems being used is when the fluorescence properties of tooth structure have been enhanced because of incorporation of tetracycline. This causes increased fluorescence which must be corrected for [130, 131]. While few irrigants or medicaments used in modern endodontics give fluorescence emissions, certain potent oxidizers can quench (reduce) fluorescence emissions by bacteria. Using scavengers such as sodium ascorbate or sodium thiosulphate can remove such quenching actions and allow reliable fluorescence readings to be obtained.

13. Conclusions

There are a range of options available for disinfection of the root canal. Recognizing that none of the existing agents used alone aligns with all the properties of an ideal agent (as listed in **Table 2**), the effort to enhance the capabilities of the existing agents and find new approaches must continue. Major directions for the future include improved formulations such as the use of non-aqueous solvents for Ca(OH)_2 for medicaments to remain in the root canal between successive appointments, as well as changes to clinical treatments delivered by the dentist during endodontic treatment. The latter include agitation of water-based disinfectant irrigants with lasers, and the combination of antimicrobial NPs with optical technologies, for synergy of effects as a final high-level disinfection step before the root canal is filled. The ideal protocol should address issues of clinical time, materials cost and complexity, as well as efficacy and safety. Using fluorescence to measure levels of pathogens can provide an endpoint to clinical interventions, to inform the practitioner when pathogens no longer remain. Including antimicrobial agents into dental materials used to fill or seal the root canal is a further avenue to explore, applying the principles discussed in this chapter.

Author details

Patricia P. Wright and Laurence J. Walsh*

*Address all correspondence to: l.walsh@uq.edu.au

School of Dentistry, The University of Queensland, Brisbane, Australia

References

- [1] Paiva SS, Siqueira JF Jr, Rôças IN, Carmo FL, Ferreira DC, Curvelo JA, Soares RM, Rosado AS. Supplementing the antimicrobial effects of chemomechanical debridement with either passive ultrasonic irrigation or a final rinse with chlorhexidine: a clinical study. *J Endod.* 2012;38:1202-6.
- [2] Manfredi M, Figini L, Gagliani M, Lodi G. Single versus multiple visits for endodontic treatment of permanent teeth. *Cochrane Database Syst Rev.* 2016;12:CD005296.
- [3] Del Fabbro M, Corbella S, Sequeira-Byron P, Tsesis I, Rosen E, Lolato A, Taschieri S. Endodontic procedures for retreatment of periapical lesions. *Cochrane Database Syst Rev.* 2016;10:CD005511.
- [4] Carrotte P. Endodontics: part 7. Preparing the root canal. *Br Dent J.* 2004;197:603-13.
- [5] Kuzekanani M, Walsh LJ, Yousefi MA. Cleaning and shaping curved root canals: Mtwo vs ProTaper instruments, a lab comparison. *Int J Dent Res.* 2009;20:268-70.

- [6] Regan JD, Fleury AA. Irrigants in non-surgical endodontic treatment. *J Ir Dent Assoc.* 2006;52:84-92.
- [7] Paiva SS, Siqueira JF Jr, Rôças IN, Carmo FL, Leite DC, Ferreira DC, Rachid CT, Rosado AS. Clinical antimicrobial efficacy of NiTi rotary instrumentation with NaOCl irrigation, final rinse with chlorhexidine and interappointment medication: a molecular study. *Int Endod J.* 2013;46:225-33.
- [8] Shen Y, Gao Y, Lin J, Ma J, Wang Z, Haapasalo M. Methods and models to study irrigation. *Endodont Topics.* 2012;27:3-34.
- [9] Clarkson R, Moule A, Podlich H, Kellaway R, Macfarlane R, Lewis D, Rowell J. Dissolution of porcine incisor pulps in sodium hypochlorite solutions of varying compositions and concentrations. *Aust Dent J.* 2006;51:245-51.
- [10] Schäfer E. Irrigation of the root canal. *ENDO.* 2007;1:11-27.
- [11] Fukuzaki S. Mechanisms of actions of sodium hypochlorite in cleaning and disinfection processes. *Biocontrol Sci.* 2006;11:147-57.
- [12] Estrela C, Estrela CR, Barbin EL, Spanó JCE, Marchesan MA, Pécora JD. Mechanism of action of sodium hypochlorite. *Braz Dent J.* 2002;13:113-7.
- [13] Radcliffe C, Potouridou L, Qureshi R, Hababbeh N, Qualtrough A, Worthington H, Drucker D. Antimicrobial activity of varying concentrations of sodium hypochlorite on the endodontic microorganisms *Actinomyces israelii*, *A. naeslundii*, *Candida albicans* and *Enterococcus faecalis*. *Int Endod J.* 2004;37:438-46.
- [14] Agrawal Vineet S, Rajesh M, Sonali K, Mukesh P. A contemporary overview of endodontic irrigants: a review. *J Dent App.* 2014;1:105-15.
- [15] Walsh LJ, Athanassiadis B. The challenge of endodontic “superbugs” in clinical practice. *Australas Dent Pract.* 2008;19:102-6.
- [16] Clegg MS, Vertucci FJ, Walker C, Belanger M, Britto LR. The effect of exposure to irrigant solutions on apical dentine biofilms in vitro. *J Endod.* 2006;32:434-7.
- [17] Naenni N, Thoma K, Zehnder M. Soft tissue dissolution capacity of currently used and potential endodontic irrigants. *J Endod.* 2004;30:785-7.
- [18] Mohammadi Z. Sodium hypochlorite in endodontics: an update review. *Int Dent J.* 2008;58:329-41.
- [19] Mohammadi Z, Shalavi S. Antimicrobial activity of sodium hypochlorite in endodontics. *J Mass Dent Soc.* 2013;62:28-31.
- [20] Al-Ali M, Sathorn C, Parashos P. Root canal debridement efficacy of different final irrigation protocols. *Int Endod J.* 2012;45:898-906.
- [21] Estrela C, Silva JA, de Alencar AH, Leles CR, Decurcio DA. Efficacy of sodium hypochlorite and chlorhexidine against *Enterococcus faecalis* – a systematic review. *J Appl Oral Sci.* 2008;16:364-8.

- [22] Clarkson R, Moule A, Podlich H. The shelf-life of sodium hypochlorite irrigating solutions. *Aust Dent J*. 2001;46:269-76.
- [23] Walsh LJ, Athanassiadis B. Endodontic aesthetic iatrodontics. *Australas Dent Pract*. 2007;18:62-64.
- [24] Hülsmann M, Hahn W. Complications during root canal irrigation – literature review and case reports. *Int Endod J*. 2000;33:186-93.
- [25] Zehnder M. Root canal irrigants. *J Endod*. 2006;32:389-98.
- [26] Xavier AC, Martinho FC, Chung A, Oliveira LD, Jorge AO, Valera MC, Carvalho CA. One-visit versus two-visit root canal treatment: effectiveness in the removal of endotoxins and cultivable bacteria. *J Endod*. 2013;39:959-64.
- [27] DeOude N. *The Handbook of Environmental Chemistry. Volume 3, Part F: Anthropogenic Compounds*. Heidelberg: Springer Verlag, 1992.
- [28] Baron A, Lindsey K, Sidow SJ, Dickinson D, Chuang A, McPherson JC. Effect of a benzalkonium chloride surfactant–sodium hypochlorite combination on elimination of *Enterococcus faecalis*. *J Endod*. 2016;42:145-9.
- [29] Wang Z, Shen Y, Ma J, Haapasalo M. The effect of detergents on the antibacterial activity of disinfecting solutions in dentin. *J Endod*. 2012;38:948-53.
- [30] Ordinola-Zapata R, Bramante C, Aprecio R, Handysides R, Jaramillo D. Biofilm removal by 6% sodium hypochlorite activated by different irrigation techniques. *Int Endod J*. 2014;47:659-66.
- [31] Peters OA, Bardsley S, Fong J, Pandher G, DiVito E. Disinfection of root canals with photon-initiated photoacoustic streaming. *J Endod*. 2011;37:1008-12.
- [32] Guneser MB, Arslan D, Usumez A. Tissue dissolution ability of sodium hypochlorite activated by photon-initiated photoacoustic streaming technique. *J Endod*. 2015;41:729-32.
- [33] DiVito E, Peters O, Olivi G. Effectiveness of the erbium: YAG laser and new design radial and stripped tips in removing the smear layer after root canal instrumentation. *Lasers Med Sci*. 2012;27:273-80.
- [34] Abraham S, Raj JD, Venugopal M. Endodontic irrigants: a comprehensive review. *J Pharm Sci Res*. 2015;7:5-9.
- [35] Chiniforush N, Pourhajibagher M, Shahabi S, Bahador A. Clinical approach of high technology techniques for control and elimination of endodontic microbiota. *J Lasers Med Sci*. 2015;6:139.
- [36] Haapasalo M, Shen Y, Wang Z, Gao Y. Irrigation in endodontics. *Brit Dent J*. 2014;216:299-303.
- [37] Plotino G, Cortese T, Grande NM, Leonardi DP, Di Giorgio G, Testarelli L, Gambarini G. New technologies to improve root canal disinfection. *Braz Dent J*. 2016;27:3-8.

- [38] Vianna ME, Gomes BP, Berber VB, Zaia AA, Ferraz CC, de Souza-Filho FJ. In vitro evaluation of the antimicrobial activity of chlorhexidine and sodium hypochlorite. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2004;97:79-84.
- [39] Portenier I, Waltimo T, Ørstavik D, Haapasalo M. Killing of *Enterococcus faecalis* by MTAD and chlorhexidine digluconate with or without cetrimide in the presence or absence of dentine powder or BSA. *J Endod.* 2006;32:138-41.
- [40] Basrani BR, Manek S, Sodhi RN, Fillery E, Manzur A. Interaction between sodium hypochlorite and chlorhexidine gluconate. *J Endod.* 2007;33:966-9.
- [41] Rølla G, Løe H, Schiøtt CR. Retention of chlorhexidine in the human oral cavity. *Arch Oral Biol.* 1971;16:1109-33.
- [42] Kanisavaran ZM. Chlorhexidine gluconate in endodontics: an update review. *Int Dent J.* 2008;58:247-57.
- [43] Athanassiadis B, Abbott PV, Walsh LJ. The use of calcium hydroxide, antibiotics and biocides as antimicrobial medicaments in endodontics. *Aust Dent J.* 2007;52(Suppl): S64-82.
- [44] Ferguson JW, Hatton JF, Gillespie MJ. Effectiveness of intracanal irrigants and medications against the yeast *Candida albicans*. *J Endod.* 2002;28:68-71.
- [45] Kampf G. Acquired resistance to chlorhexidine – is it time to establish an ‘antiseptic stewardship’ initiative? *J Hosp Infect.* 2016;94:213-27.
- [46] Saatchi M, Shokraneh A, Navaei H, Maracy MR, Shojaei H. Antibacterial effect of calcium hydroxide combined with chlorhexidine on *Enterococcus faecalis*: a systematic review and meta-analysis. *J Appl Oral Sci.* 2014;22:356-65.
- [47] del Carpio-Perochena AE, Bramante CM, Duarte MA, Cavenago BC, Villas-Boas MH, Graeff MS, Ordinola-Zapata R. Biofilm dissolution and cleaning ability of different irrigant solutions on intraorally infected dentin. *J Endod.* 2011;37:1134-8.
- [48] Grant PB, Walsh LJ. Problems and adverse effects associated with the use of chlorhexidine in dental practice. *Periodont.* 1985;6:33-8.
- [49] Mohammadi Z, Abbott PV. The properties and applications of chlorhexidine in endodontics. *Int Endod J.* 2009;42:288-302.
- [50] George R, Walsh LJ. Apical extrusion of root canal irrigants when using Er:YAG and Er,Cr:YSGG lasers with optical fibers: an in vitro dye study. *J Endod.* 2008;34:706-8.
- [51] Chou K, George R, Walsh LJ. Effectiveness of different intra-canal irrigation techniques in removing intra-canal paste medicaments. *Aust Endod J.* 2014;40:21-5.
- [52] Nagendran V, Wicking J, Ekbote A, Onyekwe T, Garvey LH. IgE-mediated chlorhexidine allergy: a new occupational hazard? *Occup Med.* 2009;59:270-2.
- [53] Tran C, Walsh LJ. Anaphylaxis to dental products in dental hygiene practice. *Hyg Today.* 2011;11:4-7.

- [54] Bui TB, Baumgartner JC, Mitchell JC. Evaluation of the interaction between sodium hypochlorite and chlorhexidine gluconate and its effect on root dentin. *J Endod.* 2008;34:181-5.
- [55] Steinberg D, Heling I, Daniel I, Ginsburg I. Antibacterial synergistic effect of chlorhexidine and hydrogen peroxide against *Streptococcus sobrinus*, *Streptococcus faecalis* and *Staphylococcus aureus*. *J Oral Rehab.* 1999;26:151-6.
- [56] Russell A, Day M. Antibacterial activity of chlorhexidine. *J Hospital Infect.* 1993;25:229-38.
- [57] Ruiz-Linares M, Aguado-Pérez B, Baca P, Arias-Moliz MT, Ferrer-Luque CM. Efficacy of antimicrobial solutions against polymicrobial root canal biofilm. *Int Endod J.* 2017;50:77-83.
- [58] de Souza CAS, Teles RP, Souto R, Chaves MAE, Colombo APV. Endodontic therapy associated with calcium hydroxide as an intracanal dressing: microbiologic evaluation by the checkerboard DNA-DNA hybridization technique. *J Endod.* 2005;31:79-83.
- [59] Mohammadi Z, Dummer PM. Properties and applications of calcium hydroxide in endodontics and dental traumatology. *Int Endod J.* 2011;44:697-730.
- [60] Estrela C, Sydney GB, Bammann LL, Felipe O. Mechanism of action of calcium and hydroxyl ions of calcium hydroxide on tissue and bacteria. *Braz Dent J.* 1995;6:85-90.
- [61] Siqueira J, Lopes H. Mechanisms of antimicrobial activity of calcium hydroxide: a critical review. *Int Endod J.* 1999;32:361-9.
- [62] Upadya M, Shrestha A, Kishen A. Role of efflux pump inhibitors on the antibiofilm efficacy of calcium hydroxide, chitosan nanoparticles, and light-activated disinfection. *J Endod.* 2011;37:1422-6.
- [63] Athanassiadis B, Abbott PV, George N, Walsh LJ. In vitro study of the inactivation by dentine of some antimicrobial endodontic medicaments and their bases. *Aust Dent J.* 2010;55:298-305.
- [64] Athanassiadis B, Abbott PV, George N, Walsh LJ. An in vitro study of the antimicrobial activity of some endodontic medicaments and their bases using an agar well diffusion assay. *Aust Dent J.* 2009;54:141-6.
- [65] Athanassiadis B, Abbott PV, George N, Walsh LJ. An in vitro study of the anti-microbial activity of some endodontic medicaments and their bases against *Enterococcus faecalis* biofilms. *Aust Dent J.* 2010;55:150-5.
- [66] de Freitas RP, Greatti VR, Alcalde MP, Cavenago BC, Vivan RR, Duarte, MAH, Weckwerth PH. Effect of the association of nonsteroidal anti-inflammatory and antibiotic drugs on antibiofilm activity and pH of calcium hydroxide pastes. *J Endod.* 2017;43:131-4.
- [67] Walsh LJ, Athanassiadis B. Non-aqueous solvents influence pH of calcium hydroxide products. *J Dent Res.* 2014;94(Spec Iss C):193435.
- [68] Walsh LJ, Athanassiadis B. PEG 400 solvent alters pH in calcium hydroxide endodontic medicaments. *J Dent Res.* 2016;96(Spec Iss C):1705.

- [69] Walsh LJ, Athanassiadis B. Alkaline compositions and their dental and medical use. US patent publication US 2014/0322144-A1.
- [70] Teoh YY, Athanassiadis B, Walsh LJ. The influence of aqueous and PEG 400 solvent vehicles on hydroxyl ion release from calcium hydroxide medicaments. *Internat Dent*. 2016;11:42-50.
- [71] Ikai H, Nakamura K, Shirato M, Kanno T, Iwasawa A, Sasaki K, Kohno M. Photolysis of hydrogen peroxide, an effective disinfection system via hydroxyl radical formation. *Antimicrob Agent Chemother*. 2010;54:5086-91.
- [72] De Moor RJG, Verheyen J, Diachuk A, Verheyen P, Meire MA, De Coster PJ, Walsh LJ. Insight in the chemistry of laser-activated dental bleaching. *Scientif World J*. 2015;2015:650492.
- [73] Walsh LJ. Safety issues relating to the use of hydrogen peroxide in dentistry. *Aust Dent J*. 2000;45:257-69.
- [74] Neyens E, Baeyens J. A review of classic Fenton's peroxidation as an advanced oxidation technique. *J Hazard Mater*. 2003;98:33-50.
- [75] Walsh LJ. Photoactivated disinfection (PAD) and endodontics. *Australas Dent Pract*. 2003;14:88.
- [76] Ibi H, Hayashi M, Yoshino F, Tamura M, Yoshida A, Kobayash Y, Ogiso B. Bactericidal effect of hydroxyl radicals generated by the sonolysis and photolysis of hydrogen peroxide for endodontic applications. *Microbial Pathogen*. 2017;103:65-70.
- [77] Walsh LJ. 'Radical' progress: ozone-boosted in-office power whitening. *ADA News Bull*. 2006;341:26-8.
- [78] Case PD, George R, Bird PS, Walsh LJ. Treatment of root canal biofilms of *Enterococcus faecalis* with ozone gas and passive ultrasound activation. *J Endod*. 2012;38:523-6.
- [79] Kobayashi Y, Hayashi M, Yoshino F, Tamura M, Yoshida A, Ibi H, Ogiso B. Bactericidal effect of hydroxyl radicals generated from a low concentration hydrogen peroxide with ultrasound in endodontic treatment. *J Clin Biochem Nutr*. 2014;54:161-5.
- [80] George R, Meyers IA, Walsh LJ. Laser activation of endodontic irrigants with improved conical laser fiber tips for removing smear layer in the apical third of the root canal. *J Endod*. 2008;34:1524-7.
- [81] George R, Chan K, Walsh LJ. Laser-induced agitation and cavitation from proprietary honeycomb tips for endodontic applications. *Lasers Med Sci*. 2015;30:1203-8.
- [82] Hmud R, Kahler WA, Walsh LJ. Temperature changes accompanying infrared diode laser treatment of wet canals. *J Endod*. 2010;36:908-11.
- [83] George R, Walsh LJ. Thermal effects from modified endodontic laser tips used in the apical third of root canals with erbium-doped yttrium aluminium garnet and erbium,

chromium-doped yttrium scandium gallium garnet lasers. *Photomed Laser Surg.* 2010;28:161-5.

- [84] Hmud R, Kahler WA, George R, Walsh LJ. Cavitation effects in aqueous endodontic irrigants generated by near infrared lasers. *J Endod.* 2010;36:275-8.
- [85] George R, Walsh LJ. Laser fiber-optic modifications and their role in endodontics. *J Laser Dent.* 2012;20:24-30.
- [86] Lagemann M, Chai L, George R, Walsh LJ. Activation of ethylenediaminetetraacetic acid by a 940 nm diode laser for enhanced removal of smear layer. *Aust Endod J.* 2014;40:72-5.
- [87] Walsh LJ. The role of natural and synthetic phenolic compounds in home oral health care. *Australas Dent Pract.* 2004;6:22-5.
- [88] Lei L, Shao M, Yang Y, Mao M, Yang Y, Hu T. Exopolysaccharide dispelled by calcium hydroxide with volatile vehicles related to bactericidal effect for root canal medication. *J Appl Oral Sci.* 2016;24:487-95.
- [89] Shrestha A, Kishen A. Antibacterial nanoparticles in endodontics: a review. *J Endod.* 2016;42:1417-26.
- [90] Samiei M, Farjami A, Dizaj SM, Lotfipour F. Nanoparticles for antimicrobial purposes in endodontics: a systematic review of in vitro studies. *Mater Sci Eng C Mater Biol Appl.* 2016;58:1269-78.
- [91] Reece JB, Meyers N, Urry LA, Cain ML, Wasserman SA, Minorsky PV, Jackson RB, Cooke BN. *Campbell Biology*, 8th edn. San Francisco: Pearson Benjamin Cummings, 2009.
- [92] Kishen A. Nanoparticles for endodontic disinfection. In: Kishen A (Ed.) *Nanotechnology in Endodontics: Current and Potential Clinical Applications*. Switzerland: Springer International Publishing, 2015. pp. 97-119.
- [93] de Paz LEC, Resin A, Howard KA, Sutherland DS, Wejse PL. Antimicrobial effect of chitosan nanoparticles on *Streptococcus mutans* biofilms. *Appl Env Microbiol.* 2011;77:3892-5.
- [94] del Carpio-Perochena A, Kishen A, Shrestha A, Bramante CM. Antibacterial properties associated with chitosan nanoparticle treatment on root dentin and 2 types of endodontic sealers. *J Endod.* 2015;41:1353-8.
- [95] Shrestha A, Kishen A. Antibiofilm efficacy of photosensitizer-functionalized bioactive nanoparticles on multispecies biofilm. *J Endod.* 2014;40:1604-10.
- [96] Pagonis TC, Chen J, Fontana CR, Devalapally H, Ruggiero K, Song X, Yamazaki H. Nanoparticle-based endodontic antimicrobial photodynamic therapy. *J Endod.* 2010;36:322-8.
- [97] Louwakul P, Saelo A, Khemaleelakul S. Efficacy of calcium oxide and calcium hydroxide nanoparticles on the elimination of *Enterococcus faecalis* in human root dentin. *Clin Oral Invest.* 2016; Apr 30. [Epub ahead of print].

- [98] Samanta A, Chanda DK, Das PS, Ghosh J, Mukhopadhyay AK, Dey A. Synthesis of nano calcium hydroxide in aqueous medium. *J Am Ceram Soc.* 2016;99:787-95.
- [99] Chernousova S, Epple M. Silver as antibacterial agent: ion, nanoparticle, and metal. *Angew Chem Int Ed Eng.* 2013;52:1636-53.
- [100] Maddinedi SB, Mandal BK, Anna KK. Environment friendly approach for size control-able synthesis of biocompatible silver nanoparticles using diastase. *Environ Toxicol Pharmacol.* 2016;49:131-6.
- [101] AbdelRahim K, Mahmoud SY, Ali AM, Almaary KS, Mustafa AE, Husseiny SM. Extracellular biosynthesis of silver nanoparticles using *Rhizopus stolonifer*. *Saudi J Biol Sci.* 2017;24:208-16.
- [102] Laroo H. Colloidal nano silver- its production method, properties, standards and its bio-efficacy as an inorganic antibiotic. *J Phys Chem Biophys.* 2013;3:130.
- [103] Knight GM, McIntyre JM, Craig GG, Mulyani. Ion uptake into demineralized dentine from glass ionomer cement following pretreatment with silver fluoride and potassium iodide. *Aust Dent J.* 2006;51:237-41.
- [104] Deutsch A. An alternate technique of care using silver fluoride followed by stan-nous fluoride in the management of root caries in aged care. *Spec Care Dentist.* 2016;36:85-92.
- [105] Takamiya AS, Monteiro DR, Bernabé DG, Gorup LF, Camargo ER, Gomes-Filho JE, Oliveira SH, Barbosa DB. In vitro and in vivo toxicity evaluation of colloidal silver nanoparticles used in endodontic treatments. *J Endod.* 2016;42:953-60.
- [106] Zhang FH, Li M, Wei ZJ, Zhao B. The effect of a combined nanoparticulate/calcium hydroxide medication on the biofilm of *Enterococcus faecalis* in starvation phase. *Shanghai Kou Qiang Yi Xue.* 2016;25:11-5.
- [107] Afkhami F, Akbari S, Chiniforush N. *Entrococcus faecalis* elimination in root canals using silver nanoparticles, photodynamic therapy, diode laser, or laser-activated nanopar-ticles: an in vitro study. *J Endod.* 2016 Dec 24. pii: S0099-2399(16)30582-9.
- [108] PathiranaRD, O'Brien-SimpsonNM, VeithPD, RileyPF, ReynoldsEC. Characterizationof proteinase-adhesin complexes of *Porphyromonas gingivalis*. *Microbiol.* 2006;152:2381-94.
- [109] Gutknecht N, van Gogswaardt D, Conrads G, Apel C, Schubert C, Lampert F. Diode laser radiation and its bactericidal effect in root canal wall dentin. *J Clin Laser Med Surg.* 2000;18:57-60.
- [110] Beer F, Buchmair A, Wernisch J, Georgopoulos A, Moritz A. Comparison of two diode lasers on bactericidity in root canals – an in vitro study. *Lasers Med Sci.* 2012;27:361-4.
- [111] Walsh LJ. The current status of low level laser therapy in dentistry. Part 1. Soft tissue applications. *Aust Dent J.* 1997;42:247-54.
- [112] Walsh LJ. The current status of low level laser therapy in dentistry. Part 2. Hard tissue applications. *Aust Dent J.* 1997;42:302-6.

- [113] Caccianiga G, Baldoni M, Ghisalberti CA, Paiusco A. A preliminary in vitro study on the efficacy of high-power photodynamic therapy (HLLT): comparison between pulsed diode lasers and superpulsed diode lasers and impact of hydrogen peroxide with controlled stabilization. *BioMed Res Int.* 2016;2016:1386158.
- [114] Diogo P, Goncalves T, Palma P, Santos JM. Photodynamic antimicrobial chemotherapy for root canal system asepsis: a narrative literature review. *Int J Dent.* 2015;2015:269205.
- [115] Walsh LJ. The current status of laser applications in dentistry. *Aust Dent J.* 2003;48:146-55.
- [116] George S, Kishen A. Augmenting the antibiofilm efficacy of advanced noninvasive light activated disinfection with emulsified oxidizer and oxygen carrier. *J Endod.* 2008;34:1119-23.
- [117] Lee MT, Bird PS, Walsh LJ. Photo-activated disinfection of the root canal: a new role for lasers in endodontics. *Aust Endod J.* 2004;30:93-8.
- [118] Siddiqui SH, Awan KH, Javed F. Bactericidal efficacy of photodynamic therapy against *Enterococcus faecalis* in infected root canals: a systematic literature review. *Photodiagnosis Photodyn Ther.* 2013;10:632-43.
- [119] Arneiro RA, Nakano RD, Antunes LA, Ferreira GB, Fontes K, Antunes LS. Efficacy of antimicrobial photodynamic therapy for root canals infected with *Enterococcus faecalis*. *J Oral Sci.* 2014;56:277-85.
- [120] Tennert C, Feldmann K, Haamann E, Al-Ahmad A, Follo M, Wrbas KT, Hellwig E, Altenburger MJ. Effect of photodynamic therapy (PDT) on *Enterococcus faecalis* bio-film in experimental primary and secondary endodontic infections. *BMC Oral Health.* 2014;14:132.
- [121] Silva LAB, Novaes AB, de Oliveira RR, Nelson-Filho P, Santamaria M, Silva RAB. Antimicrobial photodynamic therapy for the treatment of teeth with apical periodontitis: a histopathological evaluation. *J Endod.* 2012;38:360-6.
- [122] Bonsor S, Nichol R, Reid T, Pearson G. An alternative regimen for root canal disinfection. *Brit Dental J.* 2006;201:101-5.
- [123] Bonsor S, Nichol R, Reid T, Pearson G. Microbiological evaluation of photo-activated disinfection in endodontics (an in vivo study). *Brit Dent J.* 2006;200:337-41.
- [124] Trindade AC, De Figueiredo JAP, Steier L, Weber JBB. Photodynamic therapy in endodontics: a literature review. *Photomed Laser Surg.* 2015;33:175-82.
- [125] Walsh LJ, George R. Surface structure modification. US patent 8,977,085. 2015.
- [126] Hislop J, Bird PS, Walsh LJ. Photoactivated disinfection of *Enterococcus faecalis* in bio-films. *Aust Dent J.* 2010;55 (Suppl):S22.
- [127] Tennert C, Drews AM, Walther V, Altenburger MJ, Karygianni L, Wrbas KT, Hellwig E, Al-Ahmad A. Ultrasonic activation and chemical modification of photosensitizers enhances the effects of photodynamic therapy against *Enterococcus faecalis* root-canal isolates. *Photodiagnosis Photodyn Ther.* 2015;12:244-51.

- [128] George S, Kishen A. Photophysical, photochemical, and photobiological characterization of methylene blue formulations for light-activated root canal disinfection. *J Biomed Optics*. 2007;12:034029.
- [129] Garcez AS, Núñez SC, Baptista MS, Daghastanli NA, Itri R, Hamblin MR, Ribeiro MS. Antimicrobial mechanisms behind photodynamic effect in the presence of hydrogen peroxide. *Photochem Photobiol Sci*. 2011;10:483-490.
- [130] Sainsbury AL, Bird PS, Walsh LJ. DIAGNOdent laser fluorescence assessment of endodontic infection. *J Endod*. 2009;35:1404-7.
- [131] Ho QV, George R, Sainsbury AL, Kahler WA, Walsh LJ. Laser fluorescence assessment of the root canal using plain and conical optical fibers. *J Endod*. 2010;36:119-22.
- [132] Walsh LJ. Applications of laser fluorescence for diagnosis of bacterial infections in the root canal. *Australas Dent Pract*. 2010;21:54-6.
- [133] Shakibaie F, George R, Walsh LJ. Applications of laser-induced fluorescence in dentistry. *Int J Dent Clin*. 2011;3:26029.
- [134] Walsh LJ. Fluorescence applications in dentistry: current status and future prospects. *Australas Dent Pract*. 2013;24:62-4.