We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Introductory Chapter: Chlorophyll Molecules and Their Technological Relevance

Maria Isabel Queiroz, Andrêssa Silva Fernandes,
Mariany Costa Deprá, Eduardo Jacob-Lopes and
Leila Queiroz Zepka

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/67953>

Chlorophylls were the major tool used, long before science was able to recognize the existence of plants and animals; our predecessors identified this distinction and slowly provided technological evidence that chlorophyll was responsible for the green color of the plants.

The name chlorophyll was proposed by Pierre Joseph Pelletier and Joseph Bienaimé Caventou in 1818 to designate the Green substance that could be extracted from the leaves with the aid of alcohol [1–3]. However, only about a century later, in 1911, the study of this interesting molecule gained more importance by Richard Willstätter, who obtained for the first-time pure chlorophyll and established its correct molecular formula $C_{55}H_{70}MgN_4O_6$ [2]. A year later, Willstätter showed that chlorophyll obtained from a wide variety of sources was a mixture of two compounds, chlorophyll-a and chlorophyll-b, to which he assigned the correct molecular formula $C_{55}H_{70}MgN_4O_5$ and $C_{55}H_{70}MgN_4O_6$, respectively. Nevertheless, the detailed understanding of the structures of these molecules was the result of the studies of Fischer and collaborators, who were the first to delineate the structure of porphyrin ring [2–5].

These compounds are complex organic molecules formed by derivatives of porphyrin, a macrocyclic structure, asymmetric, totally unsaturated. Basically, chlorophyll molecules are conjugated tetrapyrroles to which a cyclopentanone ring jointly with the third ring linked together by bridges methylene, with the central atom, magnesium (**Figure 1**). This structure also contains at the C-17, a propionic acid chain esterified with the phytol, diterpene alcohol [3, 6–8].

Chlorophyll-b differs from chlorophyll-a by the presence of aldehyde residue instead of the methyl group at position 7. The synthesis of this compound is given by the oxidation of the methyl group to aldehyde [9, 10] through the enzyme oxygenase, which catalyzes this conversion [11].

Structurally, the chlorophyll molecules differ from each other, due to the degree of saturation of the pyrrolic rings. For example, chlorophyll-c contains fully unsaturated phytylporphyrin (double alloys C17–C18), whereas the other chlorophylls are C17–C18, phytylchlorins (Figure 1). These differences in chlorophyll macrocycle saturation have profound consequences on the absorbance spectrum. For example, chlorophyll-a, chlorophyll-d, and chlorophyll-f phytylchlorins have approximately equal intensities of absorption in blue, red, and green. On the other hand, the phytylporphyrins of chlorophyll-c absorb themselves weakly in red and more intensely around 450 nm [6, 12, 13].

The spectral differences of the chlorophyll molecules are reflected in the key in which bodies are present that the chlorophyll-a appears more bluish green, chlorophyll-b bright green, chlorophyll-c yellowish green, chlorophyll d bright forest green, and chlorophyll-f emerald green [6]. This diversity of shades of green makes these organisms potential source of natural colorants [14].

Figure 1. Structure of the side chain variables of chlorophyll molecules.

In addition, a chlorophyll molecule consists of two distinct parts: hydrophile, the macrocycle, and a hydrophobic part, phytol chain. The most hydrophilic segment of the macrocycle is the cyclopentanone ring and the propionic ester group (17 positions). The hydrophilic and hydrophobic character directly influences the choice of solvent for extraction, which, from a quantitative or qualitative point of view, is an extremely important aspect when it focuses on the technological relevance of chlorophyll molecules [15–17].

Given this scenario, analysis techniques such as liquid chromatography coupled to ultraviolet and mass spectrum allowed demonstrating that currently a total of five molecules, so-called chlorophyll, have been found in nature. All of them occur in the cyanobacterium, which is cosmopolitan microalgae, prokaryotic, Gram negative, with metabolism preferentially photoautotrophs, able to perform photosynthesis oxygen production, eukaryotic algae, and higher plants. These microorganisms demonstrate considerable biocatalytic potential in biotechnological processes due to their robustness and require nutritional simplicity [18]. Chlorophyll-a is present in all organisms that perform photosynthesis and chlorophyll-b oxygen (**Figure 1**), which are the most abundant pigment in green algae, is found in higher plants in nature, and other similar structures are used as accessory pigments in photosynthetic process, called chlorophyll-c, chlorophyll-d, chlorophyll-e, and chlorophyll-f [6–8, 13]. These specific pigments present in the cells of the photosynthetic organisms influence efficiency of dispersion of light, because each species has its own distinct characteristic pigment [9].

Chlorophyll-a is the most abundant and important structure of all chlorophylls, corresponding to approximately 75% of the green pigment found in nature. This molecule has bioactive properties with extensive use in pharmaceutical and food industry, is considered a high added value, and may reach high values on the market. The bioactive potential of the cells is associated with a complex structure of conjugated connections in pyrrolic rings, allowing an oxidant and antioxidant, acting without free radical sequestration [7]. In addition, it is related to nutraceutical properties, playing an important role in health, through its anti-inflammatory and vascular constrictor, which makes this molecule an important ingredient in food formulations, cosmetics, and drugs [16].

Another molecule that differs from chlorophyll-a, to present the first pyrrole ring, a substitution of the vinyl C3 group for the chlorophyll-a molecule for a formyl group, was called chlorophyll-d. This molecule was discovered 70 years ago, but only in 1996, an important property of this molecule was observed [6, 9, 13], performing a function equivalent to chlorophyll-a in oxygenic photosynthesis [7, 13]. The properties of chlorophyll-d, like its photosynthetic power associated with the difference in wavelength of absorption in relation to chlorophyll-a, make this a potential component molecule to be employed in a biorefinery. In the last years, it has been argued that there is a strong interest in the development of technologies for the production of natural molecules [16, 19, 20].

On the other hand, chlorophyll biomolecules are highly prone to change, because their chemical structure is rich in double bonds combined. In this sense, in order to increase stability, chlorophyll undergoes changes in its molecule, replacing the atom of Mg^{2+} for Cu^{2+} , resulting in the call copper chlorophyllin, which is stable and can be used in hydro- or lipo-soluble formulations [21].

In summary, the broad application of chlorophyll makes it a molecule of great importance for the global market. However, there is still doubt about its application due to its chemical instability. In this way, the extension of studies to elucidate the open gaps will be of extreme relevance to expand the scientific knowledge base and its relation to the industrial application.

Author details

Maria Isabel Queiroz¹, Andrêssa Silva Fernandes², Mariany Costa Deprá², Eduardo Jacob-Lopes² and Leila Queiroz Zepka^{2*}

*Address all correspondence to: zepkaleila@yahoo.com.br

1 School of Chemistry and Food, Federal University of Rio Grande, UFRGS, Rio Grande, RS, Brazil

2 Food Science and Technology Department, Federal University of Santa Maria (UFSM), Santa Maria, RS, Brazil

References

- [1] Pelletier, F.; Caventou, J. B. Sur la matièri verte dès feuilles. *Annales de C himieet de Physique*, 9, 194–196, 1818.
- [2] Gossauer, A.; Engel N. Chlorophyll catabolisms—Structures, mechanisms conversions. *Journal of Photochemistry and Photobiology*, 32, 141–151, 1996.
- [3] Schoefs, B. Chlorophyll and carotenoids analysis in food products properties of the pigments and methods of analysis. *Trends in food Science & Technology*, 13, 361–371, 2002.
- [4] Fischer, H.; Orth. H. Die chemie des pyrrols. Akaemie Verlagsgesellschaft, Vol. 2, Part 1, Leipzig, Germany, 1937.
- [5] Fischer, H.; Stern, A. Die chemies des pyrrols. Akademie Verlagsgesellschaft, Part 2, Leipzig, Germany, 1940.
- [6] Chen, M.; Blankenship, R. E. Expanding the solar spectrum used by photosynthesis. *Trends in Plant Science*, 16, 488–496, 2010.
- [7] Chen, M.; Yaqiong, L.; Birch, D.; Willows, R. D. A cyanobacterium that contains chlorophyll-f a red absorbing photopigment. *FFBS Letters*, 586, 3249–3254, 2012.
- [8] Fu, W.; Magnusdóttir, M; Palson, B. UPLC – UV – MS analysis for quantification and chlorophyll species in algae. *Analytical and Bioanalytical Chemistry*, 404, 3145–3154, 2012.
- [9] Harada, J.; Mizoguchi, T.; Tsukatani, Y.; Noguchi, M.; Tamiaki, H. A seventh Bacterial chlorophyll driving a large light-harvesting antenna. *Scientific Reports*, 2(671), 1–5, 2012.

- [10] Yen, H. W.; Hu, I. C.; Chen, C. Y.; Ho, S. H.; Lee, D. J.; Chang, J. Microalgae based biorefinaria – from biofuels to natural products. *Bioresource Technology*, 35, 166–174, 2013.
- [11] Xu, W.; Tang, H.; Wang, Y.; Chitins, P. Proteins of the cyanobacterial photosystem I. *Biochimica et Biophysica Acta*, 1507, 32–40, 2001.
- [12] Zapata, M.; Rodriguez, A.; Garrido, J. L. Separation of chlorophyll and carotenoids from marine phytoplankton a new HPLC method using a reversed phase C8 column and pyridine containing mobile phases. *Marine Ecology Progress Series*, 195, 29–45, 2000.
- [13] Loughlin, P.; Lin, Y.; Chen, M. Chlorophyll d and *Acaryochloris marina*: current status. *Photosynthesis Research*, 116(2–3), 277–293, 2013.
- [14] Pelayo, R. D.; Galhardo-Guerreiro, L.; Homero, M. Chlorophyll and carotenoid pigments in the peel and flesh of commercial Apple fruit varieties. *Food Research International*, 65, 272–281, 2014.
- [15] Rosseló-Soto, E.; Galanakis, C. M.; Brncic, C. M.; Orlens, V. Trugillo, F. J. R.; Mawson, R.; Barba, F. J. Clean recovery of antioxidant compounds from plants food, byproducts and algae assisted by ultrasound processing. Modelling approaches to optimize processing conditions. *Trends in Foods Science & Technology*, 42, 134–149, 2015.
- [16] Fernandes, A.; Nogara, G.; Menezes, C. R.; Cichoski, A. J.; Mercadante, A. Z.; Jacob-Lopes, E.; Zepka, L. Q. Identification of chlorophyll molecules with peroxy radical scavenger capacity in microalgae *Phormidium autumnale* using ultrasound-assisted extraction. *Food Research International*, *In Press*, 2016.
- [17] Streit, N.; Merida, L. G. R.; Zepka, L. Q.; Jacob-Lopes, E.; Queiroz, M. I. Production de pigmentos por *Aphanotheca microscopica* nãglia partir de resíduos industrialis lácteos. *Ingeniare Revista Chilena de Ingenieria*, 25, 11–18, 2017.
- [18] Guiry, M. D.; Guiry, G. M. *Algae Base*: world-wide electronic publication National University of Ireland. Available at: <http://www.algaebase.org>. Accessed Oct 13, 2016.
- [19] Queiroz, M. I.; Hornes, M. O.; Silva-Manetti, A. G.; Zepka, L. Q.; Jacob-Lopes, E. Fish processing wastewater a platform of the microalgal biorefineries. *Biosystem Engineering*, 115, 95–202, 2013.
- [20] Kim, S.; Kim, M. S.; Lee, B. Y.; Lee, P. Generation of structurally novel short carotenoids and study of their biological activity. *Scientific Reports*, 6, 1–11, 2016.
- [21] Sant’Anna, V.; Gurak, P. D.; Marczak, L. D. F.; Tessaro, I. C. Tracking bioactive compounds with colour changes in foods–A review. *Dyes and Pigments*, 98(3), 601–608, 2013.