We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Chlorophyll-a and the Supply Side Ecology: Lessons from the Rocky Shores

Ana Carolina de Azevedo Mazzuco and
Paula Kasten

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/68044>

Abstract

The aims of this study were to summarize and describe the influences of phytoplankton on the larval cycle of rocky shore invertebrates, and to assess the relationship between fluctuations in chlorophyll-a concentration and the rates of larval processes. We carried out a mini review of the published data regarding the theme of the chapter, in which we described the ecological trends for the most common taxa and key species at small and larger spatiotemporal scales. The following topics were addressed: (i) the influence of phytoplankton on larval development, rhythms of larval release, larval quality, larval transport, settlement, and recruitment; (ii) the relationships between variations in chlorophyll-a concentration and the rates of larval processes; (iii) climate change on phytoplankton larva dynamics. The information presented here highlights the role of phytoplankton on rocky shore communities, as well as the importance of chlorophyll-a as a tool for modeling and forecasting the supply side ecology in rocky shore communities.

Keywords: phytoplankton, chlorophyll-a, supply side ecology, marine invertebrates, rocky shores, benthic-pelagic coupling

1. Introduction

Larval supply is the main source of new individuals to the populations of rocky shore invertebrates [1–3]. In these communities, larval success regulates how energy is transferred through the trophic web [4–6]; consequently, variations in the supply of propagules are the basis of trophic interactions at rocky shores [7, 8]. Since phytoplankton is the main food source for planktonic larvae of marine invertebrates [9], variations in phytoplankton biomass and diversity have significant influences on the larval cycle. Larval responses to the variability in phytoplankton abundance and diversity are species-specific. Larval fitness is influenced by environmental conditions experienced by adults

and larvae [10, 11]. The effects of phytoplankton on larval dynamics depend on the phase of larval development [12–15] and may be stronger when variations in phytoplankton occur on temporal scales that larvae or breeding adults are able to respond [16]. The direct interaction between phytoplankton and the larval stages have short-term consequences for larval dynamics (e.g., Ref. [14]), and it might have long-term effects as well. Because of that, variations in the rates of the ecological processes of rocky shore invertebrates are commonly correlated with fluctuations in chlorophyll-a concentration in the ocean (e.g., Refs. [17–20]). These numerical relationships are important tools to ecological modeling, and may be used to improve stock management in some extent [21].

2. The role of phytoplankton blooms in reproduction timing and in the rhythms of larval release

In the rocky shore communities, filter feeders depend greatly on phytoplankton as their main source of food and its consumption results in energy for growth and reproduction [22]. It is common to find larger animals with higher fecundity rates at rocky shores located in areas of high primary productivity, as a response to the higher concentrations of phytoplankton, and thus, food availability [19, 23–25]. Different types of phytoplankton present distinct physiological qualities as food particles [26], thereby both the amount of phytoplankton in the water column and their diversity influence the reproductive traits in marine invertebrates.

But not only adults on the rocky shore depend on phytoplankton in order to survive, larvae produced by those organisms also rely on these microorganisms to develop and reach the juvenile phase [27]. As evolution drives maximum reproductive activity to happen when environmental conditions are the best for offspring development, food availability is one of the most important factors regulating reproduction and allowing adults to produce viable offspring. Thus, it is common to observe peaks of larval release by rocky shore invertebrates synchronized with phytoplankton blooms (e.g., Refs. [28, 29]). Some metabolites produced by phytoplankton are signs of favorable environmental conditions for the larval development, triggering the spawning activity of green sea urchins and blue mussels, for instance Ref. [28]. These animals perceive such chemical compounds as an indication of good food abundance, so synchronizing the timing of larval release with high abundance of phytoplankton would promote higher offspring survival. Barnacles, on the other hand, just need a physical contact with phytoplankton cells to trigger their spawning activity, and larger the phytoplankton cell is, the stronger is the response [28].

Therefore, the presence of phytoplankton may overcome other environmental factors in the regulation of reproduction timing and larval release [30]. Spring and summer are the main reproductive periods for rocky shore invertebrates at temperate and upwelling regions [31], as it is during these seasons that phytoplankton blooms occur. Mussels from the Baltic sea, for example, start to develop their gonads when temperature starts to drop in the beginning of winter; but its maturation and ripening processes proceed in a way that the animals are ready to reproduce at the same time that phytoplankton blooms occur in the beginning of spring [32]. Some barnacles are even able to maintain their fully developed nauplii in the mantle

cavity until a high abundance of phytoplankton is perceived by the adults and only then, the nauplii will be released, a strategy that enhances the offspring survival due to the higher chance of facing a favorable feeding environment [33].

Similar reproductive timing was registered in the Indian coasts, where phytoplankton blooms occur during the monsoons and barnacles spawn their nauplii short after a break of the monsoon conditions [34]. However, these are not the best conditions for nauplii development, as these breaks stop and unfavorable monsoon conditions for larval development return soon after. Such misleading cue could result in lower recruitment rates for barnacles in this region. In subtropical coasts, peaks of larval production in intertidal barnacles are also preceded by high concentrations of chlorophyll-a in the water column [35]. On the daily scale, phytoplankton diversity might be as important as biomass in the regulation of larval release [36]. The presence of phytoplankton may overcome other environmental factors known to act as synchronization cues for reproduction timing and larval release [30].

3. How do changes in phytoplankton affect larval development from release to competency?

As seen in the previous section, phytoplankton has an important role in the reproductive success of marine invertebrates inhabiting the rocky shores. Part of this reproductive success involves the survival of larvae up to the juvenile stage, and a successful return to the benthic habitat is essential to the maintenance of rocky shore populations [2, 37]. It is straightforward to think that larval development is strictly linked to changes in phytoplankton community, since these cells are the main food items for marine planktotrophic larvae [9]. Because of that, the physiological quality of a larva would be determined in the plankton during its development and influenced directly by the phytoplankton in the water column. However, phytoplankton may change larval physiological quality much before that same larva is produced, through maternal effects, that is, when maternal individuals have the capacity to perceive the environment and manipulate the energy allocated for propagule production [38].

The amount of energetic reserves allocated to each propagule produced depends on the amount of energy the maternal individual can provide to its offspring. This capacity, in turn, is limited by the food available for the mothers, their perception of it, and their competency to gather and assimilate energy [38, 39]. For those marine organisms that produce lecithotrophic larvae, maternal effects are extremely important for shaping larval physiological quality because these larvae depend exclusively on the energetic resources from embryogenesis to survive [40]. If food ration is low, mothers can either preserve the energy acquired for their own metabolism and produce lower quality larvae (a selfish strategy, Ref. [39]) or invest all energy possible into their propagules, enhancing the survival potential of that higher quality larvae (an anticipatory strategy, Ref. [39]). In a scenario where maternal individuals are feeding mainly on phytoplankton, as the majority of filter feeding invertebrates in the rocky shores are, it is possible to understand the effect that oscillations in the quantity and type of phytoplankton available for these animals to feed has on larval quality.

However, most invertebrates that inhabit the rocky shores produce planktotrophic larvae. These larvae are submitted to transport and dispersion; they will feed in the plankton and will probably not experience the same conditions of the maternal environment, hypothetically reducing the necessity of energy transfer from mother to larvae. Thus, one could assume that the food environment experienced by mother would not impact the quality of the larvae produced. Interestingly, few authors have shown that, under stressful temperatures and low phytoplankton concentrations, maternal individuals of a tropical barnacle are able to manipulate the transfer of different types of fatty acids to their nauplii, a possible strategy to guarantee higher survival rates until this same nauplii encounters better food conditions in the water column [41]. Variations in the amount and type of phytoplankton available for planktotrophic larvae during development cycle interfere in the different larval traits, including in the success of metamorphosis into the juvenile stage. Larvae of gastropods [15, 16, 42], bivalves [36, 43, 44], and barnacles [45, 46] vary in size, development rate, and survival to the juvenile stage, in direct association with the quality and amount of phytoplankton offered them during their development.

Larvae must be able to survive from pelagic to benthic conditions and return to the rocky shore communities, in order to reach the adult phase. Settlement success and post-settlement survivorship are also matters of larval history [12, 15, 21], and many more. Contrary of what has been accepted for a long time, settlement of larvae in the benthic environment, and its metamorphose to the juvenile stage do not result in a “new beginning” for those individuals, but the feeding conditions experienced by larvae and its results on their physiological quality can be carried over to the next stage, and those individuals who faced low phytoplankton concentrations during its life in the plankton might become juveniles with lower growth and survival potential, influencing directly on the fate of that population [46–51].

4. Larval transport, settlement, and recruitment

Phytoplankton and larval abundances are sometimes controlled by the same oceanographic processes. Phytoplankton grows and reproduces under very specific environmental conditions, driven mainly by turbulence and nutrient availability [52]. Ocean movements, such as turbulence, vertical mixing, and currents, also affect larval abundance at small (e.g., Ref. [53]) and larger scales (e.g., Ref. [54]). Marine larvae take advantage of meso- and large-scale oceanographic features for transport and dispersion. These larvae have different responses depending on the velocity at that depth, assuming a specific swimming or orientation pattern (e.g., Ref. [55]). Besides, larvae are able to control their position in the water column and move together with the main current at that specific depth [56–58], what in turn might result in variability of larval supply in time and space [59]. Some oceanographic features that accumulate and transport marine invertebrate larvae are responsible for disturbing phytoplankton as well. For example, upwelling currents, which cause phytoplankton blooms by injecting cold nutrient-rich waters in the photic zone, may move larvae of rocky shore invertebrate to shallower waters (e.g., Refs. [60, 61]). Storms are other meteorological-oceanographic

phenomena that disturb both chlorophyll-a concentration at the nearshore environments (e.g., Ref. [62]) and the larval abundances close to the rocky shores [63].

Settlement is a function of larval supply [64]. Consequently, successful settlement relies on larvae, which need to find suitable settlement sites and be able to metamorphose. In this phase of the larval cycle, biochemical and physical cues either stimulate or block settlement. The presence of biofilm on the rocks is very important for settling larvae, in particular for the sessile larvae, because biofilm may define if that is a favorable settlement spot. Biofilm characteristics control larval behavior during settlement [65]; as a result, settlement rates and the chlorophyll-a content in the biofilm are correlated [66]. Settlement may also be correlated with fluctuations in chlorophyll-a concentration just as a consequence of the coupling between phytoplankton blooms and larval release [12, 28]. When the latter situation is true, fluctuations in chlorophyll-a concentration and variations in settlement rates are time lagged in several days [35], what may depend on the time that the larva takes to fully develop. On the other hand, if larval supply and phytoplankton dynamics are controlled by the same features, as it was explained in the previous paragraph, peaks in chlorophyll-a concentrations and settlement rates will occur simultaneously (e.g., Ref. [20]).

Recruitment rates are regulated by fluctuations in the pelagic environment affecting larval supply [67]. Recruitment success means that settled larvae survived until they are able to reproduce. In the post-settlement period, phytoplankton availability in the benthos and pelagic can control the survivorship of settlers in rocky shore communities. Although most early recruits of rocky shore invertebrates are filter feeders, they do not have the same diet and they may be very selective [68], choosing determinate phytoplankton species as food items depending on their size. Changes in the phytoplankton community might benefit one or the other species depending on their feeding behavior [68]. Although the relationship between recruitment and chlorophyll-a concentration is influenced by species-specific characteristics, information on this subject is still relatively scarce for rocky shore invertebrates. Small- and large-scale spatial variability in recruitment of rocky shore invertebrates are related to local and regional gradients of chlorophyll-a concentration in the surface waters. Geographic barriers that restrict phytoplankton abundance are also responsible for setting geographical limits for recruitment at the rocky shores. Recruitment rates may vary in several orders of magnitude among regions and sites, potentially due to persistent gradients in phytoplankton availability, and in turn gradients in chlorophyll-a concentration (e.g., Refs. [69, 70]. Even sites within the same bay or just less than 1 km apart may present high contrasts in recruitment rates as a consequence of differences in the phytoplankton dynamics [71].

5. The numerical relationships between chlorophyll-a concentration and larval processes

Phytoplankton is a limiting resource to the survival of marine invertebrate larvae, as it was described throughout the chapter; consequently, chlorophyll-a concentration is a key factor

regulating larval dynamics in rocky shore communities. Variations in larval processes and fluctuations in chlorophyll-a concentration tend to be highly correlated (e.g., trends of recruitment rates [69]). These correlations could be incorporated to ecological and numerical models to predict larval processes based on the values of chlorophyll-a concentration in the water (e.g., Ref. [72]). Although there are daily measurements of chlorophyll-a concentration in the ocean surfaces at a global scale, the levels of correlation between chlorophyll-a and larval dynamics are described only for a few species and some coastal areas.

Trends may be divided in groups according to the relationship between larval and phytoplankton dynamics. If the oceanographic processes promoting larval supply and settlement are also responsible for enabling phytoplankton growth and reproduction, variations in larval processes and in chlorophyll-a concentration may be positively correlated. On the other hand, if larval supply and settlement are enabled by less favorable conditions for phytoplankton, the fluctuations in the rates of larval processes may be negatively related to the concentrations of chlorophyll-a. Evidences of both trends were registered for rocky shore invertebrates in several regions [20, 21, 73]. Although the oceanographic and ecological processes that affect community dynamics are similar at the rocky shores, the correlation degrees between phytoplankton abundance and larval processes vary among sites and taxa. Correlations are stronger when reproduction and larval processes are regulated by the same mechanisms controlling phytoplankton blooms. For instance, in upwelling regions, these correlations are expected to be stronger [74], but may not be significant depending on the site (e.g., Ref. [75]). Barnacle and mussel recruits that occupy the same intertidal zone are not necessarily affected by fluctuations in chlorophyll-a concentration in similar ways, even presenting opposite trends in recruitment [21].

6. Climate change on phytoplankton larval dynamics

Climate change has important consequences for benthic-pelagic dynamics. Global warming has already caused alterations in the patterns of sea surface temperature and ocean currents, which in turn directly influenced the trends of phytoplankton abundance. Larvae and recruits of rocky shore invertebrates have to cope with such alterations in food availability concomitant to other climatic changes. The effects of phytoplankton and other climatic factors, such as water temperature, tend to be synergic [76]. Global warming conditions might not be positive for marine invertebrate larvae which, on one hand, survive under a wide range of conditions, but their fitness is highly influenced by changes in food availability. Short- and long-term consequences of climate change on phytoplankton larval dynamics were already detected for rocky shore communities. On the scale of decades, longer events of upwelling in the recent 20 years doubled the recruitment rates in some shores [77]. Results showed that, in small scale conditions, variability in phytoplankton has different effects on larval performance under different levels of climate change (Kasten, personal communication). However, how species will respond to multiple factors under *in situ* oceanic climatic conditions are hard to forecast, since information in larval dynamics are not available for most species and rocky shore systems.

7. Final considerations

Phytoplankton has a high regulatory potential in larval dynamics in the rocky shore communities. Rates of larval processes in rocky shore invertebrates are highly correlated with spatiotemporal fluctuations in chlorophyll-a concentration in the sea surfaces. The role of phytoplankton in larval dynamics at the community levels is not known, because information for most species is incipient. It is important to highlight that scientific improvements are needed to allow that use of variations chlorophyll-a concentration as a tool for modeling and forecasting the supply side ecology in rocky shore communities.

Author details

Ana Carolina de Azevedo Mazzuco^{1,2*} and Paula Kasten²

*Address all correspondence to: ac.mazzuco@me.com

1 Federal University of Espírito Santo, Dept. of Oceanography, Vitória, ES, Brazil

2 Federal University of São Paulo, Institute of Ocean Sciences, Santos, SP, Brazil

References

- [1] Roughgarden J, Iwasa Y, Blaxter C (1985) Demographic theory for an open population with space-limited recruitment. *Ecology* 66:54–67. doi: 10.2307/1941306
- [2] Roughgarden J, Gaines S, Possingham H (1988) Recruitment dynamics in complex life cycles. *Science* 241:1460–1466. doi: 10.1126/science.11538249
- [3] Gaines S, Bertness M (1992) Dispersal of juveniles and variable recruitment in sessile marine species. *Nature* 360:579–580. doi:10.1038/360579a0
- [4] Bustamante RH, Branch GM, Eekhout S, Robertson B, Zou-tendyk P, Schleyer M, Dye A, Hanekom N, Keats D, Jurd M, McQuaid C (1995) Gradients of intertidal productivity around the coast of South Africa and their relationships with consumer biomass. *Oecologia* 102:189–201. doi: 10.1007/BF00333251
- [5] Bustamante RH, Branch GM, and Eekhout S (1995) Maintenance of an exceptional intertidal grazer biomass in South Africa: subsidy by subtidal kelps. *Ecology* 76:2314–2329. doi: 10.2307/1941704
- [6] Nielsen KG, Navarrete SA (2004) Mesoscale regulation comes from the bottom-up: intertidal interactions between consumers and upwelling. *Ecol Lett* 7:31–41. doi: 10.1046/j.1461-0248.2003.00542.x
- [7] Menge BA, Lubchenco J, Bracken MES, Chan F, Foley MM, Freidenburg TL, Gaines SD, Hudson G, Krenz C, Leslie H, Menge DNL, Russell R, Webster MS (2003) Coastal

- oceanography sets the pace of rocky intertidal community dynamics. *Proc Natl Acad Sci USA* 100 (21):12229–12234. doi: 10.1073/pnas.1534875100
- [8] Navarrete SA, Wieters EA, Broitman BR, Castilla JC (2005) Scales of benthic–pelagic coupling and the intensity of species interactions: from recruitment limitation to top down control. *Proc Natl Acad Sci USA* 102:18046–18051
- [9] Thorson G (1950) Reproductive and larval ecology of marine bottom invertebrates. *Biol Rev* 25:1–45. doi: 10.1111/j.1469-185X.1950.tb00585.x
- [10] Giménez L (2006) Phenotypic links in complex life cycles: conclusions from studies with decapod crustaceans. *Integr Comp Biol* 46:615–622. doi: 10.1093/icb/icl010
- [11] Shima J, Swearer S (2009) Larval quality is shaped by matrix effects: implications for connectivity in a marine metapopulation. *Ecology* 90:1255–1267. doi: 10.1890/08-0029.1
- [12] Basch L, Pearse J (1995) Consequences of larval feeding environment for settlement and metamorphosis of a temperate echinoderm. *Oceanol Acta* 19:273–285.
- [13] Pechenik J, Estrella M, Hammer K (1996) Food limitation stimulates metamorphosis of competent larvae and alters postmetamorphic growth rate in the marine prosobranch gastropod *Crepidula fornicata*. *Mar Biol* 127:267–275. doi: 10.1007/BF00942112
- [14] Anil A, Desai D, Khandeparker L (2001) Larval development and metamorphosis in *Balanus amphitrite* Darwin (Cirripedia; Thoracica): significance of food concentration, temperature and nucleic acids. *J Exp Mar Bio Ecol* 263:125–141. doi: 10.1016/S0022-0981(01)00280-5
- [15] Pechenik J, Tyrell A (2015) Larval diet alters larval growth rates and post-metamorphic performance in the marine gastropod *Crepidula fornicata*. *Mar Biol* 162(8):1597–1610. doi: 10.1007/s00227-015-2696-7
- [16] Pechenik J, Jarrett J, Rooney J (2002) Relationships between larval nutritional experience, larval growth rates, juvenile growth rates, and juvenile feeding rates in the prosobranch gastropod *Crepidula fornicata*. *J Exp Mar Bio Ecol* 280:63–78. doi: 10.1016/S0022-0981(02)00367-2
- [17] Menge B (1992) Community regulation: under what conditions are bottom-up factors important on rocky shores? *Ecology* 73:755–765. doi: 10.2307/1940155
- [18] Gaines S, Hudson G, Krenz C, et al (2003) Coastal oceanography sets the pace of rocky intertidal community dynamics. *Proc Natl Acad Sci USA* 100:12229–12234. doi: 10.1073/pnas.1534875100
- [19] Leslie H, Breck E, Chan F, et al (2005) Barnacle reproductive hotspots linked to near-shore ocean conditions. *Proc Natl Acad Sci USA* 102:10534–10539. doi: 10.1073/pnas.0503874102
- [20] Burrows MT, Jenkins SR, Robb L, Harvey R (2010) Spatial variation in size and density of adult and post-settlement *Semibalanus balanoides*: effects of oceanographic and local conditions. *Mar Ecol Prog Ser* 398:207–219. doi: 10.3354/meps08340

- [21] Mazzuco, ACA, Pineda, J, Starczak, VC, Christofolletti, RA, Ciotti, AM (2015) Temporal variation in intertidal community recruitment and its relationships to physical forcings, chlorophyll-a concentration and sea surface temperature. *Mar Biol* 162:1705–1725. doi: 10.1007/s00227-015-2689-6
- [22] Menge B, Daley B, Wheeler P, Strub P (1997) Rocky intertidal oceanography: an association between community structure and nearshore phytoplankton concentration. *Limnol Oceanogr* 42:57–66. doi: 10.4319/lo.1997.42.1.0057
- [23] Page HM, Hubbard DM (1987) Temporal and spatial patterns of growth in mussels *Mytilus edulis* on an offshore platform: relationships to water temperature and food availability. *J Exp Mar Bio Ecol* 111:159–179. doi: 10.1016/0022-0981(87)90053-0
- [24] Sanford E, Menge B (2001) Spatial and temporal variation in barnacle growth in a coastal upwelling system. *Mar Ecol Prog Ser* 209:143–157. doi: 10.3354/meps209143
- [25] Figueiras FG, Labarta U, Fernández Reiriz MJ (2002) Coastal upwelling, primary production and mussel growth in the Rías Baixas of Galicia. *Hydrobiologia* 484:121–131. doi: 10.1023/A
- [26] Stone C (1989) A comparison of algal diets for cirripede nauplii. *J Exp Mar Bio Ecol* 132: 17–40. [https://doi.org/10.1016/0022-0981\(89\)90174-3](https://doi.org/10.1016/0022-0981(89)90174-3)
- [27] Olson R, Olson M (1989) Food limitation of planktotrophic marine invertebrate larvae: does it control recruitment success? *Annu Rev Ecol Syst* 20:225–247.
- [28] Starr M, Himmelman J, Therriault J (1991) Coupling of nauplii release in barnacles with phytoplankton blooms: a parallel strategy to that of spawning in urchins and mussels. *J Plankton Res* 13:561–571. doi: 10.1093/plankt/13.3.561
- [29] Bueno M, Moser G, Tocci B, Flores A (2010) Retention-favorable timing of propagule release in barnacles and periwinkles. *Mar Ecol Prog Ser* 414:155–165. doi: 10.3354/meps 08715
- [30] Kasten P, Flores A (2013) Disruption of endogenous tidal rhythms of larval release linked to food supply and heat stress in an intertidal barnacle. *Mar Ecol Prog Ser* 472:185–198. doi: 10.3354/meps10005
- [31] Hines AH (1979) The comparative reproduction ecology of three species of intertidal barnacles. In: Stancyk S.E., ed. *Reproductive ecology of marine invertebrates*. pp. 213–234. Belle Baruch Library In Marine Sciences, Univ. S. Carolina Press, Columbia.
- [32] Kautsky N (1982) Quantitative studies on gonad cycle, fecundity, reproductive output and recruitment in a baltic *Mytilus edulis* population. *Mar Biol* 68:143–160. doi: 10.1007/BF00397601
- [33] Barnes H (1962) Note on variations in the release of nauplii of *Balanus balanoides* with special reference to the spring diatom outburst. *Crustaceana* 4:118–122.
- [34] Desai D, Anil A (2005) Recruitment of the barnacle *Balanus amphitrite* in a tropical estuary: implications of environmental perturbation, reproduction and larval ecology. *J Mar Biol Assoc UK* 85:909–920. doi: 10.1017/S0025315405011884

- [35] Barbosa A, Gomes C, Pereira G, et al (2016) Local biological drivers, not remote forcing, predict settlement rate to a subtropical barnacle population. *Mar Ecol Prog Ser* 543:201–208. doi: 10.3354/meps11589
- [36] Toupoint N, Gilmore-Solomon L, Bourque F, et al (2012) Match/mismatch between the *Mytilus edulis* larval supply and seston quality: effect on recruitment. *Ecology* 93:1922–1934. doi: 10.1890/11-1292.1
- [37] Pineda J, Reynolds N, Starczak V (2009) Complexity and simplification in understanding recruitment in benthic populations. *Popul Ecol* 51:17–32. doi: 10.1007/s10144-008-0118-0
- [38] Bernardo J (1996) Maternal effects in animal ecology. *Integr Comp Biol* 36 (2): 83–105. <https://doi.org/10.1093/icb/36.2.83>
- [39] Marshall J, Uller T (2007) When is a maternal effect adaptive? *Oikos* 116:1957–1963. doi: 10.1111/j.2007.0030-1299.16203.x
- [40] McEdward LR (1995), Jaeckle WB (1995) Variation in the size, energy content, and biochemical composition of invertebrate eggs: correlates to the mode of larval development. In: McEdward L (ed) *Ecology of marine invertebrate larvae*. CRC Press, Boca Raton, FL, p 49–77.
- [41] Freuchet F, Tremblay R, Flores A (2015) Interacting environmental stressors modulate reproductive output and larval performance in a tropical intertidal barnacle. *Mar Ecol Prog Ser* 532:161–175. doi: 10.3354/meps11377
- [42] McCann M, Padilla D (2015) Effects of a patchy food environment across life history stages. *J Exp Mar Bio Ecol* 472:135–141. doi: 10.1016/j.jembe.2015.07.009
- [43] Pernet F, Tremblay R, Langdon C, Bourget E (2004) Effect of additions of dietary triacylglycerol microspheres on growth, survival, and settlement of mussel (*Mytilus* sp.) larvae. *Mar Biol* 144:693–703. doi: 10.1007/s00227-003-1234-1
- [44] Gagné R, Tremblay R, Pernet F, et al (2010) Lipid requirements of the scallop *Pecten maximus* (L.) during larval and post-larval development in relation to addition of *Rhodomonas salina* in diet. *Aquaculture* 309:212–221. doi: 10.1016/j.aquaculture.2010.09.040
- [45] Qiu J, Gosselin L, Qian P (1997) Effects of short-term variation in food availability on larval development in the barnacle *Balanus amphitrite amphitrite*. *Mar Ecol Prog Ser* 161:83–91. doi: 10.3354/meps161083
- [46] Thiagarajan V, Harder T, Qian P (2002) Relationship between cyprid energy reserves and metamorphosis in the barnacle *Balanus amphitrite* Darwin (Cirripedia; Thoracica). *J Exp Mar Bio Ecol* 280:79–93. doi: 10.1016/S0022-0981(02)00415-X
- [47] Pechenik J, Wendt D, Jarrett J (1998) Metamorphosis is not a new beginning larval experience influences juvenile performance. *Bioscience* 48:901–910. doi: 10.2307/1313294
- [48] Harder T, Thiagarajan V, Qian P (2001) Combined effect of cyprid age and lipid content on larval attachment and metamorphosis of *Balanus amphitrite* Darwin. *Biofouling* 17:257–262. doi: 10.1080/08927010109378486

- [49] Thiyagarajan V, Harder T, Qiu J, Qian P (2003) Energy content at metamorphosis and growth rate of the early juvenile barnacle *Balanus amphitrite*. Mar Biol 143:543–554. doi: 10.1007/s00227-003-1077-9
- [50] Pechenik J (2006) Larval experience and latent effects—metamorphosis is not a new beginning. Integr Comp Biol 46:323–333. doi: 10.1093/icb/icj028
- [51] Chiu J, Wang H, Thiyagarajan V, Qian P (2008) Differential timing of larval starvation effects on filtration rate and growth in juvenile *Crepidula onyx*. Mar Biol 154:91–98. doi: 10.1007/s00227-007-0902-y
- [52] Cullen JJ, Franks PJS, Karl DM, Longhurst A (2002) Physical influences on marine ecosystem dynamics. In: Robinson A.R., McCarthy J.J. and Rothchild B.J. (eds.), The Sea - Volume 12. pp. 297–336. Wiley, New York.
- [53] Eckman JE, Wener FE, Gross TF (1994) Modeling some effects of behavior on larval settlement in a turbulent boundary-layer. Deep Sea Res Part II Trop Stud Oceanogr 41 (1):185–208. doi: 10.1016/0967-0645(94)90067-1
- [54] Gaylord B, Hodin J, Ferner MC (2013) Turbulent shear spurs settlement in larval sea urchins. Proc Natl Acad Sci USA 110 (17): 6901–6906. doi: 10.1073/pnas.1220680110
- [55] Wheeler JD, Chan KYK, Anderson EJ, Mullineaux LS (2016) Ontogenetic changes in larval swimming and orientation of pre-competent sea urchin *Arbacia punctulata* in turbulence. J Exp Biol 219(9): 1303–1310. doi: 10.1242/jeb.129502
- [56] Queiroga H, Cruz T, dos Santos A, Dubert J, Gonzalez-Gordillo JL, Paula J, Peliz A, Santos AMP (2007) Oceanographic and behavioural processes affecting invertebrate larval dispersal and supply in the western Iberia upwelling ecosystem. Prog Oceanogr 74(2–3): 174–191.
- [57] Saunders MI, Metaxas A (2010). Physical forcing of distributions of bryozoan cyphonautes larvae in a coastal embayment. Mar Ecol Prog Ser 418:131–145. doi:10.3354/meps08842
- [58] Tapia FJ, DiBacco C, Jarrett J, Pineda J (2010) Vertical distribution of barnacle larvae at a fixed nearshore station in southern California: stage-specific and diel patterns. Estuar Coast Shelf Sci 86 (2): 265–270. doi: 10.1016/j.ecss.2009.11.003
- [59] Queiroga H, Blanton J (2005) Interactions between behavior and physical forcing in the control of horizontal transport of decapods crustacean larvae. Adv Mar Biol 47:107–214. doi: 10.1016/S0065-2881(04)47002-3
- [60] Graham WM, Field JG, Potts DC (2013) Persistent upwelling shadows and their influence on zooplankton distributions. Mar Biol 114 (4): 561–570. doi: 10.1007/BF00357253
- [61] Morgan SG, Fisher JL (2010) Larval behavior regulates nearshore retention and offshore migration in an upwelling shadow and along the open coast. Mar Ecol Prog Ser 404: 109–126. doi: 10.3354/meps08476

- [62] Farfan LM, D'Sa EJ, Liu KB, Rivera-Monroy VH (2014) Tropical cyclone impacts on coastal regions: the case of the Yucatan and the Baja California Peninsulas, Mexico. *Estuar Coast* 37(6):1388–1402. doi: 10.1007/s12237-014-9797-2
- [63] Gyory J, Pineda J (2011) High-frequency observations of early-stage larval abundance: do storms trigger synchronous larval release in *Semibalanus balanoides*? *Mar Biol* 158(7): 1581–1589. doi: 10.1007/s00227-011-1671-1
- [64] Sutherland JP (1990) Recruitment regulates demographic variation in a tropical intertidal barnacle. *Ecology* 71:955–972. doi: 10.2307/1937365
- [65] Thompson RC, Norton TA, Hawkins SJ (1998) The influence of epilithic microbial films on the settlement of *Semibalanus balanoides* cyprids – a comparison between laboratory and field experiments. *Hydrobiologia* 375: 203–216. doi: 10.1023/A:1017036301082
- [66] Yang J, Li X, Liang X, Bao W, Shen H, Li J (2014) Effects of natural biofilms on settlement of plantigrades of the mussel *Mytilus coruscus*. *Aquaculture* 424: 228–233. doi: 10.1016/j.aquaculture.2014.01.007
- [67] Pineda J, Starczak V, Stueckle T (2006) Timing of successful settlement: demonstration of a recruitment window in the barnacle *Semibalanus balanoides*. *Mar Ecol Prog Ser* 320:233–237. doi: 10.3354/meps320233
- [68] Vargas CA, Manríquez PH, Navarrete SA (2006) Feeding by larvae of intertidal invertebrates: assessing their position in pelagic food webs. *Ecology* 87(2):444–457. doi: 10.1890/05-0265
- [69] Krenz C, Menge BA, Freidenburg TL, Lubchenco J, Chan F, Foley MM, Nielsen KJ (2011) Ecological subsidies to rocky intertidal communities: linear or non-linear changes along a consistent geographic upwelling transition? *J Exp Mar Bio Ecol* 409 (1–2): 361–370. doi: 10.1016/j.jembe.2011.10.003
- [70] Pfaff MC, Branch GM, Wieters EA, Branch RA, Broitman BR (2011) Upwelling intensity and wave exposure determine recruitment of intertidal mussels and barnacles in the southern Benguela upwelling region. *Mar Ecol Prog Ser* 425:141–152. doi: 10.3354/meps09003
- [71] Reddin CJ, Docmac F, O'Connor NE, Bothwell JH, Harrod C (2015) Coastal upwelling drives intertidal assemblage structure and trophic ecology. *Plos One* 10(7): e0130789. doi: 10.1371/journal.pone.0130789
- [72] Menge BA, Blanchette C, Raimondi P, Freidenburg T, Gaines S, Lubchenco J, Lohse D, Hudson G, Foley M, Pamplin J (2004) Species interaction strength: testing model predictions along an upwelling gradient. *Ecol Monogr* 74(4): 663–684. doi: 10.1890/03-4060
- [73] Shanks AL, Mcculloch A, Miller J (2003) Topographically generated fronts, very near-shore oceanography and the distribution of larval invertebrates and holoplankters. *J Plank Res* 25(10):1251–1277. doi: 10.1093/plankt/fbg090
- [74] Menge BA, Gouhier TC, Freidenburg T, Lubchenco J (2011). Linking long-term, large-scale climatic and environmental variability to patterns of marine invertebrate recruitment: toward explaining “unexplained” variation. *J Exp Mar Biol Ecol*, 400(1–2): 236–249. doi: 10.1016/j.jembe.2011.02.003

- [75] Blanchette CA, Broitman BR, Gaines SD (2006). Intertidal community structure and oceanographic patterns around Santa Cruz Island, CA, USA. *Mar Biol*, 149: 689–701. doi: 10.1007/s00227-005-0239-3
- [76] Byrne M, Przeslawski R. (2013) Multistressor impacts of warming and acidification of the ocean on marine invertebrates' life histories. *Integr Comp Biol* 53(4): 582–596. doi: 10.1093/icb/ict049
- [77] Menge BA, Chan F, Nielsen KJ, Di Lorenzo E, Lubchenco J (2009). Climatic variation alters supply-side ecology: impact of climate patterns on phytoplankton and mussel recruitment. *Ecol Monogr*, 79(3): 379–395. doi: 10.1890/08-2086.1