We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Caffeine Dose-Response Relationship and Behavioral Screening in Zebrafish

Luana C. Santos, Julia Ruiz-Oliveira,
Priscila F. Silva and Ana C. Luchiari

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.68341>

Abstract

It has been centuries since humans consume coffee and get the benefits of this bean. Many researches worldwide continue to show healthful properties of coffee, while others suggest a number of side effects. In fact, anything consumed in excess may cause disturbance of the body functioning, whereas caffeine is a central nervous system stimulant that increases focus and improves performance, its high concentration can cause insomnia, dizziness, and vomiting. Thus, the question is: which coffee dose promotes benefits and prevents risks? To answer it, we used the zebrafish, a popular animal model that is at the vanguard of psychopharmacological research due to its unique combination of complexity and simplicity, translational relevance and applicability to high throughput behavioral drug screens. In the current study, we examine time-course and dose-dependent changes in zebrafish following exposure to caffeine. Our data show an inverted U-shaped path for the locomotor parameters and crescent path for the anxiety-like parameters. High doses are harmful to the individual, because the stimulating effect disappears and anxiogenic effects take place. We conclude that temporal analysis of zebrafish behavior is a sensitive method for the study of acute caffeine exposure-induced functional changes in the vertebrate brain.

Keywords: *Danio rerio*, anxiety-like, drug therapy, pharmacology, biphasic effect

1. Introduction

Caffeine is a psychostimulant substance worldwide used, which can potentially increase alertness and decrease fatigue and drowsiness [1–5]. Coffee, the beverage in which caffeine is most representative, is known to be rich in biologically active compounds that possess a

variety of therapeutic and functional effects. However, heavy coffee consumption may provoke systemic damages such as irregular heart rate, increased ventilation, anxiety, and due to its psychoactive properties, caffeine is likely to have addictive properties [6, 7]. Unlike other psychoactive drugs, caffeine consumption is legal and does not present any form of regulation. Moreover, caffeine consumption is not restricted to coffee and tea, but it has commonly been combined with other food products, such as chocolates, sodas, potato chips, and also bottled water. Even more risky, caffeine has been associated with other psychoactive drugs, as alcohol (i.e., energy drinks). While there is no specific recommendation on the amount of caffeine used or an indication of a critical value that may cause health problems, the U.S. Food and Drug Administration (FDA) has suggested 400 mg of caffeine/day for health adults [8]. However, there is not a clear picture of the overtime and dose-dependent effects of caffeine, demanding attention on behavioral pharmacological studies on this issue.

During the past 2 decades, numerous studies have approached the effects of psychoactive compounds used by humans in other mammals (rodents) [9–11]. Caffeine has gained attention because of its multiple targets in the brain. For instance, adenosine, ryanodine, and γ -aminobutyric acid receptors and cyclic nucleotide phosphodiesterase isoenzymes [12] seem to be related to stimulant effects of caffeine. However, as other drugs, caffeine empowers the central nervous system functioning when it reaches the therapeutic range; otherwise, it is too low concentration to cause an effect or too high concentration that causes intoxication. In this sense, several studies show contradictory behavioral effects following caffeine exposure: both increase in locomotor activity [13] and decrease in motor response [14, 15] were observed. The divergences on results may be related to caffeine dose and observation period in each study. Therefore, it is urgent to present effect of different doses and a short time-scale evaluation of caffeine induced changes in order to establish its therapeutic range, and then, how it can be properly used when in combination with other stimulant or depressant drugs.

Instead of using the most common animal model in pharmacological research, we propose the use of zebrafish to fill this gap in caffeine research. This small vertebrate is at the vanguard of neuroethological research and has been suggested for behavioral screening of drugs. The zebrafish has gained attention in behavioral brain research due to its ideal balance between the complexity of the physiological system and the simplicity of the biological model. It includes the fact that the zebrafish presents several molecular pathways, proteins, and protein products also found in mammals [16–22], besides the genome homology of about 70–80% [23]. Also, its brain structure [24, 25] and neurochemistry [26] offer translational relevance to humans [23] and allow exploring the model for a thorough understanding of the effects of substances used/abused by humans. In fact, various studies have shown that zebrafish respond similar to mammals when treated with many pharmacological compounds [27–29]. For example, benzodiazepine medication causes sedative effects in mouse [30, 31] and zebrafish [32]. For caffeine, it is not different: both rodents and zebrafish present anxiety-like behavior following caffeine exposure [33–36]. Furthermore, caffeine is water soluble and can be delivered to the fish via noninvasive method.

Zebrafish is not only an ideal model for behavioral screening, but also the majority of the genes identified in this species is conserved and has homologs in mammals [23, 37], which

allows for the examination of brain function and the development of brain diseases [27]. The zebrafish is an important model for research on psychoactive substances; in this sense, to know the effect of different doses of caffeine in their behavior is an important step for the development of methodologies to assess the effect of the substance in physiology and cognition. Our overtime dose-response analyses are one of the most detailed studies of caffeine in zebrafish and serves as a behavioral screen for future studies on the neural effects of caffeine or its effects when combined with other drugs.

2. Materials and methods

2.1. Animals and housing

Adult zebrafish (wild-type, both sexes) was obtained from a local fish farm (Natal-RN) and held in 50 L tanks forming a closed recirculating high-density system at the vivarium of the Fish Laboratory (Physiology Department—UFRN). The system maintained water quality by a multistage filtration, in which four filters processed the water in a flow of 3200 L/h, including a mechanical, a biological, activated carbon, and a UV light sterilizing filter. Water temperature was maintained at 28 C, pH in 7.1. Photoperiod was set at 12:12 light:dark cycle, with light intensity of 250 lx.

Fishes were fed two times a day with live brine shrimp and flaked food. Experimental procedures were revised and approved by the Ethical Committee for Animal Use of Federal University of Rio Grande do Norte (CEUA 045/2015).

2.2. Caffeine exposure

To determine overtime effect of caffeine doses in zebrafish, 144 animals (both sexes; 4.87 ± 1.35 g) were randomly assigned to different experimental groups that corresponded to each caffeine concentration ($n = 12$ for each group). This experimental design utilized 12 acute challenge doses: 0.0 (control), 0.5, 1.0, 5.0, 10.0, 15.0, 25.0, 50.0, 65.0, 75.0, 100.0, and 150.0 mg/L caffeine (Sigma Aldrich, 1,3,7-trimethylxanthine, Cat#C0750).

Fishes were initially held in groups of 12 in glass tanks (50 cm × 30 cm × 25 cm, width × depth × height; 37 L) for 7 days to acclimatize the fish to the test room. The bottom and back side of the holding tanks were covered with white paper to provide a uniform environment. During this period, water quality was kept the same as in the stock condition, with filtration and oxygen renovation given by a 140 Bio Wheel power filters. Food was offered twice daily.

For the behavioral assay, smaller tanks (40 cm × 20 cm × 25 cm, 15 L) were used. Caffeine was added directly to the testing water, to achieve each testing dose. Fishes were individually transferred to the testing tanks and behavior was recorded during 60 min using an HD camcorder (Sony Digital Video Camera Recorder; DCR-SX45) positioned 1 m away and in front of the tanks.

Fish behavior records were tracked using the Zebtrack software developed in MatLab [38]. The behavioral variables measured were average swimming speed, total distance travelled, duration of immobility (freezing), and time spent at the bottom of the tank (up to 5 cm from the bottom).

2.3. Statistical analysis

Data were first evaluated in search for outliers, homogeneity, normality, zero trouble, collinearity, and variables independency by inferential statistics [39]. After that, a Mixed Effects Model Analysis was applied considering the behavioral response as the response variable and the time (60 min records) and caffeine dose as the explanatory variables. The repeated measures characteristic of the data (over time data sampling) required longitudinal data analysis [39].

The exploratory analysis showed abnormal distribution and over dispersion of the residuals, and thus, a glmmPQL command (MASS package [40]) was used to develop the mixed model in the R program [41]. The mixed model showed random effect factors, which was the variation in behavior between groups, fixed effect factors, which was the caffeine doses used, and error.

The response variables were positive continuous quantitative data, not including zero ($Y > 0$); thus, a Goodness-of-fittest was performed to verify the best distribution function. If the response variable is continuous, then the normal and gamma are the best options [39]. Our response variable data best distribution function fitted gamma distribution (link function = inverse). The post-hoc comparisons between treatments, of each model, were made using the Tukey test in “lsmeans” package [42].

Average speed, distance traveled, freezing, and time at the bottom of the tank were also compared between the groups (caffeine doses) using One-Way ANOVA. For all comparison, the probability level considered for significance was $p < 0.05$.

3. Results

Figure 1 depicts zebrafish average swimming speed during acute caffeine exposure. Graphs referent to the very small doses (0.5 and 1.0 mg/L) are not presented to make it clear and due to its similarity to the doses below (0 mg/L) and above (5 mg/L). The mixed model comparison showed that average speed changed over time (GLMM, $\chi^2 = 7.33$, $df = 1$, $p < 0.006$; **Table 1**) and due to the caffeine dose used (GLMM, $\chi^2 = 651.49$, $df = 10$, $p < 0.001$; **Table 1**). The post-hoc comparison test (lsmeans) between caffeine doses is shown in **Table 1**: 10 and 25 mg/L of caffeine lead to an increase in average speed, while higher doses (50–150 mg/L) cause a decrease in swimming speed compared to low doses (0–5 mg/L).

Figure 2 shows zebrafish total distance traveled along the 60-min period of caffeine exposure. Graphs denoting the behavior for 0.5 and 1.0 mg/L are not shown. The mixed model

Figure 1. Time-course path of the average swimming speed during 60-min caffeine exposure in zebrafish. Mean $\pm$ SEM are shown for every 1-min intervals of the total 60-min recording. The caffeine doses (group designations) are shown above the graphs. Sample sizes (n) were 12 for each dose. Note the elevated activity in the group of fish exposed to 10 and 25 mg/L caffeine as compared to control. Also note the decreased activity in the fish that was exposed to doses of 50 mg/L caffeine and above it. For statistical analysis see Section 3 and **Table 1**.

comparison showed that total distance traveled changed over time (GLMM, $\chi^2 = 11.68$, $df = 1$, $p < 0.0006$; **Table 1**) and due to the caffeine dose used (GLMM, $\chi^2 = 271.49$, $df = 10$, $p < 0.001$; **Table 1**). The post-hoc comparison test (lsmeans) between caffeine doses is shown in **Table 1**. Caffeine doses higher than 50 mg/L decreased distance traveled compared to doses from 0 to 25 mg/L.

Figure 3 illustrates freezing behavior presented by zebrafish during caffeine challenge, a behavior related to fear and anxiety. Graphs referent to doses 0.5 and 1.0 mg/L are omitted. The mixed model comparison showed that freezing did not change over time (GLMM, $\chi^2 = 2.13$, $df = 1$, $p < 0.14$; **Table 1**) but changed according to the caffeine dose used (GLMM, $\chi^2 = 214.66$, $df = 10$, $p < 0.001$; **Table 1**). The post-hoc comparison test (lsmeans) between caffeine doses is shown in **Table 1**. Caffeine doses higher than 50 mg/L increased freezing behavior in zebrafish compared to doses from 0 to 25 mg/L.

Figure 4 presents zebrafish time spent at the bottom of the testing tank, another behavior associated to fear and anxiety response. Again, graphs representing doses 0.1 and 1.0 mg/L were not presented. The mixed model comparison showed that distance from the bottom did not change over time (GLMM, $\chi^2 = 0.18$, $df = 1$, $p < 0.66$; **Table 1**) but changed due to the caffeine exposure (GLMM, $\chi^2 = 170.91$, $df = 10$, $p < 0.001$; **Table 1**). The post-hoc comparison test (lsmeans) between caffeine doses is shown in **Table 1**. Caffeine doses of 50 mg/L and higher increased the time fish spent at the bottom of the tank.

Explanatory variable	Behavioral parameters											
	Average speed			Total distance travelled			Freezing			Distance from bottom		
	Chi-squared	<i>p</i> -Value		Chi-squared	<i>p</i> -Value		Chi-squared	<i>p</i> -Value		Chi-squared	<i>p</i> -Value	
60 min time course	7.33	0.006		11.68	0.0006		2.12	0.14		0.18	0.66	
Doses	651.49	>0.001		271.49	<0.001		214.66	<0.001		170.91	<0.001	
Pairwise comparison	lsmeans ± SEM	<i>t</i> -Value	<i>p</i> -Value	lsmeans ± SEM	<i>t</i> -Value	<i>p</i> -Value	lsmeans ± SEM	<i>t</i> -value	<i>p</i> -value	lsmeans ± SEM	<i>t</i> -value	<i>p</i> -value
Dose 0.0 vs. dose 0.5	-0.00 ± 0.01	-0.22	1.0	-4.17 ± 0.00	-0.83	0.99	0.01 ± 0.02	0.73	0.99	-0.01 ± 0.01	-0.84	0.99
Dose 0.0 vs. dose 1.0	-0.00 ± 0.01	-0.28	1.0	-3.11 ± 0.00	-0.62	0.99	0.01 ± 0.02	0.42	1.00	-0.01 ± 0.01	-0.80	0.99
Dose 0.0 vs. dose 5.0	-0.00 ± 0.01	-0.40	1.0	-3.45 ± 0.00	-0.68	0.99	0.03 ± 0.02	1.34	0.95	-0.00 ± 0.01	-0.33	1.00
Dose 0.0 vs. dose 10	0.06 ± 0.01	4.52	0.00	7.32 ± 0.00	1.46	0.92	-0.06 ± 0.02	-2.45	0.34	-0.00 ± 0.01	-0.64	0.99
Dose 0.0 vs. dose 25	0.06 ± 0.01	4.29	0.00	4.17 ± 0.00	0.83	0.99	-0.06 ± 0.02	-2.43	0.45	-0.00 ± 0.01	-0.07	1.00
Dose 0.0 vs. dose 50	-0.09 ± 0.01	-0.55	<0.00	-3.08 ± 0.00	-6.03	<0.00	0.14 ± 0.02	5.34	<0.00	0.04 ± 0.01	3.38	0.03
Dose 0.0 vs. dose 65	-0.13 ± 0.02	-8.07	<0.00	-3.74 ± 0.00	-6.69	<0.00	0.14 ± 0.02	5.25	<0.00	0.06 ± 0.01	4.50	0.00
Dose 0.0 vs. dose 75	-0.18 ± 0.01	-12.26	<0.00	-4.78 ± 0.00	-9.19	<0.00	0.15 ± 0.02	5.87	<0.00	0.08 ± 0.01	6.34	<0.00
Dose 0.0 vs. dose 100	-0.13 ± 0.01	-9.20	<0.00	-3.29 ± 0.00	-6.42	<0.00	0.07 ± 0.02	5.80	0.01	0.06 ± 0.01	5.08	0.00
Dose 0.0 vs. dose 150	-0.12 ± 0.01	-8.18	<0.00	-2.64 ± 0.00	-5.18	<0.00	0.17 ± 0.02	6.72	<0.00	0.07 ± 0.01	5.32	<0.00
Dose 0.5 vs. dose 1.0	-0.00 ± 0.01	-0.06	1.0	1.05 ± 0.00	0.20	1.00	-0.00 ± 0.02	0.30	1.00	0.00 ± 0.01	0.03	1.00

	Behavioral parameters											
	Average speed			Total distance travelled			Freezing			Distance from bottom		
Dose 0.5 vs. dose 5.0	-0.00 ± 0.01	-0.18	1.0	7.18 ± 0.00	0.14	1.00	0.01 ± 0.02	0.61	0.99	0.00 ± 0.01	0.50	1.00
Dose 0.5 vs. dose 10	0.07 ± 0.01	4.74	0.00	1.14 ± 0.00	2.29	0.44	-0.08 ± 0.02	-3.18	0.06	0.00 ± 0.01	0.19	1.00
Dose 0.5 vs. dose 25	0.06 ± 0.01	4.51	0.00	8.35 ± 0.00	1.66	0.85	-0.08 ± 0.02	-3.17	0.06	0.01 ± 0.00	0.76	0.99
Dose 0.5 vs. dose 50	-0.09 ± 0.01	-6.33	<0.00	-2.67 ± 0.00	-5.21	<0.00	0.12 ± 0.02	4.61	0.00	0.05 ± 0.01	4.23	0.00
Dose 0.5 vs. dose 65	-0.12 ± 0.02	-7.88	<0.00	-4.36 ± 0.01	-8.38	<0.00	0.12 ± 0.02	4.57	0.00	0.07 ± 0.01	5.29	<0.00
Dose 0.5 vs. dose 75	-0.18 ± 0.01	-12.04	<0.00	-2.87 ± 0.01	-5.60	<0.00	0.13 ± 0.02	5.13	0.00	0.09 ± 0.01	7.19	<0.00
Dose 0.5 vs. dose 100	-0.13 ± 0.01	-8.98	<0.00	-2.22 ± 0.01	-4.35	0.00	0.05 ± 0.02	2.07	0.60	0.08 ± 0.01	5.93	<0.00
Dose 0.5 vs. dose 150	-0.12 ± 0.01	-7.96	<0.00	-3.35 ± 0.01	-0.06	0.00	0.15 ± 0.02	5.99	<0.00	0.08 ± 0.01	6.17	<0.00
Dose 1.0 vs. dose 5.0	-0.00 ± 0.01	-0.12	1.0	1.04 ± 0.01	2.08	0.59	0.02 ± 0.01	0.92	0.99	0.00 ± 0.01	0.46	1.00
Dose 1.0 vs. dose 10	0.07 ± 0.01	4.80	0.00	7.29 ± 0.01	1.45	0.93	-0.07 ± 0.01	-2.87	0.14	0.00 ± 0.01	0.16	1.00
Dose 1.0 vs. dose 25	0.06 ± 0.01	4.57	0.00	-2.77 ± 0.01	-5.42	<0.00	-0.07 ± 0.01	-2.06	0.14	0.01 ± 0.01	0.72	0.99
Dose 1.0 vs. dose 50	-0.09 ± 0.01	-6.27	<0.00	-3.43 ± 0.01	-6.12	<0.00	0.12 ± 0.02	4.92	0.00	0.05 ± 0.01	4.19	0.00
Dose 1.0 vs. dose 65	-0.12 ± 0.02	-7.82	<0.00	-4.47 ± 0.01	-8.58	<0.00	0.13 ± 0.02	4.85	0.00	0.07 ± 0.01	5.26	<0.00
Dose 1.0 vs. dose 75	-0.18 ± 0.01	-11.98	<0.00	-2.98 ± 0.00	-5.80	<0.00	0.14 ± 0.01	5.44	<0.00	0.09 ± 0.01	7.16	<0.00
Dose 1.0 vs. dose 100	-0.13 ± 0.01	-8.92	<0.00	-2.33 ± 0.00	-4.56	0.00	0.06 ± 0.02	2.38	0.38	0.08 ± 0.01	5.90	<0.00

	Behavioral parameters											
	Average speed			Total distance travelled			Freezing			Distance from bottom		
Dose 1.0 vs. dose 150	-0.11 ± 0.01	-7.90	<0.00	1.07 ± 0.00	2.15	0.04	0.16 ± 0.01	6.30	<0.00	0.08 ± 0.01	6.14	<0.00
Dose 5.0 vs. dose 10	0.07 ± 0.01	4.93	0.00	1.07 ± 0.00	2.15	0.54	-0.10 ± 0.00	-3.80	0.01	-0.00 ± 0.01	-0.30	1.00
Dose 5.0 vs. dose 25	0.06 ± 0.01	4.69	0.00	7.63 ± 0.00	1.52	0.90	-0.09 ± 0.01	-3.78	0.01	0.00 ± 0.01	0.25	1.00
Dose 5.0 vs. dose 50	-0.09 ± 0.01	-6.14	<0.00	-2.74 ± 0.00	-5.35	<0.00	0.10 ± 0.01	4.00	0.00	0.05 ± 0.01	3.73	0.01
Dose 5.0 vs. dose 65	-0.12 ± 0.02	-7.71	<0.00	-3.39 ± 0.00	-6.06	<0.00	0.11 ± 0.02	4.00	0.00	0.07 ± 0.01	4.83	0.00
Dose 5.0 vs. dose 75	-0.18 ± 0.01	-11.86	<0.00	-4.43 ± 0.00	-8.52	<0.00	0.11 ± 0.02	4.52	0.00	0.09 ± 0.01	6.70	<0.00
Dose 5.0 vs. dose 100	-0.13 ± 0.01	-8.80	<0.00	-2.94 ± 0.00	-5.74	<0.00	0.03 ± 0.00	1.45	0.93	0.07 ± 0.01	5.43	<0.00
Dose 5.0 vs. dose 150	-0.11 ± 0.01	-7.78	<0.00	-2.29 ± 0.01	-4.49	0.00	0.14 ± 0.02	5.38	<0.00	0.07 ± 0.01	5.68	<0.00
Dose 10 vs. dose 25	0.00 ± 0.01	-0.23	1.00	-3.14 ± 0.00	-0.63	0.99	0.00 ± 0.00	0.01	1.00	0.00 ± 0.01	0.56	1.00
Dose 10 vs. dose 50	-0.16 ± 0.01	-11.04	<0.00	-3.82 ± 0.00	-7.48	<0.00	0.20 ± 0.02	7.79	<0.00	0.05 ± 0.01	4.03	0.00
Dose 10 vs. dose 65	-0.19 ± 0.02	-12.19	<0.00	-4.47 ± 0.00	-8.01	<0.00	0.21 ± 0.02	7.52	<0.00	0.07 ± 0.01	5.11	0.00
Dose 10 vs. dose 75	-0.25 ± 0.01	-16.68	<0.00	-5.51 ± 0.01	-10.62	<0.00	0.21 ± 0.02	8.32	<0.00	0.09 ± 0.01	7.00	<0.00
Dose 10 vs. dose 100	-0.20 ± 0.01	-13.66	<0.00	-4.02 ± 0.00	-7.87	<0.00	0.13 ± 0.00	5.25	<0.00	0.07 ± 0.01	5.74	<0.00
Dose 10 vs. dose 150	-0.19 ± 0.01	-12.66	<0.00	-3.37 ± 0.00	-6.63	<0.00	0.24 ± 0.02	9.17	<0.00	0.08 ± 0.01	5.98	<0.00
Dose 25 vs. dose 50	-0.16 ± 0.01	-10.81	<0.00	-3.50 ± 0.01	-6.86	<0.00	0.20 ± 0.02	7.78	<0.00	0.04 ± 0.01	3.47	0.02

	Behavioral parameters											
	Average speed			Total distance travelled			Freezing			Distance from bottom		
Dose 25 vs. dose 65	-0.19 ± 0.02	-11.98	<0.00	-4.16 ± 0.00	-7.44	<0.00	0.21 ± 0.02	7.50	<0.00	0.06 ± 0.01	4.59	0.00
Dose 25 vs. dose 75	-0.25 ± 0.01	-16.45	<0.00	-5.20 ± 0.01	-10.00	<0.00	0.21 ± 0.00	8.30	<0.00	0.08 ± 0.01	6.44	<0.00
Dose 25 vs. dose 100	-0.20 ± 0.01	-13.43	<0.00	-3.71 ± 0.00	-7.24	<0.00	0.13 ± 0.01	5.24	<0.00	0.07 ± 0.01	5.18	<0.00
Dose 25 vs. dose 150	-0.18 ± 0.01	-12.42	<0.00	-3.05 ± 0.00	-6.00	<0.00	0.24 ± 0.01	9.16	<0.00	0.17 ± 0.01	5.42	<0.00
Dose 50 vs. dose 65	-0.03 ± 0.02	-2.05	0.60	-6.55 ± 0.00	-1.15	0.98	0.00 ± 0.02	0.29	1.00	0.01 ± 0.01	1.35	0.95
Dose 50 vs. dose 75	-0.08 ± 0.01	-5.76	0.08	-1.69 ± 0.00	-3.19	0.06	0.01 ± 0.00	0.52	1.00	0.03 ± 0.01	2.98	0.11
Dose 50 vs. dose 100	-0.04 ± 0.01	-2.67	0.22	-2.05 ± 0.00	-0.39	1.00	-0.06 ± 0.02	-2.54	0.29	0.02 ± 0.01	1.70	0.82
Dose 50 vs. dose 150	-0.02 ± 0.01	-1.64	0.85	4.47 ± 0.01	0.86	0.99	0.03 ± 0.02	1.38	0.95	0.02 ± 0.01	1.95	0.67
Dose 65 vs. dose 75	-0.05 ± 0.02	-3.28	0.05	-1.03 ± 0.01	-1.80	0.77	0.00 ± 0.02	0.18	1.00	0.02 ± 0.01	1.40	0.94
Dose 65 vs. dose 100	-0.00 ± 0.02	-0.41	1.00	4.50 ± 0.00	0.79	0.99	-0.07 ± 0.02	-2.65	0.23	0.00 ± 0.01	0.22	1.00
Dose 65 vs. dose 150	0.00 ± 0.02	0.53	1.00	1.10 ± 0.01	1.94	0.68	0.02 ± 0.02	0.98	0.99	0.00 ± 0.01	0.45	1.00
Dose 75 vs. dose 100	0.04 ± 0.01	3.09	0.08	1.48 ± 0.00	2.80	0.16	-0.08 ± 0.01	-3.06	0.09	-0.01 ± 0.01	-1.28	0.97
Dose 75 vs. dose 150	0.06 ± 0.01	4.12	0.07	2.14 ± 0.00	4.05	0.07	0.02 ± 0.02	0.85	0.99	-0.01 ± 0.01	-1.02	0.99
Dose 100 vs. dose 150	0.01 ± 0.01	1.03	0.99	6.52 ± 0.00	1.25	0.97	0.10 ± 0.02	3.92	0.00	0.00 ± 0.01	0.25	1.00

Dose values are expressed in mg/L.

Table 1. Estimates of mixed effect model for the behavioral parameters measured during caffeine exposure.

Figure 2. Time-course path of the total distance traveled during 60-min caffeine exposure in zebrafish. Mean $\pm$ SEM are shown for every 1-min intervals of the total 60 min recording. The caffeine doses (group designations) are shown above the graphs. Sample sizes (n) were 12 for each dose. Note the elevated activity in the group of fish exposed to 10 and 25 mg/L caffeine as compared to control. Also note the decreased activity in the fish that was exposed to doses of 50 mg/L caffeine and above it. For statistical analysis see Section 3 and **Table 1**.

Figure 3. Time-course path of the freezing behavior during 60-min caffeine exposure in zebrafish. Mean $\pm$ SEM are shown for every 1-min intervals of the total 60 min recording. The caffeine doses (group designations) are shown above the graphs. Sample sizes (n) were 12 for each dose. Note the decreased freezing in the group of fish exposed to 5, 10 and 25 mg/L caffeine as compared to control. On the contrary, note the increased freezing in the fish that was exposed to doses above 50 mg/L caffeine. For statistical analysis, see Section 3 and **Table 1**.

Figure 4. Time-course path of the time spent at the bottom of the tank (up to 5 cm from the bottom) during 60-min caffeine exposure in zebrafish. Mean $\pm$ SEM are shown for every 1-min intervals of the total 60 min recording. The caffeine doses (group designations) are shown above the graphs. Sample sizes (n) were 12 for each dose. Note the decreased time at the bottom in the group of fish exposed to 5, 10, and 25 mg/L caffeine as compared to control. Also note the increased time at the bottom in the fish exposed to doses above 50 mg/L caffeine. For statistical analysis, see Section 3 and **Table 1**.

Figure 5 displays the median and range of variation of the 60-min caffeine exposure in zebrafish, both for the parameters related to locomotion (**Figure 5a** and **b**) and the parameters related to anxiety-like behavior (**Figure 5c** and **d**). One-Way ANOVA between groups (caffeine doses) showed that average speed did not differ from the control condition when fish is exposed to doses up to 5 mg/L caffeine, but it is increased with doses of 10 and 25 mg/L and decreased with doses above 50 mg/L ($F = 1087.97$, $df = 10$, $p < 0.001$, **Figure 5a**) indicating an inverted U shape. The same patterns were observed for total distance travelled, in which the lower doses (0.5–5 mg/L) did not differ from the control, doses of 10 and 25 mg/L increased distance traveled and doses above 50 mg/L decreased distance traveled (One-Way ANOVA, $F = 374.82$, $df = 10$, $p < 0.001$, **Figure 5b**). For the freezing behavior, One-Way ANOVA showed a slight decrease with increasing doses, with the lowest values of freezing registered at 10 and 25 mg/L, and a sharp increase with doses above 50 mg/L ($F = 462.15$, $df = 10$, $P < 0.001$, **Figure 5c**), suggesting the anxiogenic effect of high caffeine doses. The time fish spent at the bottom of the tank was reduced by doses from 0.5 to 25 mg/L and highly increased by doses above 50 mg/L (One-Way ANOVA, $F = 427.27$, $df = 10$, $p < 0.001$). **Figure 5d** depicts the comparison between caffeine doses and the tendency line indicating an increasing in time at the bottom concomitant to the increase in caffeine dose.

Figure 5. Box plot shows median and interquartile range of locomotor behavior: (a) average swimming speed, (b) total distance traveled, (c) freezing, and (d) time spent at the bottom of the tank, whiskers represent the range. Tendency line indicates the inverted U shape observed for increasing doses of caffeine on locomotor behavioral parameters (swimming speed and distance traveled), and the ascendant pattern observed for increase doses of caffeine on anxiety-like behavioral parameters (freezing and time at the bottom of the tank). For statistical analysis, see Section 3.

4. Discussion

This study characterizes variations in the locomotor pattern and anxiety-like behavior derived from acute exposure to caffeine in zebrafish. We evaluated a wide spectrum of caffeine doses, from 0.5 to 150 mg/L, every 1-min interval along 60-min period of the drug exposure. This detailed analysis of different doses overtime allowed to the observation that caffeine has a biphasic effect, stimulating locomotion and decreasing anxiety at low levels, and diminishing activity and increasing anxiety-like behavior at doses above 50 mg/L.

While caffeine is accepted to act as a stimulant on the central nervous system [43], it is worth noting that in fact this drug exerts distinct responses depending on the amount used. Initially, caffeine has little or no impact on behavior, followed by the most evident effect caused by an intermediate dose, and then a remarkable suppression at the level of behavioral activity, as observed in **Figure 5**. High caffeine doses (50 and 100 mg/L) were previously shown to depress locomotor activity in zebrafish [15, 44], but our study is the first to present a time-course analyses of the effects of several doses of caffeine in adult zebrafish. The same dose-dependent effects observed herein in rodents, suggesting the models similarity in terms of the mechanism by which caffeine produces behavioral effects [45].

Adenosine receptor blockade seems to be the prevalent action of caffeine in the brain. The increase on locomotor activity after caffeine exposure derives from the blockade of A_{2A} adenosine receptors, preventing the inhibitory action of adenosine on the nervous system [46]. The antagonist role of caffeine usually stimulates the central nervous system and also activates dopaminergic transmission [47–49], which is consistent with the drug reinforcing properties. Another event that occurs at the time of caffeine ingestion and which causes increased

locomotor activity is the inhibition of phosphodiesterase (an enzyme that hydrolyses cAMP), which promotes the release of calcium from intracellular reserves and interferes with the sensitivity of GABA receptors [50].

Caffeine is a substance widely used by the society [1] and, if used in moderation (up to 200 mg/day/person on average), it may lead to several benefits, such as improved performance on tasks that require attention and focus [51, 52]. For instance, it was observed that zebrafish improves object discrimination when treated with moderate doses of caffeine [53]. The most notable effects of low-to-moderate doses of caffeine include increased alertness, energy, and ability to concentrate [54]. On the other hand, high and abusive use of caffeine may inhibit these effects [55]. At high concentrations, caffeine is suggested to increase glucose utilization in the CNS, what also seems to be related to its stimulatory effects [56]. The elevated sugar level (main CSN substrate) together with the blockage of adenosine inhibitory effects, in turn, is responsible to the caffeine side-effects on the motor system and sleep-wake cycle, two functions highly susceptible to caffeine.

Moreover, higher caffeine doses induce negative effects such as increased sympathetic response (tachycardia, higher ventilation), restlessness, insomnia, and anxiety [57]. The high acute dose of caffeine exacerbates anxiety-like behavior, reduces locomotion, and in many cases, causes behavior similar to seizure [58], very high doses of caffeine may also cause intoxication and death of the individuals [54, 59]. The same pattern of effect was observed in zebrafish larvae under the action of high doses of caffeine [60]. It is known that adenosine regulates the activity of several neurons, such as glutamatergic; thus, if the effect of adenosine is blocked, glutamatergic transmission increases and may turn to an extremely high excitatory response [12, 56].

Finally, our results reinforce the zebrafish as a valuable model organism for throughput screening of behavioral-related drugs. While caffeine is a legal and widely consumed substance, the amount of caffeine ingested should be taken into consideration since negative effects are observed after high doses consumption. We found that moderate caffeine intake ameliorates performance, but a robust anxiety-related response occurs following exposure to high doses of caffeine. Taken together, these results confirm zebrafish as an accurate, reliable, and efficient model for basic translational research of psychoactive drugs on physiology and behavior.

Acknowledgements

The authors are thankful to J.J. Oliveira for technical support and F. Freire for statistical assistance.

Author details

Luana C. Santos, Julia Ruiz-Oliveira, Priscila F. Silva and Ana C. Luchiari*

*Address all correspondence to: analuchiari@yahoo.com.br

Department of Physiology, Bioscience Center, Federal University of Rio Grande do Norte, Natal, Rio Grande do Norte, Brazil

References

- [1] Fredholm BB, Bättig K, Holmén J, Nehlig A, Zvartau EE. Actions of caffeine in the brain with special reference to factors that contribute to its widespread use. *Journal of Pharmacology and Experimental Therapeutics*. 1999;**51**(1):83–133. DOI: 0031-6997/99/5101-0083\$03.00/0
- [2] Frary CD, Johnson RK, Wang MQ. Food sources and intakes of caffeine in the diets of persons in the United States. *Journal of the American Dietetic Association*. 2005;**105**(1):110–113. DOI: 10.1016/j.jada.2004.10.027
- [3] Mitchell DC, Knight CA, Hockenberry J, Teplansky R, Hartman TJ. Beverage caffeine intakes in the U.S. *Food and Chemical Toxicology*. 2014;**63**:136–142. DOI: 10.1016/j.fct.2013.10.042
- [4] Heckman MA, Weil J, Mejia EG. Caffeine (1,3,7-trimethylxanthine) in foods: A comprehensive review on consumption, functionality, safety, and regulatory matters. *Journal of Food Science*. 2010;**75**(3):77–87. DOI: 10.1111/j.1750-3841.2010.01561.x
- [5] Beydoun MA, Gamaldo AA, Beydoun HA, Tanaka T, Tucker KL, Talegawkar SA, Ferrucci L, Zonderman AB. Caffeine and alcohol intakes and overall nutrient adequacy are associated with longitudinal cognitive performance among U.S. adults. *The Journal of Nutrition*. 2014;**144**(6):890–901. DOI: 10.3945/jn.113.189027
- [6] Budygin EA, Weiner JL. Exploring the neurochemical basis of alcohol addiction-related behaviors: Translational research. *Translational Biomedicine*. 2015;**6**(Suppl Spec). <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4710378/pdf/nihms723979.pdf>
- [7] Ferré S. Mechanisms of the psychostimulant effects of caffeine: Implications for substance use disorders. *Psychopharmacology*. 2016;**233**(10):1963–1979. DOI: 10.1007/s00213-016-4212-2
- [8] U.S. Food and Drug Administration. Citizen Petition for New Labeling Requirement for Caffeine in Food [Internet]. 2016. Available from: <http://www.fda.gov/ohrms/dockets/dockets/06p0039/06p-0039-cp00001-01-vol1.pdf> [Accessed: 05-01-2017]
- [9] Molloy JW, Calcagno CJ, Williams CD, Jones FJ, Torres DM, Harrison SA. Association of coffee and caffeine consumption with fatty liver disease, nonalcoholic steatohepatitis, and degree of hepatic fibrosis. *Hepatology*. 2012;**55**(2):429–446. DOI: 10.1002/hep.24731
- [10] Sepkowitz KA. Energy drinks and caffeine-related adverse effects energy drinks and caffeine-related adverse effects. *JAMA*. 2013;**309**(3):243–244. DOI: 10.1001/jama.2012.173526
- [11] Denoeud F, Carretero-Paulet L, Dereeper A, Droc G, Guyot R, Pietrella M, Zheng C, Alberti A, Anthony F, Aprea G, Aury JM, Bento P, Bernard M, Bocs S, Campa C, Cenci A, Combes MC, Crouzillat D, Da Silva C, Daddiego L, De Bellis F, Dussert S, Garsmeur O, Gayraud T, Guignon V, Jahn K, Jamilloux V, Joët T, Labadie K, Lan T, Leclercq J, Lepelley M, Leroy T, Li LT, Librado P, Lopez L, Muñoz A, Noel B, Pallavicini A, Perrotta G, Poncet V, Pot D,

- Priyono, Rigoreau M, Rouard M, Rozas J, Tranchant-Dubreuil C, VanBuren R, Zhang Q, Andrade AC, Argout X, Bertrand B, de Kochko A, Graziosi G, Henry RJ, Jayarama, Ming R, Nagai C, Rounsley S, Sankoff D, Giuliano G, Albert VA, Wincker P, Lashermes P. The coffee genome provides insight into the convergent evolution of caffeine biosynthesis. *Science*. 2014;**345**(6201):1181–1184. DOI: 10.1126/science.1255274
- [12] Fisone G, Borgkvist A, Usiello A. Caffeine as a psychomotor stimulant: Mechanism of action. *Cellular and Molecular Life Sciences*. 2004;**61**(7–8):857–872. DOI: 10.1007/s00018-003-3269-3
- [13] Maximino C, da Silva AW, Gouveia AJ, Herculano AM. Pharmacological analysis of zebrafish (*Danio rerio*) scototaxis. *Progress in Neuro-Psychopharmacology & Biological Psychiatry*. 2011;**35**(2):624–631. DOI: 10.1016/j.pnpbp.2011.01.006
- [14] Pechlivanova D, Tchekalarova J, Nikolov R, Yakimova K. Dose-dependent effects of caffeine on behavior and thermoregulation in a chronic unpredictable stress model of depression in rats. *Behavioural Brain Research*. 2010;**209**(2):205–211. DOI: 10.1016/j.bbr.2010.01.037
- [15] Ladu F, Mwaffo V, Li J, Macrì S, Porfiri M. Acute caffeine administration affects zebrafish response to a robotic stimulus. *Behavioural Brain Research*. 2015;**289**:48–54. DOI: 10.1016/j.bbr.2015.04.020
- [16] Faraco JH, Appelbaum L, Marin W, Gaus SE, Mourrain P, Mignot E. Regulation of hypocretin (orexin) expression in embryonic zebrafish. *Journal of Biological Chemistry*. 2006;**281**(40):29753–29761. DOI: 10.1074/jbc.M605811200
- [17] Holzschuh J, Ryu S, Aberger F, Driever W. Dopamine transporter expression distinguishes dopaminergic neurons from other catecholaminergic neurons in the developing zebrafish embryo. *Mechanisms of Development*. 2001;**101**(1–2):237–243. DOI: 10.1016/S0925-4773(01)00287-8
- [18] Kaslin J, Nystedt JM, Ostergård M, Peitsaro N, Panula P. *Journal of Neuroscience*. 2004;**24**(11):2678–2689. DOI: 10.1523/JNEUROSCI.4908-03.2004
- [19] Kaslin J, Panula P. Comparative Anatomy of the Histaminergic and Other Aminergic Systems in zebrafish (*Danio rerio*). *Journal of Comparative Neurology*. 2001;**440**(4):342–377. DOI: 10.1002/cne.1390
- [20] McLean DL, Fetcho JR. Ontogeny and innervation patterns of dopaminergic, noradrenergic, and serotonergic neurons in larval zebrafish. *Journal of Comparative Neurology*. 2004;**480**(1):38–56. DOI: 10.1002/cne.20280
- [21] Mueller T, Vernier P, Wullmann MF. The adult central nervous cholinergic system of a neurogenetic model animal, the zebrafish *Danio rerio*. *Brain Research*. 2004;**1011**(2):156–169.
- [22] Prober DA, Rihel J, Onah AA, Sung RJ, Schier AF. Hypocretin/orexin overexpression induces an insomnia-like phenotype in Zebrafish. *Journal of Neuroscience*. 2006;**26**(51):13400–13410. DOI: 10.1523/JNEUROSCI.4332-06.2006

- [23] Crollius HR, Weissenbach J. Fish genomics and biology. *Genome Research*. 2005;**15**(12):1675–1682. DOI: <http://www.genome.org/cgi/doi/10.1101/gr.3735805>
- [24] Klee EW, Schneider H, Clark KJ, Cousin MA, Ebbert JO, Hooten WM, Karpyak VM, Warner DO, Ekker SC. Zebrafish: A model for the study of addiction genetics. *Human Genetics*. 2012;**131**(6):977–1008. DOI: 10.1007/s00439-011-1128-0
- [25] Kolb B, Whishaw IQ. Brain plasticity and behavior. *Annual Review of Psychology*. 1998;**49**(1):43–64. DOI: 10.1146/annurev.psych.49.1.43
- [26] Gerlai R, Chatterjee D, Pereira T, Sawashima T, Krishnannair R. Acute and chronic alcohol dose: Population differences in behavior and neurochemistry of zebrafish. *Genes, Brain, and Behavior*. 2009;**8**(6):586–599. DOI: 10.1111/j.1601-183X.2009.00488.x
- [27] Kalueff AV, Stewart AM, Gerlai R. Zebrafish as an emerging model for studying complex brain disorders. *Trends in Pharmacological Sciences*. 2014;**35**(2):63–75. DOI: 10.1016/j.tips.2013.12.002
- [28] Abreu MS, Giacomini AC, Gusso D, Rosa JG, Koakoski G, Kalichak F, Idalêncio R, Oliveira TA, Barcellos HH, Bonan CD, Barcellos LJ. Acute exposure to waterborne psychoactive drugs attract zebrafish. *Comparative Biochemistry and Physiology. Toxicology & Pharmacology*. 2016;**179**:37–43. DOI: 10.1016/j.cbpc.2015.08.009
- [29] Howarth DL, Lindtner C, Vacaru AM, Sachidanandam R, Tsedensodnom O, Vasilkova T, Buettner C, Sadler KC. Activating transcription factor 6 is necessary and sufficient for alcoholic fatty liver disease in zebrafish. *PLoS Genetics*. 2014;**10**(5):e1004335. DOI: 10.1371/journal.pgen.1004335
- [30] Nishino T, Takeuchi T, Takechi K, Kamei C. Evaluation of anxiolytic-like effects of some short-acting benzodiazepine hypnotics in mice. *Journal of Pharmacological Sciences*. 2008;**107**(3):349–354. DOI: 10.1254/jphs.08107FP
- [31] Figueredo YN, Rodríguez EO, Reyes YV, Domínguez CC, Parra AL, Sánchez JR, Hernández RD, Verdecia MP, Pardo AGL. Characterization of the anxiolytic and sedative profile of JM-20: A novel benzodiazepine-dihydropyridine hybrid molecule. *Neurological Research*. 2013;**35**(8):804–812. DOI: 10.1179/1743132813Y.0000000216
- [32] Giacomini AC, Abreu MS, Giacomini LV., Siebel AM, Zimerman FF, Rambo CL, Mocelin R, Bonan CD, Piato AL, Barcellos LJ. Fluoxetine and diazepam acutely modulate stress induced-behavior. *Behavioural Brain Research*. 2016;**296**:301–310. DOI: 10.1016/j.bbr.2015.09.027
- [33] Egan RJ, Bergner CL, Hart PC, Cachat JM, Canavello PR, Elegante MF, Elkhayat SI, Bartels BK, Tien AK, Tien DH, Mohnot S, Beeson E, Glasgow E, Amri H, Zukowska Z, Kalueff AV. Understanding behavioral and physiological phenotypes of stress and anxiety in zebrafish. *Behavioural Brain Research*. 2009;**205**(1):38–44. DOI: 10.1016/j.bbr.2009.06.022G

- [34] Maximino C, Silva AWB, Araujo J, Lima MG, Miranda V, Puty B, Benzecry R, Picanço-Diniz DLW, Gouveia A Jr, Oliverira KRM, Herculano. Fingerprinting of psychoactive drugs in zebrafish anxiety-like behaviors. PLoS One. 2014;**9**(7):e103943. DOI: 10.1371/journal.pone.0103943
- [35] Gupta P, Khobragade SB, Shingatgeri VM, Rajaram SM. Assessment of locomotion behavior in adult Zebrafish after acute exposure to different pharmacological reference compounds. Drug Development and Therapeutics. 2014;**5**(2):127–33. DOI: 10.4103/2394-2002.139626
- [36] Richendrfer H, Pelkowski SD, Colwill RM, Creton R. On the edge: Pharmacological evidence for anxiety-related behavior in zebrafish larvae. Behavioural Brain Research. 2012;**228**(1):99–106. DOI: 10.1016/j.bbr.2011.11.041
- [37] Cerdà J, Conrad M, Markl J, Brand M, Herrmann H. Zebrafish vimentin: Molecular characterization, assembly properties and developmental expression. European Journal of Cell Biology. 1998;**77**(3):175-187. DOI: 10.1016/S0171-9335(98)80105-2
- [38] Pinheiro-da-Silva J, Silva PF, Nogueira MB, Luchiari AC. Sleep deprivation effects on object discrimination task in zebrafish (*rerio*). Animal Cognition. 2017; **20**(2):159–169. DOI: 10.1007/s10071-016-1034-x
- [39] Zuur AF, Ieno EN, Elphick CS. A protocol for data exploration to avoid common statistical problems. Methods in Ecology and Evolution. 2010;**1**(1):3–14. DOI: 10.1111/j.2041-210X.2009.00001.x
- [40] Ripley B, Venables B, Hornik K, Gebhardt A, Firth D. Package ‘MASS’[Internet]. 2013. Available from: http://www.trichord-inc.com/pricing/frames/content/solar_power.pdf [Accessed: 05-01-2017]
- [41] Team RC. R: A language and environment for statistical computing [Internet]. R Foundation for Statistical Computing. 2015. Available from: <https://www.r-project.org/> [Accessed: 05-01-2017]
- [42] Lenth RV, Hervé M. lsmeans: Least-Squares Means. R package version 2.11 [Internet]. 2014. Available from: <http://CRAN.R-project.org/package=lsmeans> [Accessed: 05-01-2017]
- [43] De Luca MA, Bassareo V, Bauer A, Di Chiara G. Caffeine and accumbens shell dopamine. Journal of Neurochemistry. 2007;**103**(1):157–163. DOI: 10.1111/j.1471-4159.2007.04754.x
- [44] Richendrfer H, Pelkowski SD, Colwill RM, Créton R. Developmental sub-chronic exposure to chlorpyrifos reduces anxiety-related behavior in zebrafish larvae. Neurotoxicology and Teratology. 2012;**34**(4):458–465. DOI: 10.1016/j.ntt.2012.04.010
- [45] Angelucci MEM, Cesário C, Hiroi RH, Rosalen PL, Da Cunha C. Effects of caffeine on learning and memory in rats tested in the Morris water maze. Brazilian Journal of Medical and Biological Research. 2002;**35**(10):1201–1208. DOI: 10.1590/S0100-879X2002001000013

- [46] Dunwiddie TV, Masino SA. The role and regulation of adenosine in the central nervous system. *Annual Review of Neuroscience*. 2001;**24**(1):31–55. DOI: 10.1146/annurev.neuro.24.1.31
- [47] Daly JW, Fredholm BB. Caffeine—an atypical drug of dependence. *Drug and Alcohol Dependence*. 1998;**51**(1–2):199–206. DOI: 10.1016/S0376-8716(98)00077-5
- [48] Kirch DG, Taylor TB, Gerhardt GA, Benowitz NL, Stephen C, Wyatt RJ. Effect of chronic caffeine administration on monoamine and monoamine metabolite concentrations in rat brain. *Neuropharmacology*. 1990;**29**(6):599–602. DOI: 10.1016/0028-3908(90)90073-Z
- [49] Nehlig A. Is caffeine a cognitive enhancer. *Journal of Alzheimer's Disease*. 2010;**20**(S1): S85–94. DOI: 10.3233/JAD-2010-091315.
- [50] Ribeiro JA, Sebastiao AM. Caffeine and adenosine. *Journal of Alzheimer's Disease*. 2010;**20**(S1):3–15. DOI: 10.3233/JAD-2010-1379.
- [51] Abreu RV, Silva-Oliveira EM, Moraes MF, Pereira GS, Moraes-Santos T. Chronic coffee and caffeine ingestion effects on the cognitive function and antioxidant system of rat brains. *Pharmacology Biochemistry and Behavior*. 2011;**99**(4):659–664. DOI: 10.1016/j.pbb.2011.06.010
- [52] de Aguiar RA, Turnes T, Cardoso TE, Vasconcelos DIC, Caputo F. Efeito da ingestão de cafeína em diferentes tarefas de tempo de reação. *Revista Brasileira de Ciências do Esporte*. 2011;**34**(2):465–476. DOI: 10.1590/S0101-32892012000200015.
- [53] Santos LC, Ruiz-oliveira J, Oliveira JJ, Silva PF, Luchiari AC. Pharmacology, biochemistry and Behavior. Irish coffee: Effects of alcohol and caffeine on object discrimination in zebrafish. *Pharmacology Biochemistry and Behavior*. 2016;**143**:34–43. DOI: 10.1016/j.pbb.2016.01.013
- [54] Glade MJ. Caffeine—not just a stimulant. *Nutrition*. 2010;**26**(10):932–938. DOI: 10.1016/j.nut.2010.08.004
- [55] Burgess JW. Chronic exposure to caffeine during early development modifies spatial behavior in juvenile jewel fish schools. *Pharmacology Biochemistry and Behavior*. 1982;**17**(1):137–140. DOI: 10.1016/0091-3057(82)90275-1
- [56] Nehlig A, Daval JL, Debry G. Caffeine and the central nervous system: Mechanisms of action, biochemical, metabolic and psychostimulant effects. *Brain Research Reviews*. 1992;**17**(2):139–170. DOI: 10.1016/0165-0173(92)90012-B
- [57] Ardais AP, Borges MF, Rocha AS, Sallaberry C, Cunha RA., Porciúncula LO. Caffeine triggers behavioral and neurochemical alterations in adolescent rats. *Neuroscience*. 2014;**270**:27–39. DOI: 10.1016/j.neuroscience.2014.04.003
- [58] Hughes RN, Hancock NJ. Strain-dependent effects of acute caffeine on anxiety-related behavior in PVG/c, Long–Evans and Wistar rats. *Pharmacology Biochemistry and Behavior*. 2016;**140**:51–61. DOI: 10.1016/j.pbb.2015.11.005

- [59] Jabbar SB, Hanly MG. Fatal caffeine overdose: A case report and review of literature. *American Journal of Forensic Medicine and Pathology*. 2013;**34**(4):321–324. DOI: 10.1097/PAF.0000000000000058
- [60] Rodriguez RS, Haugen R, Rueber A, Huang CC. Reversible neuronal and muscular toxicity of caffeine in developing vertebrates. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*. 2014;**163**:47–54.