We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Etiology, Immunopathogenesis and Biomarkers in Behçet's disease

Fahd Adeeb, Maria Usman Khan,
Austin G. Stack and Alexander D. Fraser

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.68342>

Abstract

Behçet's disease (BD) is a type of vasculitis with many distinctive clinical manifestations and multifactorial immunopathogenesis. The cause of BD remains unknown, but it has been postulated that in a genetically predisposed or susceptible population, exogenous agents trigger the dysregulation of both autoinflammatory and autoimmune responses resulting in multisystem vasculitis. There are robust ongoing efforts across the globe to elucidate and identify signature markers to improve and assist in rapid diagnosis of the disease and to tailor the best therapy accordingly. While association of human leukocyte antigen (HLA)-B*51 (B*51:01 subtype) allele is well recognized as the strongest genetic susceptibility gene so far among genetically predisposed BD patients, further investigations using the latest technology have led to the identification of several novel single nucleotide polymorphisms (SNPs) and other associated genes involved in the pathogenesis. There are several "established" cytokines known to be involved in the pathogenesis of BD, which have been further implicated in the genome-wide association studies (GWAS)-based cytokine/receptor gene loci studies, as well as numerous "novel" cytokines, which are currently being studied and identified. This chapter offers insights into current knowledge and thoughts regarding the future of biomarkers in BD.

Keywords: Behçet's disease, Behçet's syndrome, biomarker, immunopathogenesis, immunogenetics, pathogenetics, etiopathogenesis, cytokines, HLA-B*51

1. Introduction

Behçet's disease (BD) is a type of vasculitis characterized by recurrent inflammatory attacks causing many distinctive clinical manifestations, most commonly affecting the orogenital mucosa, skin, and the eyes [1–3]. The etiology has yet to be fully established, but it has been

postulated that a genetically predisposed or susceptible population, exposed to exogenous agents, may result in the dysregulation of both autoinflammatory and autoimmune responses. It is a complex disease that has been the subject of intense research and clinical interest.

In this era of precision medicine, there is a need to integrate biomarkers into clinical practice, which may serve as valuable predictive and prognostic tools to assist practicing physicians in making clinical decisions while managing complex diseases. This may also improve diagnostic capability, risk stratification, prediction of disease progression, assist in targeted therapy, and monitoring of response to treatment to improve patients' overall clinical outcome.

This review chapter will look into the most relevant articles that have defined and clarified BD over the years, as well as review the most recent publications offering new insights in order to help fill the gaps in further understanding the disease. The chapter will focus on the etiology, immunopathogenesis, recent advances in search of potential biomarkers and targets for further research.

We will examine recent advances made that have shed new light on an old disease and explore the potential genetic (including genome-wide association studies (GWAS) and next-generation DNA sequencing, looking into the human leukocyte antigen (HLA) and non-HLA genetic associations) and molecular markers (including innate immune lymphoid cells and adaptive immune cells, cytokines, chemokines, other circulating biomarkers, and signaling molecules of inflammation) and their potential correlations with disease activity and therapy.

2. The pursuit for genetic susceptibility markers in Behçet's disease

BD is a genetically complex and heterogeneous disease. The pursuit of gene discovery for causative genetic factors in BD spans over more than four decades since Professor Shigeaki Ohno first described the association of HL-A5 antigen observed in his Japanese BD cohort that was later renamed human leukocyte antigen (HLA)-B5 [4]. Other early evidence stems from the observation of familial clustering in BD families where more than one family member developed the disease [5–8], which provided further clues to a strong genetic predisposition to the disease.

2.1. The MHC I and HLA-B*51 and its association with BD

The major histocompatibility complex (MHC) has an expansive immune component including the HLA and plays a pivotal role in the genetic influences on susceptibility to autoimmunity. The ~3.5-Mb region has the highest density of genes in the human genome, the majority of which have fundamental roles in immunity [9].

A remarkably consistent body of evidence demonstrates association of HLA-B*51 (B*51:01 subtype) allele as the strongest genetic susceptibility gene so far among genetically predisposed BD patients. However, certain indigenous Amerindians have a high prevalence of HLA-B*51 but virtually no reported cases of BD. High level of recombination within the MHC is known

to have occurred in these Eastern populations before their migration into Beringia and it was suggested that disruption of the genetic loci in linkage disequilibria with HLA-B*51 might be one reason for the absence of disease in these high HLA-B*51-bearing populations [10]. These findings emphasize the fundamental roles and interplay of both genetic and environmental components in the development of the disease.

There have been conflicting views, however, on whether the disease association with HLA-B*51 is attributed to a role of MHC class I variant itself or if the association is found due to its linkage disequilibrium (LD) with another variant in the region [11].

2.2. The unifying concept of MHC-I-opathy

The understanding of the pathophysiology of BD is challenging, as it is at the crossroads between autoimmune and autoinflammatory syndromes [12]. A new perspective of a unifying concept of MHC-I-opathy was proposed recently comprising of BD and several clinically distinct spondyloarthropathies (such as ankylosing spondylitis and psoriasis)—all associated with MHC Class I alleles, such as HLA-B*51, HLA-B*27, and HLA-C*0602, and epistatic endoplasmic reticulum aminopeptidase 1 (ERAP-1) interactions.

McGonagle et al. have proposed that the MHC-I-opathies share an immunopathogenetic basis including barrier dysfunction in environmentally exposed organs such as the skin and aberrant innate immune reactions at sites of mechanical stress. This they argue can often trigger secondary adaptive immune CD8⁺ T cell responses with prominent neutrophilic inflammation that culminate in the initiation and chronicity of these diseases [13]. Further research and understanding into this unifying concept of MHC I driven inflammatory response may provide further targets for disease management.

2.3. Other MHC/HLA associations

The complexity and strong LD with the HLA-B*51 allele make it difficult to explore additional independent susceptibility loci within this region. Despite having the strongest genetic association, it remains unclear to this day whether disease susceptibility in BD is due to the HLA-B*51 itself or due to the genes located around the HLA which is in LD with it.

The MHC Class I chain-related gene A (MICA) is located in proximity and in between the HLA-B and tumor necrosis factor (TNF) genes on the short arm of chromosome 6. It has long been considered a major genetic susceptibility gene for BD and has been studied in many different populations since the first observation of a possible association by Mizuki et al. [14]. Lee et al. conducted a meta-analysis on the associations of MICA and BD and found statistically significant association in various ethnic populations [15]. However, due to its strong LD with HLA-B*51, it has been difficult to prove MICA as a primary susceptibility gene for BD. Furthermore, different GWAS did not find an independent association between MICA and BD [16–18].

Ombrello et al. performed stepwise conditional analysis and found independent genetic associations for HLA-B*15 and HLA-B*27 and risk, while HLA-B*49 and HLA-A*03 were protective

[19]. Montes-Cano et al. demonstrated HLA-B*57 as a marker for risk in the Spanish population [20], while Meguro et al. showed that HLA-A*26 is a risk marker in the Japanese population [16].

Using a custom platform (ImmunoChip) that includes 8572 single nucleotide polymorphisms (SNPs) in the HLA extended region, Hughes et al. demonstrated that the robust HLA-B*51 association in BD is due to a strong association signal of an SNP rs116799036 in two independent BD cohorts (Turkish and Italian) from two ancestry groups, while also identifying two additional independent genetic associations with genome-wide significance ($P < 5 \times 10^{-8}$) in the HLA region: rs12525170 and rs114854070 [21].

2.4. Genomic strategies for a complex genetic disease beyond the typical Mendelian inheritance

It was difficult to venture beyond the MHC in the past: the tools to do so were unavailable as linkage studies were only suitable for Mendelian disorders. Intriguing new techniques have surfaced in recent times accelerating the pace for gene discovery, especially for complex diseases such as BD.

2.4.1. Genome-wide association studies (GWAS)

Being a complex disease, BD does not follow the typical Mendelian law of inheritance, rather of a dichotomous nature conforming the “polygenic threshold model” where the phenotypic expression is resultant upon genetic variation at multiple rather than a single loci, with the majority of the cases occurring sporadically. This is the basis foundation for the development of GWAS, which was initially thought to be the “comprehensive” option designed to identify genetic variants associated with such complex disease [22].

The advent of GWAS earlier in this century has dramatically improved our ability to identify and map successfully susceptibility loci associated with complex diseases such as BD, usually as single nucleotide polymorphisms (SNPs). It seems that these common genetic variants contribute to polygenic disease manifestations where phenotypic variance depends on contributions from several genetic variance.

Fei et al. [23] performed the first GWAS study in BD in a relatively small Turkish population and identified several novel candidate genetic loci (KIAA1529, CPVL, LOC100129342, UBASH3B, and UBAC2) that are associated with increased susceptibility to BD. In the same year, Meguro et al. [16] found that the main susceptibility locus in BD Japanese population remains in the MHC itself, wherein reside two independent loci: HLA-B*51 and HLA-A*26. Two large GWAS conducted in Turkey and Japan followed in 2010: Remmers et al. confirmed the association of HLA-B*51, identified a second, independent association within the MHC Class I region and also found association at IL10 [17], while Mizuki et al. identified IL23R-IL12RB2 and IL10 as Behçet's disease susceptibility loci [18].

In 2012, GWAS performed by Kirino et al. identified novel susceptibility loci at chemokine receptors CCR1-CCR3, signal transducer and activator of transcription 4 (STAT4), killer cell lectin-like receptor K1 subfamily K, member 1 (KLRK-1), killer cell lectin-like receptor

subfamily C, member 4 (KLRC4), and ERAP1 in a Turkish population [24], thus apparently supporting the emerging concept delineating common pathogenic mechanisms for BD, ankylosing spondylitis, and psoriasis. In the same year, Hou et al. also identified STAT4 as a novel susceptibility locus for BD in the Chinese population in their GWAS and functional studies [25].

One of the difficulties of using GWAS in cohorts of mixed ethnicity is due to the rarity and unequal distribution of disease prevalence among different ethnic background. This was overcome by novel statistical approaches, demonstrated by Kappan et al. [26] who confirmed the central role of the HLA region in the disease and validated the association of IL2A gene by meta-analysis with previous work.

2.4.2. Next-generation sequencing (NGS) and candidate gene analysis

Despite discoveries of many unimpeachable associations with GWAS, it became apparent that the approach alone could not explain the full range of heritability or genetic susceptibilities to complex diseases [27] and at best could only identify moderate proportions of genetic variants contributing to the disease heritability. Among the growing menu of techniques, targeted next-generation sequencing is the latest promising technology in search of rare genetic variants with fewer alleles (minor-allele frequency), which likely carry a greater larger impact on disease manifestations and with larger deleterious biological effects [28, 29]. Next-generation exome sequencing has the ability to generate millions of short reads of sequence, ranging from 50 to 500 bp, in parallel. It can be targeted in key regions of the genome-to make quick discoveries [29].

2.5. Non-MHC I susceptibility genes

Early data revealing the contributory influence of non-HLA susceptibility genes came from studies in the 1990s; however, some of the associations were weak or inconclusive. In 2009, Karasneh et al. published the first systematic whole genome linkage analysis from 28 Turkish BD families, which provided evidence for non-HLA susceptibility loci with the strongest evidence seen for 12p12–13 and 6p22–24 [30].

The revelation of GWAS unfolded associations with genome-wide significance ($P < 5 \times 10^{-8}$) in the interleukin 23 receptor (IL23R)-IL12RB2, IL12A, IL10, ubiquitin-associated domain containing 2 (UBAC2), STAT4, CCR1-CCR3, KLRC4, ERAP1, TNF alpha-induced protein 3 (TNFAIP3), and fucosyltransferase 2 (FUT2) loci [17, 18, 22–26]. Some of these results have been replicated since in case-control candidate gene studies from Iranian, Chinese and Spanish European populations [30–36].

Targeted next-generation sequencing revealed the additional involvement of rare non-synonymous variants in toll-like receptor 4 (TLR4), nucleotide-binding oligomerization domain-containing protein 2 (NOD2), and the Mediterranean Fever Gene (MEFV) [37]. Recently, Oggenovski et al. used whole exome sequencing in BD of European descent for the first time and identified and replicated two novel putative protein-damaging genetic variants within LIMK2 and NEIL1, which may influence cytoskeletal regulation and DNA repair [38].

The associations with ERAP1, IL23R, IL10, and MEFV variations suggest that BD may share susceptibility genes and inflammatory pathways with spondyloarthritis [39], while the TLR and FUT polymorphisms that affect response to invasive pathogens have led to an increase interest in responses to microbiomes [40].

3. Molecular markers of BD

3.1. The cytokines network of BD

The term “cytokine” was first introduced in 1974 by Cohen et al. [41] to describe a polypeptide mediator superfamily central in the immune system generation and regulation. An entwined network comprising of interleukins (ILs), interferons (IFNs), tumor necrosis factor (TNF), chemokines, and other mediators primed to regulate the immune system, however, due to several factors such as imbalance of its receptor expression and dysregulation of its functions, generates the pathologic systemic inflammatory and/or immune responses seen in various autoimmune and autoinflammatory disorders.

Pro-inflammatory and anti-inflammatory cytokines have been shown to be involved in patients with BD (as discussed in more detailed below). Several studies demonstrated elevated levels of cytokines in local lesions indicating its involvement in the disease local immune responses [42–46]. Evidence from the GWASs further implicated several cytokines underlying the pathogenesis of BD [17, 18, 22–26]. Moreover, the successful use of various anti-cytokine therapies in BD patients provides additional evidence that cytokines play a crucial role in its pathogenesis [47–64]. These overall observations highlight the fundamental role of cytokines as key players in the pathogenesis of BD.

There are several established cytokines that are known to be involved including IL1 β , TNF α , IL6, IL10, and IL23. Various new promising candidate's cytokines identified to be associated with BD include IL21, IL22, IL33, IL37, and several others, all of which be described as detailed below.

3.1.1. The main pro-inflammatory cytokines

Despite the pleiotropic nature of most cytokines, this group of cytokines primarily promotes inflammation. Several pro-inflammatory cytokines have been implicated in BD.

3.1.1.1. The interleukin-1 (IL1) family: IL1 β , IL18, IL33

All cells of the innate immune system express and/or are affected by IL1 family members, which play a key role in the differentiation and function of polarized innate and adaptive lymphoid cells [65]. Among the 11 cytokines in the family, IL1 β is the principal pro-inflammatory cytokine, leading to the expression of many chemokines and secondary mediators of inflammation and upregulating innate immunity in response to infectious agents [66]. The levels of IL1 β have been shown to be elevated in several studies [45, 46, 49, 67, 68], including in synovial

fluids of BD patients [45, 46]. A proof-of-concept study by Güll et al. strongly implicated IL1 β and BD with significant improvement seen especially in patients with uveitis treated with IL1 β -regulating antibody [47]. Recently Tugal-Tutkun et al. demonstrated rapid control of uveitis in BD patients without the need for high-dose corticosteroid in a prospective, open-label, randomized phase II trial [48] supporting several other previous studies of the proven efficacy of IL1 β to induce stable clinical remission among BD patients [49–52].

Other pro-inflammatory cytokines in the IL1 family implicated in BD especially in the last decade include IL18 [42, 69–71], an important component of polarized Th1 cell and natural killer (NK) cell responses and of the interplay between macrophages and NK cells [65]. Identified in 2005 by Schmitz et al, IL33 was noted to have the capability to activate NF- κ B and MAP kinases, and induces the expression of IL-4, IL-5, and IL-13 in vivo, leading to severe pathological changes in mucosal organs [72]. In several of his studies, Hamzaoui et al. demonstrated higher levels of IL33 in sera of active BD patients compared to BD patients in remission [73–75], and this was supported by Kim et al. [76] who found elevated IL33 in BD patients with erythema nodosum (EN) and EN-like skin lesions. Surprisingly, Koca et al. found contrasting results of lower IL33 levels in active BD Turkish patients compared to the inactive patients and healthy controls (HC) but did find significantly higher levels of IL33 among BD patients with uveitis [77].

3.1.1.2. *The tumor necrosis factor (TNF) superfamily*

Among the 19 TNF superfamily cytokines that has so far been identified, the first member of the family, TNF α , which was the first to be discovered, is the most highly investigated. Levels have been shown to be elevated in studies from different populations [49, 69, 78–83]. Meta-analysis by Touma et al. [84] found TNF (-238A/G, -1031C/T, and -857T/C) polymorphisms are associated with susceptibility to BD, while an updated meta-analysis by Zhang et al. confirmed a significant association between the TNF-308A/G polymorphism and BD susceptibility [85]. Treatment has also been shown to be highly effective [53, 54].

There is not much information about the other members of the TNF superfamily and its association with BD, but despite the limitation, isolated studies have shown several other TNF family member to be associated with BD in different ethnic populations: Cantarini et al. found significantly higher serum soluble TNFR and soluble CD40L [86], Shaker et al. demonstrated higher levels of B cell activating factor (BAFF), A proliferation producing ligand (APRIL), and B cell maturation antigen (BCMA) in BD patients [87] and Düzgün et al. found elevated soluble CD30 levels in active BD patients compared to controls [88].

Besides TNF α , other members of the TNF superfamily may offer options as potential targets and therapeutic candidates in BD; however, more research is needed in this field to prove its efficacy and safety profile.

3.1.1.3. *Interleukin-6 (IL6)*

Several studies have shown higher levels of serum IL6 in active BD compared to inactive BD and HC [71, 89, 90], and interestingly in neuro-BD patients, IL6 was noted to be markedly

elevated in the cerebrospinal fluid (CSF), but not in the sera [91, 92]. Blockage of IL6 signaling with tocilizumab in BD patients despite looking promising in the treatment of neuro-BD [55–58] has revealed mixed results for non-neurological manifestations [58, 59, 93, 94] and is currently undergoing further evaluation in controlled clinical trials.

3.1.2. *The main anti-inflammatory cytokine: type I interferon (IFN α)*

Being the oldest cytokine discovered exactly 60 years ago, interferon- α (IFN α) has been the scope of investigation in many inflammatory diseases including BD. Despite initially thought to have mainly pro-inflammatory effects, it is becoming clearer that IFN α display a more complex function and its anti-inflammatory properties have led to its use as one of the treatment modalities in BD since the mid-1980s. Different studies by Hamzaoui, Kötter, and Pay et al. in their respective Tunisian, German and Turkish populations demonstrated higher levels of IFN α among BD patients [71, 95, 96]. In 2010, Liu et al. published their in vitro experiments demonstrating the ability of IFN α to inhibit IL17 expression and increase IL10 production by PBMCs and CD4⁺ T cells [97]. Successful uses of IFN α -2a and -2b as treatment modalities have been reported [60–62], and more recently, Lightman et al. reported the successful use of pegylated IFN α -2b in BD resulting significant reduction in corticosteroid use and improvement of quality of life [63]. The exact mechanism of IFN α , however, is still largely unknown.

3.1.3. *Helper and regulatory T lymphocytes involvement in BD*

There have been remarkable advances leading to our current understanding on the lineage commitment and plasticity of helper CD4⁺ T cells. The “naïve” CD4⁺ T cells in the presence of its associated cytokines differentiates into distinct T helper (Th) cells populations-Th1, Th2, Th17 or Th22: tailoring their responses to address specific threats accordingly. On the other hand, CD4⁺ CD25⁺ regulatory T cells (Tregs), derived from the thymus or differentiation from naïve T cells, downregulate Th responses and are critical for the preservation of immune tolerance and maintaining balance in the immune system.

3.1.3.1. *Th1, Th17, and Th22 cell-associated cytokines*

In the early 1980s, Ohno et al. demonstrated for the first time significantly higher levels of IFN γ in Japanese BD population [98]. Ahn et al. in their case series showed that the levels of IFN γ were elevated in aqueous humor and serum in BD patients with uveitis, which was then suppressed with combined low-dose cyclosporine/prednisone treatment [99]. Many other studies similarly found elevated serum IFN γ in active BD patients [44, 71, 83, 100–106], especially in BD patients with uveitis [44, 83, 102–105].

Other Th1 cell-associated cytokines associated with BD include IL2 and IL12. Despite conflicting results for IL2 levels in the ocular fluid of active BD patients with uveitis [106, 107], it was found to be significantly elevated in the serum of BD patients [108] and in active disease [109]. The alpha-chain of the IL2R that is shed from the surface of T cells by proteolytic enzymes to form the soluble sIL2R, which retains affinity to IL2, is also found to be significantly higher in active BD [46, 79, 110–113] and specifically in BD patients with uveitis [112, 113]. Serum IL12

levels were also found to be elevated in BD patients with active uveitis [114–116] and other active manifestations [42, 117, 118].

IL23 influences Th17 cell responses but shares a common p40 subunit with IL12 [119], and like IL12 has also been shown to be elevated in BD patients with active disease [101, 105, 120, 121]. Ustekinumab, a therapeutic agent, targeting both IL12 and IL23 cytokines has been shown to be therapeutic in BD [122], and subsequently a phase 2 open-label study to evaluate the proof-of-concept of ustekinumab in BD (STELABEC) has been recently registered in France. Both IL12 and IL23 stimulates nonreceptor Janus kinase 2 (JAK2) and tyrosine kinase 2 (TYK2) activity, leading to phosphorylation of STAT family members, with IL12 particularly activating STAT4 homodimers and IL23 predominantly activating STAT3 [119, 123–125]. Tulunay et al. demonstrated that the JAK1/STAT3 signaling pathway is activated in BD, and several other studies have shown similar findings [126]. Tulunay further suggested that more direct therapies aimed at JAK/STAT-associated cytokines such as ustekinumab (anti-IL12/23) and recently the approved tofacitinib that specifically inhibits JAK1/3, may be new therapeutic options for BD [126].

Th17 and Th22 are the “newer” helper-T cell subsets that secrete pro-inflammatory cytokines IL17 and IL22, respectively. Both are also implicated in the pathogenesis of BD, and their levels were markedly increased in BD patients [97, 105, 127–133] including active uveitis [107, 127, 128]. Chi et al. in their study demonstrated that production of IL17 was successfully inhibited by treatment with cyclosporine [127]. Another interesting finding in one of the studies above is increased levels of CCL20, an essential potent chemoattractant for the recruitment of Th17 lymphocytes [129]. Sugita et al. established Th22-type T cell clones from ocular samples taken from BD patients with active uveitis, which produced large amounts of IL22 and TNF α [106]. Sugita also demonstrated that IL22 in the presence of retinal antigens were able to produce high levels of IL22 in mice with experimental autoimmune uveitis [106]. From the therapeutic point of view, Liu et al. demonstrated significantly higher levels of IL17 in active BD patients, and stimulation with IFN α significantly decreases this IL17 production [97].

IL21 is one of the more recently identified type I cytokines that has been shown to tilt the balance between Th17 cells and regulatory T cells (Tregs) [134]. Geri et al. found markedly increased IL21 in active BD patients' sera and in the CSF of active neuro-BD patients. He further demonstrated increased Th17 and Th1 differentiation and decreased frequency of Tregs cells after stimulation of CD4⁺ T cells with IL21. Conversely, IL21 blockade with an IL21R-Fc restored the Th17 and Tregs homeostasis in BD patients, which might represent a potential target for novel therapy [135].

3.1.3.2. Th2 cell-associated cytokines and Tregs

The studies on Th2 cell-associated cytokines and their contribution in the pathogenesis of BD have been rather conflicting. Several studies found lower or no significant differences of the related cytokines in BD patients compared to HC [45, 106, 109, 119, 136, 137]. However, studies by Hamzaoui et al. found increased serum levels of Th2 (IL4 and IL13) cytokines, and Takeuchi et al. found elevated IL4 and IL10 in BD patients [71, 83], while studies from Raziuddin and Aridogan et al. demonstrated high levels of IL4, IL10 and IL13 in active BD

patients [138, 139]. Liu et al. and Guenane et al. in separate studies found significantly higher levels of IL10 in uveitis patients with BD compared to HC and idiopathic uveitis, respectively [97, 114]. Dalghous and his colleagues observed the presence of IL4 cytokines in oral lesions only from BD patients compared to RAS patients [140], while Ben Ahmed et al. found elevated levels of IL10 comparable to the increased IFN γ levels in active lesions of BD patients [141].

The possible role of Tregs in the pathogenesis of BD has gained considerable interest in recent times. Hamzaoui and Gündüz both demonstrated decreased Tregs level in clinically active BD patients [43, 142], Nanke et al. suggested that a decreased percentage of Tregs in peripheral blood of BD patients might be a predictive marker of ocular attack [143], while Sugita et al. demonstrated that Tregs level increased significantly with infliximab therapy but not with colchicine or cyclosporine in BD patients with uveitis [144]. Another subset of Tregs expresses high levels of CD52 glycoprotein [145], and together with other CD52-bearing cells (T cells, B cells, monocytes, macrophages, NK cells, dendritic cells, and granulocytes) are molecular targets of CAMPATH-1. A humanized antibody of IgG1 CAMPATH-1H/alemtuzumab has been successfully used in BD [146, 147] including in an open trial involving 18 BD patients with complete or partial remission achieved in 84% of patients [146].

3.1.3.3. *Gammadelta ($\gamma\delta$) T cells*

Gammadelta ($\gamma\delta$) T cells are innate-like lymphocytes that express a unique T cell receptor (TCR) γ and δ chain. Despite constituting only a small proportion (1–5%) of lymphocytes, they are more widespread within epithelial-rich tissues, such as the skin, intestine and reproductive tract, where they can comprise up to 50% of T cells [148]. They are a unique population of T cells that have features of both innate and adaptive immunity and express characteristics of conventional T cells, natural killer cells, and myeloid antigen presenting cells [149].

The relationship between $\gamma\delta$ T cells and BD has been noted in several studies since the early 1990s. Increased $\gamma\delta$ T cells levels were seen in BD patients compared to HC [99, 150–155] and in active BD compared to inactive BD [99, 151, 155, 156]; however conversely, several studies did not show any significant difference with HC [157–159]. Hasan et al. postulated that these discrepancies might be due to the activation status of the disease, as a reflection of local tissue inflammation compared to peripheral blood $\gamma\delta$ T cells and such variation might be dependent on several other factors including disease severity, usage of medications such as immunomodulatory agents, and perhaps other variables, namely, age, gender, ethnicity, and/or environmental factors [149]. Their roles in the pathogenesis and potentially as therapeutic targets remain to be elucidated.

3.1.4. *Other cytokines*

IL37 is part of the IL1 family (discovered in 2009 and formerly identified as IL1F7) but has emerged as an inhibitor of innate immunity [160]. It has been shown to be significantly lower in BD patients compared to HC, with pronounced inhibition in active patients, and was associated with increased production of IL1 β , IL6, and TNF α in LPS-stimulated PBMCs [161, 162]. Furthermore, in vitro experiments revealed that supplementing IL37 in BD patients significantly

suppresses these three pro-inflammatory cytokines [163]. There have also been suggestions of associations between BD and other cytokines such as IL15 and IL27; however, the data are still inadequate and sparse [44, 164–166].

3.1.5. Chemokines

The attraction of leukocytes to tissues is essential for inflammation and is controlled by chemokines, which are chemotactic cytokines [167]. Saruhan-Direskineli et al. observed significantly higher α -chemokine CXCL10/IP10 CSF levels in neuro-BD patients compared to patients with non-inflammatory neurological disease (NIN) and multiple sclerosis, whereas CXCL8/IL8 was increased in neuro-BD compared to NIN [168]. El-Asrar et al. found higher levels of CXCL9/MIG, CXCL10/IP10 and CXCL11 in BD patients' serum with uveitis [169], while its receptors CXCR3 expression were observed by Dalghous et al. to be higher in oral lesions biopsied from BD patients [140]. Recently, Ambrose et al. demonstrated significantly higher production of CXCL10/IP10 in blood monocytes of BD patients stimulated with IFN γ compared to HC, rheumatoid arthritis, and systemic lupus erythematosus controls [170]. There is even more robust evidence for CXCL8/IL8, a potent neutrophil chemoattractant, being implicated in BD pathogenesis, with some of the authors proposing that it could be a marker for vascular involvement and a more reliable marker for disease activity than the C-reactive protein or erythrocyte sedimentation rate [171–174].

In regard to β -chemokines, Ozer et al. found significantly elevated levels of MCP1/CCL2, MIP1 α /CCL3, and RANTES/CCL5 in active BD serum than in HC [175]. Similarly, Kökçam and Kim and their respective colleagues in two separate studies demonstrated high levels of MIP1 α /CCL3 [176, 177] while Kaburaki et al. and Do et al. found higher levels of MCP1/CCL2 in BD patients compared to HC [178, 179]. In CSF of neuro-BD patients, Saruhan-Direskineli et al. and Miyagishi et al. both demonstrated significantly higher levels of MIP1 α /CCL3 compared to NIN [168, 180].

The emerging evidence of a complex cytokine and chemokine network interplay involved in the pathogenesis of BD, the identification of candidate gene including cytokine polymorphisms and the proven potency of anti-cytokines treatment shed more light on the fundamental role of cytokines and chemokines in BD. Perhaps cytokine and/or chemokine gene therapy, which has been used in cancer therapy, though not extremely impressive but nonetheless promising, may offer a novel yet powerful approach in the treatment of BD in the future.

3.2. The innate immune network

3.2.1. Neutrophil hyperfunction and endothelial cell activation

Becatti et al. demonstrated significant enhancement in leukocyte reactive oxygen species (ROS) production particularly by neutrophils in BD patients and only neutrophil-derived ROS (but not lymphocyte- or monocyte-derived ROS) showed a significant correlation with fibrinogen carbonyl content, highlighting neutrophil activation as the promoter of fibrinogen oxidation and thrombus formation in BD [181] supporting similar finding in several previous

studies [182, 183]. Increasing evidence supports a role for neutrophil/lymphocyte ratio (NLR) as a cheap and simple disease activity marker in BD. NLR has been proposed as a surrogate marker for endothelial dysfunction and inflammation, and several authors have proposed the use of NLR as part of evaluation for disease activity in BD [184–187].

3.2.2. Innate lymphoid cells (ILCs)

Groundbreaking studies over recent years have formally identified innate lymphoid cells (ILCs) as a distinct arm of the innate immune system, comprising of the classic cytotoxic natural killer (NK) cells, lymphoid tissue inducer cells, and non-cytotoxic ILC populations [188]. Besides controlling tissue homeostasis, it has the ability to promote inflammation at mucosal and surface barriers [188]. Yamaguchi et al. reported that NK cells are actively involved in the induction and maintenance of disease remission in BD patients, through NK2 polarization [189] while Takeno et al. from their study concluded that abnormal killer inhibitory receptor (KIR) expression of NK cells may be associated with the development of BD [190]. Furthermore, ILCs have been recently shown as an important source of cytokines production [188, 191, 192]. The recent discovery of this latest group of diverse immune cells with its many emerging diverse roles in autoimmunity and inflammation may redefine or rather perhaps reinforce our understanding of the pathogenesis of BD in the future.

3.3. Inflammasomes

The inflammasome has been shown to be a key regulator of IL1 β and IL18 via direct activation of caspase-1 [173–194]. Liang et al. demonstrated production of IL-1 β was significantly decreased in ocular BD patients after the Nod-like receptor protein 3 (NLRP3) inflammasome was downregulated [195] while Kim et al. showed that the basal and LPS-induced expressions of NLRP3 inflammasome components were significantly increased at both mRNA and protein levels in BD patients [196]. Conversely, Türe-Özdemir and his team were not able to find any difference in DC and neutrophils of BD patients compared to HC after stimulating caspase-1 activation [197].

3.4. Autoantibodies

There have been several studies implicating certain autoantibodies in BD, but all are neither non-specific nor sensitive and are of limited clinical significance. Among them the most described were anti-*Saccharomyces cerevisiae* antibodies (ASCAs) [198–201] and anti-endothelial cell antibodies (AECAs) [202–205]. ASCAs were linked more to gastrointestinal manifestations in BD [198–202] and their healthy relatives [201]. AECA has been implicated with vascular [202, 204] and intestinal [205] involvement in BD and several studies demonstrated that alpha-enolase is the target antigen of AECA in BD patients [203–205]. These positive results, however, were not replicated in certain other ethnic populations [206–208].

3.5. The host-microbe interaction in BD

The role of microbial triggers in BD has long been postulated since the disease was first described. Microbial heat shock proteins (HSPs) show significant homology with human

mitochondrial HSP and molecular mimicry is suggested as the mechanism of pathology exacerbating BD when patients were exposed to these foreign antigens. While both streptococci and herpes simplex virus have garnered the most particular interest among researchers [209–214], evidence of exposure to other microbes such as staphylococci and mycobacteria has also been reported [215, 216]. Nonetheless, conflicting reports have been published and a specific pathogen has yet to be identified.

3.5.1. Microbiome: the rapidly re-emerging hypothesis in inflammatory disorders

Microbiome is a term first described by Lederberg in 2001 describing the microbial ecosystem [217], but it was Metchnikoff more than a century ago who hypothesized that the microbiota might influence the balance between pro-inflammatory and regulatory host responses and that alterations in the composition of the microbiota (a process that is known as dysbiosis) could jeopardize host immune responses [218]. Recent evidence indicates the possible contribution of the intestinal microbiota to immunological diseases outside the gut [218, 219].

The advent of the 16S ribosomal RNA (16S rRNA) sequencing technology over the past quarter century has identified a comprehensive human microbiota far more comprehensive than we ever imagined. Despite the emergence of newer application such as metagenomics [220], due to the nature of the rRNA genes that are highly conserved and evolutionarily stable but differ in their hypervariable regions enables identification of species, and owing to a confluence of methodological advancements, 16S rRNA has re-emerged as a stand-alone molecular tool [221].

BD patients seem to exhibit specific microbiome signature. Consolandi et al. compared fecal microbiota of BD patients and HC and found significantly depleted *Roseburia* and *Subdoligranulum* genera and butyrate production in BD patients [222] while Shimizu et al. also demonstrated gut dysbiosis in BD patients with significantly increased genera *Bifidobacterium* and *Eggerthella* and decreased genera *Megamonas* and *Prevotella* compared to HC [223].

Several authors also identified salivary dysbiosis among BD patients. Seoudi et al. found in BD patients an increased colonization of *Rothia dentocariosa* at non-ulcer oral sites, while the ulcer sites were highly colonized with *Strep salivarius* compared to recurrent aphthous stomatitis (RAS) patients, and with *Strep sanguinis* compared to HC, who were more highly colonized with *Neisseria* and *Veillonella* [224]. Coit et al. found significantly less diverse microbial structure in stimulated saliva samples in BD patients both with and without immunosuppressant [225].

3.6. Other possible markers

There are other possible molecular markers that have been or are still under investigations. Fecal calprotectin (FC) were demonstrated to be significantly elevated in intestinal BD in several studies [226–228], and interestingly Özşeker et al. demonstrated high fecal FC levels in asymptomatic but endoscopically proven BD patients with intestinal involvement [226]. Vascular endothelial growth factor (VEGF) levels have been observed in BD patients, particularly in active BD [229–232]. Several studies provided evidence for the increased levels of markers for endothelial activation or dysfunction such as vascular and intercellular adhesion molecules VCAM1, ICAM1, Selectins, and YKL40 in BD [233–238]. Various authors explored the association between certain genetic mutations and thrombosis in BD; however, results

have been inconclusive [239–244], and current data indicate that the pathogenesis of thrombosis in BD is not due to a coagulation abnormality [245].

The immunomodulatory role of vitamin D is of increasing interest, and several *in vitro* studies have demonstrated downregulation of inflammation by vitamin D [246–249]. Moreover, hypovitaminosis D had been implicated in various inflammatory disorders including BD [250–255]. Further investigations from different ethnic populations may provide further insights to this potentially clinically relevant knowledge of vitamin D as a potential suppressor of inflammatory response in BD.

4. Gaps and future directions

Efforts to develop biomarkers in BD have been confounded by substantial impediments and challenges and the greatest is probably due to the rare nature of the disease, while others include the complex role of the susceptibility genes and related cytokines, chemokines and other signaling molecules, variability of duration and severity of the disease, as well as variations between different geographical areas and the limited number of patient samples.

Standardization and quality assurance are significant hurdles and collaboration between laboratories at different centers to standardize protocols and assays is essential. There are clear similarities and differences across different ethnic groups phenotypically and at a genetic or molecular level. So far, clinical data trials support the critical role of innate cytokines TNF, IL1, and IL6 in the development of inflammatory episodes of BD, and targeting T cells or B cells may provide favorable results [256]. In this era of personalized and precision medicine, collaborative efforts nationally or internationally are needed to assemble adequately powered cohorts to perform further population- or regional-based molecular and genetic studies.

One possible way to move forward is broadening the classification criteria to combine objective clinical indicators and biomarkers, but despite the emergence of these candidate markers, there is still a lack of sufficient widespread evidence to support their implementation and incorporation into the contemporary classification criteria. In the meantime, it must be noted that BD remains fundamentally a clinical diagnosis.

Many questions remain a conundrum including (1) which patients will develop a more severe form of disease, (2) who will be resistant to certain therapy, (3) which patients with recurrent aphthous ulcers will progress to develop BD, and (4) who will benefit the most from a particular therapy. The search remains a highly scientific priority, but until we find the biomarkers, likelihood is, many of these questions will remain uncertain.

5. Conclusions

It has been a long, challenging journey in search of biomarkers in Behçet's disease. There are clear genetic and molecular similarities and variability between different ethnic populations.

Collaborative efforts nationally or internationally are needed to assemble sufficiently powered sample size to perform further population- or regional-based molecular and/or genetic studies in the search for the elusive “magic bullet” as the signature marker that will revolutionize the field of BD.

Author details

Fahd Adeeb^{1,3*}, Maria Usman Khan^{1,3}, Austin G. Stack^{2,3} and Alexander D. Fraser^{1,3}

*Address all correspondence to: fahd_adeeb@yahoo.com

1 Department of Rheumatology, University Hospital Limerick, Limerick, Ireland

2 Department of Nephrology, University Hospital Limerick, Limerick, Ireland

3 Graduate Entry Medical School, University of Limerick, Limerick, Ireland

References

- [1] Davatchi F, Chams-Davatchi C, Shams H, Shahram F, Nadji A, Akhlaghi M, et al. Behçet's disease: Epidemiology, clinical manifestations, and diagnosis. *Expert Review of Clinical Immunology*. 2017;**13**:57–65
- [2] Fitzgerald CW, Adeeb F, Timon CV, Shine NP, Fraser AD, Hughes JP. Significant laryngeal destruction in a northern European cohort of Behçet's disease patients. *Clinical and Experimental Rheumatology*. 2015;**33**(6 Suppl 94):S123–S128
- [3] Emmungil H, Yaşar Bilge NŞ, Küçükşahin O, Kılıç L, Okutucu S, Gücenmez S, et al. A rare but serious manifestation of Behçet's disease: Intracardiac thrombus in 22 patients. *Clinical and Experimental Rheumatology*. 2014;**32**(4 Suppl 84):S87–S92
- [4] Ohno S, Aoki K, Sugiura S, Nakayama E, Itakura K. Letter: HL-A5 and Behçet's disease. *Lancet*. 1973;**2**:1383–1384
- [5] Vaiopoulos G, Sfrikakis PP, Hatzinikolaou P, Stamatelos G, Kaklamanis P. Adamantiadis- Behçet's disease in sisters. *Clinical Rheumatology*. 1996;**15**:382–384
- [6] Woodrow JC, Graham DR, Evans CC. Behçet's syndrome in HLA-identical siblings. *British Journal of Rheumatology*. 1990;**29**:225–227
- [7] Koné-Paut I, Geisler I, Wechsler B, Ozen S, Ozdogan H, Rozenbaum M, et al. Familial aggregation in Behçet's disease: High frequency in siblings and parents of pediatric probands. *Journal of Pediatrics*. 1999;**135**:89–93
- [8] Gül A, Inanç M, Ocal L, Aral O, Koniçe M. Familial aggregation of Behçet's disease in Turkey. *Annals of the Rheumatic Diseases*. 2000;**59**:622–625.

- [9] Muers M. Complex disease: Ups and downs at the MHC. *Nature Reviews Genetics*. 2011;**12**:456–457
- [10] Verity DH, Marr JE, Ohno S, Wallace GR, Stanford MR. Behçet's disease, the Silk Road and HLA-B51: Historical and geographical perspectives. *Tissue Antigens*. 1999;**54**:213–220
- [11] Takeuchi M, Kastner DL, Remmers EF. The immunogenetics of Behçet's disease: A comprehensive review. *Journal of Autoimmunity*. 2015;**64**:137–148
- [12] McGonagle D, McDermott MF. A proposed classification of the immunological diseases. *PLoS Medicine*. 2006;**3**(8):e297
- [13] McGonagle D, Aydin SZ, Gül A, Mahr A, Direskeneli H. 'MHC-I-opathy'-unified concept for spondyloarthritis and Behçet disease. *Nature Reviews Rheumatology*. 2015;**11**:731–740
- [14] Mizuki N, Ota M, Kimura M, et al. Triplet repeat polymorphism in the transmembrane region of the MIC-A gene: A strong association of six GCT repetitions with Behcet disease. *Proceedings of the National Academy of Sciences of the United States of America*. 1997;**94**:1298–1303
- [15] Lee YH, Song GG. Associations between major histocompatibility complex class I chain-related gene A polymorphisms and susceptibility to Behcet's disease. A meta-analysis. *Zeitschrift für Rheumatologie*. 2015;**74**:714–721
- [16] Meguro A, Inoko H, Ota M, Katsuyama Y, Oka A, Okada E, et al. Genetics of Behçet disease inside and outside the MHC. *Annals of the Rheumatic Diseases*. 2010;**69**:747–754
- [17] Remmers EF, Cosan F, Kirino Y, Ombrello MJ, Abaci N, Satorius C, et al. Genome-wide association study identifies variants in the MHC class I, *IL10*, and *IL23R-IL12RB2* regions associated with Behçet's disease. *Nature Genetics*. 2010;**42**:698–702
- [18] Mizuki N, Meguro A, Ota M, Ohno S, Shiota T, Kawagoe T, et al. Genome-wide association studies identify *IL23R-IL12RB2* and *IL10* as Behçet's disease susceptibility loci. *Nature Genetics*. 2010;**42**:703–706
- [19] Ombrello MJ, Kirino Y, de Bakker PI, Gül A, Kastner DL, Remmers EF. Behcet disease-associated MHC class I residues implicate antigen binding and regulation of cell-mediated cytotoxicity. *Proceedings of the National Academy of Sciences of the United States of America*. 2014;**111**:8867–8872
- [20] Montes-Cano MA, Conde-Jaldón M, García-Lozano JR, Ortiz-Fernández L, Ortego-Centeno N, Castillo-Palma MJ, et al. HLA and non-HLA genes in Behçet's disease: A multicentric study in the Spanish population. *Arthritis Research & Therapy*. 2013;**15**(5):R145
- [21] Hughes T, Coit P, Adler A, Yilmaz V, Aksu K, Düzgün N, et al. Identification of multiple independent susceptibility loci in the HLA region in Behçet's disease. *Nature Genetics*. 2013;**45**:319–324
- [22] Hirschhorn JN, Daly MJ. Genome-wide association studies for common diseases and complex traits. *Nature Reviews Genetics*. 2005;**6**:95–108

- [23] Fei Y, Webb R, Cobb BL, Direskeneli H, Saruhan-Direskeneli G, Sawalha AH. Identification of novel genetic susceptibility loci for Behçet's disease using a genome-wide association study. *Arthritis Research & Therapy*. 2009;**11**(3):R66
- [24] Kirino Y, Bertsias G, Ishigatsubo Y, Mizuki N, Tugal-Tutkun I, Seyahi E, et al. Genome-wide association analysis identifies new susceptibility loci for Behçet's disease and epistasis between HLA-B51 and ERAP1. *Nature Genetics*. 2013;**45**:202–207
- [25] Hou S, Yang Z, Du L, Jiang Z, Shu Q, Chen Y, et al. Identification of a susceptibility locus in STAT4 for Behçet's disease in Han Chinese in a genome-wide association study. *Arthritis & Rheumatology*. 2012;**64**:4104–4113
- [26] Kappen JH, Medina-Gomez C, Hagen PMv, Stolk L, Estrada K, Rivadeneira F, et al. Genome-wide association study in an admixed case series reveals *IL12A* as a new candidate in Behçet disease. *PLoS One*. 2015;**10**(3):e0119085
- [27] Kilpinen H, Barrett JC. How next-generation sequencing is transforming complex disease genetics. *Trends Genetics*. 2013;**29**:23–30
- [28] Nelson RM, Pettersson ME, Carlborg Ö. A century after Fisher: Time for a new paradigm in quantitative genetics. *Trends Genetics*. 2013;**29**:669–676
- [29] Tennessen JA, Bigham, AW, O'Connor TD, Fu W, Kenny EE, Gravel S, et al. Evolution and functional impact of rare coding variation from deep sequencing of human exomes. *Science*. 2012;**337**:64–69
- [30] Karasneh J, Gul A, Ollier WE, Silman AJ, Worthington J. Whole-genome screening for susceptibility genes in multicase families with Behcet's disease. *Arthritis & Rheumatology*. 2005;**52**:1836–1842
- [31] Lee YJ, Horie Y, Wallace GR, Choi YS, Park JA, Choi JY, et al. Genome-wide association study identifies GIMAP as a novel susceptibility locus for Behcet's disease. *Annals of the Rheumatic Diseases*. 2013;**72**:1510–1516
- [32] Xavier JM, Shahram F, Sousa I, Davatchi F, Matos M, Abdollahi BS, et al. FUT2: Filling the gap between genes and environment in Behçet's disease? *Annals of the Rheumatic Diseases*. 2015;**74**:618–624
- [33] Wu Z, Zheng W, Xu J, Sun F, Chen H, Li P, et al. IL10 polymorphisms associated with Behcet's disease in Chinese Han. *Human Immunology*. 2014;**75**:271–276
- [34] Xavier JM, Shahram F, Davatchi F, Rosa A, Crespo J, Abdollahi BS, et al. Association study of IL10 and IL23R–IL12RB2 in Iranian patients with Behcet's disease. *Arthritis & Rheumatology*. 2012;**64**:2761–2772
- [35] Li H, Liu Q, Hou S, Du L, Zhou Q, Zhou Y, et al. TNFAIP3 gene polymorphisms confer risk for Behcet's disease in a Chinese Han population. *Human Genetics*. 2013;**132**:293–300
- [36] Hou S, Shu Q, Jiang Z, Chen Y, Li F, Chen F, et al. Replication study confirms the association between UBAC2 and Behçet's disease in two independent Chinese sets of patients and controls. *Arthritis Research & Therapy*. 2012;**14**(2):R70

- [37] Kirino Y, Zhou Q, Ishigatsubo Y, Mizuki N, Tugal-Tutkun I, Seyahi E, et al. Targeted resequencing implicates the familial Mediterranean fever gene MEFV and the toll-like receptor 4 gene TLR4 in Behçet disease. *Proceedings of the National Academy of Sciences of the United States of America*. 2013;**110**:8134–8139
- [38] Ognenovski M, Renauer P, Gensterblum E, Kötter I, Xenitidis T, Henes JC, et al. Whole exome sequencing identifies rare protein-coding variants in Behçet's disease. *Arthritis & Rheumatology*. 2016;**68**:1272–1280
- [39] Gül A. Genetics of Behçet's disease: Lessons learned from genomewide association studies. *Current Opinion in Rheumatology*. 2014;**26**:56–63
- [40] Morton LT, Situnayake D, Wallace GR. Genetics of Behçet's disease. *Current Opinion in Rheumatology*. 2016;**28**:39–44
- [41] Cohen S, Bigazzi PE, Yoshida T. Similarities of T cell function in cell-mediated immunity and antibody production. *Cellular Immunology*. 1974;**12**:150–159
- [42] Nagafuchi H, Takeno M, Yoshikawa H, Kurokawa MS, Nara K, Takada E, Masuda C, Mizoguchi M, Suzuki N. Excessive expression of Txk, a member of the Tec family of tyrosine kinases, contributes to excessive Th1 cytokine production by T lymphocytes in patients with Behçet's disease. *Clinical & Experimental Immunology*. 2005;**139**:363–370
- [43] Hamzaoui K, Borhani Haghighi A, Ghorbel IB, Houman H. RORC and Foxp3 axis in cerebrospinal fluid of patients with neuro-Behçet's disease. *Journal of Neuroimmunology*. 2011;**233**:249–253
- [44] Ahn JK, Yu HG, Chung H, Park YG. Intraocular cytokine environment in active Behçet uveitis. *American Journal of Ophthalmology*. 2006;**142**:429–434
- [45] Pay S, Erdem H, Pekel A, Simsek I, Musabak U, Sengul A, et al. Synovial proinflammatory cytokines and their correlation with matrix metalloproteinase-3 expression in Behçet's disease. Does interleukin-1beta play a major role in Behçet's synovitis? *Rheumatology International*. 2006;**26**:608–613
- [46] Ertenli I, Kiraz S, Calgüneri M, Celik I, Erman M, Haznedaroglu IC. Synovial fluid cytokine levels in Behçet's disease. *Clinical and Experimental Rheumatology*. 2001;**19**(5 Suppl 24):S37–S41
- [47] Gül A, Tugal-Tutkun I, Dinarello CA, Reznikov L, Esen BA, Mirza A, et al. Interleukin-1 β -regulating antibody XOMA 052 (gevokizumab) in the treatment of acute exacerbations of resistant uveitis of Behçet's disease: An open-label pilot study. *Annals of the Rheumatic Diseases*. 2012;**71**:563–566
- [48] Tugal-Tutkun I, Kadayifcilar S, Khairallah M, Lee SC, Ozdal P, Özyazgan Y, et al. Safety and efficacy of gevokizumab in Patients with Behçet's disease uveitis: Results of an exploratory phase 2 study. *Ocular Immunology and Inflammation*. 2016;**30**:1–9
- [49] Düzgün N, Ayaşlıoğlu E, Tutkak H, Aydintuğ OT. Cytokine inhibitors: Soluble tumor necrosis factor receptor 1 and interleukin-1 receptor antagonist in Behçet's disease. *Rheumatology International*. 2005;**25**:1–5

- [50] Emmi G, Talarico R, Lopalco G, Cimaz R, Cantini F, Viapiana O, et al. Efficacy and safety profile of anti-interleukin-1 treatment in Behçet's disease: A multicenter retrospective study. *Clinical Rheumatology*. 2016;**35**:1281–1286
- [51] Cantarini L, Vitale A, Scalini P, Dinarello CA, Rigante D, Franceschini R, et al. Anakinra treatment in drug-resistant Behçet's disease: A case series. *Clin Rheumatol* 2015; **34**: 1293–301
- [52] Vitale A, Rigante D, Caso F, Brizi MG, Galeazzi M, Costa L, et al. Inhibition of interleukin-1 by canakinumab as a successful mono-drug strategy for the treatment of refractory Behçet's disease: A case series. *Dermatology*. 2014;**228**:211–214
- [53] Vallet H, Riviere S, Sanna A, Deroux A, Moulis G, Addimanda O, et al. Efficacy of anti-TNF alpha in severe and/or refractory Behçet's disease: Multicenter study of 124 patients. *Journal of Autoimmunity*. 2015;**62**:67–74
- [54] Arida A, Fragiadaki K, Giavri E, Sfrikakis PP. Anti-TNF agents for Behçet's disease: Analysis of published data on 369 patients. *Seminars in Arthritis & Rheumatology*. 2011;**41**:61–70
- [55] Addimanda O, Pipitone N, Pazzola G, Salvarani C. Tocilizumab for severe refractory neuro-Behçet: Three cases IL-6 blockade in neuro-Behçet. *Seminars in Arthritis & Rheumatology*. 2015;**44**:472–475
- [56] Shapiro LS, Farrell J, Borhani Haghighi A. Tocilizumab treatment for neuro-Behçet's disease, the first report. *Clinical Neurology and Neurosurgery*. 2012;**114**:297–298
- [57] Urbaniak P, Hasler P, Kretzschmar S. Refractory neuro-Behçet treated by tocilizumab: A case report. *Clinical and Experimental Rheumatology*. 2012;**30**(3 Suppl 72):S73–S75
- [58] Deroux A, Chiquet C, Bouillet L. Tocilizumab in severe and refractory Behçet's disease: Four cases and literature review. *Seminars in Arthritis & Rheumatology*. 2016;**45**:733–737
- [59] Calvo-Rio V, de la Hera D, Beltran-Catalan E, Blanco R, Hernandez M, Martinez-Costa L, et al. Tocilizumab in uveitis refractory to other biologic drugs: A study of 3 cases and a literature review. *Clinical and Experimental Rheumatology*. 2014;**32**(4 Suppl 84):S54–S57
- [60] Kotter I, Zierhut M, Eckstein A, Vonthein R, Ness T, Günaydin I, et al. Human recombinant interferon-alpha2a (rhIFN alpha2a) for the treatment of Behçet's disease with sight-threatening retinal vasculitis. *Advances in Experimental Medicine and Biology*. 2003;**528**:521–523
- [61] Alpsoy E, Durusoy C, Yilmaz E, Ozgurel Y, Ermis O, Yazar S, et al. Interferon alfa-2a in the treatment of Behçet disease: A randomized placebo-controlled and double-blind study. *Archives of Dermatology*. 2002;**138**:467–471
- [62] Calguneri M, Ozturk MA, Ertenli I, Kiraz S, Apraş S, Ozbalkan Z, et al. Effects of interferon alpha treatment on the clinical course of refractory Behçet's disease: An open study. *Annals of the Rheumatic Diseases*. 2003;**62**:492–493

- [63] Lightman S, Taylor SR, Bunce C, Longhurst H, Lynn W, Moots R, et al. Pegylated interferon- α -2b reduces corticosteroid requirement in patients with Behçet's disease with upregulation of circulating regulatory T cells and reduction of Th17. *Annals of the Rheumatic Diseases*. 2015;**74**:1138–1144
- [64] Arida A, Sfrikakis PP. Anti-cytokine biologic treatment beyond anti-TNF in Behçet's disease. *Clinical and Experimental Rheumatology*. 2014;**32**(4 Suppl 84):S149–S155
- [65] Garlanda C, Dinarello CA, Mantovani A. The interleukin-1 family: Back to the future. *Immunity*. 2013;**39**:1003–1018
- [66] Dinarello CA, van der Meer JW. Treating inflammation by blocking interleukin-1 in humans. *Seminars in Immunology*. 2013;**25**:469–484
- [67] Yosipovitch G, Shohat B, Bshara J, Wysenbeek A, Weinberger A. Elevated serum interleukin 1 receptors and interleukin 1B in patients with Behçet's disease: Correlations with disease activity and severity. *Israel Journal of Medical Sciences*. 1995;**31**:345–348
- [68] Hamzaoui K, Hamza M, Ayed K. Production of TNF- α and IL-1 in active Behcet's disease. *Journal of Rheumatology*. 1990;**17**:1428–1429
- [69] Oztas MO, Onder M, Gurer MA, Bukan N, Sancak B. Serum interleukin 18 and tumour necrosis factor-alpha levels are increased in Behcet's disease. *Clinical and Experimental Dermatology*. 2005;**30**:61–3
- [70] Musabak U, Pay S, Erdem H, Simsek I, Pekel A, Dinc A. Serum interleukin-18 levels in patients with Behçet's disease. Is its expression associated with disease activity or clinical presentations? *Rheumatology International*. 2006;**26**:545–550
- [71] Hamzaoui K, Hamzaoui A, Guemira F, Bessioud M, Hamza M, Ayed K. Cytokine profile in Behçet's disease patients. Relationship with disease activity. *Scandinavian Journal of Rheumatology*. 2002;**31**:205–210
- [72] Schmitz J, Owyang A, Oldham E, Song Y, Murphy E, McClanahan TK, et al. IL-33, an interleukin-1-like cytokine that signals via the IL-1 receptor-related protein ST2 and induces T helper type 2-associated cytokines. *Immunity*. 2005;**23**:479–490
- [73] Hamzaoui K, Kaabachi W, Fazaa B, Zakraoui L, Mili Boussen I, Haj Sassi F. Serum IL-33 levels and skin mRNA expression in Behcet's disease. *Clinical and Experimental Rheumatology*. 2013;**31**(Suppl. 77):6–14
- [74] Hamzaoui K, Bouali E, Hamzaoui A. Interleukin-33 and Behçet disease: Another cytokine among others. *Human Immunology*. 2015;**76**:301–306
- [75] Hamzaoui K, Borhani-Haghighi A, Kaabachi W, Hamzaoui A. Increased interleukin 33 in patients with neuro-Behcet's disease: Correlation with MCP-1 and IP-10 chemokines. *Cellular & Molecular Immunology*. 2014;**11**:613–616
- [76] Kim DJ, Baek SY, Park MK, Park KS, Lee JH, Park SH, et al. Serum level of interleukin-33 and soluble ST2 and their association with disease activity in patients with Behcet's disease. *Journal of Korean Medical Science*. 2013;**28**:1145–1153

- [77] Koca SS, Kara M, Deniz F, Ozgen M, Demir CF, Ilhan N, et al. Serum IL-33 level and IL-33 gene polymorphisms in Behçet's disease. *Rheumatology International*. 2015;**35**:471–477
- [78] Mege JL, Dilsen N, Sanguedolce V, Gul A, Bongrand P, Roux H, et al. Overproduction of monocyte derived tumor necrosis factor alpha, interleukin (IL) 6, IL-8 and increased neutrophil superoxide generation in Behçet's disease: A comparative study with familial Mediterranean fever and healthy subjects. *Journal of Rheumatology*. 1993;**20**:1544–1549
- [79] Sayinalp N, Ozcebe OI, Ozdemir O, Haznedaroğlu IC, Dünder S, Kirazli S. Cytokines in Behçet's disease. *Journal of Rheumatology*. 1996;**23**:321–322
- [80] El Menyawi M, Fawzy M, Al-Nahas Z, Edris A, Hussein H, Shaker O, et al. Serum tumor necrosis factor alpha (TNF- α) level in patients with Behçet's disease: Relation to clinical manifestations and disease activity. *Egypt Rheumatologist*. 2014;**36**:139–143
- [81] Kwok SK, Park SH, Park MK, Cho ML, Seo SH, Ju JH, et al. Upregulation of macrophage migration inhibitory factor (MIF) production from peripheral blood mononuclear cells (PBMCs) stimulated by tumor necrosis factor (TNF)- α in patients with Behçet's syndrome. *Journal of the Korean Rheumatism Association*. 2007;**14**:112–117
- [82] Jiang Z, Hennein L, Tao Y, Tao L. Interleukin-23 receptor gene polymorphism may enhance expression of the IL-23 receptor, IL-17, TNF- α and IL-6 in Behçet's disease. *PLoS One*. 2015;**10**(7):e0134632
- [83] Takeuchi M, Karasawa Y, Harimoto K, Tanaka A, Shibata M, Sato T, et al. Analysis of Th cell-related cytokine production in Behçet disease patients with uveitis before and after infliximab treatment. *Ocular Immunology and Inflammation*. 2016;**12**:1–10
- [84] Touma Z, Farra C, Hamdan A, Shamseddeen W, Uthman I, Hourani H, et al. TNF polymorphisms in patients with Behçet disease: A meta-analysis. *Archives of Medical Research*. 2010;**41**:142–146
- [85] Zhang M, Xu WD, Wen PF, Liang Y, Liu J, Pan HF, et al. Polymorphisms in the tumor necrosis factor gene and susceptibility to Behçet's disease: An updated meta-analysis. *Molecular Vision*. 2013;**19**:1913–1924
- [86] Cantarini L, Pucino V, Vitale A, Talarico R, Lucherini OM, Magnotti F, et al. Immunometabolic biomarkers of inflammation in Behçet's disease: Relationship with epidemiological profile, disease activity and therapeutic regimens. *Clinical & Experimental Immunology*. 2016;**184**:197–207
- [87] Shaker OG, Tawfic SO, El-Tawdy AM, El-Komy MH, El Menyawi M, Heikal AA. Expression of TNF- α , APRIL and BCMA in Behçet's disease. *Journal of Immunology Research*. 2014;**2014**:380405
- [88] Düzgün N, Ayaslioglu E, Tutkak H. Serum soluble CD30 levels in Behçet's disease. *Clinical and Experimental Rheumatology*. 2004;**22**(4 Suppl 34):S17–S20
- [89] Hamzaoui K, Hamzaoui A, Kahan A, Hamza M, Chabbou A, Ayed K. Interleukin-6 in peripheral blood and inflammatory sites in Behçet's disease. *Mediators of Inflammation*. 1992;**1**:281–285

- [90] Yamakawa Y, Sugita Y, Nagatani T, Takahashi S, Yamakawa T, Tanaka S, et al. Interleukin-6 (IL-6) in patients with Behçet's disease. *Journal of Dermatological Science*. 1996;**11**:189–195
- [91] Hirohata S, Isshi K, Oguchi H, Ohse T, Haraoka H, Takeuchi A, et al. Cerebrospinal fluid interleukin-6 in progressive Neuro-Behçet's syndrome. *Clinical Immunology and Immunopathology*. 1997;**82**:12–17
- [92] Akman-Demir G, Tüzün E, İçöz S, Yeşilot N, Yentür SP, Kürtüncü M, et al. Interleukin-6 in neuro-Behçet's disease: Association with disease subsets and long-term outcome. *Cytokine*. 2008;**44**:373–376
- [93] Diamantopoulos AP, Hatemi G. Lack of efficacy of tocilizumab in mucocutaneous Behçet's syndrome: Report of two cases. *Rheumatology*. 2013;**52**:1923–1924
- [94] Cantarini L, Lopalco G, Vitale A, Coladonato L, Rigante D, Lucherini OM, et al. Paradoxical mucocutaneous flare in a case of Behçet's disease treated with tocilizumab. *Clinical Rheumatology*. 2015;**34**:1141–1143
- [95] Kötter I, Koch S, Vonthein R, Rückwaldt U, Amberger M, Günaydin I, et al. Cytokines, cytokine antagonists and soluble adhesion molecules in patients with ocular Behçet's disease treated with human recombinant interferon-alpha2a. Results of an open study and review of the literature. *Clinical and Experimental Rheumatology*. 2005;**23**(4 Suppl 38):S20–S26
- [96] Pay S, Simsek I, Erdem H, Pekel A, Musabak U, Sengul A, et al. Dendritic cell subsets and type I interferon system in Behçet's disease: Does functional abnormality in plasmacytoid dendritic cells contribute to Th1 polarization? *Clinical and Experimental Rheumatology*. 2007;**25**(4 Suppl 45):S34–S40
- [97] Liu X, Yang P, Wang C, Li F, Kijlstra A. IFN-alpha blocks IL-17 production by peripheral blood mononuclear cells in Behcet's disease. *Rheumatology*. 2011;**50**:293–298
- [98] Ohno S, Kato F, Matsuda H, Fujii N, Minagawa T. Detection of gamma interferon in the sera of patients with Behçet's disease. *Infection and Immunity*. 1982;**36**:202–208
- [99] Ahn JK, Seo JM, Yu J, Oh FS, Chung H, Yu HG. Down-regulation of IFN-gamma-producing CD56+ T cells after combined low-dose cyclosporine/prednisone treatment in patients with Behçet's uveitis. *Investigative Ophthalmology & Visual Science*. 2005;**46**:2458–2464
- [100] Djaballah-Ider F, Chaib S, Belguendouz H, Talbi D, Touil-Boukoffa C. T cells activation and interferon- γ /nitric oxide production during Behçet disease: A study in Algerian patients. *Ocular Immunology and Inflammation*. 2012;**20**:215–217
- [101] Na SY, Park MJ, Park S, Lee ES. Up-regulation of Th17 and related cytokines in Behçet's disease corresponding to disease activity. *Clinical and Experimental Rheumatology*. 2013;**31**(3 Suppl 77):32–40
- [102] Shen H, Xia LP, Lu J. Elevated levels of interleukin-27 and effect on production of interferon- γ and interleukin-17 in patients with Behçet's disease. *Scandinavian Journal of Rheumatology*. 2013;**42**:48–51

- [103] Belguendouz H, Messaoudène D, Lahmar K, Ahmedi L, Medjeber O, Hartani D, et al. Interferon- γ and nitric oxide production during Behçet uveitis: Immunomodulatory effect of interleukin-10. *Journal of Interferon & Cytokine Research*. 2011;**31**:643–651
- [104] Mesquida M, Molins B, Llorenç V, Sainz de la Maza M, Hernandez MV, Espinosa G, et al. Proinflammatory cytokines and C-reactive protein in uveitis associated with Behçet's disease. *Mediators of Inflammation*. 2014;**2014**:396204
- [105] Chi W, Zhu X, Yang P, Liu X, Lin X, Zhou H, et al. Upregulated IL-23 and IL-17 in Behçet patients with active uveitis. *Investigative Ophthalmology & Visual Science*. 2008;**49**:3058–3064
- [106] Sugita S, Kawazoe Y, Imai A, Yamada Y, Horie S, Mochizuki M. Inhibition of Th17 differentiation by anti-TNF-alpha therapy in uveitis patients with Behçet's disease. *Arthritis Research & Therapy*. 2012;**14**:R99
- [107] Türkçüoğlu P, Arat YO, Kan E, Kan EK, Chaudhry IA, Koca S, et al. Association of disease activity with serum and tear IL-2 levels in Behçet disease. *Ocular Immunology and Inflammation*. 2016;**24**:313–318
- [108] Akdeniz N, Esrefoglu M, Keleş MS, Karakuzu A, Atasoy M. Serum interleukin-2, interleukin-6, tumour necrosis factor-alpha and nitric oxide levels in patients with Behçet's disease. *Annals of the Academy of Medicine, Singapore*. 2004;**33**:596–599.
- [109] Sugi-Ikai N, Nakazawa M, Nakamura S, Ohno S, Minami M. Increased frequencies of interleukin-2- and interferon-gamma-producing T cells in patients with active Behçet's disease. *Investigative Ophthalmology & Visual Science*. 1998;**39**:996–1004.
- [110] BenEzra D, Maftzir G, Kalichman I, Barak V. Serum levels of interleukin-2 receptor in ocular Behçet's disease. *American Journal of Ophthalmology*. 1993;**115**:26–30
- [111] Uchio E, Matsumoto T, Tanaka SI, Ohno S. Soluble intercellular adhesion molecule-1 (ICAM-1), CD4, CD8 and interleukin-2 receptor in patients with Behçet's disease and Vogt-Koyanagi-Harada's disease. *Clinical and Experimental Rheumatology*. 1999;**17**:179–184
- [112] Alpsoy E, Cayirli C, Er H, Yilmaz E. The levels of plasma interleukin-2 and soluble interleukin-2R in Behçet's disease: A marker of disease activity. *Journal of Dermatology*. 1998;**25**:513–516
- [113] Evereklioglu C1, Er H, Türköz Y, Cekmen M. Serum levels of TNF-alpha, sIL-2R, IL-6, and IL-8 are increased and associated with elevated lipid peroxidation in patients with Behçet's disease. *Mediators of Inflammation*. 2002;**11**:87–93
- [114] Guenane H, Hartani D, Chachoua L, Lahlou-Boukoffa OS, Mazari F, Touil-Boukoffa C. Production of Th1/Th2 cytokines and nitric oxide in Behçet's uveitis and idiopathic uveitis. *Journal Francais D'Ophtalmologie*. 2006;**29**:146–152
- [115] Belguendouz H, Messaoudene D, Hartani D, Chachoua L, Ahmedi ML, Lahmar-Belguendouz K, et al. Effect of corticotherapy on interleukin-8 and -12 and nitric oxide production during Behçet and idiopathic uveitis. *Journal Francais D'Ophtalmologie*. 2008;**31**:387–395

- [116] Ahmedi ML, Belguendouz H, Messaoudene D, Mesbah-Amroun H, Terahi M, Lahlou-Boukoffa OS, et al. Influence of steroid hormones on the production of two inflammatory markers, IL-12 and nitric oxide, in Behçet's disease. *Journal Francais D'Ophthalmologie*. 2016;**39**:333–340
- [117] Frassanito MA, Dammacco R, Cafforio P, Dammacco F. Th1 polarization of the immune response in Behçet's disease: A putative pathogenetic role of interleukin-12. *Arthritis & Rheumatology*. 1999;**42**:1967–1974
- [118] Turan B, Gallati H, Erdi H, Gürler A, Michel BA, Villiger PM. Systemic levels of the T cell regulatory cytokines IL-10 and IL-12 in Behçet's disease; soluble TNFR-75 as a biological marker of disease activity. *Journal of Rheumatology*. 1997;**24**:128–132
- [119] Teng MW, Bowman EP, McElwee JJ, Smyth MJ, Casanova JL, Cooper AM, et al. IL-12 and IL-23 cytokines: from discovery to targeted therapies for immune-mediated inflammatory diseases. *Nature Medicine*. 2015;**21**:719–729
- [120] Habibagahi Z, Habibagahi M, Heidari M. Raised concentration of soluble form of vascular endothelial cadherin and IL-23 in sera of patients with Behçet's disease. *Modern Rheumatology*. 2010;**20**:154–159
- [121] Gheita TA, Gamal SM, Shaker I, El Fishawy HS, El Sisi R, Shaker OG, et al. Clinical significance of serum interleukin-23 and A/G gene (rs17375018) polymorphism in Behçets disease: Relation to neuro-Behçet, uveitis and disease activity. *Joint Bone Spine*. 2015;**82**:213–215
- [122] Baerveldt EM, Kappen JH, Thio HB, van Laar JA, van Hagen PM, Prens EP. Successful long-term triple disease control by ustekinumab in a patient with Behcet's disease, psoriasis and hidradenitis suppurativa. *Annals of the Rheumatic Diseases*. 2013;**72**:626–627
- [123] Watford, WT, Hissong BD, Bream JH, Kanno Y, Muul L, O'Shea JJ. Signaling by IL-12 and IL-23 and the immunoregulatory roles of STAT4. *Immunological Reviews*. 2004;**202**:139–156
- [124] Trinchieri, G. Interleukin-12 and the regulation of innate resistance and adaptive immunity. *Nature Reviews Immunology*. 2003;**3**:133–146
- [125] Parham C, Chirica M, Timans J, Vaisberg E, Travis M, Cheung J, et al. A receptor for the heterodimeric cytokine IL-23 is composed of IL-12R β 1 and a novel cytokine receptor subunit, IL-23R. *Journal of Immunology*. 2002;**168**:5699–5708
- [126] Tulunay A, Dozmorov MG, Ture-Ozdemir F, Yilmaz V, Eksioglu-Demiralp E, Alibaz-Oner F, et al. Activation of the JAK/STAT pathway in Behcet's disease. *Genes & Immunity*. 2015;**16**:176
- [127] Chi W, Yang P, Zhu X, Wang Y, Chen L, Huang X, et al. Production of interleukin-17 in Behçet's disease is inhibited by cyclosporin A. *Molecular Vision*. 2010;**16**:880–886
- [128] Cai T, Wang Q, Zhou Q, Wang C, Hou S, Qi J, et al. Increased expression of IL-22 is associated with disease activity in Behcet's disease. *PLoS One*. 2013;**8**:e59009

- [129] Yasuoka H, Chen Z, Takeuchi T, Kuwana M. Th17 is involved in the pathogenesis of Behçet's disease via CCL20-CCR6 axis. *Arthritis Research & Therapy*. 2012;**14**(Suppl 1):P79
- [130] Cordero-Coma M, Calleja S, Llorente M, Rodriguez E, Franco M, Ruiz de Morales JG. Serum cytokine profile in adalimumab-treated refractory uveitis patients: Decreased IL-22 correlates with clinical responses. *Ocular Immunology and Inflammation*. 2013;**21**:212–219
- [131] Chi W, Zhou S, Yang P, Chen L. CD4+ T cells from Behcet patients produce high levels of IL-17. *Eye Science*. 2011;**26**:65–69
- [132] Shimizu J, Takai K, Fujiwara N, Arimitsu N, Ueda Y, Wakisaka S, et al. Excessive CD4+ T cells co-expressing interleukin-17 and interferon- γ in patients with Behçet's disease. *Clinical & Experimental Immunology*. 2012;**168**:68–74
- [133] Al-Zifzaf DS, Mokbel AN, Abdelaziz DM. Interleukin-17 in Behçet's disease: Relation with clinical picture and disease activity. *Egyptian Rheumatology and Rehabilitation*. 2015;**42**:34–38
- [134] Nurieva R, Yang XO, Martinez G, Zhang Y, Panopoulos AD, Ma L, et al. Essential autocrine regulation by IL-21 in the generation of inflammatory T cells. *Nature*. 2007;**448**:480–483
- [135] Geri G, Terrier B, Rosenzweig M, Wechsler B, Touzot M, Seilhean D, et al. Critical role of IL-21 in modulating T(H)17 and regulatory T cells in Behçet disease. *Journal of Allergy and Clinical Immunology*. 2011;**128**:655–664
- [136] Houman H, Hamzaoui A, Ben Ghorbal I, Khanfir M, Feki M, Hamzaoui K. Abnormal expression of chemokine receptors in Behçet's disease: Relationship to intracellular Th1/Th2 cytokines and to clinical manifestations. *Journal of Autoimmunity*. 2004;**23**:267–273
- [137] Koarada S, Haruta Y, Tada Y, Ushiyama O, Morito F, Ohta A, et al. Increased entry of CD4+ T cells into the Th1 cytokine effector pathway during T cell division following stimulation in Behcet's disease. *Rheumatology*. 2004;**43**:843–851
- [138] Raziuddin S, al-Dalaan A, Bahabri S, Siraj AK, al-Sedairy S. Divergent cytokine production profile in Behçet's disease. Altered Th1/Th2 cell cytokine pattern. *Journal of Rheumatology*. 1998;**25**:329–333
- [139] Aridogan BC, Yildirim M, Baysal V, Inaloz HS, Baz K, Kaya S. Serum levels of IL-4, IL-10, IL-12, IL-13 and IFN-gamma in Behçet's disease. *Journal of Dermatology*. 2003;**30**:602–607
- [140] Dalghous AM, Freysdottir J, Fortune F. Expression of cytokines, chemokines, and chemokine receptors in oral ulcers of patients with Behcet's disease (BD) and recurrent aphthous stomatitis is Th1-associated, although Th2-association is also observed in patients with BD. *Scandinavian Journal of Rheumatology*. 2006;**35**:472–475
- [141] Ben Ahmed M, Houman H, Miled M, Dellagi K, Louzir H. Involvement of chemokines and Th1 cytokines in the pathogenesis of mucocutaneous lesions of Behçet's disease. *Arthritis & Rheumatology*. 2004;**50**:2291–2295

- [142] Gündüz E, Teke HU, Bilge NS, Cansu DU, Bal C, Korkmaz C, et al. Regulatory T cells in Behçet's disease: Is there a correlation with disease activity? Does regulatory T cell type matter? *Rheumatology International*. 2013;**33**:3049–3054
- [143] Nanke Y, Kotake S, Goto M, Ujihara H, Matsubara M, Kamatni N. Decreased percentages of regulatory T cells in peripheral blood of patients with Behçet's disease before ocular attack: A possible predictive marker of ocular attack. *Modern Rheumatology*. 2008;**18**:354–358
- [144] Sugita S, Yamada Y, Kaneko S, Horie S, Mochizuki M. Induction of regulatory T cells by infliximab in Behçet's disease. *Investigative Ophthalmology & Visual Science*. 2011;**52**:476–484
- [145] Bandala-Sanchez E, Zhang Y, Reinwald S, Dromey JA, Lee BH, Qian J, et al. T cell regulation mediated by interaction of soluble CD52 with the inhibitory receptor Siglec-10. *Nature Immunology*. 2013;**14**:741–748
- [146] Mohammad AJ, Smith RM, Chow YW, Chaudhry AN, Jayne DR. Alemtuzumab as remission induction therapy in Behçet disease: A 20-year experience. *Journal of Rheumatology*. 2015;**42**:1906–1913
- [147] Perez-Pampin E, Campos-Franco J, Blanco J, Mera A. Remission induction in a case of refractory Behçet disease with alemtuzumab. *Journal of Clinical Rheumatology*. 2013;**19**:101–103
- [148] Carding SR, Egan PJ. Gammadelta T cells: Functional plasticity and heterogeneity. *Nature Reviews Immunology*. 2002;**2**:336–345
- [149] Hasan MS, Bergmeier LA, Petrushkin H, Fortune F. Gamma delta ($\gamma\delta$) T cells and their involvement in Behçet's disease. *Journal of Immunology Research*. 2015;**2015**:705831
- [150] Fortune F, Walker J, Lehner T. The expression of $\gamma\delta$ T cell receptor and the prevalence of primed, activated and IgA-bound T cells in Behçet's syndrome. *Clinical & Experimental Immunology*. 1990;**82**:326–332
- [151] Suzuki Y, Hoshi K, Matsuda T, Mizushima Y. Increased peripheral blood gamma delta+ T cells and natural killer cells in Behçet's disease. *Journal of Rheumatology*. 1992;**19**:588–592
- [152] Hamzaoui K, Hamzaoui A, Hentati F, Kahan A, Ayed K, Chabbou A, et al. Phenotype and functional profile of T cells expressing gamma delta receptor from patients with active Behçet's disease. *Journal of Rheumatology*. 1994;**21**:2301–2306
- [153] Hasan A, Fortune F, Wilson A, Warr K, Shinnick T, Mizushima Y, et al. Role of gamma delta T cells in pathogenesis and diagnosis of Behçet's disease. *Lancet*. 1996;**347**:789–794
- [154] Freysdottir J, Lau S, Fortune F. Gammadelta T cells in Behçet's disease (BD) and recurrent aphthous stomatitis (RAS). *Clinical & Experimental Immunology*. 1999;**118**:451–457
- [155] Bank I, Duvdevani M, Livneh A. Expansion of gammadelta T-cells in Behçet's disease: Role of disease activity and microbial flora in oral ulcers. *Journal of Laboratory and Clinical Medicine*. 2003;**141**:33–40

- [156] Verjans GM, van Hagen PM, van der Kooi A, Osterhaus AD, Baarsma GS. Vgamma9Vdelta2 T cells recovered from eyes of patients with Behçet's disease recognize non-peptide prenyl pyrophosphate antigens. *Journal of Neuroimmunology*. 2002;**130**:46–54
- [157] Triolo G, Accardo-Palumbo A, Dieli F, Ciccia F, Ferrante A, Giardina E, et al. Vgamma9/Vdelta2 T lymphocytes in Italian patients with Behçet's disease: Evidence for expansion, and tumour necrosis factor receptor II and interleukin-12 receptor beta1 expression in active disease. *Arthritis Research & Therapy*. 2003;**5**:R262–R268
- [158] Clemente A, Cambra A, Munoz-Saá I, Crespi C, Pallarés L, Juan A, et al. Phenotype markers and cytokine intracellular production by CD8+ gammadelta T lymphocytes do not support a regulatory T profile in Behçet's disease patients and healthy controls. *Immunology Letters*. 2010;**129**:57–63
- [159] Parlakgul G, Guney E, Erer B, Kılıcaslan Z, Direskeneli H, Gul A, et al. Expression of regulatory receptors on $\gamma\delta$ T cells and their cytokine production in Behçet's disease. *Arthritis Research & Therapy*. 2013;**15**:R15
- [160] Nold MF, Nold-Petry CA, Zepp JA, Palmer BE, Bufler P, Dinarello CA. IL-37 is a fundamental inhibitor of innate immunity. *Nature Immunology*. 2010;**11**:1014–1022
- [161] Bouali E, Kaabachi W, Hamzaoui A, Hamzaoui K. Interleukin-37 expression is decreased in Behçet's disease and is associated with inflammation. *Immunology Letters*. 2015;**167**:87–94
- [162] Ye Z, Wang C, Kijlstra A, Zhou X, Yang P. A possible role for interleukin 37 in the pathogenesis of Behçet's disease. *Current Molecular Medicine*. 2014;**14**:535–542
- [163] Hamzaoui K, Hamzaoui A. The anti-inflammatory activity of interleukin-37 in Behçet's disease. *Inflammation and Cell Signaling*. 2016;**3**:e1452
- [164] Wang C, Tian Y, Ye Z, Kijlstra A, Zhou Y, Yang P. Decreased interleukin 27 expression is associated with active uveitis in Behçet's disease. *Arthritis Research & Therapy*. 2014;**16**:R117
- [165] Curnow SJ, Pryce K, Modi N, Knight B, Graham EM, Stewart JE, et al. Serum cytokine profiles in Behçet's disease: Is there a role for IL-15 in pathogenesis? *Immunology Letters*. 2008;**121**:7–12
- [166] Hamzaoui K, Hamzaoui A, Ghorbel I, Khanfir M, Houman H. Levels of IL-15 in serum and cerebrospinal fluid of patients with Behçet's disease. *Scandinavian Journal of Immunology*. 2006;**64**:655–660
- [167] Luster AD. Chemokines—Chemotactic cytokines that mediate inflammation. *New England Journal of Medicine*. 1998;**338**:436–445
- [168] Saruhan-Direskeneli G, Yentür SP, Akman-Demir G, Işık N, Serdaroğlu P. Cytokines and chemokines in neuro-Behçet's disease compared to multiple sclerosis and other neurological diseases. *Journal of Neuroimmunology*. 2003;**145**:127–134
- [169] El-Asrar AM, Al-Obeidan SS, Kangave D, et al. CXC chemokine expression profiles in aqueous humor of patients with different clinical entities of endogenous uveitis. *Immunobiology*. 2011;**216**:1004–1009

- [170] Ambrose N, Khan E, Ravindran R, Lightstone L, Abraham S, Botto M, et al. The exaggerated inflammatory response in Behçet's syndrome: Identification of dysfunctional post-transcriptional regulation of the IFN- γ /CXCL10 IP-10 pathway. *Clinical & Experimental Immunology*. 2015;**181**:427–433
- [171] Durmazlar SP, Ulkar GB, Eskioglu F, Tatlican S, Mert A, Akgul A. Significance of serum IL-8 levels in patients with Behçet's disease: High levels may indicate vascular involvement. *International Journal of Dermatology*. 2009;**48**:259–264
- [172] Mantas C, Direskeneli H, Oz D, Yavuz S, Akoglu T. IL-8 producing cells in patients with Behçet's disease. *Clinical and Experimental Rheumatology*. 2000;**18**:249–251
- [173] Katsantonis J, Adler Y, Orfanos CE, Zouboulis CC. Adamantiades-Behçet's disease: Serum IL-8 is a more reliable marker for disease activity than C-reactive protein and erythrocyte sedimentation rate. *Dermatology*. 2000;**201**:37–39
- [174] Erdem H, Pay S, Serdar M, Simşek I, Dinç A, Muşabak U, et al. Different ELR (+) angiogenic CXC chemokine profiles in synovial fluid of patients with Behçet's disease, familial Mediterranean fever, rheumatoid arthritis, and osteoarthritis. *Rheumatology International*. 2005;**26**:162–167
- [175] Ozer HT, Erken E, Gunesacar R, Kara O. Serum RANTES, MIP-1 α , and MCP-1 levels in Behçet's disease. *Rheumatology International*. 2005;**25**:487–488
- [176] Kökçam I, Turgut D, İlhan NF, Çiçek D. The levels of serum chemokines in patients with Behçet's disease. *Turkish Journal of Medical Sciences*. 2012;**42**:1105–1110
- [177] Kim WU, Do JH, Park KS, Cho ML, Park SH, Cho CS, et al. Enhanced production of macrophage inhibitory protein-1 α in patients with Behçet's disease. *Scandinavian Journal of Rheumatology*. 2005;**34**:129–135.
- [178] Kaburaki T, Fujino Y, Kawashima H, Merino G, Numaga J, Chen J, et al. Plasma and whole-blood chemokine levels in patients with Behçet's disease. *Graefe's Archive for Clinical and Experimental Ophthalmology*. 2003;**241**:353–358
- [179] Do JH, Jung JH, Park CS, Ko JS, Kim SS, Choi HC, et al. Elevated monocyte chemoattractant protein-1 in patients with Behçet's disease. *Korean Journal of Medicine*. 2003;**65**:458–466
- [180] Miyagishi R, Kikuchi S, Fukazawa T, Tashiro K. Macrophage inflammatory protein-1 α in the cerebrospinal fluid of patients with multiple sclerosis and other inflammatory neurological diseases. *Journal of the Neurological Sciences*. 1995;**129**:223–227
- [181] Becatti M, Emmi G, Silvestri E, Bruschi G, Ciucciarelli L, Squatrito D, et al. Neutrophil activation promotes fibrinogen oxidation and thrombus formation in Behçet disease. *Circulation*. 2016;**133**:302–311
- [182] Köse K, Yazici C, Cambay N, Aşcıoğlu O, Doğan P. Lipid peroxidation and erythrocyte antioxidant enzymes in patients with Behçet's disease. *The Tohoku Journal of Experimental Medicine*. 2002;**197**:9–16

- [183] Yazici C, Köse K, Çalış M, Demİr M, Kirnap M, Ateş F. Increased advanced oxidation protein products in Behçet's disease: A new activity marker? *British Journal of Dermatology*. 2004;**151**:105–111
- [184] Yuksel M, Yildiz A, Oylumlu M, Turkcu FM, Bilik MZ, Ekinci A, et al. Novel markers of endothelial dysfunction and inflammation in Behçet's disease patients with ocular involvement: Epicardial fat thickness, carotid intima media thickness, serum ADMA level, and neutrophil-to-lymphocyte ratio. *Clinical Rheumatology*. 2016;**35**:701–708
- [185] Ozturk C, Balta S, Balta I, Demirkol S, Celik T, Turker T, et al. Neutrophil-lymphocyte ratio and carotid-intima media thickness in patients with Behçet disease without cardiovascular involvement. *Angiology*. 2015;**66**:291–296
- [186] Acikgoz N. The neutrophil-lymphocyte ratio and Behcet disease. *Angiology*. 2016;**67**:297
- [187] Rifaioglu EN, Bülbül Şen B, Ekiz Ö, Cigdem Dogramaci A. Neutrophil to lymphocyte ratio in Behçet's disease as a marker of disease activity. *Acta Dermatovenerologica Alpina, Pannonica et Adriatica*. 2014;**23**:65–67
- [188] Artis D, Spits H. The biology of innate lymphoid cells. *Nature*. 2015;**517**:293–301
- [189] Yamaguchi Y, Takahashi H, Satoh T, Okazaki Y, Mizuki N, Takahashi K, et al. Natural killer cells control a T-helper 1 response in patients with Behçet's disease. *Arthritis Research & Therapy*. 2010;**12**:R80
- [190] Takeno M, Shimoyama Y, Kashiwakura J, Nagafuchi H, Sakane T, Suzuki N. Abnormal killer inhibitory receptor expression on natural killer cells in patients with Behçet's disease. *Rheumatology International*. 2004;**24**:212–216
- [191] Diefenbach A, Colonna M, Koyasu S. Development, differentiation, and diversity of innate lymphoid cells. *Immunity*. 2014;**41**:354–365
- [192] Walker JA, Barlow JL, McKenzie AN. Innate lymphoid cells—How did we miss them? *Nature Reviews Immunology*. 2013;**13**:75–87
- [193] Schroder K, Tschopp, J. The inflammasomes. *Cell*. 2010;**140**:821–832
- [194] Strowig T, Henao-Mejia J, Elinav E, Flavell R. Inflammasomes in health and disease. *Nature*. 2012;**481**:278–286
- [195] Liang L, Tan X, Zhou Q, Zhu Y, Tian Y, Yu H, et al. IL-1 β triggered by peptidoglycan and lipopolysaccharide through TLR2/4 and ROS-NLRP3 inflammasome-dependent pathways is involved in ocular Behçet's disease. *Investigative Ophthalmology & Visual Science*. 2013;**54**:402–414
- [196] Kim EH, Park MJ, Park S, Lee ES. Increased expression of the NLRP3 inflammasome components in patients with Behçet's disease. *Journal of Inflammation*. 2015;**12**:41
- [197] Türe-Özdemir F, Tulunay A, Elbasi MO, Tatli I, Maurer AM, Mumcu G, et al. Pro-inflammatory cytokine and caspase-1 responses to pattern recognition receptor activation of neutrophils and dendritic cells in Behcet's disease. *Rheumatology*. 2013;**52**:800–805

- [198] Krause I, Monselise Y, Milo G, Weinberger A. Anti-Saccharomyces cerevisiae antibodies—A novel serologic marker for Behçet's disease. *Clinical and Experimental Rheumatology*. 2002;**20**(suppl 26):S21–S24
- [199] Fresko I, Ugurlu S, Ozbakir F, Celik A, Yurdakul S, Hamuryudan V, et al. Anti-Saccharomyces cerevisiae antibodies (ASCA) in Behçet's syndrome. *Clinical and Experimental Rheumatology*. 2005;**23**(4 Suppl 38):S67–S70
- [200] Choi CH, Kim TI, Kim BC, Shin SJ, Lee SK, Kim WH, et al. Anti-Saccharomyces cerevisiae antibody in intestinal Behçet's disease patients: Relation to clinical course. *Diseases of the Colon & Rectum*. 2006;**49**:1849–1859
- [201] Monselise A, Weinberger A, Monselise Y, Fraser A, Sulkes J, Krause I. Anti-Saccharomyces cerevisiae antibodies in Behçet's disease—A familial study. *Clinical and Experimental Rheumatology*. 2006;**24**(5 Suppl 42):S87–S90
- [202] Aydıntug AO, Tokgöz G, D'Cruz DP, Gürler A, Cervera R, Düzgün N, et al. Antibodies to endothelial cells in patients with Behçet's disease. *Clinical Immunology and Immunopathology*. 1993;**67**:157–162
- [203] Lee KH, Chung HS, Kim HS, Oh SH, Ha MK, Baik JH, et al. Human alpha-enolase from endothelial cells as a target antigen of anti-endothelial cell antibody in Behçet's disease. *Arthritis & Rheumatology*. 2003;**48**:2025–2035
- [204] Lee JH, Cho SB, Bang D, Oh SH, Ahn KJ, Kim J, et al. Human anti-alpha-enolase antibody in sera from patients with Behçet's disease and rheumatologic disorders. *Clinical and Experimental Rheumatology*. 2009;**27**(2 Suppl 53):S63–S66
- [205] Shin SJ, Kim BC, Kim TI, Lee SK, Lee KH, Kim WH. Anti-alpha-enolase antibody as a serologic marker and its correlation with disease severity in intestinal Behçet's disease. *Digestive Diseases and Sciences*. 2011;**56**:812–818
- [206] Vaiopoulos G, Lakatos PL, Papp M, Kaklamanis F, Economou E, Zevgolits V, et al. Serum anti-Saccharomyces cerevisiae antibodies in Greek patients with Behçet's disease. *Yonsei Medical Journal*. 2011;**52**:347–350
- [207] Filik L, Biyikoglu I. Differentiation of Behçet's disease from inflammatory bowel diseases: Anti-Saccharomyces cerevisiae antibody and anti-neutrophilic cytoplasmic antibody. *World Journal of Gastroenterology*. 2008;**14**:7271
- [208] Rhee SH, Kim YB, Lee ES. Comparison of Behçet's disease and recurrent aphthous ulcer according to characteristics of gastrointestinal symptoms. *Journal of Korean Medical Science*. 2005;**20**:971–976
- [209] Isogai E, Ohno S, Kotake S, Isogai H, Tsurumizu T, Fujii N, et al. Chemiluminescence of neutrophils from patients with Behçet's disease and its correlation with an increased proportion of uncommon serotypes of Streptococcus sanguis in the oral flora. *Archives of Oral Biology*. 1990;**35**:43–8

- [210] Yokota K, Hayashi S, Fujii N, et al. Antibody response to oral streptococci in Behçet's disease. *Microbiology and Immunology*. 1992;**36**:815–822
- [211] Mumcu G, Inanc N, Aydin SZ, Ergun T, Direskeneli H. Association of salivary *S. mutans* colonisation and mannose-binding lectin deficiency with gender in Behçet's disease. *Clinical and Experimental Rheumatology*. 2009;**27**:S32–S36
- [212] Hirohata S, Oka H, Mizushima Y. Streptococcal-related antigens stimulate production of IL6 and interferon-gamma by T cells from patients with Behcet's disease. *Cellular Immunology*. 1992;**140**:410–419
- [213] Studd M, McCance DJ, Lehner T. Detection of HSV-1 DNA in patients with Behçet's syndrome and in patients with recurrent oral ulcers by the polymerase chain reaction. *Medical Microbiology*. 1991;**34**:39–43
- [214] Kim DY, Cho S, Choi MJ, Sohn S, Lee E-S, Bang D. Immunopathogenic role of herpes simplex virus in Behçet's disease. *Genetics Research International*. 2013;**2013**:638273
- [215] Hatemi G, Bahar H, Uysal S, Mat C, Gogus F, Masatlioglu S, et al. The pustular skin lesions in Behcet's syndrome are not sterile. *Annals of the Rheumatic Diseases*. 2004;**63**:1450–1452
- [216] Zouboulis CC, Turnbull JR, Mühlradt PF. Association of *Mycoplasma fermentans* with Adamantiades-Behçet's disease. *Advances in Experimental Medicine and Biology*. 2003;**528**:191–194
- [217] Lederberg J, McCray AT. 'Ome Sweet' omics—A genealogical treasury of words. *Scientist*. 2001;**15**:8
- [218] Cerf-Bensussan N, Gaboriau-Routhiau V. The immune system and the gut microbiota: Friends or foes? *Nature Reviews Immunology*. 2010;**10**:735–744
- [219] Cua DJ, Sherlock JP. Autoimmunity's collateral damage: Gut microbiota strikes 'back'. *Nature Medicine*. 2011;**17**:1055–1056
- [220] Tringe SG, Rubin EM. Metagenomics: DNA sequencing of environmental samples. *Nature Reviews Genetics*. 2005;**6**:805–814
- [221] Tringe SG, Hugenholtz P. A renaissance for the pioneering 16S rRNA gene. *Current Opinion in Microbiology*. 2008;**11**:442–446
- [222] Consolandi C, Turrone S, Emmi G, Severgnini M, Fiori J, Peano C, et al. Behçet's syndrome patients exhibit specific microbiome signature. *Autoimmunity Reviews*. 2015;**14**:269–276
- [223] Shimizu J, Kubota T, Takada E, Takai K, Fujiwara N, Arimitsu N, et al. Bifidobacteria abundance-featured gut microbiota compositional change in patients with Behcet's disease. *PLoS ONE*. 2016;**11**:e0153746

- [224] Seoudi N, Bergmeier LA, Drobniowski F, Paster B, Fortune F. The oral mucosal and salivary microbial community of Behçet's syndrome and recurrent aphthous stomatitis. *Journal of Oral Microbiology*. 2015;**7**:27150
- [225] Coit P, Mumcu G, Ture-Ozdemir F, Unal AU, Alpar U, Bostanci N, et al. Sequencing of 16S rRNA reveals a distinct salivary microbiome signature in Behçet's disease. *Clinical Immunology*. 2016;**169**:28–35
- [226] Özşeker B, Şahin C, Özşeker HS, Efe SC, Kav T, Bayraktar Y. The role of fecal calprotectin in evaluating intestinal involvement of Behçet's disease. *Disease Markers*. 2016;**2016**:5423043
- [227] Hatemi I, Hatemi G, Çelik AF. Systemic vasculitis and the gut. *Current Opinion in Rheumatology*. 2017;**29**:33–38
- [228] Kim DH, Park Y, Kim B, Kim SW, Park SJ, Hong SP, et al. Fecal calprotectin as a non-invasive biomarker for intestinal involvement of Behçet's disease. *J Gastroenterol Hepatol* 2017; **32**: 595–601
- [229] Bae JH, Lee SC. Effect of intravitreal methotrexate and aqueous humor cytokine levels in refractory retinal vasculitis in Behcet disease. *Retina*. 2012;**32**:1395–1402
- [230] Ozdamar Y, Berker N, Bahar G, Soykan E, Bicer T, Ozkan SS, et al. Inflammatory mediators and posterior segment involvement in ocular Behcet disease. *European Journal of Ophthalmology*. 2009;**19**:998–1003
- [231] Bozoglu E, Dinc A, Erdem H, Pay S, Simsek I, Kocar IH. Vascular endothelial growth factor and monocyte chemoattractant protein-1 in Behçet's patients with venous thrombosis. *Clinical and Experimental Rheumatology*. 2005;**23**(4 Suppl 38):S42–S48
- [232] Paroli MP, Teodori C, D'Alessandro M, Mariani P, Iannucci G, Paroli M. Increased vascular endothelial growth factor levels in aqueous humor and serum of patients with quiescent uveitis. *European Journal of Ophthalmology*. 2007;**17**:938–942
- [233] Kose O, Stewart J, Waseem A, Lalli A, Fortune F. Expression of cytokeratins, adhesion and activation molecules in oral ulcers of Behçet's disease. *Clinical and Experimental Dermatology*. 2008;**33**:62–69
- [234] Saglam K, Yilmaz MI, Saglam A, Ulgey M, Bulucu F, Baykal Y. Levels of circulating intercellular adhesion molecule-1 in patients with Behçet's disease. *Rheumatology International*. 2002;**21**:146–148
- [235] Haznedaroglu E, Karaaslan Y, Büyükaşık Y, Koşar A, Ozcebe O, Haznedaroglu BC, et al. Selectin adhesion molecules in Behçet's disease. *Annals of the Rheumatic Diseases*. 2000;**59**:61–63
- [236] Sari RA, Kiziltunç A, Taysi S, Akdemir S, Gündoğdu M. Levels of soluble E-selectin in patients with active Behcet's disease. *Clinical Rheumatology*. 2005;**24**:55–59

- [237] Seo J, Ahn Y, Zheng Z, Kim BO, Choi MJ, Bang D, et al. Clinical significance of serum YKL-40 in Behçet disease. *British Journal of Dermatology*. 2016;**174**:1337–1344
- [238] Bilen H, Altinkaynak K, Sebin E, Aksoy H, Akcay F. Serum YKL-40 and MDA levels in Behcet disease. *Journal of the Pakistan Medical Association*. 2016;**66**:1299–1302
- [239] Ar MC, Hatemi G, Ekizoğlu S, Bilgen H, Saçlı S, Buyru AN, et al. JAK2 (V617F) mutation is not associated with thrombosis in Behcet syndrome. *Clinical and Applied Thrombosis/Hemostasis*. 2012;**18**:421–426
- [240] Adeeb F, Tayel M, El Kaffash DM, Idris KM, Hassan MF, Fraser AD. Janus kinase 2 V617F mutation and thrombotic events in Behçet's disease: The Alexandria experience. *European Journal of Rheumatology*. 2016;**3**:73–74
- [241] Silingardi M, Salvarani C, Boiardi L, Accardo P, Iorio A, Olivieri I, et al. Factor V Leiden and prothrombin gene G20210A mutations in Italian patients with Behçet's disease and deep vein thrombosis. *Arthritis & Rheumatology*. 2004;**51**:177–183
- [242] Leiba M, Seligsohn U, Sidi Y, Harats D, Sela BA, Griffin JH, et al. Thrombophilic factors are not the leading cause of thrombosis in Behçet's disease. *Annals of the Rheumatic Diseases*. 2004;**63**:1445–1449
- [243] Gurgey A, Balta G, Boyvat A. Factor V Leiden mutation and PAI-1 gene 4G/5G genotype in thrombotic patients with Behcet's disease. *Blood Coagulation & Fibrinolysis*. 2003;**14**:121–124
- [244] Toydemir PB, Elhan AH, Tükün A, Toydemir R, Gürler A, Tüzüner A, et al. Effects of factor V gene G1691A, methylenetetrahydrofolate reductase gene C677T, and prothrombin gene G20210A mutations on deep venous thrombogenesis in Behçet's disease. *Journal of Rheumatology*. 2000;**27**:2849–2854
- [245] Seyahi E, Yurdakul S. Behçet's syndrome and thrombosis. *Mediterranean Journal of Hematology and Infectious Diseases*. 2011;**3**:e2011026
- [246] Guillot X, Semerano L, Saidenberg-Kermanac'h N, Falgarone G, Boissier MC. Vitamin D and inflammation. *Joint Bone Spine*. 2010;**77**:552–557
- [247] Hamzaoui K, Ben Dhifallah I, Karray E, Sassi FH, Hamzaoui A. Vitamin D modulates peripheral immunity in patients with Behçet's disease. *Clinical and Experimental Rheumatology*. 2010;**28**(4 Suppl 60):S50–S57
- [248] Bscheider M, Butcher EC. Vitamin D immunoregulation through dendritic cells. *Immunology*. 2016;**148**:227–236
- [249] Szymczak I, Pawliczak R. The active metabolite of vitamin D3 as a potential immunomodulator. *Scandinavian Journal of Immunology*. 2016;**83**:83–91
- [250] Gatenby P, Lucas R, Swaminathan A. Vitamin D deficiency and risk for rheumatic diseases: An update. *Current Opinion in Rheumatology*. 2013;**25**:184–191

- [251] Karatay S, Yildirim K, Karakuzu A, Kiziltunc A, Engin RI, Eren YB, et al. Vitamin D status in patients with Behcet's disease. *Clinics (Sao Paulo)* 2011;**66**:721–723
- [252] Can M, Gunes M, Haliloglu OA, Haklar G, Inanç N, Yavuz DG, Direskeneli H. Effect of vitamin D deficiency and replacement on endothelial functions in Behçet's disease. *Clinical and Experimental Rheumatology*. 2012;**30**(3 Suppl 72):S57–S61
- [253] Khabbazi A, Rashtchizadeh N, Ghorbanihaghjo A, Hajialiloo M, Ghojazadeh M, Taei R, et al. The status of serum vitamin D in patients with active Behcet's disease compared with controls. *International Journal of Rheumatic Diseases*. 2014;**17**:430–434
- [254] Do JE, Kwon SY, Park S, Lee ES. Effects of vitamin D expression of Toll-like receptor of monocytes from patients with Behcet's disease. *Rheumatology (Oxford)*. 2008;**47**:840–848
- [255] Kechida M, Harzallah O, Hellara I, Klii R, Na eti F, Najjar MF, et al. Vitamin D status of Behcet's patients. Analysis of correlation with activity and severity of the disease as well as with the quality of life of patients. *Int Arch Med* 2015; **8**: 1–7
- [256] Gül A. Pathogenesis of Behçet's disease: Autoinflammatory features and beyond. *Seminars in Immunopathology*. 2015;**37**:413–418