We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Transcriptome Analysis in Chickpea (*Cicer arietinum* L.): Applications in Study of Gene Expression, Non-Coding RNA Prediction, and Molecular Marker Development

Chandra Kant, Vimal Pandey, Subodh Verma,
Manish Tiwari, Santosh Kumar and Sabhyata Bhatia

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.69884>

Abstract

Extensive analyses of transcriptome have been carried out in chickpea, which is the third most important legume valued as a source of dietary protein and micronutrients. Over the last two decades, several laboratories have used a wide range of techniques encompassing expressed sequence tag (EST) analysis, serial analysis of gene expression (SAGE), microarray and next-generation sequencing (NGS) technologies for analysing the chickpea transcriptomes. However, chickpea transcriptome analysis witnessed significant progress with the advent of the NGS platforms. Gene expression analyses using NGS platforms were carried out in the vegetative and reproductive tissues such as shoot, root, mature leaf, flower bud, young pod, seed and nodule by various groups which resulted in identification of several tissue-specific transcripts. Some laboratories have utilized transcriptomics to explore the response of chickpea to abiotic and biotic stresses such as drought, salinity, heat, cold, *Fusarium oxysporum* and *Ascochyta rabiei* differentially expressed genes and also established crosstalk between biotic and abiotic stress responses. Transcriptome analysis has been utilized extensively to identify non-coding RNAs such as miRNAs and long intergenic non-coding (LINC) RNAs. Transcriptome analysis has facilitated the development of molecular markers such as simple sequence repeats (SSRs), single-nucleotide polymorphisms (SNPs) and potential intron polymorphisms (PIPs) that are being used to expedite the chickpea breeding programmes. The available chickpea transcriptomes will continue to serve as the foundation for devising strategies for chickpea improvement.

Keywords: transcriptome, chickpea, next-generation sequencing (NGS), gene expression, molecular markers

1. Introduction

Chickpea (*Cicer arietinum* L.) is a diploid ($2n = 16$), self-pollinated plant which is grown in the cool season and has a genome size of 738 Mb [1]. It is the third most produced pulse crop in the world (13.73 million tons) after beans (26.52 million tons) and green pea (17.43 million tons) (FAOSTAT 2014). It is considered to be an ideal crop for the semiarid and arid regions as it exhibits an extensive tap root system. Chickpea seeds are an excellent source of nutrition as they contain $\approx 40\%$ carbohydrates, $\approx 6\%$ oil and 20–30% protein and good source of minerals and trace elements such as calcium, magnesium, phosphorus, iron and zinc [2]. Moreover, chickpea contributes to improvement of soil fertility since it has the capability to establish symbiotic association with *Mesorhizobium ciceri* that helps in fixing atmospheric nitrogen to the reduced nitrogen (NH_3). Chickpea, through symbiotic nitrogen fixation (SNF), can fulfil up to 80% of its nitrogen requirement [3]. All these qualities make chickpea an economically important crop as it is an affordable source that can fulfil the dietary protein requirement of the masses.

2. Challenges in chickpea production

The world average of chickpea productivity is 982.1 kg/ha (FAOSTAT 2014); however, a simulated study showed that potential productivity of chickpea in rain-fed situations ranged from 1390 to 4590 kg/ha [4]. There is a huge yield gap of 408–3608 kg/ha. A number of biotic and abiotic factors affect chickpea plant growth and, therefore, are responsible for poor productivity.

Chickpea is mostly raised on conserved soil moisture under rain-fed conditions [5]. Therefore, drought stress generally affects the crop at terminal stage [6] and leads to productivity loss of up to 50% [7]. Drought reduces overall biomass, reproductive growth and seed yield and increases flower abortion, pod abscission and number of empty pods [8]. Soil salinity affects productivity by delaying the flowering leading to decrease in reproductive success of chickpea [9]. Since chickpea is a cool season crop, high temperatures adversely affect the development of the plant [10]. Chander [11] reported a decline in yield of chickpea by about 301 kg/ha per 1°C increase in mean seasonal temperature in India [12, 13]. Biotic factors also adversely affect the yield of chickpea crop. *Fusarium* wilt, caused by *Fusarium oxysporum* f.sp. *ciceri*; *Ascochyta* blight, caused by *Ascochyta rabiei* and *Botrytis* grey mould, caused by *Botrytis cinerea* mainly affect the leaves of chickpea, whereas *Pythium ultimum* causes root and seed rot and is common in the areas where the chickpea growing season is cool and humid [14, 15]. A number of other fungi, such as *Alternaria* sp., *Ascochyta pisi*, *Uromyces* sp., *Botrytis* sp., *Phytophthora medicaginis* and so on, cause considerable damage to chickpea crops. Pod borer (*Helicoverpa armigera* Hubner) is the major pest affecting chickpea worldwide [15–17]. Therefore, improvement in yield, nutritional quality and stress tolerance are the major targets of chickpea research and breeding programmes which may be facilitated by detailed understanding of biological processes occurring in tissue-specific and developmental pathways. Moreover, responses to various stresses at molecular level also need to be elucidated in detail.

3. Legume genomics

With the advent of next-generation sequencing technologies, there has been a rapid increase in the efficiency of DNA and RNA sequencing and decrease in the cost involved. *Leguminosae* is a very important family known due to the economic and nutritional value of its members [18]. The recent years have witnessed a spurt in the number of studies utilizing genomic approaches to understand the biology of several agronomic traits in legumes.

The advances in DNA sequencing have led to whole genome sequencing of important legumes such as *Glycine max* [19], *Medicago truncatula* [20], *Lotus japonicus* [21], *Cajanus cajan* [22], *Phaseolus vulgaris* [23], diploid ancestors of peanut *Arachis duranensis* and *Arachis ipaensis* [24] and *C. arietinum* [1, 25]. Moreover, whole genome resequencing has been carried out for soybean [26, 27], *Medicago* [28] and chickpea [1] in order to understand the genetic variability, evolution and domestication in greater depth. Simultaneously, in order to unravel the functional aspects of legume biology, several NGS-based studies of transcriptomes were carried out. These studies have made significant contributions towards understanding of gene expression, alternative splicing events and small RNA identification. Gene expression atlases have been developed for soybean [29, 30], *Medicago* [31], *L. japonicus* [32] and pigeon pea [33]. Moreover, in chickpea a number of transcriptome studies have been performed. These include exploring the overall transcriptome of various tissues [34–37], specifically understanding of the development of flower [38], seed [39] and root nodule [40]. Transcriptome analysis of chickpea under different abiotic and biotic stresses such as drought, desiccation, salinity, cold and *Fusarium* wilt has also been carried out [41–43].

Next-generation sequencing (NGS)-based plant genomics has also assisted in understanding of genetic variation within and between species mostly through identification of single-nucleotide polymorphisms (SNPs). In chickpea, a number of studies have been performed to identify SNPs and utilized for various applications such as construction of linkage maps, synteny analysis, anchoring of whole genome sequencing and quantitative trait loci (QTL) analysis [44–49]. A CicArVarDB has also been developed which includes SNP and InDel variations in chickpea [50].

4. Transcriptome

A cell undergoing a functional or developmental process has a specific set of genes undergoing transcription at a particular time and is collectively called the 'transcriptome'. Thus, a transcriptome represents up to an extant physical, biochemical and developmental status of a cell. A transcriptome represents a pool of protein coding as well as nonprotein-coding RNAs; moreover, there may be the presence of variants of genes originating from alternative splicing and RNA editing, making the transcriptome more complex than a genome. Study of transcriptome may reveal information regarding spatial and temporal expression patterns of genes, and therefore it is possible to generate global expression profiles of genes representing developmental stages of an organism [34].

5. Methods for transcriptome analysis

Transcriptome analysis was initiated with the generation of expressed sequence tags (ESTs) that are 200–800 nucleotide long cDNA sequences, synthesised from mRNA through reverse transcription. ESTs represent the expressed part of an organism's genome and hence are an excellent resource for the study of gene expression at a genome-wide level. Conventionally, EST resources have been developed through Sanger sequencing. Although this process is used to generate and sequence longer fragments of cDNA, it is tedious and labour intensive and offers poor coverage of the transcriptome. These limitations of EST-based transcriptome analysis inspired scientists to develop microarray and other tag-based methods for gene expression analysis. Therefore, tools such as microarrays and serial analysis of gene expression (SAGE) continued to be used for several years for analysis of global gene expression patterns. However, with the advent of NGS and the simultaneous development of *in silico* analytical tools, global genome and transcriptome analysis has become a standard practice for deriving information to relate genotype to phenotype. However, it is not possible to sequence the transcripts to the full length due to technological limitations. Transcriptome analysis is based on the principle that the depth of coverage of a sequence is proportional to the level of expression of the corresponding gene. Therefore, by mapping and counting the sequenced reads onto the given transcript, expression can be measured, thereby translating sequence information to some biologically significant information. A host of NGS technologies such as sequencing by synthesis (Illumina Inc., USA), SOLiD (ThermoFisher Scientific) and pyrosequencing (454 biosciences/Roche) has provided unprecedented opportunities for high-throughput functional genomic research [51–53]. Moreover, a number of technologies for transcriptome sequencing are emerging such as The Ion Torrent (ThermoFisher Scientific), single-molecule real time (SMRT) (Pacific Biosciences, USA) and Nanopore (Oxford Nanopore Technologies, UK).

6. Using transcriptome analysis for studying biological processes in chickpea

Extensive transcriptome analysis has been carried out in chickpea in order to gain insights into the numerous biological processes. Techniques, such as EST sequencing, SAGE and most importantly the NGS, have been used to analyse the transcriptomes of root, shoot, flower, seed and nodule tissues in order to understand the tissue-specific development and function. Several groups undertook EST sequencing, and till date (March 2017) 53,333 chickpea ESTs are reported in the NCBI database. In another earlier study of the root transcriptome, an EST library was constructed by subtractive suppressive hybridization (SSH) of two related chickpea varieties, ICC 4958 and Annigeri, as they show different root traits. Sequences of more than 2800 ESTs were reported and used to develop the 'Chickpea Root Expressed Sequence Tag Database' [54].

A major advancement in transcriptome analysis for understanding developmental and biological processes occurred with the advent of the NGS platform. Several large-scale NGS-based transcriptome analyses were carried out in chickpea [34–36]. In one of the first NGS-based

studies, the Illumina sequencing of transcriptome of chickpea genotype ICC 4958 root and shoot followed by de novo assembly resulted in generation of 53,409 transcripts. Of these 34,676 transcripts were annotated, and 6577 transcripts were identified as transcription factors (TFs) belonging to 57 families. Another study by Garg et al. reported the Roche/454-based transcriptomes of 'shoot', 'root', 'mature leaf', 'flower bud' and 'young pod' of chickpea genotype ICC 4958 [34]. These sequence reads generated by the Roche/454 platform were merged with the Illumina reads from the previous study, and a hybrid assembly was generated [34], which resulted in 34,760 tentative consensus (TC) transcripts. Of these, 1851 transcripts were annotated as transcription factors belonging to 84 families. This analysis also led to the identification of 1132, 695, 513, 408 and 126 TCs specifically expressed in flower bud, young pod, shoot, root and mature leaf, respectively. The complete data were integrated leading to the development of the 'Chickpea Transcriptome Data Base' (CTDB) which provides a searchable interface to the chickpea transcriptome data [34]. Further, transcriptome analysis of the wild progenitor of chickpea, i.e. *Cicer reticulatum* PI489777, was also performed by Jhanwar et al. [37]. Moreover, transcriptomes of the kabuli, desi and wild chickpeas were compared [55] and used to create an improved version of the Chickpea Transcriptome Data Base V2.0 [56].

Flower development is an important and specialized process that takes place in angiosperms. Hence, in order to gain insights into the molecular mechanisms responsible for flower development in chickpea, transcriptome analysis was carried out using the Illumina sequencing platform [38]. Transcriptome sequencing of eight successive developing stages of flower (flower buds at sizes 4, 6, 8 and 8–10 mm and flowers with closed petals, partially opened petals, opened and faded petals and senescing petals) along with young leaf, germinating seedling and shoot apical meristem was carried out. Differential expression analysis revealed 1572 genes to be differentially expressed in at least one stage of flower development. A number of 1118 genes (908 upregulated and 201 downregulated) and 966 genes (857 upregulated and 109 downregulated) were found to be differentially regulated in flower bud and flower developmental stages, respectively [38]. The majority of the differentially expressed genes were found to be involved in various flower developmental pathways such as floral organ identity; development of corolla, androecium and gynoecium and gametophyte development. Moreover, genes related to cell wall development and transport were also found to be differentially expressed. In addition, 111 TF genes were found differentially expressed in floral bud and flower.

Chickpea is most valued for its seeds since they serve as a source of protein, especially for vegetarian population. Therefore, a thorough understanding of the transcriptional flux during seed development is important in order to get insights into the biological processes that define the seed. Towards this, an NGS-based deep transcriptome analysis of chickpea seed at four developmental stages, i.e. 10 days after anthesis (DAA), 20 DAA, 30 DAA and 40 DAA, was carried out [39]. The transcriptome was sequenced using the 454 pyrosequencing on the GS-FLX Titanium platform followed by its assembly into 51,099 transcripts. A gene ontology enrichment of seed-specific genes revealed genes related to reproductive structure development, fruit development and embryonic and post-embryonic development to be highly represented. Many metabolic pathways such as proteolysis, lipid metabolic process, regulation of RNA metabolic

process, regulation of transcription, terpenoid metabolic process and gibberellin metabolic processes were also found to be significantly represented [39]. In another study, sequencing of ESTs from the chickpea embryo resulted in identification of 1480 unigenes expressed during embryo development [57]. The analysis also identified 12 genes encoding for F-box proteins, of which 2 F-box genes (*CarFbox_PP2* and *CarF-box_LysM*) were predicted to be involved in seed development [57].

Another important distinctive feature of chickpea is its ability to form symbiotic relationship with *M. ciceri* which results in the formation of specialized structures called root nodules. These are formed by the host plant and protect the oxygen-sensitive, bacterial nitrogen fixing machinery. It is a complex phenomenon and a detailed understanding of the molecular pathways governing that the process of nodule development and nitrogen fixation would certainly help plant scientists in developing sustainable farming strategies for chickpea. Towards this, a DeepSuperSAGE-based transcriptome analysis led to the identification of 71 genes being differentially expressed in root nodules [58]. Further, in order to understand the root nodulation in greater depth, a deep transcriptome analysis of the chickpea root nodule at different developing stages was carried out using the 454 pyrosequencing [40]. Sequencing of transcriptomes of uninfected root and three developing stages of nodules followed by reference-based assembly resulted in 83,405 transcripts. Of these 3760 were found to be differentially expressed in at least one of the stages of nodule when compared to uninfected root. Also, 1606 transcripts were identified as transcription factors, of which 171 TFs were found to be differentially expressed during nodulation.

7. Using transcriptome analysis for study of stress response in chickpea

Transcriptome analysis has been utilized exclusively to study different abiotic and biotic stress responses in chickpea. Drought and salinity are the major factors that limit the growth and productivity of the plants. Terminal drought is thought to be a major constraint affecting productivity of chickpea as it can lower the yield of chickpea by about 50% [59]. Cold stress also affects susceptible chickpea mainly at the reproductive stage where it leads to pollen sterility and flower abortion [60]. Thus, it is important to study the response of chickpea under these stress conditions in order to devise strategies for development of stress-tolerant chickpea. Earlier studies based on EST sequencing, SAGE and microarray provided preliminary evidence for drought responses of chickpea at transcriptome level [61–66]. An EST sequencing-based study of drought and salinity stress in chickpea resulted in generation of 20,162 ESTs, of which 105 were found to have differential expression during one of the stresses [65]. In another comparison between ESTs generated from chickpea, ICC 4958 (drought tolerant) and ICC 1882 (drought resistant) varieties resulted in identification of 5494 drought-responsive ESTs [61]. A microarray-based transcriptome analysis of root and leaf of chickpea under drought stress resulted in identification of 4815 differentially expressed genes. Approximately 2623 and 3969 genes were found to be differentially expressed, whereas 88 and 52 genes were found to be specifically expressed during drought stress in root and leaf tissues, respectively [66]. Another microarray analysis in chickpea revealed 109, 210 and 386

genes to be differentially expressed in drought, cold and high-salinity stresses, respectively [62]. A SuperSAGE-based transcriptome analysis of chickpea drought stressed and control tissues gave rise to 17,493 unique transcripts (UniTags) of which 7532 were differentially expressed in drought stress [63]. Another SuperSAGE followed by 454 sequencing of root nodule transcriptome of salt-tolerant variety INRAT-93 identified 363 and 106 genes to be upregulated and downregulated, respectively, in root and nodule tissues [64].

The more global view of stress response in chickpea was provided by the study of Garg et al. [41] in which the transcriptome of chickpea root and shoot under desiccation, salinity and cold stress was analysed. The Illumina sequencing-based transcriptome and comparison revealed 11,640 transcripts to be differentially expressed during at least one of the stresses. Seven hundred forty-five transcription factors (TFs) were also found to be differentially regulated in at least one stress condition. Moreover 3536 unannotated genes from the chickpea transcriptome were also identified [41]. A more detailed transcriptome analysis of drought-tolerant (ICC 4958), drought-sensitive (ICC 1882), salinity-tolerant (JG 62) and salinity-sensitive (ICCV2) chickpea varieties resulted in identification of 18,462 transcripts representing 13,964 unique loci in at least one sample/stress condition. The study also revealed 4954 and 5545 genes exclusively regulated in drought-tolerant and salinity-tolerant varieties. A number of 775 TFs encoding genes belonging to 80 families were also found differentially regulated in stress conditions. Members of the bHLH, WRKY, NAC, AP2-EREBP and MYB were found among the top differentially expressed TFs in stress condition [42]. In order to understand the effect of cold stress, AFLP-based transcript profiling (cDNA-AFLP) approach was used [67], which showed that in cold-tolerant chickpea, 102 transcript-derived fragments (TDFs) were differentially expressed during cold stress. Moreover, transcriptome analysis of cold-tolerant chickpea ICC 16349 using cDNA differential display (DDRT-PCR) resulted in identification of 127 ESTs as differentially expressed in anthers during cold stress conditions.

Ascochyta blight caused by *A. rabiei* and *Fusarium* wilt caused by *F. oxysporum* are major fungal diseases of chickpea. In order to understand the response of chickpea to *A. rabiei*, an EST library sequencing of blight-resistant chickpea variety ICC 3996 infected with *A. rabiei* was performed by Coram and Pang [68]. The study reported 516 genes of which 4% were related to defense and found to encode for lignin and phytoalexin biosynthesis enzymes, pathogenesis-related proteins, signalling proteins and putative-defensive proteins [68]. For further identification of resistance-related genes, transcriptome analysis of four genotypes, *C. arietinum* ICC 3996, *C. arietinum* Lasseter, *C. arietinum* FLIP94-508C and *Cicer echinospermum* ILWC245, was performed using 756 featured microarrays. The study revealed 97 genes to be differentially expressed upon infection with *A. rabiei*. A comparison between resistant and susceptible varieties identified many genes such as pathogenesis-related proteins, SNAKIN2 antimicrobial peptide, proline-rich protein, disease resistance response protein DRRG49-C, environmental stress-inducible protein, leucine-zipper protein, polymorphic antigen membrane protein and Ca-binding protein, which might be responsible for imparting resistance to the tolerant varieties [69]. On the other hand, in order to identify genes involved in wilt resistance in chickpea, EST sequencing followed by microarray analysis of chickpea wilt susceptible genotype (JG-62) and resistant genotype (WR-315) was performed after infecting them with *F. oxysporum*

ciceri. The analysis resulted in identification of 257 differentially expressed genes associated with the early signalling pathway [70]. In order to understand the differential response of susceptible and tolerant/resistant chickpea varieties to *F. oxysporum*, transcriptomes of wilt susceptible (JG62) and wilt-tolerant/wilt-resistant (ICCV2, K850 and WR315) chickpea varieties were analysed using the Illumina platform. Comparison among the transcriptomes led to identification of 303 polymorphic SSRs, 14,462 SNPs and 1864 insertions/deletions (InDels). Moreover, a large number of SNPs and/or InDels were found to be present in defence-related genes [43].

In order to identify common genes between biotic and abiotic responses in chickpea, Mantri et al. [71] performed microarray analysis of chickpea ICC 3996 under three abiotic stresses (drought, cold and high salinity) and biotic stress (infection with *A. rabiei*). This analysis revealed 46, 54, 266 and 51 genes to be differentially regulated in drought, cold, high salinity and *A. rabiei* stresses, respectively. *A. rabiei* stress response was found to be more similar to that of high-salinity stress [71].

8. Transcriptome analysis for non-coding RNA studies in chickpea

Non-coding RNAs usually act as regulatory elements that have a decisive role in fine regulation of gene activity. Non-coding transcripts comprise of small and long non-coding RNAs. Small non-coding RNAs regulate diverse developmental processes by controlling gene expression at transcriptional and post-transcriptional level [72, 73]. MicroRNAs (miRNAs) constitute the major class of small non-coding RNAs and are 20–24 nucleotides long key regulatory elements. They are highly conserved and play an important part in various developmental processes in plants such as leaf development, flowering, formation and maintenance of the shoot, floral and axillary meristems, establishment of organ polarity, root nodule symbiosis, vegetative to reproductive phase transition and response to biotic and abiotic stresses [73–77]. In chickpea, small RNA libraries were sequenced from normal tissues and those under different stress conditions [78–80]. Small RNA sequence data were filtered and processed for miRNA prediction using miRDeep pipeline resulting in identification of distinct conserved miRNAs from shoot (302, including Cat-miR156b-5p, Cat-miR156j.1, Cat-miR159.1, Cat-miR169b-5p), root (280, including Cat-miR156c.1, Cat-miR169n, Cat-miR171k-3p), mature leaf (248, including Cat-miR156k, Cat-miR172d.2, Cat-NovmiR319b, Cat-miR167a, Cat-miR167d.2), stem (268, including Cat-miR172c-3p, Cat-miR159.3, Cat-NovmiR319d, Cat-miR171k-3p), flower bud (247, Cat-miR319g.2, Cat-miR167c.2, Cat-miR167d.1, Cat-miR171b-3p.2), flower (293, Cat-miR159.4, Cat-miR159e, Cat-miR171m) and young pod (274, Cat-miR172d.1, Cat-NovmiR159a, Cat-miR167-5p). By ab initio prediction, a total of 109, 76, 123, 100, 106, 98 and 120 novel candidate miRNAs were identified from the above tissues, respectively. Overall 618 miRNAs were identified from all the tissues with the maximum being 373 miRNAs from the shoot and minimum 303 from flower buds. Of the 618 miRNAs predicted, 158 were present in all the tissues, and 29% of the miRNAs were found to be tissue specific. Of the 618 miRNAs, 421 were clustered to 73 miRNA families, and 197 could not find similarity to any miRNA family and were termed putative novel. Chickpea miRNAs targeted a wide range of transcripts involved

in diverse cellular processes including protein turnover and modification, metabolism, transcriptional regulation and signal transduction [78]. A similar kind of study performed in leaf and flower tissue resulted in the prediction of 96 highly conserved miRNAs belonging to 38 miRNA families and 20 novel miRNAs belonging to 17 miRNA families in chickpea [80]. In addition to identification of miRNA from different tissues, studies were also conducted for characterization of miRNA in response to different biotic and abiotic stresses. In one such kind of study, three libraries were sequenced for small RNA identification [79]. Libraries were constructed from fungal-infected (*F. oxysporum* f.sp. *ciceris*), salt-treated and untreated seedlings of chickpea and were sequenced using the Illumina GAIIx platform. The analysis identified 122 conserved miRNAs belonging to 25 different families along with 59 novel miRNAs. miR156, miR396 and miR319 were upregulated in response to salt stress. miR156 and miR396 expression was found to be 1.5 times upregulated in both wilt and salt stresses, indicating a common mechanism implied by chickpea involving these miRNAs to cope up with both the stresses. miR530 was found to be significantly upregulated during wilt stress and may be involved in defence to fungal infection. Three legume-specific miRNAs, miR2111, miR2118 and miR5213, were also indicated to play a critical role in defence to pathogen attack. Targets of miR2111 include F-box protein and TIR (Toll/Interleukin-1 Receptor) domain-containing NBS-LRR disease-resistance proteins, and miR2118 and miR5213 also target the same class of R genes. Interestingly, miR2118 is upregulated following wilt infection and downregulated following salt stress [79].

Long intergenic non-coding (linc) RNAs belong to a class of non-coding transcripts which have a length of at least 200bp lacking coding potential and are transcribed from intergenic region of protein coding genes [81, 82]. Linc RNAs control gene regulation at transcriptional and post-transcriptional level by mechanisms including chromatin modification, promoter binding complex attachment and shielding mRNA degradation by acting as sponge against miRNA [83–85]. RNA-seq data from 11 different tissues of chickpea were used for mining linc-RNA [86]. RNA-seq data were processed using TopHat2 and Cufflinks program using chickpea genome as the reference. From 32,984 transcripts obtained, 5782 putative intergenic transcripts were extracted out and subjected to the optimized pipeline for identification of linc-RNA. After removing potential coding transcripts and transcripts having similarity to protein domains, finally a total of 2248 transcripts were retained as putative chickpea linc-RNAs. About 79%, i.e. 1790 linc-RNAs, could be assigned a putative function. Through expression profiling it was evident that a large number of linc-RNAs have tissue-specific expression in distinct tissues. Along with this several linc-RNAs were found to be targets of miRNAs and were involved in various developmental and reproductive processes [86].

9. Expanding transcriptome data to aid development of molecular markers

A DNA-based molecular marker is a DNA sequence with an identifiable location on the genome that can be transmitted from one generation to the next following the standard laws

of inheritance [87]. Recent years have witnessed an immense interest in generation and utilization of molecular markers, as they provide the essential tools for a variety of genomic applications such as QTL mapping, map-based cloning, marker-assisted breeding, association mapping and genetic diversity assessment. These approaches can be applied to understand the genomic architecture of the crop and can expand the efficiency of breeding programmes, thereby aiding to expedite agricultural research. The advent of NGS has enabled the exploration of thousands of markers across entire genomes and transcriptomes. Although transcriptomics has been majorly used for gene expression analysis, it has also been utilized to identify molecular markers such as SSRs and SNPs especially in the genic regions. Such gene-based markers located in coding regions of the genes greatly enhance the opportunity of precise mapping of genes linked to important traits. Transcriptome sequencing offers another advantage for those crops in which a reference genome is not available. To identify SSRs from transcriptome data, several bioinformatics tools have been developed such as MISA (MicroSatellite identification tool) (pgrc.ipk-gatersleben.de/misa/), RISA (Rapid Identification of SSRs and Analysis of primers) (<http://sol.kribb.re.kr/RISA/>) and RepeatAnalyzer [88]. In chickpea, initially a large number of molecular markers were derived from ESTs. Buhariwalla et al. [89] reported 106 EST-based markers developed from an EST library of root tissue from chickpea. In another study by Choudhary et al. [90], 2131 ESTs were utilized for development of 246 EST-SSR markers. Apart from SSRs, several types of markers such as ESTPs, PIPs and EST-SNPs were developed in chickpea using transcriptome data. For instance, Choudhary et al. [91] reported 125 EST-SSRs, 109 ESTPs, 102 SNPs and 151 ITPs. Gupta et al. [57] reported 367 novel EST-derived functional markers which included 187 EST-SSRs, 130 potential intron polymorphisms (PIPs) and 50 expressed sequence tag polymorphisms (ESTPs). In another study, 71 gene-based SNP markers were developed utilizing candidate chickpea transcripts [92]. However, currently transcriptomic resources can be easily generated by high-throughput NGS technologies and utilized to identify molecular markers very rapidly and cost-effectively. Hiremath et al. [36] utilizing the Roche platform generated about 3000 gene-based markers from a large subset of transcripts derived from different chickpea tissues. Currently, SNPs are the markers of choice and are preferred over the SSRs and other markers because of their genome-wide presence and amenability to high-throughput genotyping. Theoretically, SNP calling may be defined as the process of identifying a single-nucleotide variation from an accession read that differs from the existing reference genome or a de novo assembly at similar nucleotide position. Read assembly files generated by mapping programs such as BWA, Bowtie and SOAP are used to perform SNP calling. Bioinformatics tools such as HaploSNPer [93], SAMtools [94, 95], POLYBAYES [96], SNVer [97] and SOAPsnp [98] have been designed to detect the variations in the NGS data. Comparison of transcriptome datasets from contrasting genotypes could help derive SNPs. To date, several studies have been carried out using NGS technology-based transcriptome sequencing to generate large sets of molecular markers in various crop species including chickpea. For instance, a report by Garg et al. [35] facilitated identification of 4816 SSRs from the de novo assembly of the chickpea transcriptome. In another study, sequencing the transcriptome of *C. reticulatum* (PI489777), the wild relative of chickpea, by GS-FLX 454 technology, generated a total of 4072 SSRs and 36,446 SNPs.

Likewise, Agarwal et al. (2012) sequenced the transcriptome of ICCV2 and identified 5409 SSRs. Amongst these, 130 and 493 SSRs were found to be polymorphic after comparing with the transcriptome of *desi* and *wild* chickpea. In addition to the SSRs, a total of 1986 and 37,954 SNPs were also identified between the *desi*, *kabuli* and *wild* genotypes. Similarly, in another study, 51,632 genic SNPs were identified by 454 transcriptome sequencing of *C. arietinum* and *C. reticulatum* genotypes [99]. In a recent study, Srivastava et al. [100] identified 11,621 differentially expressed genes in root vs shoot tissues using RNA sequencing of a wild perennial *Cicer microphyllum* and integrated above transcriptome profiling with high-resolution QTL mapping in order to identify drought-responsive root-specific genes. The transcriptomic resources, therefore, clearly have remarkable potential to expedite the development of large numbers of molecular markers which can be used in genomic-assisted breeding for developing improved varieties of chickpea.

10. Future perspectives

The last few years have witnessed legume genomics attaining new heights as genomes, and transcriptomes of many model legumes (*M. truncatula*, *L. japonicus*) and crop legumes (*G. max*, *C. cajan*, *P. vulgaris*, *Arachis hypogaea*, *Vignas*, etc.) were sequenced. Transcriptomes of both types of cultivated chickpea (*desi* and *kabuli*) and its wild progenitor (*C. reticulatum*) have also been sequenced. Several studies have been carried out to analyse the transcriptome of chickpea which have led to genome-wide determination of transcript levels in various tissues and developmental pathways as well as during biotic and abiotic stresses. This comprehensive analysis of the chickpea transcriptome has advanced the understanding of the molecular mechanisms underlying several critical biological pathways in chickpea. Moreover, analyses of the non-coding RNAs have revealed potential regulators of important pathways affecting the overall development and stress tolerance in chickpea. Further, transcriptome analysis has also facilitated the development of large sets of genic molecular markers such as SSRs and SNPs that will serve as excellent tools for advancing the chickpea breeding programmes. Overall, the transcriptome sequencing of chickpea has not only provided a deep insight into the gene space and quantitation of gene expression but also an opportunity to isolate genes of interest and functional markers for use in chickpea improvement.

Author details

Chandra Kant, Vimal Pandey, Subodh Verma, Manish Tiwari, Santosh Kumar and Sabhyata Bhatia*

*Address all correspondence to: sabhyatabhatia@nipgr.ac.in

National Institute of Plant Genome Research, New Delhi, India

References

- [1] Varshney RK, Song C, Saxena RK, Azam S, Yu S, Sharpe AG, et al. Draft genome sequence of chickpea (*Cicer arietinum*) provides a resource for trait improvement. *Nature Biotechnology*. 2013;**31**(3):240-246
- [2] Jukanti AK, Gaur PM, Gowda CLL, Chibbar RN. Nutritional quality and health benefits of chickpea (*Cicer arietinum* L.): A review. *British Journal of Nutrition*. 2012;**108**(S1):S11-S26
- [3] Dixon RA, Sumner LW. Legume natural products: Understanding and manipulating complex pathways for human and animal health. *Plant Physiology*. 2003;**131**(3):878-885
- [4] Singh P, Vijaya D, Srinivas K, Wani SP. Potential productivity, yield gap, and water balance of soybean-chickpea sequential system at selected benchmark sites in India. *Global Theme 3: Water, Soil, and Agrobiodiversity Management for Ecosystem Health*. Report No. 1. Patancheru, Andhra Pradesh, India: International Crops Research Institute for the Semi-Arid Tropics. 2002.
- [5] Leport L, Turner NC, French RJ, Barr MD, Duda R, Davies SL, et al. Physiological responses of chickpea genotypes to terminal drought in a Mediterranean-type environment. *European Journal of Agronomy*. 1999;**11**(3-4):279-291
- [6] Johansen C, Krishnamurthy L, Saxena NP, Sethi SC. Genotypic variation in moisture response of chickpea grown under line-source sprinklers in a semi-arid tropical environment. *Field Crops Research*. 1994;**37**(2):103-112
- [7] Varshney RK, Thudi M, Nayak SP, Gaur PM, Kashiwagi J, Krishnamurthy L. Genetic dissection of drought tolerance in chickpea (*Cicer arietinum* L.). *Theoretical and Applied Genetics*. 2014;**127**:445-462
- [8] Pang J, Turner NC, Khan T, Du Y-L, Xiong J-L, Colmer TD, et al. Response of chickpea (*Cicer arietinum* L.) to terminal drought: Leaf stomatal conductance, pod abscisic acid concentration, and seed set. *Journal of Experimental Botany*. 2017; **68**(8):1973-1985
- [9] Pushpavalli R, Quealy J, Colmer TD, Turner NC, Siddique KHM, Rao MV, et al. Salt stress delayed flowering and reduced reproductive success of chickpea (*Cicer arietinum* L.), a response associated with Na⁺ accumulation in leaves. *Journal of Agronomy and Crop Science*. 2016;**202**(2):125-138
- [10] Summerfeld R, Virmani S, Roberts E, Ellis R. Adaptation of chickpea to agroclimatic constraints. *Chickpea in the Nineties: proceedings of the Second International Workshop on Chickpea Improvement*. ICRISAT, Patancheru, India. 1990:61-72
- [11] Kalra N, Chakraborty D, Sharma A, Rai HK, Jolly M, Chander S, et al. Effect of increasing temperature on yield of some winter crops in northwest India. *Current Science*. 2008;**94**(1):82-88

- [12] Singh D, Peters D, Singh P, Singh M. Diurnal patterns of canopy photosynthesis, evapo-transpiration and water use efficiency in chickpea (*Cicer arietinum* L.) under field conditions. *Photosynthesis Research*. 1987;**11**(1):61-69
- [13] Summerfield R, Hadley P, Roberts E, Minchin F, Rawsthorne S. Sensitivity of chickpeas (*Cicer arietinum*) to hot temperatures during the reproductive period. *Experimental Agriculture*. 1984;**20**(01):77-93
- [14] Kaiser WJ. Epidemiology of *Ascochyta rabiei*. In: Singh KB, Saxena MC, editors. *Disease Resistance Breeding in Chickpea*. Aleppo, Syria: ICARDA; 1992. pp. 117-134
- [15] Smithson JB, Thompson JA, Summerfield RJ. Chickpea (*Cicer arietinum* L.). In: Summerfield RJ, Roberts EH, editors. *Grain Legume Crops*. London, UK: Collins; 1985. pp. 312-390
- [16] Duke JA. *Handbook of Legumes of World Economic Importance*. New York: Plenum Press; 1981. pp. 52-57
- [17] Van Emden HF, Ball SL, Rao MR. Pest disease and weed problems in pea lentil and faba bean and chickpea. In: Summerfield RJ, editor. *World Crops: Cool Season Food Legumes*. Dordrecht, The Netherlands: Kluwer Academic Publishers; 1988. pp. 519-534
- [18] Graham PH, Vance CP. Legumes: Importance and constraints to greater use. *Plant Physiology*. 2003;**131**(3):872-877
- [19] Schmutz J, Cannon SB, Schlueter J, Ma J, Mitros T, Nelson W, et al. Genome sequence of the palaeopolyploid soybean. *Nature*. 2010;**463**(7278):178-183
- [20] Young ND, Debelle F, Oldroyd GED, Geurts R, Cannon SB, Udvardi MK, et al. The *Medicago* genome provides insight into the evolution of rhizobial symbioses. *Nature*. 2011;**480**(7378):520-524
- [21] Sato S, Nakamura Y, Kaneko T, Asamizu E, Kato T, Nakao M, et al. Genome structure of the legume, *Lotus japonicus*. *DNA Research*. 2008;**15**(4):227-239
- [22] Varshney RK, Chen W, Li Y, Bharti AK, Saxena RK, Schlueter JA, et al. Draft genome sequence of pigeon pea (*Cajanus cajan*), an orphan legume crop of resource-poor farmers. *Nature Biotechnology*. 2012;**30**(1):83-89
- [23] Schmutz J, McClean PE, Mamidi S, Wu GA, Cannon SB, Grimwood J, et al. A reference genome for common bean and genome-wide analysis of dual domestications. *Nature Genetics*. 2014;**46**(7):707-713
- [24] Bertoli DJ, Cannon SB, Froenicke L, Huang G, Farmer AD, Cannon EKS, et al. The genome sequences of *Arachis duranensis* and *Arachis ipaensis*, the diploid ancestors of cultivated peanut. *Nature Genetics*. 2016;**48**(4):438-446
- [25] Jain M, Misra G, Patel RK, Priya P, Jhanwar S, Khan AW, et al. A draft genome sequence of the pulse crop chickpea (*Cicer arietinum* L.). *The Plant Journal*. 2013;**74**(5):715-729

- [26] Kim MY, Lee S, Van K, Kim TH, Jeong SC, Choi IY, et al. Whole-genome sequencing and intensive analysis of the undomesticated soybean (*Glycine soja* Sieb. and Zucc.) genome. *Proceedings of the National Academy of Sciences of the United States of America*. 2010;**107**(51):22032-22037
- [27] Lam HM, Xu X, Liu X, Chen W, Yang G, Wong FL, et al. Resequencing of 31 wild and cultivated soybean genomes identifies patterns of genetic diversity and selection. *Nature Genetics*. 2010;**42**(12):1053-1059
- [28] Branca A, Paape TD, Zhou P, Briskine R, Farmer AD, Mudge J, et al. Whole-genome nucleotide diversity, recombination, and linkage disequilibrium in the model legume *Medicago truncatula*. *Proceedings of the National Academy of Sciences of the United States of America*. 2011;**108**(42):E864-E870
- [29] Libault M, Farmer A, Joshi T, Takahashi K, Langley RJ, Franklin LD, et al. An integrated transcriptome atlas of the crop model *Glycine max*, and its use in comparative analyses in plants. *The Plant Journal: For Cell and Molecular Biology*. 2010;**63**(1):86-99
- [30] Severin AJ, Woody JL, Bolon YT, Joseph B, Diers BW, Farmer AD, et al. RNA-Seq Atlas of *Glycine max*: A guide to the soybean transcriptome. *BMC Plant Biology*. 2010;**10**:160
- [31] Benedito VA, Torres-Jerez I, Murray JD, Andriankaja A, Allen S, Kakar K, et al. A gene expression atlas of the model legume *Medicago truncatula*. *The Plant Journal: For Cell and Molecular Biology*. 2008;**55**(3):504-513
- [32] Verdier J, Torres-Jerez I, Wang M, Andriankaja A, Allen SN, He J, et al. Establishment of the *Lotus japonicus* Gene Expression Atlas (LjGEA) and its use to explore legume seed maturation. *The Plant Journal: For Cell and Molecular Biology*. 2013;**74**(2):351-362
- [33] Pazhamala LT, Purohit S, Saxena RK, Garg V, Krishnamurthy L, Verdier J, et al. Gene expression atlas of pigeon pea and its application to gain insights into genes associated with pollen fertility implicated in seed formation. *Journal of Experimental Botany*. 2017;**68**(8):2037-2054
- [34] Garg R, Patel RK, Jhanwar S, Priya P, Bhattacharjee A, Yadav G, et al. Gene discovery and tissue-specific transcriptome analysis in chickpea with massively parallel pyrosequencing and web resource development. *Plant Physiology*. 2011;**156**(4):1661-1678
- [35] Garg R, Patel RK, Tyagi AK, Jain M. De novo assembly of chickpea transcriptome using short reads for gene discovery and marker identification. *DNA Research*. 2011;**18**(1):53-63
- [36] Hiremath PJ, Farmer A, Cannon SB, Woodward J, Kudapa H, Tuteja R, et al. Large-scale transcriptome analysis in chickpea (*Cicer arietinum* L.), an orphan legume crop of the semi-arid tropics of Asia and Africa. *Plant Biotechnology Journal*. 2011;**9**(8):922-931
- [37] Jhanwar S, Priya P, Garg R, Parida SK, Tyagi AK, Jain M. Transcriptome sequencing of wild chickpea as a rich resource for marker development. *Plant Biotechnology Journal*. 2012;**10**(6):690-702

- [38] Singh VK, Garg R, Jain M. A global view of transcriptome dynamics during flower development in chickpea by deep sequencing. *Plant Biotechnology Journal*. 2013;**11**(6):691-701
- [39] Pradhan S, Bandhiwal N, Shah N, Kant C, Gaur R, Bhatia S. Global transcriptome analysis of developing chickpea (*Cicer arietinum* L.) seeds. *Frontiers in Plant Science*. 2014;**5**:698
- [40] Kant C, Pradhan S, Bhatia S. Dissecting the root nodule transcriptome of chickpea (*Cicer arietinum* L.). *PLoS One*. 2016;**11**(6):e0157908
- [41] Garg R, Bhattacharjee A, Jain M. Genome-scale transcriptomic insights into molecular aspects of abiotic stress responses in chickpea. *Plant Molecular Biology Reporter*. 2015;**33**(3):388-400
- [42] Garg R, Shankar R, Thakkar B, Kudapa H, Krishnamurthy L, Mantri N, et al. Transcriptome analyses reveal genotype- and developmental stage-specific molecular responses to drought and salinity stresses in chickpea. *Scientific Reports*. 2016;**6**:19228
- [43] Jain M, Pole AK, Singh VK, Ravikumar RL, Garg R. Discovery of molecular markers for Fusarium wilt via transcriptome sequencing of chickpea cultivars. *Molecular Breeding*. 2015;**35**(10):198
- [44] Bajaj D, Upadhyaya HD, Khan Y, Das S, Badoni S, Shree T, et al. A combinatorial approach of comprehensive QTL-based comparative genome mapping and transcript profiling identified a seed weight-regulating candidate gene in chickpea. *Scientific Reports*. 2015;**5**:9264
- [45] Das S, Upadhyaya HD, Bajaj D, Kujur A, Badoni S, Laxmi, et al. Deploying QTL-seq for rapid delineation of a potential candidate gene underlying major trait-associated QTL in chickpea. *DNA Research*. 2015;**22**(3):193-203
- [46] Gaur R, Jeena G, Shah N, Gupta S, Pradhan S, Tyagi AK, et al. High density linkage mapping of genomic and transcriptomic SNPs for synteny analysis and anchoring the genome sequence of chickpea. *Scientific Reports*. 2015;**5**:13387
- [47] Kale SM, Jaganathan D, Ruperao P, Chen C, Punna R, Kudapa H, et al. Prioritization of candidate genes in "QTL-hotspot" region for drought tolerance in chickpea (*Cicer arietinum* L.). *Scientific Reports*. 2015;**5**:15296
- [48] Kujur A, Upadhyaya HD, Bajaj D, Gowda CLL, Sharma S, Tyagi AK, et al. Identification of candidate genes and natural allelic variants for QTLs governing plant height in chickpea. *Scientific Reports*. 2016;**6**:27968
- [49] Pushpavalli R, Krishnamurthy L, Thudi M, Gaur PM, Rao MV, Siddique KH, et al. Two key genomic regions harbour QTLs for salinity tolerance in ICCV 2 × JG 11 derived chickpea (*Cicer arietinum* L.) recombinant inbred lines. *BMC Plant Biology*. 2015;**15**(1):124
- [50] Doddamani D, Khan AW, Katta MAVSK, Agarwal G, Thudi M, Ruperao P, et al. CicArVarDB: SNP and InDel database for advancing genetics research and breeding applications in chickpea. *Database*. 2015;**2015**:bav078-bav

- [51] Mardis ER. Next-generation DNA sequencing methods. *Annual Review of Genomics and Human Genetics*. 2008;**9**:387-402
- [52] Morozova O, Marra MA. Applications of next-generation sequencing technologies in functional genomics. *Genomics*. 2008;**92**(5):255-264
- [53] Schuster SC. Next-generation sequencing transforms today's biology. *Nature Methods*. 2008;**5**(1):16-18
- [54] Jayashree B, Buhariwalla HK, Shinde S, Crouch JH. A legume genomics resource: The Chickpea Root Expressed Sequence Tag Database. *Electronic Journal of Biotechnology*. 2005;**8**(2):128-133
- [55] Agarwal G, Jhanwar S, Priya P, Singh VK, Saxena MS, Parida SK, et al. Comparative analysis of kabuli chickpea transcriptome with desi and wild chickpea provides a rich resource for development of functional markers. *PLoS One*. 2012;**7**(12):e52443
- [56] Verma M, Kumar V, Patel RK, Garg R, Jain M. CTDB: An integrated chickpea transcriptome database for functional and applied genomics. *PLoS One*. 2015;**10**(8):e0136880
- [57] Gupta S, Garg V, Bhatia S. A new set of ESTs from chickpea (*Cicer arietinum* L.) embryo reveals two novel F-box genes, CarF-box_PP2 and CarF-box_LysM, with potential roles in seed development. *PLoS One*. 2015;**10**(3):e0121100
- [58] Afonso-Grunz F, Molina C, Hoffmeier K, Rycak L, Kudapa H, Varshney RK, et al. Genome-based analysis of the transcriptome from mature chickpea root nodules. *Frontiers in Plant Science*. 2014;**5**:325
- [59] Ahmad F, Gaur P, Croser J. Chickpea (*Cicer arietinum* L.). In: Singh R, Jauhar P, editors. *Genetic Resources, Chromosome Engineering and Crop Improvement—Grain Legumes*. USA: CRC Press; 2005. pp. 185-214
- [60] Sharma KD, Nayyar H. Cold stress alters transcription in meiotic anthers of cold tolerant chickpea (*Cicer arietinum* L.). *BMC Research Notes*. 2014;**7**:717
- [61] Deokar AA, Kondawar V, Jain PK, Karuppayil SM, Raju NL, Vadez V, et al. Comparative analysis of expressed sequence tags (ESTs) between drought-tolerant and -susceptible genotypes of chickpea under terminal drought stress. *BMC Plant Biology*. 2011;**11**(1):70
- [62] Mantri NL, Ford R, Coram TE, Pang EC. Transcriptional profiling of chickpea genes differentially regulated in response to high-salinity, cold and drought. *BMC Genomics*. 2007;**8**:303
- [63] Molina C, Rotter B, Horres R, Udupa SM, Besser B, Bellarmino L, et al. SuperSAGE: The drought stress-responsive transcriptome of chickpea roots. *BMC Genomics*. 2008;**9**:553
- [64] Molina C, Zaman-Allah M, Khan F, Fatnassi N, Horres R, Rotter B, et al. The salt-responsive transcriptome of chickpea roots and nodules via deepSuperSAGE. *BMC Plant Biology*. 2011;**11**:31

- [65] Varshney RK, Hiremath PJ, Lekha P, Kashiwagi J, Balaji J, Deokar AA, et al. A comprehensive resource of drought- and salinity- responsive ESTs for gene discovery and marker development in chickpea (*Cicer arietinum* L.). *BMC Genomics*. 2009;**10**:523
- [66] Wang X, Liu Y, Jia Y, Gu H, Ma H, Yu T, et al. Transcriptional responses to drought stress in root and leaf of chickpea seedling. *Molecular Biology Reports*. 2012;**39**(8):8147-8158
- [67] Dinari A, Niazi A, Afsharifar AR, Ramezani A. Identification of upregulated genes under cold stress in cold-tolerant chickpea using the cDNA-AFLP approach. *PLoS One*. 2013;**8**(1):e52757
- [68] Coram TE, Pang ECK. Isolation and analysis of candidate ascochyta blight defence genes in chickpea. Part I. Generation and analysis of an expressed sequence tag (EST) library. *Physiological and Molecular Plant Pathology*. 2005;**66**(5):192-200
- [69] Coram TE, Pang EC. Expression profiling of chickpea genes differentially regulated during a resistance response to *Ascochyta rabiei*. *Plant Biotechnology Journal*. 2006;**4**(6):647-666
- [70] Ashraf N, Ghai D, Barman P, Basu S, Gangisetty N, Mandal MK, et al. Comparative analyses of genotype dependent expressed sequence tags and stress-responsive transcriptome of chickpea wilt illustrate predicted and unexpected genes and novel regulators of plant immunity. *BMC Genomics*. 2009;**10**(1):415
- [71] Mantri NL, Ford R, Coram TE, Pang ECK. Evidence of unique and shared responses to major biotic and abiotic stresses in chickpea. *Environmental and Experimental Botany*. 2010;**69**(3):286-292
- [72] Bartel DP. MicroRNAs: Genomics, biogenesis, mechanism, and function. *Cell*. 2004;**116**(2):281-297
- [73] Jones-Rhoades MW, Bartel DP, Bartel B. MicroRNAs and their regulatory roles in plants. *Annual Review of Plant Biology*. 2006;**57**:19-53
- [74] Dezulian T, Palatnik JF, Huson D, Weigel D. Conservation and divergence of microRNA families in plants. *Genome Biology*. 2005;**6**(11):P13
- [75] Rogers K, Chen X. Biogenesis, turnover, and mode of action of plant microRNAs. *Plant Cell*. 2013;**25**(7):2383-2399
- [76] Simon SA, Meyers BC, Sherrier DJ. MicroRNAs in the rhizobia legume symbiosis. *Plant Physiology*. 2009;**151**(3):1002-1008
- [77] Sunkar R. MicroRNAs with macro-effects on plant stress responses. *Seminars in Cell & Developmental Biology*. 2010;**21**(8):805-811
- [78] Jain M, Chevala VV, Garg R. Genome-wide discovery and differential regulation of conserved and novel microRNAs in chickpea via deep sequencing. *Journal of Experimental Botany*. 2014;**65**(20):5945-5958

- [79] Kohli D, Joshi G, Deokar AA, Bhardwaj AR, Agarwal M, Katiyar-Agarwal S, et al. Identification and characterization of Wilt and salt stress-responsive microRNAs in chickpea through high-throughput sequencing. *PLoS One*. 2014;**9**(10):e108851
- [80] Srivastava S, Zheng Y, Kudapa H, Jagadeeswaran G, Hivrale V, Varshney RK, et al. High throughput sequencing of small RNA component of leaves and inflorescence revealed conserved and novel miRNAs as well as phasiRNA loci in chickpea. *Plant Science*. 2015;**235**:46-57
- [81] Guttman M, Amit I, Garber M, French C, Lin MF, Feldser D, et al. Chromatin signature reveals over a thousand highly conserved large non-coding RNAs in mammals. *Nature*. 2009;**458**(7235):223-227
- [82] Ulitsky I, Bartel DP. lincRNAs: Genomics, evolution, and mechanisms. *Cell*. 2013;**154**(1):26-46
- [83] Cabili MN, Trapnell C, Goff L, Koziol M, Tazon-Vega B, Regev A, et al. Integrative annotation of human large intergenic noncoding RNAs reveals global properties and specific subclasses. *Genes & Development*. 2011;**25**(18):1915-1927
- [84] Imig J, Brunschweiler A, Brummer A, Guennewig B, Mittal N, Kishore S, et al. miR-CLIP capture of a miRNA targetome uncovers a lincRNA H19-miR-106a interaction. *Nature Chemical Biology*. 2015;**11**(2):107-114
- [85] Rinn JL, Chang HY. Genome regulation by long noncoding RNAs. *Annual Review of Biochemistry*. 2012;**81**:145-166
- [86] Khemka N, Singh VK, Garg R, Jain M. Genome-wide analysis of long intergenic non-coding RNAs in chickpea and their potential role in flower development. *Scientific Reports*. 2016;**6**:33297
- [87] Semagn K, Bjørnstad Å, Ndjiondjop M. Principles, requirements and prospects of genetic mapping in plants. *African Journal of Biotechnology*. 2006;**5**(25):2569-2587
- [88] Catanese HN, Brayton KA, Gebremedhin AH. RepeatAnalyzer: A tool for analysing and managing short-sequence repeat data. *BMC Genomics*. 2016;**17**(1):422
- [89] Buhariwalla HK, Jayashree B, Eshwar K, Crouch JH. Development of ESTs from chickpea roots and their use in diversity analysis of the *Cicer* genus. *BMC Plant Biology*. 2005;**5**(1):16
- [90] Choudhary S, Sethy NK, Shokeen B, Bhatia S. Development of chickpea EST-SSR markers and analysis of allelic variation across related species. *Theoretical and Applied Genetics*. 2009;**118**(3):591-608
- [91] Choudhary S, Gaur R, Gupta S. EST-derived genic molecular markers: Development and utilization for generating an advanced transcript map of chickpea. *Theoretical and Applied Genetics*. 2012;**124**(8):1449-1462

- [92] Nayak SN, Zhu H, Varghese N, Datta S, Choi H-K, Horres R, et al. Integration of novel SSR and gene-based SNP marker loci in the chickpea genetic map and establishment of new anchor points with *Medicago truncatula* genome. Theoretical and Applied Genetics. 2010;**120**(7):1415-1441
- [93] Tang J, Leunissen JA, Voorrips RE, van der Linden CG, Vosman B. HaploSNPer: A web-based allele and SNP detection tool. BMC Genetics. 2008;**9**:23
- [94] Li H. A statistical framework for SNP calling, mutation discovery, association mapping and population genetical parameter estimation from sequencing data. Bioinformatics. 2011;**27**(21):2987-2993
- [95] Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The sequence alignment/map format and SAMtools. Bioinformatics. 2009;**25**(16):2078-2079
- [96] Marth GT, Korf I, Yandell MD, Yeh RT, Gu Z, Zakeri H, et al. A general approach to single-nucleotide polymorphism discovery. Nature Genetics. 1999;**23**(4):452-456
- [97] Wei Z, Wang W, Hu P, Lyon GJ, Hakonarson H. SNVer: A statistical tool for variant calling in analysis of pooled or individual next-generation sequencing data. Nucleic Acids Research. 2011;**39**(19):e132
- [98] Li R, Li Y, Fang X, Yang H, Wang J, Kristiansen K, et al. SNP detection for massively parallel whole-genome resequencing. Genome Research. 2009;**19**(6):1124-1132
- [99] Deokar AA, Ramsay L, Sharpe AG, Diapari M, Sindhu A, Bett K, et al. Genome wide SNP identification in chickpea for use in development of a high density genetic map and improvement of chickpea reference genome assembly. BMC Genomics. 2014;**15**(1):708
- [100] Srivastava R, Bajaj D, Malik A, Singh M, Parida SK. Transcriptome landscape of perennial wild *Cicer microphyllum* uncovers functionally relevant molecular tags regulating agronomic traits in chickpea. Scientific Reports. 2016; 6:33616