We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Vitamin C: Sources, Functions, Sensing and Analysis

Sudha J. Devaki and Reshma Lali Raveendran

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.70162>

Abstract

Vitamin C is a water-soluble compound found in living organisms. It is an essential nutrient for various metabolism in our body and also serves as a reagent for the preparation of many materials in the pharmaceutical and food industry. In this perspective, this chapter can develop interest and curiosity among all practicing scientists and technologists by expounding the details of its sources, chemistry, multifunctional properties and applications.

Keywords: vitamin C, biomarker, antioxidant, sensors

1. Introduction

Vitamin C also known as ascorbic acid (AA) is an essential nutrient in many multicellular organisms, especially in humans. Ascorbic acid is a water-soluble vitamin and is found in variable quantities in fruits and vegetables and organ meats (e.g. liver and kidney). Deficiency of vitamin C causes scurvy, widespread connective tissue weakness and capillary fragility. Among chemists, it is used as a reagent for the preparation of fine chemicals, enzymatic reagent and nanomaterials. Consequently, the detection and quantification of ascorbic acid in food samples, products and nutraceuticals is receiving overwhelming importance among researchers, medical practitioners and also in the pharmaceutical and food industry. **Figure 1** shows the schematic representation of the sources and multifunctional applications of vitamin C in the metabolism of our body.

2. Sources

Vitamin C (**Figure 2**) is abundantly available in many natural sources, including fresh fruits and vegetables. The richest sources of ascorbic acid including Indian gooseberry, citrus fruits

Figure 1. Sources and multifunctional role of vitamin C in metabolic processes in human body.

such as limes, oranges and lemons, tomatoes, potatoes, papaya, green and red peppers, kiwi-fruit, strawberries and cantaloupes, green leafy vegetables such as broccoli, fortified cereals and its juices are also rich sources of vitamin C.

Another source of vitamin C is animals. They usually synthesize their own vitamin C and are highly concentrated in the liver part [1, 2]. Sources of vitamin C and its content are given in **Figure 3**. Average daily recommended amounts of vitamin C for different ages are also given in **Table 1**.

Figure 2. Structure of vitamin C.

Figure 3. Amount of vitamin C in different sources.

Infants					
Age			Adequate intake (AI) mg/day		Upper intake level (UL)
0–6 months ^a			25		–
7–12 months ^b			30		–
Children and adolescents					
Age	EAR ^c		RDI ^d		
	mg/day		mg/day		mg/day
1–3	25		35		400
4–8	25		35		650
9–14	28		40		1200
14–18	28		40		1800
Adults					
Men			Women		Both
Age	EAR ^c	RDI ^d	EAR ^c	RDI ^d	
	mg/day	mg/day	mg/day	mg/day	mg/day
19–30	30	45	30	45	2000
31–50	30	45	30	45	2000
51–70	30	45	30	45	2000
>70	30	45	30	45	2000

Infants					
Age			Adequate intake (AI) mg/day		Upper intake level (UL)
Motherhood					
Pregnancy			Lactation		Both
Age	EAR ^c	RDI ^d	EAR ^c	RDI ^d	
	mg/day	mg/day	mg/day	mg/day	mg/day
14–18	38	55	58	80	2000
19–30	40	60	60	85	2000
31–50	40	60	60	85	2000

^aCalculated as per the average intake of breast milk.

^bCalculated on a body weight basis of infants.

^cEstimated average requirement.

^dRecommended dietary intake.

Table 1. Dietary intake of vitamin C [3–5].

3. Functions

Vitamin C plays an important role in many physiological processes in humans. It is needed for the repair of tissues in all parts of the body. The important functions of vitamin C include the formation of protein used to make skin, tendons, ligaments, and blood vessels for healing wounds and forming scar tissue, for repairing and maintaining cartilage, bones, and teeth and aid in the absorption of iron. It can also act as a reducing and capping agent for metal nanoparticles.

3.1. As reducing and capping agent

Ascorbic acid acts as a reducing and capping agent for the synthesis of metal nanoparticles such as silver, gold, copper, etc. Ascorbic acid molecules can cap or surround the particle and prevent the uncontrolled growth of the particles to micron-sized dimensions. A study by Khan et al., in 2016 reported the synthesis of copper nanoparticle by using ascorbic acid as both the reducing agent [6]. Sun et al. reported in *Journal of Materials Science* in 2009 that gold nanoparticles can be synthesised in reverse micelles without the addition or introduction of any other reducing or capping reagent [7]. In analytical methods in 2014, D’souza et al. reported the use of AA-Au NPs as a colorimetric probe for detection of dichlorvos in water and wheat samples. The influence of AA concentration on the aggregation induced by dichlorvos in AA-AuNPs (**Figure 4**) and the optical property of the AA-Au NPs was investigated by UV-Vis spectroscopy [8].

Figure 4. Analytical process for detecting dichlorvos using AA-Au NPs as a colorimetric probe [8].

3.2. Antioxidant activity

One of the important properties of vitamin C is its antioxidant activity. Antioxidant activity of vitamin C helps to prevent certain diseases such as cancer, cardiovascular diseases, common cold, age-related muscular degeneration and cataract.

3.3. In cancer treatment

Since 1970, it has been known that high dose of vitamin C has beneficial effects on the survival time in patients with terminal cancer, which was reported by Cameron, Campbell, and Pauling. Research is undergoing in detail for using vitamin C in cancer treatment [9–11]. One of the studies suggests that pharmacologic doses of vitamin C might show promising effects on the treatment of tumours [12]. Vitamin C can act as pro-oxidant and it can generate hydrogen peroxide [13, 14]. Administration of high dose of vitamin C gives long survival times for patients with advanced cancers.

The continuing attack of DNA by unquenched reactive oxygen species is believed to cause cancer. As a physiological antioxidant, ascorbic acid plays a role in the prevention of oxidative damage to DNA, which is elevated in cells at sites of chronic inflammation and in many pre-neoplastic lesions. Most DNA damage is repaired metabolically; however, the frequency of elevated steady-state levels of oxidized DNA bases is estimated to be sufficient to cause mutational events. The base damage product 8-hydroxy-2'-deoxyguanosine has been found to be elevated in individuals with severe vitamin C deficiency and to be reduced by supplementation with vitamins C and E [15].

Oxidative stress, arising as a result of an imbalance between free radical production and antioxidant defences, causes damage to a wide range of molecular species including lipids, proteins and nucleic acids. Oxidative stress caused by free radicals is shown in **Figure 5**. The best way to ensure adequate intake of the antioxidant nutrients, such as vitamins C and E and various kinds of minerals, is through a balanced diet consisting of 5–8 servings of fruits and vegetables per day.

3.4. In cardiovascular diseases

The antioxidant property of vitamin C helps for the treatment of cardiovascular diseases. Vitamin C has the capability for reducing monocyte adherence to the endothelium, improving endothelium-dependent nitric oxide production and vasodilation and reducing vascular smooth-muscle-cell apoptosis, which prevents plaque instability in atherosclerosis [15]. The oxidative damage including the oxidative modification of low-density lipoproteins is a major cause of cardiovascular disease. The antioxidant property of vitamin C helps to reduce this to a certain extent [16, 17].

3.5. In common cold

Pauling in 1970 suggested that vitamin C can be used for the treatment of common cold [18]. There are so many reports in Cochrane Database Syst. Review showing the use of prophylactic

Figure 5. Oxidative stress by free radicals.

vitamin C reduces the cold duration in adults and children [19]. The use of vitamin C might reduce the duration of common cold due to its anti-histamine effect of high dose of vitamin C [20]. However, the results are inconsistent and still research is undergoing in this field. There are Database Syst. Review showing the use of prophylactic vitamin C reduces the cold duration in adults and children [19]. The use of vitamin C might reduce the duration of common cold due to its anti-histamine effect of high dose of vitamin C [20].

3.6. In age-related macular degeneration (AMD) and cataract

Age-related macular degeneration (AMD) and cataracts are two of the main causes of vision loss in older. Oxidative stress might contribute to the aetiology of both conditions. Thus, researchers have taken interest in the role of vitamin C and other antioxidants in the development and treatment of these diseases. There are many reports to study the role of vitamin C in AMD and cataract [21–23]. Results from two studies indicate that vitamin C intakes greater than 300 mg/day reduce the risk of cataract formation by 75% [16, 24, 25].

All the studies indicate that the vitamin C formulations might slow AMD progression and reduce the high risk of developing advanced AMD. AMD is shown in **Figure 6**.

3.7. Antioxidant mechanism

Vitamins C can protect the body against the destructive effects of free radicals. Antioxidants neutralize free radicals by donating one of their own electrons, ending the electron-stealing reaction, as shown in **Figure 7**. The antioxidant nutrients themselves do not become free radicals by donating an electron because they are stable in either form or act as scavengers, helping to prevent cell and tissue damage that could lead to cellular damage and disease.

Ascorbic acid reacts with free radicals undergoing single-electron oxidation to produce a relatively poor reactive intermediate, the ascorbyl radical, which disproportionates to ascorbate and dehydroascorbate. Thus, ascorbic acid can reduce toxic, reactive oxygen species superoxide anion ($O_2^{\bullet-}$) and hydroxyl radical ($OH^{\bullet}$), as well as organic ($RO_2^{\bullet}$) and nitrogen ($NO_2^{\bullet}$) oxy

Figure 6. Age-related macular degeneration.

Figure 7. Antioxidant mechanism.

radicals. Those reactions are likely to be of fundamental importance in all aerobic cells. This reaction is the basis of most of the biological functions of ascorbic acid. The mechanism of free radical action on DNA is shown in Figure 8.

Vitamin C also has role in protecting other vitamins (vitamin A and vitamin E) from the harmful effects of oxidation. Vitamin C helps in protecting gums and retards ageing. It strengthens the general physical condition by removing toxic metals from the body. Vitamin C reduces the formation of cataract and hence useful in the treatment of glaucoma.

Figure 8. Mechanism of free radical action on DNA.

3.8. Synthesis of protein

Another important function of vitamin C is its role in the synthesis of protein. Vitamin C helps in the synthesis of collagen. Collagen protects our skin from wrinkling and makes our skin firm and strong. Collagen also protects and supports organs and other soft tissues. One of the amino acids used to build collagen—hydroxyproline—is only synthesized when vitamin C is available. Functions of vitamin C on skin.

3.9. Functions of vitamin C on skin

3.9.1. Photoprotection

Vitamin C reduces the damage caused by UV-light exposure. It cannot act as a sunscreen since it cannot absorb UV light. But the antioxidant activity of vitamin C helps to protect UV damage caused by free radicals [26]. In response to UV light, vitamin C transport proteins are increased, suggesting an increased need for vitamin C uptake for adequate protection [27, 28]. Addition of keratinocytes on vitamin C reduces damages caused by UV light and lipid peroxidation, limits the release of pro-inflammatory cytokines and protects against apoptosis [29]. Many studies have suggested that vitamin C consumption alone will not reduce the effect of UV exposure; however, a combination of vitamin C and E effectively increases minimal erythema dose (MED) (the lowest dose of ultraviolet radiation (UVR) that will produce a detectable erythema 24 hours after UVR exposure) and decreases erythema-induced blood flow to damaged areas of skin [30]. Thus, interactions between the two antioxidant vitamins may be necessary to achieve UV protection. The topical application of vitamin C also reduces the effect of UV exposure and skin wrinkling and skin tumour [31]. Vitamin C reduced the number of sunburned cells, decreased erythema response and reduced DNA damage induced by UV exposure [27]. The combination of antioxidant vitamins decreased the immunosuppressive effects of UV exposure, increased MED and decreased cell damage [32, 33].

3.9.2. Wound healing

Vitamin C plays a key role in healing wound by the formation of collagen, connective tissue [34–36]. The new tissue is rebuilt with the help of collagen framework. This function is supported by its co-factor vitamin C. Besides this, vitamin C performs as a strong antioxidant and immune system modulator [37, 38].

3.10. Deficiency of vitamin C

Deficiency of vitamin C in humans causes a major disease called scurvy. The major signs of the disease occur primarily in mesenchymal tissues. It leads to impaired wound healing; oedema; haemorrhage (due to deficient formation of intercellular substance) in the skin, mucous membranes, internal organs, and muscles; and weakening of collagenous structures in bone, cartilage, teeth and connective tissues. Those who suffer from scurvy have swollen, bleeding gums with tooth loss. They also show lethargy, fatigue, rheumatic pains in the legs, muscular atrophy and skin lesions, massive sheet hematomas in the thighs,

and ecchymoses and haemorrhages in many organs, including the intestines, sub-periosteal tissues and eyes. All these features are accompanied by psychological changes: hysteria, hypochondria and depression. In children, the syndrome is called Moeller-Barlow disease; it is seen in non-breastfed infants usually at about 6 months of age and is characterized by widening of bone-cartilage boundaries, stressed epiphyseal cartilage of the extremities, severe joint pain and frequently, anaemia and fever. Children having this disease present with a limp or inability to walk, tenderness of the lower limbs, bleeding of the gums and petechial haemorrhages.

4. Detection and sensing

Many analytical techniques are used for the determination of vitamin C in different matrices, such as titrimetric [39], fluorimetric [40], spectrophotometric [41], high-performance liquid chromatography [42], enzymatic [43], kinetic [44, 45] and electrochemical, etc.

4.1. High-performance liquid chromatography (HPLC)

High-performance liquid chromatography (HPLC) methods are preferred earlier because they are faster and more effective than spectrophotometric, titration or enzymatic methods, and they do not usually need derivatization [46]. In pharmaceutical and cosmetic industries, HPLC is used, which is considered as a sensitive and selective method.

Detection of vitamin C through HPLC has been done by many groups. Racz et al. in 1990 reported that HPLC used the HPLC method for the determination of ascorbic acid in fruits and vegetables. The method has been used to deep-frozen raspberry cream analysis together with three commonly used chemical methods of vitamin C analysis. The HPLC method has been compared with the chemical methods from several aspects, and the superiority of HPLC method has been concluded [47].

Snezana et al. reported the HPLC method for the determination of vitamin C in pharmaceutical samples in *Tropical Journal of Pharmaceutical Research* in 2011 [48]. The simplicity of this low-cost, rapid technique and its high specificity to ascorbic acid, even in the presence of a variety of excipients, demonstrate that this HPLC method would be particularly suitable for the determination of ascorbic acid (**Figure 9**). It can be used in pharmaceutical/veterinary formulations without prior sample preparation.

Another work carried out by Franco et al., which was to optimize and validate a new analytic strategy for the determination of vitamin C in strawberries by UV-HPLC [49].

For the accurate and reliable measurement of ascorbic acid and dehydroascorbic acid, HPLC can be used in combination with electrochemical or ultraviolet detection. Line and Hoffer have reported a method for the determination of vitamin C in plasma. It can be analysed by HPLC in connection with electrochemical or ultraviolet light detection. Electrochemical HPLC has advantages over UV-HPLC for plasma total vitamin C analysis [50].

Figure 9. HPLC plot of ascorbic acid detection.

4.2. Spectrophotometric method

Among many analytical methods, spectrophotometric methods are very simple and low-cost. Several studies used the spectrophotometric method for the determination of ascorbic acid. Güçlü et al. [51] have proposed a spectrophotometric method based on ascorbic acid oxidation to dehydroascorbic acid, by using the Cu(II)-neocuproine complex, which is reduced to Cu(I)-bis(neocuproine), the absorbance of the latter being determined at 450 nm. Other optical methods for vitamin C estimation include spectrophotometrical determination of iodine reacted with ascorbic acid [52] and chemiluminescence [53].

A sensitive, simple and low-cost spectrophotometric method was introduced by Kobra and Somayye in 2015. The present method was successfully applied to determine the ascorbic acid in food and pharmaceutical samples. The samples were multivitamin tablet, effervescent tablet, vitamin C injection, natural orange juice, orange syrup powdered and commercial orange liquid. The method is based on the reaction of AgNO_3 with ascorbic acid in the presence of polyvinyl pyrrolidone (PVP) and slightly basic medium to prepare silver nanoparticles [54].

Kapur et al. in 2015 reported a method that is based on the oxidation of ascorbic acid to dehydroascorbic acid by bromine water in the presence of acetic acid. In this method, the total ascorbic acid (ascorbic acid + dehydroascorbic acid) has been determined in 21 different samples of fruits and vegetables by the spectrophotometric method [55]. Mohammed and Hazim in 2016 reported a UV-spectrophotometric method for the determination of ascorbic acid in fruits and vegetables from hill region with 2,4-dinitrophenylhydrazine [56].

A new sensitive colorimetric method for the determination of ascorbic acid tablet in aqueous solution was reported by Ahmed and Mohamed in 2013. The method is based on the formation of coloured azo dye by diazotization of 2,4-dichloroaniline, followed by azo-coupling reaction between the resulting product and ascorbic acid [57].

4.3. Electrochemical sensing

Electrochemical sensing methods are more widely applied as they are simple, sensitive and moderate methods. Ascorbic acid (AA), dopamine and catecholamine are important neurotransmitters in the human body. Hence, the low-level detection of ascorbic acid is very important. All the methods mentioned above involve complicated pretreatment techniques and expensive instruments. Hence, researchers have developed a more simple, sensitive and accurate electrochemical method for the detection of ascorbic acid. Several studies have reported the electrochemical sensing methods for detection of ascorbic acid. Attempts to simplify such methods have resulted in the development of new methods such as the nickel hexacyanoferrate film-modified aluminium electrode [58] and graphite epoxy electrode [59]. These methods use specific and modified working electrode systems that are complicated [60]. These electrochemical methods are widely used to monitor AA metabolites *in vivo* or *in vitro* [61, 62]. For example, batch injection amperometric methods can achieve a detection limit as low as 450 mg/L [63]. The metallic nickel electrode (2610–6 M), chemiluminometric flow method (5l M) [64], carbon paste mixed electrode (1.08610–5 M) [65], polyviologen-modified technique (0.38l M) [66] and NiGCNF plating-modified electrode (2610–6 M) [67] achieve very sensitive detection ranges.

Suw et al. in 2004 developed a stripping voltammetric technique that manifests faster response, more cost-effective and sensitive preconcentration techniques, which was used along with a simpler glassy carbon electrode [68]. In this method, low ascorbic acid concentrations were detected by the square wave stripping voltammetry and a glassy carbon electrode. The lower detection limit reported as 0.30l g/L (S/N = 3). This method can be used to detect biological materials, pharmaceuticals, food and drugs.

Many reports are there for the simultaneous electrochemical detection of ascorbic acid in the presence of dopamine and uric acid on the glassy carbon electrode. From our own group, we have synthesized electrically conducting poly (3,4-ethylenedioxythiophene) nanospindles (PEDOTs) and used for the electrochemical sensing of ascorbic acid [69] (**Figure 10**).

Figure 10. Cyclic voltammogram of electrochemical sensing of ascorbic acid using a polyaniline-modified platinum electrode [70].

In another study, our group developed a low-cost electrochemical sensor based on a platinum electrode for ascorbic acid conductive polyaniline-based composite. This unique low-cost and user-friendly sensor was validated for the nanomolar detection of AA. The lower detection limit for AA was observed at 0.1 nM. The aqueous PPICS/Pt electrode could serve as a prospective low-cost and efficient electrochemical sensor and provide promising and outstanding contributions to the food, beverage and medical industries [70].

Electrochemical methods have attracted much attention from clinical diagnostic perspectives because of their easy operation, low cost, rapid response, high sensitivity and good selectivity.

5. Conclusion

Vitamin C plays a pivotal role in body-building process and in disease prevention. The various functions of vitamin C, including the antioxidant activity, formation of protein, tendons, ligaments and blood vessels, for healing wounds and form scar tissue, for repairing and maintaining cartilage, bone, and teeth, and aiding in the absorption of iron, were discussed. This chapter will definitely benefit the students, researchers and technologists globally.

Author details

Sudha J. Devaki* and Reshma Lali Raveendran

*Address all correspondence to: sudhajd2001@yahoo.co.in

Chemical Sciences and Technology Division, CSIR-National Institute for Interdisciplinary Science and Technology (CSIR-NIIST), Thiruvananthapuram, India

References

- [1] Food and Nutrition Board. Dietary Reference Intakes for Vitamin C, Vitamin E, Selenium, and Carotenoids. Washington, DC: National Academy Press; 2000
- [2] Nutrient Data Laboratory. USDA National Nutrient Database for Standard Reference, Release 24 [Internet]. Available from <http://www.ars.usda.gov/ba/bhnrc/nd>
- [3] Committee on Medical Aspects of Food Policy; 1991
- [4] Food and Nutrition Board: Institute of Medicine; 2000
- [5] Food and Agricultural Organization, World Health Organization; 2002
- [6] Khan A, Rashid A, Younas R, Chong R. A chemical reduction approach to the synthesis of copper nanoparticles. *International Nano Letters*. 2016;6:21-26.

- [7] Sun K, Qiu J, Liu J, Miao Y. Preparation and characterization of gold nanoparticles using ascorbic acid as reducing agent in reverse micelles. *J Mater Sci.* 2009;**44**:754
- [8] D'souza SL, Pati RK, Kailasa SK. Ascorbic acid functionalized gold nanoparticles as a probe for colorimetric and visual read-out determination of dichlorvos in environmental samples. *Analytical Methods.* 2014;**6**:9007-9014
- [9] Cameron E, Campbell A. The orthomolecular treatment of cancer. II. Clinical trial of high-dose ascorbic acid supplements in advanced human cancer. *Chem Biol Interact.* 1974;**9**:285-315
- [10] Cameron E, Pauling L. Supplemental ascorbate in the supportive treatment of cancer: Prolongation of survival times in terminal human cancer. *Proc Natl Acad Sci U S A.* 1976;**73**:3685-9
- [11] Cameron E, Pauling L. Supplemental ascorbate in the supportive treatment of cancer: reevaluation of prolongation of survival times in terminal human cancer. *Proc Natl Acad Sci U S A.* 1978;**75**:4538-42
- [12] Chen Q, Espey MG, Sun AY, Pooput C, Kirk KL, Krishna MC. Pharmacologic doses of ascorbate act as a prooxidant and decrease growth of aggressive tumor xenografts in mice. *Proceedings of the National Academy of Sciences of the United States of America.* 2008;**105**:11105-11109
- [13] Chen Q, Espey MG, Krishna MC, Mitchell JB, Corpe CP, Buettner GR. Pharmacologic ascorbic acid concentrations selectively kill cancer cells: Action as a pro-drug to deliver hydrogen peroxide to tissues. *Proceedings of the National Academy of Sciences of the United States of America.* 2005;**102**:13604-13609
- [14] Chen Q, Espey MG, Sun AY, Lee JH, Krishna MC, Shacter E. Ascorbate in pharmacologic concentrations selectively generates ascorbate radical and hydrogen peroxide in extracellular fluid in vivo. *Proceedings of the National Academy of Sciences of the United States of America.* 2007;**104**:8749-8754
- [15] Honarbakhsh S, Schachter M. Vitamins and cardiovascular disease. *British Journal of Nutrition.* 2008;**101**:1113-1131
- [16] Jacob RA, Sotoudeh G. Vitamin C function and status in chronic disease. *Nutrition in Clinical Care.* 2002;**5**:66-74
- [17] Willcox BJ, Curb JD, Rodriguez BL. Antioxidants in cardiovascular health and disease: Key lessons from epidemiologic studies. *American Journal of Cardiology.* 2008;**101**:75-86D
- [18] Pauling L. The significance of the evidence about ascorbic acid and the common cold. *Proceedings of the National Academy of Sciences of the United States of America.* 1971;**68**:2678-2681
- [19] Douglas RM, Hemilä H, Chalker E, Treacy B. Vitamin C for preventing and treating the common cold. *Cochrane Database Systematic Review.* 2007;**3**:CD000980

- [20] Johnston CS. The antihistamine action of ascorbic acid. *Subcellular Biochemistry*. 1996;**25**:189-213
- [21] Evans JR, Lawrenson JG. Antioxidant vitamin and mineral supplements for slowing the progression of age-related macular degeneration. *Cochrane Database Systematic Review*. 2012;11:CD000254
- [22] Age-Related Eye Disease Study Research Group. A randomized, placebo-controlled, clinical trial of high-dose supplementation with vitamins C and E, beta carotene, and zinc for age-related macular degeneration and vision loss: AREDS report no. 8. *Archives of Ophthalmology*. 2001;**119**:1417-1436
- [23] The Age-Related Eye Disease Study 2 (AREDS2) Research Group. Lutein + zeaxanthin and omega-3 fatty acids for age-related macular degeneration: The Age-Related Eye Disease Study 2 (AREDS2) randomized clinical trial. *Journal of American Medical Association*. 2013;**309**:2005-2015
- [24] Carr AC, Frei B. Toward a new recommended dietary allowance for vitamin C based on antioxidant and health effects in humans. *American Journal of Clinical Nutrition*. 1999;**69**:1086-1107
- [25] Rautiainen S, Lindblad BE, Morgenstern R, Wolk A. Vitamin C supplements and the risk of age-related cataract: A population-based prospective cohort study in women. *American Journal of Clinical Nutrition*. 2010;**91**(2):487-493
- [26] Darr D, Combs S, Dunston S, Manning T, Pinnell S. Topical vitamin C protects porcine skin from ultraviolet radiation-induced damage. *British Journal of Dermatology*. 1992;**127**:247-253
- [27] Steiling H, Longet K, Moodycliffe A. Sodium-dependent vitamin C transporter isoforms in skin: Distribution, kinetics, and effect of UVB-induced oxidative stress. *Radical Biology and Medicine*. 2007;**43**:752-762
- [28] Kang JS, Kim HN, Jung da J. Regulation of UVB-induced IL-8 and MCP-1 production in skin keratinocytes by increasing vitamin C uptake via the redistribution of SVCT-1 from the cytosol to the membrane. *Journal of Investigative Dermatology*. 2007;**127**:698-706
- [29] Tebbe B, Wu S, Geilen CC, Eberle J, Kodelja V, Orfanos CE. L-ascorbic acid inhibits UVA-induced lipid peroxidation and secretion of IL-1 α and IL-6 in cultured human keratinocytes in vitro. *Journal of Investigative Dermatology*. 1997;**108**:302-306
- [30] Eberlein-Konig B, Placzek M, Przybilla B. Protective effect against sunburn of combined systemic ascorbic acid (vitamin C) and d-alpha-tocopherol (vitamin E). *Journal of the American Academy of Dermatology*. 1998;**38**:45-48
- [31] Bissett DL, Chatterjee R, Hannon DP. Photoprotective effect of superoxide-scavenging antioxidants against ultraviolet radiation-induced chronic skin damage in the hairless mouse. *Photodermatology, Photoimmunology & Photomedicine*. 1990;**7**:56-62
- [32] Lin FH, Lin JY, Gupta RD. Ferulic acid stabilizes a solution of vitamins C and E and doubles its photoprotection of skin. *Journal of Investigative Dermatology*. 2005;**125**:826-832

- [33] Lin JY, Selim MA, Shea CR. UV photoprotection by combination topical antioxidants vitamin C and vitamin E. *Journal of the American Academy of Dermatology*. 2003;**48**:866-874
- [34] Savini I, Catani MV, Rossi A, Duranti G, Melino G, Avigliano L. Characterization of keratinocyte differentiation induced by ascorbic acid: Protein kinase C involvement and vitamin C homeostasis. *Journal of Investigative Dermatology*. 2002;**118**:372-379
- [35] Boyce ST, Supp AP, Swope VB, Warden GD. Vitamin C regulates keratinocyte viability, epidermal barrier, and basement membrane in vitro, and reduces wound contraction after grafting of cultured skin substitutes. *Journal of Investigative Dermatology*. 2002;**118**:565-572
- [36] Ponc M, Weerheim A, Kempenaar J. The formation of competent barrier lipids in reconstructed human epidermis requires the presence of vitamin C. *Journal of Investigative Dermatology*. 1997;**109**:348-355
- [37] Silverstein RJ, Landsman AS. The effects of a moderate and high dose of vitamin C on wound healing in a controlled guinea pig model. *Journal of Foot and Ankle Surgery*. 1999;**38**:333-338
- [38] Vaxman F, Olender S, Lambert A. Effect of pantothenic acid and ascorbic acid supplementation on human skin wound healing process. A double-blind, prospective and randomized trial. *Journal of Surgical Research*. 1995;**27**:158-166
- [39] Verma KK, Jain A, Sahasrabuddhey B, Gupta K, Mishra S. Solid-phase extraction cleanup for determining ascorbic acid and dehydroascorbic acid by titration with 2,6-dichlorophenolindophenol. *Journal of AOAC International*. 1996;**79**:1236
- [40] Xia W, Yuxia D, Changxia S, Jinghe Y, Yuebo W, Shuna S. Fluorimetric determination of ascorbic acid with o-phenylenediamine. *Talanta*. 2003;**59**:95-99
- [41] Rahman Khan MM, Rahman MM, Islam MS, Begum SA. A simple UV-spectrophotometric method for the determination of vitamin C content in various fruits and vegetables at Sylhet area in Bangladesh. *Journal of Biological Sciences*. 2006;**6**:388-392
- [42] Nyssönen K, Salonen JT, Parviainen MT. Ascorbic acid. In: De Leenheer AP, Lambert WE, Van Bocxlaer JF, editors. *Modern Chromatographic Analysis of Vitamins*. 3rd ed. Marcel Dekker: New York; 2000
- [43] Casella L, Gullotti M, Marchesini M, Petrarulo M. Rapid enzymatic method for vitamin C assay in fruits and vegetables using peroxidase. *Journal of Food Science*. 2006;**54**:374
- [44] Ensafi AA, Rezaei B, Movahedinia H. Kinetic-spectrophotometric determination of ascorbic acid by inhibition of the hydrochloric acid-bromate reaction. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*. 2002;**58**:2589-2594
- [45] Safavi A, Fotouhi L. Kinetic spectrophotometric determination of ascorbic acid by reduction of toluidine blue. *Talanta*. 1994;**41**:1225-1228

- [46] Nováková L, Solich P, Solichová D. HPLC methods for simultaneous determination of ascorbic and dehydroascorbic acids. *Trends in Analytical Chemistry*. 2008;**27**:942
- [47] Racz E, Parlagh-Huszar K, Kecskes T. HPLC method for determination of ascorbic acid in fruits and vegetables. *Periodica Polytechnica Chemical Engineering*. 1991;**35**(1-2):23-30
- [48] Snezana SM, Danijela AK, Danijela CN, Milan NM. Rapid and reliable HPLC method for the determination of vitamin C in pharmaceutical samples. *Tropical Journal of Pharmaceutical Research*. 2011;**10**(1):105-111
- [49] Franco VV, María EP, María SC, Daniel RG, del H. Bernardi CM. Optimization and validation of a UV-HPLC method for vitamin C determination in strawberries (*Fragaria ananassa* Duch.), using experimental designs. *Food Analytical Methods*. 2012;**5**:1097-1104
- [50] Line R, Hoffer LJ. A simple method for plasma total vitamin C analysis suitable for routine clinical laboratory use. *Nutrition Journal*. 2016;**15**:40
- [51] Güçlü K, Sözgen K, Tütem E, Özyürek M, Apak R. Spectrophotometric determination of ascorbic acid using copper(II)-neocuproine reagent in beverages and pharmaceuticals. *Talanta*. 2005;**65**(5):1226-1232
- [52] Danet AF, David V, Oancea M. Dispozitiv de analiza in flux cu injectarehidrodinamica. Determinareavitaminei C, *Revista de Chimie*. 1994;**45**(11):1000-1006
- [53] Danet BM, Aboul-Enein HY. Flow injection system with chemiluminometric detection for enzymatic determination of ascorbic acid. *Luminescence*. 2000;**15**:305-309
- [54] Kobra Z, Somayye M. Sensitive spectrophotometric determination of ascorbic acid in drugs and foods using surface plasmon resonance band of silver nanoparticles. *Cogent Chemistry*. 2015;**1**:1109172
- [55] Kapur A, Hasković A, Čopra-Janićijević A, Klepo L, Topčagić A, Tahirović I, Sofić E. Spectrophotometric analysis of total ascorbic acid content in various fruits and vegetables. *Bulletin of the Chemists and Technologists of Bosnia and Herzegovina*. 2012;**38**:39-42
- [56] Mohammed IHM, Hazim YA. Determination of Vitamin C (ascorbic acid) Contents in various fruit and vegetable by UV-spectrophotometry and titration methods. *Journal of Chemical and Pharmaceutical Sciences*. 2016;**9**:2972
- [57] Ahmed Z, Mohamed H. Spectrophotometric determination of ascorbic acid in aqueous solutions and in pharmaceuticals formulations. *Journal of Al-Nahrain University*. 2013;**16**:65-71
- [58] Cooper JA, Woodhouse KE, Chippindale AM, Compton RG. Photoelectrochemical determination of ascorbic acid using methylene blue immobilized in α -zirconium phosphate. *Electroanalysis*. 1999;**11**:1259-1265
- [59] Morais H, Rodrigues P, Ramos C, Forges E, Cserhti T, Oliveira J. Effect of ascorbic acid on the stability of β -carotene and capsanthin in paprika (*Capsicum annuum*) powder. *Nahrung/Food*. 2002;**46**:308-310

- [60] Julio CBF, Lauro TK, Graciliano DON. Potentiometric sensor for L-ascorbic acid based on EVA membrane doped with copper (II). *Electroanalysis*. 1999;**11**:475-480
- [61] Song Z, Qun X, Wen Z, Litong J, Ji YJ. In vivo monitoring of the monoamine neurotransmitters in rat brain using microdialysis sampling with liquid chromatography electrochemical detection. *Analytica Chimica Acta*. 2001;**427**:45-53
- [62] Joseph W, Alain W. Zeolite-modified carbon paste electrode for selective monitoring of dopamine. *Journal of Electroanalytical Chemistry*. 1996;**407**:183-187
- [63] Adriana DD, Jairo JP, Ivano GRG. A batch injection analysis system for ascorbic acid determination using amperometric detection on a sessile mercury drop electrode. *Electroanalysis*. 1999;**11**:1124-1129
- [64] Andrei DF, Mihaela B, Hassan YAE. Flow injection system with chemiluminometric detection for enzymatic determination of ascorbic acid. *Lumin*. 2000;**15**:305-309
- [65] Jahan BR, Reza O, Abolfazl K. Carbon paste electrode spiked with ferrocene carboxylic acid and its application to the electrocatalytic determination of ascorbic acid. *Journal of Electroanalytical Chemistry*. 2001;**515**:45-51
- [66] Jyh MZ, Dong MT, Hsueh HY. Direct determination of vitamin C in fruit juices using a polyviologen-modified electrode. *Electroanalysis*. 2002;**14**:1597-1600
- [67] Pournaghi AMH, Razmi NH, Hafezi B. Amperometric determination of ascorbic acid in real samples using an aluminum electrode, modified with Nickel Hexacyanoferrate Films by simple electroless dipping method. *Electroanalysis*. 2002;**14**:206-219
- [68] Suw YL, Jung IC, Young SJ, Woon WJ, Hye JL, Seong HL. Electrochemical detection of ascorbic acid (vitamin C) using a glassy carbon electrode. *Nahrung/Food*. 2004;**3**:201-204
- [69] Sudha JD, Neethu KS, Renjith S, Hans-Juergen PA, Pich A. Water dispersible electrically conductive poly (3,4-ethylenedioxythiophene) nanospindles by liquid crystalline template assisted polymerization. *Journal of Materials Chemistry C*. 2014;**2**:6991
- [70] Reena VL, Sudha JD, Rohini KN, Neethu KS. Design of a nanostructured electromagnetic polyaniline-keggin iron-clay composite modified electrochemical sensor for the nanomolar detection of ascorbic acid. *Journal of Applied Polymer Science*. 2014;**131**:40936-4094