We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Organic Reactions Promoted by Metal-Free Organic Dyes Under Visible Light Irradiation

Hideto Miyabe

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.70507>

Abstract

Although dyes have received much attention as the visible light-activated photocatalysts, the use of metal-free organic dyes in synthetic organic chemistry is still limited. This chapter summarizes the recent progress in the visible light photocatalysis promoted by metal-free organic dyes. Eosin Y is the typical organic dyes to induce the photoredox catalysis. Recently, other organic dyes such as Rose Bengal, fluorescein, and methylene blue have been studied as photocatalysts to promote the single-electron transfer processes.

Keywords: photocatalyst, organocatalyst, dye, catalysis, visible light, radical

1. Introduction

The use of abundant sunlight as a clean source of energy is an important aim of green chemistry. In recent years, dyes have attracted a great deal of attention as the visible light-activated photocatalysts in synthetic organic chemistry. However, these studies have mainly concentrated on the redox transformations using transition metal dyes such as ruthenium or iridium photocatalysts [1–9]. In contrast, the use of metal-free dyes still remains rather underdeveloped, although organic dyes are more environmentally friendly and cheaper. Eosin Y is the typical organic dyes to induce the photoredox catalysis [10]. Recently, Rose Bengal, fluorescein, methylene blue, and other organic dyes have been studied as photocatalysts to promote the single-electron transfer processes [11–13]. Additionally, 3-cyano-1-methylquinolinium, 9-mesityl-10-methylacridinium ion, and acridinium salts were developed as organic photocatalysts [12, 13].

The photoredox cycle is initiated by the visible light irradiation of dye in the ground state to produce the high-energy excited state of dye (Dye*) (**Figure 1**). Two distinctive pathways

Figure 1. Dye-catalyzed photoredox cycle.

from dye in the excited state (Dye^*) are described for the mechanism of visible light photoredox catalysis. The reductive property of Dye^* can be used in the presence of a sacrificial electron acceptor. In other words, Dye^* serves as an electron donor leading the radical cation species of Dye. In contrast, Dye^* also acts as an electron acceptor in the presence of a sacrificial electron donor.

2. Eosin Y and eosin B

Eosin Y (EY) is the typical organic dye to induce the synthetically useful photoredox transformations [14–30]. EY that absorbed visible light populates in the lowest excited singlet state. The subsequent spin-forbidden singlet-triplet intersystem crossing affords EY in the excited triplet state. A variety of photoredox transformations are induced by the single-electron transfer from EY in the excited triplet state. In 2014, König provides the review article concerning the utility of EY as a photocatalyst in synthetic organic chemistry [10]. Therefore, this section highlights the recent remarkable progress in the EY-catalyzed photoredox transformations.

The EY-catalyzed generation of aryl radicals from aryl diazonium salts was studied by König's group [14]. The direct C–H bond arylation of heteroarenes with aryl diazonium salts was achieved by employing only 1 mol% of EY (**Figure 2**). The arylation of furan with diazonium salt **1** in dimethyl sulfoxide (DMSO) proceeded smoothly under the visible light irradiation to give the desired coupling product **2** in 85% yield. This transformation proceeds through the radical mechanism. Initially, aryl radical is produced by the single-electron transfer from the excited EY (EY^*) to aryl diazonium salt **1**. The addition of aryl radical to furan leads to the formation of the radical intermediate **A**, which is further oxidized to cation intermediate **B**. Final deprotonation gives the coupling product **2**. Next, the EY-catalyzed arylation of simple arenes with fluorinated aryl bromides was developed [15]. In the presence of EY (5 mol%) and triethylamine as an electron donor, the direct arylation using 1-bromo-2,3,4,5,6-pentafluorobenzene **3** and benzene gave the coupling product **4** in 85% yield. The mechanistic investigations reveal that the photooxidation of

Figure 2. EY-catalyzed generation of aryl radicals and coupling reactions.

triethylamine by the excited EY (EY*), and the subsequent single-electron transfer from the radical anion species of EY to **3** leads to the formation of the polyfluorinated aryl radical. The mild visible light-mediated generation of aryl radicals from diazonium salts was also investigated by Wangelin's group [16–18]. The coupling reaction catalyzed by EY was investigated with no use of any sacrificial oxidants [19, 20].

The vinyl sulfones were synthesized by the EY-catalyzed reaction of alkenes with sodium aryl sulfinates [21]. The reaction of 1,2-dihydronaphthalene **6** with sodium benzenesulfinate **5** was performed in the presence of EY (10 mol%) and nitrobenzene as a terminal oxidant (**Figure 3**).

Figure 3. EY-catalyzed reaction using sodium benzenesulfinate.

The desired vinyl sulfone **7** was obtained in 99% yield. In this reaction, sodium sulfinate **5** is oxidized by the excited EY (EY^*) to give the sulfonyl radical, which attacks the double bond of **6** to form the radical intermediate **C**. Nitrobenzene oxidizes the radical cation species of EY to give EY in the ground state and the radical anion species **D** of nitrobenzene, which reacts with radical intermediate **C** to give the vinyl sulfone **7**.

The oxidative cyclization reaction between 3-phenylpropiolate **8** and 4-methylbenzenesulfinic acid **9** was studied (Figure 4) [22]. In the presence of EY (1 mol%) and *tert*-butyl hydroperoxide (TBHP), the reaction between **8** and **9** was performed in MeCN– H_2O (1:1, v/v) under the

Figure 4. EY-catalyzed reaction using 4-methylbenzenesulfinic acid.

visible light irradiation. The desired coumarin **10** was obtained in 78% isolated yield. Initially, *tert*-butoxyl radical is produced by the single-electron transfer from the excited EY (EY*) to TBHP. The cyclization reaction is promoted by the addition of sulfonyl radical, generated from sulfinic acid **9** and *tert*-butoxyl radical, to alkyne moiety of **8**. The coumarin **10** is formed *via* the oxidation of the cyclized radical intermediate **E** by the radical cation species of EY.

The EY-catalyzed cyclization of 2-isocyanobiphenyls with arylsulfonyl chlorides took place under the oxidant-free visible light irradiation conditions [23]. In the presence of K₂HPO₄ as a base, the EY-catalyzed reaction of benzenesulfonyl chloride **11** and 2-isocyanobiphenyl **12** proceeded smoothly to give the 6-phenyl-substituted phenanthridine **13** in 79% yield (Figure 5). Initially, the single-electron transfer from the excited EY (EY*) to sulfonyl chloride **11** gives the phenyl radical, which adds to isocyanide **12** to form the imidoyl radical intermediate **F**. The subsequent cyclization gives the cyclized radical intermediate **G**, which is oxidized by the radical cation species of EY. Finally, the deprotonation leads to **13**.

EY could be used as the photocatalyst for the 5-*exo*-trig cyclization of iminyl radicals generated from *O*-aryl oximes [24]. Among several aryl oximes evaluated, 2,4-dinitro-substituted aryl oxime **14** has the excellent reactivity due to its low reduction potential (Figure 6). In the presence of cyclohexadiene (CHD) as a H-donor, EY-catalyzed photoreaction of **14** gave the cyclized product **15** in 78% yield. In this transformation, the iminyl radical **I** is generated *via* the reduction of **14** by the excited EY (EY*) followed by the fragmentation of radical anion **H**. The cyclization of **I** gives the cyclized C-centered radical **J**, which abstracts H-atom from CHD to give the desired product **15**. Furthermore, the formation of product **15** was observed even in the absence of EY, when the MeCN solution of **14** was treated with Et₃N under the visible

Figure 5. EY-catalyzed cyclization of 2-isocyanobiphenyl.

Figure 6. EY-catalyzed cyclization of 2,4-dinitro-substituted aryl oxime.

light irradiation. In this case, the visible light-mediated electron transfer would be induced by the formation of donor-acceptor complex between Et_3N and 2,4-dinitrophenyl group of **14**.

The EY-induced photocatalysis was applied to the radical cascade cyclization of polyenes [25]. The photocatalytic cascade cyclization of polyene **16** proceeded by employing EY (**Figure 7**). Hexafluoro-2-propanol (HFIP) was identified as the optimal solvent. The cyclized product **17** was obtained in 93% yield with the excellent diastereoselectivity *via* the radical cation intermediate **K** generated by the single-electron transfer from **16** to the excited EY (EY*). In this process, the OH moiety of **16** would act as a terminator.

Eosin B is also the active catalyst under the visible light irradiation [31]. The C–H functionalization of thiazole derivatives with diarylphosphine oxides was achieved by the eosin B-catalyzed photoredox process. When eosin B was employed as a photocatalyst, the phosphorylation of benzothiazole **18** with diphenylphosphine oxide proceeded effectively to give the phosphorylation product **19** in 87% yield (**Figure 8**). In this transformation, hydrogen (H_2) is the only by-product.

Figure 7. EY-catalyzed cascade cyclization of polyene.

Figure 8. Eosin B-catalyzed phosphorylation of benzothiazole.

3. Rose Bengal

Rose Bengal (RB) was widely used as a visible light-activated photocatalyst [32–44]. Tan's group studied the photoredox catalysis using RB [32–36]. RB was a good catalyst for the dehydrogenative coupling reaction between tetrahydroisoquinolines and nitroalkanes (**Figure 9**) [33]. In the presence of RB (5 mol%), the reaction of *N*-phenyl-tetrahydroisoquinoline **20** with nitromethane gave the adduct **21** in 92% yield. In the absence of O_2 , a much lower yield was obtained relative to the reaction performed in open air; thus, air is important for this reaction as an oxidant. Additionally, this reaction was expanded to the dehydrogenative Mannich reaction using enamine nucleophiles generated from ketones and pyrrolidine.

Figure 9. RB-catalyzed reaction of tetrahydroisoquinoline.

Next, the combination of graphene oxide and RB was studied in the reaction between tetrahydroisoquinolines with TMSCN or TMSCF₃ [34]. In the presence of RB (5 mol%) and graphene oxide (50 wt%), the reaction of **20** with TMSCN proceeded effectively to give the adduct **22** in 99% yield, while the yield of **22** decreased to 45% in the absence of graphene oxide. The use of graphene oxide as a cocatalyst improves the reaction rates and yields.

New method for the synthesis of Meyers's bicyclic lactams was developed by using RB photocatalysis [37–39]. This cascade transformation is the one-pot reaction which begins from furan substrates (**Figure 10**) [37]. Despite the extraordinary complexity of reaction cascade, the reaction between 2-methylfuran **23** and L-serine ethyl ester **24** led to the formation of bicyclic lactam **25** in 68% yield. At first, RB promotes the photooxidation of methylfuran **23** with singlet oxygen in MeOH. The intermediate **L** is formed by the in situ reduction of hydroperoxy with Me₂S. The next reaction of **L** with L-serine ethyl ester **24** gives the intermediate **M**, which is converted to 2-pyrrolidinone **N** *via* imino enal. Actually, 2-pyrrolidinone **N** could be isolated by the flash column chromatography using silica gel neutralized by trimethylamine. Finally, treatment of **N** with TFA gives the bicyclic lactam **25** as the final product of one-pot reaction cascade.

The aerobic visible light-promoted indole C3 formylation reaction was achieved by using RB as a photocatalyst and *N,N,N',N'*-tetramethylethylenediamine (TMEDA) as a one-carbon source (**Figure 11**) [40]. Upon the irradiation of visible light, the reaction of *N*-methylindole **26** with TMEDA in the presence of RB (5 mol%) and KI as an additive under air afforded 3-formyl-*N*-methylindole **27** in 70% yield. This transformation proceeds *via* the addition of *N*-methylindole **26** to iminium ion **O** generated by the oxidation of TMEDA. Next, the C3 thiocyanation reaction of indoles was developed by using ammonium thiocyanate (NH₄SCN) as a thiocyanate radical source [41]. In the presence of RB (1 mol%), the reaction of indole **28** with NH₄SCN gave the adduct **29** in 98% yield. In this reaction, thiocyanate radical is generated by the single-electron transfer between thiocyanate anion and the excited RB (RB*). The thiocyanate radical adds to indole **28**. The subsequent oxidation leads to **29**.

Figure 10. RB-catalyzed synthesis of bicyclic lactam.

Figure 11. RB-catalyzed functionalization of C3 in indoles.

Figure 12. RB-catalyzed oxidative cyclization of benzimidamide.

The new method for the synthesis of quinazolines was developed by using RB as a photocatalyst (**Figure 12**) [42]. In the presence of RB (1 mol%), CBr_4 as an oxidant, and K_2CO_3 as a base, the oxidative carbon-carbon bond-forming cyclization of *N*-benzyl-*N'*-phenyl benzimidamide **30** proceeded smoothly under the visible light irradiation. The desired quinazoline **31** was obtained in 88% yield. In this transformation, CBr_3 radical is generated by the reaction between CBr_4 and the radical anion species of RB. Next, the iminium ion intermediate **Q** is formed from the radical cation intermediate **P** by the association of CBr_3 radical. Finally, the intramolecular Friedel-Craft reaction of iminium ion **Q** leads to quinazoline **31**.

4. Fluorescein and rhodamine B

The utility of fluorescein was demonstrated in the alkoxy-carboxylation of aryldiazonium salts using CO gas [45]. In the presence of 0.5 mol% of fluorescein as a photocatalyst, treatment of diazonium tetrafluoroborate **32** with CO (80 atm pressure) in methanol under the irradiation of visible light gave methyl ester **33** in 80% yield (**Figure 13**). Initially, the phenyl radical is generated from diazonium **32** by the single-electron transfer from the excited state of fluorescein (Dye^*). Next, benzoyl radical **R** is formed *via* trapping of CO molecule by phenyl radical. The methyl ester **33** is obtained *via* the oxidation of benzoyl radical **R** by the reactive radical cation species of dye followed by trapping of the resulting benzylidyneoxonium **S** with methanol. Furthermore, the dual catalytic system using photocatalyst and gold catalyst was studied by Glorius's group [46].

Figure 13. Fluorescein-catalyzed alkoxy-carboxylation of aryldiazonium.

The utility of rhodamine B as a water-soluble photocatalyst was demonstrated in the aqueous-medium carbon-carbon bond-forming radical reactions [47]. In the presence of $(i\text{-Pr})_2\text{NEt}$ as a reductive quencher, the rhodamine B-catalyzed reaction of alkene **34** with $i\text{-C}_3\text{F}_7\text{I}$ in H_2O proceeded smoothly to give the product **35** in 90% yield (**Figure 14**). In this transformation, the photo-induced electron transfer from the excited singlet state (S_1) of rhodamine B to $i\text{-C}_3\text{F}_7\text{I}$ was proposed. This electron transfer process was supported by the fluorescence quenching of rhodamine B with addition of $i\text{-C}_3\text{F}_7\text{I}$. Additionally, the aqueous-medium radical addition-cyclization-trapping reaction of **36** proceeded effectively even in the absence of $(i\text{-Pr})_2\text{NEt}$. In this transformation, the single-electron transfer from iodine ion (I^-) to the radical cation species of rhodamine B in an ion pair would proceed to give $\text{I}_2^\bullet$, because the oxidation potential of rhodamine B is positive enough to oxidize I^- into $\text{I}_2^\bullet$.

Figure 14. Rhodamine B-catalyzed aqueous-medium radical reactions.

5. Methylene blue and acridine red

Methylene blue (MB) is a member of the thiazine dye family. Scaiano's group used MB as a photocatalyst under the visible light irradiation [48]. The radical trifluoromethylation of electron-rich heterocycles was studied by the use of Togni's reagent **39** as a CF_3 radical source (Figure 15). The trifluoromethylation of 3-methylindole **38** proceeded with good yield at low catalyst concentration, when *N,N,N',N'*-tetramethylethylenediamine (TMEDA) was used as an electron donor. In the presence of 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) as an electron donor, the reaction of terminal alkene **41** with Togni's reagent **39** also gave the hydrotrifluoromethylation product **42** in 67% yield as a major product, because the fully reduced form of MB, leuco-MB, acts as a hydrogen source [49]. The possible mechanism for the catalytic formation of CF_3 radical is shown. The visible light-excited MB (MB^*) is readily quenched by aliphatic amines such as TMEDA or DBU to form the semi-reduced MB as a radical anion and an α -amino radical. CF_3 radical is generated *via* the reduction of Togni's reagent **39** with semi-reduced MB and/or an α -amino radical.

The one-pot transformation of furans into 5-hydroxy-1*H*-pyrrol-2(5*H*)-ones was investigated by using MB as a photocatalyst [50]. In the presence of MB (2 mol%) and oxygen, the reaction of furan **43** with benzylamine gave lactam **44** in 72% yield *via* the reduction of the intermediate by

Figure 15. MB-catalyzed trifluoromethylation using Togni's reagent.

Me₂S (**Figure 16**). The use of 2-(3,4-dimethoxyphenyl)ethanamine **45** instead of benzylamine led to the formation of tricycle **46** *via* Pictet-Spengler cyclization process. For Pictet-Spengler cyclization, HCOOH was added as an acid leading to *N*-acyliminium ion, which spontaneously cyclized to form **46**.

New phenothiazine-based organic dye was also developed as a visible light-activated photocatalyst [51].

Acridine red was used as a photocatalyst for the visible light-induced direct thiolation of ethers (**Figure 17**) [52]. The thiolation of tetrahydrofuran (THF) using diphenyl disulfide **47** was carried out in the presence of acridine red (2 mol%) and *tert*-butyl hydroperoxide (TBHP) as an oxidant. The reaction occurred at ambient conditions to give α -arylthioether **48** in 82% yield. This transformation proceeds *via* the generation of radical intermediate from THF, which reacts with diphenyl disulfide **47** to afford 2-(phenylthio)-tetrahydrofuran **48**.

Figure 16. MB-catalyzed reactions of furan with amines.

Figure 17. Acridine red-catalyzed thiolation of tetrahydrofuran.

6. Riboflavin tetraacetate

Riboflavin tetraacetate (RFT) is an effective photocatalyst for the visible light-driven organic reactions. The aerobic oxidation of alkyl benzenes to ketones and carboxylic acids was investigated through a dual catalysis using RFT and the tris(2-pyridylmethyl)amine-iron complex [Fe(TPA)(MeCN)₂](ClO₄)₂ (TPA=tris(2-pyridylmethyl)amine) [53]. When a mixture of RFT (10 mol%) and [Fe(TPA)(MeCN)₂](ClO₄)₂ (2 mol%) was employed, the oxidation of 4-ethylanisole

49 proceeded effectively under the visible light irradiation to give 4-acetylanisole **50** in 80% yield (**Figure 18**). In this oxidation, the iron complex acts as a catalyst for not only oxidation of **49** but also disproportionation of hydrogen peroxide H_2O_2 which is obtained in the RFT-catalyzed oxidation of **49**.

RFT also catalyzed the aerobic oxidation of sulfides to sulfoxides without overoxidation to sulfones [54]. In the presence of RFT (2 mol%), sulfide **51** was transformed chemoselectively to the corresponding sulfoxide **52** in 91% yield (**Figure 19**).

Figure 18. RFT-catalyzed oxidation of 4-ethylanisole.

Figure 19. RFT-catalyzed oxidation of sulfide.

7. Concluding remarks

Organic dyes that absorbed visible light induce the synthetically valuable photochemical transformations. The metal-free photocatalysis using organic dyes rapidly progresses in the last few years. In addition to the organic dyes shown in this chapter, Fukuzumi' group has developed 3-cyano-1-methylquinolinium and 9-mesityl-10-methylacridinium ions as photocatalysts [13]. More recently, Nicewicz' group has studied the photocatalysis using acridinium salts [12]. These visible light-induced catalysis disclosed a broader aspect of the utility of organic photocatalysts for synthetic organic chemistry. This chapter will inspire creative new contributions to organic chemists.

Author details

Hideto Miyabe

Address all correspondence to: miyabe@huhs.ac.jp

School of Pharmacy, Hyogo University of Health Sciences, Minatojima, Chuo-ku, Japan

References

- [1] Zeitler K. Photoredox catalysis with visible light. *Angewandte Chemie International Edition*. 2009;**48**:9785-9786
- [2] Narayanam JMR, Stephenson CRJ. Visible light photoredox catalysis: Applications in organic synthesis. *Chemical Society Reviews*. 2011;**40**:102-113
- [3] Tucker JW, Stephenson CRJ. Shining light on photoredox catalysis: Theory and synthetic applications. *Journal of Organic Chemistry*. 2012;**77**:1617-1622
- [4] Yoon TP. Visible light photocatalysis: The development of photocatalytic radical ion cycloadditions. *ACS Catalysis*. 2013;**3**:895-902
- [5] Angnes RA, Li Z, Correia CRD, Hammond GB. Recent synthetic Additions to the visible light photoredox catalysis toolbox. *Organic and Biomolecular Chemistry*. 2015;**13**:9152-9167
- [6] Meggers E. Asymmetric catalysis activated by visible light. *Chemical Communications*. 2015;**51**:3290-3301
- [7] Xuan J, Zhang ZG, Xiao WJ. Visible-light-induced decarboxylative functionalization of carboxylic acids and their derivatives. *Angewandte Chemie International Edition*. 2015;**54**:15632-15641
- [8] Amador AG, Yoon TP. A chiral metal photocatalyst architecture for highly enantioselective photoreactions. *Angewandte Chemie International Edition*. 2016;**55**:2304-2306
- [9] Gurry M, Aldabbagh F. A new era for homolytic aromatic substitution: Replacing Bu_3SnH with efficient light-induced chain reactions. *Organic and Biomolecular Chemistry*. 2016;**14**:3849-3862
- [10] Hari DP, König B. Synthetic applications of eosin Y in photoredox catalysis. *Chemical Communications*. 2014;**50**:6688-6699
- [11] Ravelli D, Fagnoni M, Albini A. Photoorganocatalysis. What for? *Chemical Society Reviews*. 2013;**42**:97-113
- [12] Nicewicz DA, Nguyen TM. Recent applications of organic dyes as photoredox catalysts in organic synthesis. *ACS Catalysis*. 2014;**4**:355-360

- [13] Fukuzumi S, Ohkubo K. Organic synthetic transformations using organic dyes as photoredox catalysts. *Organic and Biomolecular Chemistry*. 2014;**12**:6059-6071
- [14] Hari DP, Schroll P, König B. Metal-free, visible-light-mediated direct C-H arylation of heteroarenes with aryl diazonium salts. *Journal of the American Chemical Society*. 2012;**134**:2958-2961
- [15] Meyer AU, Slanina T, Yao CJ, König B. Metal-free perfluoroarylation by visible light photoredox catalysis. *ACS Catalysis*. 2016;**6**:369-375
- [16] Majek M, Jacobi von Wangelin A. Organocatalytic visible light mediated synthesis of aryl sulfides. *Chemical Communications*. 2013;**49**:5507-5509
- [17] Majek M, Jacobi von Wangelin A. Metal-free carbonylations by photoredox catalysis. *Angewandte Chemie International Edition*. 2015;**54**:2270-2274
- [18] Majek M, Filace F, Jacobi von Wangelin A. Visible light driven hydro-/deuterodefunctionalization of anilines. *Chemistry A European Journal*. 2015;**21**:4518-4522
- [19] Meng QY, Zhong JJ, Liu Q, Gao XW, Zhang HH, Lei T, Li ZJ, Feng K, Chen B, Tung CH, Wu LZ. A cascade cross-coupling hydrogen evolution reaction by visible light catalysis. *Journal of the American Chemical Society*. 2013;**135**:19052-19055
- [20] Zhong JJ, Wu CJ, Meng QY, Gao XW, Lei T, Tung CH, Wu LZ. A cascade cross-coupling and *in situ* hydrogenation reaction by visible light catalysis. *Advanced Synthesis & Catalysis*. 2014;**356**:2846-2852
- [21] Meyer AU, Jäger S, Hari DP, König B. Visible light-mediated metal-free synthesis of vinyl sulfones from aryl sulfinates. *Advanced Synthesis & Catalysis*. 2015;**357**:2050-2054
- [22] Yang W, Yang S, Li P, Wang L. Visible-light initiated oxidative cyclization of phenyl propiolates with sulfinic acids to coumarin derivatives under metal-free conditions. *Chemical Communications*. 2015;**51**:7520-7523
- [23] Gu L, Jin C, Liu J, Ding H, Fan B. Transition-metal-free, visible-light induced cyclization of arylsulfonyl chlorides with 2-isocyanobiphenyls to produce phenanthridines. *Chemical Communications*. 2014;**50**:4643-4645
- [24] Davies J, Booth SG, Essafi S, Dryfe RAW, Leonori D. Visible-light-mediated generation of nitrogen-centered radicals: Metal-free hydroimination and iminohydroxylation cyclization reactions. *Angewandte Chemie International Edition*. 2015;**54**:14017-14021
- [25] Yang Z, Li H, Zhang L, Zhang MT, Cheng JP, Luo S. Organic photocatalytic cyclization of polyenes: A visible-light-mediated radical cascade approach. *Chemistry A European Journal*. 2015;**21**:14723-14727
- [26] Jadhav SD, Bakshi D, Singh A. Visible light mediated organocatalytic activation of ethyl bromofluoroacetate: Coupling with indoles and anilines. *Journal of Organic Chemistry*. 2015;**80**:10187-10196
- [27] Zhang J, Wang L, Liu Q, Yang Z, Huang Y. Synthesis of α,β -unsaturated carbonyl compounds *via* a visible-light-promoted organocatalytic aerobic oxidation. *Chemical Communications*. 2013;**19**:11662-11664

- [28] Zhou C, Li P, Zhu X, Wang L. Merging photoredox with palladium catalysis: Decarboxylative *ortho*-acylation of acetanilides with α -oxocarboxylic acids under mild reaction conditions. *Organic Letters*. 2015;**17**:6198-6201
- [29] Li J, Zhang J, Tan H, Wang DZ. Visible-light-promoted vinylation of tetrahydrofuran with alkynes through direct C-H bond functionalization. *Organic Letters*. 2015;**17**:2522-2525
- [30] Mitra S, Ghosh M, Mishra S, Hajra A. Metal-free thiocyanation of imidazoheterocycles through visible light photoredox catalysis. *Journal of Organic Chemistry*. 2015;**80**:8275-8281
- [31] Luo K, Chen YZ, Yang WC, Zhu J, Wu L. Cross-coupling hydrogen evolution by visible light photocatalysis toward C(sp²)-P formation: Metal-free C-H functionalization of thiazole derivatives with diarylphosphine oxides. *Organic Letters*. 2016;**18**:452-455
- [32] Liu H, Feng W, Kee CW, Zhao Y, Leow D, Pan Y, Tan CH. Organic dye photocatalyzed α -oxyamination through irradiation with visible light. *Green Chemistry*. 2010;**12**:953-956
- [33] Pan Y, Kee CW, Chen L, Tan CH. Dehydrogenative coupling reactions catalysed by rose Bengal using visible light irradiation. *Green Chemistry*. 2011;**13**:2682-2685
- [34] Pan Y, Wang S, Kee CW, Dubuisson E, Yang Y, Loh KP, Tan CH. Graphene oxide and rose Bengal: Oxidative C-H functionalization of tertiary amines using visible light. *Green Chemistry*. 2011;**13**:3341-3344
- [35] Liu H, Feng W, Kee CW, Zhao Y, Leow D, Pan Y, Tan CH. Organic dye photocatalyzed α -oxyamination through irradiation with visible light. *Green Chemistry*. 2012;**14**:953-956
- [36] Teo YC, Pan Y, Tan CH. Organic dye-photocatalyzed acylnitroso ene reaction. *Chem CatChem*. 2013;**5**:235-240
- [37] Kalaitzakis D, Montagnon T, Alexopoulou I, Vassilikogiannakis G. A versatile synthesis of Meyers' bicyclic lactams from furans: Singlet-oxygen-initiated reaction cascade. *Angewandte Chemie International Edition*. 2012;**51**:8868-8871
- [38] Kalaitzakis D, Antonatou E, Vassilikogiannakis G. One-pot synthesis of 1-azaspiro frameworks initiated by photooxidation of simple furans. *Chemical Communications*. 2014;**50**:400-402
- [39] Triantafyllakis M, Tofi M, Montagnon T, Kouridaki A, Vassilikogiannakis G. Singlet oxygen-mediated synthesis of bis-spiroketals found in azaspiracids. *Organic Letters*. 2014;**16**:3150-3153
- [40] Li X, Gu X, Li Y, Li P. Aerobic transition-metal-free visible-light photoredox indole C-3 formylation reaction. *ACS Catalysis*. 2014;**4**:1897-1900
- [41] Fan W, Yang Q, Xu F, Li P. A visible-light-promoted aerobic metal-free C-3 thiocyanation of indoles. *Journal of Organic Chemistry*. 2014;**79**:10588-10592
- [42] Shen Z, Yang P, Tang Y. Transition metal-free visible light-driven photoredox oxidative annulation of arylamidines. *Journal of Organic Chemistry*. 2016;**81**:309-317

- [43] Fidaly K, Ceballos C, Falguières A, Veitia MSI, Guy A, Ferroud C. Visible light photoredox organocatalysis: A fully transition metal-free direct asymmetric α -alkylation of aldehydes. *Green Chemistry*. 2012;**14**:1293-1297
- [44] Barata-Vallejo S, Yerien DE, Postigo A. Benign perfluoroalkylation of aniline derivatives through photoredox organocatalysis under visible-light irradiation. *European Journal of Organic Chemistry*. 2015;**2015**:7869-7875
- [45] Guo W, Lu LQ, Wang Y, Wang YN, Chen JR, Xiao WJ. Metal-free, room-temperature, radical alkoxycarbonylation of aryldiazonium salts through visible-light photoredox catalysis. *Angewandte Chemie International Edition*. 2015;**54**:2265-2269
- [46] Hopkinson MN, Sahoo B, Glorius F. Dual photoredox and gold catalysis: Intermolecular multicomponent oxyarylation of alkenes. *Advanced Synthesis & Catalysis*. 2014;**356**: 2794-2800
- [47] Yoshioka E, Kohtani S, Jichu T, Fukazawa T, Nagai T, Kawashima A, Takemoto Y, Miyabe H. Aqueous-medium carbon-carbon bond-forming radical reactions catalyzed by excited rhodamine B as a metal-free organic dye under visible light irradiation. *Journal of Organic Chemistry*. 2016;**81**:7217-7229
- [48] Pitre SP, McTiernan CD, Ismaili H, Scaiano JC. Metal-free photocatalytic radical trifluoromethylation utilizing methylene blue and visible light irradiation. *ACS Catalysis*. 2014;**4**:2530-2535
- [49] Yamamoto S, Fujiyama Y, Shiozaki M, Sueishi Y, Nishimura N. Hydride transfer reactions of leuco methylene blue and leuco thionine with some *p*-benzoquinones. *Journal of Physical Organic Chemistry*. 1995;**8**:805-809
- [50] Kalaitzakis D, Kouridaki A, Noutsias D, Montagnon T, Vassilikogiannakis G. Methylene blue as a photosensitizer and redox agent: Synthesis of 5-hydroxy-1*H*-pyrrol-2(5*H*)-ones from furans. *Angewandte Chemie International Edition*. 2015;**54**:6283-6287
- [51] Park JH, Ko KC, Kim E, Park N, Ko JH, Ryu DH, Ahn TK, Lee JY, Son SU. Photocatalysis by phenothiazine dyes: Visible-light-driven oxidative coupling of primary amines at ambient temperature. *Organic Letters*. 2012;**14**:5502-5505
- [52] Zhu X, Xie X, Li P, Guo J, Wang L. Visible-light-induced direct thiolation at α -C(sp³)-H of ethers with disulfides using acridine red as photocatalyst. *Organic Letters*. 2016;**18**:1546-1549
- [53] Mihldorf B, Wolf R. C-H photooxygenation of alkyl benzenes catalyzed by riboflavin tetraacetate and a non-heme iron catalyst. *Angewandte Chemie International Edition*. 2016;**55**:427-430
- [54] Nevesely T, Svobodová E, Chudoba J, Sikorski M, Cibulka R. Efficient metal-free aerobic photooxidation of sulfides to sulfoxides mediated by a vitamin B₂ derivative and visible light. *Advanced Synthesis & Catalysis*. 2016;**358**:1654-1663