We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Tandem-, Domino- and One-Pot Reactions Involving Wittig- and Horner-Wadsworth-Emmons Olefination

Fatima Merza, Ahmed Taha and Thies Thiemann

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.70364>

Abstract

The Wittig olefination utilizing phosphoranes and the related Horner-Wadsworth-Emmons (HWE) reaction using phosphonates transform aldehydes and ketones into substituted alkenes. Because of the versatility of the reactions and the compatibility of many functional groups towards the transformations, both Wittig olefination and HWE reactions are a mainstay in the arsenal of organic synthesis. Here, an overview is given on Wittig- and Horner-Wadsworth-Emmons (HWE) reactions run in combination with other transformations in one-pot procedures. The focus lies on one-pot oxidation Wittig/HWE protocols, Wittig/HWE olefinations run in concert with metal catalyzed cross-coupling reactions, Domino Wittig/HWE—cycloaddition and Wittig-Michael transformations.

Keywords: Wittig olefination, one-pot reactions, Domino reactions, tandem reactions, Horner-Wadsworth-Emmons olefination

1. Introduction

The Wittig olefination utilizing phosphoranes and the related Horner-Wadsworth-Emmons (HWE) reaction using phosphonates transform aldehydes and ketones into substituted alkenes. Because of the versatility of the reactions and the compatibility of many functional groups in the transformations, both Wittig olefination and HWE reactions are a mainstay in the arsenal of organic synthesis. The mechanism of the Wittig olefination has been the subject of intense debate [1]. While initially it was supposed that all Wittig olefination reactions lead via 1,2-addition to betaine structures **4** as zwitterionic intermediates that would form oxaphosphetane **3** with a final release of alkene and phosphine oxide by ring opening (*syn*-cycloreversion process), it has been seen more recently that especially under salt-free, aprotic conditions, many ylides undergo

a π^2s/π^2a [2+2]-cycloaddition with the carbonyl component leading to the oxaphosphetane **3** directly [2], which in certain cases can be in equilibrium with betaine structures **4** (Scheme 1). In HWE reactions, the deprotonated phosphonate **6a** undergoes a nucleophilic addition to the carbonyl compound (e.g., **7**), which usually is the rate limiting step [3]. The elimination to the final products proceeds through oxaphosphetane **9** (Scheme 2). The Wittig olefination has been used industrially in the synthesis of terpenoids [4]. Recently, a one-pot synthesis of the vasodilator and anti-platelet agent Beraprost sodium, a prostacyclin analog, was communicated with the HWE reaction as the key transformation with the idea of using the approach in an industrial synthesis of the pharmaceutical [5].

For years after the discovery of the Wittig olefination [6, 7], most Wittig transformations were carried out under inert atmosphere using dry solvents such as THF [8], DME [9], diethyl ether [10] and benzene [11]. Later it was realized that stabilized and semi-stabilized Wittig reagents can be reacted in non-de-aerated solvents, where the solvents need not be dried specifically. Most of these conjugated Wittig reagents are thermally stable and tolerate water, air and mild oxidants, while maintaining reactivity towards aldehydes and often also towards ketones. This allows for a plethora of reaction conditions for many Wittig olefination reactions such as obviating solvents altogether [12, 13] or running the reactions in aqueous solutions [14, 15] or in mixed solvents [16]. Also, it permits one-pot transformations of Wittig olefinations in combination with other reactions, also because the stabilized and to some extent the semi-stabilized phosphoranes are inert to mild oxidizing and reducing agents. However, also with non-stabilized phosphoranes, where reactions have to be performed under the exclusion of air and moisture, Wittig reactions can be performed in conjunction with further transformations [17].

This chapter is to give an insight into the types of transformations that can be combined with Wittig- and Horner-Wadsworth-Emmons olefinations in Domino-, tandem and one-pot

Scheme 1. Schematic presentation of the reaction mechanism of the Wittig olefination.

Scheme 2. General reaction mechanism of the HWE reaction.

reaction strategies. These include the preparation of phosphoranes and their reaction *in situ*, one-pot oxidation of alcohols to aldehydes and Wittig-olefination, *in situ*-recycling of phosphine oxides and catalytic Wittig reactions, one-pot Wittig-olefination metal catalyzed C–C bond forming reactions such as Suzuki-Miyaura, Sonogashira- and Heck reactions, Wittig and Horner-Emmons reactions in combination with polar cyclizations, Wittig-reactions carried out in combination with electrocyclic reactions, one-pot Wittig and Horner Emmons-addition reactions; cascade reactions featuring (triphenylphosphoranylidene)-ethenone and similar phosphoranes.

2. Wittig and Horner-Wadsworth-Emmons (HWE) olefination reactions with phosphoranes and phosphonates prepared *in situ*

Primarily, phosphoranes as Wittig reagents are prepared by the reaction of a triarylphosphine, usually triphenylphosphine, or, more seldom, a trialkylphosphine, and an alkyl halide with subsequent dehydrohalogenation of the triaryl(alkyl)alkylphosphonium halide produced. Non-stabilized Wittig reagents are not stable enough to be stored over longer periods of time; therefore, it is the norm that the Wittig-ylide is formed *in situ* from the oftentimes stable phosphonium salt, usually with a strong base, and then reacted directly with the carbonyl compound. In the case of stabilized phosphoranes, they are often stable enough to store, and the dehydrohalogenation necessitates only a weak base such as sodium carbonate or even sodium bicarbonate [18]. Nevertheless, this likewise allows the preparation of the phosphorane and the subsequent Wittig olefination in one pot [19], where even protic solvents can be used, such

as water. Similarly, semi-stabilized phosphoranes can be obtained *in situ* from their respective phosphonium salts, also even in aqueous medium, where LiCl promotes the Wittig olefination and suppresses the decomposition of the phosphoranes [14, 15]. Furthermore, all the catalytic Wittig reactions (see below) rely on the fact that the phosphorane is produced *in situ*.

Perhaps more interesting is the one-pot reaction of an alkyl halide, a phosphine and a carbonyl compound (**Scheme 3**). This can be achieved by consecutive addition of the components, when one or more of the components are sensitive, or by mixing of all components simultaneously. A consecutive addition of components in one pot was pursued by McNulty and Das who reacted air-sensitive triethylphosphine with benzyl bromides to the respective benzyltriethylphosphonium bromides, which were transformed to the phosphoranes with aq. NaOH, before being reacted with benzaldehydes to give (*E*)-stilbenes in an aqueous Wittig olefination [20]. Here, the triethylphosphine oxide by-product is water soluble. This reaction procedure has been diversified further by a one-pot preparation of benzyltriethylphosphonium bromides from the air-stable triethylphosphine hydrobromide and benzyl alcohols and subsequent Wittig olefination with aromatic aldehydes in aqueous medium [21]. Simultaneous mixing of alkyl halide such as α -haloesters (e.g., **13**), α -halonitriles, α -halocarbonyl compounds and α -alkyl- α -halocarbonyl compounds, triphenylphosphine (**12**), and carbonyl compound (e.g., **11**, **15**, **18**) in the presence either of a base [17, 22–26] or an alkene [27] was shown to give α,β -unsaturated esters [17, 22–27] (e.g., **14**, **17**, **19**), α,β -unsaturated nitriles [23, 26] and enones [27], respectively (**Scheme 3**). Epoxides are stable under these reaction conditions as can be seen in the transformation of **18** to **19** (**Scheme 1**). A one-pot, fluoride catalyzed Wittig-olefination has also been devised, where ethyl bromoacetate is reacted with carbaldehydes in the presence of tri-*n*-butylphosphine and tetrabutylammonium fluoride (Bu_4NF) to give (*E*)-configured α,β -unsaturated esters in good yield [28]. The synthesis of α,β -unsaturated esters has also been achieved from their alkyl halide and aldehyde constituents using tributylarsine [29] or a substituted triarylarsine instead of triphenylphosphine [30]. The use of tributylarsine in the presence triphenyl phosphite [29] led to the creation of a catalytic system which was

Scheme 3. *In situ* preparation of phosphoranes and subsequent Wittig olefination.

developed further with one-pot transformations that were managed with catalytic amounts (2 mol%) of poly(ethylene glycol) and (PEG)-supported tellurides in the presence of K_2CO_3 as base [31–34]. Also, micellar reaction systems such as micellar solutions of sodium dodecyl sulfate (SDS) in water have been used, in which Wittig olefinations were carried out between aldehydes and phosphoranes, synthesized *in situ* [35, 36]. A. Galante has performed Wittig reactions in the fluorous phase with *in situ* pre-formed perfluorinated ylides [37].

Traditionally, stabilized halophosphoranes have been prepared by the halogenation of the nonhalogenated parent phosphoranes and a subsequent dehydrohalogenation of the halogenated phosphonium salt obtained. Karama et al. have combined this *in situ* halogenation: dehydrohalogenation step with the Wittig reaction itself. Additionally, an *in situ* alcohol oxidation to provide the aldehyde starting material was integrated into many of these reaction sequences (**Scheme 4**) [38–42].

Scheme 4. One-pot oxidation, halogenation, and Wittig reaction to 2-haloacrylates.

3. *In situ* alcohol oxidation—Wittig/HWE reactions; other *in situ* aldehyde preparations run with subsequent Wittig/HWE sequences in one pot

The tolerance of stabilized phosphoranes towards mild oxidants allows for the oxidation of an alcohol to an aldehyde and its Wittig reaction in one-pot (**Schemes 5 and 6**). As oxidants, activated MnO_2 [43–46], barium permanganate [47, 48], tetra-*n*-propylammonium perruthenate (TPAP)/*N*-methylmorpholine *N*-oxide (NMO) [49–54] and TPAP/*N,N,N',N'*-tetramethylethylenediamine dioxide (TMEDAO₂) [55], *o*-iodoxybenzoic acid (IBX) [56–58], Dess-Martin periodinane [59–61], DMSO-oxalyl chloride (Swern conditions) [62–64], DMSO- SO_3 -pyridine (Parikh-Doering oxidation) [38, 39] or DMSO- SO_3 -triethylamine [65], pyridinium chlorochromate (PCC) or PCC/celite [66–69] as well as pyridinium dichromate (PDC) [70] such as PDC encapsulated in sol gel [71] have been used. In addition, metal catalyzed aerobic oxidation reactions of aldehydes with concomitant olefination reactions are known, where [(eta-*p*-cymene)RuCl₂]₂ (**27**) [72], nanoparticulate ruthenium supported on highly porous aluminum oxyhydroxide [73] or on silica gel [74], and nickel nanoparticles [75, 76] (**Scheme 6**) have been used as catalyst in the case of a concomitant Wittig reaction and gold/palladium bimetallic nanoparticles in the case of a concomitant Horner-Wadsworth-Emmons (HWE) reaction [77], Cu(I)-phenanthroline as a catalyst in an oxidation: HWE: sequential procedure [78].

Taylor et al. give a good overview of the tandem oxidation-Wittig processes developed until 2005, focusing especially on the tandem oxidation process (TOP) developed by his group [43–46],

Scheme 5. One-pot MnO_2 -mediated oxidation—Wittig olefination.

Scheme 6. One-pot metal catalyzed oxidation of alcohols utilizing oxygen—Wittig reaction.

using activated MnO_2 [79]. Over the years, this process has been used more often [40, 80–97] than the other processes shown above. Recently, also MnO_2 derived molecular sieve material such as OMS-2 [$\text{KMn}^{4+}\text{Mn}^{3+}\text{O}_{16} \cdot n\text{H}_2\text{O}$] has been used with success in aerobic, catalytic one-pot oxidation Wittig reactions of benzylic and allylic alcohols to the respective cinnamates [98]. Overall, the

Wittig transformations of the aldehydes produced *in situ* allows for the manipulation of aldehydes that are inherently instable such as of silyl substituted aldehydes, propargyl aldehyde [97], and chiral γ -aminoaldehydes, the latter without loss of stereochemical integrity (**Scheme 5**) [89]. In the case of Wittig transformations of chiral α -aminoaldehydes, β -aminoalcohols were oxidized to α -aminoaldehydes with NaOCl in the presence of AcNH-TEMPO, where the crude α -aminoaldehydes gained from the oxidation were subjected directly to olefination to give Wittig products without loss of stereochemical integrity [99–101].

Other preparation methods of aldehydes in conjunction with Wittig olefinations or HWE reactions have been reported. Thus, an oxidative cleavage of a glycol can be carried out in combination with a subsequent Wittig-olefination [102–105] (**Scheme 7**). Also a one-pot carboxylic acid to aldehyde reduction and Wittig reaction is known [106]. Finally, a Domino hydroformylation/Wittig olefination procedure has been developed, starting from allylamines (**Scheme 8**). The aldehyde is not isolated [107]. Domino/hydroformylation/Wittig olefination protocols have been introduced with other olefinic starting materials, also [108–110].

Scheme 7. Oxidative glycol cleavage—Wittig reaction.

Scheme 8. Hydroformylation—Wittig reaction.

4. Wittig- and HWE reactions and C–C-coupling reactions in one-pot procedures

Wittig- and Horner-Wadsworth-Emmons reactions can be combined with C–C-coupling reactions such as Suzuki cross-coupling [111–113], Mizoroki-Heck reaction [113–118] and Sonogashira-reaction [119]. Initially, it was observed that conjugated phosphoranes were stable under reaction conditions used for Heck- or Suzuki reactions (**Scheme 9**). Thus, phosphoranes themselves could be functionalized by Suzuki- [120], Mizoroki-Heck- [121], or Sonogashira-type [119] cross-coupling reactions, either in solution or when polymer-bound [122]. These phosphoranes could then be subjected to normal Wittig-olefination reactions with ketones or aldehydes [120–122]. The one-pot Wittig-Heck-reaction strategy can be extended to include an *O*-alkylation, where the Wittig reaction of a *p*-hydroxybenzaldehyde (**43**) with methylenetriphenylphosphorane, obtained *in situ* from phosphonium salt **44** provides the *p*-hydroxystyrene as the olefin component in the Mizoroki-Heck reaction in the presence of an alkyl bromide (e.g., **45**), which *O*-alkylates the phenoxy-function to give alkoxystilbenes **46** (**Scheme 9**) [123].

Scheme 9. One-pot Heck cross-coupling/Wittig reaction.

5. One-pot Wittig- and HWE olefination/cycloaddition reaction

One can easily visualize that an alkene prepared by a Wittig olefination can easily be used as a 2- π component in cycloaddition reactions, in one pot (**Scheme 10**). A typical such cycloaddition is [4+2]-cycloaddition, such as the classical Diels Alder reaction, which can be performed both inter-[69, 124] and intramolecularly [125–132] in tandem with a Wittig-reaction.

Hilt and Hengst have published a cobalt(I)-catalyzed Diels Alder reaction of alkynyltriphenylphosphonium and 1,3-dienes with a consecutive Wittig reaction of the cycloadduct with

Scheme 10. Oxidation—Wittig-olefination—Diels-Alder reaction sequence.

various aldehydes in one pot that lead after a further dehydrogenative step to substituted stilbenes and styrenes (**Scheme 11**) [133].

Interesting is the cycloaddition of *in situ* produced benzyne (**55**) to 1,4-diphenylbutadiene, prepared *in situ* by HWE reaction from cinnamaldehyde, (**15**) give 1,4-diphenylnaphthalene (**56**) (**Scheme 12**) [134].

The transformation sequence Diels-Alder/Wittig can be part of a more complex reaction chain. Thus, Ramachary and Barbas III [135] have forwarded a Domino Wittig/Knoevenagel/

Scheme 11. Cobalt (I)-catalyzed Diels Alder reaction—Wittig reaction.

Scheme 12. One-pot HWE reaction—cycloaddition of *in situ* produced benzyne.

Diels-Alder sequence to spirotriones **58** (**Scheme 13**) and a Wittig/Knoevenagel/Diels-Alder/Huisgen cycloaddition sequence to polysubstituted triazoles **61** (**Scheme 14**).

Oxidation of benzyl alcohols to benzaldehydes can be incorporated into a Wittig-Diels Alder sequence [69]. Also, hetero-Diels-Alder reactions can be run in tandem with a Wittig olefination as shown by Ramachary et al. in their synthesis of tetrahydropyrans **64** (**Scheme 15**) [136]. Here, diamine **63** is used as a catalyst. The reaction, however, gives the product only in low enantiomeric excess (**Scheme 15**).

Huisgen type [3+2]-cycloaddition reactions can be run also in a simple tandem process rather than incorporated in a more complex reaction chain (see above). A typical example is shown in **Scheme 16**, where azidoethyl-tetrahydro-hydroxyfuran **66** is treated with phosphorane **21** to give triazoline **68** alongside diazoamine **69** [137]. Further such approaches are known [138, 139].

Scheme 13. Domino Wittig/Knoevenagel/Diels-Alder sequence.

Scheme 14. Wittig/Knoevenagel/Diels-Alder/Huisgen cycloaddition sequence.

Scheme 15. Wittig-reaction/hetero-Diels Alder reaction.

Scheme 16. Wittig reaction—intramolecular Huisgen type [3+2]-cycloaddition.

6. One-pot Wittig- and HWE olefination/addition reaction

Electrophiles can be added to the alkene function obtained, in a one-pot reaction with the Wittig olefination. A typical example is the stereoselective bromination of the Wittig product with oxalyl bromide (**71**), where triphenylphosphine oxide (**70**) as side product of the olefination step acts as a catalyst in the bromination (Scheme 17) [140]. Hamza and Blum have developed a sol-gel entrapped tertiary phosphine by co-polycondensation of tetramethoxysilane, 2-diphenyl(phosphino)ethyltri(ethoxy)silane and *N*-2-(aminoethyl)-3-aminopropyltri(methoxy)silane. This could be reacted in a Wittig type olefination with benzyl chlorides (e.g., **76**) and benzaldehydes, prepared *in situ* from benzyl alcohols (e.g., **75**). The strategy allows for the combination of the process with a bromination step in one pot by addition of sol-gel-bound pyridinium hydrobromide perbromide after completion of the Wittig reaction (Scheme 18) [71].

Alternatively, the process can be combined with a hydrogenation step by the addition of hydrogen in the presence of an added heterogenized Wilkinson catalyst (Scheme 19) [141]. A further Wittig olefination—hydrogenation sequence was developed by Zhou et al. who obtained α -CF₃- γ -ketoesters **82** by adding trichlorosilane to the reaction mixture where triphenylphosphine oxide (again as side product of the Wittig olefination) acts as a Lewis base and activates the silane as hydrogenating agent (Scheme 20) [142]. The routine was expanded to other aldehydes including alkanals as educts [143]. This reaction was also carried out with

Scheme 17. Wittig olefination— Ph_3PO -catalyzed addition of bromine.

Scheme 18. Wittig olefination—addition of bromine.

Scheme 19. Wittig olefination—hydrogenation.

glyoxal derivatives **84** as starting materials, where after conjugate addition of trichlorosilane a few drops of methanol were added to the solution resulting in conversion of the trichlorosilylenol ether (**86**) to the keto compound **87** while at the same time generating HCl, which then promoted a Paal-Knorr reaction of **87** to the furan **88** (Scheme 21) [144].

Scheme 20. Wittig reaction—triphenylphosphine oxide catalyzed hydrogenation.

Scheme 21. Furan synthesis by one-pot Wittig olefination—hydrogenation—Paal-Knorr reaction.

Lu and Toy showed that the Wittig-olefination—trichloromethylsilane conjugate addition sequence can be coupled with the initial preparation of the phosphorane in one pot [145]. The conjugate addition to furnish the silyl enol ether can be combined with a reductive Aldol reaction, where for the Wittig reaction and for the reductive Aldol reaction two separate aldehydes can be used (**Scheme 22**) [145]. The reactions above can be run with a triarylphosphine-tertiary amine bifunctional polymeric reagent (Rasta-Resin-PPh₃-NBnⁱPr₂), where the polymer bound triarylphosphine oxide also exerts a catalyzing effect on the addition of Cl₃SiH while making it possible to recycle the polymer [146].

As many Wittig olefinations can be performed in aqueous medium, it is possible to combine the reaction with an enzymatic step. One such sequence is the enzymatic reduction of the olefinic moiety by a recombinant enoate reductase from *Gluconobacter oxydans*, carried out with an enzyme-coupled *in situ* cofactor regeneration with a glucose dehydrogenase as enzyme component and D-glucose as co-substrate (**Scheme 23**) [147].

Interestingly, a Wittig reaction can also be run in combination with an enzymatic reduction, where the *in situ* prepared enone **93** is transformed to the alkenol **94** (**Scheme 24**) [148].

Scheme 22. Wittig olefination—reductive Aldol reaction.

Scheme 23. Wittig-olefination—enzymatic ene-hydrogenation.

Scheme 24. Wittig-olefination—enzymatic keto-reduction.

The possibility of a combination of a Wittig/HWE olefination and a Michael addition has been studied by a number of research groups. Thus, Piva and Comesse have added phosphonoesters to copper enolates derived from the 1,4 addition of cuprates **97** to enones **96** with the idea that the enolate would deprotonate the phosphonoester **98** producing the reactive ketone and phosphonate, which undergo HWE reaction. Products **99** of the tandem Michael-HWE reaction are produced in acceptable yield (Scheme 25) [149, 150]. This strategy was used with *p*-methylcinnamaldehyde (**100**) as carbonyl component in the total synthesis of ($\pm$)-*ar*-turmerone (**105**), a bisabolane-type natural product found in *Zingiber* and *Curcuma* species (Scheme 26) [151].

Scheme 25. One-pot Michael addition—HWE reaction.

Scheme 26. Synthetic route to utilizing a one-pot Michael addition—HWE reaction.

7. One-pot Wittig-olefination/functional group interconversion

Wittig reactions can be performed with alkoxy carbonylmethylidenetriphenylphosphorane (**21**) in aq. NaOH, where the cinnamates formed are hydrolysed *in situ* to cinnamic acids **106** (**Scheme 27**) [152]. After completion of the reaction, triphenylphosphine oxide can be filtered off from the strongly basic, aqueous solution, and the cinnamic acids are isolated by simple filtration after acidification of the filtrate. Pinacol-acetal tripropylphosphonium salt **107** has been reacted in aq. 1 M NaOH with different benzaldehydes **37**; the cinnamaldehyde *O,O*-pinacol acetal can be hydrolyzed directly to the cinnamaldehydes **108** with 25w% aq. H_3PO_4 (**Scheme 28**) [153].

This procedure provides a nice alternative to the reaction of benzaldehydes with triphenylphosphoranylideneacetaldehyde, which often produces dienals and trienals as side-products. A tandem Wittig-cyanosilylation was developed by Zhou et al., where again Ph_3PO as side product of the Wittig olefination acts as Lewis base to catalyze TMS-CN in the cyanosilylation step. Chiral salen

Scheme 27. One-pot Wittig reaction—ester hydrolysis.

Scheme 28. One-pot Wittig reaction—acetal hydrolysis.

aluminum catalyst **109** was used as Lewis acid to activate the keto function in the cyanosilylation. Products were obtained with high enantioselectivity [68–93%ee]. TMSCN and chiral catalyst **109** were added after completion of the Wittig reaction, albeit in one pot (Scheme 29) [143].

As Wittig reactions can be carried out in aqueous medium, enzymatic reactions can be integrated into the process (*vide supra*). In this regard, M. Krauß et al. showed that 4-phenylbut-3-en-2-ones (**93**), obtained by Wittig olefination, are reduced to the corresponding 4-phenylbut-3-en-2-ols (**94**) in >99 ee(%) using (*S*)-alcohol dehydrogenase [(*S*)-ADH] from *Rhodococcus* sp. or (*R*)-ADH from *Lactobacillus kefir* [148].

Scheme 29. Wittig reaction—cyanosilylation.

8. One-pot Wittig- and HWE olefination/cyclization

Michael type cyclization—cyclic hemiacetals can be used efficiently as substrates in Wittig olefination reactions with stabilized Wittig reagents. After the Wittig reaction, the tethered alcohol

function induces a cyclization through a Michael reaction. This reaction sequence has been used especially in the construction of functionalized C-glycosides such as in the stereospecific synthesis of ω -amino- β -D-furanoribosylacetic acid derivative **115** (Scheme 30) [154].

In their synthesis to C-glycoside amphiphiles, Ranoux et al. followed a similar strategy, reacting non-protected sugars with HWE reagents in aqueous or solventless conditions, leading to C-glucosides **117** and **121** (Scheme 31) [155].

A different mechanism to C-glucosides operates when 5,6-dideoxy-5,6-anhydro-6-nitro-D-glucofuranose **122** is reacted with an excess of phosphorane **21**. Here, **21** acts as a base and **122** experiences an anion driven ring opening to **123**, which undergoes an oxy-Michael addition to **124** with concomitant Wittig reaction, resulting in C-vinyl glycoside **125** (Scheme 32) [156].

A highly stereoselective tandem Wittig-reaction-Michael addition has been developed by Liu et al. [157] when reacting 3-carboxy-2-oxopropylidene)triphenylphosphorane **126** with

Scheme 30. Synthesis of ω -amino- β -D-furanoribosylacetic acid derivative **115** utilizing a Wittig olefination-ring closure reaction en route.

Scheme 31. Synthesis of C-glucosides with a HWE—ring closure reaction.

Scheme 32. Tandem oxy-Michael addition–Wittig reaction.

enaldehydes (e.g., **15**), using a chiral pyrrolidine-based catalyst such as **128** (**Scheme 33**). Most likely, the asymmetric Michael addition proceeds by the reaction of **15** with the iminium compound **129** (**Scheme 33**), formed from **15** with catalyst **128**.

Beltrán-Rodil et al. have elaborated a retro-aldol initiated Wittig-olefination-Michael addition sequence leading to an exchange of the hydroxyl function in **130** for a carbalkoxymethyl group in **134**. The retro-aldol reaction is effected by the commercially available trimethylamine *N*-oxide (TMAO, **131**) [158] (**Scheme 34**).

Electrocyclizations, incl. photocyclizations, and pericyclic reactions: Electrocyclization can be run in concert with Wittig reactions. One such example is shown in **Scheme 35**, where allylic bromide **135**, the product of a Morita-Baylis-Hillman transformation, is converted with triphenylphosphine to the corresponding phosphonium salt, which is reacted with benzaldehyde (**11**) to give triene **136**. **136**, heated under aeration, undergoes a 6π -electrocyclization—base catalyzed aerobic oxidation to *o*-terphenyl derivative **137** (**Scheme 35**) [159].

Scheme 33. Asymmetric Michael-addition-Wittig-olefination.

Scheme 34. Retro-aldol-Wittig-olefination-Michael addition cascade.

Scheme 35. One-pot phosphorane synthesis—Wittig-reaction—6 π -electrocyclization—oxidative dehydrogenation.

Similarly, Hamza and Blum [71], who developed a Wittig olefination with a sol-gel entrapped tertiary phosphine derived phosphorane (*vide supra*, **Schemes 18 and 19**) showed that the Wittig reaction can be run in concert with a photochemical cyclization under aerobic conditions to produce phenanthrene (**138**) (**Scheme 36**) [71].

A number of tandem Wittig/HWE reaction—Claisen/Cope rearrangements have been reported [160–173]. A typical example is shown in **Scheme 37**, where neat (4-fluorophenoxyacetyl)cyanomethylene)triphenylphosphorane **139** is subjected to microwave irradiation at 450 W in a sealed tube to undergo an intramolecular Wittig reaction—Claisen rearrangement to furnish benzofuran **134** (**Scheme 37**) [173].

Mali et al. achieved the synthesis of seselin and angelicin derivatives (e.g., **148** and **150**) by a tandem Wittig-olefination—Claisen rearrangement from propargyl and chloroalkyl ethers of 2,4-dihydroxybenzaldehyde and 2,4-dihydroxyacetophenone (e.g., **146** and **149**) (**Scheme 38**) [164].

Nevertheless, sometimes, these reactions are not easy to control. Thus, a cascade of Wittig reaction and double Claisen and Cope rearrangements starting from 2,4-prenyloxybenzaldehyde

Scheme 36. Wittig olefination—photocyclization—oxidative dehydrogenation.

Scheme 37. Intramolecular Wittig reaction—Claisen rearrangement.

Scheme 38. One-pot syntheses of seselin and angelicin derivatives.

151 leads to a plethora of products through the range of reactions that are possible with the intermediate **153**, itself produced through the Wittig reaction and a first Claisen rearrangement. The final products found include gravelliferone (**154**, 10%), balsamiferone (**155**, 5%), and 6,8-diprenylumbelliferone (**156**, 15%) (**Scheme 39**) [169].

Less common is the tandem Wittig and ene reaction. Tilve et al. have published such a combination of Wittig and ene reaction in their total synthesis of (±)-kainic acid (**160**), an amino

Scheme 39. Wittig reaction—double Claisen and cope rearrangements.

acid found in different species of red algae [174]. Here, the product was formed in 65% yield as a mixture of diastereoisomers **159a**/**159b** in a ratio of 1:5. Previously, the authors had synthesized ($\pm$)-kainic acid (**160**) utilizing a Wittig—Michael reaction as the key step (**Scheme 40**) [175].

Finally, the possibility of a tandem Wittig-olefination—aza-Wittig rearrangement should be mentioned—this combination was carried out on 2-benzoylaziridine **161** to give stereoisomeric dehydropiperidines **163**/**164** (**Scheme 41**) [176].

Scheme 40. Wittig-ene cascade as a key step towards the synthesis of kainic acid (**160**).

Scheme 41. Tandem Wittig-olefination—aza-Wittig-rearrangement.

9. Other transformations

A wealth of further transformations have been found to be possible in combination with Wittig/HWE reactions. Thus, cyclopropanation of alkenes using sulfur-ylide reagent **166** can be run in tandem with a Wittig reaction with a conjugated phosphorane such as **21**. This combination of reactions can be performed with the preparation of the aldehyde as the Wittig substrate by oxidation of the corresponding alcohol **165** with MnO_2 in one pot (**Scheme 42**) [177].

Scheme 42. One-pot oxidation—Wittig-olefination—cyclopropanation.

Generally, non-stabilized phosphoranes are basic. This basicity has been used by Knüppel et al. in the transformation of α,α -dibromoene **168** with excess methylenetriphenylphosphorane, where the phosphorane induces a Corey-Fuchs-reaction-type dehydrobromination/debromination to generate a terminal alkyne, which together with the concomitantly run Wittig-olefination delivers **169**, an intermediate to the trisnorsesquiterpene (–)-clavukerin A (**171**) (**Scheme 43**)

Scheme 43. Wittig-olefination—Corey-Fuchs-reaction-type dehydrobromination/debromination.

[178]. A metathesis reaction completes the sequence to **171**. In this case, the metathesis reaction is not run in one pot with the previous transformations.

Nevertheless, one-pot Wittig—metathesis reactions are well known from the literature [179–181]. A typical example is shown in **Scheme 44**, where catalyst **174** serves both as a catalyst for the metathesis as well as for the Wittig olefination, when the *in situ* produced aldehyde **175** is treated with triphenylphosphine and ethyl diazoacetate (**176**) in one pot (**Scheme 44**).

Scheme 44. One-pot Wittig—Metathesis reaction.

10. Conclusion

Due to the fact that phosphoranes and phosphonates are stable under more diverse conditions than was initially realized, it has become possible to perform reaction cascades and one-pot reactions with Wittig- and HWE reactions as an integral part of the reaction sequence. Frequently, Wittig olefination reactions are carried out with *in situ* prepared phosphonium salts and phosphoranes [17, 22–27]. One-pot oxidation—Wittig olefination reactions are also quite common [40, 43–110], especially when the carbonyl component is labile [89, 97]. Often, the oxidant of choice is MnO_2 [40, 43–46, 79–97], although a number of reactions are known where transformations were carried with air oxygen using metals and metal oxides as catalysts [72–78]. As many Wittig- and HWE reactions tolerate metal catalysts, this allows the running of Wittig/HWE reactions in combination with metal catalyzed cross coupling reactions and olefinations such as Heck [114–123], Suzuki [111–113], Sonogashira [119–121], and metathesis reactions [179–181]. The alkenes gained in the olefination reactions can be submitted to cycloaddition reactions, including Diels Alder reactions [69]. Furthermore, the alkenes lend themselves to 1,2-addition reactions [71, 140] in one-pot procedures. In cases where enones or enaldehydes are produced in the olefination reaction, a 1,4-addition becomes a possibility; this includes the Michael addition [149–151]. Also, the combination of ring opening of cyclic hemiacetals or acetals, olefination reaction and a 1,4-addition leading to ring closure is quite common [154–156]. The outcome of one-pot sequences of olefination reaction—electrocyclic rearrangement can be predicted less easily. Nevertheless, one-pot Wittig olefination—Claisen- [173], Wittig olefination—Cope- [169], and Wittig olefination—aza Wittig [176] rearrangement reactions have been published. Lastly, Wittig olefination and HWE reactions have been combined with functional group transformations, including the hydrolysis of an ester function [152] and the reduction of a carbonyl group [148].

The prospects of multi-step, one-pot reactions and reaction cascades incorporating Wittig reagents can be seen in the rich chemistry of ketylenetriphenylphosphorane (**178**) (Scheme 45) [182–185], which has been reviewed earlier [182, 186, 187]. Lastly, catalytic Wittig reactions can be seen as a subset of tandem reactions involving phosphoranes. Further research in specifically this area will help make the Wittig olefination more atom-economical and environmentally sustainable, so that this reliable alkene forming reaction will remain a competitive olefination strategy of choice.

Scheme 45. Cascade reactions with ketylenetriphenylphosphorane (**178**).

Author details

Fatima Merza¹, Ahmed Taha² and Thies Thiemann^{1*}

*Address all correspondence to: thies@uaeu.ac.ae

¹ Department of Chemistry, College of Science, United Arab Emirates

² UAEU Libraries, United Arab Emirates University, Al Ain, United Arab Emirates

References

- [1] Maryanoff BE, Reitz AB. The Wittig olefination reaction and modifications involving phosphoryl stabilized carbanions, stereochemistry, mechanism, and selected synthetic aspects. *Chemical Reviews*. 1989;**89**:863-927
- [2] Vedejs E, Marth CF. Mechanism of the Wittig reaction: The role of substituents at phosphorus. *Journal of the American Chemical Society*. 1988;**110**:3948-3958
- [3] Larsen EO, Aksnes G. Kinetic studies of the Horner reaction I. Phosphorus Silicon. 1983;**15**:219-228
- [4] Pommer H. The Wittig reaction in industrial practice. *Angewandte Chemie (International Ed. in English)*. 1977;**16**:423-429
- [5] Wang Y, Hou DL, Hu HZ, Zhang BZ, Wang ES. Efficient synthesis of beraprost sodium. *Chemical Research in Chinese Universities*. 2016;**32**:581-585
- [6] Wittig G, Schöllkopf U. Über Triphenylphosphinmethylene als olefinbildende Reagenzien (1. Mitteil.). *Chemische Berichte*. 1954;**87**:1318-1330
- [7] Wittig G, Haag W. Ueber Triphenylphosphinmethylene als olefinbildende Reagenzien (2. Mitteil.). *Chemische Berichte*. 1955;**88**:1654-1666
- [8] Taber C, Nelson CG. Potassium hydride in paraffin: A useful base for organic synthesis. *The Journal of Organic Chemistry*. 2006;**71**:8973-8974
- [9] Corey EJ, Clark DA, Goto G, Marfat A, Mioskowski C, Samuelsson B, Hammerström B. Stereospecific total synthesis of a "slow reacting substance" of anaphylaxis, leukotriene C-1. *Journal of the American Chemical Society*. 1980;**192**:1436-1439
- [10] Wittig G, Schöllkopf U. Methylenecyclohexane. *Organic Syntheses*. 1960;**40**:66-66
- [11] Aksnes G, Berg TJ, Gramstad T. Temperature and solvent effects in Wittig reactions. *Phosphorus, Sulfur, and Silicon*. 1995;**106**:79-84
- [12] Thiemann T, Watanabe M, Tanaka Y, Mataka S. Solvent-free Wittig olefination with stabilized phosphoranes—scope and limitations. *New Journal of Chemistry*. 2004;**28**:578-584
- [13] Thiemann T. Solventless Wittig reactions with fluorinated benzaldehydes. *Journal of Chemical Research*. 2007;**31**:336-341
- [14] Wu J, Zhang D, Wei S. Wittig reactions of stabilized phosphorus ylides with aldehydes in water. *Synthetic Communications*. 2005;**35**:1213-1222
- [15] Wu J, Zhang D. Aqueous Wittig reactions of aldehydes with *in situ* formed semistabilized phosphorus ylides. *Synthetic Communications*. 2005;**35**:2543-2551
- [16] Watanabe M, Ribeiro Morais G, Mataka S, Ideta K, Thiemann T. Two variations of solvent-reduced Wittig olefination reactions—comparison of solventless Wittig reactions to Wittig reactions under ultrasonication with minimal work-up. *Zeitschrift für Naturforschung B*. 2005;**60**:909-915

- [17] Parvartur PT, Torney PS, Tilve SG. Recent developments of Wittig reaction in organic synthesis through tandem or sequential processes. *Current Organic Synthesis*. 2013;**10**: 288-317
- [18] Kelly MJB, Fallot LB, Gustaffson JL, Bergdahl BM. Water mediated Wittig reactions of aldehydes in the teaching laboratory: Using sodium bicarbonate for the in situ formation of stabilized ylides. *Journal of Chemical Education*. 2016;**93**:1631-1636
- [19] Al Jasem Y, El-Esawi R, Thiemann T. Wittig- and Horner-Wadsworth-Emmons (HWE) olefination reactions with stabilized and semi-stabilized phosphoranes and phosphonates under non-classical conditions. *Journal of Chemical Research*. 2014;**38**:453-463
- [20] McNulty J, Das P. Highly stereoselective and general synthesis of (*E*)-stilbenes and alkenes by means of an aqueous Wittig reaction. *European Journal of Organic Chemistry*. 2009;4031-4035
- [21] McNulty J, Das P, McLeod D. Microwave-assisted, aqueous Wittig reactions: Organic-solvent- and protecting-group-free chemoselective synthesis of functionalized alkenes. *European Journal of Chemistry*. 2010;**16**:6756-6760
- [22] El-Batta A, Jiang C, Zhao W, Anness R, Cooksy AL, Bergdahl M. Wittig reactions in water media employing stabilized ylides with aldehydes. Synthesis of α,β -unsaturated esters from mixing aldehydes, α -bromoesters, and Ph_3P in aqueous NaHCO_3 . *The Journal of Organic Chemistry*. 2007;**72**:5244-5259
- [23] Wu J, Yue C. One-pot Wittig reactions in aqueous media: A rapid and environmentally benign synthesis of α,β -unsaturated carboxylic esters and nitriles. *Synthetic Communications*. 2006;**36**:2939-2947
- [24] Westman J. An efficient combination of microwave dielectric heating and the use of solid-supported triphenylphosphine for Wittig reactions. *Organic Letters*. 2001;**3**:3745-3747
- [25] Choudhary BM, Mahendar K, Kantam ML, Ranganath KVS, Athar T. The one-pot Wittig reaction: A facile synthesis of α,β -unsaturated esters and nitriles by using nanocrystalline magnesium oxide. *Advanced Synthesis and Catalysis*. 2006;**348**:1977-1985
- [26] Kantam ML, Kumar KBS, Balasubramanyam V, Venkanna GT, Figueras F. One-pot Wittig reaction for the synthesis of α,β -unsaturated esters using highly basic magnesium/lanthanum mixed oxide. *Journal of Molecular Catalysis A: Chemical*. 2010;**321**:10-14
- [27] Liu D-N, Tian S-K. Stereoselective synthesis of polysubstituted alkenes through a phosphine-mediated three-component system of aldehydes, α -halo carbonyl compounds, and terminal alkenes. *Chemistry – A European Journal*. 2009;**15**:4538-4542
- [28] Fumagalli T, Sello G, Orsini F. One-pot, fluoride promoted Wittig reaction. *Synthetic Communications*. 2009;**39**:2178-2195
- [29] Shi L, Wang W, Wang Y, Huang Y. The first example of a catalytic Wittig-type reaction. Tri-*n*-butylarsine catalysed olefination in the presence of triphenyl phosphite. *The Journal of Organic Chemistry*. 1989;**54**:2027-2028

- [30] Wang C, Yang L, Zhou R, Mei G. Solvent-free, one-pot synthesis of α,β -unsaturated esters in the presence of a C3-symmetric arsine. *Synthetic Communications*. 2016;**46**:1074-1079
- [31] Huang Z-Z, Ye S, Xia W, Yu Y-H, Tang Y. Wittig-type olefination catalysed by PEG-telluride. *The Journal of Organic Chemistry*. 2002;**67**:3096-3103
- [32] Huang Z-Z, Ye S, Xia W, Tang Y. A practical catalytic Wittig-type reaction. *Journal of the Chemical Society, Chemical Communications*. 2001;1384-1385
- [33] Huang Z-Z, Tang Y. Unexpected catalyst for Wittig-type and dehalogenation reactions. *The Journal of Organic Chemistry*. 2002;**67**:5320-5326
- [34] Tang Y, Ye S, Huang Z-Z, Huang Y-Z. Telluronium ylides in cyclopropanation and catalytic olefination. *Heteroatom Chemistry*. 2002;**13**:463-466
- [35] Orsini F, Sello G, Fumigalli T. One-pot Wittig reactions in water and in the presence of a surfactant. *Synlett*. 2006;1717-1718
- [36] Agarkar SV, Pathan RU. Green synthesis of morpholino cinnamides using sodium lauryl sulphate and water. *Chemical Science Review and Letters*. 2014;**4**:1-3
- [37] Galante A, Lihoste P, Sinou D. Wittig reaction using perfluorinated ylides. *Tetrahedron Letters*. 2001;**42**:5425-5427
- [38] Karama U, Mahfouz R, Al-Othman Z, Warad I, Almansoor A. One-pot combination of the Wittig olefination with bromination and oxidation reactions. *Synthetic Communications*. 2013;**43**:893-898
- [39] Karama U, Alshamari H, Abdelall H, Sultan MA. A facile one-pot synthesis of (Z)- α -chloro- α,β -unsaturated esters from alcohols. *Arabian Journal of Chemistry*. in press
- [40] Karama U, Al-Othman Z, Al-Majid A, Almansour A. A facile one-pot synthesis of α -bromo- α,β -unsaturated esters from alcohols. *Molecules*. 2010;**15**:3276-3280
- [41] Karama U. One-pot synthesis of (E)- α -chloro- α,β -unsaturated esters. *Journal of Chemical Research*. 2009;405-406
- [42] Karama U. One-pot approach to the conversion of alcohols into α -iodo- α,β -unsaturated esters. *Synthetic Communications*. 2010;**40**:3447-3451
- [43] Wei X, Taylor RJK. *In situ* oxidation – Wittig reactions. *Tetrahedron Letters*. 1998;**39**:1815-3818
- [44] Blackburn L, Wei X, RJK T. Manganese oxide can oxidize unactivated alcohols under in situ oxidation-Wittig conditions. *Journal of the Chemical Society, Chemical Communications*. 1999;1337-1338
- [45] Wei X, Taylor RJK. In situ manganese dioxide alcohol oxidation – Wittig reactions: Preparation of bifunctional dienyl building blocks. *The Journal of Organic Chemistry*. 2000;**65**:616-620

- [46] Taylor RJK, Campbell L, McAllister GD. ($\pm$)-trans-3,3'-(1,2-cyclopropanediyl)bis-2-(*E*)-propenoic acid, diethyl ester: Tandem oxidation procedure (TOP) using MnO₂ oxidation – Stabilized phosphorane trapping. *Organic Syntheses*. 2008;**85**:15-26
- [47] Shuto S, Niizuma S, Matsuda S. One-pot conversion of α,β -unsaturated alcohols into the corresponding carbon-elongated dienes with a stable phosphorus-ylide – BaMnO₄ system. Synthesis of 6'-methylene derivatives of neplanocin A as potential antiviral nucleosides. New neplanocin analogues. 11. *The Journal of Organic Chemistry*. 1998;**63**:4489-4493
- [48] Gholinejad M, Firouzabadi H, Bahrami M, Nájera C. Tandem oxidation – Wittig reaction using nanocrystalline barium manganate (BaMnO₄): An improved one-pot protocol. *Tetrahedron Letters*. 2016;**57**:3773-3775
- [49] MacCoss RN, Balkus EP, Ley SV. A sequential tetra-*n*-propylammonium perruthenate (TPAP)-Wittig oxidation olefination protocol. *Tetrahedron Letters*. 2003;**44**:7779-7781
- [50] Hoshi M, Kaneko O, Nakajima M, Arai S, Nishida A. Total synthesis of $\pm$ -lundurine B. *Organic Letters*. 2014;**16**:768-771
- [51] Lagouette R, Šebesta P, Jiroš P, Kalinová B, Jirošová A, Straka J, Černá K, Šobotník J, Cvačka J, Jahn U. Total synthesis, proof of absolute configuration, and biosynthetic origin of stylpsal, the first isolated sex pheromone of *Strepsiptera*. *Chemistry – A European Journal*. 2013;**19**:8515-8524
- [52] Candy M, Tomas L, Parat S, Hernan V, Bienaymé H, Pons J-M, Bressy C. A convergent approach to (–)-callystatin A based on local symmetry. *Chemistry – A European Journal*. 2012;**18**:14267-14271
- [53] Matovic NJ, Hayes PY, Penman K, Lehmann RP, De Voss JJ. Polyunsaturated alkyl amides from *Echinacea*: Synthesis of diynes, enynes, and dienes. *The Journal of Organic Chemistry*. 2011;**76**:4467-4481
- [54] Boyer A, Isono N, Lackner S, Lautens M. Domino rhodium(I)-catalysed reactions for the efficient synthesis of substituted benzofurans and indoles. *Tetrahedron*. 2010;**66**:6468-6482
- [55] Read CDG, Moore PW, Williams CM. *N,N,N',N'*-Tetramethylenediamine dioxide (TMEDAO₂) facilitates atom economical/open atmosphere Ley-Griffith (TPAP) tandem oxidation – Wittig reactions. *Green Chemistry*. 2015;**17**:4537-4540
- [56] Crich D, Mo X-S. One pot selective 5'-oxidation/olefination of 2'-deoxynucleosides. *Synlett*. 1999;67-68
- [57] Maiti A, Yadav JS. One-pot oxidation and Wittig olefination of alcohols using *o*-iodoxybenzoic acid and stable Wittig ylide. *Synthetic Communications*. 2001;**31**:1499-1506
- [58] Chen J, Fu X-G, Zhou L, Zhang J-T, Qi X-L, Cao X-P. A convergent route for the total synthesis of malynamides O, P, Q, and R. *The Journal of Organic Chemistry*. 2009;**74**:4149-4157

- [59] Huang CC. Synthesis of C-14 labelled 1-(4-methoxybenzoyl)-5-oxo-2-pyrrolidinepropanoic acid (CI-933). *Journal of Labelled Compounds and Radiopharmaceuticals*. 1987;**24**:676-681
- [60] Barrett AGM, Hamprecht D, Ohkubo M. Dess-Martin periodinane oxidation of alcohols in the presence of stabilized phosphorus ylides: A convenient method for the homologation of alcohols via unstable aldehydes. *The Journal of Organic Chemistry*. 1997;**62**:9376-9378
- [61] Mori K. Pheromone synthesis. Part 257: Synthesis of methyl (2*E*,4*Z*,7*Z*)-2,4,7-decatrienoate and methyl (*E*)-2,4,5-tetradecatrienoate, the pheromone components of the male dried bean beetle, *Acanthoscelides obtectus* (Say). *Tetrahedron*. 2015;**71**:5589-5596
- [62] Ireland RE, Norbeck DW. Application of the Swern oxidation to the manipulation of highly reactive carbonyl compounds. *The Journal of Organic Chemistry*. 1985;**50**:2198-2200
- [63] Salgado A, Mann E, Sanchez-Sancho F, Herradon B. Synthesis of heterocyclic γ -amino- α,β -unsaturated acid derivatives and peptide-heterocycle hybrids. *Heterocycles*. 2003;**60**:57-71
- [64] Lin S, Deiana L, Tseggai A, Córdova A. Concise total synthesis of dihydrocorynanthenol, Protoemetinol, protoemetine, 3-*epi*-protoemetinol and emetine. *European Journal of Organic Chemistry*. 2012;**398**-408
- [65] Crisóstomo FRP, Carrillo R, Martín T, García-Tellado F, Martín VS. A convenient one-pot oxidation/Wittig reaction for the C2-homologation of carbohydrate-derived glycols. *The Journal of Organic Chemistry*. 2005;**70**:10099-10101
- [66] Bressette AR, Glover IV LC. A convenient one-pot PCC oxidation Wittig reaction of alcohols. *Synlett* 2004:738-740.
- [67] Shet J, Desai V, Tilve S. Domino primary alcohol oxidation-Wittig reaction: Total synthesis of ABT-418 and (*E*)-4-oxonon-2-enoic acid. *Synthesis*. 2004;1859-1863
- [68] Majik MS, Parameswaran PS, Tilve SG. Total synthesis of (-)- and (+)-tedanalactam. *The Journal of Organic Chemistry*. 2009;**74**:6378-6381
- [69] Harris GH, Graham AE. Efficient oxidation-Wittig olefination-Diels-Alder multicomponent reactions of α -hydroxyketones. *Tetrahedron Letters*. 2010;**51**:6890-6892
- [70] Dhumaskar KL, Bhat C, Tilve SG. PDC-mediated tandem oxidative Wittig olefination. *Synthetic Communications*. 2014;**44**:1501-1506
- [71] Hamza K, Blum J. One-pot combination of the Wittig olefination with oxidation, hydrogenation, bromination and photocyclization reactions. *Tetrahedron Letters*. 2007;**48**:293-295
- [72] Kim G, Lee DG, Chang S. *In situ* aerobic alcohol oxidation-Wittig reactions. *Bulletin of the Korean Chemical Society*. 2001;**22**:943-944

- [73] Lee EY, Kim Y, Lee JS, Park J. Ruthenium-catalyzed, one-pot alcohol oxidation-Wittig reaction producing α,β -unsaturated esters. *European Journal of Organic Chemistry*. 2009;2943-2946
- [74] Carillo AI, Schmidt LC, Marín ML, Scaiano JC. Mild synthesis of mesoporous silica supported ruthenium nanoparticles as heterogeneous catalysts in oxidative Wittig coupling reactions. *Catalysis Science & Technology*. 2014;4:435-440
- [75] Alonso F, Riente P, Yus M. Wittig-type olefination of alcohols promoted by nickel nanoparticles: Synthesis of polymethoxylated and polyhydroxylated stilbenes. *European Journal of Organic Chemistry*. 2009;6034-6042
- [76] Alonso F, Riente P, Yus M. One-pot synthesis of stilbenes from alcohols through a Wittig-type olefination reaction promoted by nickel nanoparticles. *Synlett*. 2009;1579-1582
- [77] Miyamura H, Suzuki A, Yasukawa T, Kobayashi S. Integrated process of aerobic oxidation-olefination-asymmetric C-C bond formation catalysed by robust heterogeneous gold/palladium and chirally modified rhodium nanoparticles. *Advanced Synthesis and Catalysis*. 2015;357:3815-3819
- [78] Reddy AGK, Mahendar L, Satyanarayana G. Simple copper(I)-catalysed oxidation of benzylic/allylic alcohols to carbonyl compounds: Synthesis of functionalized cinnamates in one pot. *Synthetic Communications*. 2014;44:2076-2087
- [79] Taylor RJK, Reid M, Foot J, Raw SA. Tandem oxidation processes using manganese dioxide. *Accounts of Chemical Research*. 2005;38:851-869
- [80] MacDonald G, Alcaraz L, Wei X, Lewis NJ, Taylor RJK. Unsaturated amides derived from 2-amino-3-hydroxycyclopentenone: A Stille approach to Asuka mABA, 2880-II, and limocrocin. *Tetrahedron*. 1998;54:9823-9836
- [81] Alcaraz L, Macdonald G, Ragot J, Lewis NJ, Taylor RJK, Manumycin A. Revision of structure and synthesis of the (+)-enantiomer. *The Journal of Organic Chemistry*. 1998;63:3526-3527
- [82] Alcaraz L, Macdonald G, Ragot J, Lewis NJ, Taylor RJK. Synthetic approaches to the manumycin A, B, and C antibiotics: The first total synthesis of (+)-manumycin-A. *Tetrahedron*. 1999;55:3707-3716
- [83] Zeng F, Negishi E. A highly efficient, selective, and general method for the synthesis of conjugated (all-*E*)-oligoenes of the $(\text{CH}=\text{CH})_n$ type via iterative hydrozirconation-palladium-catalysed cross-coupling. *Organic Letters*. 2002;4:703-706
- [84] Petrowski RJ. Straightforward preparation of (2*E*,4*Z*)-2,4-heptadien-1-ol and (2*E*,4*Z*)-2,4-heptadienal. *Synthetic Communications*. 2003;33:3233-3241
- [85] Nicolaou KC, Li Y, Fylaktakidou KC, Monenschein H, Li Y, Weyershausen B, Mitchell HJ, Wei H-X, Guntupalli P, Hepworth D, Sugita K. Total synthesis of apopotolidin. *Journal of the American Chemical Society*. 2003;125:15433-15442

- [86] Aitken DJ, Faure S, Roche S. Synthetic approaches to the southern part of cyclotheonamide C. *Tetrahedron Letters*. 2003;**44**:8827-8830
- [87] Mladenova M, Tavlinova M, Momchev M, Alami M, Ourévitch M, Brion JD. An iterative procedure for the synthesis of conjugated omega-chlorotrienoic and -tetraenoic esters and related derivatives. *European Journal of Organic Chemistry*. 2003;2713-2718
- [88] Phillips DJ, Pillinger KS, Li W, Taylor AE, Graham AE. Desymmetrization of diols by a tandem oxidation/Wittig olefination reaction. *Journal of the Chemical Society, Chemical Communications*. 2006;2280-2282
- [89] Davies SB, McKervery MA. Convenient in-situ synthesis of nonracemic N-protected β -amino aldehydes from β -amino acids. Applications in Wittig reaction and heterocycle synthesis. *Tetrahedron Letters*. 1999;**40**:1229-1232
- [90] Lang S, Taylor RJK. Tandem oxidation-Wittig-Wittig sequences for the preparation of functionalized dienoates. *Tetrahedron Letters*. 2006;**47**:5489-5492
- [91] Blackburn L, Pei C, Taylor RJK. *In situ* alcohol oxidation-Wittig reactions using non-stabilized phosphoranes. *Synlett*. 2002;215-218
- [92] Blackburn L, Kanno H, Taylor RJK. In situ alcohol oxidation-Wittig reactions using *N*-methoxy-*N*-methyl-2-(triphenylphosphoranylidene)acetamide: Application to the synthesis of a novel analogue of 5-oxo-eicosatetraenoic acid. *Tetrahedron Letters*. 2003;**44**:115-118
- [93] Runcie KA, Taylor RJK. The *in situ* oxidation-Wittig reaction of α -hydroxyketones. *Journal of the Chemical Society, Chemical Communications*. 2002;974-975
- [94] Quesada E, Raw SA, Reid M, Roman E, Taylor RJK. One-pot conversion of activated alcohols into 1,1-dibromoalkenes and terminal alkynes using tandem oxidation processes with manganese dioxide. *Tetrahedron*. 2006;**62**:6673-6680
- [95] McAllister GA, Oswald MF, Paxton RJ, Raw SA, Taylor RJK. The direct preparation of functionalized cyclopropanes from allylic alcohols or α -hydroxyketones using tandem oxidation processes. *Tetrahedron*. 2006;**62**:6681-6694
- [96] Raw SA, Reid M, Roman E, Taylor RJK. A tandem oxidation procedure for the conversion of alcohols into 1,1-dibromoalkenes. *Synlett*. 2004;819-822
- [97] Pavlakos E, Georgiou T, Tofi M, Montagnon T, Vassilikogiannakis G. γ -Spiroketal γ -lactones from 2-(γ -hydroxyalkyl)furans: Syntheses of epi-pyrenolides D and crassalactone D. *Organic Letters*. 2009;**11**:4556-4559
- [98] Kona J, King'ondou CK, Howell AR, Suib SL. OMS-2 for aerobic, catalytic, one-pot alcohol oxidation-Wittig reactions: Efficient access to α,β -unsaturated esters. *ChemCatChem*. 2014;**6**:749-752
- [99] Loukas V, Noula C, Kokotos G. Efficient protocol for the synthesis of enantiopure gamma-amino acids with proteinogenic side chains. *Journal of Peptide Science*. 2003;**9**:312-319

- [100] Loukas V, Markidis T, Kokotos G. Amino acid based synthesis of chiral long chain diamines and tetramines. *Molecules*. 2002;**7**:767-776
- [101] Magrioti V, Antonopoulou G, Pantoleon E, Kokotos G. Synthesis of 2-amino alcohols and unnatural amino acids from serine. *ARKIVOC*. 2002;**Xiii**:51-55
- [102] Dunlap NK, Mergo W, Jones JM, Carrick JD. A general procedure for a one-pot oxidative cleavage/Wittig reaction of glycols. *Tetrahedron Letters*. 2002;**43**:3923-3925
- [103] Terrell LR, Ward JS II, Maleczka RE Jr. Synthetic studies toward amphidinolide A: A synthesis of fully functionalized subunits. *Tetrahedron Letters*. 1999;**40**:3097-3100
- [104] Ray PC, Roberts SM, Juliá-Colonna SM. Stereoselective epoxidation of some α,β -unsaturated enones possessing a stereogenic centre at the γ -position: Synthesis of a protected galactonic acid derivative. *Journal of the Chemical Society, Perkin Transactions 1*. 2001;149-153
- [105] Outram HS, Raw SA, Taylor RJK. *In situ* oxidative diol cleavage-Wittig process. *Tetrahedron Letters*. 2002;**43**:6185-6187
- [106] Duan Y, Yao P, Du Y, Feng J, Wu Q, Zhu D. Synthesis of α,β -unsaturated esters via a chemo-enzymatic chain elongation approach by combining carboxylic acid reduction and Wittig reaction. *Beilstein Journal of Organic Chemistry*. 2015;**11**:2245-2251
- [107] Farwick A, Helmchen G. Stereoselective synthesis of β -proline derivatives from allyl-amines via Domino hydroformylation/Wittig olefination and Aza-Michael addition. *Advanced Synthesis and Catalysis*. 2010;**352**:1023-1032
- [108] Ruan Q, Zhou L, Breit B. New phosphorus self-assembling ligands for the tandem hydroformylation-Wittig olefination reaction of homoallylic alcohols—A key step for stereoselective pyran synthesis. *Catalysis Communications*. 2014;**53**:87-90
- [109] Breit B, Zahn SK. Domino hydroformylation-Wittig reactions. *Angewandte Chemie, International Edition*. 1999;**38**:969-971
- [110] Wong GW, Landis CR. Iterative asymmetric hydroformylation/Wittig olefination sequence. *Angewandte Chemie, International Edition*. 2013;**52**:1564-1567
- [111] Thiemann T, Watanabe M, Tanaka Y, Mataka S. One-pot Wittig-olefination/Suzuki-reaction—The compatibility of conjugated phosphoranes in Pd(0) catalysed C-C-bond forming reactions. *New Journal of Chemistry*. 2006;**30**:359-369
- [112] Thiemann T, Watanabe M, Tanaka Y, Mataka S. One pot Suzuki coupling—Wittig olefination reactions. *Journal of Chemical Research*. 2004;**28**:723-727
- [113] Chaudhary AR, Bedekar AV. 1-(α -aminobenzyl)-2-naphthol as phosphine-free ligand Pd-catalysed Suzuki and one-pot Wittig-Suzuki reaction. *Applied Organometallic Chemistry*. 2012;**26**:430-437
- [114] Burmester C, Mataka S, Thiemann T. Synthesis of non-symmetric Divinylarenes by a Heck/Wittig reaction combination. *Synthetic Communications*. 2010;**40**:3196-3208

- [115] Saiyed AS, Bedekar AV. One-pot synthesis of stilbenes by dehydrohalogenation–Heck olefination and multicomponent Wittig–Heck reaction. *Tetrahedron Letters*. 2010;**51**:6227-6231
- [116] Saiyed AS, Patel KN, Kamath BV, Bedekar AV. Synthesis of stilbene analogues by one-pot oxidation-Wittig and oxidation-Wittig–Heck reaction. *Tetrahedron Letters*. 2012;**53**:4692-4696
- [117] Patel HA, Patel AL, Bedekar AV. Polyaniline-anchored palladium catalyst-mediated Mizoroki-Heck and Suzuki-Miyaura reactions and one-pot Wittig-Heck and Wittig-Suzuki reactions. *Applied Organometallic Chemistry*. 2015;**29**:1-6
- [118] Patel KN, Bedekar AV. One-pot synthesis and study of spectroscopic properties of oligo(phenylenevinylene)s. *Tetrahedron Letters*. 2015;**56**:6617-6621
- [119] Watanabe M, Mataka S, Thiemann T. One pot Sonogashira-coupling/Wittig olefination procedures. *Journal of Chemical Research*. 2005;**29**:636-639
- [120] Thiemann T, Umeno K, Ohira D, Inohae E, Sawada T, Mataka S. Suzuki-Kumada coupling with Bromoaryl methylidenephosphoranes. *New Journal of Chemistry*. 1999;**23**:1067-1070
- [121] Thiemann T, Umeno K, Inohae E, Mataka S. Novel elongated Phosphoranes by Heck-reaction and Pd(0)-catalysed Alkynylation and their use in C-7 group functionalisation in Estrones. *The Reports of Institute of Advanced Material Study, Kyushu University*. 2000;**14**(1):17-29 *Chem. Abstr.* 2000;133:335380w
- [122] Thiemann T, Umeno K, Wang J, Arima K, Watanabe M, Tabuchi Y, Tanaka Y, Gorohmaru H, Mataka S. Elongated phosphoranes by C-C coupling of haloaryl methylidene-phosphoranes—Synthesis and applications. *Journal of the Chemical Society, Perkin Transactions 1*. 2002;2090-2110
- [123] Patel KN, Kamath BV, Bedekar AV. Synthesis of alkoxy stilbenes by one-pot *O*-alkylation-Wittig and *O*-alkylation Wittig-Heck reaction sequence. *Tetrahedron Letters*. 2013;**54**: 80-84
- [124] Thiemann T, Ohira D, Li YQ, Sawada T, Mataka S, Rauch K, Noltemeyer M, de Meijere A. [4+2]-Cycloaddition of Thiophene-S-monoxides onto activated Methylene-cyclopropanes. *Journal of the Chemical Society, Perkin Transactions 1*. 2000;**17**:2968-2976
- [125] Jarosz S, Szewczyk K. Stability of regioisomeric sugar allyltins. Cleavage of the carbon-oxygen bond under radical conditions. *Tetrahedron Letters*. 2001;**42**:3021-3024
- [126] Jarosz S, Kozłowska E, Jeżewski A. Intramolecular Diels-Alder reaction of chiral, highly oxygenated trienoates derived from sugar allyltins. *Tetrahedron*. 1997;**53**:10775-10782
- [127] Jarosz S, Skóra S. A convenient route to enantiomerically pure highly oxygenated decalins from sugar allyltin derivatives. *Tetrahedron: Asymmetry*. 2000;**11**:1433-1448

- [128] Jarosz S. Tandem Wittig-Diels-Alder reaction between sugar-derived phosphoranes and sugar aldehydes. An easy route to optically pure highly oxygenated decalins. *Journal of the Chemical Society, Perkin Transactions 1*. 1997;3579-3580
- [129] Tilve SG, Torney PS, Patre RE, Kamat DP, Srinivasan BR, Zubkov FI. Domino Wittig-Diels Alder reaction: Synthesis of carbazole lignans. *Tetrahedron Letters*. 2016;57:2266-2268
- [130] Torney P, Patre R, Tilve S. A rapid assembly of furo[3,4-b]- and pyrrolo[3,4-b]carbazolones by Domino Wittig Diels Alder reaction. *Synlett*. 2011;639-642
- [131] Wu J, Sun L, Dai W-M. Microwave-assisted tandem Wittig-intramolecular Diels-Alder cycloaddition. Product distribution and stereochemical assignment. *Tetrahedron*. 2006;62:8360-8372
- [132] Wu J, Jiang X, Xu J, Dai W-M. Tandem Wittig-intramolecular Diels-Alder cycloaddition of ether-tethered 1,3,9-decatrienes under microwave heating. *Tetrahedron*. 2011;67:179-192
- [133] Hilt G, Hengst C. A concise synthesis of substituted stilbenes and styrenes from propargylic phosphonium salts by a cobalt-catalysed Diels-Alder/Wittig olefination reaction sequence. *The Journal of Organic Chemistry*. 2007;72:7337-7342
- [134] Chen Z, Shou W, Wang Y. One-pot synthesis of 1,4-diarylnaphthalenes via a Wittig-Horner reaction/[4+2]-cycloaddition/dehydrogenation sequence. *Synthesis*. 2009;1075-1080
- [135] Ramachary DB, Barbas CF. Towards organo-click chemistry: Development of organocatalytic multicomponent reactions through combinations of Aldol, Wittig, Knoevenagel, Michael, Diels-Alder and Huisgen cycloaddition reactions. *Chemistry—A European Journal*. 2004;10:5323-5331
- [136] D. B. Ramachary, R. Mondal, S. Jain, Direct organocatalytic Wittig/hetero-Diels-Alder reactions in one-pot: Synthesis of highly-substituted tetrahydropyranes. *ARKIVOC* 2016;ii:98-115.
- [137] Herdeis C, Schiffer T. Synthesis of nonracemic 2,3,6-trisubstituted piperidine derivatives from sugar lactones via tandem Wittig [2+3] cycloaddition reaction. A novel entry to prosopis and cassia alkaloids. *Tetrahedron*. 1999;55:1043-1056
- [138] Herdeis C, Telser J. A stereoselective synthesis of nonracemic (+)-desoxoprosophylline by a tandem Wittig [2+3]-cycloaddition reaction. *European Journal of Organic Chemistry*. 1999;1407-1414
- [139] Herdeis C, Schiffer T. Synthesis of chiral nonracemic homo-1-deoxyzasugars with D-talo- and L-allo-configuration via tandem Wittig [2+3] cycloaddition reaction. *Tetrahedron*. 1999;52:14745-14756
- [140] Yu T-Y, Wei H, Luo Y-C, Wang Y, Wang Z-Y, Xu P-F. PPh_3O as an activating reagent for one-pot stereoselective syntheses of di- and polybrominated esters from simple aldehydes. *The Journal of Organic Chemistry*. 2016;81:2730-2736

- [141] Gelman F, Blum J, Avnir D. One-pot reactions with opposing reagents: Sol-gel entrapped catalyst and base. *Journal of the American Chemical Society*. 2000;**122**:11999-12000
- [142] Chen L, Shi T-D, Zhou J. Waste as catalyst: Tandem Wittig/conjugate addition reduction sequence to α -CF₃ γ -keto esters that uses Ph₃PO as catalyst for the chemoselective conjugate reduction. *Chemistry, An Asian Journal*. 2013;556-559
- [143] Cao J-J, Zhou F, Zhou J. Improving the atom efficiency of the Wittig reaction by a "waste as catalyst/co-catalyst" strategy. *Angewandte Chemie, International Edition*. 2010;**49**:4976-4980
- [144] Chen L, Du Y, Zeng X-P, Shi T-D, Zhou F, Zhou J. Successively recycle waste as catalyst: A one-pot Wittig/1,4-reduction/Paal-Knorr sequence for modular synthesis of substituted furans. *Organic Letters*. 2015;**17**:1557-1560
- [145] Lu J, Toy PH. Tandem one-pot Wittig/reductive aldol reactions in which the waste from one catalyzes a subsequent reaction. *Chemistry, An Asian Journal*. 2011;**6**:2251-2254
- [146] Teng Y, Lu J, Toy PH. Rasta resin-PPh₃-NBnPr₂ and its use in one-pot Wittig cascades. *Chemistry, An Asian Journal*. 2012;**7**:351-359
- [147] Kraußer M, Winkler T, Richter N, Dommer S, Fingerhut A, Hummel W, Gröger H. Combination of C=C bond formation by Wittig reaction and enzymatic C=C bond reduction in a one-pot process in water. *ChemCatChem*. 2011;**3**:293-296
- [148] Kraußer M, Hummel W, Gröger H. Enantioselective one-pot two-step synthesis of hydrophobic allylic alcohols in aqueous medium through the combination of a Wittig reaction and an enzymatic ketone reduction. *European Journal of Organic Chemistry*. 2007;5175-5179
- [149] Piva O, Comesse S. Tandem Michael-Wittig-Horner reaction: One-pot synthesis of δ -substituted α,β -unsaturated esters. *Tetrahedron Letters*. 1997;**38**:7191-7194
- [150] Piva O, Comesse S. Tandem Michael-Wittig-Horner reaction: One-pot synthesis of δ -substituted α,β -unsaturated carboxylic acid derivatives – Application to a concise synthesis of (Z)- and (E)-ochtoden-1-al. *European Journal of Organic Chemistry*. 2000;2417-2424
- [151] Chuzel O, Piva O. Tandem Michael-Wittig-Horner reaction: Application to the synthesis of bisabolenes. *Synthetic Communications*. 2003;**33**:393-402
- [152] Thiemann T, Elshorbagy MW, Salem M, Ahmadani SAN, Al-Jasem Y, Al Azani M, Al-Sulaibi M, Al-Hindawi B. Facile, direct reaction of benzaldehydes to 3-arylprop-2-enoic acids and 3-arylprop-2-ynoic acids in aqueous medium. *International Journal of Organic Chemistry*. 2016;**6**:26-141
- [153] McNulty J, Zepeda-Velazquez C, McLeod D. Development of a robust reagent for the two-carbon homologation of aldehydes to (E)- α,β -unsaturated aldehydes. *Green Chemistry*. 2013;**15**:3146-3149

- [154] Ma D, Li X, Hao L, Zhang P, Chen H, Zhao Y. Stereospecific synthesis of ω -amino- β -D-furanoribosyl acetic acid based on microwave assisted Wittig-Michael tandem reaction. *Chemical Journal of Chinese Universities*. 2014;**35**:959-964
- [155] Ranoux A, Lemiègre L, Benoit M, Guégan J-P, Benvegu T. Horner-Wadsworth-Emmons reaction of unprotected sugars in water or in the absence of any solvent: One-step access to C-glycoside amphiphiles. *European Journal of Organic Chemistry*. 2010;**32**:1314-1323
- [156] Kumar RS, Karthikeyan K, Kumar BVNP, Muralidharan D, Perumal PT. Synthesis of densely functionalized C-glycosides by a tandem oxy Michael addition-Wittig olefination pathway. *Carbohydrate Research*. 2010;**345**:457-461
- [157] Liu Y-K, Ma C, Jiang K, Liu T-Y, Chen Y-C. Asymmetric tandem Michael addition – Wittig reaction to cyclohexenone annulation. *Organic Letters*. 2009;**11**:2848-2851
- [158] Beltrán-Rodil S, Donald JR, Edwards MG, Raw SA, Taylor RJK. Tandem retro-aldol/Wittig/Michael and related cascade processes. *Tetrahedron Letters*. 2009;**50**:3378-3380
- [159] Lim CH, Kim SH, Kim KH, Kim JN. An efficient synthesis of *o*-terphenyls from Morita-Baylis-Hillman adducts of cinnamaldehydes: A consecutive bromination, Wittig reaction, 6π -electrocyclization, and an aerobic oxidation process. *Tetrahedron Letters*. 2013;**54**:2476-2479
- [160] Quesada E, Taylor RJK. Tandem Horner-Wadsworth-Emmons olefination/Claisen rearrangement/hydrolysis sequence: Remarkable acceleration in water with microwave irradiation. *Synthesis*. 2005;**3**:3193-3195
- [161] Ajmeri AA, Rao SSM. Studies in synthesis and characterization of tandem Wittig and Claisen reactions of cinnamylated and prenylated 2, 4-dihydroxybenzaldehyde. *Indian Journal of Heterocyclic Chemistry*. 2014;**24**:175-180
- [162] Mali RS, Sandhu PK. One-pot Wittig reaction and Claisen rearrangement of 2-hydroxy-4-prenyloxybenzaldehyde and 2-hydroxy-4-prenyloxyacetophenone: Synthesis of 2',3',3'-trimethyl-2',3'-dihydroangelicins. *Journal of Chemical Research (S)*. 1996;**148**:149
- [163] Mali RS, Joshi P. Useful syntheses of prenylated and pyrano-3-arylcoumarins. *Synthetic Communications*. 2001;**31**:2883-2891
- [164] Mali RS, Pandhare NA, Sindkhedkar MD. Convenient two-step syntheses of seselin and angelicin derivatives. *Tetrahedron Letters*. 1995;**36**:7109-7110
- [165] Mali RS, Walture AS, Babu KN. Synthesis of 7-desoxy-8-allylcoumarins. *Organic Preparations and Procedures International*. 1996;**28**:217-221
- [166] Mali RS, Joshi PP, Sandhu PK, Manekar-Tilve A. Efficient syntheses of 6-prenylcoumarins and linear pyranocoumarins: Total synthesis of suberosin, toddaculin, O-methylapigravin (O-methylbrosiperin), O-methylbalsamiferone, dihydroxanthyletin, xanthyletin, and luvangetin. *Journal of the Chemical Society, Perkin Transactions 1*. 2002;**371**:376

- [167] Rehman H, Rao JM. Tandem intramolecular Wittig and Claisen rearrangement reactions in the thermolysis of 2-methyl-2-phenoxypropionylcyanomethylenetriphenylphosphoranes: Synthesis of substituted 2H-1-benzopyrans and benzofurans. *Tetrahedron*. 1987;**43**:5335-5340
- [168] Yadla R, Rao JM. Thermolysis of phenoxyacetylcyanomethylenetriphenylphosphoranes – Tandem intramolecular Wittig and Claisen rearrangement reactions. *Heterocycles*. 1987;**26**:329-331
- [169] Patre RE, Shet JB, Parameswaran PS, Tilve SG. Cascade Wittig reaction-double Claisen and Cope rearrangements: One-pot synthesis of diprenylated coumarins gravelliferone, balsamiferone, and 6,8-diprenylumbelliferone. *Tetrahedron Letters*. 2009;**50**:6488-6490
- [170] Schmidt B, Riemer M. Synthesis of allyl- and prenylcoumarins via microwave-promoted tandem Claisen rearrangement/Wittig olefination. *Synthesis*. 2016;**48**:141-149
- [171] Kawasaki T, Terashima R, Sakaguchi K-e, Sekiguchi H, Sakamoto M. A short route to “reverse-prenylated” pyrrolo[2,3-*b*]indoles via tandem olefination and Claisen rearrangement of 2-(3,3-dimethylallyloxy)indol-3-ones: First total synthesis of flustramine C. *Tetrahedron Letters*. 1996;**37**:7525-7528
- [172] Rama Rao VVVNS, Reddy GV, Yadla R, Narsaiah B, Rao PS. Synthesis of fluorine containing 3-cyano/ethoxycarbonyl-2-ethyl-benzo[*b*]furans via microwave assisted tandem intramolecular Wittig and Claisen rearrangement. *ARKIVOC*. 2005;**3**:211-220
- [173] Rama Rao VVVNS, Reddy GV, Maitraie D, Ravikanth S, Yadla R, Narsaiah B, Rao PS. One-pot synthesis of fluorine containing 3-cyano/ethoxycarbonyl-2-methyl-benzo[*b*]furans. *Tetrahedron*. 2004;**60**:12231-12237
- [174] Majik MS, Parameswaran PS, Tilve SG. Tandem Wittig-Ene reaction approach to Kainic acid. *The Journal of Organic Chemistry*. 2009;**74**:3591-3594
- [175] Bhat C, Tilve SG. Tandem approach for the synthesis of functionalized pyrrolidones: Efficient routes toward allokalnic acid and kainic acid. *Tetrahedron Letters*. 2013;**54**:245-248
- [176] Coldham I, Collis AI, Mould RJ, Rathmell RE. Synthesis of 4-phenylpiperidines by tandem Wittig Olefination – aza- Wittig rearrangement of 2-benzoylaziridines. *Journal of the Chemical Society, Perkin Transactions 1*. 1995;2739-2745
- [177] Huang WH, Wang LL. One-pot synthesis of cyclopropane derivatives with a *cis/trans* stereoselectivity by Wittig olefination - sulfur ylide cyclopropanation sequence. *Journal of Chemical Research*. 2013;380-384
- [178] Knüppel S, Rogachev VO, Metz P. A concise catalytic route to the marine sesquiterpenoids (–)-clavukerin A and (–)-isoclavukerin A. *European Journal of Organic Chemistry*. 2010;**(32)**:6145-6148
- [179] Sirasani G, Paul T, Andrade RB. Sequencing cross-metathesis and non-metathesis reactions to rapidly access building blocks for synthesis. *Tetrahedron*. 2011;**67**:2197-2205

- [180] Paul T, Andrade RB. Sequential cross-metathesis/phosphorus-based olefination: Stereoselective synthesis of 2,4-dienoates. *Tetrahedron Letters*. 2007;**48**:5367-5370
- [181] Murelli RP, Snapper ML. Ruthenium-catalyzed tandem cross-metathesis/Wittig olefination: Generation of conjugated dienoic esters from terminal olefins. *Organic Letters*. 2007;**9**:1749-1752
- [182] Schobert R, Gordon GJ. Bioactive heterocycles from Domino Wittig-pericyclic reactions. *Current Organic Chemistry*. 2002;**6**:1181-1196
- [183] Schobert R, Löffler J, Siegfried S. Phosphacumulene ylides as versatile C₂ building blocks: New pathways to tetronates, coumarins and azacycles. *Targets in Heterocyclic Systems*. 1999;**3**:245
- [184] Schobert R, Dietrich M, Mullen G, Urbina-Gonzalez J-M. Phosphorus ylide based functionalizations of tetronic and tetramic acids. *Synthesis*. 2006;3902-3914
- [185] Marcos IS, Pedrero AB, Sexmero MJ, Diez D, Basabe P, Hernandez FA, Urones JG. Synthesis and absolute configuration of three natural *ent*-halimanoilides with biological activity. *Tetrahedron Letters*. 2003;**44**:369-372
- [186] Schobert R. Domino syntheses of bioactive tetronic and tetramic acids. *Naturwissenschaften*. 2007;**94**:1-11
- [187] Schobert R, Hölzl C. Heterocycles from unsaturated phosphorus ylides. *Topics in Heterocyclic Chemistry*. 2008;**12**:193-218