We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Biomolecules Oxidation by Hydrogen Peroxide and Singlet Oxygen

Kazutaka Hirakawa

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.71465>

Abstract

Hydrogen peroxide (H_2O_2) and singlet oxygen ($^1\text{O}_2$) are important reactive oxygen species (ROS) for biological and medicinal fields. Oxidation processes of chemical materials by molecular oxygen are important H_2O_2 source, whereas photochemical reaction is important for $^1\text{O}_2$ production. Reactivity and biomolecule damage by these ROS depend on the surrounding conditions and targeting molecules. In this chapter, production mechanisms of H_2O_2 and $^1\text{O}_2$, biomolecule oxidation by these ROS, their detection methods, and production control of $^1\text{O}_2$ are briefly reviewed.

Keywords: hydrogen peroxide, singlet oxygen, DNA damage, protein damage, photooxidation

1. Introduction

Biomolecule damage, for example, oxidation of DNA and/or protein, by reactive oxygen species (ROS) is closely related to carcinogenicity [1–3] and/or toxicity [4–6]. Furthermore, oxidative damage to unwanted tissue can be applied to the treatment of disease including cancer treatment [7–9], and similar reaction is applied to sterilization [10–14]. Hydrogen peroxide (H_2O_2) is a relatively long-lived ROS compared with a short-lived ROS such as superoxide anion radicals ($\text{O}_2^{\bullet-}$) [15]. One of the most important producing mechanisms of H_2O_2 is a dismutation of $\text{O}_2^{\bullet-}$, which is easily formed through oxidation of various materials by dioxygen molecule (O_2). Various carcinogenic chemical compounds produce H_2O_2 through their oxidation processes. Relationship among molecular oxygen and ROS is shown in **Figure 1**. Oxygen molecules are easily reduced by surrounding materials, and various ROS and the intermediates are formed (**Figure 1A**). In the case of photosensitized reaction, excited states of oxygen molecules are produced (**Figure 1B**). Singlet oxygen ($^1\text{O}_2$), which is also an important ROS, can

Figure 1. Relationship among ground-state oxygen molecule ($^3\text{O}_2$) and ROS (A) and the energy levels of oxygen molecule (B). HOMO and SOMO are the abbreviations of highest occupied molecular orbital and semi-occupied molecular orbital, respectively. The “arrows” in (B) indicate the electron spin.

be easily generated via photosensitized reaction [16–18]. The $^1\Sigma_g^+$ state ($^1\text{O}_2(^1\Sigma_g^+)$) is mainly produced through the excitation energy transfer from the excited state, in general triplet excited (T_1) state, of photosensitizer [16–18]. The $^1\text{O}_2(^1\Sigma_g^+)$ has higher energy, 1.6 eV, corresponding to the ground state of oxygen molecule ($^3\text{O}_2$). The lifetime of $^1\text{O}_2(^1\Sigma_g^+)$ is several picoseconds, and $^1\text{O}_2(^1\Sigma_g^+)$ is rapidly converted to the $^1\Delta_g$ state ($^1\text{O}_2(^1\Delta_g)$) [16–18]. Because the lifetime of $^1\text{O}_2(^1\Delta_g)$ (several microseconds) is markedly longer than that of $^1\text{O}_2(^1\Sigma_g^+)$, $^1\text{O}_2(^1\Delta_g)$ is a more important ROS. After that, $^1\text{O}_2$ indicates $^1\text{O}_2(^1\Delta_g)$ without explanation in this chapter. Visible light, other than ultraviolet radiation, has sufficient energy to produce $^1\text{O}_2$ from the ground state of oxygen molecule. Therefore, $^1\text{O}_2$ production is an important mechanism of phototoxicity and/or photo-carcinogenicity under strong light illumination with phototoxic materials. The purpose of this chapter is a review of the ROS-mediated biomolecule damage and the related topics.

2. Hydrogen peroxide

Hydrogen peroxide itself is not strongly ROS. However, other ROS including hydroxyl radicals ($\cdot\text{OH}$) are produced from H_2O_2 . In general, H_2O_2 is produced from the dismutation of $\text{O}_2^{\bullet -}$, and, in vivo, production of H_2O_2 and $\text{O}_2^{\bullet -}$ occurs in mitochondria [19]. In this section, H_2O_2 formation from compounds, specifically artificial materials, is introduced.

2.1. Hydrogen peroxide formation through oxidation of chemical compounds

One of the most important processes of H_2O_2 production is a dismutation of $\text{O}_2^{\bullet -}$. Various chemical compounds or metals can be oxidized by oxygen molecules. In the case of a simple electron transfer-mediated oxidation, $\text{O}_2^{\bullet -}$ is produced by the electron extraction from chemical compounds or metals. The lifetime of $\text{O}_2^{\bullet -}$ in aqueous solution is about several milliseconds [15]. The produced $\text{O}_2^{\bullet -}$ in aqueous media is converted to H_2O_2 through the dismutation by proton (H^+) as follows:

For example, hydroquinone, which is one of the metabolites of benzene, can produce H_2O_2 through the autoxidation process (**Figure 2**) [20]. This process is markedly enhanced by the presence of metal ions, specifically Cu^{2+} ions [20]. In the presence of sacrificial reductants, for example, nicotinamide adenine dinucleotide (NADH), the oxidized form of hydroquinone, *p*-benzoquinone, is reduced to the parent hydroquinone. Consequently, the redox cycle is formed, leading to the production of H_2O_2 abundantly. It has been also reported that hydrazine analogues produce H_2O_2 through their autoxidation processes (**Figure 3**) [21–23].

2.2. Hydrogen peroxide production through photochemical processes

Photochemical processes also contribute to the formation of H_2O_2 . Because the reorganization energy of the reduction of small molecule, such as O_2 molecules, through electron transfer becomes large due to the Marcus theory [24, 25], the $\text{O}_2^{\bullet-}$ production through photoinduced electron transfer is energetically difficult [26, 27]. However, ultraviolet radiation to reductive photosensitizer, such as NADH (**Figure 4**), can produce $\text{O}_2^{\bullet-}$ as follows [28]:

Figure 2. Autoxidation process of hydroquinone and ROS production.

Figure 3. Autoxidation process of hydrazine analogues and ROS production.

where NADH* is the photoexcited state of NADH and NAD[•] is the radical form. NAD[•] undergoes further oxidation by oxygen molecules to NAD⁺, the final oxidized product. The formed O₂^{•−} is also converted to H₂O₂ through the dismutation process of Eq. (1).

Photocatalytic reaction can also produce H₂O₂ [29–34]. For example, the surface of titanium dioxide (TiO₂) can reduce relatively oxidative molecules under ultraviolet A (UVA; wavelength, 315–400 nm) irradiation [29–32]. Two crystalline forms of TiO₂, anatase and rutile with band gap energies of 3.26 and 3.06 eV, respectively, are well-known semiconducting photocatalyst [29–32]. The adsorbed oxygen molecules on the TiO₂ surface is reduced to O₂^{•−} by the electron of conduction band, which is excited from the valence band by UVA energy (**Figure 5**).

Figure 4. Structure of NADH.

Figure 5. Photocatalytic production of ROS by TiO_2 .

Similarly to the abovementioned reaction, $\text{O}_2^{\bullet-}$ is also converted to H_2O_2 through the dismutation process of Eq. (1). In addition, oxidation reaction of TiO_2 photocatalyst also produces H_2O_2 . The formed hole (h^+) in the valence band by UVA irradiation oxidizes water molecules on the surface of TiO_2 to $\bullet\text{OH}$. The reaction of two $\bullet\text{OH}$ species can produce H_2O_2 as follows:

Although TiO_2 particles are barely incorporated into cell nucleus [35], cellular DNA damage was reported [36–39]. Because H_2O_2 has a transparency for nuclear membrane, the cellular DNA damage can be explained by H_2O_2 -mediated mechanism [32]. The activation of H_2O_2 and DNA damage by H_2O_2 are described later.

2.3. Secondary formation of hydrogen peroxide through photocatalytic reaction

Photocatalytic reaction can produce oxidized intermediates other than final oxidized products of chemical compounds. For example, photooxidized amino acids [40] and sugars [41] by TiO_2 photocatalyst produce H_2O_2 through secondary oxidation reaction in the presence of metal ions (**Figure 6**). Titanium dioxide can photocatalyze the production of $\bullet\text{OH}$, a strong oxidant, through the decomposition of H_2O . The formed h^+ in the valence band by UVA irradiation can also oxidize various materials adsorbed on TiO_2 surface. Hydroxyl radicals and h^+ can oxidize these biomolecules, resulting in the production of oxidized intermediates. The formation of partly oxidized molecules leads to the secondary H_2O_2 production in the presence of metal ions. This H_2O_2 production process may cause a remote H_2O_2 generation in cells.

It has been reported that the photooxidized phenylalanine and tyrosine by TiO_2 produce H_2O_2 in the presence of copper(II) ion [40]. Since TiO_2 photocatalysis induces a hydroxylation of

Figure 6. The secondary production of ROS by TiO_2 photocatalysis.

aromatic compounds [42], the formation of benzenediol derivatives from these aromatic amino acids is possible (**Figure 7**). As mentioned above, hydroquinone can produce H_2O_2 through the autooxidation (**Figure 2**) [20, 43]. The amount of H_2O_2 production from the photooxidized phenylalanine is significantly larger than that from tyrosine [40]. The difference of their autooxidation rates should affect the H_2O_2 production. It has been reported that the autooxidation rate of 1,4-form of benzenediol is markedly faster than that of 1,2-form [20]. Phenylalanine can be oxidized into various types of benzenediol, including the 1,4-form; however, tyrosine cannot be converted to the 1,4-form. Consequently, phenylalanine can produce relatively large amount of secondary H_2O_2 through the photocatalysis of TiO_2 . Furthermore, other amino acids can be also oxidized by TiO_2 photocatalyst and induce the secondary ROS production. Specifically, photocatalyzed cysteine by anatase form produces significantly large amount of secondary H_2O_2 . In the case of sugar oxidation by TiO_2 photocatalyst, the activity of secondary H_2O_2 production by anatase form TiO_2 is larger than that of rutile form [41].

Figure 7. Tyrosine oxidation by TiO_2 photocatalysis and the copper ion mediated ROS production. Phenylalanine is also oxidized by the similar processes, leading to the secondary ROS production.

3. DNA damage by hydrogen peroxide

Hydrogen peroxide itself barely induces DNA damage; however, it can oxidize nucleobases and cleave sugar-phosphate backbone in the presence of metal ions. In this section, the sequence-specific DNA damage by the H₂O₂-derived ROS and its biological effect are briefly introduced.

3.1. Sequence-specific DNA damage by hydrogen peroxide

Hydrogen peroxide causes alkali-labile products at guanine, thymine, and cytosine in the presence of copper ion (Cu²⁺) [44]. Since copper ions are associated with chromatin [45] to form stable complexes with DNA [46–49], Cu²⁺ can play an important role in the activation of H₂O₂ in cell nucleus. Polyacrylamide gel electrophoresis studies demonstrated that H₂O₂ itself cannot cleave and oxidize DNA [44]. However, the incubation of DNA with H₂O₂ and Cu²⁺ induce base modifications at guanine, thymine, and cytosine residues. These base modification sites can be cleaved by hot piperidine treatment [20–22, 44]. The derived reactive species from H₂O₂, for example, copper-peroxyl species (Cu(I)-OOH), are responsible for this DNA damage:

Cu(I)-OOH is not strongly reactive compared with •OH; however, its lifetime is relatively long to induce DNA base modification. Single-stranded DNA is easier oxidized by these ROS. Therefore, DNA damage by H₂O₂ is enhanced by denaturation of DNA [44]. Abovementioned chemical compounds, benzenediol [20] and hydrazine [21], induce these base modification in the presence of Cu²⁺. In the case of relatively low concentration of TiO₂ particles, similar sequence-specific DNA damage was observed after UVA irradiation with Cu²⁺ [32]. DNA damage mediated by H₂O₂ is effectively inhibited by catalase [50], which is an enzyme to decompose H₂O₂ to H₂O and O₂. Chelating molecules for copper ions also effectively suppress this DNA damage. In addition, 3-methylthiopropional (methional) is an effective inhibitor of Cu(I)-OOH [20, 32, 44]. Cu(I)-OOH cannot be scavenged by free •OH scavengers, such as sugars and alcohols [20, 22, 32, 44]. In the presence of Cu²⁺, UVA-irradiated NADH also induces DNA damage by the similar process through H₂O₂ production [28]. In general, photosensitized DNA damage could be explained by ¹O₂ formation mechanism or electron transfer-mediated oxidation [51]. The H₂O₂-mediated DNA is a rare case in the photochemical DNA damage.

Hydrogen peroxide and Cu²⁺ can induce tandem lesion at guanine and thymine residues [32]. Clustered DNA lesions including tandem damage have important mutagenic potential [52–54]. Furthermore, the repair of such DNA damage is more difficult than single-base damage [55–60]. Therefore, oxidative DNA damage through H₂O₂ production may play an important role in carcinogenesis.

In the presence of iron ions (Fe^{2+}), $\cdot\text{OH}$ is formed as follows:

Formed $\cdot\text{OH}$ induces base oxidation with non-sequence specificity, because $\cdot\text{OH}$ can oxidize all nucleobases [44, 61]. In addition, direct cleavage of sugar-phosphate backbone is caused by $\cdot\text{OH}$. Hydroxyl radical-mediated DNA damage was reported by the case of ascorbate with Cu^{2+} [62]. As mentioned above, in the case of TiO_2 photocatalysis, $\cdot\text{OH}$ is directly produced from water decomposition [29–32], and DNA damage without sequence specificity can be induced in the absence of metal ions [32]. Relatively high concentration of anatase form of TiO_2 induce non-sequence-specific DNA damage under UVA irradiation without metal ions through $\cdot\text{OH}$ production [32]. DNA damage by $\cdot\text{OH}$ is effectively inhibited by sugars and alcohols [32, 44]. However, in the presence of metal ions, the addition of $\cdot\text{OH}$ scavengers rather enhances DNA damage through the secondary generation of H_2O_2 from the oxidized products of scavengers themselves by $\cdot\text{OH}$ [32, 41]. Base modifications can cause carcinogenesis. Because H_2O_2 can penetrate into nuclear membrane, DNA modification can be induced by H_2O_2 originally formed in the sphere of outer cell nucleus through the assistance of metal ions.

3.2. Mutagenicity and cytotoxicity caused by hydrogen peroxide production

As oxidized products of nucleobases by the H_2O_2 -mediated mechanism, 8-oxo-7,8-dihydroguanine (8-oxo-G; oxidized guanine, **Figure 8**) [63–65]; 5,6-dihydroxy-5,6-dihydrothymine (OH-thy; oxidized thymine, **Figure 9**) [58, 66, 67]; 5-hydroxyuracil (OH-Ura; oxidized cytosine, **Figure 9**) [67, 68]; 5-hydroxyhydantoin (OH-Hyd; oxidized cytosine, **Figure 9**) [68], and 5-hydroxycytosine (OH-Cys; oxidized cytosine, **Figure 9**) [67] are well-known compounds.

Figure 8. Guanine oxidation by ROS. This scheme is an example of the guanine oxidation by $^1\text{O}_2$ to 8-oxo-G. Other H_2O_2 -derived ROS, $\cdot\text{OH}$ and $\text{Cu}(\text{I})\text{-OOH}$, also produce 8-oxo-G through the oxidation of guanine.

Figure 9. Oxidized products of thymine and cytosine by $^1\text{O}_2$.

In a certain case, oxidative DNA damage induces cell death [69, 70]. As a minor oxidized product of guanine, 2,6-diamino-4-hydroxy-5-formamidopyrimidine (Fapy-G, **Figure 10**) can be formed by H_2O_2 and metal ions [63, 71]. Mutagenicity of Fapy-G is low [72]; however, a related product, methyl-Fapy-G formation, is a lethal lesion [73]. Furthermore, a theoretical study suggested that the formation of Fapy-G contributes to mutation [74]. Cytotoxicity of TiO_2 photocatalyst can be explained by oxidative damage of membrane protein [75–77]. In addition, cellular DNA damage was also reported [78, 79]. Because H_2O_2 has a transparency for nuclear membrane, the cellular DNA damage by TiO_2 photocatalysis can be explained by H_2O_2 production. The formed H_2O_2 through TiO_2 photocatalysis is incorporated into cell nucleus and activated by endogenous metal ions, leading to oxidative DNA damage [32]. Examples of the mutations caused by the oxidized guanines are described in Section 4.

Figure 10. Structure of Fapy-G.

4. Singlet oxygen

In general, the production mechanism of $^1\text{O}_2$ involves photochemical processes. Various photooxidation processes can be explained by $^1\text{O}_2$ production. In this section, the production mechanism of $^1\text{O}_2$, its application, and biomolecule oxidation by $^1\text{O}_2$ are briefly introduced.

4.1. General property of singlet oxygen

Singlet oxygen is an excited state of $^3\text{O}_2$, ground triplet state of molecular oxygen [16–18]. In general, singlet excited (S_1) states of O_2 are $^1\Delta_g$ and $^1\Sigma_g^+$; they have excitation energy of 0.98 eV and 1.63 eV above $^3\text{O}_2$, respectively [16–18]. Because of the short lifetime of $^1\Sigma_g^+$ (a few picoseconds), $^1\Delta_g$, the lower S_1 state of O_2 , plays an important role in various oxidation reactions. In this chapter, $^1\Delta_g$ is denoted throughout as $^1\text{O}_2$. The highest occupied molecular orbital (HOMO) of $^3\text{O}_2$ is a semi-occupied molecular orbital (SOMO), whereas this molecular orbital of $^1\text{O}_2$ becomes the lowest unoccupied molecular orbital (LUMO) (**Figure 1B**). The oxidative activity of $^1\text{O}_2$ is stronger than that of $^3\text{O}_2$ due to the vacant molecular orbital. Commonly, $^1\text{O}_2$ is produced through photosensitized reaction. Since the excitation energy of $^1\text{O}_2$ is relatively small, which corresponds to the energy of photon with the wavelength of 1270 nm (smaller than that of visible light photon), photoexcited states of various dyes can sensitize the generation of $^1\text{O}_2$ under visible light or ultraviolet irradiation. Various molecules become photosensitizer (PST) to generate $^1\text{O}_2$. In general, the photosensitized reaction of $^1\text{O}_2$ generation is an electron exchange energy transfer (the Dexter mechanism) [80]. These processes are presented as follows:

where $\text{PST}^*(S_1)$ and $\text{PST}^*(T_1)$ are the S_1 and T_1 states of PST, respectively. In general, since the lifetime of $\text{PST}^*(T_1)$ is markedly longer (several microseconds) than that of $\text{PST}^*(S_1)$ (several nanoseconds), $^1\text{O}_2$ is produced by $\text{PST}^*(T_1)$. However, the formation of $^1\text{O}_2$ by $\text{PST}^*(S_1)$ is not impossible. The lifetime of $^1\text{O}_2$ (τ_Δ) is relatively long (**Table 1**). Generated $^1\text{O}_2$ can oxidize various materials, including biomolecules, within its long lifetime. The τ_Δ strongly depends on the surroundings, and a solvent deuterium effect on the reactivity of $^1\text{O}_2$ is significant (**Table 1**). For example, the τ_Δ in deuterium oxide (D_2O) is markedly longer than that in H_2O , and the biomolecule oxidation by $^1\text{O}_2$ is significantly enhanced in D_2O compared with that in H_2O .

Solvent	Photosensitizer	$\tau_{\Delta}/\mu\text{s}$	Reference
Water (H_2O)	Cationic porphyrin	3.5	[81]
	Rose bengal	3.77	[82]
Phosphate buffer (pH 7.6)	P(V) porphyrin	3.5	[83]
Ethanol ($\text{C}_2\text{H}_5\text{OH}$)	Rose bengal	15.4	[82]
Ethanol/ H_2O (1/1)	Rose bengal	6.37	[82]
Water (D_2O)	Berberine with DNA	72	[84]
	Methylene blue	32	[85]
	Phenalenone	64.4	[86]
	Tris(bipyridine)Ru(II)	59.47	[82]
Chloroform (CHCl_3)	Phenalenone	232	[86]
Tetrachloromethane (CCl_4)	Phenalenone	34,000	[86]

Table 1. Solvent dependence of the lifetime of singlet oxygen.

4.2. Photodynamic therapy

One of the most important medicinal applications of $^1\text{O}_2$ is photodynamic therapy (PDT) (**Figure 11**) [7–9]. Photodynamic therapy is a promising and less invasive treatment for cancer [7–9] and photosterilization [10–14]. For cancer PDT, in general, porphyrins are used for photosensitizers, for example, porfimer sodium [87] and talaporfin sodium [88]. Photosterilization, antimicrobial PDT, is also carried out using dyes, for example, methylene blue (MB) [11, 14, 89]. The important mechanism of PDT processes including photosterilization is oxidation of biomolecules of cancer cell or bacteria through $^1\text{O}_2$ production under visible light irradiation. Visible light, especially longer wavelength visible light (wavelength > 650 nm), is less harmful for the human body and can penetrate into the tissue deeply. As mentioned above, $^1\text{O}_2$ can be generated by longer wavelength visible light. Administered photosensitizers, porphyrins, or other dyes produce $^1\text{O}_2$ through energy transfer to oxygen molecules with relatively large quantum yield (Φ_{Δ}).

Figure 11. Scheme of the general procedure of PDT.

4.3. Photocatalytic singlet oxygen generation

As mentioned above, TiO_2 photocatalyzes the generation of various ROS. Singlet oxygen can be also produced through the photocatalysis of TiO_2 [31, 90–96]. In general, photogenerated electron in the conduction band reduces the surface-adsorbed oxygen molecules to $\text{O}_2^{\bullet-}$. Through the reoxidation of $\text{O}_2^{\bullet-}$, $^1\text{O}_2$ is formed. The possible reactions of photocatalytic $^1\text{O}_2$ productions are as follows:

and

The photogenerated h^+ in the valence band and $\bullet\text{OH}$ can act as the oxidants to produce $^1\text{O}_2$. In addition, hydroperoxyl radical ($\bullet\text{OOH}$) generated from $\text{O}_2^{\bullet-}$ and H^+ also produces $^1\text{O}_2$ as follows:

The reported values of Φ_{Δ} are depending on the experimental condition, for example, around 0.2 (0.2, Degussa P25 in water [92], and 0.22, rutile particle in chloroform [95]). Other cases reported relatively small values, for example, 0.003 [96] and 0.02 [94]. In the cases of airborne $^1\text{O}_2$, quite small value (10^{-8} – 10^{-9}) was reported [93]. It has been reported that the τ_{Δ} value of $^1\text{O}_2$ produced by Degussa P25 aqueous suspension is 5 μs [92]. Other photocatalytic materials, for example, zinc oxide (ZnO) can photocatalyze $^1\text{O}_2$ production through the similar reaction of TiO_2 photocatalysis [97]. Recently, carbon quantum dots, which have been paid attention as interesting nano-materials, also photocatalyze $^1\text{O}_2$ production [33].

Singlet oxygen is an important ROS for PDT. Other than $^1\text{O}_2$, H_2O_2 production can be also applied for PDT mechanism. Photocatalytic materials can produce these ROS under photoirradiation. Therefore, application of photocatalysts, specifically TiO_2 nanoparticles, for PDT has been also studied [29, 98–101]. To realize the TiO_2 -utilized PDT, direct administration of small TiO_2 powders into tumor assisted with an optical fiber was proposed [29]. In addition, it was reported that oral-administrated TiO_2 nanoparticles are transported into the tumor of nude mouse skin transplanted from a human prostate cancer cell line [98]. As mentioned above, in general, TiO_2 nanoparticles can be excited by UVA irradiation. To utilize visible light for TiO_2 excitation, upconversion technique was also studied [100].

4.4. DNA oxidation by $^1\text{O}_2$ and mutation

Singlet oxygen can oxidize only guanines without sequence specificity; however, it does not have the ability to induce the oxidation of other nucleobases or to cleave the sugar-phosphate backbone [44]. The main oxidized product of guanine by $^1\text{O}_2$ is 8-oxo-G (**Figure 8**) [63–65]. Guanines undergo the Diels-Alder reaction by photoproducted $^1\text{O}_2$, leading to the formation

of [4 + 2] cycloaddition product with the imidazole ring to produce an endoperoxide. Through the subsequent proton transfer, this peroxide is converted to 8-hydroperoxyguanine [102, 103], which becomes 8-hydroxyguanine [63]. The keto-enol tautomerism produces 8-oxo-G from 8-hydroxyguanine. Because single-stranded DNA is easily oxidized by ROS, 8-oxo-G formation by $^1\text{O}_2$ is increased by DNA denaturation [44]. The 8-oxo-G formation causes DNA misreplication (**Figure 12**), which can lead to mutations such as G-C:T-A transversion caused by the stable base-pair formation between 8-oxo-G and adenine [104, 105]. Since 8-oxo-G is more easily oxidized than guanine, 8-oxo-G undergoes further reaction, leading to the formation of imidazolone and oxazolone (**Figure 13**) [63, 106, 107]. Imidazolone forms more stable base pair with guanine than cytosine [106, 107]. Therefore, guanine oxidation by $^1\text{O}_2$ may cause G-C:C-G transversion [108, 109] through imidazolone formation, a further oxidized product of 8-oxo-G. Indeed, it has been reported that UVA can induce these mutations [110].

4.5. Protein oxidation by $^1\text{O}_2$

Protein oxidation is also induced by $^1\text{O}_2$. The following amino acids, tryptophan, tyrosine, cysteine, histidine, and methionine, can be oxidized by $^1\text{O}_2$ [111]. In the case of tryptophan oxidation by $^1\text{O}_2$, *N*-formylkynurenine (**Figure 14**) is a major oxidized product [112, 113]. The reported reaction rate coefficient between tryptophan and $^1\text{O}_2$ is $3.0 \times 10^7 \text{ s}^{-1} \text{ M}^{-1}$ [114]. Oxidation of tryptophan residue in a certain protein can be examined with a fluorometer [115]. For example, human serum albumin (HSA) has one tryptophan residue, and the intrinsic fluorescence of tryptophan at around 350 nm can be diminished by the oxidative damage. Porphyrin phosphorus(V) complexes (**Figure 15**), of which the Φ_{Δ} is larger than 0.5, can induce oxidative damage to the tryptophan residue of HSA [116]. Photosensitized HSA damage is enhanced in D_2O , in which the lifetime of $^1\text{O}_2$ is markedly elongated compared in H_2O (**Table 1**). Furthermore, sodium azide (NaN_3), a strong physical quencher of $^1\text{O}_2$ [117], effectively suppresses this HSA damage. From the analysis of the effect of NaN_3 on the HSA damage, the contribution of $^1\text{O}_2$ -mediated oxidation to the total quantum yield of protein damage

Figure 12. Hydrogen bonding between 8-oxo-G and adenine.

Figure 13. Structures of imidazolone and oxazolone and the hydrogen bonding between guanine and imidazolone.

Figure 14. Structures of tryptophan and *N*-formylkynurenine, an oxidized product of tryptophan by $^1\text{O}_2$.

Figure 15. Example of P(V)porphyrin photosensitizer.

can be determined [115]. Photosensitized $^1\text{O}_2$ production by porphyrin phosphorus(V) complexes induces the damage of tyrosinase, which is an enzyme to catalyze the hydroxylation of tyrosine, resulting in the deactivation of tyrosinase [118]. Oxidation of the amino acid residue by $^1\text{O}_2$ can cause the deactivation of protein function. The protein oxidation photosensitized by porphyrins through ROS production is an important mechanism of PDT.

Photocatalyzed $^1\text{O}_2$ production by TiO_2 may not play an important role in the oxidation reaction [31, 94]. Formed $^1\text{O}_2$ on the TiO_2 surface is quenched by TiO_2 itself with relatively large quenching rate coefficient (e.g., $2.4 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ [95]). In the presence of bovine serum albumin, $^1\text{O}_2$ produced by TiO_2 photocatalysis is effectively quenched, suggesting the protein oxidation [94]. However, in the case of TiO_2 photocatalyst, other ROS are more important for protein oxidation than $^1\text{O}_2$ -mediated reaction [29–32].

5. Detection of ROS

ROS detection is an important theme to investigate a biological effect of ROS or evaluation of the activity of PDT photosensitizers [119–122]. Fluorometry is one of the most important and effective methods of ROS detection. For example, 5-carboxyfluorescein-based probe has been developed (**Figure 16**) [123]. This probe can detect H_2O_2 in the living cell. As an inexpensive method, the fluorometry using folic acid (**Figure 17**) was reported [23, 119, 124]. Folic acid can be decomposed by H_2O_2 in the presence of Cu^{2+} , resulting in the fluorescence enhancement. The limit of detection (LOD, at signal/noise = 3) for this method was $0.5 \mu\text{M H}_2\text{O}_2$. This method is based on the oxidative decomposition of folic acid by $\text{Cu}(\text{I})\text{-OOH}$. In the presence of Fe^{2+} , $\cdot\text{OH}$ slightly induces the folic acid decomposition; however, the effect of $\cdot\text{OH}$ on this folic acid decomposition is negligibly small because of the very short lifetime [125, 126]. In addition, $\text{O}_2^{\cdot-}$ does not

Figure 16. Structure of 5-carboxyfluorescein-based fluorescence probe for H_2O_2 [123].

Figure 17. Structures of folic acid and methotrexate and the fluorometry of ROS [119, 124].

have the activity of folic acid decomposition. Using folic acid or its analogue, 1O_2 can be also detected [124]. Specifically, in D_2O , folic acid or methotrexate (**Figure 17**), an analogue of folic acid, is effectively decomposed by 1O_2 , resulting in the fluorescence enhancement [124]. Using this method, the values of Φ_Δ of various water-soluble photosensitizers can be determined.

6. Control of singlet oxygen production

Control of photosensitized 1O_2 is an important theme for biology or medicine, for example, to realize target-selective PDT [127] or “theranostics” (therapy and diagnosis) [128]. The pH-dependent control [129] and target-selective control [127, 128, 130–132] methods have been reported. It has been reported that free base porphyrins were synthesized to control their photosensitized 1O_2 generating activity by pH (**Figure 18**) [129]. The S_1 state of this porphyrin

Figure 18. Example of the reported pH-responsive porphyrin [129].

Figure 19. The examples of DNA-targeting porphyrins: Ant-P [81], Pyr-P [130], Phen-P [131], and Nap-Ps [132].

is quenched by the electron-donating moiety in neutral or alkali solution. However, protonation of this electron-donating moiety under acidic condition suppresses the electron transfer, leading to the recovery of the $^1\text{O}_2$ production activity of porphyrin ring. Because cancer cell is slightly a more acidic condition compared with normal cells [133–135], this pH-based control of photosensitized $^1\text{O}_2$ production can be applied to cancer-selective PDT. DNA-targeting control of photosensitized $^1\text{O}_2$ generation has been also reported [127, 128]. For example, electron donor-connecting porphyrins have been studied (**Figure 19**) [81, 130–132]. These compounds can be photoexcited by visible light irradiation, and their S_1 states are effectively quenched through intramolecular electron transfer. The charge-transfer state energy can be raised through the binding interaction with DNA, an anionic polymer, resulting in the inhibition of the intramolecular electron transfer and enhancement of $^1\text{O}_2$ generation.

7. Conclusions

Hydrogen peroxide is easily produced from the oxidation processes of chemical compounds by oxygen molecules. In addition, UVA-irradiated NADH and semiconductor photocatalytic materials can also produce H_2O_2 . Formed H_2O_2 in cells can be incorporated into cell nucleus and activated by endogenous metal ions. Copper ion induces Cu(I)-OOH formation from H_2O_2 ,

whereas $\cdot\text{OH}$ is produced from H_2O_2 and iron ion. These ROS cause base oxidation, and $\cdot\text{OH}$ can induce strand break of DNA. Base modifications lead to carcinogenesis or lethal effect. Photoirradiation to various sensitizing materials induces $^1\text{O}_2$ production. Visible light has sufficient energy to produce $^1\text{O}_2$. Therefore, $^1\text{O}_2$ is easily produced by various dyes under photoirradiation. Photocatalytic $^1\text{O}_2$ formation through reoxidation of $\text{O}_2^{\cdot-}$ is also possible. Formed $^1\text{O}_2$ can oxidize guanine residues of DNA without sequence specificity and several amino acid residues of protein within its lifetime, which depends on the surroundings. Various detection methods of these ROS have been developed. In addition, the target-selective or condition-selective productions of ROS become important strategies for PDT and cancer “theranostics.”

Acknowledgements

These works were partially supported by the Grants-in-Aid for Scientific Research (KAKENHI) from the Japan Society for the Promotion of Science (JSPS).

Author details

Kazutaka Hirakawa^{1,2*}

*Address all correspondence to: hirakawa.kazutaka@shizuoka.ac.jp

1 Applied Chemistry and Biochemical Engineering Course, Department of Engineering, Graduate School of Integrated Science and Technology, Shizuoka University, Hamamatsu, Shizuoka, Japan

2 Department of Optoelectronics and Nanostructure Science, Graduate School of Science and Technology, Shizuoka University, Hamamatsu, Shizuoka, Japan

References

- [1] Ziecha D, Francob R, Pappac A, Panayiotidis MI. Reactive oxygen species (ROS)-induced genetic and epigenetic alterations in human carcinogenesis. *Mutation Research Fundamental and Molecular Mechanisms of Mutagenesis*. 2011;**711**:167-173. DOI: 10.1016/j.mrfmmm.2011.02.015
- [2] Wu Q, Ni X. ROS-mediated DNA methylation pattern alterations in carcinogenesis. *Current Drug Targets*. 2015;**16**:13-19. DOI: 10.2174/1389450116666150113121054
- [3] Kawanishi S, Ohnishi S, Ma N, Hiraku Y, Murata M. Crosstalk between DNA damage and inflammation in the multiple steps of carcinogenesis. *International Journal of Molecular Sciences*. 2017;**18**:1808. DOI: 10.3390/ijms18081808
- [4] Auten RL, Davis JM. Oxygen toxicity and reactive oxygen species: The devil is in the details. *Pediatric Research*. 2009;**66**:121-127. DOI: 10.1203/PDR.0b013e3181a9eafb

- [5] Liochev SI. Reactive oxygen species and the free radical theory of aging. *Free Radical Biology and Medicine*. 2013;**60**:1-4. DOI: 10.1016/j.freeradbiomed.2013.02.011
- [6] Dixon SJ, Stockwell BR. The role of iron and reactive oxygen species in cell death. *Nature Chemical Biology*. 2014;**10**:9-17. DOI: 10.1038/nchembio.1416
- [7] Dolmans DEJGJ, Fukumura D, Jain RK. Photodynamic therapy for cancer. *Nature Reviews Cancer*. 2003;**3**:380-387. DOI: 10.1038/nrc1071
- [8] Castano AP, Mroz P, Hamblin MR. Photodynamic therapy and anti-tumour immunity. *Nature Reviews Cancer*. 2006;**6**:535-545. DOI: 10.1038/nrc1894
- [9] Chilakamarthi U, Giribabu L. Photodynamic therapy: Past, present and future. *The Chemical Records*. 2017;**17**:1-29. DOI: 10.1002/tcr.201600121
- [10] Arenas Y, Monro S, Shi G, Mandel A, McFarland S, Lilge L. Photodynamic inactivation of *Staphylococcus aureus* and methicillin-resistant *Staphylococcus aureus* with Ru(II)-based type I/type II photosensitizers. *Photodiagnosis and Photodynamic Therapy*. 2013;**10**:615-625. DOI: 10.1016/j.pdpdt.2013.07.001
- [11] Diogo P, Gonçalves T, Palma P, Santos JM. Photodynamic antimicrobial chemotherapy for root canal system asepsis: A narrative literature review. *International Journal of Dentistry*. 2015;**2015**:269205. DOI: 10.1155/2015/269205
- [12] Oruba Z, Łabuz P, Macyk W, Chomyszyn-Gajewska M. Antimicrobial photodynamic therapy-A discovery originating from the pre-antibiotic era in a novel periodontal therapy. *Photodiagnosis and Photodynamic Therapy*. 2015;**12**:612-618. DOI: 10.1016/j.pdpdt.2015.10.007
- [13] Tim M. Strategies to optimize photosensitizers for photodynamic inactivation of bacteria. *Journal of Photochemistry and Photobiology B: Biology*. 2015;**150**:2-10. DOI: 10.1016/j.jphotobiol.2015.05.010
- [14] Wainwright M, McLean A. Rational design of phenothiazinium derivatives and photoantimicrobial drug discovery. *Dyes and Pigments*. 2017;**136**:590-600. DOI: 10.1016/j.dyepig.2016.09.015
- [15] Jajic I, Sarna T, Strzalka K. Senescence, stress, and reactive oxygen species. *Plants*. 2015;**4**:393-411. DOI: 10.3390/plants4030393
- [16] DeRosa MC, Crutchley RJ. Photosensitized singlet oxygen and its applications. *Coordination Chemistry Reviews*. 2002;**233-234**:351-371. DOI: 10.1016/S0010-8545(02)00034-6
- [17] Schweitzer C, Schmidt R. Physical mechanisms of generation and deactivation of singlet oxygen. *Chemical Reviews*. 2003;**103**:1685-1758. DOI: 10.1021/cr010371d
- [18] Ogilby PR. Singlet oxygen: There is indeed something new under the sun. *Chemical Society Reviews*. 2010;**39**:3181-3209. DOI: 10.1039/B926014P
- [19] Wong HS, Dighe PA, Mezera V, Monternier PA, Brand MD. Production of superoxide and hydrogen peroxide from specific mitochondrial sites under different bioenergetic conditions. *Journal of Biological Chemistry*. 2017;**292**:16804-16809. DOI: 10.1074/jbc.R117.789271

- [20] Hirakawa K, Oikawa S, Hiraku Y, Hirosawa I, Kawanishi S. Catechol and hydroquinone have different redox properties responsible for their differential DNA-damaging ability. *Chemical Research in Toxicology*. 2002;**15**:76-82. DOI: 10.1021/tx010121s
- [21] Ito K, Yamamoto K, Kawanishi S. Manganese-mediated oxidative damage of cellular and isolated DNA by isoniazid and related hydrazines: Non-Fenton-type hydroxyl radical formation. *Biochemistry*. 1992;**31**:11606-11613
- [22] Hirakawa K, Midorikawa K, Oikawa S, Semi KS. Carcinogenic semicarbazide induces sequence-specific DNA damage through the generation of reactive oxygen species and the derived organic radicals. *Mutation Research Genetic Toxicology and Environmental Mutagenesis*. 2003;**536**:91-101. DOI: 10.1016/S1383-5718(03)00030-5
- [23] Hirakawa K. Fluorometry of hydrogen peroxide using oxidative decomposition of folic acid. *Analytical and Bioanalytical Chemistry*. 2006;**386**:244-248. DOI: 10.1007/s00216-006-0649-1
- [24] Marcus RA. On the theory of oxidation-reduction reactions involving electron transfer. I. *The Journal of Chemical Physics*. 1956;**24**:966-978. DOI: 10.1063/1.1742723
- [25] Marcus RA, Sutin N. Electron transfers in chemistry and biology. *Biochimica et Biophysica Acta*. 1985;**811**:265-322. DOI: 10.1016/0304-4173(85)90014-X
- [26] Kikuchi K, Sato C, Watabe M, Ikeda H, Takahashi Y, Miyashi Y. New aspects on fluorescence quenching by molecular oxygen. *Journal of the American Chemical Society*. 1993;**115**:5180-5184. DOI: 10.1021/ja00065a033
- [27] Sato C, Kikuchi K, Okamura K, Takahashi Y, Miyashi T. New aspects on fluorescence quenching by molecular oxygen. 2. Inhibition of long-distance electron transfer in acetonitrile. *The Journal of Physical Chemistry*. 1995;**99**:16925-16931. DOI: 10.1021/j100046a018
- [28] Ito K, Hiraku Y, Kawanishi S. Photosensitized DNA damage induced by NADH: Site specificity and mechanism. *Free Radical Research*. 2007;**41**:461-468. DOI: 10.1080/10715760601145240
- [29] Fujishima A, Rao TN, Tryk DA. Titanium dioxide photocatalysis. *Journal of Photochemistry and Photobiology C: Photochemistry Reviews*. 2000;**1**:1-21. DOI: 10.1016/S1389-5567(00)00002-2
- [30] Liu K, Cao M, Fujishima A, Jiang L. Bio-inspired titanium dioxide materials with special wettability and their applications. *Chemical Reviews*. 2014;**114**:10044-10094. DOI: 10.1021/cr4006796
- [31] Nosaka Y, Nosaka AY. Generation and detection of reactive oxygen species in photocatalysis. *Chemical Reviews*. 2017;**117**:11302-11336. DOI: 10.1021/acs.chemrev.7b00161
- [32] Hirakawa K, Mori M, Yoshida M, Oikawa S, Kawanishi S. Photo-irradiated titanium dioxide catalyzes site specific DNA damage via generation of hydrogen peroxide. *Free Radical Research*. 2004;**38**:439-447. DOI: 10.1080/1071576042000206487

- [33] Yen YC, Lin CC, Chen PY, Ko WY, Tien TR, Lin KJ. Green synthesis of carbon quantum dots embedded onto titanium dioxide nanowires for enhancing photocurrent. *Royal Society Open Science*. 2017;**4**:161051. DOI: 10.1098/rsos.161051
- [34] Donat F, Corbel S, Alem H, Pontvianne S, Balan L, Medjahdi G, Schneider R. ZnO nanoparticles sensitized by CuInZn_xS_{2+x} quantum dots as highly efficient solar light driven photocatalysts. *The Beilstein Journal of Nanotechnology*. 2017;**8**:1080-1093. DOI: 10.3762/bjnano.8.110
- [35] Cai R, Hashimoto K, Itoh K, Kubota Y, Fujishima A. Photokilling of malignant cells with ultra-fine TiO₂ powder. *Bulletin of the Chemical Society of Japan*. 1991;**64**:1268-1273. DOI: [org/10.1246/bcsj.64.1268](http://dx.doi.org/10.1246/bcsj.64.1268)
- [36] Wamer WG, Yin JJ, Wei RR. Oxidative damage to nucleic acids photosensitized by titanium dioxide. *Free Radical Biology and Medicine*. 1997;**23**:851-858. DOI: 10.1016/S0891-5849(97)00068-3
- [37] Meena R, Rani M, Pal P, Rajamani P. Nano-TiO₂-induced apoptosis by oxidative stress-mediated DNA damage and activation of p53 in human embryonic kidney cells. *Applied Biochemistry and Biotechnology*. 2012;**167**:791-808. DOI: 10.1007/s12010-012-9699-3
- [38] Nakagawa Y, Wakuri S, Sakamoto K, Tanaka N. The photogenotoxicity of titanium dioxide particles. *Mutation Research Genetic Toxicology and Environmental Mutagenesis*. 1997;**394**:125-132. DOI: 10.1016/S1383-5718(97)00126-5
- [39] Kashige N, Kakita Y, Nakashima Y, Miake F, Watanabe K. Mechanism of the photocatalytic inactivation of *Lactobacillus casei* phage PL-1 by titania thin film. *Current Microbiology*. 2001;**42**:184-189. DOI: 10.1007/s002840010201TiO2
- [40] Hirakawa K, Suzuki T. Amino acids photocatalyzed by titanium dioxide can produce secondary hydrogen peroxide. *Trends in Photochemistry and Photobiology*. 2014;**16**:63-69
- [41] Hirakawa K. Titanium dioxide photocatalyzes DNA damage via the secondary generation of hydrogen peroxide in the presence of sugars. *Trends in Photochemistry and Photobiology*. 2012;**14**:69-73
- [42] Emeline AV, Zhang X, Murakami T, Fujishima A. Activity and selectivity of photocatalysts in photodegradation of phenols. *Journal of Hazardous Materials*. 2012;**211-212**:154-160. DOI: 10.1016/j.jhazmat.2011.11.078
- [43] Hirakawa K, Sano S. Platinum nanoparticle catalyst scavenges hydrogen peroxide generated from hydroquinone. *Bulletin of the Chemical Society of Japan*. 2009;**82**:1299-1303. DOI: 10.1246/bcsj.82.1299
- [44] Kawanishi S, Hiraku Y, Oikawa S. Mechanism of guanine-specific DNA damage by oxidative stress and its role in carcinogenesis and aging. *Mutation Research Reviews in Mutation Research*. 2001;**488**:65-76. DOI: 10.1016/S1383-5742(00)00059-4
- [45] Agarwal K, Sharma A, Talukder G. Effects of copper on mammalian cell components. *Chemico-Biological Interactions*. 1989;**69**:1-16. DOI: 10.1016/0009-2797(89)90094-X

- [46] Bach D, Miller IR. Polarographic investigation of binding of Cu⁺⁺ and Cd⁺⁺ by DNA. *Biopolymers*. 1967;**5**:161-172. DOI: 10.1002/bip.1967.360050204
- [47] Bryan SE, Frieden E. Interaction of copper(II) with deoxyribonucleic acid below 30 degrees. *Biochemistry*. 1967;**6**:2728-2734. DOI: 10.1021/bi00861a012
- [48] Stoewe R, Prutz WA. Copper-catalyzed DNA damage by ascorbate and hydrogen peroxide: Kinetics and yield. *Free Radical Biology and Medicine*. 1987;**3**:97-105. DOI: 10.1016/S0891-5849(87)80003-5
- [49] Prutz WA, Butler J, Land EJ. Interaction of copper(I) with nucleic acids. *International Journal of Radiation Biology*. 1990;**58**:215-234. DOI: 10.1080/09553009014551581
- [50] Glorieux C, Calderon PB. Catalase, a remarkable enzyme: targeting the oldest antioxidant enzyme to find a new cancer treatment approach. *Biological Chemistry*. 2017;**398**:1095-1108. DOI: 10.1515/hsz-2017-0131
- [51] Hirakawa K. DNA damage through photo-induced electron transfer and photosensitized generation of reactive oxygen species. In: Kimura H, Suzuki A, editors. *New Research on DNA Damage*. Nova Science Publishers Inc., New York; 2008. pp. 197-219. ISBN: 978-1-60456-581-2
- [52] Gentil A, Le Page F, Cadet J, Sarasin A. Mutation spectra induced by replication of two vicinal oxidative DNA lesions in mammalian cells. *Mutation Research Fundamental and Molecular Mechanisms of Mutagenesis*. 2000;**452**:51-56. DOI: 10.1016/S0027-5107(00)00034-8
- [53] Kalam MA, Basu AK. Mutagenesis of 8-oxoguanine adjacent to an abasic site in simian kidney cells: Tandem mutations and enhancement of GfT transversions. *Chemical Research in Toxicology*. 2005;**18**:1187-1192. DOI: 10.1021/tx050119r
- [54] Yuan B, Jiang Y, Wang Y, Wang Y. Efficient formation of the tandem thymine glycol/8-oxo-7,8-dihydroguanine lesion in isolated DNA and the mutagenic and cytotoxic properties of the tandem lesions in *Escherichia coli* cells. *Chemical Research in Toxicology*. 2010;**23**:11-19. DOI: 10.1021/tx9004264
- [55] Chaudhry MA, Weinfeld M. The action of *Escherichia coli* endonuclease III on multiply damaged sites in DNA. *Journal of Molecular Biology*. 1995;**249**:914-922. DOI: 10.1006/jmbi.1995.0348
- [56] Venkhataraman R, Donald CD, Roy R, You HJ, Doetsch PW, Kow YW. Enzymatic processing of DNA containing tandem dihydrouracil by endonucleases III and VIII. *Nucleic Acids Research*. 2001;**29**:407-414. DOI: 10.1093/nar/29.2.407
- [57] Budworth H, Dianova II, Podust VN, Dianov GL. Repair of clustered DNA lesions. Sequence-specific inhibition of long patch base excision repair by 8-oxoguanine. *Journal of Biological Chemistry*. 2002;**277**:21300-21305. DOI: 10.1074/jbc.M201918200
- [58] Budworth H, Dianov GL. Mode of inhibition of short-patch base excision repair by thymine glycol within clustered DNA lesions. *Journal of Biological Chemistry*. 2003;**278**:9378-9381. DOI: 10.1074/jbc.M212068200

- [59] Lomax ME, Cunniffe S, O'Neill P. 8-OxoG retards the activity of the ligase III/XRCC1 complex during the repair of a single-strand break, when present within a clustered DNA damage site. *DNA Repair*. 2004;**3**:289-299. DOI: 10.1016/j.dnarep.2003.11.006
- [60] Eot-Houllier G, Eon-Marchais S, Gasparutto D, Sage E. Processing of a complex multiply damaged DNA site by human cell extracts and purified repair proteins. *Nucleic Acids Research*. 2005;**33**:260-271. DOI: 10.1093/nar/gki165
- [61] Oikawa S, Kawanishi S. Distinct mechanisms of site-specific DNA damage induced by endogenous reductants in the presence of iron(III) and copper(II). *Biochimica et Biophysica Acta*. 1998;**1399**:19-30. DOI: 10.1016/S0167-4781(98)00092-X
- [62] Kobayashi S, Ueda K, Morita J, Sakai H, Komano T. DNA damage induced by ascorbate in the presence of Cu²⁺. *Biochimica et Biophysica Acta*. 1998;**949**:143-147. DOI: 10.1016/0167-4781(88)90065-6
- [63] Burrows CJ, Muller JG. Oxidative nucleobase modifications leading to strand scission. *Chemical Reviews*. 1998;**98**:1109-1151. DOI: 10.1021/cr960421s
- [64] Kasai H, Yamaizumi Z, Berger M, Cadet J. Photosensitized formation of 7,8-dihydro-8-oxo-2'-deoxyguanosine (8-hydroxy-2'-deoxyguanosine) in DNA by riboflavin: A non singlet oxygen-mediated reaction. *Journal of the American Chemical Society*. 1992;**114**: 9692-9694. DOI: 10.1021/ja00050a078
- [65] Cullis PM, Malone ME, Merson-Davies LA. Guanine radical cations are precursors of 7,8-dihydro-8-oxo-2'-deoxyguanosine but are not precursors of immediate strand breaks in DNA. *Journal of the American Chemical Society*. 1996;**118**:2775-2788. DOI: 10.1021/ja9536025
- [66] Hayes RC, Petrullo LA, Huang HM, Wallace SS, LeClerc JE. Oxidative damage in DNA. Lack of mutagenicity by thymine glycol lesions. *Journal of Molecular Biology*. 1988;**201**:239-246. DOI: 10.1016/0022-2836(88)90135-0
- [67] D'Ham C, Romieu A, Jaquinod M, Gasparutto D, Cadet J. Excision of 5,6-dihydroxy-5,6-dihydrothymine, 5,6-dihydrothymine, and 5-hydroxycytosine from defined sequence oligonucleotides by *Escherichia coli* endonuclease III and Fpg proteins: Kinetic and mechanistic aspects. *Biochemistry*. 1999;**38**:3335-3344. DOI: 10.1021/bi981982b
- [68] Samson-Thibault F, Madugundu GS, Gao S, Cadet J, Wagner JR. Profiling cytosine oxidation in DNA by LC-MS/MS. *Chemical Research in Toxicology*. 2012;**25**:1902-1911. DOI: 10.1021/tx300195f
- [69] Dizdaroglu M. Oxidatively induced DNA damage and its repair in cancer. *Mutation Research Reviews in Mutation Research*. 2015;**763**:212-245. DOI: 10.1016/j.mrrev.2014.11.002
- [70] Kasai H. What causes human cancer? Approaches from the chemistry of DNA damage. *Genes Environment*. 2016;**38**:19. DOI: 10.1186/s41021-016-0046-8
- [71] Delaney S, Jarem DA, Volle CB, Yennie CJ. Chemical and biological consequences of oxidatively damaged guanine in DNA. *Free Radical Research*. 2012;**46**:420-441. DOI: 10.3109/10715762.2011.653968

- [72] Tudek B. Imidazole ring-opened DNA purines and their biological significance. *Journal of Biochemistry and Molecular Biology*. 2003;**36**:12-19. DOI: 10.5483/BMBRep.2003.36.1.012
- [73] Asagoshi K, Terato H, Ohyama Y, Ide H. Effects of a guanine-derived formamidopyrimidine lesion on DNA replication: Translesion DNA synthesis, nucleotide insertion, and extension kinetics. *Journal of Biological Chemistry*. 2002;**277**:14589-14597. DOI: 10.1074/jbc.M200316200
- [74] Hirakawa K, Yoshida M. Theoretical study of the effects of amino acids on one-electron oxidation of a nucleobase: Adenine residue can be a hole-trapping site. *Pure and Applied Chemical Sciences*. 2014;**2**:41-48. DOI: 10.12988/pacs.2014.424
- [75] Cai R, Hashimoto K, Kubota Y, Fujishima A. Increment of photocatalytic killing of cancer cells using TiO₂ with the aid of superoxide dismutase. *Chemistry Letters*. 1992:427-430. DOI: 10.1246/cl.1992.427
- [76] Cai R, Kubota Y, Shuin T, Sakai H, Hashimoto K, Fujishima A. Induction of cytotoxicity by photoexcited TiO₂ particles. *Cancer Research*. 1992;**52**:2346-2348
- [77] Kubota Y, Shuin T, Kawasaki C, Hosaka M, Kitamura H, Cai R, Sakai H, Hashimoto K, Fujishima A. Photokilling of T-24 human bladder cancer cells with titanium dioxide. *British Journal of Cancer*. 1994;**70**:1107-1111
- [78] Dunford R, Salinaro A, Cai L, Serpone N, Horikoshi S, Hidaka H, Knowland J. Chemical oxidation and DNA damage catalysed by inorganic sunscreen ingredients. *FEBS Letters*. 1997;**418**:87-90. DOI: 10.1016/S0014-5793(97)01356-2
- [79] Petković J, Küzma T, Rade K, Novak S, Filipič M. Pre-irradiation of anatase TiO₂ particles with UV enhances their cytotoxic and genotoxic potential in human hepatoma HepG2 cells. *Journal of Hazardous Materials*. 2011;**196**:145-152. DOI: 10.1016/j.jhazmat.2011.09.004
- [80] Dexter DL. A theory of sensitized luminescence in solids. *The Journal of Chemical Physics*. 1953;**21**:836-850. DOI: 10.1063/1.1699044
- [81] Hirakawa K, Nishimura Y, Arai T, Okazaki S. Singlet oxygen generating activity of an electron donor-connecting porphyrin photosensitizer can be controlled by DNA. *The Journal of the Physical Chemistry B*. 2013;**117**:13490-13496. DOI: org/10.1021/jp4072444
- [82] Ohara K, Kikuchi K, Origuchi T, Nagaoka S. Singlet oxygen quenching by trolox C in aqueous micelle solutions. *Journal of Photochemistry and Photobiology B: Biology*. 2009;**97**:132-137. DOI: 10.1016/j.jphotobiol.2009.08.010
- [83] Hirakawa K, Azumi K, Nishimura Y, Arai T, Nosaka Y, Okazaki S. Photosensitized damage of protein by fluorinated diethoxyphosphorus(V)porphyrin. *Journal of Porphyrins and Phthalocyanines*. 2013;**17**:56-62. DOI: 10.1142/S1088424612501258
- [84] Hirakawa K, Hirano T, Nishimura Y, Arai T, Nosaka Y. Dynamics of singlet oxygen generation by DNA-binding photosensitizers. *The Journal of the Physical Chemistry B*. 2012;**116**:3037-3044. DOI: 10.1021/jp300142e

- [85] Matheson IBC, Lee J, King AD. The lifetime of singlet oxygen ($^1\Delta_g$) in heavy water, a revised value. *Chemical Physics Letters*. 1978;**55**:49-51. DOI: 10.1016/0009-2614(78)85129-X
- [86] Shimizu O, Watanabe J, Imakubo K, Naito S. Absolute quantum yields and lifetimes of photosensitized phosphorescence of singlet oxygen O_2 ($^1\Delta_g$) in air-saturated aqueous and organic solutions of phenalenone. *Chemistry Letters*. 1999;**28**:67-68. DOI: 10.1246/cl.1999.67
- [87] Moghissi K, Dixon K, Stringer M, Thorpe JA. Photofrin PDT for early stage oesophageal cancer: Long term results in 40 patients and literature review. *Photodiagnosis and Photodynamic Therapy*. 2009;**6**:159-166. DOI: 10.1016/j.pdpdt.2009.07.026
- [88] Wang S, Bromley E, Xu L, Chen JC, Keltner L. Talaporfin sodium. *Expert Opinion on Pharmacotherapy*. 2010;**11**:133-140. DOI: 10.1517/14656560903463893
- [89] Hirakawa K, Ishikawa T. Phenothiazine dyes photosensitize protein damage through electron transfer and singlet oxygen generation. *Dyes and Pigments*. 2017;**142**:183-188. DOI: 10.1016/j.dyepig.2017.03.035
- [90] Konaka R, Kasahara E, Dunlap WC, Yamamoto Y, Chien KC, Inoue M. Irradiation of titanium dioxide generates both singlet oxygen and superoxide anion. *Free Radical Biology and Medicine*. 1999;**27**:294-300. DOI: 10.1016/S0891-5849(99)00050-7
- [91] Konaka R, Kasahara E, Dunlap WC, Yamamoto Y, Chien KC, Inoue M. Ultraviolet irradiation of titanium dioxide in aqueous dispersion generates singlet oxygen. *Redox Report*. 2001;**6**:319-325. DOI: 10.1179/135100001101536463
- [92] Nosaka Y, Daimon T, Nosaka AY, Murakamia Y. Singlet oxygen formation in photocatalytic TiO_2 aqueous suspension. *Physical Chemistry Chemical Physics*. 2004;**6**:2917-2918. DOI: 10.1039/B405084C
- [93] Naito K, Tachikawa T, Cui S, Sugimoto A, Fujitsuka M, Majima T. Single-molecule detection of airborne singlet oxygen. *Journal of the American Chemical Society*. 2006;**128**:16430-16431. DOI: 10.1021/ja066739b
- [94] Hirakawa K, Hirano T. Singlet oxygen generation photocatalyzed by TiO_2 particles and its contribution to biomolecule damage. *Chemistry Letters*. 2006;**35**:832-833. DOI: 10.1246/cl.2006.832
- [95] Li W, Gandra N, Courtney SN, Gao R. Singlet oxygen production upon two-photon excitation of TiO_2 in chloroform. *Chemphyschem*. 2009;**10**:1789-1793. DOI: 10.1002/cphc.200900155
- [96] Buchalska M, Łabuz P, Bujak Ł, Szewczyk G, Sarna T, MaćKowski S, Macyk W. New insight into singlet oxygen generation at surface modified nanocrystalline TiO_2 —The effect of near-infrared irradiation. *Dalton Transactions*. 2013;**42**:9468-9475. DOI: 10.1039/C3DT50399B
- [97] Dunlap WC, Yamamoto Y, Inoue M, Kashiba-Iwatsuki M, Yamaguchi M, Tomita K. Uric acid photo-oxidation assay: In vitro comparison of sunscreens agents. *International Journal of Cosmetic Science*. 1998;**20**:1-18. DOI: 10.1046/j.1467-2494.1998.171731.x

- [98] Miyoshi N, Kume K, Tsutumi K, Fukunaga Y, Ito S, Imamura Y, Bibin AB. Application of titanium dioxide (TiO₂) nanoparticles in photodynamic therapy (PDT) of an experimental tumor. *AIP Conference Proceedings*. 2011;**1415**:21. DOI: 10.1063/1.3667210
- [99] Zhang H, Shan Y, Dong L. A comparison of TiO₂ and ZnO nanoparticles as photosensitizers in photodynamic therapy for cancer. *Journal of Biomedical Nanotechnology*. 2014;**10**:1450-1457. DOI: 10.1166/jbn.2014.1961
- [100] Lucky SS, Muhammad Idris N, Li Z, Huang K, Soo KC, Zhang Y. Titania coated upconversion nanoparticles for near-infrared light triggered photodynamic therapy. *ACS Nano*. 2015;**9**:191-205. DOI: 10.1021/nn503450t
- [101] Jukapli NM, Bagheri S. Recent developments on titania nanoparticle as photocatalytic cancer cells treatment. *Journal of Photochemistry and Photobiology B: Biology*. 2016;**163**:421-430. DOI: 10.1016/j.jphotobiol.2016.08.046
- [102] Shu C, Kang P, Khan S, Foote CS. Low-temperature photosensitized oxidation of a guanosine derivative and formation of an imidazole ring-opened product. *Journal of the American Chemical Society*. 2002;**124**:3905-3913. DOI: 10.1021/ja011696e
- [103] Kang P, Foote CS. Formation of transient intermediates in low-temperature photosensitized oxidation of an 8-(13)C-guanosine derivatives. *Journal of the American Chemical Society*. 2002;**124**:4865-4873. DOI: 10.1021/ja012038x
- [104] Sugiyama H, Saito I. Theoretical studies of GG-specific photocleavage of DNA via electron transfer: Significant lowering of ionization potential and 5'-localization of HOMO of stacked GG bases in B-form DNA. *Journal of the American Chemical Society*. 1996;**118**:7063-7068. DOI: 10.1021/ja9609821
- [105] Yoshioka Y, Kitagawa Y, Takano Y, Yamaguchi K, Nakamura T, Saito I. Experimental and theoretical studies on the selectivity of GGG triplets toward one-electron oxidation in B-form DNA. *Journal of the American Chemical Society*. 1999;**121**:8712-8719. DOI: 10.1021/ja991032t
- [106] Kino K, Saito I, Sugiyama H. Product analysis of GG-specific photooxidation of DNA via electron transfer: 2-aminoimidazolone as a major guanine oxidation product. *Journal of the American Chemical Society*. 1998;**120**:7373-7374. DOI: 10.1021/ja980763a
- [107] Kino K, Sugiyama H. Possible cause of GC→CG transversion mutation by guanine oxidation product, imidazolone. *Chemistry and Biology*. 2001;**8**:369-378. DOI: 10.1016/S1074-5521(01)00019-9
- [108] McBride TJ, Schneider JE, Floyd RA, Loeb LA. Mutation induced by methylene blue plus light in single-stranded M13mp2. *Proceedings of the National Academy of Sciences of the United States of America*. 1992;**89**:6866-6870
- [109] Negishi K, Hao W. Spectrum of mutations in single-stranded DNA phage M13mp2 exposed to sunlight: Predominance of G-to-C transversion. *Carcinogenesis*. 1992;**13**:1615-1618. DOI: 10.1093/carcin/13.9.1615

- [110] Drobeta EA, Turcotte J, Chateaneuf A. A role for ultraviolet A in solar mutagenesis. *Proceedings of the National Academy of Sciences of the United States of America*. 1995;**92**:2350-2354
- [111] Michaeli A, Feitelson J. Reactivity of singlet oxygen toward amino acids and peptides. *Photochemistry and Photobiology*. 1994;**59**:284-289. DOI: 10.1111/j.1751-1097.1994.tb05035.x
- [112] Ehrenshaft M, Silva SO, Perdivara I, Bilski P, Sik RH, Chignell CF, Tomer KB, Mason RP. Immunological detection of N-formylkynurenine in oxidized proteins. *Free Radical Biology and Medicine*. 2009;**4**:1260-1266. DOI: 10.1016/j.freeradbiomed.2009.01.020
- [113] Thomas AH, Serrano MP, Rahal V, Vicendo P, Claparols C, Oliveros E, Lorente C. Tryptophan oxidation photosensitized by pterin. *Free Radical Biology and Medicine*. 2013;**63**:467-475. DOI: 10.1016/j.freeradbiomed.2013.05.044
- [114] Jensen RL, Arnbjerg J, Ogilby PR. Reaction of singlet oxygen with tryptophan in proteins: A pronounced effect of the local environment on the reaction rate. *Journal of the American Chemical Society*. 2012;**134**:9820-9826. DOI: 10.1021/ja303710m
- [115] Hirakawa K, Umemoto H, Kikuchi R, Yamaguchi H, Nishimura Y, Arai T, Okazaki S, Segawa H. Determination of singlet oxygen and electron transfer mediated mechanisms of photosensitized protein damage by phosphorus(V)porphyrins. *Chemical Research in Toxicology*. 2015;**28**:262-267. DOI: 10.1021/tx500492w
- [116] He XM, Carter DC. Atomic structure and chemistry of human serum albumin. *Nature*. 1992;**358**:209-215. DOI: 10.1038/358209a0
- [117] Li MY, Cline CS, Koker EB, Carmichael HH, Chignell CF, Bilski P. Quenching of singlet molecular oxygen ($^1\text{O}_2$) by azide anion in solvent mixtures. *Photochemistry and Photobiology*. 2001;**74**:760-764. DOI: 10.1562/0031-8655(2001)0740760QOSMOO2.0.CO2
- [118] Ouyang D, Hirakawa K. Photosensitized enzyme deactivation and protein oxidation by axial-substituted phosphorus(V) tetraphenylporphyrins. *Journal of Photochemistry and Photobiology B: Biology*. 2017;**175**:125-131. DOI: 10.1016/j.jphotobiol.2017.08.036
- [119] Hirakawa K. Using folic acids to detect reactive oxygen species. In: Taylor JC. editors. *Advances in Chemistry Research*. Volume 26. Nova Science Publishers Inc., New York; 2015. pp. 111-126. ISBN: 978-1-63463-630-8
- [120] Garcia-Diaz M, Huang YY, Hamblin MR. Use of fluorescent probes for ROS to tease apart Type I and Type II photochemical pathways in photodynamic therapy. *Methods*. 2016;**109**:158-166. DOI: 10.1016/j.ymeth.2016.06.025
- [121] Kalyanaraman B, Hardy M, Podsiadly R, Cheng G, Zielonka J. Recent developments in detection of superoxide radical anion and hydrogen peroxide: Opportunities, challenges, and implications in redox signaling. *Archives Biochemistry and Biophysics*. 2017;**617**:38-47. DOI: 10.1016/j.abb.2016.08.021

- [122] Guo H, Aleyasin H, Dickinson BC, Haskew-Layton RE, Ratan RR. Recent advances in hydrogen peroxide imaging for biological applications. *Cell & Bioscience*. 2014;**4**:64. DOI: 10.1186/2045-3701-4-64
- [123] Abo M, Urano Y, Hanaoka K, Terai T, Komatsu T, Nagano T. Development of a highly sensitive fluorescence probe for hydrogen peroxide. *Journal of the American Chemical Society*. 2011;**133**:10629-10637. DOI: 10.1021/ja203521e
- [124] Hirakawa K. Fluorometry of singlet oxygen generated via a photosensitized reaction using folic acid and methotrexate. *Analytical and Bioanalytical Chemistry*. 2009;**393**:999-1005. DOI: 10.1007/s00216-008-2522-x
- [125] Land EJ, Ebert M. Pulse radiolysis studies of aqueous phenol. Water elimination from dihydroxycyclohexadienyl radicals to form phenoxyl. *Transactions of the Faraday Society*. 1967;**63**:1181-1190. DOI: 10.1039/TF9676301181
- [126] Liao JC, Roider J, Jay DG. Chromophore-assisted laser inactivation of proteins is mediated by the photogeneration of free radicals. *Proceedings of the National Academy of Sciences of the United States of America*. 1994;**91**:2659-2663
- [127] Tørring T, Helmig S, Ogilby PR, Gothelf KV. Singlet oxygen in DNA nanotechnology. *Accounts of Chemical Research*. 2014;**47**:1799-1806. DOI: 10.1021/ar500034y
- [128] Hirakawa K. Control of fluorescence and photosensitized singlet oxygen-generating activities of porphyrins by DNA: Fundamentals for “theranostics”. In: Yilmaz Y, editor. *Phthalocyanines and Some Current Applications*. InTechOpen, London; 2017. p. 169-188. ISBN: 978-953-51-3255-4. DOI: 10.5772/67882
- [129] Horiuchi H, Kuribara R, Hirabara A, Okutsu T. pH-Response optimization of amino-substituted tetraphenylporphyrin derivatives as pH-activatable photosensitizers. *The Journal of Physical Chemistry A*. 2016;**120**:5554-5561. DOI: 10.1021/acs.jpca.6b05019
- [130] Hirakawa K, Harada M, Okazaki S, Nosaka Y. Controlled generation of singlet oxygen by a water-soluble meso-pyrenylporphyrin photosensitizer through interaction with DNA. *Chemical Communications*. 2012;**48**:4770-4772. DOI: 10.1039/c2cc30880k
- [131] Hirakawa K, Ito Y, Yamada T, Okazaki S. Relaxation process of the photoexcited state and singlet oxygen generating activity of water-soluble meso-phenanthrylporphyrin in a DNA microenvironment. *Rapid Communication in Photoscience*. 2014;**3**:81-84. DOI: 10.5857/RCP.2014.3.4.81
- [132] Hirakawa K, Taguchi M, Okazaki S. Relaxation process of photoexcited meso-naphthylporphyrins while interacting with DNA and singlet oxygen generation. *The Journal of Physical Chemistry B*. 2015;**119**:13071-13078. DOI: 10.1021/acs.jpcc.5b08025
- [133] Kuin A, Aalders M, Lamfers M, van Zuidam DJ, Essers M, Beijnen JH, Smets LA. Potentiation of anti-cancer drug activity at low intratumoral pH induced by the mitochondrial inhibitor m-iodobenzylguanidine (MIBG) and its analogue benzylguanidine (BG). *British Journal of Cancer*. 1999;**79**:793-801. DOI: 10.1038/sj.bjc.6690127

- [134] Gupta SC, Singh R, Asters M, Liu J, Zhang X, Pabbidi MR, Watabe K, Mo YY. Regulation of breast tumorigenesis through acid sensors. *Oncogene*. 2016;**35**:4102-4111. DOI: 10.1038/onc.2015.477
- [135] Shi R, Huang L, Duan X, Sun G, Yin G, Wang R, Zhu JJ. Selective imaging of cancer cells with a pH-activatable lysosome-targeting fluorescent probe. *Analytica Chimica Acta*. 2017;**988**:66-73. DOI: 10.1016/j.aca.2017.07.055