

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



***Taraxacum* Genus: Potential Antibacterial and Antifungal Activity**

María Eugenia Martínez Valenzuela,
Katy Díaz Peralta, Lorena Jorquera Martínez and
Rolando Chamy Maggi

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.71619>

Abstract

Plants have been used in traditional medicine for centuries as antibacterial and antifungal agents. *Taraxacum* spp., commonly known as dandelion, is a well-known herbal remedy with a long history; however, limited scientific information is available to explain its traditional use. This review aims to provide current information and a general overview of the available literature concerning the antibacterial and antifungal properties of the *Taraxacum* genus to support its potential as a powerful herbal medicine. Though *Taraxacum* has demonstrated that it is capable of inhibiting the growth of a wide range of bacteria and fungi, the technical aspects of methodology lack standardization, and, therefore, the overall results of processing are difficult to compare between studies. Phytochemical composition and antimicrobial activity in *Taraxacum* are neither directly related, nor does the published data provide sufficient information for identifying the group of unique extraction conditions that are optimal against specific microorganisms. Antimicrobial research indicates that this plant is a promising species for treating several common infections in humans, animals, and plants.

Keywords: antimicrobial, antifungal, ethnopharmacology, extraction, *Taraxacum* species

1. Introduction

Plants have been used in traditional medicine for centuries due to the synthesis of several molecules that provide antibacterial and antifungal properties, the majority of which probably evolved as defenses against infection or predation [1]. The medicinal potential of many plants is still largely unexplored. Among the estimated 250,000–500,000 plant species, a relatively small percentage have been investigated phytochemically and the fraction submitted for

biological or pharmacological screening is even smaller [2]; approximately 20% of the plant species in the world have been investigated for these properties [3]. In this context, dandelion serves as an interesting species with which to unify decades-old information regarding its biological potential against diverse microorganisms. This review gathers the existing results to advance the search for products that could strengthen the domestication and mass production of this plant.

The *Taraxacum* spp. commonly called dandelion is an herbaceous perennial plant of the *Asteraceae* (Compositae) family. This common weed is found worldwide, though originally introduced from Eurasia, and can be found growing in parks, gardens, pastures, orchards, roadsides, vegetable gardens, and among agricultural and horticultural crops [4]. Primarily used as food, the role of *Taraxacum* in traditional medicine was mentioned during ancient times by the Greek physician Dioscorides in the first century and during the renaissance by monks in Cyprus [5]. This plant has been used to treat cystitis, liver and gastric ailments, hepatic and renal detoxification, diabetes, as an anti-inflammatory and anticarcinogenic agent, and, to a lesser extent, as an antimicrobial and antiviral agent, as described in several reviews [6, 7]. Ethnopharmacologically, its use as an antimicrobial agent has been known worldwide among varying cultures, though it has always been administered as a cataplasm (poultice) or infusion. The traditional antimicrobial uses of *Taraxacum* worldwide are displayed in **Table 1**.

Asia and Europe have an important historical background regarding the traditional uses of *Taraxacum*, primarily *T. officinale*, *T. mongolicum*, and *T. coreanum*. This traditional knowledge has been the principal reason for studying the potential uses and crop requirements of *Taraxacum*; studies in America remain scarce [18]. Due to the unscientific approach often present in oral traditions, uncertainty surrounds whether *Taraxacum* use effectively treats microbial infection or, instead, treats only the symptoms. Therefore, scientific research is extremely important in avoiding misinterpretation and myths regarding *Taraxacum* or any other plant.

The first antibacterial scientific study for *Taraxacum* was reported a mere 35 years ago [19]. More than a decade later, studies related to *Taraxacum* antimicrobial activity gained significant

Species	Common use	Country	Part used	Consumption	References
<i>T. cyprium</i>	Catarrh and common cold, cough	Greece	Roots, leaves	—	[5]
<i>T. mongolicum</i>	Urinary tract infections	China	Leaf	Infusion	[8]
<i>T. officinale</i>	Malaria	Venezuela	Roots, leaves	Decoction	[9]
	Tuberculosis	Italy	—	—	[10]
	Cough	Italy	—	—	[11]
	Bacterial infection	Mexico	—	—	[12]
	Diuretic	Chikar	Roots	Tonico	[13]
<i>T. panalpinum</i>	Malaria	Portugal	Roots, leaves, juice	—	[14, 15]
<i>T. platycarpum</i>	Pleurodynia	Korea	Leaf, stem	Infusion Brewing	[16, 17]

Table 1. Ethnopharmacological information of *Taraxacum* genus used as an antimicrobial traditional medicine.

relevance as part of an Italian program between the University of Ferrara and the University of Naples for screening medicinal plants [20]. Nowadays, this plant is becoming a promising species in the treatment of several bacterial and fungal diseases due to the results of various antimicrobial-related studies. This chapter seeks to elucidate both the traditional uses and current state of *Taraxacum* in antimicrobial research to determine the potential that this genus has to become an industrial medicinal crop worldwide. Due to the high potential value that could be derived from the use of new technologies and industrial products developed from this type of plant species, the conservation and protection of the crop should be considered and sustainable global production strategies are developed in accordance with assessments of ecological, economic, and social factors.

2. Antimicrobial properties of the *Taraxacum* genus

Literature reviews providing information on the antimicrobial aspects of natural products, which had until now only been considered empirical, have been recently scientifically confirmed as a means of countering the increasing reports of pathogenic microorganisms resistant to synthetic antimicrobial agents. Some plant-derived compounds can control microbial growth, either separately or in association with conventional antimicrobials [21]. Currently, numerous studies seek to improve pathogen prevention by combining the application of medicinal herb extracts with an antibiotic or effective antipathogenic pesticide to reduce the active synthetic ingredient and resistant pathogenic strains.

2.1. *Taraxacum* species tested for antimicrobial properties

Among the *Taraxacum* genus, *T. officinale* is the most frequently reported species, with almost 80% of mentions in documents related to antimicrobial properties (see **Table 2**), followed by *T. mongolicum* and *T. coreanum*, though over 2500 *Taraxacum* species are currently identified [67]. Other, less studied species include *T. platycarpum*, *T. farinosum*, *T. ohwianum*, and *T. phaleratum*; however, the relevance of these species is confined to specific areas (mostly in Asia) in which they grow naturally since they are not deliberately cultivated for medicinal benefit. This indicates that the microbial properties of less than 1% of all *Taraxacum* species discovered have been studied, revealing the enormous research potential of this genus.

2.2. Bacterial and fungi strains tested

Taraxacum extracts have been tested on different bacterial and fungal strains affecting humans, animals, and plants to determine its antimicrobial profile, confirm its traditional usage, and expand its known uses. Antimicrobial agents are categorized based on the spectrum of action, namely “narrow” and “broad” spectrum, which indicates whether its use is specific for certain bacterial strains or active on a wider range. Bacterial infections can result in mild to life-threatening illnesses that require immediate antibiotic intervention. Alternatively, a superficial fungal infection is rarely life-threatening but can have debilitating effects and may spread to other people or become invasive or systemic, resulting in a life-threatening infection. The widespread, and sometimes inappropriate, use of chemical compounds can create antibiotic

Species	Autentification/ Voucher	C: Collected P: Purchase	Zone	Season	Plant part*	Sample manipulation	Ratio	Solvent	Extraction time	Temp.	Agitation	Inhibition activity**	Ref.
<i>T. officinale</i> Wigg.	No/No	NI	NI	NI	Flower	NI	1:10	Acetic acid 10%	1 h	RT	Homog.	+	[22]
<i>T. officinale</i> Wigg.	No/No	C	NI	Yes	Seeds	Grounded	1:10	Acetic acid 10%	1 h	RT	NI	+	[23]
<i>T. officinale</i> Weber	Yes/No	C	Yes	Yes	NI	Dried	NI	Water, ethanol and ethyl acetate	1 h	80°C	Maceration	+	[24]
<i>T. officinale</i>	Yes/Yes	C	Yes	NI	NI	Air-dried	1:14	Acetone	30 min	NI	NI	+	[25]
<i>Taraxacum</i> spp.	No/No	C	Yes	NI	NI	NI	1:10	Dichloromethane	3 h	20°C	Homog.	+	[26]
<i>T. officinale</i>	No/Yes	C	Yes	Yes	NI	NI	NI	Dichloromethane	3 days	NI	Homog.	+	[27]
<i>T. coreanum</i>	No/No	NI	NI	NI	NI	NI	1:3.3	Ethanol 75%	9 h	60°C	Reflux	+	[28]
<i>T. mongolicum</i>	No/No	C	NI	NI	Aerial	Freeze-dried and grounded 20-mesh	1:5	Ethanol 75%	2 days	NI	Soaked	+	[29]
<i>T. officinale</i> Weber	No/Yes	C	Yes	Yes	Root	Dried and grounded	NI	Ethanol 80%	NI	NI	Reflux	+	[20]
<i>T. officinale</i> F. H. Wigg	Yes/Yes	C	NI	NI	Aerial	Air-dried and crushed	1:1	Ethanol 90%	2 days	RT	Intermitent shaking	+	[30]
<i>T. mongolicum</i> H.					NI		1:10	Ethanol 95%	3 h	80°C		+	[31]
<i>T. ohwianum</i>	No/No	NI	NI	NI	NI	Freeze-dried, air-dried (40°C, 24 h), grounded 24-mesh	1:16	Ethanol 95%	24 h	RT (23°C)	Shaking	+	[32]
<i>T. officinale</i>	Yes/No	NI	NI	NI	Leaves	Air-dried	1:5	Ethylacetate	24 h	RT	150 rpm	+	[22]
<i>T. officinale</i> F.H. Wigg.	Yes/Yes	P	NI	NI	Root	Freeze-dried and blended	1:10	Hexane	Overnight	RT	70 rpm	+	[33]

Species	Autentification/ Voucher	C: Collected P: Purchase	Zone	Season	Plant part*	Sample manipulation	Ratio	Solvent	Extraction time	Temp.	Agitation	Inhibition activity**	Ref.
<i>T. officinale</i>	No/No	NI	NI	NI	Leaves	Air-dried 1 month and grounded	1:1.4	Methanol 75%	NI	NI	NI	+	[34]
<i>T. officinale</i> <i>Weber</i>	No/No	C	Yes	NI	Aerial	Air-dried a 40°C (36–48 h) and grounded	1:5	Methanol 80%	1 h	100°C	Reflux	+	[35]
<i>T. officinale</i> <i>Weber ex. F.</i> <i>H. Wigg</i>	No/Yes	NI	NI	NI	Leaves	NI	1:4	Methanol	5 days	RT	NI	+	[36]
<i>T. officinale</i> <i>Weber</i>	No/No	C	NI	NI	Aerial	Dry under shade and ground	1:10	Methanol	3 weeks	25°C	Homog.	+	[37]
<i>T.</i> <i>platycarpum</i>	No/No	NI	NI	NI	NI	Dried	NI	Methanol	3 h	80°C	NI	+	[38]
<i>T.</i> <i>platycarpum</i>	No/No	NI	NI	NI	NI	Dried	NI	Methanol	3 h	80°C	NI	+	[39]
<i>T. officinale</i>					NI		NI	Methanol	16 h	50°C		+	[40]
<i>T. officinale</i>	No/No	C	NI	NI	Leaves	Dried under shade and grounded	1:2.5	Methanol	24 h	37°C	120 rpm	+	[41]
<i>T.</i> <i>mongolicum</i> <i>Hand-</i> <i>Mazz</i>	Yes/Yes	C	Yes	Yes	NI	Air-dried and grounded	NI	Water	3 h	100°C	By boiling	+	[36]
<i>T. officinale</i>					NI		1:05	Water	NI	NI	Homog.	+	[42]
<i>T. officinale</i>	No/No	NI	NI	NI	NI	Dried at 25– 30°C for 1 week., ground with a mortar	1:20	Water	24 h	35°C	Shaking	+	[43]
<i>T. officinale</i> <i>F.H. (Webb)</i>	No/No	C/P	NI	NI	Root	Cleaning prior freeze-dried, grounded	1:10	Water	3 h	RT	170 rpm	+	[44]

Species	Autentification/ Voucher	C: Collected P: Purchase	Zone	Season	Plant part*	Sample manipulation	Ratio	Solvent	Extraction time	Temp.	Agitation	Inhibition activity**	Ref.
<i>T. officinale</i>	No/No	NI	NI	NI	NI	NI	1:04	Water	45 min	100°C	NI	+	[45]
<i>T. officinale</i> Weber ex Wigger	No/No	NI	NI	NI	leaves	Grounded	1:01	Water	5 min	NI	NI	+	[46]
<i>T. mongolicum</i>	No/No	NI	NI	NI	NI	Grounded	NI	Water	1 h	100°C	By boiling	+	[47]
<i>T. officinale</i>	No/No	NI	NI	NI	NI	Dried at 60°C × 2 h and grounded 60- mesh	1:10	Water	NI	NI	NI	+	[48]
<i>T. officinale</i> H.	NI/NI	NI	NI	NI	NI	NI						+	[49]
<i>T. officinale</i> F.H. Wigg	NI/NI	C	Yes	Yes	Honey	NI	NI	NI	NI	NI	NI	+	[50]
<i>T. farinosum</i> Hausskn. & Bornm	NI/NI	C	NI	Yes	Root	NI	NI	NI	NI	NI	NI	+	[51]
<i>T. officinale</i>	Yes/No	C	Yes	Yes	NI	Dried 40°C × 5 days and grounded	NI	NI	NI	NI	Reflux	+	[52]
<i>T. officinale</i>	NI/NI	NI	NI	NI	NI	NI	NI	NI	NI	NI	NI	+W	[53]
<i>T. officinale</i>	No/No	P	NI	NI	NI	NI	NI	Ethanol 35%	NI	NI	NI	+W	[54]
<i>T. platycarpum</i>	No/No	P	NI	NI	NI	Grounded 50- mesh	1:10	Ethanol	24 h	RT	Homog.	+W	[55]
<i>Taraxacum</i> sp.	No/No	NI	NI	NI	Aerial	Grounded	1:10	Water	4 h	100°C	By boiling	+W	[56]
<i>T. officinale</i> F.H. Wigg.	No/No	C	Yes	Yes	Aerial	Frozen, cut and grounded	1:01	Ethanol 20%	24 h	RT	NI	–	[57]
<i>T. officinale</i>	No/No	NI	NI	NI	Leaves	Dried	NI	Ethanol 40%	NI	NI	NI	–	[58]

Species	Autentification/ Voucher	C: Collected P: Purchase	Zone	Season	Plant part*	Sample manipulation	Ratio	Solvent	Extraction time	Temp.	Agitation	Inhibition activity**	Ref.
<i>T. officinale</i>	No/No	Extract (P)	NI	NI	NI	Diluted	NI	ethanol 45%	NI	NI	NI	–	[59]
<i>T. phaleratum</i> G. Hagl et Rech	Yes/Yes	C	Yes	NI	Aerial	Air-dried and grounded	NI	Ethanol 70%	NI	RT	NI	–	[60]
<i>T. officinale</i> Cass.	No/No	P	NI	NI	Root	Dried	1:04	Ethanol	24 h	NI		–	[61]
<i>T. officinale</i>	No/No	C	Yes	NI	flower	Chopped and frozen	NI	Methanol 90%	30 min	4°C	Homog.	–	[62]
<i>T. officinale</i> Weber	Yes/Yes	NI	NI	NI	NI	Dried and grounded	1:40	Methanol	Overnight	RT	NI	–	[63]
<i>T. mongolicum</i> Hand- Mazz	No/Yes	C	NI	NI	Whole	NI	1:10	Water	Overnight	NI	Homog.	–	[64]
<i>T. officinale</i>	No/No	P	NI	NI	Root	Grounded	1:8.3	Water	30 min	100°C	By boiling	–	[65]
<i>T. officinale</i>	No/No	C	Yes	Yes	Leafs, roots	NI	1:03	Water		RT	Homog.	–	[66]

*NI, No indicated.

Table 2. Physical parameters on *Taraxacum* extracts for testing antimicrobial activity.

resistance. Due to this issue, the potential of *Taraxacum* as a useful, broad-spectrum antimicrobial and antifungal agent that can be “easily and worldwide grown,” is highly valuable. A list of the strains against which *Taraxacum*’s antimicrobial activity has been tested is displayed in **Table 3**.

Bacterial strains	Fungi strains
<i>Aeromonas hydrophila</i> (–) [22]	<i>Alternaria alternata</i> (+) [46, 68]
<i>Agrobacterium tumefaciens</i> (+) [24]	<i>Aspergillus carbonarius</i> (+) [35]
<i>Bacillus cereus</i> (+) [22, 33, 36, 44] (–) [66]	<i>A. niger</i> (+) [23, 35, 37, 68, 69] (–) [27, 66]
<i>B. pumilus</i> (–) [66]	<i>A. flavus</i> (+) [37] (–) [66]
<i>B. subtilis</i> (+) [20, 24, 25, 27, 29, 34, 38, 39, 41, 48, 69] (–) [37, 64, 66]	<i>A. fumigatus</i> (+) [37] (–) [66]
<i>Campylobacter jejuni</i> (+) [54, 59]	<i>Bipolaris sorokiniana</i> (+) [23, 67] (–) [68]
<i>Chromobacterium violaceum</i> (+) [66] (–) [65]	<i>Botrytis cinerea</i> (+) [23, 35, 67]
<i>Clavibacter michiganense</i> (+) [69]	<i>Candida albicans</i> (+) [27, 34, 36, 52] (–) [55, 57, 66]
<i>Cupriavidus</i> sp. (–) [66]	<i>C. glabrata</i> (–) [55, 66]
<i>Enterobacter coccus</i> (–) [37]	<i>C. krusei</i> (–) [66]
<i>Enterococcus faecalis</i> (+) [53] (–) [37, 66]	<i>C. parapsilesis</i> (–) [55, 66]
<i>Erwinia carotovora</i> (+) [24]	<i>C. utilis</i> (–) [55]
<i>Escherichia coli</i> (+) [22, 24, 25, 27, 29, 34, 36, 38, 39, 41, 43, 45, 47, 48, 58, 70] (–) [20, 32, 33, 37, 44, 57, 62, 64, 66]	<i>C. tropicalis</i> (+) [55]
<i>Helicobacter pylori</i> (+) [31, 54]	<i>Cladosporium herbarum</i> (+) [71]
<i>Klebsiella aerogenes</i> (–) [66]	<i>Cochliobolus sativus</i> (+) [68]
<i>K. pneumoniae</i> (+) [29, 36] (–) [20, 37, 45, 66]	<i>Colletotrichum gloeosporoides</i> (–) [68]
<i>Listeria monocytogenes</i> (+) [38, 39] (–) [66]	<i>C. lagenarium</i> (+) [42]
<i>Micrococcus kristinae</i> (+) [25]	<i>Cryptococcus neoformans</i> (+) [36]
<i>M. luteus</i> (+) [37, 41]	<i>Exophiala (Wangiella) dermatitidis</i> (–) [66]
<i>Mycobacterium aurum</i> (–) [63]	<i>Fusarium avenaceum</i> (+) [68]
<i>M. bovis</i> (–) [63]	<i>F. graminearum</i> (–) [69]
<i>M. smegmatis</i> (–) [63]	<i>F. oxysporum</i> (+) [23, 56, 69]
<i>M. tuberculosis</i> (–) [60]	<i>Microsporum canis</i> (+) [51]
<i>Propionihacterium acnes</i> (+) [49]	<i>Monilinia laxa</i> (+) [35]
<i>Proteus mirabilis</i> (+) [43] (–) [20]	<i>Mucor piriformis</i> (+) [46]
<i>P. vulgaris</i> (+) [25, 29] (–) [70]	<i>Penicillium</i> sp. (–) [66]
<i>Pseudomonas</i> sp. (–) [50]	<i>P. digitatum</i> (+) [35]
<i>P. aeruginosa</i> (+) [24, 27, 29, 36, 41, 49, 70] (–) [20, 37, 57, 64, 66]	<i>P. expansum</i> (+) [26, 46] (–) [35]
	<i>P. italicum</i> (+) [35]

Bacterial strains	Fungi strains
<i>P. fluorescens</i> (+) [24]	<i>Ph. betae</i> (+) [23, 68]
<i>P. syringae</i> (+) [69]	<i>Phytophthora infestans</i> (+) [69]
<i>Serratia/Rahnella</i> sp. (–) [66]	<i>Pityrosporum ovale</i> (+) [49]
<i>Salmonella typhimurium</i> (+) [36] (–) [33, 44]	<i>Pythium debaryanum</i> (+) [69]
<i>S. abony enterica</i> (+) [58]	<i>Rhizoctonia solani</i> (+) [37, 56]
<i>S. poona</i> (–) [66]	<i>Saccharomyces cerevisiae</i> (+) [34]
<i>S. typhi</i> (+) [44, 51] (–) [20]	<i>Saprolegnia australis</i> (–) [61]
<i>Sarcina lutea</i> (+) [24]	<i>Scedosporium apiospermum</i> (–) [66]
<i>Serratia marcescens</i> (+) [25] (–) [66]	<i>Trichophyton longifusus</i> (+) [51]
<i>Shigella flexeri</i> (–) [70]	<i>T. mentagrophytes</i> (+) [27]
<i>S. sonnei</i> (+) [36]	<i>Verticillium albo-atrum</i> (+) [23] (–) [68]
<i>Staphylococcus aureus</i> (+) [22, 24, 25, 28, 29, 32–34, 36, 38, 39, 41, 43–45, 48–52, 70] (–) [20, 27, 37, 57, 58, 62, 64, 66]	
<i>S. epidermidis</i> (+) [28] (–) [66]	
<i>Streptococcus haemolyticus</i> (+) [20]	
<i>S. agalactiae</i> (+) [47]	
<i>S. dysgalactiae</i> (+) [47]	
<i>Vibrio cholerae</i> (+) [37]	
<i>V. parahaemolyticus</i> (+) [38, 39]	
<i>Xanthomonas campestris</i> (+) [69]	

(+) Extracts of *Taraxacum* active against the pathogen; (–) extracts of *Taraxacum* inactive against the pathogen.

Table 3. Bacterial and fungal strains on which *Taraxacum* extracts have been tested.

2.2.1. Human pathogens

In the study of antibacterial properties of these plants, most attention has been focused on human pathogenic strains, including *S. aureus*, *E. faecalis*, *V. cholerae*, *B. subtilis*, *P. aeruginosa*, *K. pneumonia*, and *E. coli*. These are the pathogens commonly responsible for infections in gastrointestinal and massive organ systems such as the lungs and skin. *Taraxacum officinale* is the species generally studied to combat these pathogens, but it has demonstrated diverse results depending on the extraction characteristics or the bioassay performed. For instance, a methanolic extract of *T. officinale* at 0.2 mg/mL was as effective as an antibacterial agent against *M. luteus* and *V. cholera* with minimum inhibitory concentration (MIC) values of 1.0 and 12.5 mg/mL, respectively, but displayed no activity against *S. aureus*, *E. faecalis*, *E. bacter*, *V. cholerae*, *B. subtilis*, *P. aeruginosa*, *K. pneumonia*, or *E. coli* [37]. In the same study, the inhibition percentages achieved for mycelial growth of *A. niger*, *A. flavus*, *A. fumigatus*, and *R. solani* were 37, 71, 85, and 78%, respectively. Other works indicate that methanolic *T. officinale* leaf extracts ranging from 0.003 to 0.5 mg/mL were active against *S. aureus*, *P. aeruginosa*, *B. cereus*, *S. sonnei*,

S. enterica serovar *typhimurium*, *E. coli*, *K. pneumonia*, *C. albicans*, and *C. neoformans* with MIC values ranging from 0.04 to 5.0 mg/mL [36]. A similar extract at 10 mg/mL displayed moderate growth diameter inhibition for *S. typhi*, but was highly active for *S. aureus*, *B. cereus*, and *E. coli*, even when no activity was observed for *A. hydrophila* [22]. Ethanolic extracts of 2.0 mg/mL were active against *A. aureus*, MRSA clinical, and *B. cereus*, with MIC values between 0.38 and 0.5 mg/mL, but were not effective against *E. coli* or *S. typhi*. In the same work, a water extract at the same concentration showed no activity against any strain tested [33]. Moreover, 21 ethanolic extracts from various plants were tested against 20 *Salmonella* serovars. *Taraxacum* inhibited only 5% of these, and was therefore not considered for additional antimicrobial studies [72].

Recently, methanolic and chloroformic leaf extracts of *T. officinale* were found to be effective against *M. luteus*, *P. aeruginosa*, *B. subtilis*, *E. coli*, and *S. aureus* with MIC values of 0.3 mg/mL and no observable activity for water extracts [41]. In this study, the highest impact was noted with methanol and chloroform extracts against *S. aureus* and *E. coli*, respectively, and the lowest with both extracts against *P. aeruginosa*. Furthermore, an ethanolic extract was effective against *E. coli* and *S. aureus*, but no activity was observed for either extract against *K. pneumonia* and *P. aeruginosa* at 50, 100, and 200 mg/mL. Nevertheless, a water extract was effective only for *E. coli* at 100 and 200 mg/mL [45]. Water and ethanolic extracts at 1.0 mg/mL exhibit effective inhibition against *S. aureus* and fewer inhibitory effects were observed for *P. mirabilis*; against *S. aureus*, an ethanolic extract was active at 0.5 mg/mL, but a water extract was only active at 1.0 mg/mL; and inhibition was not achieved for either extracts at 0.1 mg/mL [73]. An ethanolic extract was slightly active against *B. subtilis* and *S. haemolyticus*, but was inactive against other Gram positive and Gram negative strains, resulting in no further studies with this extract [20]. Furthermore, only weak activity was achieved by methanolic extracts of this plant against *P. syringae* [74].

Both ethanolic and water extracts of *T. officinale* were active for *S. marcescens* and *M. kristinae*. The ethanolic extract alone was active on *P. vulgaris*, *E. coli*, *B. subtilis*, and *S. aureus* with MIC values ranging from 1.0 to 7.0 mg/mL for all strains tested [25]. Similar extracts had antimicrobial effects on four species that induce acne (*P. ovale*, *P. acnes*, *P. aeruginosa*, and *S. aureus*) in broth dilution tests with effects depending on the extract concentration, but no further information was available [49]. Moreover, a leaf extract (0.04 mg/well) was reported as a bactericidal agent against *S. aureus* and fungistatic against *C. albicans* [52]. Contrarily, extracts of 130 and 200 mg/mL from aerial parts were unable to prevent the growth of 34 microorganisms from genera *Bacillus*, *Enterobacter*, *Klebsiella*, *Listeria*, *Pseudomonas*, *Salmonella*, *Staphylococcus*, *Aspergillus*, and *Candida*, among others; therefore, it was considered inactive at these concentrations in a disc diffusion assay [66]. A methanolic *T. officinale* flower extract was not active against *E. coli* or *S. aureus* at 1.0 mg/mL in a diffusion agar assay [62] and no activity was found on *S. aureus*, *E. coli*, *P. aeruginosa*, or *C. albicans* using a leaf ethanolic extract when 0.05 mL were placed in sterile discs [57]. Furthermore, an ethanolic extract of leaves displayed no activity against *S. aureus*, *E. coli*, or *S. abony* by the serial dilution method [58], with the same results for root and leaf extracts on *M. aurum* and *M. smegmatis* at 0.5 mg/mL [63].

Raw extracts of *T. officinale* have been widely tested, as well as solvent fractions. In a study in which the methanolic leaf extract was fractioned by different solvents, the methyl chloride,

ethyl acetate, and butanol fractions were active on *E. coli*, *S. aureus*, *B. subtilis*, *C. albicans*, and *S. cerevisiae* at 50 mg/mL, with inhibition percentages ranging from 13 to 76%. The water fraction showed moderate inhibition via the broth dilution method (10 and 14% for *E. coli* and *B. subtilis*, respectively) but no effect on the disc diffusion assay [34]. The only report in which a *Taraxacum* extract was compared to another natural antibacterial substance besides other plants extracts evaluated the use of *T. officinale* extract as an irrigation agent in endodontic treatments against *E. faecalis* in root canal infections. Leaf and root extracts at 0.7% were slightly active but propolis was more effective for this purpose [53]. In the case of commercial preparations, high activity has been reported for a commercial *T. officinale* ethanolic extract, showing antibacterial activity against *H. pylori* at 20 mg/mL with 26% inhibition but no observable activity for *C. jejuni* [54].

Considering other *Taraxacum* species, *T. platycarpum* anticandidal activity was determined against five different *Candida* sp. by agar diffusion assay. An ethanolic extract at 0.2 mg/mL weakly inhibited *C. tropicalis* but no other *Candida* strains [55]. A methanolic extract was active against *B. subtilis*, *S. aureus*, *L. monocytogenes*, *E. coli*, and *V. parahaemolyticus* at concentrations ranging from 0.5 to 2.0 mg/mL, with growth inhibition ranging from 5.1 to 100%, correlating to the concentration. In that study, chloroform, butanol, and ethyl acetate fractions were active in the disc diffusion assay for almost every strain tested, but an aqueous extract was inactive [38, 39].

An ethanolic extract of *T. mongolicum* at 0.2 mg/mL was not able to achieve growth inhibition in a microdilution assay for *B. subtilis*, *S. aureus*, *E. coli*, or *P. aeruginosa* [64]. In contrast, an ethanolic extract of this species was active for *E. coli*, *S. aureus*, and *P. aeruginosa* in the disc diffusion assay with MIC values between 0.05 and 0.1 mg/mL, which was three times higher than the values obtained for erythromycin. However, no activity was achieved for *S. flexneri* or *P. vulgaris* [75]. Another report indicated that only the butanol fraction of an ethanolic extract of this plant was active on *H. pylori*, but water and methyl chloride fractions were inactive. Nevertheless, a different report indicated that a butanol fraction exerted higher inhibition (13%) than the aqueous fraction, possibly due to the flavonoid and luteolin content (28 and 1.1%, respectively) [31]. Against *S. aureus* and *S. epidermis*, an acetyl acetate fraction of an ethanolic *T. coreanum* extract was active at 0.5, 1.0, and 3.0 mg/disc, a chloroform fraction was active at 1.0 and 3.0 mg/disc, and a butanol fraction at 1.0 mg/mL, but displayed no activity against MRSA displayed [28]. An ethanolic *T. ohwianum* extract was active against *E. coli* at 240 and 320 mg/mL, but not against *S. aureus* [32]. These authors indicate that the pH and temperature of the bioassay were important parameters in the antimicrobial performance of the extract. An extract of the aerial parts of *T. phaleratum* was inactive at 0.2 mg/mL against *M. tuberculosis*, even when several solvent fractions were tested [60].

Limited studies have been conducted on humans establishing the antimicrobial potential of *Taraxacum* extracts. Chinese language studies have reported the effects of various formulas containing *T. mongolicum* for medical treatment. An herbal formula known as “fu zheng qu xie” was just as effective as the antibiotic gentamycin in 75 cases of gastric disease caused by *H. pylori*. Furthermore, an herbal formula called “jie du yang gan gao,” which includes *T. mongolicum*, was significantly more effective than another botanical formulation in lowering elevated liver enzymes and curing patients with hepatitis B in a 96-person, double-blind trial [76].

2.2.2. Plant pathogens

Plant extracts have also been tested on bacteria and fungi that affect fruits and vegetables, causing rot diseases during postharvest handling, to find an alternative to chemical pesticides, which are harmful to the environment and human health. An aqueous *T. officinale* root extract (S) at different dilutions (S, S/2 to S/100) caused significant inhibition to mycelial growth in *A. alternata* (70% for S to 17% for S/100), *P. expansum* (67% for S to 5.3% for S/100), and *M. piriformis* (70% for S to 16% for S/100) [46]. In the case of *R. solani* and *C. sativus*, a *Taraxacum* acetyl acetate extract at a concentration of 100 mg/mL exhibited a weak effect on the growth of these plant pathogens and no inhibition of *F. oxysporum* [22]. A methanolic extract of *Taraxacum* at 0.2 mg/mL was not effective against *A. niger*, *A. flavus*, *A. fumigates*, or *R. solani* [37]. A methanolic extract of *Taraxacum* sp. displayed weak activity against *C. sativus*, *F. oxysporum*, and *R. solani* at 5 mg/disc and a water extract displayed no activity at all [56].

A *T. officinale* hydro-methanolic extract tested the inhibition of conidial germination and inhibition of germ tube elongation for several plant pathogens at several dilutions (0.25×, 0.5×, and 0.75×) using a microassay method on slides. Dilution at 0.75× showed inhibition of conidial germination values of 2, 3, 4, 9, 11, and 12% for *P. italicum*, *A. niger*, *A. carbonarius*, *B. cinerea*, *M. laxa*, and *P. digitatum*, respectively. For these same strains, excluding *A. carbonarius*, inhibition of germ tube elongation values were 56, 45, 38, 5 and 42%, respectively. For *P. expansum*, the plant extract did not show positive results for inhibition of conidial germination or inhibition of germ tube elongation. In artificially inoculated fruits, the extract applied to nectarines was not protective against brown rot development from *M. laxa*, while for apricots effects were similar to those of the negative control for *P. digitatum* [35]. Dichloromethane and diethyl ether *T. officinale* extracts were tested on *P. expansum* by applying either a solution or its vapor to paper discs. The dichloromethane extract was more active of the two models, though direct inoculation in apples offered no observable inhibition [26]. Water extracts of *T. officinale* and *T. platycarpum* were tested against *C. lagenarium* in cucumber, exhibiting inhibition rates of the anthracnose lesions of 1.9 and 13% in treated leaves, and 11 and 5.3% in untreated leaves, respectively. These results were not significant compared to other plant extracts [42]. *In vivo* evaluation of protective effects in plant tissue has not been as successful as the *in vitro* assays, which is typical in cases of inhibitory activity validation. To avoid these ineffective results, concentrations are increased to demonstrate the pathogen control effect.

2.2.3. Animal pathogens

Regarding animal pathogens, *Saprolegnia* infections can account for significant salmonid losses. Treatment is difficult and there are reservations regarding efficacy, prompting a search for suitable alternatives. A *T. officinale* root extract was not as effective as a fungistatic at 10, 100, 1000, or 10,000 mg/mL [61]. The effects of *Taraxacum* polysaccharides were studied on the preservation of white shrimp (*Penaeus vannamei*) during refrigerated storage (10 days at 4°C) by soaking the shrimps in aqueous extracts (1–3% w/v). Samples were periodically evaluated for total viable count, pH value, and total volatile basic nitrogen, which resulted in 2–3% of shrimp in fresh conditions (<30 mg/100 mg of total volatile basic nitrogen) and a total viable count that only increased slightly during storage. This indicated that the treatment effectively

retarded bacterial growth during refrigerated storage, prolonging shrimp shelf life for up to 10 days [76].

In the case of the meat industry, an herb mixture including *T. officinale* as a substitute for fodder antibiotics in pig feeding revealed positive growth of the animal and no change in meat quality, confirming the possibility of using herbs as an antibiotic substitute in pig feed [77, 78]. Aqueous and ethanolic extracts of *T. mongolicum* could also inhibit four pathogenic bacteria responsible for cow mastitis, a serious disease in the cow industry, at concentrations of 0.13, 0.25, and 0.5 g/mL. In this case, the ethanolic extracts displayed slightly better antibacterial activities than aqueous extracts. For *E. coli*, *S. aureus*, *S. agalactiae*, and *S. dysgalactiae*, inhibition zone diameters were slightly larger for aqueous than for ethanolic extracts but showed between medium and high sensitivity [79]. Dandelion extract can not only be used to control pathogens but also to supplement the diet of animals, which could result in increased meat, milk, whey, and other yields, contributing to the food industry. Alternatively, the extracts could be utilized in the agricultural industry as biofertilizers to promote plant growth and strengthen the plant against biotic and abiotic stress.

2.3. *Taraxacum* antimicrobial action mechanisms

Innate plant immunity involves various defense responses, including cell wall reinforcements, lytic enzyme biosynthesis, secondary metabolite production, and pathogenesis-related proteins. To protect themselves from non-beneficial microorganisms, plants accumulate secondary metabolites that form chemical barriers to microbial attacks (phytoanticipins) and produce antimicrobials (phytoalexins) [80]. Phenolics and terpenoids are considered the primary mechanisms for plant defenses because these reduce microbial attacks by disrupting the cell membranes in microorganisms, bind to adhesins and cell wall compounds, and inactivate enzymes, among other roles [81]. The action mechanisms of natural compounds are related to the disintegration of the cytoplasmic membrane and destabilization of the proton motive force, electron flow, active transport, coagulation of the cell content, inhibition of protein synthesis, inhibition of DNA synthesis, and the synthesis of metabolites used for DNA synthesis [82]. Some action mechanisms are specific to certain targets and some targets may also be affected by more than one mechanism [83]. A general scheme of the action's sites and antimicrobial potential mechanism is presented in **Figure 1** of Supporting Information.

Even though *Taraxacum* is a plant with extremely high pathogen resistance, the underlying molecular mechanisms of antimicrobial activity are poorly studied [68]. Until now, most of the research on *Taraxacum* has focused on elucidating the compounds present in the extract, and, to a lesser extent, on the mechanism involved in the antimicrobial activity itself. One study specifically illustrated the effect of four proteins from *T. officinale* flowers on fungi by light microscopy and distinguished two modes of antimicrobial action, depending on the fungus tested. *Taraxacum* proteins completely blocked conidia germination or induced thickening of multiple local hyphae and irreversible cytoplasm plasmolysis [68, 69]. Different extracts from this genus showed positive inhibitory activity in controlled studies and were characterized by protein synthesis inhibition (e.g. chloramphenicol, tetracycline, gentamicin, and kanamycin) and cell wall synthesis (e.g. amphotericin, cefixime, cephalothin, and penicillin). These




Figure 1. Main action mechanisms for antimicrobial agents (adapted from Mulvey and Simor [84]).

mechanisms need to be addressed to elucidate the *Taraxacum* active compound action mechanisms because a direct relation with the positive controls cannot be pursued.

Another response that has been studied is the modulation of microbe adherence to body tissues. Adhesion to epithelial cells has been represented as the first step in the subsequent bacterial invasion of host cells [59]. These authors reported the partial inhibition of intestinal adherence of *C. jejuni* HT-29 cells using a commercial ethanolic *Taraxacum* extract. Cytotoxic activity was less than 10%, but no antibacterial activity was observed. Moreover, *Taraxacum* has been tested with the aim of controlling bacterial diseases by inhibiting communication between bacteria. An ethanolic extract of *T. officinale* aerial parts disturbed bacterial communication systems (or quorum sensing) for *C. violaceum*, showing the moderately positive effect of the extract on the attenuation of microbial pathogenicity [30]. In contrast, an ethanolic and water extract of the rhizomes of the same plant showed no significant activity in the same assay [65].

2.4. *Taraxacum* compounds related to antimicrobial action

Several studies have named a wide range of compounds, including terpenes, flavonoids, and phenolic compounds, as responsible for the medicinal activity of different plants [85, 86]. For *Taraxacum*, only a few studies concerning its antimicrobial properties have considered

chemical identification of the obtained extracts and this identification is chiefly qualitative (e.g. using colorimetric methods indicating presence or absence). Authors report the presence of terpenoids, triterpenoids, steroids, coumarins, phenols, saponins, flavonoids, flavones, flavonols, chalcones, phlobatannins, and cardiac glycosides in antimicrobial extracts [22, 27, 34, 36, 37, 43–45, 87, 88] but neither compound isolation nor further identification were performed.

Taraxasterol acetate, lupeol acetate, tranexamic acid, and squalene, among others, were identified in the dichloromethane extract of *T. officinale* leaves, which show low activity against *E. coli*, *P. aeruginosa*, *B. subtilis*, *C. albicans*, and *T. mentagrophytes* in an agar well assay at 30 µg but no observed activity against *S. aureus* or *A. niger* [27]. Terpenoids and flavonoids were identified in the ethanolic extracts of the *T. farinosum* root, which displayed antibacterial activity against *S. aureus*, *S. typhi*, *M. canis*, and *T. longifusus* in an agar well diffusion and agar tube dilution, while the herb extract was active only against the latter two strains [51]. Fractions of a methanolic root extract indicated the significant presence of phenolic-based compounds and hydroxyl-fatty acids with liquid and mass spectrometry, and were active against *S. aureus*, MRSA clinical, and *B. cereus* at 2 mg/mL, with MIC values ranging from 0.05 to 0.19 mg/mL, and crude extracts indicating values of 0.25–0.5 mg/mL [33]. An oligosaccharide extract (DOs) from this species exhibited high antibacterial activity against *E. coli*, *B. subtilis*, and *S. aureus* at 100 mg/mL, indicating that these oligosaccharides could potentially be used as antibacterial agents [48].

Concerning specific compounds, isolated *Taraxacum* peptides displayed antimicrobial activity at 6 µg/µL, corresponding to 52–79% of kanamycin activity against *P. syringae*, *B. subtilis*, and *X. campestris* at the same concentration [69], which is a promising value that warrants further experiments. These authors indicated that though *A. niger* appeared sensitive to four proteins (ToAMP1, ToAMP2, ToAMP3, and ToAMP4) from *T. officinale* flowers, *F. graminearum* was not susceptible to any of these proteins. All proteins displayed inhibition activity against *B. cinerea*, *B. sorokiniana*, *A. niger*, *P. debaryanum*, *F. oxysporum*, and *P. infestans*, with IC₅₀ values ranging between 1.2 and 5.8 µM. The ToAMPs were also active against *P. syringae*, *B. subtilis*, and *X. campestris*, similar to a kanamycin control. Additionally, ToAMP2 was active against *C. michiganensis* at up to 0.5 µg/µL. The disease development of *P. infestans* was inhibited by ToAMP2 at 1.3 µM (20–40%) to 5.2 µM (10–20%). In further studies, *B. sorokiniana*, *C. gloeosporioides*, and *V. albo-atrum* were insensitive to ToAMP4, another peptide isolated from the seed extract of *T. officinale*, at concentrations below 15 mM. The IC₅₀ values for the agent-sensitive fungi *A. alternata*, *A. niger*, *F. avenaceum*, and *P. betae* ranged from 2.9 to 13.1 mM, with MIC values from 1.0 to 8.0 mM; no activity was observed for *P. syringae*, *B. subtilis*, *E. coli*, or *C. michiganensis* [68, 69]. Peptides supposedly have broad-spectrum activity, lack of microbial resistance, and high efficacy [69], but some action mechanisms in these molecules are still poorly defined [89]. Peptides related to albumin 2S from *Taraxacum* seeds are active against phytopathogenic fungi and bacteria. Antifungal assays displayed different activities for the 2S isoforms (ToA1, ToA2, and ToA3). The spore germination of *B. cinerea*, *A. niger*, and *P. debaryanum* were the most tolerant, and *H. sativum*, *P. betae*, and *V. albo-atrum* were the most sensitive at concentrations ranging from 0.063 mg/mL to 0.25 mg/mL. *H. sativum* and *P. betae* were inhibited by ToA1, ToA2, and ToA3, but *F. oxysporum* and *V. albo-atrum* were only inhibited by ToA2 and ToA3, respectively. In potato tubers, *P. infestans* was inhibited by ToA3 at 0.06 mg/mL at 96 and 120 h, but at 144 h ToA2 inhibited better at 0.13 mg/mL [23].

Interestingly, an antimicrobial filtrate isolated from the fungal strains of *P. betae* (PG23) from *T. mongolicum* was proven active against *E. coli*, *S. aureus*, *A. hydrophila*, *E. tarda*, and *P. multocida*, and proposed as a potential antimicrobial product for poultry and aquatic disease control [88].

3. Driving forces and tendencies in *Taraxacum* antimicrobial research

Between 2000 and 2010, approximately 40 new drugs originating from terrestrial plants, terrestrial microorganisms, marine organisms, and terrestrial vertebrates and invertebrates against different bacteria, fungi, and viruses were launched on the market [90]. This follows distinct research tendencies. Studies related to antimicrobial and antifungal properties generally aim, in developing and developed countries, to respond to the necessity of finding new drugs or products based on traditional medicine at a low cost, confirming already established activity originating from oral tradition. The driving force behind studying new antimicrobial alternatives is the necessity of finding new drugs or natural products that act against diseases due to the increased drug resistance in the latter. Furthermore, the toxicity of synthetic compounds currently utilized in farming and agricultural industries has created a market for natural compounds that are safer, cheaper, and more effective against pathogens.

Modern phytochemistry, scientific equipment, and technology have had a significant impact on natural product chemistry, including isolation, extraction, purification, and structure determination. However, this discipline still demands that research investigators establish the clinical significance of natural compounds and recognize them as drugs or industrial products (pesticides, bactericides, pharmaceutical products, etc.) [91]. Bioactive compounds in botanical drugs are purportedly superior to monosubstances because of synergistic effects. Similarly, multidrug therapy is highly important against resistant microbial strains due to the enhanced efficacy, reduced toxicity, decreased adverse side effects, increased bioavailability, lowered dosage, and reduced evolution of antimicrobial resistance [92].

Even when antibiotics have been effective in treating infectious diseases, resistance to the action mechanisms has led to the emergence of new and the re-emergence of old infectious diseases. Several plant extracts exhibit synergistic activity against microorganisms, with natural products (including flavonoids and essential oils) and synthetic drugs effectively combating bacterial, fungal, and mycobacterial infections. The mode of action of combinations differs significantly from the individual use of the same drugs; hence, isolating a single component may not highlight its importance, simplifying the task of the pharmacological industries [93].

Acknowledgements

This work has been supported by Innova Chile CORFO Code FCR-CSB 09CEII-6991 and a doctoral fellowship awarded by the Pontifical Catholic University of Valparaíso, Chile Project DI Iniciación COD 039.454/2017 Pontificia Universidad Católica de Valparaíso.

Author details

María Eugenia Martínez Valenzuela¹, Katy Díaz Peralta², Lorena Jorquera Martínez³ and Rolando Chamy Maggi^{1*}

*Address all correspondence to: rolando.chamy@pucv.cl

1 Escuela de Ingeniería Bioquímica, Facultad de Ingeniería, Pontificia Universidad Católica de Valparaíso, Valparaíso, Chile

2 Departamento de Química, Universidad Técnico Federico Santa María, Valparaíso, Chile

3 Escuela de Ingeniería en Construcción, Facultad de Ingeniería, Pontificia Universidad Católica de Valparaíso, Valparaíso, Chile

References

- [1] Modi C, Modi S, Patel H, Dudhatra G, Kumar A, Awale M. Herbal Antibacterials: A review. *Journal Intercultural Ethnopharmacology*. 2012;1:52-61. DOI: 10.5455/jice.20120207113423
- [2] Mahesh B, Satish S. Antimicrobial activity of some important medicinal plant against plant and human pathogens. *World Journal Agrícola Science*. 2008;4:839-843. DOI: 10.1556/EuJMI.3.2013.4.6
- [3] Vashegyi I, Marozsán-Tóth Z, Galiba G, Dobrev P, Vankova R, Tóth B. Cold response of dedifferentiated barley cells at the gene expression, hormone composition, and freezing tolerance levels: Studies on callus cultures. *Molecular Biotechnology*. 2013;54:337-349. DOI: 10.1007/s12033-012-9569-9
- [4] Stewart-Wade S, Neumann S, Collins L, Boland G. The biology of Canadian weeds. 117. *Taraxacum officinale* G.H. Weber ex Wiggers. *Canadian Journal Plant Science*. 2002;82:825-853. DOI: org/10.4141/P01-010
- [5] Lardos A. The botanical materia medica of the Iatrosophikon – A collection of prescriptions from a monastery in *Cyprus*. *Journal of Ethnopharmacology*. 2006;104:387-406. DOI: 10.1016/j.jep.2005.12.035
- [6] González-Castejón M, Visioli F, Rodríguez-Casado A. Diverse biological activities of dandelion. *Nutrition Reviews*. 2012;70:534-547. DOI: 10.1111/j.1753-4887.2012.00509.x
- [7] Schütz K, Carle R, Schieber A. *Taraxacum* – A review on its phytochemical and pharmacological profile. *Journal of Ethnopharmacology*. 2006;107:313-323. DOI: 10.1016/j.jep.2006.07.021
- [8] Li G. Pharmacology, toxicity and clinic of traditional Chinese medicine. In: Tianjin Science and Technique. Tianjin, China: Translation Publishing House; 1992. pp. 207-208

- [9] Ou M, Li YW. The Traditional Chinese Drug and Its Usage. Hong Kong, China: Hai Feng Press; 1994. pp. 49-51
- [10] Zhu M, Wong P, Li R. Effects of *Taraxacum mongolicum* on the bioavailability and disposition of ciprofloxacin in rats. *Journal of Pharmaceutical Sciences*. 1999;**88**:632-634. DOI: 10.1021/js980367q
- [11] Caraballo A, Caraballo B, Rodríguez-Acosta A. Preliminary assessment of medicinal plants used as antimalarials, in the southeastern Venezuelan Amazon. *Revista da Sociedade Brasileira de Medicina Tropical*. 2004;**37**:186-188. DOI: 10.1590/S0037-86822004000200016
- [12] Guarino C, De Simone L, Santoro S. Ethnobotanical study of the Sannio area, Campania, Southern Italy. *Ethnobotany Research and Applications*. 2008;**6**:255-317. DOI: 10.17348/era.6.0.255-317
- [13] Saghir I, Awan A, Majid S, Khan M, Qureshi S, Bano S. Ethnobotanical studies of Chikar and its allied areas of district Muzaffarabad. *Journal of Biological Sciences*. 2001;**1**:1165-1170
- [14] Guarrera P, Savo V. Perceived health properties of wild and cultivated food plants in local and popular traditions of Italy: A review. *Journal of Ethnopharmacology*. 2013;**146**:659-680. DOI: 10.1016/j.jep.2013.01.036
- [15] Neves J, Matos C, Moutinho C, Queiroz G, Gomes L. Ethnopharmacological notes about ancient uses of medicinal plants in Trás-os-Montes (northern of Portugal). *Journal of Ethnopharmacology*. 2009;**124**:270-283. DOI: 10.1016/j.jep.2009.04.041
- [16] Rodriguez-Fragoso L, Reyes-Esparza J, Burchiel S, Herrera-Ruiz D, Torres E. Risks and benefits of commonly used herbal medicines in Mexico. *Toxicology and Applied Pharmacology*. 2008;**227**:125-135. DOI: 10.1016/j.taap.2007.10.005
- [17] Song M, Kim H. Ethnomedicinal application of plants in the western plain region of North Jeolla Province in Korea. *Journal of Ethnopharmacology*. 2011;**137**:167-175. DOI: 10.1016/j.jep.2011.05.001
- [18] Martinez M, Poirrier P, Chamy R, Prüfer D, Schulze-Gronover C, Jorquera L, Ruiz G. *Taraxacum officinale* and related species – An ethnopharmacological review and its potential as a commercial medicinal plant. *Journal of Ethnopharmacology*. 2015;**169**:244-262. DOI: 10.1016/j.jep.2015.03.067
- [19] Wat C, Johns T, Towers G. Phototoxic and antibiotic activities of plants of the Asteraceae used in folk medicine. *Journal of Ethnopharmacology*. 1980;**2**:279-290. DOI: 10.1016/S0378-8741(80)81006-3
- [20] Izzo A, Di Carlo G, Biscardi D, De Fusco R, Mascolo N, Borrelli F, Capasso F, Fasulo M, Autore G. Biological screening of Italian medicinal plants for antibacterial activity. *Phytotherapy Research*. 1995;**9**:281-286. DOI: 10.1002/ptr.2650090410
- [21] Silva N, Fernandes A. Biological properties of medicinal plants: A review of their antimicrobial activity. *Journal of Venomous Animals and Toxins including Tropical Diseases*. 2010;**16**:402-413. DOI: 10.1590/S1678-91992010000300006

- [22] Ghaima K, Hashim N, Ali S. Antibacterial and antioxidant activities of ethyl acetate extract of nettle (*Urtica dioica*) and dandelion (*Taraxacum officinale*). Journal of Applied Pharmaceutical Science. 2013;**3**:96-99. DOI: 10.7324/JAPS.2013.3518
- [23] Odintsova T, Rogozhin E, Sklyar I, Musolyamov A, Kudryavtsev A, Pukhalsky V, Smirnov A, Grishin E, Egorov T. Antifungal activity of storage 2S albumins from seeds of the invasive weed dandelion *Taraxacum officinale* Wigg. Protein and Peptide Letters. 2010;**17**:522-529. DOI: 10.2174/092986610790963591
- [24] Stanojković A, Ceković J, Čomić L, Pivić R, Stanojković A. Antibacterial properties of some plants from the family *Asteraceae* growing wild in Serbia. Lekovite sirovine. 2008;**26/27**:11-20
- [25] Jaca T, Kambizi L. Antibacterial properties of some wild leafy vegetables of the Eastern Cape Province, South Africa. Journal of Medicinal Plants. 2011;**5**:2624-2628. DOI: <http://www.academicjournals.org/journal/JMPR/article-abstract/3CA6E7916006>
- [26] Ikeura H, Somsak N, Kobayashi F, Kanlayanarat S, Hayata Y. Application of selected plant extracts to inhibit growth of *Penicillium expansum* on apple fruits. Plant Pathology Journal. 2011;**10**:79-84. DOI: 10.3923/ppj.2011.79.84
- [27] Ragasa C, Apuada M, Rideout J. Terpenoids from *Taraxacum officinale*. NRCP Research Journal. 2009;**10**:17-26
- [28] Im D, Lee K. Antioxidative and antibacterial activity and Tyrosinase inhibitory activity of the extract and fractions from *Taraxacum coreanum* Nakai. Korean Journal of Medicinal Crop Science. 2011;**19**:238-245. DOI: 10.7783/KJMCS.2011.19.4.238
- [29] Qiao H, Sun T. Antibacterial activity of ethanol extract and fractions obtained from *Taraxacum mongolicum* flower. Research Journal of Pharmacognosy. 2014;**1**:35-39. Available from: http://rjpharmacognosy.ir/article_6336_609.html
- [30] Zaki A, Shaaban M, Hashish N, Amer M, Lahloub M. Assessment of anti-quorum sensing activity for some ornamental and medicinal plants native to Egypt. Scientia Pharmaceutics. 2013;**81**:251-258. DOI: 10.3797/scipharm.1204-26
- [31] Han S, Hwang J, Park S, Lee K, Ko K, Kim K, Kim K. Potential effect of solvent fractions of *Taraxacum mongolicum* H. on protection of gastric mucosa. Korean Journal of Medicinal Crop Science. 2005;**37**:84-89
- [32] Sun H, Li Z, Liu W, Wang X, Zhang T, Feng Z. *In vitro* antibacterial activities of antibiotics and traditional chinese medicinal herb extracts on *Escherichia coli* and *Staphylococcus aureus*. Asian Journal of Chemistry. 2014;**26**:298-302. DOI: 10.14233/ajchem.2014.16174
- [33] Kenny O, Brunton N, Walsh D, Hewage C, Mcloughlin P, Smyth T. Characterisation of antimicrobial extracts from dandelion root (*Taraxacum officinale*) using LC-SPE-NMR. Phytotherapy Research. 2015;**29**:526-532. DOI: 10.1002/ptr.5276
- [34] Tetley C, Ocloo A, Nagajyothi P, Lee K. An *in vitro* analysis of antiproliferative and antimicrobial activities of solvent fractions of *Taraxacum officinale* (dandelion) leaf. Journal of Applied Pharmaceutical Science. 2014;**4**:41-45. DOI: 10.7324/JAPS.2014.40309

- [35] Gatto M, Ippolito A, Linsalata V, Cascarano N, Nigro F, Vanadia S, Di Venere D. Activity of extracts from wild edible herbs against postharvest fungal diseases of fruit and vegetables. *Postharvest Biology and Technology*. 2011;**61**:72-82. DOI: 10.1016/j.postharvbio.2011.02.005
- [36] Fabri R, Nogueira M, Dutra L, Bouzada M, Scio E. Potencial antioxidante e antimicrobiano de espécies da família Asteraceae. *Revista Brasileira de Plantas Medicinai*s. 2011;**13**:183-189. DOI: 10.1590/S1516-05722011000200009
- [37] Khan A, Qureshi R, Gillani S, Ullah F. Antimicrobial activity of selected medicinal plants of Margalla Hills, Islamabad, Pakistan. *Journal of Medicinal Plants Research*. 2011;**5**:4665-4670
- [38] Kim K, Chun H, Han Y. Screening of antimicrobial activity of the dandelion (*Taraxacum platycarpum*) extract. *Korean Society of Food Science and Nutrition*. 1998;**14**:114-118
- [39] Kim K, Min K, Lee S, Han Y. Isolation and identification of antimicrobial compound from dandelion (*Taraxacum platycarpum* D.). *Journal of the Korean Society of Food Science and Nutrition*. 1999;**28**:822-829
- [40] Park J, Jang J, Lee M, Park E. Effect of dandelion (*Taraxacum officinale*) extracts on the intestinal microorganisms of streptozotocin-induced diabetic rats. *Journal of the Korean Society of Food Science and Nutrition*. 2002;**31**:1112-1118. DOI: 10.3746/jkfn.2002.31.6.1112
- [41] Sohail Z, Afzal M, Afzal A, Rahman I, Shad S, Ahmed B, Anjum N, Qureshi K, Bibi A. *In vitro* antibacterial study of *Taraxacum officinale* leaves extracts against different bacterial pathogenic strains. *Journal of Pharmacognosy and Phytochemistry*. 2014;**3**:15-17
- [42] Inagaki H, Yamaguchi A, Kato K, Kageyama C, Iyozumi H, Oki Y. Screening of weed extracts for antifungal properties against *Colletotrichum lagenarium*, the causal agent of anthracnose in cucumber. *Weed Biology and Management*. 2008;**8**:276-283. DOI: 10.1111/j.1445-6664.2008.00308.x
- [43] Jassim A, Farhan S, Noori O. Identification of dandelion *Taraxacum officinale* leaves components and study its extracts effect on different microorganisms. *Journal of Al-Nahrain University – Science*. 2012;**15**:7-14
- [44] Kenny O, Smyth T, Walsh D, Kelleher C, Hewage C, Brunton N. Investigating the potential of under-utilised plants from the Asteraceae Family as a source of natural antimicrobial and antioxidant extracts. *Food Chemistry*. 2014;**161**:79-86. DOI: 10.1016/j.foodchem.2014.03.126
- [45] Lateef O, Issah Y. Screening ethanolic and aqueous leaf extracts of *Taraxacum officinale* for in vitro bacteria growth inhibition. *Journal of Pharmaceutical and Biomedical Sciences*. 2012;**20**:06
- [46] Parveen S, Wani A, Ganie A, Pala S, Mir R. Antifungal activity of some plant extracts on some pathogenic fungi. *Archives of Phytopathology and Plant Protection*. 2013;**47**:279-284. DOI: 10.1080/03235408.2013.808857
- [47] Peng S, Dai W, Yu H, Wang Y, Wang X, Sun S. Antibacterial activity of aqueous and ethanolic extracts of *Portulaca oleracea* L and *Taraxacum mongolicum* against pathogenic

- bacteria of cow mastitis. International Journal of Applied Research in Veterinary Medicine. 2014;**12**:210-213. DOI: 10.18805/ijar.5960
- [48] Qian Z, Jiang L. Preparation and antibacterial activity of the oligosaccharides derived from *Rhizoma phragmites*. Carbohydrate Polymers. 2014;**111**:356-358. DOI: 10.1016/j.carbpol.2014.04.043
- [49] Chan G, Kim H. Antimicrobial effects of extracts of *Taraxacum officinale* H. on acnes strains. 국제통합보완대체의학회지 [International Journal of Integrated CAM]. 2011;**7**:3-16. DOI: 10.4172/1948-5948.1000287
- [50] Redzic S, Kurtagic H, Prazina N, Tuka M, Avdagic T. The antimicrobial activity of honey in relation to the composition of pollen (Bosnia-Herzegovina, W. Balkan). Planta Medicinal. 2011;**77**:SL62. DOI: 10.1055/s-0031-1282185
- [51] Sarer E, Sahin A, Gökbulut A. The investigation on the terpenoids and flavonoids of an endemic *Taraxacum* species from Turkey. Planta Medicinal. 2010;**76**:P303. DOI: 10.1055/s-0030-1264601
- [52] Sharma V, Sharma H, Mehta D, Chhabra B, Thakur D, Sourirajan A, Dev K. Comparative analysis of antibacterial and antifungal properties of traditional medicinal plants of Shimla and Solan, Himachal Pradesh. International Journal of Pharmacy and Pharmaceutical Research. 2014;**6**:18-26. DOI: 10.1016/j.jtcme.2017.04.002
- [53] Shafiq N, Al-Hashimi M. The antibacterial evaluation of dandelion extracts as root canal irrigating solutions (A comparative study). Journal of Bagdhdad College of Dentistry. 2014;**26**:35-40
- [54] Cwikla C, Schmidt K, Matthias A, Bone K, Lehmann R, Tiralongo E. Investigations into the antibacterial activities of phytotherapeutics against *Helicobacter pylori* and *Campylobacter jejuni*. Phytotherapy Research. 2010;**24**:649-656. DOI: 10.1002/ptr.2933
- [55] Kim J, Yi Y, Lim Y. Biological and antifungal activity of herbal plant extracts against *Candida* species. Korean Journal of Microbiology and Biotechnology. 2009;**37**:42-48
- [56] Bahraminejad S, Abbasid S, Fazlali M. In vitro antifungal activity of 63 Iranian plant species against three different plant pathogenic fungi. African Journal of Biotechnology. 2011;**10**:16193-16201. DOI: 10.5897/AJB11.1203
- [57] Borchardt J, Wyse D, Sheaffer C, Kauppi K, Fulcher R, Ehlke N, Biesboer D, Bey R. Antimicrobial activity of native and naturalized plants of Minnesota and Wisconsin. Journal of Medicinal Plants Research. 2008;**2**:98-110
- [58] Ionescu D, Mihele D, Predan G, Rizea G, Dune A, Ivopol G. Antimicrobial activity of some hydroalcoholic extracts of artichoke (*Cynara scolymus*), burdock (*Arctium lappa*) and dandelion (*Taraxacum officinale*). Agricultural Food Engineering. 2013;**6**:113-120 Available from: <http://webbut.unitbv.ro/Bulletin/Seri>
- [59] Bensch K, Tiralongo J, Schmidt K, Matthias A, Bone KM, Lehmann R, Tiralongo E. Investigations into the antiadhesive activity of herbal extracts against *Campylobacter jejuni*. Phytotherapy Research. 2011;**25**:1125-1132. DOI: 10.1002/ptr.3384

- [60] Tosun F, Vural M, Palittapongarnpim P. Antimycobacterial screening of some Turkish plants. *Journal of Ethnopharmacology*. 2004;**95**:273-275. DOI: 10.1016/j.jep.2004.07.011
- [61] Caruana S, Yoon G, Freeman M, Mackie J, Shinn A. The efficacy of selected plant extracts and bioflavonoids in controlling infections of *Saprolegnia australis* (Saprolegniales; Oomycetes). *Aquaculture*. 2012;**358–359**:146-154. DOI: 10.1016/j.aquaculture.2012.06.035
- [62] López-García J, Kuceková Z, Humpolíček P, Mlcek J, Sáha P, Humpolíček P, Mlček J, Sáha P. Polyphenolic extracts of edible flowers incorporated onto atelocollagen matrices and their effect on cell viability. *Molecules*. 2013;**18**:13435-13445. DOI: 10.3390/molecules181113435
- [63] Newton S, Lau C, Gurcha S, Besra G, Wright C. The evaluation of forty-three plant species for *in vitro* antimycobacterial activities; isolation of active constituents from *Psoralea corylifolia* and *Sanguinaria canadensis*. *Journal of Ethnopharmacology*. 2002;**79**: 57-67. DOI: 10.1016/S0378-8741(01)00350-6
- [64] Liu Y, Nielsen M, Staerk D, Jäger A. High-resolution bacterial growth inhibition profiling combined with HPLC-HRMS-SPE-NMR for identification of antibacterial constituents in Chinese plants used to treat snakebites. *Journal of Ethnopharmacology*. 2014;**155**: 1276-1283. DOI: 10.1016/j.jep.2014.07.019
- [65] Tolmacheva A, Rogozhin E, Deryabin D. Antibacterial and quorum sensing regulatory activities of some traditional Eastern-European medicinal plants. *Acta Pharmaceutica*. 2014;**64**:173-186. DOI: 10.2478/acph-2014-0019
- [66] Woods-Panzaru S, Nelson D, McCollum G, Ballard L, Millar B, Maeda Y, Goldsmith C, Rooney P, Loughrey A, Rao J, Moore J. An examination of antibacterial and antifungal properties of constituents described in traditional Ulster cures and remedies. *Ulster Medical Journal*. 2009;**78**:13-15. PMCID: PMC2629014
- [67] The Plant List. Version 1.1. Published on the Internet. Available from: <http://www.theplantlist.org/> [Accessed: May 2015]
- [68] Astafieva A, Rogozhin E, Andreev Y, Odintsova T, Kozlov S, Grishin E, Egorov T. A novel cysteine-rich antifungal peptide ToAMP4 from *Taraxacum officinale* Wigg. flowers. *Plant Physiology and Biochemistry*. 2013;**70**:93-99. DOI: 10.1016/j.plaphy.2013.05.022
- [69] Astafieva A, Rogozhin E, Odintsova T, Khadeeva N, Grishin E, Egorov T. Discovery of novel antimicrobial peptides with unusual cysteine motifs in dandelion *Taraxacum officinale* Wigg. flowers. *Peptides*. 2012;**36**:266-271. DOI: 10.1016/j.peptides.2012.05.009
- [70] Gao D. Analysis of nutritional components of *Taraxacum mongolicum* and its antibacterial activity. *Pharmacognosy Journal*. 2010;**2**:502-505. DOI: 10.1016/S0975-3575(10)80039-2
- [71] Hanawa F, Kanauchi M, Tahara S, Mizutani J. Lettucenin A as a phytoalexin of dandelion and its elicitation in dandelion cell cultures. *Journal of the Faculty of Agriculture, Hokkaido University*. 1995;**66**:151-162

- [72] Voss-Rech D, Klein C, Techio V, Scheuermann G, Rech G, Fiorentin L. Antibacterial activity of vegetal extracts against serovars of *Salmonella*. *Ciência Rural*, Santa Maria. 2011;**41**:314-320. DOI: 10.1590/S0103-84782011000200022
- [73] Al-kalby J, Alwan N. Effects of ethanolic extract of *Taraxacum officinale* leaves on some physiological and reproductive parameters of local adult male rabbits. *Basrah Journal of Veterinary Research*. 2007;**6**:42-55
- [74] Elkhalfi B, Essari A, Serrano A, Soukri A. Antibacterial activity of plant methanolic extracts on a field isolate of *Pseudomonas syringae* pv tomato from the Casablanca region (Morocco). *Advances in Bioscience of Biotechnology*. 2013;**4**:1-9. DOI: 10.4236/abb.2013.47A2001
- [75] Chen Z. Clinical study of 96 cases with chronic hepatitis B treated with jiedu yanggan gao by a double-blind method. *Zhong Xi Yi Jie He Za Zhi*. 1990;**10**:71-74
- [76] Wang H. Effect of dandelion polysaccharides on the retardation of the quality changes of white shrimp. *International Journal of Biological Macromolecules*. 2014;**68**:205-208. DOI: 10.1016/j.ijbiomac.2014.05.002
- [77] Urbanczyk J, Hanczakowska E, Swiatkiewicz M. Herb mixture as an antibiotic substitute in pig feeding. *Medycyna Weterynaryjna*. 2002;**58**:887-889
- [78] Yan L, Meng QW, Kim IM. The effects of dietary *Houttuynia cordata* and *Taraxacum officinale* extract powder on growth performance, nutrient digestibility, blood characteristics and meat quality in finishing pigs. *Livestock Science*. 2011;**141**:188-193. DOI: 10.1016/j.livsci.2011.05.017
- [79] Yang L, Lü J, Feng Y. Partition and extraction of total flavonoids from *Taraxacum mongolicum* in PEG/(NH₄)₂SO₄ aqueous two-phase system. *Chemical Industry and Engineering Progress*. 2014;**33**:1992-1996
- [80] González-Lamothe R, Mitchell G, Gattuso M, Diarra M, Malouin F, Bouarab K. Plant antimicrobial agents and their effects on plant and human pathogens. *International Journal Molecular of Science*. 2009;**10**:3400-3419. DOI: 10.3390/ijms10083400
- [81] Cowan M. Plant products as antimicrobial agents. *Clinical Microbiology Reviews*. 1999;**12**:564-582. PMCID: PMC88925
- [82] Singh S, Barrett J. Empirical antibacterial drug discovery – Foundation in natural products. *Biochemical Pharmacology*. 2006;**71**:1006-1015. DOI: 10.1016/j.bcp.2005.12.016
- [83] Burt S. Essential oils: Their antibacterial properties and potential applications in foods – A review. *International Journal of Food Microbiology*. 2004;**94**:223-253. DOI: 10.1016/j.ijfoodmicro.2004.03.022
- [84] Mulvey M, Simor A. Antimicrobial resistance in hospitals: How concerned should we be? *Canadian Medical Association Journal*. 2009;**180**:408-415. DOI: 10.1503/cmaj.080239

- [85] Chadwick M, Trewin H, Gawthrop F, Wagstaff C. Sesquiterpenoids lactones: Benefits to plants and people. *International Journal of Molecular Science*. 2013;**14**:12780-12805. DOI: 10.3390/ijms140612780
- [86] Mir M, Sawhney S, Jassal M. Qualitative and quantitative analysis of phytochemicals of *Taraxacum officinale* Wucpecker. *Journal of Pharmacy and Pharmacology*. 2013;**2**:1-5
- [87] Khan A, Qureshi R, Ullah F, Gilani S, Nosheen A, Sahreen S, Laghari M, Laghari M, Hussain I, Murad W. Phytochemical analysis of selected medicinal plants of Margalla Hills and surroundings. *Journal of Medicinal Plants Research*. 2011;**5**:6017-6023. DOI: 10.5897/JMPR11.869
- [88] Zhang H, Xiong Y, Zhao H, Yi Y, Zhang C, Yu C, Xu C. An antimicrobial compound from the endophytic fungus *Phoma* sp. isolated from the medicinal plant *Taraxacum mongolicum*. *Journal of the Taiwan Institute of Chemical Engineers*. 2013;**44**:177-181. DOI: 10.1016/j.jtice.2012.11.013
- [89] Barbosa P, Del Sarto R, Silva O, Franco O, Grossi-De-Sa M. Antibacterial peptides from plants: What they are and how they probably work. *Biochemistry Research International*. 2011. 9 pp. Article ID: 250349. DOI: 10.1155/2011/250349
- [90] Brahmachari G. Natural products in drug discovery: Impacts and opportunities – An assessment. *Bioactive Natural Products: Opportunities and Challenges in Medicinal Chemistry*. 2012;**1**:1-199
- [91] Mahmood Z, Mahmood S. Antibiotic natural products: Opportunities and challenges. In: *Microbial Pathogens and Strategies for Combating Them: Science, Technology and Education*. Formatex Research Center. 2013;**2**:823-833
- [92] Radulović N, Blagojević P, Stojanović-Radić Z, Stojanović N. Antimicrobial plant metabolites: Structural diversity and mechanism of action. *Current Medicinal Chemistry*. 2013;**20**:932-952. DOI: 10.2174/092986713805219136
- [93] Hemaiswarya S, Kruthiventi A, Doble M. Synergism between natural products and antibiotics against infectious diseases. *Phytomedicine*. 2008;**15**:639-652. DOI: 10.1016/j.phymed.2008.06.008