We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Bioaccumulation and Toxic Profiling of Nanostructured Particles and Materials

Subas Chandra Dinda

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.74802>

Abstract

Use of nanotechnological based formulations and nanomaterials are increasing day-by-day in wide range covering a broad typology of applications, from design and development of targeted drug delivery systems, manufacturing of pesticides, domestic appliances, textiles, to bioremediation engineering. There are therefore concerns about the environmental risks or bioaccumulation-related issues that may arise particularly resulting from the application of drug-loaded nanocarriers or effect of pesticides that reach the natural ecosystems. This is a major threat in the present era and needs to be balanced against their undoubted benefits to human society. The assessment of the physical and chemical properties of nanoparticles and nanomaterials influencing their toxic manifestation due to accumulation in human or in animal organs is still poorly investigated. This chapter reviews the possibilities of bioaccumulation of different nanoscale particles and materials, their potential acute and subacute toxicological profile and their identification and characterization in different organs and tissues of vertebrates.

Keywords: nanomaterials, nanoparticles, bioaccumulation, characterization, toxicity, analytical methods, in vitro studies, in vivo studies

1. Introduction

Nanostructured materials can be easily accessible to the cellular level of a tissue or an organ in comparison to macro scale particles given their small size that makes them prone to passive entrance. On the other hand the functional performance of nanoparticles depends on their size, morphology and chemical nature of its surface which influence the triggering of phagocytosis through the cellular membranes. Moreover, large pay loads may conduce to stress conditions, and thereby increase inflammation and reduce defense action against pathogens

[1, 2]. It is also known that nanoparticles made up of non-degradable or slowly degradable materials may accumulate at the cellular level hindrance the enzymatic activity of functional proteins [3]. The toxic manifestations may also arise due to their chemical composition and structure, surface charge, solubility and aggregation on the site of elimination or filtration such as at the glomerular filtration of nephrons. In this context, it is fundamental to investigate the aggregation and agglomeration of the nanostructured particles and materials, followed in the ionic environment of the biological fluids to elicit its functional activity and resulting toxicity.

Among available manufactured nanomaterials, titanium dioxide (TiO_2) is one of the most widely used to date [4]. Previous studies suggested that TiO_2 nanoparticles production ranged from 7800 to 38,000 metric tons per year in the USA alone and will be approximately 2.5 million metric tons by 2025 [5, 6]. The fate and bioaccumulation of TiO_2 based nanomaterials (nanotubes TiO_2 -NTs) and nanoparticles (TiO_2 -NPs) in a paddy microcosm over a period of 17 days showed that the Ti levels were the highest in biofilms during the exposure period [7].

Bioaccumulation factors indicated that TiO_2 -NPs and TiO_2 -NTs were largely transferred from low to superior trophic levels [8]. Considering the potential entries of TiO_2 based nanomaterials in organisms, their bioaccumulation throughout the food chain should be regarded with great concern in terms of the overall health of the natural ecosystems [9].

Aquatic floras play a vital role as primary producers of biomass which constitutes the elementary level of food in trophic chains. Environmental exposure of algae to nanoparticles at potential toxic concentrations may affect natural ecosystems by interfering at the basic energy source of the food webs. As an example, Chandler et al. demonstrated that increased alga mortality occurred upon exposure to nano-ZnO and C60 at the size of 1 ppm in comparison to 10 ppm [10].

At the level of primary consumers it was revealed that bioaccumulation of nanomaterials in *Daphnia magna* through alga (used as food source), presented a more toxic profile for zinc oxide in comparison to carbon based nanomaterials which may be explained by its higher solubility [10]. In addition Teer and colleagues investigated the effects of nano-Ag derived from coated AgNPs and AgNO_3 on phosphate availability given its potential as Ag + ligand and as determinant of phytoplankton productivity [11]. It was observed that both nanoparticles accumulated at similar concentrations into the primary producer during high phosphate insult. Yet, AgNO_3 accumulation was only increased for low phosphate condition [11].

Regarding studies at superior trophic level, it was shown that goldfish *Carassius auratus* exposed to 10–100 mg/L of TiO_2 -NPs significantly accumulated these nanoparticles on tissues and organs, namely: intestine from 42.71 to 110.68 ppb and gills from 4.10 to 9.86 ppb. In addition, overall growth inhibition induced by lipid oxidation in the liver was recorded [12]. In other work, Hanna and colleagues investigated the potential ecological damage of an essential trace element - Cu - derived from CuO engineered nanoparticles (ENPs) exposed marine mussels exposed to 3 mg of CuO ENPs for 4 weeks resulting its clearance of ionic Cu which decreased to 48% leading to a reduction in the invertebrates growth rate in comparison

to the control animals, suggesting that CuO ENPs are less toxic than ionic Cu probably due to the slow dissolution rate of the former [13].

Investigation on top predators including humans demonstrated that inhaled nanoscale particles are less cleared by macrophages than large sized particles [14], and are able to translocate to other organs through circulatory or lymphatic drainage which may increase cytokine production and imbalance in redox potential [15] toward oxidation, leading to inflammation or cell death [16]. Nano sized particles and materials were also demonstrated to be taken up by mitochondria and nucleus of the cell [17] causing DNA mutation [18].

To surmount the health risk and potential toxic manifestations or disease associated conditions resulting from nano-devices or nano-structured materials exposure, efficient multifunctional designs are required to make the most of their interesting features while avoid adverse effects.

2. Mechanism of nanoparticle bioaccumulation and toxic action

Metallic elements such as Ni, Cu, Ti and Ag are increasingly considered as nanoparticles components that are being applied to biocides and antimicrobials [19]. Once release to the natural environment their safety profile is of concern. Studies on their toxicological potential were greatly investigated in aquatic species from invertebrates to vertebrates given that these ecosystems compose the ultimate sink of their fate: zooplankton [20, 21], fish [22, 23], algae [24], marine [25] and fresh water crabs [26]. Overall, these works point to three principal factors as potentially responsible for the metallic nanoparticles toxicity, namely: (i) dissolution rate, (ii) cellular uptake and (iii) induced level of oxidative stress and subsequent cellular damage [27].

Nonmetallic carbon-based nanomaterials such as fullerenes and carbon nanotubes, which are under exploration in cancer drug delivery, were reported to be cytotoxic [28]. Obtained data on cell proliferation of human lung cancer cells and human keratinocytes upon exposure to carbon-based nanomaterials revealed decreased cellular viability [29]. Intravenous administration of single-walled carbon nanotubes on mice showed their long term accumulation in the mammalian organs such as liver, lungs and spleen, which was observed using Raman spectroscopy and TEM technique [30]. In addition, decreased levels glutathione (GSH), and increased malondialdehyde (MDA) levels suggested that the toxicity of the carbon nanotubes is due to oxidative stress [31].

Increased levels of reactive oxygen species (ROS) and malondialdehyde (MDA) were further reported after administration of silica nanoparticles (15–46 nm) at a dose of 10–100 µg/mL in human bronchoalveolar carcinoma cells [32].

2.1. Toxicity of nanoparticles due to dissolution

The toxicity of metallic nanoparticles was found to be dependent on the rate of dissolution of the metal ions from its respective nanoparticulate formulations. Bondarenko and colleagues demonstrated that Zn-ions released from ZnO NPs did not exerted significantly different toxic effects in vivo on different aquatic models, neither in vitro on mammalian cells [33].

Differences were only found on its dissolution rate, which were revealed to be non-species dependent [33]. Under aerobic conditions it was also found a positive correlation among the dissolution rate of AgNPs and their toxicological profile upon exposure of bacteria and zooplankton models [34, 35]. In addition, sub-toxic effects of CuO NPs observed in bacteria seem to be associated with dissolved Cu ions liberated from the respective nanoformulation, triggering ROS production and causing cell death [36].

Regarding non-metallic nanoscale particles, biodegradable polymeric nanoparticles made up of poly-(D, L-lactide-co-glycolide) i.e. PLGA and its derivatives it is known that their drug release is based on the dissolution rate of its surface coating polymeric material PLGA [37]. Nevertheless, it was demonstrated in macrophages that the surface coating of these polymeric nanoparticles may be linked to induced toxicity by increasing oxidative stress levels [38].

2.2. Uptake of nanoparticles and their effects on biological membranes

Metallic nano-sized particles as ZnO NPs, with size inferior to 10 nm, were demonstrated to be highly internalized by prokaryotes in vitro causing cellular damage to both Gram negative *Escherichia coli* and Gram positive *Staphylococcus aureus* [39, 40]. This bactericidal potential was also observed for Ag NPs in the size range of 1–10 nm which was shown to be size-dependent [41, 42]. TEM microscopy confirmed bacteria internalization and showed a uniform pattern of distribution [40]. Even at non cytotoxic concentrations cellular uptake occurred [43].

Regarding the mechanisms of internalization, it was proposed that non-specific diffusion and membrane damage, and porins-specific intake are the potential modes through which the NPs could transverse bacterial wall [44]. Cellular uptake depends on multiple intrinsic (e.g. specific and dependent on the types of cell, tissue or organs) and extrinsic (e.g. NPs size and coating) variables [45]. ZnO NPs functionalization was demonstrated to affect bacteria membrane permeability increasing cellular uptake levels [46]. The hydroxyl groups of polyvinylalcohol (PVA) macromolecule existent on the surface of the coated ZnO NPs was found to disrupt the cellular membrane given its alkaline nature [46]. In case of Ag NPs, it has been shown that certain sizes of these metallic nanoparticles attached to Gram-negative *Escherichia coli* wall, resulting in its perforation which lead to the cell death [47], given that their direct contact with bacteria facilitated their dissolution at cell-NP interface and thus, enhanced their antibacterial effects [48].

In this regard, cationic-NPs have been reported as potential nanoparticles to cause pronounced disruption of plasma membrane integrity leading to mitochondrial and lysosomal damage, and production of autophagosomes, more than anionic-NPs [49]. Also, nonphagocytic cells were demonstrated to ingest cationic NPs to higher extent than anionic NPs [50]. Taking into account their surface charge nanoparticles can influence the selectivity and efficacy of drug delivery and imaging by selecting either a phagocytic or non-phagocytic pathway [51].

To evaluate cellular uptake of polymeric nanoparticles composed of poly-(lactic-co-glycolic acid) (PLGA) and coated with PVA and vitamin E TPGS, human colon adenocarcinoma cells were exposed in vitro [52]. It was found that vitamin E TPGS-coated PLGA NPs showed 1.4 folds higher cellular uptake than that of PVA-coated PLGA nanoparticles [53]. Cryo-SEM and

TEM analysis confirmed the cellular internalization, indicating that NPs surface modification with vitamin E TGPS and PLGA enhance loading of chemotherapeutic agents to be administered through oral route for cancer therapy [54].

2.3. Oxidative stress induced via nanoparticle and nanomaterial cellular uptake

Oxidative damage, caused in consequence of cellular internalization of NPs, has been considered one of the main causes of NPs cytotoxicity [55]. ZnO, CuO and Ag NPs, were already reported as ROS levels disruptors in aquatic microorganisms [56].

Also, the toxic effects of ROS generating potential of engineered Ag NPs were tested on different strains of recombinant *Escherichia coli* mutants and that of wild type strain [57]. The results showed that mutant strains of recombinant type were 15 folds more responsive to Ag NPs than the wild type [58]; although, analogous effects were observed for Ag ions [59]. In eukaryotes, induction of ROS by Ag NPs exposure was also observed in zebrafish *Danio rerio* [60].

It is well known that ions of redox-active metals, including Cu, may yield free radicals via the Fenton-type reaction and inflict intracellular oxidative stress [61, 62]. The Cu (II) ions can be transformed to Cu (I) ions in the presence of biological reducing agents such as ascorbic acid or glutathione (GSH), and consequently generate reactive hydroxyl radicals from hydrogen peroxide [63]. Using luminescent bacterial tests in recombinant *Escherichia coli* strains, ROS generating potential of aqueous suspensions of CuO NPs was demonstrated [58]. It has also been shown that CuO NPs induced oxidative stress and DNA damage in these recombinant bacteria strains at low subtoxic concentrations of 0.1 mg Cu/L [58].

In vivo investigation of the exposure effects of ZnO NPs on isolated rat liver mitochondria showed increased mitochondrial membrane permeability prone to energy dissipation [64], ROS levels production caused impairment of the mitochondrial respiratory chain and even apoptosis [65]. Similarly, the experiments on isolated human liver cells in-vitro showed ROS triggered mitochondrial pathway resulting apoptosis and cell death [65].

Moreover, it was also demonstrated a concentration dependent genotoxicity derived from exposure of several nanoparticles composed of α -alumina, β -alumina, SiO_2 , SbO_3 , and Fe_2O_3 through generation of excess ROS, which led to DNA strand breakage [66]. DNA cleavage, an indicator of irreversible completion of apoptosis, occurred in organisms exposed to 500 $\mu\text{g/kg}$ of the above-mentioned precipitated nanoparticles [67]. Moreover, cleavage of inter-nucleosomal DNA ladder bands occurred upon exposure to 500 $\mu\text{g/kg}$ of γ -alumina and α -alumina [66].

3. Conclusion

Nano-scale particles and materials applications are increasing significantly and thus their potential environmental and human health risks urges consideration. Analytical methods are required to reliably detect and characterize nanoparticles and nanomaterials in the natural environments, and their properties once interacting with complex matrices such as air,

soil and water, as well as food and consumer products. Advances in engineering particles and materials testing, either qualitative or quantitative determinations, should be pursued to ensure a safety profile, including the development of standard guidelines. In addition to the toxicological studies, various uptake paths have to be investigated, including dermal, oral and intestinal routes, as well as bioaccumulation trials and potential long-term effects consideration. Research into new analytical methods is further required to address the special properties for the establishment of ecological and human health risk factors.

Disclosure statement

No conflicts of interest.

Author details

Subas Chandra Dinda

Address all correspondence to: subas.dinda@rediffmail.com

School of Pharmacy, C.H.S., Mekelle University, Mekelle, Ethiopia

References

- [1] Krol S, Macrez R, Docagne F, Defer G, Laurent S, Rahman M, Hajipour MJ, Kehoe PG, Mahmoudi M. Therapeutic benefits from nanoparticles: The potential significance of nanoscience in diseases with compromise to the blood brain barrier. *Chemical Reviews*. 2013;**113**(3):1877-1903
- [2] Dykman LA, Khlebtsov NG. Uptake of engineered gold nanoparticles into mammalian cells. *Chemical Reviews*. 2014;**114**(2):1258-1288
- [3] Singh R, Lillard JW Jr. Nanoparticle-based targeted drug delivery. *Experimental and Molecular Pathology*. 2009;**86**(3):215-223
- [4] Shi H, Magaye R, Castranova V, Zhao J. Titanium dioxide nanoparticles: A review of current toxicological data. *Particle and Fibre Toxicology*. 2013;**10**:15. DOI: 10.1186/1743-8977-10-15
- [5] Hendren CO, Mesnard X, Dröge J, Wiesner MR. Estimating production data for five engineered nanomaterials as a basis for exposure assessment. *Environmental Science & Technology*. 2011;**45**(7):2562-2569
- [6] Robichaud CO, Uyar AE, Darby MR, Zucker LG, Wiesner MR. Estimates of upper bounds and trends in nano-TiO₂ production as a basis for exposure assessment. *Environmental Science & Technology*. 2009;**43**(12):4227-4233

- [7] Kim JI, Park HG, Chang KH, Nam DH, Yeo MK. Trophic transfer of nano-TiO₂ in a paddy microcosm: A comparison of single-dose versus sequential multi-dose exposures. *Environmental Pollution*. 2016;**212**:316-324. DOI: 10.1016/j.envpol.2016.01.076
- [8] Borgå K, Fisk AT, Hargrave B, Hoekstra PF, Swackhamer D, Muir DCG. Bioaccumulation factors for PCBs revisited. *Environmental Science & Technology*. 2005;**39**:4523-4532
- [9] Yeo M-K, Nam D-H. Influence of different types of nanomaterials on their bioaccumulation in a paddy microcosm: A comparison of TiO₂ nanoparticles and nanotubes. *Environmental Pollution*. 2013;**178**:166-172
- [10] Chandler JL. Toxicity and bioaccumulation of nanomaterial in aquatic species. *Journal of the U.S. SJWP*. 2007. DOI: 10.2175/SJWP(2007)1:01
- [11] McTeer J, Dean AP, White KN, Pittman JK. Bioaccumulation of silver nanoparticles into *Daphnia magna* from a fresh water algal diet and the impact of phosphate availability. *Journal Nanotoxicology*. 2014;**8**(3):305-316. DOI: 10.3109/17435390.2013.778346
- [12] Ates M, Demir V, Adiguzel R, Arslan Z. Bioaccumulation, subacute toxicity, and tissue distribution of engineered titanium dioxide nanoparticles in goldfish (*Carassius auratus*). *Journal of Nanomaterials*. 2013; Article ID 460518:1-6. DOI: 10.1155/2013/460518
- [13] Hanna SK, Miller RJ, Lenihan HS. Accumulation and toxicity of copper oxide engineered nanoparticles in a marine mussel. *Nanomaterials*. 2014;**4**:535-547. DOI: 10.3390/nano4030535
- [14] Zhu M-T, Feng W-Y, Wang Y, Wang B, Wang M, Ouyang H, Zhao Y-L, Chai Z-F. Particokinetics and extrapulmonary translocation of Intratracheally instilled ferric oxide nanoparticles in rats and the potential health risk assessment. *Toxicological Sciences*. 2009;**107**(2):342-351
- [15] Hussain T, Tan B, Yin Y, Blachier F, Tossou MCB, Rahu N. Oxidative stress and inflammation: What polyphenols can do for us? *Oxidative Medicine and Cellular Longevity*. 2016;**2016**:1-9 <http://dx.doi.org/10.1155/2016/7432797>
- [16] Oberdörster G et al. Nanotoxicology: An emerging discipline evolving from studies of ultrafine particles. *Environmental Health Perspectives*. 2005;**113**(7):823-839. PMC 1257642 . PMID 16002369. DOI: 10.1289/ehp.7339
- [17] Li N, Sioutas C, Cho A, et al. Ultrafine particulate pollutants induce oxidative stress and mitochondrial damage. *Environmental Health Perspectives*. 2003;**111**(4):455-460. PMC 1241427. PMID 12676598. DOI: 10.1289/ehp.6000
- [18] Geiser Met al. Ultrafine particles cross cellular membranes by nonphagocytic mechanisms in lungs and in cultured cells. *Environmental Health Perspectives*. 2005;**113**(11):1555-1560. PMC 1310918. PMID 16263511. DOI: 10.1289/ehp.8006
- [19] Raghunath A, Perumal E. Metal oxide nanoparticles as antimicrobial agents: A promise for the future. *International Journal of Antimicrobial Agents*. 2017;**49**(2):37-152. DOI: 10.1016/j.ijantimicag.2016.11.011

- [20] Park S-Y, Choi J. Geno- and ecotoxicity evaluation of silver nanoparticles in fresh water crustacean *Daphnia magna*. Environmental Engineering Research. 2010;**15**(1):23-27. DOI: 10.4491/eer.2010.15.1.428
- [21] Heinlaan M, Kahru A, Kasemets K, Arbeille B, Prensier G, Dubourguier H-C. Changes in the *Daphnia magna* midgut upon ingestion of copper oxide nanoparticles: A transmission electron microscopy study. Water Research. 2011;**45**:179-190. DOI: 10.1016/j.watres.2010.08.026
- [22] Scown TM, Santaos EM, Johnston BD, GAiser B, Baalousha M, Mitov S, Lead JR, Stone V, Fernandes TF, Jepson M, van Aerle R, Tyler CR. Effects of aqueous exposure to silver nanoparticles of different sizes in rainbow trout. Toxicological Sciences. 2010;**115**(2):521-534. DOI: 10.1093/toxsci/kfq076
- [23] AshaRani PV, Wu YL, Gong Z, Valiyaveetil S. Toxicity of silver nanoparticles in zebrafish models. Nanotechnology. 2008;**19**(25):279-290. DOI: 10.1088/0957-4484/19/255102
- [24] Oukarroum A, Bras S, Perreault F, Popovic R. Inhibitory effects of silver nanoparticles in two green algae, *Chlorella vulgaris* and *Dunaliella tertiolecta*. Ecotoxicology and Environmental Safety. 2012;**78**(1):80-85. DOI: 10.1016/j.ecoenv.2011.11.012
- [25] Pan L, Zhang H. Metallothionein, antioxidant enzymes and DNA strand breaks as biomarkers of Cd exposure in a marine crab, *Charybdis japonica*. Comparative Biochemistry and Physiology, Part C. 2006;**144**(1):67-75. DOI: 10.1016/j.cbpc.2006.06.001
- [26] Walters CR, Cheng P, Pool E, Somerset V. Effect of temperature on oxidative stress parameters and enzyme activity in the tissues of cape river crab (*Potamonautes perlatus*) following exposure to silver nanoparticles (AgNP). Journal of Toxicology and Environmental Health, Part A. 2016;**79**(2):1-11. DOI: 10.1080/15287394.2015.1106357
- [27] Ivask A, Juganson K, Bondarenko O, Mortimer M, Aruoja V, Kasemets K, Blinova I, Heinlaan M, Slaveykova V, Kahru A. Mechanisms of toxic action of Ag, ZnO and CuO nanoparticles to selected ecotoxicological test organisms and mammalian cells in vitro: A comparative review. Nanotoxicology. 2014;**8**(S1):57-71. DOI: 10.3109/17435390.2013.855831
- [28] Magrez A, Kasas S, Salicio V, Pasquier N, Seo JW, Celio M, Catsicas S, Schwaller B, Forro L. Cellular toxicity of carbon-based nanomaterials. Nano Letters. 2006;**6**(6):1121-1125
- [29] Herzog E, Casey A, Lyng FM, Chambers G, Byrne HJ, Davoren M. A new approach to the toxicity testing of carbon-based nanomaterials the clonogenic assay. Toxicology Letters. 2007;**174**(1-3):49-60
- [30] Principi E, Girardello R, Bruno A, Manni I, Gini E, Pagani A, Grimaldi A, Ivaldi F, Congiu T, De Stefano D, Piaggio G, de Eguileor M, Noonan DM, Albini A. Systemic distribution of single-walled carbon nanotubes in a novel model: Alteration of biochemical parameters, metabolic functions, liver accumulation, and inflammation in vivo. International Journal of Nanomedicine. 2016;**11**:4299-4316. DOI: 10.2147/IJN.S109950

- [31] Yang ST, Wang X, Jia G, Gu Y, Wang T, Nie H, Ge C, Wang H, Liu Y. Long-term accumulation and low toxicity of single-walled carbon nanotubes in intravenously exposed mice. *Toxicology Letters*. 2008;**181**(7):182-189
- [32] Lin W, Huang YW, Zhou XD, Ma Y. In vitro toxicity of silica nanoparticles in human lung cancer cells. *Toxicology and Applied Pharmacology*. 2006;**217**(3):252-259
- [33] Bondarenko O, Ivask A, Kakin A, Kurvet I, Kahru A. Particle cell contact enhances antibacterial activity of silver nanoparticles. *PLoS One*. 2013;**8**:e64060
- [34] Boenigk J, Beisser D, Zimmermann S, Bock C, Jakobi J, Grabner D, Großmann L, Rahmann S, Barcikowski S, Sures B. Effects of silver nitrate and silver nanoparticles on a planktonic community: General trends after short-term exposure. *PLoS One*. 2014;**9**(4):e95340. DOI: 10.1371/journal.pone.0095340
- [35] Hoheisel SM, Diamond S, Mount D. Comparison of nanosilver and ionic silver toxicity in *Daphnia magna* and *Pimephales promelas*. *Environmental Toxicology and Chemistry*. 2012;**31**:2557-2563
- [36] Bondarenko O, Ivask A, Kakin A, Kahru A. Sub-toxic effects of CuO nanoparticles on bacteria: Kinetics, role of Cu ions and possible mechanisms of action. *Environmental Pollution*. 2012;**169**:81-89
- [37] Panyam J, Labhasetwar V. Biodegradable nano-particles for drug and gene delivery to cells and tissue. *Advanced Drug Delivery Reviews*. 2003;**55**(3):329-347
- [38] Grabowski N, Hillaireau H, Vergnaud J, Tsapis N, Pallardy M, Kerdine-Rö me S, Fattal E. Surface coating mediates the toxicity of polymeric nanoparticles towards human-like macrophages. *International Journal of Pharmaceutics*. 2015;**482**(1-2):75-83
- [39] Applerot G, Lipovsky A, Dror R, Perkas N, Nitzan Y, Lubart R, Gedanken A. Enhanced antibacterial activity of nanocrystalline ZnO due to increased ROS-mediated cell injury. *Advanced Functional Materials*. 2009;**19**:842-852
- [40] Brayner R, Ferrari-Iliou R, Brivois N, Djediat S, Benedetti MF, Fievet F. Toxicological impact studies based on *Escherichia coli* bacteria in ultrafine ZnO nanoparticles colloidal medium. *Nano Letters*. 2006;**6**:866-870. DOI: 10.1021/nl052326h
- [41] Lok C-N, Ho C-M, Chen R, He Q-Y, Yu W-Y, Sun H, et al. Silver nanoparticles: Partial oxidation and antibacterial activities. *Journal of Biological Inorganic Chemistry*. 2007;**12**:527-534
- [42] Morones JR, Elechiguerra JL, Camacho A, Holt K, Kouri JB, Ramirez JT, Yacaman MJ. The bactericidal effect of silver nanoparticles. *Nanotechnology*. 2005;**16**:2346-2353
- [43] Gliga AR, Skoglund S, Wallinder IO, Fadeel B, Karlsson HL. Size-dependent cytotoxicity of silver nanoparticles in human lung cells: The role of cellular uptake, agglomeration and Ag release. *Particle and Fibre Toxicology*. 2014;**11**:11. DOI: 10.1186/1743-8977-11-11

- [44] Kumar A, Pandey AK, Singh SS, Shanker R, Dhawan A. Cellular uptake and mutagenic potential of metal oxide nanoparticles in bacterial cells. *Chemosphere*. 2011;**83**:1124-1132
- [45] Treuel L, Jiang X, Nienhaus GU. New views on cellular uptake and trafficking of manufactured nanoparticles. *Journal of the Royal Society Interface*. 2013;**10**(82):20120939. DOI: 10.1098/rsif.2012.0939
- [46] Huang Z, Zheng X, Yan D, Yin G, Liao X, Kang Y, Yao Y, Huang D, Hao B. Toxicological effect of ZnO nanoparticles based on bacteria. *Langmuir*. 2008;**24**(8):4140-4144. DOI: 10.1021/la7035949
- [47] Aminul HM. Experiment-based quantitative modeling for the antibacterial activity of silver nanoparticles. [Theses and Dissertations] 2017. p. 2445 <http://scholarworks.uark.edu/etd/2445>
- [48] Gogoi SK, Gopinath P, Paul A, Ramesh A, Ghosh SS, Chattopadhyay A. Green fluorescent protein-expressing *Escherichia coli* as a model system for investigating the antimicrobial activities of silver nanoparticles. *Langmuir*. 2006;**22**:9322-9328
- [49] Ruenraroengsak P, Tetley TD. Differential bioreactivity of neutral, cationic and anionic polystyrene nanoparticles with cells from the human alveolar compartment: Robust response of alveolar type 1 epithelial cells. *Particle and Fibre Toxicology*. 2015;**12**:19. DOI: 10.1186/s12989-015-0091-7
- [50] Fleischer CC, Payne CK. Nanoparticle-cell interactions: Molecular structure of the protein corona and cellular outcomes. *Accounts of Chemical Research*. 2014;**47**(8):2651-2659. DOI: 10.1021/ar500190q
- [51] Fröhlich E. The role of surface charge in cellular uptake and cytotoxicity of medical nanoparticles. *International Journal of Nanomedicine*. 2012;**7**:5577-5591
- [52] Feng SS, Zeng W, Teng Lim Y, Zhao L, Yin Win K, Oakley R, Hin Teoh S, Hang Lee RC, Pan S. Vitamin E TPGS-emulsified poly (lactic-co-glycolic acid) nanoparticles for cardiovascular restenosis treatment. *Nanomedicine (London, England)*. 2007;**2**(3):333-344
- [53] Win KY, Feng S-S. Effects of particle size and surface coating on cellular uptake of polymeric nanoparticles for oral delivery of anticancer drugs. *Biomaterials*. 2005;**26**(15):2713-2722
- [54] Win KY, Feng SS. In vitro and in vivo studies on vitamin E TPGS emulsified poly (D,L-lactic-co-glycolic acid) nanoparticles for paclitaxel formulation. *Biomaterials*. 2006;**27**(10):2285-2222
- [55] Blokhina O, Virolainen E, Fagerstedt KV. Antioxidants, oxidative damage and oxygen deprivation stress: A review. *Annals of Botany (London)*. 2003;**91**:179-194
- [56] Von Moos N, Slaveykova VI. Oxidative stress induced by inorganic nanoparticles in bacteria and aquatic microalgae – State of the art and knowledge gaps. *Nanotoxicology*. 2014;**8**(6):605-630

- [57] Jensen KF. The *Escherichia coli* K-12 “wild types” W3110 and MG1655 have an rph frame-shift mutation that leads to pyrimidine starvation due to low pyrE expression levels. *Journal of Bacteriology*. 1993;**175**(11):3401-3407. DOI: 10.1128/jb.175.11.3401-3407.1993
- [58] Ivask A, Bondarenko O, Jephthina N, Kahru A. Profiling of the reactive oxygen species-related ecotoxicity of CuO, ZnO, TiO₂, silver, and fullerene nanoparticles using a set of recombinant luminescent *Escherichia coli* strains: Differentiating the impact of particles and solubilised metals. *Analytical and Bioanalytical Chemistry*. 2010;**398**:701-716
- [59] Ivask A, ElBadawy A, Kaweeteerawat C, Boren D, Fischer H, Zhaoxia CHC, Liu R, Tolaymat T, Telesca D, Zink JJ, Cohen Y, Holden PA, Godwin HA. Toxicity mechanisms in *Escherichia coli* vary for silver nanoparticles and differ from ionic silver. *ACS Nano*. 2014;**8**(1):374-386. DOI: 10.1021/nn4044047
- [60] Choi O, Hu Z. Size dependent and reactive oxygen species related nanosilver toxicity to nitrifying bacteria. *Environmental Science & Technology*. 2008;**42**:4583-4588
- [61] Valko M, Morris H, Cronin MTD. Metals, toxicity and oxidative stress. *Current Medicinal Chemistry*. 2005;**12**:1161-1208
- [62] Thomas C, Mackey MM, Diaz AA, Cox DP. Hydroxyl radical is produced via the fenton reaction in submitochondrial particles under oxidative stress: Implications for diseases associated with iron accumulation. *Redox Report*. 2009;**14**(3):102-108
- [63] Vidrio E, Jung H, Anastasio C. Generation of hydroxyl radicals from dissolved transition metals in surrogate lung fluid solutions. *Atmospheric Environment*. 2008;**42**(18):4369-4379
- [64] Li J-h, Liu X-r, Zhang Y, Tian F-f, Zhao G-y, Yu Q-l-y, Jiang F-l, Liu Y. Toxicity of nano zinc oxide to mitochondria. *Toxicology Research*. 2012;**1**:137-144. DOI: 10.1039/C2TX20016C
- [65] Arora S, Jain J, Rajwade JM, Paknikar KM. Cellular responses induced by silver nanoparticles: In vitro studies. *Toxicology Letters*. 2008;**179**:93-100
- [66] Oberholster PJ, Musee N, Noth A-M, Chelule PK, Focke WW, Ashton PJ. Assessment of the effect of nanomaterials on sediment-dwelling invertebrate *Chironomus tentans* larvae. *Ecotoxicology and Environmental Safety*. 2011;**74**:416-423. DOI: 10.1016/j.ecoenv.2010.12.012
- [67] Lim HK, Asharani PV, Prakash Hande M. Enhanced genotoxicity of silver nanoparticles in DNA repair deficient mammalian cells. *Frontiers in Genetics*. 2012;**3**:104. DOI: 10.3389/fgene.2012.001049