We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

p53 and Vascular Dysfunction: MicroRNA in Endothelial Cells

Munekazu Yamakuchi, Sushil Panta and Teruto Hashiguchi

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.75461>

Abstract

In many cancer cells, p53 gene is mutated and accumulated, which is considered as a mechanistical target of tumorigenesis. The role of p53 in non-cancerous cells has been focused on, since p53 activation diversely affects as human diseases, including vascular dysfunctions. p53 regulates vascular events, including vascular inflammation and senescence as well as cardiac dysfunction. Many researchers also have paid attention to the role of noncoding RNAs (ncRNAs), especially small-sized microRNAs (miRNAs) for the last decade and their noble biological cellular functions have been discovered. miRNAs expressed in endothelial cells (endothelial miRNAs) have been shown to control vascular events. Firstly, the importance of p53 in a variety of vascular events, such as vascular inflammation and senescence, are summarized. Secondly, the way to regulate miRNAs by p53 and the involvement of miRNAs on p53 function are demonstrated. Finally, several endothelial miRNAs that have important roles are focused on. The aim of this chapter is to understand the role of p53 in vascular diseases in the view of endothelial cell biology and the contribution of miRNAs related to p53.

Keywords: endothelial cells, miRNAs, p53, Dicer, Drosha

1. Introduction

1.1. p53 overview

p53 is one of well-known tumor suppressor protein and plays crucial roles in inhibiting tumor progression [1]. Tumor suppression by p53 might be carried out mostly through genotoxic stress, however, recent studies revealed that p53 is activated by oncogene activation, oxidized

lipoproteins, and hypoxic condition [2]. In general, p53 is rapidly degraded by the interaction with MDM2 and these stimuli increase p53 levels and activate antiproliferative or proapoptotic responses via downstream signaling molecules [3]. The structure of p53 consists of amino terminal transactivation domain linked to the DNA-binding domain by proline-rich region (**Figure 1**) [4]. The DNA-binding domain on the other end is bound to the tetramerization domain by another proline-rich residue and this tetramerization domain is linked to carboxyl terminus [5]. The core domain (residue 94–312) is naturally unstable and is prone to have mutation [6, 7]. Once bound to the DNA, the whole structure closes around the DNA double helix. The whole process is facilitated by flexible proline-rich region between the core and the tetramerization domain [5]. Although the expression of p53 is in lower level during normal condition, upon activation, p53 increases its level along with the increase of its half-life [8] and gets translocated to the nucleus [9]. p53 is activated mainly by any signals that could damage the DNA [10]. Further p53 undergoes phosphorylation, acetylation, methylation, ubiquitination or SUMOylation to exert its respective activity [11]. p53 interacts with p300/CBP to get acetylated which stimulates the binding of p53 to the DNA, however, p53 requires only p300 but not CBP for the well-known G1 arrest [12]. Two members of p53 family, p73 and p63, are also involved in this p53 world [13], which are not mentioned in this chapter. Regulation and function of p53 in cancer really become complex.

1.2. p53 and endothelial cells

In the complex network of cellular signaling, p53 is a transcription factor that plays an important role in controlling angiogenesis and it is a hub for cellular signaling [14]. p53 itself controls angiogenesis by taking cells under apoptosis or by downregulating mediators of angiogenesis [15]. The role of p53 in vasculature is the same as the other tissues, including cell cycle, apoptosis, senescence, and angiogenesis.

Mice genetically deleted p53 can develop normally, however, these p53 knockout mice had spontaneous tumors [16]. Conditional knockout of p53 in endothelial cells improves angiogenesis of hindlimb ischemia mice model [17]. When mice were fed diet with high calorie, p53 expression increased in endothelial cells [18]. High calorie diet impaired the activation of endothelial nitric oxide synthase (eNOS), which was restored in endothelial p53 disruption. Knockdown of p53 in endothelial cells increased the expression of eNOS and thrombomodulin in vitro [19]. Therefore, accumulating evidence suggested that p53 is one of the key transcriptional factors to regulate endothelial cell function.

Figure 1. Linear structure of p53. p53 consists of 393 amino acid sequence. The protein is divided into the following domains. The transcriptional activation domain 1/2 (TAD1/2), DNA-binding domain, and the tetramerization domains (Tet) are lysine-rich basic C-terminal domain (CTD). PRR, proline-rich region; L, the linker region; Tet, tetramerization domains.

2. p53 and vascular function

Normal cells including endothelial cells keep p53 levels quite low. Low grade upregulation of p53 is not apoptotic but it is engaged in other functions like inhibition of endothelial cell migration through downregulation of beta-3 integrin [20] and inhibition of cell survival by causing reversible cell cycle arrest [21]. The modulation of p53 varies vascular function, such as vascular inflammation, senescence, and remodeling.

2.1. Vascular inflammation

Vascular inflammation leads to form atherosclerotic lesions, in which many cells are orchestrating [22]. Role of p53 in atherosclerosis has been investigated by many researchers. Guevara et al. performed the experiments using double knockout mice with apolipoprotein E (apoE) and p53. This apoE^{-/-}, p53^{-/-} double knockout mice fed with high fat diet showed significant increase of bulky, hypercellular lesion in aorta, suggesting that p53 is involved in atherosclerotic change [23]. van Vlijmen et al. demonstrated that the role of p53 in subendothelial macrophages is one of the major components of atherosclerosis [24]. This study indicated that deficiency of p53 in macrophage increased atherosclerotic lesions. Oxidative stress induces p53 accumulation in human macrophage, which is prevented by nitric oxide (NO) [25]. NO blocked the secretion of von Willebrand factor in endothelial cells and inhibited vascular inflammation [26]. The molecular mechanism by which p53 regulates atherosclerosis has been aggressively investigated.

2.2. Senescence

Aging is an independent risk factor for atherosclerosis-related diseases and impairment of vascular function is involved in systematic senescence. The molecular difference between senescence and cell death is not an easy question. Disturbed blood flow (d-flow) causes atherosclerosis. Heo KS et al. identified protein kinase zeta (PKC zeta) as a d-flow-activated protein in endothelial cells [27]. d-Flow promotes endothelial cells apoptosis through p53 SUMOylation. Apoptosis in aortic endothelial cells by d-flow decreased in p53^{-/-} mice compared to wild type mice. Endothelial cells constitutively express Nox2 and Nox4, two important isoforms of catalytic subunit of NADPH, which are a major source of reactive oxygen species. Nox2 especially affects endothelial cell cycle arrest and cell death by modulating p53 and p21cip1 [28]. In turn, cellular senescence is a stress-induced phenomenon as well. Senescent cells delay or lose the ability to proliferate. In endothelial cells, hydrogen peroxide or frequent passage induces cellular senescence via p53 and NAD-dependent deacetylase sirtuin-1 (SIRT1) [29]. The expression of endothelial SIRT-1 is reduced during aging process [30, 31]. Reduced SIRT-1 in endothelial cells accumulates genomic instability, resulting in p53 activation and promoting more senescence [32, 33]. AMPK and mTOR signaling is thought to be important for endothelial aging [34, 35]. These molecules are also connected to p53 signals, suggesting p53 as a key regulator of senescence of endothelial cells.

2.3. Vascular and heart remodeling

Vascular remodeling is a process of structural change of vascular walls, involving changes of cellular function, including growth and death. In this process, p53 is an important player.

Chronic hypoxia promotes pulmonary vascular remodeling, causing pulmonary hypertension. Mizuno S et al. demonstrated that p53 suppress hypoxia-induced pulmonary arterial remodeling and pulmonary arterial smooth muscle cell proliferation [36]. Kruppel-like factor 4 (KLF4) controls vascular smooth muscle cell proliferation through p53 induction [37]. Cardiac remodeling and development occur during embryogenesis but stop in postnatal life due to the reduction of the genes responsible for cell cycle progression and growth factors [38, 39]. For remodeling, new cardiomyocytes are derived for pre-existing cardiomyocytes. The rate of the pre-existing cells differentiation is very low (less than 1% per year) and it decreases with age [40] and lesser than 50% of the cells are replaced during a lifespan [41]. One important molecule for cardiomyocyte division is survivin [42]. Downregulation of survivin contributes to cardiac development in spinal muscular atrophy mice model [43]. Survivin is negatively regulated by p53. Survivin expression was downregulated at mRNA and protein level by p53 through histone acetylation. While overexpression of survivin inhibited p53-induced apoptosis [44]. One of MAPK, p38, has been shown to be an important molecule that negatively regulates cell cycle in cardiomyocyte cell [45]. Treatment with FGF1 and p38 inhibitor enhanced heart regeneration by increasing cardiomyocyte proliferation and angiogenesis [46]. Repression of cyclin D1 result into downregulation of cardiac cell proliferation [47] and C reactive protein, besides downregulating cyclin D1 has been shown to accumulate and phosphorylate p53 which leads to cell cycle arrest [48]. Since p53 controls actin cytoskeleton through mechanoresponsive molecules, remodeling may be processing via p53 in mechanical environment-dependent manner.

3. p53 and miRNA in endothelial cells

3.1. miRNA overview: general information

MicroRNAs (miRNAs) are small noncoding RNAs (about 20–24 nucleotides in length) that controls gene expressions mainly by binding to 3' untranslated region (3' UTR) of their messenger RNAs (mRNAs). The biogenesis of miRNAs in animals is very unique (**Figure 2**). Primary miRNAs (pri-miRNAs) are transcribed from miRNA-encoding genes. miRNAs are encoded in any place; some are located on protein-coding region, and some are in noncoding region or intron [49]. The pri-miRNAs are cleaved into hairpin-structured small size RNAs (precursor miRNAs; pre-miRNAs) by microprocessor complex containing RNase III, Drosha and DiGeorge critical region 8 (DGCR8) [50]. Exportin 5 (XPO5) and Ran-GTP transported pre-miRNAs into the cytoplasm from the nucleus, then pre-miRNAs are cleaved in double-stranded smaller RNAs (miRNA duplexes) by another RNase III, Dicer [51]. One of the strand (mature miRNAs) are incorporated into miRNA-induced silencing complex containing Argonaute 2 (Ago2) and transactivation response RNA-binding protein (TRBP) in human and these miRNAs are ready to bind to target mRNAs [52] (**Figure 2**).

How do miRNAs inhibit the expression of target protein? In general, miRNAs use two ways of silencing: repression of translation and mRNA decay [53]. The seed sequence of miRNA (2–7 position from 5' end) can bind to 3'UTR of target mRNA with incomplete match in animals. This miRNA-mRNA binding leads to repress the translation or destroy miRNA [54]. More than 60% of protein is regulated by miRNAs in human [55, 56]. Therefore, miRNAs are involved in modifying ubiquitous cellular functions.

Figure 2. p53 regulation of miRNA biogenesis.

3.2. miRNAs and p53

The relationship between miRNA and cancer was first described by Calin GA et al. in 2002. They described downregulation of miR-15a and miR-16-1 in B cell chronic lymphocytic leukemia patients [57]. miRNAs associated with cancer are called ‘oncomiR’, which have been identified in many types of cancer [58]. Some oncomiRs decrease in cancerous tissue. In contrast, increased oncomiRs are also found in cancer, which in case inhibit tumor suppressor genes, following the proliferation of cancer cells. The proto-oncogene c-Myc is a transcriptional factor and dysregulation of c-Myc was found in many cancers. The studies for regulation of transcriptional factors by miRNAs have been started since O’Donnell et al. identified mir-17-92, a polycistronic miRNA transcript that yields six individual miRNA components, as c-Myc-regulated miRNAs in human B cell line [59]. It was not hard to assume that the next target of ‘transcription factors’ regulating miRNAs in cancer was p53.

3.3. Direct regulation of miRNAs by p53

In 2007, several articles about p53 regulation of miRNAs have been published independently from different research groups. All these studies revealed that p53 upregulated the expression of miR-34 family in different cells [60–64]. The miR-34 family comprises three members (**Figure 4C**). miR-34a is generated from the large transcript on chromosome 1p36 and miR-34b and miR-34c are generated from bicistronic transcript on chromosome 11q23 [65]. Though the expression levels of miR-34a, -34b, and -34c were not consistent in non-small cell lung cancers (NSCLCs) compared to the adjacent normal tissue, lower expressions of three miR-34 family members are lower in many cancer cell lines; H1299 (lung cancer), MCF-7 (breast cancer), U-2OS (osteosarcoma), HCT116 (colon cancer), and many pancreatic cancer cell lines such as PANC1 [60, 64]. In addition to cancer, p53 regulates miR-34 family in non-cancerous cells, such as mouse embryonic fibroblasts (MEFs) and human fetal lung fibroblasts (IMR-90 cells) [63].

Forced expression of miR-34 family induce growth arrest and apoptosis in a variety of cell lines, whatever cancers or non-cancerous cells [62]. A lot of target genes of miR-34 family have been identified, including cyclin E2 (CCNE2), cyclin-dependent kinase 4 (CDK4) and the hepatocyte growth factor receptor (MET), B cell CLL/lymphoma 2 (BCL2), baculoviral IAP repeat-containing 3 (BIRC3), and decoy receptor 3 (DcR3 also known as TNFRSF6B). Many miRNAs directly induced by p53 have been identified in cancer cell lines. As described above, miR-34a might be the most fascinating one. Among these p53-induced miRNAs, several miRNAs that affect endothelial function are demonstrated in **Figure 5A**.

3.3.1. *miR-34 family*

The expression of miR-34 family, which consists of miR-34a, -34b, and -34c are induced by p53 activation [60, 64]. In many cancers, miR-34a-promoted apoptosis as described in the previous section. In primary normal human cells, miR-34 family can change cellular senescence. A series of miRNAs, including miR-34a, were upregulated in hydrogen peroxide-induced premature senescence in human fibroblasts [66]. There are two human p53 isoforms, p53 beta which lacks C-terminal oligomerization domain and delta133 p53 which lacks N-terminal transactivation and proline-rich domains. Human fibroblasts (MBC-5 and WI-38) at early passage had many delta133 p53 but not p53 beta. In contrast, p53 beta expressed well in fibroblasts at late passage. In fibroblasts, miR-34a control replicative cellular senescence and delta133 p53 repressed miR-34a expression, extending cellular replicative lifespan [67].

Aging of endothelial cells is one of the factors for cardiovascular diseases. miR-34a, expressed relatively higher in late-passage endothelial cells, modulated endothelial cell survival and senescence [30]. Overexpression of miR-34a triggers endothelial senescence mainly by blocking SIRT1. In mice, miR-34a expression also increases in heart and spleen from older ones. Endothelial progenitor cells (EPCs) are essential for many physiological processes such as wound healing. miR-34a impairs EPC-mediated angiogenesis by increasing the number of senescent EPC probably through SIRT1 inhibition [68]. SIRT1, the major target of miR-34a, was known to deacetylate p53. Activation of p53 increased miR-34a expression, which inhibit SIRT1 expression, causing accumulation of acetylated p53. Acetylated p53 induces cell cycle arrest and apoptosis, and this increase of p53 activity induced more miR-34a expression. This suggests that p53 – miR-34a – SIRT1 works as a positive feedback loop (**Figure 3B**) [69].

Notch signaling has crucial role in artery-vein differentiation, blood vessel sprouting, and branching. Dysregulation of Notch signaling causes cardiovascular diseases [70]. miR-34a could regulate Notch signaling pathway in vascular inflammation. In the case of placental dysfunction, miR-34a exacerbated vascular endothelial inflammation via suppression of regulator of calcineurin 1 (RCAN1) [71]. Shear stress is one of the central regulators of endothelial inflammatory responses. The expression of miR-34a decreased by anti-inflammatory physiological high shear stress, in turn, inflammatory oscillatory shear stress-induced miR-34a expression in endothelial cells [72]. Increased miR-34a promoted acetylation of NF- κ B p65 subunit (Lys310) by downregulating SIRT1, which lead to upregulate vascular cell adhesion molecule-1 (VCAM-1) and intercellular adhesion molecule-1 (ICAM-1) protein expression. miR-34a also contributed to shear stress-induced EPC differentiation through a novel target Forkhead box j2 (Foxj2) [73]. The important molecules targeting by endothelial miR-34a are listed in **Figure 3A**.

Figure 3. The role of miR-34a in regulating endothelial functions. (A) miR-34a target in endothelial cells. (B) miR-34a – p53 feedback loop.

3.3.2. miR-103 and miR-107

P53 positively regulates expressions of miR-103 and miR-107 in colorectal cancer cell lines [74]. miR-107 is encoded within an intron of the gene for pantothenate kinase enzyme 1, PANK1, while miR-103 is produced from primary miRNAs which are on miR-103-1 and miR-103-2 locus (within introns of PANK2 and PANK3, respectively) (**Figure 4A**). The seed sequences of miR-103 and miR-107 are the same, therefore, these miRNAs should have similar function [75]. miR-103 and miR-107 (miR-103/107) were originally recognized as a key regulator of metabolism and a hypoxia responsible miRNA [76]. The levels of miR-103/107 increased in liver of obese mice, ob/ob mice and diet-induced obese (DIO) mice, and knockdown of miR-103/107 improved insulin sensitivity [77]. Caveolin-1 was one of miR-103/107 targets that altered the level of insulin receptor on lipid rafts.

miR-103/107 also affected angiogenesis as members of hypoxia-responsive microRNAs (HRMs) induced by HIF1 α under hypoxia in endothelial cells [78, 79]. The crucial proteins for miRNA biogenesis, Dicer-1 and Ago-1, were identified as miR-107 and miR-103/107 targets, respectively. In both cases, miR-107 provided translational de-suppression of vascular endothelial growth factor (VEGF) mRNA and increased VEGF expression. AGO1 levels regulated by miR-103/107 were associated with higher survival rate in human hepatocellular carcinoma [78]. Antagomir-107 decreased the number of capillaries in ischemic boundary zone after permanent middle cerebral artery occlusion (pMCAO) in rats, which was caused by miR-107 – Dicer-1 – VEGF axis [79].

In sepsis, miR-107 plays an important role in endothelial cells. One of the major complications of sepsis is the development of acute kidney injury (AKI) [80]. Septic AKI activates renal endothelial cells and leads to inflammation and breakdown of endothelial barrier in kidney [81]. Wang et al. isolated circulating endothelial cells (CECs) from septic AKI patients and prepared CEC-conditioned media. Human tubule epithelial cells (HK2 cells) treated with this CEC-conditioned media became more apoptotic, which was regulated by miR-107 [82].

Figure 4. Scheme of miRNA cluster (miR-103/107, miR-200/141, miR-34, and miR-17-92). (A) miR-103/107 cluster. (B) miR-200/141 cluster. (C) miR-34 cluster. (D) miR-17-92 cluster.

In brain, miR-107 is enriched in neuron and the expression of miR-107 decreased in cerebral cortical gray matter of patients with Alzheimer's disease (AD) [83]. The authors demonstrated that beta-site amyloid precursor protein-cleaving enzyme 1 (BACE1) was identified as a miR-107 target. The cerebrovascular deposition of the amyloid beta protein, the key molecule in Alzheimer's disease, causes the disruption of blood-brain barrier (BBB) and brain microvascular endothelial cell dysfunction [84]. Another studies showed that miR-107 prevented amyloid beta-induced endothelial cells dysfunction by targeting endophilin-1 [85]. In a transgenic mouse model of AD, miR-107 expression in brain was lower compared to that in wild type mice [86]. Cofilin, which maintains the structure and function of cytoskeleton, was proved to be regulated by miR-107 in this mouse model. These data from AD patients and mice model suggest that relative high level of miR-107 in neurons and endothelial cells might negatively control the onset and progression of AD.

3.3.3. miR-143/145

miR-143 and miR-145 forms a bicistronic cluster (miR-143/145 cluster) in 5q33.1. The miR-143/145 cluster has been recognized as a tumor suppressor [87]. In cervical cancer, overexpression of Musashi RNA-binding protein 2 (MSI-2) correlated with poor survival. MSI-2 was repressed by p53 regulated miRNAs, miR-143 and miR-107, resulting in the prevention from proliferation and invasion of cervical cancer cells [88]. miR-143 and miR-145 have also potential roles in differentiation of vascular smooth muscle cells [89, 90]. The expression of miR-143/145 cluster decreased in aortic aneurysms and coronary artery diseases [91, 92]. miR-145-5p controls vascular neointimal lesion formation in balloon-injured rat carotid arteries [93]. The expression of miR-145 was upregulated in the lung of bone morphogenetic protein receptor type 2 (BMPR2)-deficient mice and pulmonary arterial hypertension (PAH) patients [94]. Deng L. et al. identified transcriptional factors that regulate miR-143 and miR-145 expression in the promoter of miR-143/145 cluster [95]. Each miRNA in this cluster has

each function; however, how this cluster or miR-143 and miR145 independently regulated has not been fully understood yet.

Shear stress suppressed angiotensin-converting enzyme (ACE) expression and increased miR-143/145 levels in HUVEC [96]. The authors have shown that shear stress elicited the AMP-activated protein kinase alpha2 (AMPK α 2)-dependent phosphorylation of p53 (Serine 15), and that p53 downregulation prevented the shear stress induced decrease in ACE expression. Since overexpression of miR-143/145 decreased ACE expression, AMPK α 2 suppresses ACE expression through p53 activation and upregulation of miR-143/145 in EC. AMPK α 2 knockout mice showed higher ACE levels and impaired bradykinin-induced vasodilation compared to wild type mice. In streptozotocin-induced diabetes mellitus (DM) mice model, phosphorylation of p53 and miR-143/145 expression increased, leading to the decrease of ACE expression. Therefore, miR-143 and miR-145 may contribute to the vascular events in atherosclerosis and DM.

miR-143 itself has been studied for the role of VSMCs as well as miR-145. The expression of miR-143 decreased in proliferating hemangiomas and miR-143 overexpression suppressed cell viability and proliferation of hemangioma-derived endothelial cells [97]. Bai Y et al. showed that miR-143 is upregulated in the brain microvessels of methamphetamine-treated mice [98]. Knockdown of miR-143 protected brain-blood barrier (BBB) damage-related vascular dysfunction by methamphetamine exposure. They identified an apoptosis inducing molecule, p53 upregulated modulator of apoptosis (PUMA), as a target of miR-143. Since the expression of miR-143 was regulated by p53 and miR-143 decreased PUMA, miR-143 might act for negative feedback of p53 signaling.

3.3.4. Others: miR-192, miR-200 family, and miR-194

Dysregulation of redox balance affects vascular homeostasis. Hydrogen peroxide treatment significantly increased miR-192 levels, which were prevented by p53 knockdown in endothelial cells [99]. Overexpression of miR-192 inhibited endothelial cell growth. Another study has shown that miR-200 family and miR-141 were upregulated in HUVEC exposed to hydrogen peroxide and in skeletal muscle in acute hindlimb ischemia mice model [100]. miRNA-200 family consists of two clusters, one encodes miR-200b, miR-200a, and miR-429 from chromosome 1 (1p33.36) and the other has miR-200c and miR-141 from chromosome 2 (12p13.31) (**Figure 4B**). These miRNAs share the similar seed sequence and mostly target the same genes. miR-200 family targets ZEB1 and ZEB2, affecting endothelial cell proliferation and senescence as well as epithelial-mesenchymal transition (EMT). Astrocytes are involved in controlling central nerve system (CNS) damage. During repair process of CNS, astrocytes undergo phenotypic changes into endothelial cells. This astrocyte-endothelial cell transition was modulated by a p53 inducible miRNA, miR-194. Therefore, miR-194 could promote angiogenesis in CNS.

3.4. Regulation of miRNA biogenesis by p53

There are two mechanisms by which p53 regulates miRNA production - control of miRNA transcription and modulation of miRNA maturation. These representative miRNAs regu-

Figure 5. Endothelial miRNA and p53. (A) Endothelial miRNAs regulated by p53. (B) Endothelial miRNAs directly target p53.

lated by p53 are summarized in Section 4.3. Several miRNAs can modulate the process of miRNA biogenesis. The impact of miRNA biogenesis by transcriptional factors has first reported about transforming growth factor beta (TGF-beta) signaling in 2008 [101]. TGF-beta family orchestrates biological processes in vascular development [102]. Davis et al. demonstrated that smads, downstream transcriptional factors of TGF-beta signaling, play a critical role in processing miRNAs by the RNase III-type protein Drosha in nucleus [101]. Similarly, p53 affects the maturation process of miRNAs. The nuclear RNase III Drosha complex contains Drosha, DiGeorge syndrome critical region gene 8 (DGCR8), and the DEAD box RNA helicases, such as p68 and p72 (DDX5 and DDX17, respectively) [103]. P53 interacts with Drosha through p68, facilitating the process of primary miRNAs into precursor miRNAs [104]. Maturation of some precursor miRNAs from primary miRNAs, such as miR-16-1, miR-143, miR-145, and miR-206, is promoted by Doxorubicin stimulated wild type p53 in colon cancer cell lines, HCT-116. Association between a set of miRNAs and Ago2 protein was controlled by p53 [105]. Activated p53 interacts with AGO2 to affect incorporation of let-7 family members. Moreover, p53 induced RNA-binding-motif protein 38 (RBM38) that determined target mRNA selection with miRNAs [106]. Interestingly, Rbm38 deficient mice were likely to accelerate senescence and prone to spontaneous tumors [107]. All these studies had no data using endothelial cells; however, basic insights would be connected to the future study about p53 and miRNAs in the cardiovascular research.

3.5. Regulation of p53 by miRNAs

Many miRNAs regulate p53 directly and indirectly. Endothelial miRNAs can target p53. More than 20 miRNAs that modulated p53 are reported. Among them, miR-92, miR-25, miR-214, and miR-638 play important roles in endothelial cell (**Figure 5B**).

3.5.1. *miR-125b*

TGF- β 2 induces endothelial-to-mesenchymal transition (EndMT) [108]. The expression of miR-125b in EndMT-derived fibroblast-like cells is significantly higher compared to that in the original mice endothelial cells [109]. In this experiment, miR-125b elevation was negatively associated with p53 expression after EndMT change. Since p53 is a direct target of miR-125b in several human cells, such as neuroblastoma cells and lung fibroblast cells, downregulated p53 by miR-125b possibly modulate TGF- β -induced profibrotic signaling in endothelial cells [110]. The expression of miR-125b was altered by cell-matrix adhesion in human mesenchymal stem cells (hMSCs) [111]. miR-125b targeted p53, which regulate survival of hMSCs and endogenous miR-125b increased during reprogramming of mouse embryo fibroblasts (MEFs) to induced pluripotent cells. Indeed, miR-125b was not increased by loss of cell adhesion in HUVEC. Sepsis damages endothelial cells, causing multiple organ failure [81]. Transfection of endothelial cells with miR-125b mimics attenuate LPS-induced ICAM-1 and VCAM-1 expression by inhibiting TRAF6 and NF- κ B activation [112]. Induction of miR-125 in mice heart attenuated cecal ligation (CLP)-induced sepsis as well and improved survival. These studies suggest that miR-125b regulates angiogenesis and vascular inflammation.

3.5.2. *miR-17-92 cluster*

Seven individual mature miRNAs (miR-17-5p, miR-17-3p, miR-18a, miR-19a, miR-19b, miR-20a, and miR-92a) are produced from primary miR-17-92, located in the open reading frame 25 (C13orf25) on chromosome 13 in human (**Figure 4D**). Mice knockout or overexpressing of miR-17-92 cluster died shortly after birth, suggested that the balance of miR-17-92 expression are involved in normal development [113, 114]. Originally, miR-17-92 has shown to be a highly conserved cluster, called oncomir-1, and extensively studied the molecular mechanism of tumorigenesis [115]. The roles of miR-17-92 for cardiovascular diseases have been investigated.

One miRNA of this cluster, miR-92a, blocked angiogenetic function in endothelial cells and inhibition of miR-92a by systemic administration of an antagomir-enhanced neovascularization and functional recovery from damaged tissue in hindlimb ischemic mice model [116]. Inhibition of miR-92a-enhanced endothelial cell proliferation and migration, probably through an increased phosphorylation of ERK1/2, JNK, and eNOS. miR-92a promotes pro-atherogenic changes in endothelial cells [117]. Disturbed flow increased miR-92a level in endothelial cells and miR-92a suppressed KLF2 and phosphatidic acid phosphatase type 2B (PPAP2B) that is involved in coronary artery disease (CAD) by genome-wide association studies (GWAS), driving inflammatory and adhesive endothelial phenotype [117]. Although no reports about miR-17-92 regulation of endothelial p53, according to accumulating data above, miR-17-92 may be involved in vascular events and p53 took some parts in them.

3.5.3. *Others: miR-98, miR-150, and miR-214 and beyond*

There are many miRNAs that directly regulate p53 in cancer; however, a few in endothelial cells. A variety of miRNAs, including miR-98, miR-150, and miR-214 has been shown to decrease p53 expression in cancer [118]. Hypoxia and reoxygenation conditions promote apoptosis and oxidized low-density lipoprotein (ox-LDL)-induced dysfunction of endothelial cells.

miR-98 rescues these phenomenon by targeting caspase-3 and lectin-like oxidized low-density lipoprotein receptor 1 (LOX-1), respectively [119, 120]. Stromal cell-derived factor 1 α (SDF-1 α) and its receptor CXCR4 control mobilization and migration of EPC. miR-150 decreased CXCR4 expression, leading to impaired EPC migration [121]. In mice studies, decreased miR-150 in EPC helped to revascularize the ischemic heart. miR-150 affected blood-brain barrier (BBB) permeability. Antagomir-150 treatment protected BBB, reduced infarct volume in post-stroke rat via angiopoietin receptor Tie-2 [122].

Targets of miRNAs are recognized by pairing between the seed sequence of miRNA and complementary sites in target mRNAs. There are many useful tools to search for miRNA target genes. Among them, Targetscan is one of the reliable resources many researchers are widely taking. Targetscan predicts four miRNAs, let-7, miR-22, miR122, and miR-150, which are broadly conserved among vertebrates to bind onto 3'UTR of human p53 mRNA (http://www.targetscan.org/vert_71/). Future studies could reveal the function of these miRNAs and their relationship to p53.

4. Conclusion

miRNAs are crucial regulators of gene expression for diverse physiological and pathological processes. Endothelial miRNAs have been intensively studied since Kuehbacher A et al. released that genetic knockout of Dicer and Drosha, miRNA-processing enzymes, inhibited capillary sprouting of endothelial cells and tube formation [123]. Recently Hratmann P et al. demonstrated that Dicer in endothelial cells promoted atherosclerosis and endothelial inflammation [124]. In contrast, p53 is involved in a variety of diseases, such as vascular remodeling, atherosclerosis, hypertension, and hypoxic pulmonary artery remodeling as well as cancer biology.

The importance of p53 and miRNAs in endothelial cells has been shown here. We demonstrated the regulation of endothelial miRNAs by p53 and the modulation of p53 by miRNAs in endothelial cells. These miRNAs play pivotal roles in vascular development and the onset of cardiovascular diseases. The ubiquitin E3 ligase Mdm2 stimulates p53 degradation, in turn, p53 promotes Mdm2 gene expression. Therefore, there is a negative feedback loop between p53 and Mdm2. miR-192, miR-194, and miR-215 targeted Mdm2 protein, which could disrupt this p53-Mdm2 feedback loop [125]. Future studies will unveil the complex and fascinating pathway and loop composed by p53 and miRNAs and develop therapeutic machinery of vascular diseases.

Author details

Munekazu Yamakuchi^{1*}, Sushil Panta² and Teruto Hashiguchi¹

*Address all correspondence to: munekazu@m.kufm.kagoshima-u.ac.jp

1 Department of Laboratory and Vascular Medicine, Graduate School of Medical and Dental Sciences, Kagoshima University, Kagoshima, Japan

2 School of Health and Allied Sciences, Faculty of Health Sciences, Pokhara University, Pokhara-Lekhnaath, Nepal

References

- [1] Vogelstein B, Kinzler KW. Cancer genes and the pathways they control. *Nature Medicine*. 2004;**10**(8):789-799. doi: 10.1038/nm1087. PubMed PMID: 15286780
- [2] Vousden KH, Prives C. Blinded by the light: The growing complexity of p53. *Cell*. 2009;**137**(3):413-431. DOI: 10.1016/j.cell.2009.04.037. PubMed PMID: 19410540
- [3] Gottlieb TM, Oren M. p53 and apoptosis. *Seminars in Cancer Biology*. 1998;**8**(5):359-368. PubMed PMID: 10101801
- [4] Fields S, Jang SK. Presence of a potent transcription activating sequence in the p53 protein. *Science*. 1990;**249**(4972):1046-1049. Epub 1990/08/31. PubMed PMID: 2144363
- [5] Joerger AC, Fersht AR. The tumor suppressor p53: From structures to drug discovery. *Cold Spring Harbor Perspectives in Biology*. 2010;**2**(6):a000919. Epub 2010/06/03. DOI: 10.1101/cshperspect.a000919. PubMed PMID: 20516128; PubMed Central PMCID: PMC2869527
- [6] Bullock AN, Henckel J, DeDecker BS, Johnson CM, Nikolova PV, Proctor MR, et al. Thermodynamic stability of wild-type and mutant p53 core domain. *Proceedings of the National Academy of Sciences of the United States of America*. 1997;**94**(26):14338-14342. Epub 1998/02/07. PubMed PMID: 9405613; PubMed Central PMCID: PMC24967
- [7] Olivier M, Eeles R, Hollstein M, Khan MA, Harris CC, Hainaut P. The IARC TP53 database: New online mutation analysis and recommendations to users. *Human Mutation*. 2002;**19**(6):607-614. Epub 2002/05/15. DOI: 10.1002/humu.10081. PubMed PMID: 12007217
- [8] Lakin ND, Jackson SP. Regulation of p53 in response to DNA damage. *Oncogene*. 1999;**18**:7644. DOI: 10.1038/sj.onc.1203015
- [9] Komarova EA, Zelnick CR, Chin D, Zeremski M, Gleiberman AS, Bacus SS, et al. Intracellular localization of p53 tumor suppressor protein in gamma-irradiated cells is cell cycle regulated and determined by the nucleus. *Cancer Research*. 1997;**57**(23):5217-5220. Epub 1997/12/11. PubMed PMID: 9393737
- [10] Harris SL, Levine AJ. The p53 pathway: Positive and negative feedback loops. *Oncogene*. 2005;**24**(17):2899-2908. Epub 2005/04/20. DOI: 10.1038/sj.onc.1208615. PubMed PMID: 15838523
- [11] Appella E, Anderson CW. Post-translational modifications and activation of p53 by genotoxic stresses. *European Journal of Biochemistry*. 2001;**268**(10):2764-2772. Epub 2001/05/19. PubMed PMID: 11358490
- [12] Gu W, Roeder RG. Activation of p53 sequence-specific DNA binding by acetylation of the p53 C-terminal Domain. *Cell*. 1997 Aug 22;**90**(4):595-606. DOI: 10.1016/S0092-8674(00)80521-8
- [13] Sheikh MS, Fornace AJ, Jr. Role of p53 family members in apoptosis. *Journal of Cellular Physiology*. 2000;**182**(2):171-181. DOI: 10.1002/(SICI)1097-4652(200002)182:2<171::AID-JCP5>3.0.CO;2-3. PubMed PMID: 10623880

- [14] Jin S, Levine AJ. The p53 functional circuit. *Journal of Cell Science*. 2001;**114**(Pt 23):4139-4140. Epub 2001/12/12. PubMed PMID: 11739646
- [15] Munshi N, Fernandis AZ, Cherla RP, Park IW, Ganju RK. Lipopolysaccharide-induced apoptosis of endothelial cells and its inhibition by vascular endothelial growth factor. *Journal of Immunology (Baltimore, MD, 1950)*. 2002;**168**(11):5860-5866. Epub 2002/05/23. PubMed PMID: 12023390
- [16] Donehower LA, Harvey M, Slagle BL, McArthur MJ, Montgomery Jr CA, Butel JS, et al. Mice deficient for p53 are developmentally normal but susceptible to spontaneous tumours. *Nature*. 1992;**356**(6366):215-221. DOI: 10.1038/356215a0. PubMed PMID: 1552940
- [17] Gogiraju R, Xu X, Bochenek ML, Steinbrecher JH, Lehnart SE, Wenzel P, et al. Endothelial p53 deletion improves angiogenesis and prevents cardiac fibrosis and heart failure induced by pressure overload in mice. *Journal of the American Heart Association*. 2015 Feb 24;**4**(2). pii: e001770. DOI: 10.1161/JAHA.115.001770. PubMed PMID: 25713289; PubMed Central PMCID: PMCPMC4345879
- [18] Yokoyama M, Okada S, Nakagomi A, Moriya J, Shimizu I, Nojima A, et al. Inhibition of endothelial p53 improves metabolic abnormalities related to dietary obesity. *Cell Reports*. 2014;**7**(5):1691-1703. DOI: 10.1016/j.celrep.2014.04.046. PubMed PMID: 24857662
- [19] Kumar A, Kim CS, Hoffman TA, Naqvi A, Dericco J, Jung SB, et al. p53 Impairs endothelial function by transcriptionally repressing Kruppel-Like Factor 2. *Arteriosclerosis, Thrombosis, and Vascular Biology*. 2011;**31**(1):133-141. DOI: 10.1161/ATVBAHA.110.215061. PubMed PMID: 20947822; PubMed Central PMCID: PMCPMC3064482
- [20] Panta S, Yamakuchi M, Shimizu T, Takenouchi K, Oyama Y, Koriyama T, et al. Low grade inflammation inhibits VEGF induced HUVECs migration in p53 dependent manner. *Biochemical and Biophysical Research Communications*. 2017 Feb 5;**483**(2):803-809. DOI: 10.1016/j.bbrc.2016.12.096. PubMed PMID: 27998768
- [21] Lukin DJ, Carvajal LA, Liu WJ, Resnick-Silverman L, Manfredi JJ. p53 Promotes cell survival due to the reversibility of its cell-cycle checkpoints. *Molecular Cancer Research: MCR*. 2015;**13**(1):16-28. Epub 2014/08/28. DOI: 10.1158/1541-7786.mcr-14-0177. PubMed PMID: 25158956; PubMed Central PMCID: PMCPMC4312522
- [22] Ross R. Cell biology of atherosclerosis. *Annual Review of Physiology*. 1995;**57**:791-804. DOI: 10.1146/annurev.ph.57.030195.004043. PubMed PMID: 7778883
- [23] Guevara NV, Kim HS, Antonova EI, Chan L. The absence of p53 accelerates atherosclerosis by increasing cell proliferation in vivo. *Nature Medicine*. 1999;**5**(3):335-339. DOI: 10.1038/6585. PubMed PMID: 10086392
- [24] van Vlijmen BJ, Gerritsen G, Franken AL, Boesten LS, Kockx MM, Gijbels MJ, et al. Macrophage p53 deficiency leads to enhanced atherosclerosis in APOE*3-Leiden transgenic mice. *Circulation Research*. 2001;**88**(8):780-786. PubMed PMID: 11325869
- [25] Heinloth A, Brune B, Fischer B, Galle J. Nitric oxide prevents oxidised LDL-induced p53 accumulation, cytochrome c translocation, and apoptosis in macrophages via guanylate cyclase stimulation. *Atherosclerosis*. 2002;**162**(1):93-101. PubMed PMID: 11947902

- [26] Matsushita K, Morrell CN, Cambien B, Yang SX, Yamakuchi M, Bao C, et al. Nitric oxide regulates exocytosis by S-nitrosylation of N-ethylmaleimide-sensitive factor. *Cell*. 2003;**115**(2):139-150. PubMed PMID: 14567912; PubMed Central PMCID: PMCPMC2846406
- [27] Heo KS, Lee H, Nigro P, Thomas T, Le NT, Chang E, et al. PKCzeta mediates disturbed flow-induced endothelial apoptosis via p53 SUMOylation. *Journal of Cell Biology*. 2011;**193**(5):867-884. DOI: 10.1083/jcb.201010051. PubMed PMID: 21624955; PubMed Central PMCID: PMCPMC3105539
- [28] Li JM, Fan LM, George VT, Brooks G. Nox2 regulates endothelial cell cycle arrest and apoptosis via p21cip1 and p53. *Free Radical Biology and Medicine*. 2007;**43**(6):976-986. DOI: 10.1016/j.freeradbiomed.2007.06.001. PubMed PMID: 17697942; PubMed Central PMCID: PMCPMC2889611
- [29] Kim KS, Kang KW, Seu YB, Baek SH, Kim JR. Interferon-gamma induces cellular senescence through p53-dependent DNA damage signaling in human endothelial cells. *Mechanisms of Ageing and Development*. 2009;**130**(3):179-188. DOI: 10.1016/j.mad.2008.11.004. PubMed PMID: 19071156
- [30] Ito T, Yagi S, Yamakuchi M. MicroRNA-34a regulation of endothelial senescence. *Biochemical and Biophysical Research Communications*. 2010;**398**(4):735-740. DOI: 10.1016/j.bbrc.2010.07.012. PubMed PMID: 20627091
- [31] Donato AJ, Magerko KA, Lawson BR, Durrant JR, Lesniewski LA, Seals DR. SIRT-1 and vascular endothelial dysfunction with ageing in mice and humans. *Journal of Physiology*. 2011;**589**(Pt 18):4545-4554. DOI: 10.1113/jphysiol.2011.211219. PubMed PMID: 21746786; PubMed Central PMCID: PMCPMC3208223
- [32] Luo J, Nikolaev AY, Imai S, Chen D, Su F, Shiloh A, et al. Negative control of p53 by Sir2alpha promotes cell survival under stress. *Cell*. 2001;**107**(2):137-148. PubMed PMID: 11672522
- [33] Vaziri H, Dessain SK, Ng Eaton E, Imai SI, Frye RA, Pandita TK, et al. hSIR2(SIRT1) functions as an NAD-dependent p53 deacetylase. *Cell*. 2001;**107**(2):149-159. PubMed PMID: 11672523
- [34] Thoreen CC, Sabatini DM. AMPK and p53 help cells through lean times. *Cell Metabolism*. 2005;**1**(5):287-288. DOI: 10.1016/j.cmet.2005.04.009. PubMed PMID: 16054073
- [35] Feng Z, Zhang H, Levine AJ, Jin S. The coordinate regulation of the p53 and mTOR pathways in cells. *Proceedings of the National Academy of Sciences of the United States of America*. 2005;**102**(23):8204-8209. DOI: 10.1073/pnas.0502857102. PubMed PMID: 15928081; PubMed Central PMCID: PMCPMC1142118
- [36] Mizuno S, Bogaard HJ, Kraskauskas D, Alhussaini A, Gomez-Arroyo J, Voelkel NF, et al. p53 gene deficiency promotes hypoxia-induced pulmonary hypertension and vascular remodeling in mice. *American Journal of Physiology. Lung Cellular and Molecular Physiology*. 2011;**300**(5):L753-L761. DOI: 10.1152/ajplung.00286.2010. PubMed PMID: 21335523

- [37] Yoshida T, Kaestner KH, Owens GK. Conditional deletion of Kruppel-like factor 4 delays downregulation of smooth muscle cell differentiation markers but accelerates neointimal formation following vascular injury. *Circulation Research*. 2008;**102**(12):1548-1557. DOI: 10.1161/CIRCRESAHA.108.176974. PubMed PMID: 18483411; PubMed Central PMCID: PMC2633447
- [38] Porrello ER, Olson EN. A neonatal blueprint for cardiac regeneration. *Stem Cell Research*. 2014;**13**(3, Part B):556-570. DOI: 10.1016/j.scr.2014.06.003
- [39] Chen HW, Yu SL, Chen WJ, Yang PC, Chien CT, Chou HY, et al. Dynamic changes of gene expression profiles during postnatal development of the heart in mice. *Heart*. 2004;**90**(8):927-934. DOI: 10.1136/hrt.2002.006734. PubMed PMID: PMC1768375
- [40] Senyo SE, Steinhauser ML, Pizzimenti CL, Yang VK, Cai L, Wang M, et al. Mammalian heart renewal by preexisting cardiomyocytes. *Nature*. 2013;**493**(7432):433-436. DOI: 10.1038/nature11682. PubMed PMID: PMC3548046
- [41] Bergmann O, Bhardwaj RD, Bernard S, Zdunek S, Barnabé-Heider F, Walsh S, et al. Evidence for cardiomyocyte renewal in humans. *Science (New York, NY)*. 2009;**324**(5923):98-102. DOI: 10.1126/science.1164680. PubMed PMID: PMC2991140
- [42] Levkau B, Schafers M, Wohlschlaeger J, von Wnuck Lipinski K, Keul P, Hermann S, et al. Survivin determines cardiac function by controlling total cardiomyocyte number. *Circulation* 2008;**117**(12):1583-1593. Epub 2008/03/12. DOI: 10.1161/circulationaha.107.734160. PubMed PMID: 18332262
- [43] Sheng L, Wan B, Feng P, Sun J, Rigo F, Bennett CF, et al. Downregulation of Survivin contributes to cell-cycle arrest during postnatal cardiac development in a severe spinal muscular atrophy mouse model. *Human Molecular Genetics*. 2018 Feb 1;**27**(3):486-498. DOI: 10.1093/hmg/ddx418
- [44] Mirza A, McGuirk M, Hockenberry TN, Wu Q, Ashar H, Black S, et al. Human survivin is negatively regulated by wild-type p53 and participates in p53-dependent apoptotic pathway. *Oncogene*. 2002;**21**(17):2613-2622. Epub 2002/04/20. DOI: 10.1038/sj.onc.1205353. PubMed PMID: 11965534
- [45] Engel FB, Schebesta M, Duong MT, Lu G, Ren S, Madwed JB, et al. p38 MAP kinase inhibition enables proliferation of adult mammalian cardiomyocytes. *Genes & Development*. 2005;**19**(10):1175-1187. DOI: 10.1101/gad.1306705. PubMed PMID: PMC1132004
- [46] Engel FB, Hsieh PCH, Lee RT, Keating MT. FGF1/p38 MAP kinase inhibitor therapy induces cardiomyocyte mitosis, reduces scarring, and rescues function after myocardial infarction. *Proceedings of the National Academy of Sciences of the United States of America*. 2006;**103**(42):15546-15551. DOI: 10.1073/pnas.0607382103. PubMed PMID: PMC1622860
- [47] Toyoda M, Shirato H, Nakajima K, Kojima M, Takahashi M, Kubota M, et al. Jumonji down-regulates cardiac cell proliferation by repressing cyclin D1 expression. *Developmental Cell*. 2003;**5**(1):85-97. Epub 2003/07/11. PubMed PMID: 12852854

- [48] Choi JW, Lee KH, Kim SH, Jin T, Lee BS, Oh J, et al. C-reactive protein induces p53-mediated cell cycle arrest in H9c2 cardiac myocytes. *Biochemical and Biophysical Research Communications*. 2011;**410**(3):525-530. Epub 2011/06/18. DOI: 10.1016/j.bbrc.2011.06.016. PubMed PMID: 21679689
- [49] Axtell MJ, Westholm JO, Lai EC. Vive la difference: biogenesis and evolution of microRNAs in plants and animals. *Genome Biology*. 2011;**12**(4):221. DOI: 10.1186/gb-2011-12-4-221. PubMed PMID: 21554756; PubMed Central PMCID: PMC3218855
- [50] Lee Y, Ahn C, Han J, Choi H, Kim J, Yim J, et al. The nuclear RNase III Drosha initiates microRNA processing. *Nature*. 2003;**425**(6956):415-419. DOI: 10.1038/nature01957. PubMed PMID: 14508493
- [51] Ketting RF, Fischer SE, Bernstein E, Sijen T, Hannon GJ, Plasterk RH. Dicer functions in RNA interference and in synthesis of small RNA involved in developmental timing in *C. Elegans*. *Genes & Development*. 2001;**15**(20):2654-2659. DOI: 10.1101/gad.927801. PubMed PMID: 11641272; PubMed Central PMCID: PMC3212808
- [52] Bernstein E, Caudy AA, Hammond SM, Hannon GJ. Role for a bidentate ribonuclease in the initiation step of RNA interference. *Nature*. 2001;**409**(6818):363-366. DOI: 10.1038/35053110. PubMed PMID: 11201747
- [53] Iwakawa HO, Tomari Y. The functions of MicroRNAs: mRNA decay and translational repression. *Trends in Cell Biology*. 2015;**25**(11):651-665. DOI: 10.1016/j.tcb.2015.07.011. PubMed PMID: 26437588
- [54] Djuranovic S, Nahvi A, Green R. miRNA-mediated gene silencing by translational repression followed by mRNA deadenylation and decay. *Science*. 2012;**336**(6078):237-240. DOI: 10.1126/science.1215691. PubMed PMID: 22499947; PubMed Central PMCID: PMC3971879
- [55] Friedman RC, Farh KK, Burge CB, Bartel DP. Most mammalian mRNAs are conserved targets of microRNAs. *Genome Research*. 2009;**19**(1):92-105. DOI: 10.1101/gr.082701.108. PubMed PMID: 18955434; PubMed Central PMCID: PMC2612969
- [56] Bartel DP. MicroRNAs: Target recognition and regulatory functions. *Cell*. 2009;**136**(2):215-233. DOI: 10.1016/j.cell.2009.01.002. PubMed PMID: 19167326; PubMed Central PMCID: PMC3794896
- [57] Calin GA, Dumitru CD, Shimizu M, Bichi R, Zupo S, Noch E, et al. Frequent deletions and down-regulation of micro- RNA genes miR15 and miR16 at 13q14 in chronic lymphocytic leukemia. *Proceedings of the National Academy of Sciences of the United States of America*. 2002;**99**(24):15524-15529. DOI: 10.1073/pnas.242606799. PubMed PMID: 12434020; PubMed Central PMCID: PMC32137750
- [58] Hammond SM. RNAi, microRNAs, and human disease. *Cancer Chemotherapy and Pharmacology*. 2006;**58**(Suppl 1):s63-s68. DOI: 10.1007/s00280-006-0318-2. PubMed PMID: 17093929

- [59] O'Donnell KA, Wentzel EA, Zeller KI, Dang CV, Mendell JT. C-Myc-regulated microRNAs modulate E2F1 expression. *Nature*. 2005;**435**(7043):839-843. DOI: 10.1038/nature03677. PubMed PMID: 15944709
- [60] Chang TC, Wentzel EA, Kent OA, Ramachandran K, Mullendore M, Lee KH, et al. Transactivation of miR-34a by p53 broadly influences gene expression and promotes apoptosis. *Molecular Cell*. 2007;**26**(5):745-752. DOI: 10.1016/j.molcel.2007.05.010. PubMed PMID: 17540599; PubMed Central PMCID: PMCPMC1939978
- [61] Raver-Shapira N, Marciano E, Meiri E, Spector Y, Rosenfeld N, Moskovits N, et al. Transcriptional activation of miR-34a contributes to p53-mediated apoptosis. *Molecular Cell*. 2007;**26**(5):731-743. DOI: 10.1016/j.molcel.2007.05.017. PubMed PMID: 17540598
- [62] Tarasov V, Jung P, Verdoodt B, Lodygin D, Epanchintsev A, Menssen A, et al. Differential regulation of microRNAs by p53 revealed by massively parallel sequencing: miR-34a is a p53 target that induces apoptosis and G1-arrest. *Cell Cycle*. 2007;**6**(13):1586-1593. DOI: 10.4161/cc.6.13.4436. PubMed PMID: 17554199
- [63] He L, He X, Lim LP, de Stanchina E, Xuan Z, Liang Y, et al. A microRNA component of the p53 tumour suppressor network. *Nature*. 2007;**447**(7148):1130-1134. DOI: 10.1038/nature05939. PubMed PMID: 17554337; PubMed Central PMCID: PMCPMC4590999
- [64] Bommer GT, Gerin I, Feng Y, Kaczorowski AJ, Kuick R, Love RE, et al. p53-mediated activation of miRNA34 candidate tumor-suppressor genes. *Current Biology*. 2007;**17**(15):1298-1307. DOI: 10.1016/j.cub.2007.06.068. PubMed PMID: 17656095
- [65] He L, He X, Lowe SW, Hannon GJ. microRNAs join the p53 network--another piece in the tumour-suppression puzzle. *Nature Reviews. Cancer*. 2007;**7**(11):819-822. DOI: 10.1038/nrc2232. PubMed PMID: 17914404; PubMed Central PMCID: PMCPMC4053212
- [66] Maes OC, Sarojini H, Wang E. Stepwise up-regulation of microRNA expression levels from replicating to reversible and irreversible growth arrest states in WI-38 human fibroblasts. *Journal of Cellular Physiology*. 2009;**221**(1):109-119. DOI: 10.1002/jcp.21834. PubMed PMID: 19475566
- [67] Fujita K, Mondal AM, Horikawa I, Nguyen GH, Kumamoto K, Sohn JJ, et al. p53 Isoforms Delta133p53 and p53beta are endogenous regulators of replicative cellular senescence. *Nature Cell Biology*. 2009;**11**(9):1135-1142. DOI: 10.1038/ncb1928. PubMed PMID: 19701195; PubMed Central PMCID: PMCPMC2802853
- [68] Zhao T, Li J, Chen AF. MicroRNA-34a induces endothelial progenitor cell senescence and impedes its angiogenesis via suppressing silent information regulator 1. *American Journal of Physiology. Endocrinology and Metabolism*. 2010;**299**(1):E110-E116. DOI: 10.1152/ajpendo.00192.2010. PubMed PMID: 20424141; PubMed Central PMCID: PMCPMC2904051
- [69] Yamakuchi M, Lowenstein CJ. MiR-34, SIRT1 and p53: The feedback loop. *Cell Cycle*. 2009;**8**(5):712-715. DOI: 10.4161/cc.8.5.7753. PubMed PMID: 19221490
- [70] Gridley T. Notch signaling in vascular development and physiology. *Development*. 2007;**134**(15):2709-2718. DOI: 10.1242/dev.004184. PubMed PMID: 17611219

- [71] Yuan HY, Zhou CB, Chen JM, Liu XB, Wen SS, Xu G, et al. MicroRNA-34a targets regulator of calcineurin 1 to modulate endothelial inflammation after fetal cardiac bypass in goat placenta. *Placenta*. 2017;**51**:49-56. DOI: 10.1016/j.placenta.2017.01.128. PubMed PMID: 28292468
- [72] Fan W, Fang R, Wu X, Liu J, Feng M, Dai G, et al. Shear-sensitive microRNA-34a modulates flow-dependent regulation of endothelial inflammation. *Journal of Cell Science*. 2015;**128**(1):70-80. DOI: 10.1242/jcs.154252. PubMed PMID: 25395581
- [73] Cheng BB, Qu MJ, Wu LL, Shen Y, Yan ZQ, Zhang P, et al. MicroRNA-34a targets Forkhead box j2 to modulate differentiation of endothelial progenitor cells in response to shear stress. *Journal of Molecular and Cellular Cardiology*. 2014;**74**:4-12. DOI: 10.1016/j.yjmcc.2014.04.016. PubMed PMID: 24792364
- [74] Yamakuchi M, Lotterman CD, Bao C, Hruban RH, Karim B, Mendell JT, et al. P53-induced microRNA-107 inhibits HIF-1 and tumor angiogenesis. *Proceedings of the National Academy of Sciences of the United States of America*. 2010;**107**(14):6334-6339. DOI: 10.1073/pnas.0911082107. PubMed PMID: 20308559; PubMed Central PMCID: PMC2851979
- [75] Finnerty JR, Wang WX, Hebert SS, Wilfred BR, Mao G, Nelson PT. The miR-15/107 group of microRNA genes: Evolutionary biology, cellular functions, and roles in human diseases. *Journal of Molecular Biology*. 2010;**402**(3):491-509. DOI: 10.1016/j.jmb.2010.07.051. PubMed PMID: 20678503; PubMed Central PMCID: PMC2978331
- [76] Rottiers V, Naar AM. MicroRNAs in metabolism and metabolic disorders. *Nature Reviews. Molecular Cell Biology*. 2012;**13**(4):239-250. DOI: 10.1038/nrm3313. PubMed PMID: 22436747; PubMed Central PMCID: PMC2978331
- [77] Trajkovski M, Hausser J, Soutschek J, Bhat B, Akin A, Zavolan M, et al. MicroRNAs 103 and 107 regulate insulin sensitivity. *Nature*. 2011;**474**(7353):649-653. DOI: 10.1038/nature10112. PubMed PMID: 21654750
- [78] Chen Z, Lai TC, Jan YH, Lin FM, Wang WC, Xiao H, et al. Hypoxia-responsive miRNAs target argonaute 1 to promote angiogenesis. *The Journal of Clinical Investigation*. 2013;**123**(3):1057-1067. DOI: 10.1172/JCI65344. PubMed PMID: 23426184; PubMed Central PMCID: PMC3582133
- [79] Li Y, Mao L, Gao Y, Baral S, Zhou Y, Hu B. MicroRNA-107 contributes to post-stroke angiogenesis by targeting Dicer-1. *Scientific Reports*. 2015;**5**:13316. DOI: 10.1038/srep13316. PubMed PMID: 26294080; PubMed Central PMCID: PMC4543985
- [80] Jacobs R, Honore PM, Joannes-Boyau O, Boer W, De Regt J, De Waele E, et al. Septic acute kidney injury: The culprit is inflammatory apoptosis rather than ischemic necrosis. *Blood Purification*. 2011;**32**(4):262-265. DOI: 10.1159/000330244. PubMed PMID: 21860231
- [81] Peters K, Unger RE, Brunner J, Kirkpatrick CJ. Molecular basis of endothelial dysfunction in sepsis. *Cardiovascular Research*. 2003;**60**(1):49-57. PubMed PMID: 14522406

- [82] Wang S, Zhang Z, Wang J, Miao H. MiR-107 induces TNF-alpha secretion in endothelial cells causing tubular cell injury in patients with septic acute kidney injury. *Biochemical and Biophysical Research Communications*. 2017;**483**(1):45-51. DOI: 10.1016/j.bbrc.2017.01.013. PubMed PMID: 28063928
- [83] Wang WX, Rajeev BW, Stromberg AJ, Ren N, Tang G, Huang Q, et al. The expression of microRNA miR-107 decreases early in Alzheimer's disease and may accelerate disease progression through regulation of beta-site amyloid precursor protein-cleaving enzyme 1. *The Journal of Neuroscience*. 2008;**28**(5):1213-1223. DOI: 10.1523/JNEUROSCI.5065-07.2008. PubMed PMID: 18234899; PubMed Central PMCID: PMCPMC2837363
- [84] Gheorghiu M, Enciu AM, Popescu BO, Gheorghiu E. Functional and molecular characterization of the effect of amyloid-beta42 on an in vitro epithelial barrier model. *Journal of Alzheimer's Disease*. 2014;**38**(4):787-798. DOI: 10.3233/JAD-122374. PubMed PMID: 24072066
- [85] Wood H. Alzheimer disease: Fibrinogen links amyloid with vascular dysfunction. *Nature Reviews. Neurology*. 2010;**6**(8):413. DOI: 10.1038/nrneurol.2010.98. PubMed PMID: 20718101
- [86] Yao J, Hennessey T, Flynt A, Lai E, Beal MF, Lin MT. MicroRNA-related cofilin abnormality in Alzheimer's disease. *PLoS One*. 2010;**5**(12):e15546. DOI: 10.1371/journal.pone.0015546. PubMed PMID: 21179570; PubMed Central PMCID: PMCPMC3002958
- [87] Kent OA, McCall MN, Cornish TC, Halushka MK. Lessons from miR-143/145: The importance of cell-type localization of miRNAs. *Nucleic Acids Research*. 2014;**42**(12):7528-7538. DOI: 10.1093/nar/gku461. PubMed PMID: 24875473; PubMed Central PMCID: PMCPMC4081080
- [88] Dong P, Xiong Y, Hanley SJB, Yue J, Watari H. Musashi-2, a novel oncoprotein promoting cervical cancer cell growth and invasion, is negatively regulated by p53-induced miR-143 and miR-107 activation. *Journal of Experimental & Clinical Cancer Research*. 2017;**36**(1):150. DOI: 10.1186/s13046-017-0617-y. PubMed PMID: 29073938; PubMed Central PMCID: PMCPMC5659032
- [89] Cordes KR, Sheehy NT, White MP, Berry EC, Morton SU, Muth AN, et al. miR-145 and miR-143 regulate smooth muscle cell fate and plasticity. *Nature*. 2009;**460**(7256):705-710. DOI: 10.1038/nature08195. PubMed PMID: 19578358; PubMed Central PMCID: PMCPMC2769203
- [90] Boettger T, Beetz N, Kostin S, Schneider J, Kruger M, Hein L, et al. Acquisition of the contractile phenotype by murine arterial smooth muscle cells depends on the Mir143/145 gene cluster. *The Journal of Clinical Investigation*. 2009;**119**(9):2634-2647. DOI: 10.1172/JCI38864. PubMed PMID: 19690389; PubMed Central PMCID: PMCPMC2735940
- [91] Elia L, Quintavalle M, Zhang J, Contu R, Cossu L, Latronico MV, et al. The knockout of miR-143 and -145 alters smooth muscle cell maintenance and vascular homeostasis in mice: Correlates with human disease. *Cell Death and Differentiation*. 2009;**16**(12):1590-1598. DOI: 10.1038/cdd.2009.153. PubMed PMID: 19816508; PubMed Central PMCID: PMCPMC3014107

- [92] Lovren F, Pan Y, Quan A, Singh KK, Shukla PC, Gupta N, et al. MicroRNA-145 targeted therapy reduces atherosclerosis. *Circulation*. 2012;**126**(11 Suppl 1):S81-S90. DOI: 10.1161/CIRCULATIONAHA.111.084186. PubMed PMID: 22965997
- [93] Cheng Y, Liu X, Yang J, Lin Y, Xu DZ, Lu Q, et al. MicroRNA-145, a novel smooth muscle cell phenotypic marker and modulator, controls vascular neointimal lesion formation. *Circulation Research*. 2009;**105**(2):158-166. DOI: 10.1161/CIRCRESAHA.109.197517. PubMed PMID: 19542014; PubMed Central PMCID: PMC2728297
- [94] Caruso P, Dempsie Y, Stevens HC, McDonald RA, Long L, Lu R, et al. A role for miR-145 in pulmonary arterial hypertension: Evidence from mouse models and patient samples. *Circulation Research*. 2012;**111**(3):290-300. DOI: 10.1161/CIRCRESAHA.112.267591. PubMed PMID: 22715469
- [95] Deng L, Blanco FJ, Stevens H, Lu R, Caudrillier A, McBride M, et al. MicroRNA-143 activation regulates smooth muscle and endothelial cell crosstalk in pulmonary arterial hypertension. *Circulation Research*. 2015;**117**(10):870-883. DOI: 10.1161/CIRCRESAHA.115.306806. PubMed PMID: 26311719; PubMed Central PMCID: PMC4620852
- [96] Kohlstedt K, Trouvain C, Boettger T, Shi L, Fisslthaler B, Fleming I. AMP-activated protein kinase regulates endothelial cell angiotensin-converting enzyme expression via p53 and the post-transcriptional regulation of microRNA-143/145. *Circulation Research*. 2013; **112**(8):1150-1158. DOI: 10.1161/CIRCRESAHA.113.301282. PubMed PMID: 23476055
- [97] Huang C, Huang J, Ma P, Yu G. microRNA-143 acts as a suppressor of hemangioma growth by targeting Bcl-2. *Gene*. 2017;**628**:211-217. DOI: 10.1016/j.gene.2017.07.046. PubMed PMID: 28716710
- [98] Bai Y, Zhang Y, Hua J, Yang X, Zhang X, Duan M, et al. Silencing microRNA-143 protects the integrity of the blood-brain barrier: Implications for methamphetamine abuse. *Scientific Reports*. 2016;**6**:35642. DOI: 10.1038/srep35642. PubMed PMID: 27767041; PubMed Central PMCID: PMC45073292
- [99] Fuschi P, Carrara M, Voellenkle C, Garcia-Manteiga JM, Righini P, Maimone B, et al. Central role of the p53 pathway in the noncoding-RNA response to oxidative stress. *Aging (Albany NY)*. 2017;**9**(12):2559-2586. DOI: 10.18632/aging.101341. PubMed PMID: 29242407; PubMed Central PMCID: PMC5764393
- [100] Magenta A, Cencioni C, Fasanaro P, Zaccagnini G, Greco S, Sarra-Ferraris G, et al. miR-200c is upregulated by oxidative stress and induces endothelial cell apoptosis and senescence via ZEB1 inhibition. *Cell Death and Differentiation*. 2011;**18**(10):1628-1639. DOI: 10.1038/cdd.2011.42. PubMed PMID: 21527937; PubMed Central PMCID: PMC3172120
- [101] Davis BN, Hilyard AC, Lagna G, Hata A. SMAD proteins control DROSHA-mediated microRNA maturation. *Nature*. 2008;**454**(7200):56-61. DOI: 10.1038/nature07086. PubMed PMID: 18548003; PubMed Central PMCID: PMC2653422
- [102] ten Dijke P, Arthur HM. Extracellular control of TGFbeta signalling in vascular development and disease. *Nature Reviews. Molecular Cell Biology*. 2007;**8**(11):857-869. DOI: 10.1038/nrm2262. PubMed PMID: 17895899

- [103] Gregory RI, Yan KP, Amuthan G, Chendrimada T, Doratotaj B, Cooch N, et al. The microprocessor complex mediates the genesis of microRNAs. *Nature*. 2004;**432**(7014):235-240. DOI: 10.1038/nature03120. PubMed PMID: 15531877
- [104] Suzuki HI, Yamagata K, Sugimoto K, Iwamoto T, Kato S, Miyazono K. Modulation of microRNA processing by p53. *Nature*. 2009;**460**(7254):529-533. DOI: 10.1038/nature08199. PubMed PMID: 19626115
- [105] Krell J, Stebbing J, Carissimi C, Dabrowska AF, de Giorgio A, Frampton AE, et al. TP53 regulates miRNA association with AGO2 to remodel the miRNA-mRNA interaction network. *Genome Research*. 2016;**26**(3):331-341. DOI: 10.1101/gr.191759.115. PubMed PMID: 26701625; PubMed Central PMCID: PMC4772015
- [106] Leveille N, Elkon R, Davalos V, Manoharan V, Hollingworth D, Oude Vrielink J, et al. Selective inhibition of microRNA accessibility by RBM38 is required for p53 activity. *Nature Communications*. 2011;**2**:513. DOI: 10.1038/ncomms1519. PubMed PMID: 22027593; PubMed Central PMCID: PMC3221330
- [107] Zhang J, Xu E, Ren C, Yan W, Zhang M, Chen M, et al. Mice deficient in Rbm38, a target of the p53 family, are susceptible to accelerated aging and spontaneous tumors. *Proceedings of the National Academy of Sciences of the United States of America*. 2014;**111**(52):18637-18642. DOI: 10.1073/pnas.1415607112. PubMed PMID: 25512531; PubMed Central PMCID: PMC4284600
- [108] Zeisberg EM, Kalluri R. Origins of cardiac fibroblasts. *Circulation Research*. 2010;**107**(11):1304-1312. DOI: 10.1161/CIRCRESAHA.110.231910. PubMed PMID: 21106947; PubMed Central PMCID: PMC3098499
- [109] Ghosh AK, Nagpal V, Covington JW, Michaels MA, Vaughan DE. Molecular basis of cardiac endothelial-to-mesenchymal transition (EndMT): Differential expression of microRNAs during EndMT. *Cellular Signalling*. 2012;**24**(5):1031-1036. DOI: 10.1016/j.cellsig.2011.12.024. PubMed PMID: 22245495; PubMed Central PMCID: PMC3298765
- [110] Le MT, Teh C, Shyh-Chang N, Xie H, Zhou B, Korzh V, et al. MicroRNA-125b is a novel negative regulator of p53. *Genes & Development*. 2009;**23**(7):862-876. DOI: 10.1101/gad.1767609. PubMed PMID: 19293287; PubMed Central PMCID: PMC2666337
- [111] Yu X, Cohen DM, Chen CS. miR-125b is an adhesion-regulated microRNA that protects mesenchymal stem cells from anoikis. *Stem Cells*. 2012;**30**(5):956-964. DOI: 10.1002/stem.1064. PubMed PMID: 22331826; PubMed Central PMCID: PMC3323671
- [112] Ma H, Wang X, Ha T, Gao M, Liu L, Wang R, et al. MicroRNA-125b prevents cardiac dysfunction in Polymicrobial sepsis by targeting TRAF6-mediated nuclear factor kappaB activation and p53-mediated apoptotic Signaling. *The Journal of Infectious Diseases*. 2016;**214**(11):1773-1783. DOI: 10.1093/infdis/jiw449. PubMed PMID: 27683819; PubMed Central PMCID: PMC47735
- [113] Ventura A, Young AG, Winslow MM, Lintault L, Meissner A, Erkland SJ, et al. Targeted deletion reveals essential and overlapping functions of the miR-17 through 92 family

- p>of miRNA clusters.
- Cell*
- . 2008;
- 132**
- (5):875-886. DOI: 10.1016/j.cell.2008.02.019. PubMed PMID: 18329372; PubMed Central PMCID: PMCPMC2323338
- [114] Lu Y, Thomson JM, Wong HY, Hammond SM, Hogan BL. Transgenic over-expression of the microRNA miR-17-92 cluster promotes proliferation and inhibits differentiation of lung epithelial progenitor cells. *Developmental Biology*. 2007;**310**(2):442-453. DOI: 10.1016/j.ydbio.2007.08.007. PubMed PMID: 17765889; PubMed Central PMCID: PMCPMC2052923
- [115] Mogilyansky E, Rigoutsos I. The miR-17/92 cluster: A comprehensive update on its genomics, genetics, functions and increasingly important and numerous roles in health and disease. *Cell Death and Differentiation*. 2013;**20**(12):1603-1614. DOI: 10.1038/cdd.2013.125. PubMed PMID: 24212931; PubMed Central PMCID: PMCPMC3824591
- [116] Bonauer A, Carmona G, Iwasaki M, Mione M, Koyanagi M, Fischer A, et al. MicroRNA-92a controls angiogenesis and functional recovery of ischemic tissues in mice. *Science* 2009;**324**(5935):1710-1713. DOI: 10.1126/science.1174381. PubMed PMID: 19460962
- [117] Wu C, Huang RT, Kuo CH, Kumar S, Kim CW, Lin YC, et al. Mechanosensitive PPAP2B regulates endothelial responses to atherorelevant hemodynamic forces. *Circulation Research*. 2015;**117**(4):e41-e53. DOI: 10.1161/CIRCRESAHA.117.306457. PubMed PMID: 26034042; PubMed Central PMCID: PMCPMC4522239
- [118] Deng G, Sui G. Noncoding RNA in oncogenesis: A new era of identifying key players. *International Journal of Molecular Sciences*. 2013;**14**(9):18319-18449. DOI: 10.3390/ijms140918319. PubMed PMID: 24013378; PubMed Central PMCID: PMCPMC3794782
- [119] Li HW, Meng Y, Xie Q, Yi WJ, Lai XL, Bian Q, et al. miR-98 Protects endothelial cells against hypoxia/reoxygenation induced-apoptosis by targeting caspase-3. *Biochemical and Biophysical Research Communications*. 2015;**467**(3):595-601. DOI: 10.1016/j.bbrc.2015.09.058. PubMed PMID: 26367177
- [120] Chen Z, Wang M, He Q, Li Z, Zhao Y, Wang W, et al. MicroRNA-98 rescues proliferation and alleviates ox-LDL-induced apoptosis in HUVECs by targeting LOX-1. *Experimental and Therapeutic Medicine*. 2017;**13**(5):1702-1710. DOI: 10.3892/etm.2017.4171. PubMed PMID: 28565756; PubMed Central PMCID: PMCPMC5443247
- [121] Rolland-Turner M, Goretti E, Bousquenaud M, Leonard F, Nicolas C, Zhang L, et al. Adenosine stimulates the migration of human endothelial progenitor cells. Role of CXCR4 and microRNA-150. *PLoS One*. 2013;**8**(1):e54135. DOI: 10.1371/journal.pone.0054135. PubMed PMID: 23326587; PubMed Central PMCID: PMCPMC3541240
- [122] Fang Z, He QW, Li Q, Chen XL, Baral S, Jin HJ, et al. MicroRNA-150 regulates blood-brain barrier permeability via Tie-2 after permanent middle cerebral artery occlusion in rats. *The FASEB Journal*. 2016;**30**(6):2097-2107. DOI: 10.1096/fj.201500126. PubMed PMID: 26887441
- [123] Kuehbach A, Urbich C, Zeiher AM, Dimmeler S. Role of dicer and Drosha for endothelial microRNA expression and angiogenesis. *Circulation Research*. 2007;**101**(1):59-68. DOI: 10.1161/CIRCRESAHA.107.153916. PubMed PMID: 17540974

- [124] Hartmann P, Zhou Z, Natarelli L, Wei Y, Nazari-Jahantigh M, Zhu M, et al. Endothelial dicer promotes atherosclerosis and vascular inflammation by miRNA-103-mediated suppression of KLF4. *Nature Communications*. 2016;**7**:10521. DOI: 10.1038/ncomms10521. PubMed PMID: 26837267; PubMed Central PMCID: PMC4742841
- [125] Pichiorri F, Suh SS, Rocci A, De Luca L, Taccioli C, Santhanam R, et al. Downregulation of p53-inducible microRNAs 192, 194, and 215 impairs the p53/MDM2 autoregulatory loop in multiple myeloma development. *Cancer Cell*. 2010;**18**(4):367-381. DOI: 10.1016/j.ccr.2010.09.005. PubMed PMID: 20951946; PubMed Central PMCID: PMC3561766