We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Quality of Life and Functionality in Patients with Flatfoot

Cristina Gonzalez-Martin,
Salvador Pita-Fernandez and Sonia Pertega-Diaz

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.76236>

Abstract

The objective of the study is to determine the prevalence of a plan, its impact on quality of life, dependence and functional limitation in a random population of 40 years and over. Cross-sectional study in a random population sample in Cambre (A Coruña-Spain) ($n = 835$) ($\alpha = 0.05$; precision = $\pm 3.4\%$). Anthropometric variables are studied, comorbidity (Charlson Score), foot functionality (FFI questionnaire), foot health questionnaire (FHSQ), quality of life (SF-36) and dependence on activities of daily living (Barthel index and Lawton). A logistic and linear multiple regression analysis was performed. The prevalence of flat feet was 26.62%. Patients with flat feet presented higher: age (65.73 ± 11.04 years), comorbidity index (0.92 ± 1.49), BMI (31.45 ± 5.55) and foot size ($25, 16 \pm 1.66$ cm). Having flat feet decreases the quality of life and function of the foot. The association of flat feet with age, Charlson index, and BMI and foot size was found. The SF-36, Barthel and Lawton questionnaires remained unchanged due to the presence of the flat foot, a difference between the FHSQ and FFI that were significantly sensitive.

Keywords: flat foot, quality of life, dependence, prevalence, functionality

---

1. Introduction

Flexible flatfoot is a common deformity in adults [1]. It is characterized by medial rotation and plantar flexion of the talus, eversion of the calcaneus, collapsed medial arch, and abduction of the forefoot [2].

In clinical practice flat foot may be diagnosed through different procedures, such as clinical diagnosis [3], radiological study [4] and footprint analysis [5].

Footprint analysis using a pedograph is a simple, swift, and cost-effective method. The three measurements habitually used in the diagnosis of flat foot using a pedograph are: Clarke's angle [6] the Chippaux-Smirak index [7] and the Staheli index [8].

Studies have found relation between these indices [9, 10] and their validity has been determined using diagnosis carried out with a podoscope on children as a reference group [11].

Prevalence changes with age, the type of population studied and the presence of other pathologies. Some studies show prevalence between 26.5% [12] and 19.0% [13] and other studies on patients with associated comorbidity report a prevalence of 37% [14].

Flat foot has been associated to family history, the use of footwear in infancy, obesity and urban residence [15], and it has also been associated with age [16], gender [17] and foot length [18].

The presence of flat foot has also been associated with the presence of different states of health [19], the presence of pain, and the fatigue in women [12]. Other studies, however, find no relationship of pain or functionality with the changes in the foot [20, 21].

We conducted this study, in order to determine the variables associated with the prevalence of flat foot in a random population sample, and the impact on quality of life, dependence, foot pain, disability and functional limitation, using specific and generic questionnaires.

2. Materials and methods

A cross-sectional study was conducted in a random population sample from 2009 to 2012 in Cambre (A Coruña-Spain).

The sample size was taken from people who lived in Cambre and were identified through the National Health System card census. People aged 40 and over were included who signed the informed consent.

The sample size is calculated of the total population of the municipality ($n = 23,649$) after stratification by age and gender. Finally, a total of 835 people were included in the study. This sample size ($n = 835$ people; 445 aged 40–64 years old and 390 aged 65 years and older) makes it possible to estimate the parameters of interest with a confidence of 95% ($\alpha = 0.05$) and a precision of $\pm 3.4\%$. The general characteristics of a different sample from the same population have already been described above [22].

For each person included in the study, the following variables were studied: anthropometric variables (age, gender, body mass index), study of chronic comorbid diseases (comorbidities) using the Charlson comorbidity index [23], quality of life (SF-36 questionnaire) [24], Foot Health Status Questionnaire (FHS) [25], Foot Function Index (FFI) [25] Barthel index [26],

Lawton index [27], podiatric examination and type of footwear. The podiatric examination was carried out by an experienced podiatrist.

The Charlson Index contains 19 categories of comorbidity, which are primarily defined using the ICD-9-CM diagnosis codes (a few procedure codes are also employed). Each category has an associated weight, taken from the original Charlson paper [20], which is based on the adjusted risk of one-year mortality. The overall comorbidity score reflects the cumulative increased likelihood of one-year mortality; the higher the score, the more severe the burden of comorbidity.

In order to study quality of life, the SF-36 health questionnaire was used, adapted and validated for Spain by Alonso et al. [21].

The questionnaire sf-36 is formed by 36 questions that evaluate the Physical Function, Physical Role, Corporal Pain, General Health, Vitality, Social Function, Emotional Role and Mental Health. The score scale varies from 0 to 100, with 100 the best state of health.

Foot Health Status Questionnaire (FHSQ) [22] is a health-related quality of life questionnaire and is specific to the foot, is divided into 4 domains that assess pain, functional capacity, footwear and overall health of the foot. The questionnaire does not provide an overall score. The score varies from 0 to 100, 0 is the worst state of health.

The questionnaire Foot function Index (FFI) [22] measures disability and pain in the feet.

The FFI consists of 23 items divided into 3 subscales: pain (9 items), disability (9 items) and functional limitation (5 items). To evaluate each item, it consists of a visual analog scale with values between 0 and 9, where 0 is the minimum score and 9 is the maximum score. To get the result, we must add all the scores made by the person and then divide this result by the maximum value that could reach. This result is then multiplied by 100 and rounded to integers. The final score will be between 0 and 100. Higher scores indicate worsening foot health and quality.

2.1. Flat foot diagnostic

The study of the footprint was obtained by a pedograph. Three measurements were used: Clarke's angle, the Chippaux-Smirak index, and the Staheli arch index [6–8].

The specific methods of measurements of these indexes was described previously [25].

For the study of the footwear, the type of footwear most used, the heel (flat, low, medium, high) and the shape (shoe, sporty, boot, clog) or type of closure (moccasin, zipper, buckle, drawstring).

2.2. Statistical analysis

A descriptive analysis of the variables collected in the study was carried out. The quantitative variables are expressed as mean $\pm$ standard deviation, median and range. The qualitative

variables are expressed as frequency (n) and percentage with the estimation of the corresponding 95% confidence interval.

The association between qualitative variables was estimated using the Chi-square test or Fisher's test as appropriate. The assumption of normality was checked by the Kolmogorov-Smirnov test, which determined the use of the Student's T test or the Mann-Whitney test for the comparison of two means.

2.3. Ethics

The study complies with the principles laid down in the Declaration of Helsinki. Informed consent was obtained from all the participants in the study. Confidentiality was preserved in accordance with the current Spanish Data Protection Law (15/1999). Patient and ethical review approval was obtained previously (code 2008/264 CEIC Galicia).

3. Results

The general characteristics of the sample studied, according to different variables are shown in **Table 1**. The mean age is 61.70 ± 11.60 years, with a prevalence of overweight of 42.2% and a median Charlson comorbidity index from 2.0.

People with flat feet use closed shoes (88.0%), followed by sports (3.8%). The most used heel was the medium heel (2–4 cm) (71.8%). The most used footwear style would be moccasin type (48.1%) followed by cord shoe (44.2%).

This study shows that the prevalence of flatfoot is 26.62% (**Table 2**).

The presence of flatfoot is significantly associated with bivariate analysis with: age, comorbidity, BMI and foot size. Among patients with flat feet, there was a higher mean age (65.73 years vs. 61.03 years), higher comorbidity (2.99 vs. 2.09), higher BMI (31.45 kg/m² vs. 28.4045 kg/m²) and have a greater average foot size (25.16 cm vs. 24.82 cm). They were not associated in the analysis bivariate with the presence of flat foot or forefoot width, or sex (**Table 3**).

After performing a multivariate logistic regression analysis, we observed that the variables that have an independent effect associated with the presence of flat feet are: BMI (OR = 1.137), age (OR = 1.029), mean foot size OR = 1.287) and comorbidity (OR = 1.217) (**Table 3**). That is, higher values of the different variables previously described increase the greater probability of flat foot.

If we study the area under the curve (AUC) to predict presence of flat feet according to each of the previously described variables, the most likely predictor is BMI (AUC = 0.683) and age (AUC = 0.614) (**Figure 1**).

Variables	n	Mean ± SD	Median	Minimum–maximum
Age (years)	835	61.70 ± 11.60	63	42–91
BMI (kg/m ²)	835	29.18 ± 4.74	28.65	19.13–64.09
Charlson comorbidity index	786	2.31 ± 1.89	2	0–14
	n	%	95% CI	
Gender				
Male	369/835	44.2%	(40.76;47.62)	
Female	466/835	55.8%	(52.38;59.34)	
Age groups				
<65 years	445/835	53.3%	(49.85;56.74)	
65 years and over	390/835	46.7%	(43.26;50.15)	
BMI categories				
Normal weight (18.5 kg/m ² ≤ BMI < 25 kg/m ²)	140/832	16.8%	(14.17;19.36)	
Overweight (25 kg/m ² ≤ BMI < 30 kg/m ²)	369/832	44.2%	(40.19;47.62)	
Obesity (BMI ≥ 30 kg/m ²)	323/832	38.7%	(35.32;42.05)	
Smoking habit				
Former smoker	212/835	25.4%	(22.38;28.40)	
Yes	136/835	16.3%	(13.72;18.52)	
No	213/835	58.3%	(22.49;28.53)	
Charlson comorbidity index				
Diabetes	100/815	12.3%	(9.71;14.24)	
COPD	55/816	6.7%	(4.84;9.33)	
Peripheral vascular disease	48/818	5.9%	(4.11;7.39)	
Peptic ulcer	46/818	5.6%	(3.69;6.85)	
Leukemia	44/812	5.4%	(3.69;6.85)	
Myocardial infarction	37/819	4.5%	(2.97;5.89)	
Liver disease	26/814	3.3%	(1.88;4.35)	
Connective tissue disease	21/818	2.6%	(1.39;3.68)	

Variables	n	Mean ± SD	Median	Minimum–maximum
Cerebrovascular disease	14/818	1.7%	(0.75;2.61)	
Moderate to severe chronic kidney disease	9/815	1.1%	(0.32;1.84)	
Congestive heart failure	7/819	0.9%	(0.16;1.52)	
Dementia	6/819	0.7%	(0.09;1.35)	
Metástatic	1/813	0.1%	(<0.01;0.66)	
AIDS	1/814	0.1%	(<0.01;0.66)	
Peripheral disease	0/819	—	—	
Hemiplegia	0/819	—	—	

Table 1. Distribution of patients according to demographic characteristics and comorbidity.

3.1. Quality of LIFE scales taking into account the foot and functionality of the foot

The scores of the different questionnaires used to measure the functionality, quality of life and dependence according to the presence or absence of flat foot in the entire sample studied and stratified by sex is shown in **Table 4**.

This table shows that patients with flat feet have significantly lower scores of the different quality of life domains of the FHSQ than those without flat feet. These values are consistent in both men and women being significantly inferior in the women and being in the men next to be significant.

It is also objected that FFI is greater in patients with flat feet than in patients who do not, and that difference is in the limit of statistical significance. This index reflects that the higher the score the worse functionality.

They are not significantly modified with the flatfoot or the dimensions of the physical and mental summary of the SF-36 questionnaire nor the Barthel index.

Although significant differences have been found between the values of the Lawton scale and whether or not having flat feet, in the bivariate analysis, dependence for instrumental activities (Lawton Scale) is not related to the presence of flat feet but to age and comorbidity (**Table 4**).

After identifying in the univariate analysis that the different FHSQ and FFI scores are modified with the presence of flat feet, the extent to which this effect is maintained after considering other variables such as age, gender and comorbidity is studied. For this, we perform different regression models presented in **Table 5**.

Anthropometric variables	n	Mean $\pm$ SD	Median	Minimum–maximum
Foot size (cm)	812	24.92 $\pm$ 1.66	24.75	20.50–29.80
Forefoot width (cm)	796	9.37 $\pm$ 0.62	9.40	7.55–11
<i>Left footprint</i>	n	%	95% IC	
Normal left footprint	413/803	51.4%	(47.91;54.95)	
Left flat footprint	174/803	21.7%	(18.76;24.59)	
Left cavus footprint	216/803	26.9%	(23.77;30.03)	
<i>Right footprint</i>				
Normal right footprint	385/793	48.50%	(45.01;52.09)	
Right flat footprint	184/793	23.20%	(20.20;26.20)	
Right cavus footprint	224/793	28.20%	(25.05;31.44)	
<i>Flat foot</i>	213/800	26.62%	(22.49;28.52)	
Unilateral	72/213	33.8%	(27.215;40.39)	
Bilateral	141/213	66.2%	(59.61;72.78)	
<i>Hallux abductus valgus</i>	325/805	40.4%	(36.92;43.82)	
Unilateral	38/325	11.7%	(8.04;15.34)	
Bilateral	287/325	88.3%	(84.66;91.95)	
<i>Hallux rigidus</i>	97/801	12.11%	(9.79;14.43)	
Unilateral	32/97	32.99%	(23.12;42.86)	
Bilateral	65/97	67.01%	(57.14;76.88)	
<i>Hallux extensus</i>	109/805	13.5%	(11.11;15.97)	
Unilateral	13/109	11.93%	(5.39;18.47)	
Bilateral	96/109	88.07%	(81.53;94.61)	
<i>One or more claw toes left</i>				
Yes	297/836	36.9%	(32.22;38.83)	
No	507/836	63.1%	(57.27;64.02)	
<i>One or more claw toes right</i>				
Yes	290/836	36.1%	(31.40;37.97)	
No	513/836	61.4%	(58.01;64.72)	

Table 2. Description of the sample according to type of footprint and presence of different foot pathologies.

	Flat foot		P	Crude OR	Adjusted OR** (95% CI)
	Yes	No			
	Mean (SD)*	Mean (SD)			
Age (years)	65.73 (11.04)	61.03 (11.45)	<0.001	1.037	1.029 (1.012–1.046)
Charlson comorbidity index adjusted for age	2.99 (2.11)	2.09 (1.75)	<0.001	1.275	
Charlson comorbidity index	0.92 (1.49)	0.50 (0.98)	<0.001	1.335	1.217 (1.042–1.421)
BMI (kg/m ²)	31.45 (5.55)	28.40 (4.17)	<0.001	1.147	1.137 (1.094–1.181)
Forefoot width (cm)	9.42 (0.64)	9.41 (2.01)	0.983	1.001	
Foot size (cm)	25.16 (1.66)	24.82 (1.65)	0.011	1.131	1.287 (1.102–1.504)
	n (%)	n (%)	p		
Age groups			<0.001		
40–64 years	86/425 (20.22%)	339/425 (79.8%)		1	
≥65 years	127/375 (33.9%)	248/375 (66.1%)		2.019	
BMI categories			<0.001		
Normal weight (18.5 kg/m ² ≤ BMI < 25 kg/m ²)	23/135 (17%)	112/135 (83%)		1	
Overweight (25 kg/m ² ≤ IMC < 30 kg/m ²)	57/351 (16.2%)	294/351 (83.8%)	0.832	0.944	
Obesity (IMC ≥ 30 kg/m ²)	133/312 (42.6%)	179/312 (57.4%)	<0.001	3.618	
Gender			0.419		
Male	99/353 (28%)	254/353 (72%)		1	1
Female	114/447 (25.5%)	333/447 (74.5%)		0.878	1.618 (0.963–2.717)

*SD: standard deviation.
** Adjusted OR: Adjusted Odds Ratio by age of the patient. Charlson’s comorbidity score. BMI, foot size and gender.
Statistical significative results are indicated in bold

Table 3. Differences between the presence or not of flatfoot and different variables.

After this regression, we objectified how the presence of flat feet continues to modify the score of the different dimensions of the FHSQ after adjusting or taking into account age, gender and comorbidity.

Figure 1. Area under the curve (AUC) to predict flatfoot according to different variables.

	Total sample (n = 835)			Female (n = 466)			Male (n = 369)		
	Flat foot			flat foot			flat foot		
	Yes	No		Yes	No		Yes	No	
	Mean (SD)	Mean (SD)	p	Mean (SD)	Mean (SD)	p	Mean (SD)	Media (SD)	p
<i>SF-36</i>									
Physical summary	53.72 (8.25)	54.55 (7.78)	0.189	53.06 (9.60)	55.48 (7.95)	0.017	54.48 (6.29)	53.34 (7.37)	0.148
Mental summary	47.25 (9.55)	48.53 (8.48)	0.086	48.14 (9.94)	48.49 (8.98)	0.744	46.22 (9.02)	48.60 (7.80)	0.015
Barthel index	97.38 (11.23)	99.43 (4.03)	0.052	96.80 (12.42)	99.41 (3.23)	0.112	97.95 (9.97)	99.46 (4.83)	0.183
Lawton index	6.14 (1.89)	6.52 (1.57)	0.040	7.54 (1.51)	7.87 (0.63)	0.104	4.74 (0.96)	4.91 (0.42)	0.188
<i>Foot Health Status Questionnaire</i>									
Foot pain domain	86.91 (29.63)	90.52 (17.62)	0.024	82.12 (22.56)	86.90 (19.97)	0.047	92.47 (10.19)	95.28 (12.49)	0.132
Function domain foot	90.30 (19.64)	94.36 (14.55)	0.006	86.51 (21.96)	92.13 (16.81)	0.014	94.71 (15.53)	97.30 (10.19)	0.129
Footwear domain	60.07 (37.38)	68.44 (35.60)	0.004	53.95 (37.79)	64.48 (35.77)	0.008	67.26 (35.75)	73.62 (34.77)	0.130

	Total sample (n = 835)			Female (n = 466)			Male (n = 369)		
	Flat foot			flat foot			flat foot		
	Yes	No	p	Yes	No	p	Yes	No	p
	Mean (SD)	Mean (SD)		Mean (SD)	Mean (SD)		Mean (SD)	Media (SD)	
General foot health domain	48.88 (21.66)	53.67 (20.89)	0.005	44.19 (22.99)	49.89 (21.02)	0.021	54.34 (18.66)	58.63 (19.67)	0.064
Foot function Index	7.63 (13.93)	5.22 (11.58)	0.055	9.76 (13.53)	12.73 (6.86)	0.082	4.91 (14.06)	2.84 (9.17)	0.178

SD, standard deviation.
Statistical significative results are indicated in bold

Table 4. Differences between the presence or not of flatfoot stratified by sex according to the questionnaires studied: SF-36, Barthel and Lawton index, Foot Health status questionnaire and Foot function index.

Variables	B	Standard error	Beta	t	p
<i>Linear regression model to predict dimension score foot pain FHSQ</i>					
Gender	-9.225	3.743	-0.249	-7.016	<0.001
Age	-0.007	0.060	-0.004	-2.134	0.913
Charlson Score	-1.284	0.602	-0.080	-2.134	0.003
Flat foot	-2.931	1.510	-0.070	-1.942	0.053
<i>Linear regression model to predict dimension score function foot FHSQ</i>					
Gender	-5.872	1.148	-0.183	-5.116	<0.001
Age	-0.054	0.053	-0.039	-1.029	0.304
Charlson Score	-1.009	0.525	-0.073	-1.922	0.055
Flat foot	-3.329	1.317	-0.092	-2.528	0.012
<i>Linear regression model to predict score footwear dimension FHSQ</i>					
Gender	-10.305	2.591	-0.142	-3.977	<0.001
Age	-0.519	0.119	-0.165	-4.351	<0.001
Charlson Score	1.286	1.185	0.041	1.086	0.278
Flat foot	-6.897	2.979	-0.084	-2.315	0.021
<i>Linear regression model to predict overall health score foot dimension FHSQ</i>					
Gender	-9.214	1.527	-0.215	-6.035	<0.001
Age	-0.094	0.070	-0.051	-1.336	0.182
Charlson Score	-1.248	0.699	-0.068	-1.786	0.074
Flat foot	-3.614	1.752	-0.075	-2.063	0.039
<i>Linear regression model to predict final score of the Foot Function Index</i>					
Sexo	4.400	1.031	0.177	4.269	<0.001
Edad	0.056	0.049	0.051	1.155	0.249
Charlson Score	1.242	0.489	0.112	2.540	0.011
Pie plano	1.821	1.187	0.066	1.534	0.126

Statistical significative results are indicated in bold

Table 5. Multiple linear regression to predict the different dimensions of foot health status questionnaire and the FFI adjusting for gender, age, comorbidity and presence of flatfoot.

As for the functionality measured by the FFI we objectify how the presence of flat foot is in turn close to being significant and has a positive regression coefficient which implies that the presence of flat foot increases the FFI score and therefore decreases the functionality.

4. Discussion

This study shows that the prevalence of flatfoot was 26.62%. This finding is practically identical to a study carried out in Japan in a sample of 242 women and 98 men, with a prevalence of 26.5%, and as this finding is related to obesity and affection of pain and function [9].

Similar findings are found in other publications regarding the prevalence of flatfoot. In other population studies (Springfield, Massachusetts) the prevalence of flatfoot was 19.0% (20.1% in women and 17.2% in men) [10]. Another study conducted in the Boston area found a prevalence of 20% in women and 17% in men [11]. There are even studies in diabetic population in a sample of 230 patients that even refer to a prevalence of 37% [11].

It is evident that the characteristics and age of the population under study are determinants of this prevalence, so we also found that among Saudi Arabian army recruits in a sample of 2100 recruits aged 18–21 found a prevalence of 5% and factors associated with their presence have been family history, use of shoes in childhood, obesity and urban residence, no differences in functionality or discomfort in the foot [12].

Some studies conducted in India indicate that the use of shoes at earlier ages increases along with obesity and ligament laxity the prevalence of flat feet [26].

Another study carried out in Nigeria in 560 children between 6 and 12 years shows that although in the univariate analysis we found association with the type of footwear and age. However, after considering both, only age remained as a variable associated with the presence of flat foot [13].

The urban residence as a risk factor for the prevalence of flatfoot has also been described in a study carried out in Congo children where it was objected after studying 1851 footprints of 906 girls and 945 children between 3 and 12 years old that the prevalence decreases with the age is higher in urban areas, in the male sex and the use of footwear has little influence on this prevalence [14].

This study shows how BMI, age, comorbidity, and foot size are associated with the prevalence of flatfoot. Some studies describe how podologic pathology increases with age [17] while other studies describe how flatfoot decreases with age, after adjusting for other covariates [18], while others indicate that neither age nor gender nor the BMI, are related to the flat foot [19].

Studies carried out in primary schools identified gender and being overweight as a risk factor for flatfoot [20, 21] while studies with adolescents [22] and preschoolers [23] identified associated flatfoot to an increase in BMI.

Foot length and the presence of flatfoot associated with flatfoot have also been referenced in the literature [24] although there are also authors who say that it is not associated with length [19].

In the adult population this pathology was also found to be associated with race and concomitant pathology of the foot [16, 27–29].

Some studies even describe radiological findings of different morphology in the foot according to different ethnic groups [30].

Others point out how the different morphology radiology (angle of talus with the first metatarsal) is related to the symptomatic presence or not of flat foot [31].

Although obesity has been repeatedly associated with obesity [32]. Not all show this association with it [33].

4.1. Related to health

Some articles indicate not only the association of the flat foot with different characteristics such as age, sex, BMI, concomitant pathology, but also as a health modifier [16].

Thus there are studies of 97,279 recruits of the armed forces, who give flat feet to localized pains in the knee [34].

As we have previously pointed out in the article that finds a flat foot prevalence identical to ours, they also objectify how this alteration is also associated with the presence of pain and fatigue in women [9].

Others performed in Australian recruits of area forces show how foot alterations are not related to pain, injury or functionality, although flatfoot is associated with a lower subjective feeling of physical health than those with normal foot [18].

In another study where the adult population ($n = 784$) was studied in Boston, there was no association between foot alteration, pain and functionality [17].

Other studies find an association between the presence of flat feet and accidents produced in the training of professionals of the armed forces [35]. Although this finding is not consistent in all publications [36].

We also found an association between flat feet with disabilities in workers with spondylarthrosis [37] and fractures of the lower limbs [38].

This study shows that the quality of life and functionality in patients with flatfoot is lower than in those who do not, and that this effect is maintained after adjusting for age, sex and comorbidity using the FHSQ and FFI questionnaires. The use of specific instruments to measure this affection is important because general health questionnaires such as the SF-36 in this study have shown no differences between those with or without flat feet. Similar results were found by other authors who did not objectify differences between patients with podiatric pathology and did not use SF-36 as a quality of life measurement instrument [39].

The SF-36 is sensitive to changes but is a generic questionnaire. The SF-36 was described as a relevant tool to detect changes in results after Hallux valgus surgery [40].

Other authors have described a progressive reduction of SF-36 components as the severity of Hallux valgus increases [41].

The use of specific questionnaires to study the quality of life and the functionality of the foot is widely documented in the literature [22, 42, 43].

The changes experienced in quality of life by the FHSQ questionnaires and the pathological pathology have also been described in the literature [42, 44, 45].

The validity of the Spanish version of the FHSQ and the FFI has been described in the literature [46, 47].

It is therefore reasonable to have objectified in this study that the use of specific questionnaires on the foot objective significant differences that other more generic questionnaires have not detected.

5. Conclusions

Age, Charlson's comorbidity index, BMI and foot size are associated with the presence of flat feet.

The questionnaires SF-36, Barthel and Lawton were not altered with the presence of flat feet, while the questionnaires FHSQ and FFI were sensitive to the presence of flat feet.

Conflict of interest

The authors declare no conflicts of interest.

Author details

Cristina Gonzalez-Martin¹, Salvador Pita-Fernandez^{2*} and Sonia Pertega-Diaz²

*Address all correspondence to: cristina.gmartin@udc.es

1 Clinical Epidemiology Research Group, Health Sciences Department, Escuela Universitaria de Enfermería y Podología, Universidade da Coruña (UDC), Ferrol, Spain

2 Clinical Epidemiology and Biostatistics Research Group, Instituto de Investigación Biomédica de A Coruña (INIBIC), Complejo Hospitalario Universitario de A Coruña (CHUAC), SERGAS, Universidade da Coruña, A Coruña, Spain

References

- [1] Richie DH. Biomechanics and clinical analysis of the adult acquired flatfoot. *Clinics in Podiatric Medicine and Surgery*. 2007;**24**(4):617-644 vii

- [2] Ozan F, Doğar F, Gençer K, Koyuncu Ş, Vatansever F, Duygulu F, et al. Symptomatic flexible flatfoot in adults: Subtalar arthroereisis. *Therapeutics and Clinical Risk Management*. 2015;**11**:1597-1602
- [3] Menz HB, Fotoohabadi MR, Wee E, Spink MJ. Visual categorisation of the arch index: A simplified measure of foot posture in older people. *Journal of Foot and Ankle Research*. 2012;**5**(1):10
- [4] Murley GS, Menz HB, Landorf KB. A protocol for classifying normal- and flat-arched foot posture for research studies using clinical and radiographic measurements. *Journal of Foot and Ankle Research*. 2009;**2**:22
- [5] Queen RM, Mall NA, Hardaker WM, Nunley JA. Describing the medial longitudinal arch using footprint indices and a clinical grading system. *Foot & Ankle International*. 2007;**28**(4):456-462
- [6] Lizis P, Posadzki P, Smith T. Relationship between explosive muscle strength and medial longitudinal arch of the foot. *Foot & Ankle International*. 2010;**31**(9):815-822
- [7] Chen KC, Tung LC, Yeh CJ, Yang JF, Kuo JF, Wang CH. Change in flatfoot of pre-school-aged children: A 1-year follow-up study. *European Journal of Pediatrics*. 2013; **172**(2):255-260
- [8] Staheli LT, Chew DE, Corbett M. The longitudinal arch. A survey of eight hundred and eighty-two feet in normal children and adults. *The Journal of Bone and Joint Surgery. American Volume*. 1987;**69**(3):426-428
- [9] Otsuka R, Yatsuya H, Miura Y, Murata C, Tamakoshi K, Oshiro K, et al. Association of flatfoot with pain, fatigue and obesity in Japanese over sixties. *Nihon Koshu Eisei Zasshi*. 2003;**50**(10):988-998
- [10] Munro BJ, Steele JR. Foot-care awareness. A survey of persons aged 65 years and older. *Journal of the American Podiatric Medical Association*. 1998;**88**(5):242-248
- [11] Lauterbach S, Kostev K, Becker R. Characteristics of diabetic patients visiting a podiatry practice in Germany. *Journal of Wound Care*. 2010;**19**(4):140. 2, 4 passim
- [12] Abdel-Fattah MM, Hassanin MM, Felembane FA, Nassaane MT. Flat foot among Saudi Arabian army recruits: Prevalence and risk factors. *Eastern Mediterranean Health Journal*. 2006;**12**(1-2):211-217
- [13] Abolarin TO, Aiyegbusi AI, Tella BA, Akinbo SR. Relationship between selected anthropometric variables and prevalence of flatfoot among urban and rural school children in south West Nigeria. *Nigerian Quarterly Journal of Hospital Medicine*. 2011;**21**(2):135-140
- [14] Echarri JJ, Forriol F. The development in footprint morphology in 1851 Congolese children from urban and rural areas, and the relationship between this and wearing shoes. *Journal of Pediatric Orthopaedics. Part B*. 2003;**12**(2):141-146
- [15] Chen JP, Chung MJ, Wang MJ. Flatfoot prevalence and foot dimensions of 5- to 13-year-old children in Taiwan. *Foot & Ankle International*. 2009;**30**(4):326-332

- [16] Shibuya N, Jupiter DC, Ciliberti LJ, VanBuren V, La Fontaine J. Characteristics of adult flatfoot in the United States. *The Journal of Foot and Ankle Surgery*. 2010;**49**(4):363-368
- [17] Badlissi F, Dunn JE, Link CL, Keysor JJ, McKinlay JB, Felson DT. Foot musculoskeletal disorders, pain, and foot-related functional limitation in older persons. *Journal of the American Geriatrics Society*. 2005;**53**(6):1029-1033
- [18] Esterman A, Pilotto L. Foot shape and its effect on functioning in Royal Australian Air Force recruits. Part 1: Prospective cohort study. *Military Medicine*. 2005;**170**(7):623-628
- [19] Pita-Fernandez S, González-Martín C, Seoane-Pillado T, Pertega-Díaz S, Perez-García S, López-Calviño B. Podiatric medical abnormalities in a random population sample 40 years or older in Spain. *Journal of the American Podiatric Medical Association*. 2014;**104**(6):574-582
- [20] Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: Development and validation. *Journal of Chronic Diseases*. 1987;**40**(5):373-383
- [21] Alonso J, Prieto L, Antó JM. The Spanish version of the SF-36 health survey (the SF-36 health questionnaire): An instrument for measuring clinical results. *Medicina Clínica (Barcelona)*. 1995;**104**(20):771-776
- [22] Riskowski JL, Hagedorn TJ, Hannan MT. Measures of foot function, foot health, and foot pain: American Academy of Orthopedic Surgeons Lower Limb Outcomes Assessment: Foot and Ankle Module (AAOS-FAM), Bristol Foot Score (BFS), Revised Foot Function Index (FFI-R), Foot Health Status Questionnaire (FHSQ), Manchester Foot Pain and Disability Index (MFPDI), Podiatric Health Questionnaire (PHQ), and Rowan Foot Pain Assessment (ROFPAQ). *Arthritis Care & Research (Hoboken)*. 2011;**63**(Suppl 11):S229-S239
- [23] Cid-Ruzafa J, Damián-Moreno J. Disability evaluation: Barthel's index. *Revista Española de Salud Pública*. 1997;**71**(2):127-137
- [24] Graf C. The Lawton instrumental activities of daily living scale. *The American Journal of Nursing*. 2008;**108**(4):52-62; quiz -3
- [25] Pita-Fernandez S, Gonzalez-Martin C, Seoane-Pillado T, Lopez-Calvino B, Pertega-Diaz S, Gil-Guillen V. Validity of footprint analysis to determine flatfoot using clinical diagnosis as the gold standard in a random sample aged 40 years and older. *Journal of Epidemiology*. 2015;**25**(2):148
- [26] Sachithanandam V, Joseph B. The influence of footwear on the prevalence of flat foot. A survey of 1846 skeletally mature persons. *Journal of Bone and Joint Surgery. British Volume (London)*. 1995;**77**(2):254-257
- [27] Golightly YM, Hannan MT, Dufour AB, Jordan JM. Racial differences in foot disorders and foot type. *Arthritis Care & Research (Hoboken)*. 2012;**64**(11):1756-1759
- [28] Dunn JE, Link CL, Felson DT, Crincoli MG, Keysor JJ, McKinlay JB. Prevalence of foot and ankle conditions in a multiethnic community sample of older adults. *American Journal of Epidemiology*. 2004;**159**(5):491-498

- [29] Hagedorn TJ, Dufour AB, Riskowski JL, Hillstrom HJ, Menz HB, Casey VA, et al. Foot disorders, foot posture, and foot function: The Framingham foot study. *PLoS One*. 2013;**8**(9):e74364
- [30] Castro-Aragon O, Vallurupalli S, Warner M, Panchbhavi V, Trevino S. Ethnic radiographic foot differences. *Foot & Ankle International*. 2009;**30**(1):57-61
- [31] Pehlivan O, Cilli F, Mahirogullari M, Karabudak O, Koksall O. Radiographic correlation of symptomatic and asymptomatic flexible flatfoot in young male adults. *International Orthopaedics*. 2009;**33**(2):447-450
- [32] Fuhrmann RA, Trommer T, Venbrocks RA. The acquired buckling-flatfoot. A foot deformity due to obesity? *Der Orthopäde*. 2005;**34**(7):682-689
- [33] Rivera-Saldívar G, Torres-González R, Franco-Valencia M, Ríos-Monroy R, Martínez-Ramírez F, Pérez-Hernández E, et al. Risk factors associated with the conformation of the medial longitudinal arch and the symptomatic flat foot in a metropolitan school population in Mexico. *Acta Ortopédica Mexicana*. 2012;**26**(2):85-90
- [34] Lakstein D, Fridman T, Ziv YB, Kosashvili Y. Prevalence of anterior knee pain and pes planus in Israel defense force recruits. *Military Medicine*. 2010;**175**(11):855-857
- [35] Wang X, Wang PS, Zhou W. Risk factors of military training-related injuries in recruits of Chinese People's Armed Police Forces. *Chinese Journal of Traumatology*. 2003;**6**(1):12-17
- [36] Cowan DN, Jones BH, Robinson JR. Foot morphologic characteristics and risk of exercise-related injury. *Archives of Family Medicine*. 1993;**2**(7):773-777
- [37] López-Rojas P, Aguilar-Salinas A, Salinas-Tovar S, Marín-Cotoñieto IA, del Carmen Martínez-García M, Garduño-Espinosa J. Disabling spondyloarthritis risk factors in valley of Mexico workers. *Archives of Medical Research*. 2002;**33**(5):495-498
- [38] Keegan TH, Kelsey JL, Sidney S, Quesenberry CP. Foot problems as risk factors of fractures. *American Journal of Epidemiology*. 2002;**155**(10):926-931
- [39] Groarke P, Galvin R, Kelly J, Stephens MM. Quality of life in individuals with chronic foot conditions: A cross sectional observational study. *Foot (Edinburgh, Scotland)*. 2012;**22**(2):66-69
- [40] Saro C, Jensen I, Lindgren U, Felländer-Tsai L. Quality-of-life outcome after hallux valgus surgery. *Quality of Life Research*. 2007;**16**(5):731-738
- [41] Menz HB, Roddy E, Thomas E, Croft PR. Impact of hallux valgus severity on general and foot-specific health-related quality of life. *Arthritis Care & Research (Hoboken)*. 2011;**63**(3):396-404
- [42] Menz HB, Auhl M, Ristevski S, Frescos N, Munteanu SE. Comparison of the responsiveness of the foot health status questionnaire and the Manchester foot pain and disability index in older people. *Health and Quality of Life Outcomes*. 2014;**12**:158

- [43] Budiman-Mak E, Conrad KJ, Roach KE. The foot function index: A measure of foot pain and disability. *Journal of Clinical Epidemiology*. 1991;**44**(6):561-570
- [44] Beeson P, Phillips C, Corr S, Ribbans WJ. Hallux rigidus: A cross-sectional study to evaluate clinical parameters. *Foot (Edinburgh, Scotland)*. 2009;**19**(2):80-92
- [45] López López D, Bouza Prego ML, Requeijo Constenla A, Saleta Canosa JL, Bautista Casasnovas A, Tajés FA. The impact of foot arch height on quality of life in 6-12 year olds. *Colombia Médica*. 2014;**45**(4):168-172
- [46] Cuesta-Vargas A, Bennett P, Jimenez-Cebrian AM, Labajos-Manzanares MT. The psychometric properties of the Spanish version of the Foot Health Status Questionnaire. *Quality of Life Research*. 2013;**22**(7):1739-1743
- [47] Paez-Moguer J, Budiman-Mak E, Cuesta-Vargas AI. Cross-cultural adaptation and validation of the Foot Function Index to Spanish. *Foot and Ankle Surgery*. 2014;**20**(1):34-39