We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Introduction: Biomedical Challenges and Socioeconomic Burden

Nikolai V. Gorbunov and Joseph B. Long

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.75743>

1. Introduction

Modern socioeconomic developments have generally resulted in a greatly improved quality of life for most, but these advances have been accompanied by the introduction of numerous health challenges arising from new diseases and casualties associated with environmental, industrial, and economical disasters, social and armed military clashes, occupational exposures, daily high-speed traffic accidents, and so on. Among the diseases of growing public and military health concern is traumatic brain injury (TBI), which has been recently recognized as a “silent epidemic” emerging globally at the transition of the twentieth and twenty-first centuries [1–8].

2. Traumatic brain injury (TBI): definition, assessment and classification

Traumatic brain injury (TBI) can be defined as alteration in brain functions due to head collision with a stationary or a moving object (e.g., a projectile) or upon coupling of an external mechanical force (e.g., g-force, blast shock wave (SW)) with the head [9–12]. The traumatic effects of these insults emerge as either open or closed head wounds yielding penetrating or closed TBI [11–14]. Clinical classification of TBI severity is widely achieved using the *Glasgow Coma Scale* (GCS)—a neurological scale designed to tally medical conditions of individuals in the disease sequelae. The severity of TBI can be classified as mild, moderate, severe, as well as vegetative state and brain death—estimated upon clinical presentation of a patient’s neurologic signs and symptoms varying from case to case. Translational and clinical observations indicate that many symptoms resolve completely upon recovery, while others, especially

those resulting from “secondary injury” due to neurological complications and reactive traumatic responses to “primary trauma,” can persist as chronic illnesses resulting in partial or permanent disability [2, 3, 13, 15–18]. Consequently, TBI should not be regarded as a single clinical entity with a defined outcome but rather represents “a spectrum of brain injuries,” where each TBI subtype can lead to a distinct clinical condition which requires case-specific medical treatment [3]. From these considerations efficient personalized therapy would need implementation of advanced diagnostic techniques (such as contrast-enhanced computed tomography, diffusion tensor MRI, TBI biomarkers) for assessment of injury and monitoring of recovery (discussed in Chapters 1.5, 2.1, 2.2 of this book).

3. TBI: Etiology, pathobiology and translational research

Statistically, the vast majority of TBI is associated with vehicle collisions, damaging assaults, falls, collision/contact sports, and combat operations [4–11, 13]. These events initiate the primary mechanisms of blunt, ballistic, and blast-associated neurotrauma that may or may not be accompanied by skull fracture. Penetrating TBI is readily apparent and generates damage localized along the projectile path through the brain that includes a site of fractured or perforated skull, ruptured meninges, and lesions of the brain tissue [13, 19]. The absence of such conspicuous hallmarks in closed TBI can result in an initial underestimation of injury severity, particularly when a TBI score is “mild” [12]. In situations producing blunt closed TBI, the damaging forces to the head induce an intracranial inertial force due to linear acceleration/deceleration or rotational momentum to the brain, so it collides inside with the skull resulting in focal “coup or contrecoup or rotational shearing injuries” [12, 20, 21]. Moreover, the same external forces to the skull can generate an array of tensor stress forces (e.g., normal and shear, tensile) through the brain tissue that cause cell compression/stretching, disorder axonal trafficking (i.e., axoplasmic transport of mitochondria, synaptic vesicles, proteins, etc., from neuron’s body through the axoplasm), and yet shear and fraction neuronal fibers and disrupt the microvasculature and meningeal structures, thus leading to different forms of intracranial hemorrhage [13, 18, 20, 22–28]. The disruption and dismantling of brain circuitry by these mechanical forces are the principal effectors of injury [20, 25]. Primary brain contusions can elicit concussions, diffuse axonal injury (DAI) and encephalopathy, dysregulation of intracranial pressure and the flow of cerebrospinal fluid (CSF), and the impairment of visceral organs and systems complicated by a variety of neurochemical and metabolic effects [12, 20, 21, 26, 27]. From an etiological perspective, the secondary injury factors can feature brain ischemia and hypoxia, hypercapnia, neuroinflammation, impairment of blood-brain barrier, cerebral edema, meningitis, seizures, and so on [12–17, 21, 25].

A devastating form of TBI is produced by shock waves generated by detonation of explosive devices [7, 11, 12]. At the center of the explosion, gaseous products instantaneously expand from a small volume at a very high-pressure state through the surrounding ambient pressure environment. The compressed gases expand outwards at a supersonic speed in a form of air shock wave (SW). When encountering the head, the SW can impart energy to skull bones, dura and arachnoid mater, CSF, and neuronal tissue through which energy is delivered and dissipated via different mechanisms, namely inertia, spalling, shearing, compression, and

cerebral air embolism. A combination of DAI, meningitis, brain edema, ischemia, and neurocognitive disorders accompanied by systemic organ system complications are the described features of blast TBI and reflect the multifactorial nature of SW effects [12, 21, 23, 28].

The pathogenesis and recoveries from these varieties of TBI are age- and phenotype-dependent, greatly adding to the complexities and challenges for the development of therapies [29–31]. A recent focus in translational research has been the roles of genetic and epigenetic polymorphism in TBI disease, giving new perspectives on TBI management and identification of potential targets for rehabilitation [30–32]. This translational research has been primarily driven by animal models that have been developed over the past decades to mimic the clinical sequelae of human TBI. As noted earlier, since human TBI is very heterogeneous, no single animal model suffices, and researchers have relied upon the use of distinct yet complementary models to capture the characteristic features of human TBI documented through clinical and postmortem examination. As extensively reviewed [23 34–36], although imperfect, these in vivo and in vitro models together have provided valuable insights into posttraumatic sequelae which can be targeted for therapeutic intervention. Nevertheless, the failure to date to successfully translate a neuroprotective drug through phase 2 and 3 clinical trials highlights the compelling need to improve models to achieve an ecological validity and a greater translational value.

4. Epidemiology and social impact of TBI-related diseases

The epidemiology of TBI is overwhelming worldwide. According to the U.S. Department of Health and Human Services, in the United States, the overall incidence of TBI (either as an isolated injury or in combination with other injuries (i.e., polytrauma)) in 2010 was estimated to be 823.7 per 100,000 population, and the cost for direct TBI medical care in U.S. was estimated at more than \$50 billion per year [4]. A lower TBI rate was reported in Europe (235 per 100,000) in 2006 [5]. It should be noted that the above numbers in U.S. did not account for those persons who received care at the U.S. military or Veterans Affairs hospitals [6]. According to the U.S. Department of Defense report of 2013, the cohort of servicemen diagnosed with a TBI from 2000 through 2011 represented 235,046 persons (or 4.2% of the 5,603,720 who served in the Army, Air Force, Navy, and Marine Corps) (<http://dvbic.dcoe.mil/dod-worldwide-numbers-tbi>) [7, 11, 33]. Overall, among TBI casualties, almost 100% of persons with severe head injury and over 50% of those with moderate head injury acquire permanent disability and will not return to their premorbid level of function, which creates a major socioeconomic burden [4, 5, 9, 33]. In addition, dramatic psychological changes can occur among the TBI survivors who experience “the invisible injuries” of brain trauma (e.g., posttraumatic stress disorder (PTSD)). These occult injuries are particularly challenging, since the changes occur in the absence of any outward manifestation of injury and alterations in patient appearance, making diagnosis, management, and prognosis extremely difficult.

In the global perspective, the recent continuous expansion of military conflicts in the Middle East, Afghanistan, and North Africa occurs with the implementation of enormous amounts of weaponized conventional explosives which when deployed and detonated inevitably affect

civilian populations in the conflict zones. The bTBI civilian casualties due to these proxy wars are poorly determined [8]. Considering massive migration of civilian populations driven by these disasters to Europe, the bTBI epidemiology in these groups requires a particular attention, since their social care is becoming a burden of host countries.

5. Conclusion

TBI disease remains a continually growing public health concern both domestically and globally and has taken on a heightened urgency with the recent recognition of a growing number of chronic traumatic encephalopathy victims among athletes and Warfighters exposed to repetitive sub-concussive insults [37–39]. Although a daunting problem, considerable progress has been made over the past decade with characterizations of TBI etiology, epidemiology, and advances in definitions of age- and genotype-specific pathobiology/pathophysiology, diagnostics, acute medical-surgical treatments (e.g., prevention of secondary injuries and maintenance of brain physiology), as well as on the development of new modalities for long-term targeted therapy, rehabilitative care, and TBI prevention (see Chapters 1.1–2.5 of this book). Despite significantly reduced TBI mortality rates at surgical emergency and acute treatments, improvements in long-term outcomes remain a great challenge, largely because as noted earlier, the disease does not represent one pathological entity but is rather a syndrome represented by a wide range of lesions that can require different patient-specific therapies in order to sustain neurological and physiological recovery. Further resolution of these problems requires a mutual effort of clinicians/surgeons, biomedical scientists, biotechnologists, pharmacologists, and biomedical engineers.

Disclaimer

The contents, opinions and assertions contained herein are private views of the authors and are not to be construed as official or reflecting the views of the Department of the Army or the Department of Defense. The authors report no conflict of interest.

Author details

Nikolai V. Gorbunov^{1*} and Joseph B. Long²

*Address all correspondence to: gorbunov.nikolaiv@gmail.com

1 Uniformed Services University of the Health Sciences, Bethesda, MD, USA

2 Blast-Induced Neurotrauma Branch, Walter Reed Army Institute of Research, Silver Spring, MD, USA

References

- [1] Goldstein M. Traumatic brain injury: A silent epidemic. *Annals of Neurology*. 1990; 27(3):327
- [2] Rusnak M. Traumatic brain injury: Giving voice to a silent epidemic. *Nature Reviews. Neurology*. 2013;9(4):186-187. DOI: 10.1038/nrneurol.2013.38
- [3] Malpass K. Read all about it! Why TBI is big news. *Nature Reviews. Neurology*. 2013;9(4): 179. DOI: 10.1038/nrneurol.2013.55
- [4] Frieden TR, Houry D, Baldwin G. Report to Congress on Traumatic Brain Injury in the United States: Epidemiology and Rehabilitation. Atlanta, GA: Centers for Disease Control and Prevention. National Center for Injury Prevention and Control; Division of Unintentional Injury Prevention; 2015. https://www.cdc.gov/traumaticbraininjury/pdf/tbi_report_to_congress_epi_and_rehab-a.pdf
- [5] Majdan M, Plancikova D, Brazinova A, Rusnak M, Nieboer D, Feigin V, Maas A. Epidemiology of traumatic brain injuries in Europe: A cross-sectional analysis. *Lancet Public Health*. 2016;1(2):e76-e83. DOI: 10.1016/S2468-2667(16)30017-2
- [6] Faul M, Xu L, Wald MM, Coronado VG. Traumatic Brain Injury in the United States: Emergency Department Visits, Hospitalizations, and Deaths 2002-2006. Atlanta, GA: Centers for Disease Control and Prevention, National Center for Injury Prevention and Control; 2010
- [7] CDC, NIH, DOD, and VA Leadership Panel. Report to Congress on Traumatic Brain Injury in the United States: Understanding the Public Health Problem among Current and Former Military Personnel. Centers for Disease Control and Prevention (CDC), the National Institutes of Health (NIH), the Department of Defense (DOD), and the Department of Veterans Affairs (VA); 2013. https://www.cdc.gov/traumaticbraininjury/pdf/report_to_congress_on_traumatic_brain_injury_2013-a.pdf
- [8] Hagopian A, Flaxman AD, Takaro TK, Esa Al Shatari SA, Rajaratnam J, Becker S, Levin-Rector A, Galway L, Hadi Al-Yasseri BJ, Weiss WM, Murray CJ, Burnham G. Mortality in Iraq associated with the 2003-2011 war and occupation: Findings from a national cluster sample survey by the university collaborative Iraq Mortality Study. *PLoS Medicine*. 2013;10(10):e1001533. DOI: 10.1371/journal.pmed.1001533
- [9] Ainsworth CR, Brown GS. Head Trauma: Background, Epidemiology, Etiology. 2015. <http://emedicine.medscape.com/article/433855-overview>
- [10] Menon DK, Schwab K, Wright DW, Maas AI. Demographics and Clinical Assessment Working Group of the International and Interagency Initiative toward Common Data Elements for Research on Traumatic Brain Injury and Psychological Health. Position statement: Definition of traumatic brain injury. *Archives of Physical Medicine and Rehabilitation*. 2010;91:1637-1640

- [11] Marshall SA, Bell R, Armonda RA, Ling GSF. Traumatic Brain Injury. Chapter 8. In: Lenhart MK, Savitsky E, Eastridge B. *Combat Casualty Care: Lessons Learned from OEF and OIF*. Fort Detrick, MD: Office of The Surgeon General Borden Institute; 2012. pp. 343-391. <http://www.cs.amedd.army.mil/borden/book/ccc/uclafrontmatter.pdf>; https://archive.org/stream/CombatCasualtyCare/CCCFull_djvu.txt
- [12] Ling G, Bandak F, Grant G, Armonda R, Ecklund J. Neurotrauma from Explosive Blast. In: Elsayed, Atkins, editors. *Explosion and Blast Related Injuries*. Amsterdam: Elsevier; 2008. pp. 91-104. ISBN: 978-0-12-369514-7
- [13] Black KL, Hanks RA, Wood DL, Zafonte RD, Cullen N, Cifu DX, Englander J, Francisco GE. Blunt versus penetrating violent traumatic brain injury: Frequency and factors associated with secondary conditions and complications. *The Journal of Head Trauma Rehabilitation*. 2002 Dec;**17**(6):489-496
- [14] Kazim SF, Shamim MS, Tahir MZ, Enam SA, Waheed S. Management of penetrating brain injury. *Journal of Emergencies, Trauma, and Shock*. 2011;**4**(3):395-402. DOI: 10.4103/0974-2700.83871. PMCID: PMC3162712
- [15] Hernandez-Ontiveros DG, Tajiri N, Acosta S, Giunta B, Tan J, Borlongan CV. Microglia activation as a biomarker for traumatic brain injury. *Frontiers in Neurology*. 2013;**4**:30. DOI: 10.3389/fneur.2013.00030. eCollection 2013
- [16] Webster KM, Sun M, Crack P, O'Brien TJ, Shultz SR, Semple BD. Inflammation in epileptogenesis after traumatic brain injury. *Journal of Neuroinflammation*. 2017;**14**(1):10. DOI: 10.1186/s12974-016-0786-1
- [17] Bae DH, Choi KS, Yi HJ, Chun HJ, Ko Y, Bak KH. Cerebral infarction after traumatic brain injury: Incidence and risk factors. *Korean Journal of Neurotrauma*. 2014;**10**(2): 35-40. DOI: 10.13004/kjnt.2014.10.2.35. Epub 2014 Oct 31
- [18] Hegde A, Prasad GL, Kini P. Neurogenic pulmonary oedema complicating traumatic posterior fossa extradural haematoma: Case report and review. *Brain Injury*. 2017;**31**(1): 127-130. DOI: 10.1080/02699052.2016.1219388. Epub 2016 Nov 23
- [19] Hegde MN. *A Coursebook on Aphasia and Other Neurogenic Language Disorders*. 3rd ed. Delmar Cengage Learning: Clifton Park, NY; 2006
- [20] AC MK, Daneshvar DH. The neuropathology of traumatic brain injury. *Handbook of Clinical Neurology*. 2015;**127**:45-66. DOI: 10.1016/B978-0-444-52892-6.00004-0. ISBN 9780444528926. PMC: 4694720. PMID: 25702209
- [21] Chandra N, Sundaramurthy A. Acute pathophysiology of blast injury—From biomechanics to experiments and computations: Implications on head and polytrauma. Chapter 18. In: Kobeissy FH, editor. *Brain Neurotrauma: Molecular, Neuropsychological, and Rehabilitation Aspects*. Boca Raton (FL): CRC Press/Taylor & Francis; 2015. ISBN-13: 978-1-4665-6598-2
- [22] Strich SJ. Shearing of nerve fibres as a cause of brain damage due to head injury. *Lancet* 1961;**ii**:443-8

- [23] Sosa MA, De Gasperi R, Paulino AJ, Pricop PE, Shaughness MC, Maudlin-Jeronimo E, Hall AA, Janssen WG, Yuk FJ, Dorr NP, Dickstein DL, McCarron RM, Chavko M, Hof PR, Ahlers ST, Elder GA. Blast overpressure induces shear-related injuries in the brain of rats exposed to a mild traumatic brain injury. *Acta Neuropathologica Communications*. 2013;**1**(1):51. DOI: 10.1186/2051-5960-1-51
- [24] Armstrong RC, Mierzwa AJ, Marion CM, Sullivan GM. White matter involvement after TBI: Clues to axon and myelin repair capacity. *Experimental Neurology*. 2016 Jan;**275**(Pt 3): 328-333. DOI: 10.1016/j.expneurol.2015.02.011
- [25] Lifshitz J. Experimental CNS trauma. In: Kobeissy FH, editor. *Brain Neurotrauma: Molecular, Neuropsychological, and Rehabilitation Aspects*. Boca Raton, FL: CRC Press/Taylor & Francis; 2015. pp. 4-11. ISBN-13: 978-1-4665-6598-2
- [26] Hulka F, Mullins RJ, Frank EH. Blunt brain injury activates the coagulation process. *Archives of Surgery*. 1996;**131**(9):923-927; discussion 927-8
- [27] Stovell MG, Yan JL, Sleigh A, Mada MO, Carpenter TA, Hutchinson PJA, Carpenter KLH. Assessing metabolism and injury in acute human traumatic brain injury with magnetic resonance spectroscopy: Current and future applications. *Frontiers in Neurology*. 2017;**8**: 426. DOI: 10.3389/fneur.2017.00426. eCollection 2017
- [28] Cernak I. Understanding blast-induced neurotrauma: How far have we come? *Concussion*. 2017;**2**(3):CNC42 www.futuremedicine.com
- [29] Li W, Risacher SL, McAllister TW, Saykin AJ. Traumatic brain injury and age at onset of cognitive impairment in older adults. *Journal of Neurology*. 2016;**263**(7):1280-1285. DOI: 10.1007/s00415-016-8093-4
- [30] Kurowski B, Martin LJ, Wade SL. Genetics and outcomes after traumatic brain injury (TBI): What do we know about pediatric TBI? *Journal of Pediatric Rehabilitation Medicine*. 2012;**5**(3):217-231
- [31] Weaver SM, Portelli JN, Chau A, Cristofori I, Moretti L, Grafman J. Genetic polymorphisms and traumatic brain injury: The contribution of individual differences to recovery. *Brain Imaging and Behavior*. 2014;**8**(3):420-434. DOI: 10.1007/s11682-012-9197-9
- [32] Rostami E, Krueger F, Zoubak S, Dal Monte O, Raymont V, Pardini M, Hodgkinson CA, Goldman D, Risling M, Grafman J. BDNF polymorphism predicts general intelligence after penetrating traumatic brain injury. *PLoS One*. 2011;**6**(11):e27389. DOI: 10.1371/journal.pone.0027389
- [33] Bagalman E. Traumatic Brain Injury Among Veterans. CRS Report for Congress; R40941. 2013. pp. 1-17. www.crs.gov
- [34] Xiong Y, Mahmood A, Chopp M. Animal models of traumatic brain injury. *Nature Reviews Neuroscience*. 2013;**14**(2):128-42. DOI: 10.1038/nrn3407. Review
- [35] Morganti-Kossmann MC, Yan E, Bye N. Animal models of traumatic brain injury: Is there an optimal model to reproduce human brain injury in the laboratory? *Injury*. 2010;**41**(Suppl 1):S10-S13. DOI: 10.1016/j.injury.2010.03.032. Epub 2010 Apr 22. Review

- [36] Morales DM, Marklund N, Lebold D, Thompson HJ, Pitkanen A, Maxwell WL, Longhi L, Laurer H, Maegele M, Neugebauer E, Graham DI, Stocchetti N, McIntosh TK. Experimental models of traumatic brain injury: Do we really need to build a better mousetrap? *Neuroscience*. 2005;**136**(4):971-989. Epub 2005 Oct 20. Review
- [37] Tagge CA, Fisher AM, Minaeva OV, Gaudreau-Balderrama A, Moncaster JA, Zhang XL, Wojnarowicz MW, Casey N, Lu H, Kokiko-Cochran ON, Saman S, Ericsson M, Onos KD, Veksler R, Senatorov VV Jr, Kondo A, Zhou XZ, Miry O, Vose LR, Gopaul KR, Upreti C, Nowinski CJ, Cantu RC, Alvarez VE, Hildebrandt AM, Franz ES, Konrad J, Hamilton JA, Hua N, Tripodis Y, Anderson AT, Howell GR, Kaufer D, Hall GF, Lu KP, Ransohoff RM, Cleveland RO, Kowall NW, Stein TD, Lamb BT, Huber BR, Moss WC, Friedman A, Stanton PK, McKee AC, Goldstein LE. Concussion, microvascular injury, and early tauopathy in young athletes after impact head injury and an impact concussion mouse model. *Brain*. 2018;**141**(2):422-458. DOI: 10.1093/brain/awx350
- [38] Mez J, Daneshvar DH, Kiernan PT, Abdolmohammadi B, Alvarez VE, Huber BR, Alosco ML, Solomon TM, Nowinski CJ, McHale L, Cormier KA, Kubilus CA, Martin BM, Murphy L, Baugh CM, Montenigro PH, Chaisson CE, Tripodis Y, Kowall NW, Weuve J, McClean MD, Cantu RC, Goldstein LE, Katz DI, Stern RA, Stein TD, McKee AC. Clinicopathological evaluation of chronic traumatic encephalopathy in players of American football. *Journal of the American Medical Association*. 2017;**318**(4):360-370. DOI: 10.1001/jama.2017.8334
- [39] Stern RA, Daneshvar DH, Baugh CM, Seichepine DR, Montenigro PH, Riley DO, Fritts NG, Stamm JM, Robbins CA, McHale L, Simkin I, Stein TD, Alvarez VE, Goldstein LE, Budson AE, Kowall NW, Nowinski CJ, Cantu RC, McKee AC. Clinical presentation of chronic traumatic encephalopathy. *Neurology*. 2013;**81**(13):1122-1129. DOI: 10.1212/WNL.0b013e3182a55f7f. Epub 2013 Aug 21

IntechOpen