We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

New Insights into Alleviating Diabetes Mellitus: Role of Gut Microbiota and a Nutrigenomic Approach

Lissé Angarita Dávila, Valmore Bermúdez Pirela,
Nadia Reyna Villasmil, Silvana Cisternas León,
Waldo Díaz-Vásquez, Ma Cristina Escobar,
Paula Carrasco, Samuel Durán, Kristian Buhring,
Rodrigo Buhring, Constanza Bugman,
Virginia Céspedes, Marcell Gatica,
Diana Rojas-Gómez, Marion Guerrero Wyss and
Francisco Valdebenito

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.76202>

Abstract

The scientific literature has shown that diet is able to modify the gut microbiota and contribute to obesity and diabetes development. This process—characterized by inflammation and gut barrier disruption—can affect the immune system and alter the adipogenesis and insulin resistance. This chapter describes the advances in nutrigenomics and Human Intestinal Microbiota (HIM) modification, and its relation with diabetes mellitus type two (DM2). In context where health and feeding are the main concerns of the human being, food innovation takes a special interest to people that look for a healthy diet or demand a functional aliments, such as nutraceutical. Some products derived from diet and interaction with HIM module the expression of many genes on the host, the so-called epigenome, with favorable effects. Novel functional fiber like low-glycemic oligosaccharides and sweeteners shows a potential prebiotic activity giving a new focus of nutritional guidelines for control and prevention of DM2. The use of prebiotics derived from functional fiber sources, such as fructo-oligosaccharides and beta-glucans as well as lignin and keffir, can contribute to the development of a healthy HIM by promoting the growth of specific bacteria, some of them associated with the prevention of obesity and diabetes.

Keywords: human intestinal, diet, fiber, prebiotic, carbohydrates, diabetes

1. Introduction

Diabetes mellitus type two (DM2) is a complex pathology, it depends of the interaction of genetic, epigenetic, environmental, and lifestyle factors [1]. This disease has generated an epidemiological worldwide impact, with a current report of 425 million adults having the disease according to the International Diabetes Federation (IDF). Currently, the last report of the year 2017, the epidemiological data showed an increment of 10 million cases diagnosed respect to 2015 [2]. According to the IDF, projection for 2045 is 650 million subjects with DM2. North America and the Caribbean have the highest prevalence of this disease (11%), where an increase of 62% is expected for the same period [3].

In the multidisciplinary treatment of this pathology, dietotherapy has been specifically considered as a critical control point in the international guidelines for DM2 [4, 5]. Recently, important advances in nutrition management have been developed associated to nutritional genomics, whose objective focuses on the interaction between the bioactive components of food and the human genome, this approach includes studies of nutrigenetics, nutrigenomics, and epigenetic modifications caused by nutrients [6].

Some investigations have used nutrigenomics to illustrate the modulation mechanism of specific fatty acids on gene expression, producing an impact on human metabolism [7, 8]. A common approach is the examination of individual levels of mRNA in relation to nutrient intake [9]. Tests with carbohydrates and dietary components such as fiber show a relationship between specific polymorphisms and the effect on insulin resistance [10]. In this sense, in a recent review, the effectiveness of the supply of fermentable carbohydrates on human metabolism is explained [11].

Furthermore, new advances in study of the composition of the human microbiota have shown an evident relationship between Human Intestinal Microbiota (HIM) and DM2 [12]. In this context, a significantly greater association of Firmicutes/Bacteroidetes in DM2 has been observed when is compared with normal weight and obese subjects [13].

New focus of nutritional treatment and its potential epigenetic effect constitute a panacea in the modification of the diabetic patient's microbiota [14]. The HIM is affected by the ingestion of bioactive compounds, showing prebiotic or probiotic effects, whose action can help to generate the growth of beneficial bacteria, such as *Bifidobacterium* and Bacteroidetes. The development of personalized nutritional methods considering the genomic information, use of prebiotics from novel sources of functional fibers (Fructo-oligosaccharides FOS, beta-glucans) [15], consumption of carbohydrates with low-glycemic index (GI), as well as the use of monosaccharide sweeteners with potential prebiotic activity, such as (tagatose) [16], would allow to generate a new therapeutic orientation for the control and prevention of this pathology. These dietary practices are important as part of the near future and will be analyzed in this chapter. Finally, a description in nutrigenomics advances and the effect of prebiotics consumption on modification of HIM are shown and its relationship with DM2 will be discussed.

2. Nutritional treatment and new perspectives

2.1. Prebiotics derived from functional fiber sources

A prebiotic is defined as “a non-digestible food ingredient that beneficially affects the host by selectively stimulating the growth and/or activity of one or a limited number of bacteria in the colon” [17]. Modification encouraged by prebiotics on the composition of HIM leads to the predominance of a few of the potentially health-promoting bacteria, especially, but not exclusively, *Lactobacilli* and *Bifidobacteria* [18]. Some prebiotics pass by the small intestine to the lower gut and become accessible for probiotic bacteria without being utilized by other intestinal bacteria [19]. Lactulose, galacto-oligosaccharides, fructo-oligosaccharides, inulin, and its hydrolysates, malto-oligosaccharides, and resistant starch are prebiotics normally used in the human diet [20, 21].

The definition used at present was given by the Food and Agriculture Organization of the United Nations World Health Organization, according to which probiotics are redefined as “live microorganisms which when are administered in adequate amounts confer a health benefit on the host.” In relation to foods, the definition can be adjusted to beneficial effect exerted by microorganisms “when are consumed in adequate amounts as part of food” [17, 21].

2.1.1. *Betaglucans*

2.1.1.1. *Cereal β -glucans*

Functional properties of β -glucans have been particularly attributed to the fact that these create viscous solutions in aqueous solution, as occurs in the digestive tract [22, 23]. This viscosity causes β -glucans to delay gastric emptying and interfere with the contact between pancreatic enzymes and their substrates in the intestinal lumen, slowing the digestion and absorption processes of nutrients [24]. This property could explain the effect of β -glucans on the reduction of plasma cholesterol concentrations and the glycemic index [25, 26].

2.1.1.2. *Yeasts and fungi β -glucans*

Another pivotal property of fungi/yeasts β -glucans is the modulation of the immune system [27, 28]. This effect could be due to the ability of β -glucans to stimulate receptors of the innate immune system present in the membrane of enterocytes, M cells and dendritic cells, improving the phagocytic activity of macrophages and antimicrobial activity of mononuclear cells and neutrophils [29–31]. This type of β -glucans would also prevent the promotion and progression of certain types of cancer, acting synergistically with monoclonal antibodies and chemotherapy [32, 33]. This stimulation of immunity would be achieved by increasing the secretion of pro-inflammatory cytokines and chemokines [34]. The main receptor involved in the effect of β -glucans immunity is Dectin-1, even though there is also a role for the receptor 3 of the complement, TLR-2, TLR-6 and the “scavengers” receptors [35, 36]. Dectin-1, known in human beings as β -glucans receptor (β GR), is a member of the pattern recognition receptors (PRR)

which fulfill an essential role in the innate immune response against viruses, bacteria, yeasts, and fungi, contributing to the recognition and elimination of pathogens [34]. This receptor is highly expressed in immune cells such as dendritic cells, neutrophils, eosinophils, and monocytes as well as in some populations of T and B cells and, to a lesser extent, in macrophages and enterocytes [31, 34, 35]. Dectin-1 acts through signal transduction activating Syk and RAF-1 [14]. It can also act synergistically with TLR which mediates the production of pro-inflammatory cytokines, such as IL-12 and TNF- α [33, 35].

2.1.2. *Fructo-oligosaccharides*

Inulin is a non-digestible carbohydrate present in many vegetables, fruits, and cereals [36]. Currently, at the industrial level it is extracted from the chicory root (*Cichorium intybus*) and is widely used as an ingredient in functional foods. Inulin and its derivatives (oligofructose, fructo-oligosaccharides) are generally called fructans, basically composed of linear chains of fructose [37]. The maximum dose allowed to be added to food formulated with inulin is up to 20 g/day for a simple dose and up to 10 g/day for multiple doses. At higher doses it can cause intolerances after consumption, such as osmotic effects (diarrhea), intestinal noises and flatulence as a result of the fermentation process [38]. Oligofructose is obtained by the partial enzymatic hydrolysis of inulin, composed of linear chains of glucosyl-fructosil. GP ranges between 2 and 8, with an average value of approximately [37]. It is present in foods such as cereals, onions, garlic, banana, and corn [38, 39]. There are promising evidences of its performance in the regulation of lipid parameters, reduction of the risk of cancer, reinforcement of the immune response and protection against intestinal disorders [40]. In a wide variety of food products, inulin and its derivatives are used as: thickener, emulsifier, gelling agent, sugar and fat substitute, moisturizer, depressor of the freezing point [37, 39].

2.1.3. *Lignin's: mucilage's (flaxseed)*

2.1.3.1. *Lignans*

Plant lignans are phenolic compounds with a skeleton of 2,3-dibenzylbutane [40]. Flaxseed is the richest food source in the precursors of lignans, secoisolariciresinol diglucoside (SDG), and materesinol, which are phytoestrogens that by action of gastric acid and bacterial glucosidase (facultative aerobics of Clostridia class) of the digestive tract transform into enterolactone and enterodiols, respectively, known as lignans of mammals [41]. These have more antioxidant capacity than their predecessors. Other lignans, such as lariciresinol, hinoquinina, arctigenin, divanillyl tetrahydrofuran nordihydroguaiaretic acid, isolariciresinol, and pinoresinol, are also present in flaxseed but the most abundant is SDG [40]. The health benefits of flaxseed lignans rely in their antioxidant capacity as retainers of hydroxyl radicals, and as estrogenic and anti-estrogenic compounds due to their structural similarity to the 17- β -estradiol [37, 41]. The antioxidant activity of flaxseed lignan (SDG) is related to the suppression of the oxidizing conditions of oxygen reactive species [41]. Secoisolariciresinol diglucoside and its aglycone secoisolariciresinol show a high antioxidant capacity and protective effects to the damage of the DNA and liposomes, especially in the epithelial cells of the colon exposed to these

compounds, during the metabolism of the colon bacteria that transform them into lignans of mammals [42, 43].

2.1.3.2. *Mucilage*

Mucilage is water-soluble polysaccharide present in many seeds, capable of absorbing 60–100 times their weight in water forming gels. They are formed by ramified arabinoxylans chains [44]. The mucilage is similar to the gums, composed of galactose, mannose, xylose, and other sugars [45]. One of the best known mucilage is psyllium (psyllium) or also called plantain, coming from the seeds of *Plantago* genus [42, 44]. The mucilage extracted from algae contains sugars somewhat different from terrestrial vegetables, such as agarobiose in the agar and sulf-sugar in the carrageenan, used in food technology [44]. Flaxseed mucilage is a complex polydisperse hydrocolloid and the different rheological behaviors observed in cultivars are caused by the differences in the ratio between neutral and acid polymers and by the molecular weight and structural conformation of polysaccharides [45, 46].

2.1.3.3. *Flaxseed*

Even though flaxseed is much known, it is not widely used in the formulation of food [47]. This seed has significant amounts of bioactive compounds, such as alpha-linolenic acid, lignans and dietary fiber, with potential effects in the prevention of some chronic diseases such as reducing the risk of cardiovascular diseases, mitigating the effects of diabetes, renal pathologies, obesity, colon and rectum cancer, reducing serum cholesterol level, and promoting the intestinal evacuation [46, 47]. These characteristics make flaxseed an attractive source of ingredients to be used in the elaboration of different functional foods [48].

2.1.4. *Kefirs*

Different *in vitro* and *in vivo* studies have demonstrated the ability of kefir to promote health through the presence of bioactive peptides. Multiple bioactivities of this beverage such as antihypertensive, antimicrobial, immune-modulating, mineral-carrying, antithrombotic, opioid, and antioxidant have been the most reported [49]. These characteristics of kefir, along to the pre- and probiotic properties, hypocholesterolemic, the bioavailability of milk components with biological activity and the presence of metabolites such as organic acids and bacteriocins, situate it as functional food [50]. This is a food that beyond the nutritional contribution of its components has been proven to benefit one or more physiological functions of the organism, improving the health, well-being, and/or reducing the risk to suffer diseases [51–53]. Additionally, there are bioactivities that have been poorly studied as the mineral fixing properties and the antithrombotic activity. There are some studies about the substrate-microorganism-metabolite-bioactivity inter-relationships based on metagenome studies [54, 55]. It would be important to carry out more investigations to determine the different bioactivities more deeply and the effective dose by trying to reach the intestinal level in sufficient quantity to implant and colonize its surface [56].

2.2. Low-glycemic index of carbohydrates and inflammatory state intestinal mucose

2.2.1. *Historic context glycemic index and glycemic load*

One of the major dietary changes of the modern world has been the high consumption of processed foods rich in carbohydrates and low in fiber; highly related to the increasing rates of obesity and diabetes [49]. In this sense, pharmacological approaches focused on large clinical trials have been useful for improving glycemic control in patients with type 2 diabetes (DM2) [51]. Similarly, a positive effect in the control of diabetes has been associated with the consumption of diets low in GI [52], this indicator determines the effect of the available carbohydrates in food on the average concentration of glucose in blood, this value is defined as the relation between the area under the curve of 50 g of available carbohydrates in a food, with the area under the curve of same amount of carbohydrates of a reference food [53].

White bread and glucose, which has been assigned a GI of 100, are considered reference foods rather than a high value for this indicator [54]. Different entities worldwide, such as the American Diabetes Association (ADA) [55], the European Association for the Study of Diabetes [56], the Canadian Diabetes Association [57], and the UK Diabetes Nutrition Subcommittee [58] have prioritized dietary treatment with a relevant approach to carbohydrate quality for the glycemic control, with special emphasis on reducing the digestion rate, absorption and metabolism of carbohydrates from foods [59].

Therefore, this indicator expresses the potential glycemia of a meal, representing the quality of foods with predominance of carbohydrates [60]. Foods with carbohydrates capable of digesting, absorbing and metabolizing quickly are considered food with high GI ($GI \geq 70$ in the glucose scale). Those between $GI = 55$ and 70 are considered in an intermediate value, while those digested, absorbed and metabolized slowly are classified as foods with low GI ($GI \leq 55$ in the glucose scale) [53, 60]. There are international tables with the published values of this indicator for a large number of products on the market. The first table was published in 1981 and was later updated in 1994 and 1995 [61, 62]. There is a marked controversy over the use of this indicator in the decade of the 80, due to an inadequate interpretation of the evidence for its determination [60, 63]. Criticism has focused on the methodological validity of the process to quantify it since a large number of factors had influenced the results [60, 64]. The position of the latest consensus of glycemic index experts in 1995 held in Europe has determined that most of the current critics are not valid, and that these reflect a failure of the knowledge translation [60]. In this context, it is important to consider that important entities such as the International Diabetes Federation have recognized the relevance of post-prandial regulation of glucose in order to achieve the objectives of HbA1C by developing specific guidelines, whose management is related to the GI concept [64].

2.2.2. *Glycemic load, glycemic index, and insulin response*

The value of glycemia and the insulin response depends on the quantity and quality of carbohydrate and the mix of food ingested, the so-called glycemic load (GL). The GL represents a relationship between the quantity and quality of carbohydrate [54, 60], and is defined as the total carbohydrate content available in an amount of food ($GL = GI \times \text{available carbohydrates} / \text{specific amount of food}$) [60], and is the result of multiplying the amount of carbohydrate

ingested in food by the value of its GI. The GL should be interpreted as a measure in the demand for insulin, this value is a good indicator of the levels of post-prandial glycemia, associated to the amount of calories in a particular portion of food or diet [65]. Thus, foods with high GL and high GI have a direct effect on the development of hyperinsulinemia [66], insulin resistance and risks to develop DM, which also have been linked to high-IG foods [67].

2.2.3. *Insulin response and inflammation mucus and glycocalyx layer*

Several studies have determined a clear link between the glycemic index and the glycemic load of food and the insulin response [52, 68, 69]. Studies suggest that carbohydrates can modify the microbiota, depending on their ability to increase glycemic and insulin response values according to glycemic and insulinemic index [70, 71]. In this sense, several studies in rodents have reported oligofructose as a recognized prebiotic, capable of modulating IM and improving insulin sensitivity [72, 73]. Similarly, inulin-type fructans have been tested to determine their ability to modulate lipid metabolism and carbohydrate in various animal models [73, 74]. It has been reported that oligofructose (OFS) decreases the intake of food, the development of fat mass and hepatic steatosis in normal and obese rodents. In addition, OFS exerts an antidiabetic effect in rats treated with streptozotocin and mice treated with high content of fat [72]. Chang et al. demonstrated that the addition of OFS also caused changes in the IM, specifically for *Bifidobacterium* and *Clostridium leptum* [75] content. These results suggest that OFS may be an effective therapeutic complement in the treatment of diabetes type 1 (DM1) by improving insulin sensitivity and beta cell function, leading to better glycemic control [76]. OFS reduced body weight, energy intake and fat mass in both phenotypes ($P < 0.05$) [76]. In another study carried out in two different groups of rodents, OFS did not modify ghrelin in plasma, but plasma levels of GIP were reduced and PYY were elevated ($P < 0.05$) [76] by OFS, reducing body weight and adiposity in prone obese phenotypes and in those insulin-resistant [76].

The changes induced by this saccharide in the profiles of IM of these animals, along with the changes of intestinal hormone levels probably contribute to lower body weights sustained [76, 77]. Milk prebiotic oligosaccharides have been reported to alter the IM and may influence the metabolism of the host. In a study performed in rats comparing diets with 15% of glucose, fructose, galactose, and methylcellulose content, daily intake of 15% galactose improved the sensitivity to hepatic insulin compared with glucose and fructose, producing an increase in the content of hepatic glycogen in the feeding state and a positive change in the IM populations; unlike the intake of galacto-oligosaccharides [78], which improved the IM profile without any effect on the insulin sensitivity. The GI of lactose, fructose and isomaltose is (=43), (=20), and (=2), respectively [62]. Further studies on these indicators are required in monosaccharides and their effect on the human microbiome.

3. New perspectives in biotechnology of foods, low-glycemic index and the microbiota

In context where health and feeding are the main concerns of the human being, food innovation takes a special interest to people that look for a healthy diet or demands a greater number

of functional products, such as nutraceutical, that often generates more contribution than nutrients, helping to improve the prevent of different diseases [79].

A functional food has been defined as a: (i) natural food, (ii) food which a component with some technology or biotechnology has been added or removed, (iii) food where the nature of one or more components has been varied, (iv) food which the bioavailability of one or more of its components has been modified, and/or (v) any combination of the above possibilities [80]. The world commercialization for functional foods and beverages have grown from \$ 33 billion in 2000 to 67.7 billion pesos in 2013, that mean the 5% of the global food commercialization, and the growth of investment in food industry as a whole. Latin America is currently a potential producer and consumer of functional foods, because of its large natural resources, a wide biodiversity of flora and fauna having a variety of plants and edible fruits with potential and beneficial effects for health [78].

Bioactive molecules works mainly modifying cellular signaling and causing changes in expression of certain genes, for instance producing a defensive response to harmful processes like differentiation and cell proliferation, inflammation, it is the base of the understanding for most prevalent diseases. New technology applied for food and nutrition sciences are closely related to the biomedical area, researchers require strong training in molecular biology, genetics and nutritional biochemistry, among others disciplines [81]. The current “omics” technologies, such as genomics, transcriptomics, proteomics, metagenomics, metatranscriptomics and metabolomics, have introduced important strides in the fields of health, biotechnology, ecology, and food [81]. The increase of the importance of (I + D) from academy, where food and pharmaceutical industry have worked together to promote healthy feeding, functional foods and nutraceuticals developing, those products when are consumed in a regular way, contribute to the prevention and/or treatments of certain diseases [82]. Genetic engineering plays an important role in the improvement of functional foods, which involves biological and technological research and also normative and ethical communication [83]. New probiotic strains isolated from natural niches and other produced by genetically engineered organisms (GMOs) have broadened the spectrum of organisms with improved probiotic properties for incorporation into functional foods [84]. More than 500 probiotic food products have been introduced into the world stores over the last couple of decades [81]. The contribution of biotechnology to production of prebiotics is remarkable. Prebiotic such as inulin and fructose polymer are produced by extraction of natural products (mainly chicory for fructose polymer), other prebiotics are produced by bioprocesses involving microorganisms or enzymes specifically conditioned for efficient synthesis of non-digestible oligosaccharides. On the other hand, inulin is the most used prebiotic, although it is probably not the most effective, actually, in the formulation of functional foods, also providing textural and rheological properties to the food matrix [83]. Another example of innovation is the design and development of product with intestinal microbiota and/or GI control effects, such as powdered additive, that incorporates also beneficial bacteria to the food. This development, achieved by researchers from National Institute Food Technology in Chile and Conicet Argentina, incorporated as an additive to certain foods—cold or lukewarm liquids—enriches the digestive system, balances the intestinal microbiota with a positive impact on the immune system [85].

3.1. Non-caloric sweeteners and gut microbiota

Non-caloric sweeteners (NCSs) are food additives widely used as sugar substitutes; these sweeteners enhance tastes and simultaneously reduce calories consumption. Some epidemiological studies have shown that artificial sweeteners are beneficial for weight loss, principally for subjects having glucose intolerance and type 2 diabetes [86]. Historically, the consumption of NCSs was restricted to people who have diseases such as diabetes; however, their consumption has increased in recent decades for general population. For their approval for human consumption, there are rigorous procedures required to consider them safe, however, today a controversy exist in its safety and it has been noted the possibility that the NCSs alter intestinal microbiota (IM). IM is involved in the metabolism of the host and plays a crucial role in food digestion and energy homeostasis. However, multiple environmental factors, such as diet, antibiotics and heavy metals, can disrupt the ecological balance of microbiota in the intestine [87]. A study in male Sprague-Dawley rats who were subjected to oral probe of 100, 300, 500, or 1000 mg/kg of Splenda for 12 weeks showed at the end of the treatment period, the number of total anaerobes, *Bifidobacteria*, *Lactobacilli*, *Bacteroides*, *Clostridia* and total aerobic bacteria decreased significantly. These changes occurred in Splenda doses containing sucralose at 1.1–11 mg/kg (FDA's acceptable daily intake for sucralose is 5 mg/kg) [88]. Other study realized in 8 weeks old C57B1 mice, two experiments were performed. Experiment 1, 4-week-old male mice were divided into three groups ($n = 8 \times \text{group}$) and treated for 8 weeks as follows: mice in control group received distilled water; mice in the low dose sucralose group (LS) a sucralose solution of 1.5 mg/kg body weight per day were given; and mice in the high-dose sucralose group (HS) received a sucralose solution of 15 mg/kg body weight per day, which is equal to the maximum IDA. In Experiment 2, 4-week-old male mice were divided into two groups and treated for 8 weeks as follows: Mice in control group received distilled water ($n = 8$); and acesulfame-K mice were given an acesulfame-K solution of 15 mg/kg body weight per day, which is equal to the ADI ($n = 9$), resulting that consumption of sucralose, but not of acesulfame-K, reduced the relative amount of *Clostridium cluster XIVa* in feces. Meanwhile, sucralose and acesulfame-K did not increase food intake [89]. Acesulfame k is genotoxic, and can inhibit the fermentation of glucose by intestinal bacteria [90]. A study in CD-1 mice (~8 weeks of age), were given a dose of 37.5 mg/kg body weight/day of acesulfame-K during 4 weeks, in males *Bacteroides* showed increased instead in females mice drastically decreased the relative abundance of multiple genres, including *Lactobacillus*, *Clostridium*, Ace-K disrupts the composition of the intestinal microbiome in a sex-dependent manner [90]. Another study in adult male C57B1/6 WT mice, gave two groups of mice a high in fat diet (60%) and commercial saccharin (equivalent to one human IDA) or glucose [91], resulting in an alteration in the glucose tolerance, the authors concluded that glucose intolerance was mediated by change in the microbiota (increase of *Bacteroidetes* and *Clostridium*). To corroborate the latter, a fecal transplantation to germ-free mice w performed, after 6 days an altered glucose tolerance was present in these mice. A similar study was carried out this time in seven humans (five men and two women), who were given 5 mg/kg/weight of saccharin (IDA equivalent) for 7 days, four of whom had altered glycemic responses. Other study, carried out in 31 humans that evaluated the consumption of aspartame and acesulfame k, showed that the consumers of these NCSs presented a different bacterial diversity to those who did not consume these

Figure 1. NCSs: sucralose, saccharin and acesulfame-K have been found to modify the balance of the HIM, either by decreasing or increasing the number of Bacteroidetes.

NCS [92]. This group also performed a smaller trial of seven healthy volunteers (five males, two females, and ages 28–36) who did not normally consume NCS and who received saccharin for 1 week at a dose of 5 mg/kg, IDA for these sweeteners. Most of these (4/7), known as “NCS responders” developed lower glucose tolerance and altered IM compared to “non-responders of NCS” [92]. Microbiome of “NCS responders” showed changes in composition by 16S rRNA analysis. Due this control group was not included in the design, it is unclear whether some healthy individuals exposed to seven consecutive tests of oral glucose tolerance (daily intake of 75 g of glucose) would have developed changes in glucose metabolism in the absence of saccharin. Palmnas et al. [107], demonstrated that 8 weeks of exposure to aspartame (at an equal dose to subjects consuming approximately 2–3 sodas/day) disrupted the intestinal microbiota; aspartame + high fat diet vs. water + high fat diet increased total bacteria; *Enterobacteraceae*, *Clostridium leptum*, and *Roseburia* spp. reduced *Bifidobacterium* sp. On the contrary, when the diet was low fat + aspartame or low-fat + water, *Clostridium leptum* increased, resulting in elevated levels of fasting glucose and insufficiency tolerance to insulin in rats [51]. However, the mechanism by which aspartame disrupted the IM is unclear, as aspartame is metabolized before it reaches the colon by intestinal esterases and peptidases in amino acids and methanol (Figure 1) [49].

3.2. Tagatose and prebiotic potential activity

D-tagatose (D-tag) is an isomer of fructose approximately 90% sweeter than sucrose. Only 20% of the oral intake of tagatose is completely metabolized, mainly in the liver [49]. The mayor part of this molecule is not digested or absorbed and passes through colon where water is absorbed and D-tag is fermented by colonic bacteria. This natural sweetener can be artificially

obtained from lactose. This sweetener with natural origin can be obtained artificially from lactose. Through food technology, glucose is separated and galactose is extracted, whose molecule is transformed into D-tagatose through an isomerization process [43].

D-tag would have an antihyperglycemic potential through its beneficial effects increasing postprandial serum glucose and hyperinsulinemia. Recent studies indicate that D-tag has a potent anti-diabetic effect and could be eventually associated with significant benefits for the treatment of obesity. The hypothesis regarding the mechanism of action proposed for this hypoglycemic effect would consider the interference with carbohydrates absorption by inhibition of intestinal disaccharidases and glucose transport, an also a mechanism of inhibition of hepatic glycogenolysis [37]. Another important characteristics of the D-tag is its low GI, considering white bread and glucose as reference foods, the D-tag GI is 3 and 4, respectively [63]. The potential applications of D-tag in the pharmaceutical industry and in food industry have reached a great boom [41]. However, the use of D-tag is limited by its high cost of production [36]. Another characteristic of D-tag is its potential prebiotic activity, and in order to preserve this effect the processing and storing of the food must ensure the maintenance of the chemical structure of the sweetener [35]. It has been determined that D-tag can be used for the formulation of diabetic beverages with minimal chance of degradation and very low loss of prebiotic activity [31, 33, 36], maintaining adequate thermal stability. Preliminary results suggest that D-tag would have an effect on the reduction of total cholesterol, VLDL, and LDL compared to sucrose in diabetic patients [53]; the contribution of D-tag to increase levels of HDL cholesterol has also been shown [54]. These clinical studies and wonderful advances in food technology make this molecule an ideal sweetener in functional products for patients with diabetes [62–64], with the ability to positively affect the intestinal microbiota of these patients, making its consumption more interesting and useful in a little explored area [85, 87]. On the other hand, the incorporation of novel functional sources of fiber, as well as oligosaccharides of potential prebiotic activity, has generated great scientific interest in the formulation of healthy foods aimed at diabetics. This new direction of science could be the anticipation of a new line of research that is beginning to emerge. Finally, future projection of personalized nutrigenomics foresees a great challenge toward the integration of different sciences as transcriptomics, epigenetics, proteomics, and metabolomics, with the purpose of positively modifying the microbiome, generating impact in the gene expression of the human organism, and avoiding manifestation of chronic diseases such as DM2.

4. Conclusions

The use of prebiotics obtained from functional fiber sources such as fructo-oligosaccharides and beta-glucans, as well as lignin and prebiotics such as keffir, can contribute to the development of a healthy HIM by promoting the growth of bacterial species that have been associated with obesity and diabetes prevention. On the other hand, it has been described that some low GI monosaccharides can positively modify the composition of the HIM in animal models, by regulating the mechanism of insulin sensitivity. More investigations are needed to evaluate the effect of saccharides, such as fructose, lactose and isomaltose in the human microbiome. Although, some NCS such as sucralose, saccharin, and acesulfame-K can modify the balance of HIM, mainly through the alteration in the number of *Bacteroidetes* species. Nevertheless, more studies in humans are required. In this sense, a new caloric sugar called D-tag has proposed

as possible hypoglycemic and probiotic effects. Finally, the new information presented in this chapter allows us to map out the near future where the integration of nutrigenomics and nutritional treatment focused on the microbiota modification will be plausible. Furthermore, the use of bioactive compounds that alter gene expression and/or affects immunity of pancreatic beta cells represent a projection toward the treatment and/or prevention of DM2.

Author details

Lissé Angarita Dávila^{1*}, Valmore Bermúdez Pirela^{2,3}, Nadia Reyna Villasmil³, Silvana Cisternas León^{1,9}, Waldo Díaz-Vásquez⁴, Ma Cristina Escobar¹, Paula Carrasco¹, Samuel Durán⁴, Kristian Buhning¹, Rodrigo Buhning⁶, Constanza Bugman¹, Virginia Céspedes⁷, Marcell Gatica¹, Diana Rojas-Gómez⁸, Marion Guerrero Wyss⁵ and Francisco Valdebenito¹

*Address all correspondence to: lisse.angarita@unab.cl

1 Carrera de Nutrición y Dietética, Facultad de Medicina, Universidad Andres Bello, Concepción, Chile

2 Universidad Simón Bolívar, Facultad de Ciencias de la Salud, Barranquilla, Colombia

3 Centro de Investigaciones Endocrino-Metabólicas Dr. Félix Gómez, Facultad de Medicina, La Universidad del Zulia, Maracaibo, Venezuela

4 Escuela de Nutrición y Dietética, Facultad de Ciencias para el Cuidado de la Salud, Universidad San Sebastián, Santiago, Chile

5 Escuela de Nutrición y Dietética, Facultad de Ciencias para el Cuidado de la Salud, Universidad San Sebastián, Valdivia, Chile

6 Departamento de Salud Pública, Universidad Católica de la Santísima Concepción, Sede Concepción, Chile

7 Servicio de Rehabilitación y Fisiatría, Hospital 12 de Octubre, Madrid, Spain

8 Escuela de Nutrición y Dietética, Facultad de Medicina, Universidad Andres Bello, Santiago, Chile

9 Area Salud. Universidad Tecnológica de Chile, Inacap, Chile

References

- [1] Schulze MB, Hu FB. Primary prevention of diabetes: What can be done and how much can be prevented? *Annual Review of Public Health*. 2005;**26**:445-467
- [2] International Diabetes Federation (IDF). The 7th Edition of the Diabetes Atlas. International Diabetes Foundation; 2015. Available from: idf.org/www.idf.org [Accessed: Sep 29, 2017]

- [3] International Diabetes Federation (IDF). The 8th Edition of the Diabetes Atlas. International Diabetes Foundation; 2017. Available from: idf.org/www.idf.org [Accessed: Dec 4, 2017]
- [4] International Diabetes Federation (IDF). Guideline for management of postmeal glucose in diabetes. *Diabetes Research and Clinical Practice*. 2014;**103**(2):256-268
- [5] American Diabetes Association. Standards of Medical Care in Diabetes. Abridged for primary care providers. *Clinical Diabetes* 2016;**39**(Suppl. 1):S1-S112
- [6] Ortega Á, Berná G, Rojas A, Martín F, Soria B. Gene-diet interactions in type 2 diabetes: The chicken and egg debate. *International Journal of Molecular Sciences*. Jun 2, 2017;**18**(6). pii: E1188. DOI: 10.3390/ijms18061188
- [7] Berná G, Oliveras-López MJ, Jurado-Ruiz E, Tejedo J, Bedoya F, Soria B, Martín F. Nutrigenetics and nutrigenomics insights into diabetes etiopathogenesis. *Nutrients*. Nov 21, 2014;**6**(11):5338-5369. DOI: 10.3390/nu6115338
- [8] Ferguson LR, De Caterina R, Görmán U, Allayee H, Kohlmeier M, Prasad C, Choi MS, Curi R, de Luis DA, Gil Á, et al. Guide and position of the international society of nutrigenetics/nutrigenomics on personalised nutrition: Part 1 – Fields of precision nutrition. *Journal of Nutrigenetics and Nutrigenomics*. 2016;**9**(1):12-27. Epub: May 12, 2016
- [9] Rostami H, Samadi M, Yuzbashian E, Zarkesh M, Asghari G, Hedayati M, Daneshafrooz A, Mirmiran P, Khalaj A. Habitual dietary intake of fatty acids are associated with leptin gene expression in subcutaneous and visceral adipose tissue of patients without diabetes. *Prostaglandins, Leukotrienes and Essential Fatty Acids*. Nov 2017;**126**:49-54. DOI: 10.1016/j.plefa.2017.09.010. Epub: Sep 14, 2017
- [10] Smith CE, Arnett DK, Corella D, Tsai MY, Lai CQ, Parnell LD, Lee YC, Ordovás JM. Perilipin polymorphism interacts with saturated fat and carbohydrates to modulate insulin resistance. *Nutrition, Metabolism & Cardiovascular Diseases*. May 2012;**22**(5):449-455 (Artículo en Inglés | MEDLINE | ID: mdl-21193293)
- [11] Korpela K. Diet, microbiota, and metabolic health: Trade-off between saccharolytic and proteolytic fermentation. *Annual Review of Food Science and Technology*. Jan 3, 2018. DOI: 10.1146/annurev-food-030117-012830 [Epub ahead of print]
- [12] Woo V, Alenghat T. Host-microbiota interactions: Epigenomic regulation. *Current Opinion in Immunology*. 2017;**16**(44):52-60. DOI: 10.1016/j.coi.2016.12.001
- [13] Remely M, Aumueller E, Jahn D, Hippe B, Brath H, Haslberger AG. Microbiota and epigenetic regulation of inflammatory mediators in type 2 diabetes and obesity. *Beneficial Microbes*. Mar 2014;**5**(1):33-43. DOI: 10.3920/BM2013.006
- [14] Bhat MI, Kapila R. Dietary metabolites derived from gut microbiota: Critical modulators of epigenetic changes in mammals. *Nutrition Reviews*. May 1, 2017;**75**(5):374-389. DOI: 10.1093/nutrit/nux001
- [15] Tamura K et al. Molecular mechanism by which prominent human gut bacteroidetes utilize mixed-linkage beta-glucans, major health-promoting cereal polysaccharides. *Cell Rep*. 2017;**21**(2):417-430. DOI: 10.1016/j.celrep.2017.09.049

- [16] Jayamuthunagai J, Srisowmeya G, Chakravarthy M, Gautam P. D-Tagatose production by permeabilized and immobilized *Lactobacillus plantarum* using whey permeate. *Bioresource Technology*. 2017;**235**:250-255. DOI: 10.1016/j.biortech.2017.03.123
- [17] FAO/WHO. Report on Joint FAO/WHO Expert Consultation on Evaluation of Health and Nutritional Properties of Probiotics in Food Including Powder Milk with Live Lactic Acid Bacteria. 2001. Available from: <http://www.fao.org/3/a-a0512e.pdf> [Accessed: Sep 7, 2017]
- [18] Madrigal L, Sangronis E. La inulina y derivados como ingredientes claves en alimentos funcionales. *Archivos Latinoamericanos de Nutrición*. 2007;**57**(4):387-396
- [19] Cani PD, Hoste S, Guiot Y, Delzenne NM. Dietary nondigestible carbohydrates promote L-cell differentiation in the proximal colon of rats. *The British Journal of Nutrition*. 2007;**98**:32-37. DOI: 10.1017/S0007114507691648
- [20] Maurer AD, Chen Q, McPherson C, Reimer RA. Changes in satiety hormones and expression of genes involved in glucose and lipid metabolism in rats weaned onto diets high in fibre or protein reflect susceptibility to increased fat mass in adulthood. *Journal of Physiology*, London. 2009;**587**:679-691. DOI: 10.1113/jphysiol.2008.161844
- [21] Tan J, McKenzie C, Potamitis M, Thorburn AN, Mackay CR, Macia L. The role of short-chain fatty acids in health and disease. *Advances in Immunology*. 2014;**121**:91-119. DOI: 10.1016/B978-0-12-800100-4.00003-9
- [22] Mandaşescu S, Mocanu V, Dăscăliţa AM, Haliga R, Nestian I, et al. Flaxseed supplementation in hyperlipidemic patients. *Revista Medico-Chirurgicală a Societăţii de Medici şi Naturalişti din Iaşi*. 2005;**109**:502-506
- [23] Daun JK, Barthet VJ, Chornick TL, Duguid S. Structure, Composition, and Variety Development of Flaxseed. New York, USA: AOCS Press; 2003. DOI: 10.1201/9781439831915.ch1
- [24] Im AR, Kim HS, Hyun JW, Chae S. Potential for tyndalized *Lactobacillus acidophilus* as an effective component in moisturizing skin and anti-wrinkle products. *Experimental and Therapeutic Medicine*. 2016;**12**(2):759-764. DOI: 10.3892/etm.2016.3406
- [25] Ivanova EP, Vysotskii MV, Svetashev VI, Nedashkovskaya OI, Gorshkova NM, Mikhailov VV, Yumoto N, Shigeri Y, Taguchi T, Yoshikawa S. Characterization of bacillus strains of marine origin. *International Microbiology*. 1999;**2**:267-271
- [26] Jarmuda S, O'Reilly N, Zaba R, Jakubowicz O, Szkaradkiewicz A KK. Potential role of Demodex mites and bacteria in the induction of rosacea. *Journal of Medical Microbiology*. 2012;**61**:1504-1510. DOI: 10.1099/jmm.0.048090-0
- [27] Kano M, Masauoka N, Kaga C, Sugimoto S, Iizuka R, Manabe K, Sone T, Oeda K, Nonaka C, Miazaki K, Ishikawa F. Consecutive intake of fermented milk containing *Bifidobacterium breve* strain yakult and galacto-oligosaccharides benefits skin condition in healthy adult women. *Bioscience of Microbiota, Food and Health*. 2013;**32**:33-39. DOI: 10.12938/bmfh.32.33

- [28] Mori N, Kano M, Masuoka N, Konno T, Suzuki Y, Miyazaki K, Ueki Y. Effect of probiotic and prebiotic fermented milk on skin and intestinal conditions in healthy young female students. *Bioscience of Microbiota, Food and Health*. 2016;**35**:105-112. DOI: 10.12938/bmfh.2015-022
- [29] Mukherjee S, Mitra R, Maitra A, Gupta S, Kumaran S, Chakraborty A, Majumder PP. Sebum and hydration levels in specific regions of human face significantly predict the nature and diversity of facial skin microbiome. *Scientific Reports*. 2016;**27**(6):36062. DOI: 10.1038/srep36062
- [30] Muraca M, Putignani L, Fierabracci A, Teti A, Perilongo G. Gut microbiota-derived outer membrane vesicles: Underrecognized major players in health and disease? *Discovery Medicine*. 2015;**19**:343-348
- [31] Kellow NJ, Coughlan MT, Reid CM. Metabolic benefits of dietary prebiotics in human subjects: A systematic review of randomised controlled trials. *The British Journal of Nutrition*. 2014;**111**:1147-1161. DOI: 10.1017/S0007114513003607
- [32] Parnell JA, Reimer RA. Weight loss during oligofructose supplementation is associated with decreased ghrelin and increased peptide YY in overweight and obese adults. *The American Journal of Clinical Nutrition*. 2009;**89**:1751-1759. DOI: 10.3945/ajcn.2009.27465
- [33] De Luis DA, de la Fuente B, Izaola O, Aller R, Gutiérrez S, Morillo M. Double blind randomized clinical trial controlled by placebo with a FOS enriched cookie on satiety and cardiovascular risk factors in obese patients. *Nutrición Hospitalaria*. 2013;**28**:78-85. DOI: 10.3305/nh.2013.28.1.6255
- [34] Abrams SA, Griffin IJ, Hawthorne KM, Ellis KJ. Effect of prebiotic supplementation and calcium intake on body mass index. *The Journal of Pediatrics*. 2007;**151**:293-298. DOI: 10.1016/j.jpeds.2007.03.043
- [35] Myles IA, Williams KW, Reckhow JD, Jammeh ML, Pincus NB, Sastalla I, Saleem D, Stone KD, Datta SK. Transplantation of human skin microbiota in models of atopic dermatitis. *JCI Insight*. Jul 7, 2016;**1**(10). pii: e86955. DOI: 10.1172/jci.insight.86955
- [36] Ni Raghallaigh S, Bender K, Lacey N, Brennan L, Powell FC. The fatty acid profile of the skin surface lipid layer in papulopustular rosacea. *The British Journal of Dermatology*. 2012;**166**:279-28758. DOI: 10.1111/j.1365-2133.2011.10662.x
- [37] Nyquist OL, McLeod A, Brede DA, Snipen L, Aakra A, Nes IF. Comparative genomics of *Lactobacillus sakei* with emphasis on strains from meat. *Molecular Genetics and Genomics*. 2011;**285**:297-311. DOI: 10.1007/s00438-011-0608-1
- [38] Uhde M, Ajamian M, Caio G, De Giorgio R, Indart A, Green PH, Verna EC, Volta U, Alaedini A. Intestinal cell damage and systemic immune activation in individuals reporting sensitivity to wheat in the absence of coeliac disease. *Gut*. 2016;**65**:1930-1937. DOI: 10.1136/gutjnl-2016-311964

- [39] Urban J, Fergus DJ, Savage AM, Ehlers M, Menninger HL, Dunn RR, Horvath JE. The effect of habitual and experimental antiperspirant and deodorant product use on the armpit microbiome. *Peer Journal*. 2016;2(4):e1605. DOI: 10.7717/peerj.16054:e1605
- [40] Van Rensburg JJ, Lin H, Gao X, Toh E, Fortney KR, Ellinger S, Zwickl B, Janowicz DM, Katz BP, Nelson DE, Dong Q, Spinola SM. The human skin microbiome associates with the outcome of and is influenced by bacterial infection. *MBio Journal*. 2015;6(5):e01315-15. DOI: 10.1128/mBio.01315-15
- [41] Assadi MM, Pourahmad R, Moazami N. Use of isolated kefir starter cultures in kefir production. *World Journal of Microbiology and Biotechnology*. 2000;16:541-543. DOI: 10.1023/A:1008939132685
- [42] Beshkova D, Simova ED, Simov ZI, Frengova GI, Spasov ZN. Pure cultures for making kefir. *Food Microbiology*. 2002;19:537-544. DOI: 10.1006/fmic.2002.0499
- [43] Bottazzi V, Bianchi F. A note on scanning electron microscopy of micro-organisms associated with the Kefir granule. *Journal of Applied Microbiology*. 1980;48:265-268
- [44] Hu C, Yuan YV, Kitts DD. Antioxidant activities of the flaxseed lignin secoisolariciresinol diglucoside, its aglycone secoisolariciresinol and the mammalian lignans enterodiols and enterolactone in vitro. *Food and Chemical Toxicology*. 2007;45:2219-2227. DOI: 10.1016/j.fct.2007.05.017
- [45] Li H et al. Effects of a flaxseed-derived lignan supplement on C-reactive protein, IL-6 and retinolbindingprotein 4 in type 2 diabetic patients. *The British Journal of Nutrition*. 2009;101:1145-1149. DOI: 10.1017/S0007114508061527
- [46] Tarpila S, Aro A, Salminen I, Tarpila A, Kleemola P, et al. The effect of flaxseed supplementation in processed foods on serum fatty acids and enterolactone. *European Journal of Clinical Nutrition*. 2002;56:157-165. DOI: 10.1038/sj.ejcn.1601298
- [47] Kuijsten A, Arts IC, van't Veer P, Hollman PC. The relative bioavailability of enterolignans in humans is enhanced by milling and crushing of flaxseed. *The Journal of Nutrition*. 2005;135:2812-2816
- [48] Paschos GK, Magkos F, Panagiotakos DB, Votteas V, Zampelas A. Dietary supplementation with flaxseed oil lowers blood pressure in dyslipidaemic patients. *European Journal of Clinical Nutrition*. 2007;61:1201-1206. DOI: 10.1038/sj.ejcn.1602631
- [49] Ueshima H, Stamler J, Elliott P, Chan Q, Brown IJ, et al. Food omega-3 fatty acid intake of individuals (total, linolenic acid, long-chain) and their blood pressure: INTERMAP study. *Hypertension*. 2007;50:313-319. DOI: 10.1161/HYPERTENSIONAHA.107.090720
- [50] Westcott ND, Muir AD. Flax seed lignan in disease prevention and health promotion. *Phytochemistry Reviews*. 2003;2:401-417. DOI: 10.1023/B:PHYT.0000046174.97809.b6
- [51] Carneiro RP. Desenvolvimento de uma cultura iniciadora para produção de kefir. Brazil: UFMG; 2010. p. 142. Available from: <http://www.bibliotecadigital.ufmg.br/dspace/handle/1843/MAFB-8EBKTH?show=full> [Accessed: Aug 26, 2017]

- [52] Chifiriuc MC, Cioaca AB, Lazar V. In vitro assay of the antimicrobial activity of kephir against bacterial and fungal strains. *Anaerobe*. 2011;**17**:433-435. DOI: 10.1016/j.anaerobe.2011.04.020
- [53] Diniz RO, Perazzo FF, Carvalho JCT, Schneenedorf JM. Atividade antiinflamatória de quefir, um probiótico da medicina popular. *Revista Brasileira de Farmacognosia*. 2003;**13**: 19-21. DOI: 10.1590/S0102-695X2003000300008
- [54] Dobson A, O'Sullivan O, Cotter PD, Ross P, Hill C. High-throughput sequence-based analysis of the bacterial composition of kefir and an associated kefir grain. *FEMS Microbiology Letters*. 2011;**320**:56-62. DOI: 10.1111/j.1574-6968.2011.02290.x
- [55] Ford ES, Mokdad AH. Epidemiology of obesity in the western hemisphere. *The Journal of Clinical Endocrinology and Metabolism*. 2008;**93**:S1-S8. DOI: 10.1210/jc.2008-1356
- [56] Farnworth ER, Mainville I. Kefir – A fermented milk product. In: Farnworth ER, editor. *Handbook of Fermented Functional Foods*. 2nd ed. Boca Raton, USA: CRC Press Taylor & Francis Group; 2003. pp. 89-127
- [57] Gradiser M, Bilic-Curcic I, Djindjic B, Berkovic MC. The effects of transition from bedtime to morning glargine administration in patients with poorly regulated type 1 diabetes mellitus: Croatian pilot study. *Diabetes Therapy*. 2015;**35**(10):675-684. DOI: 10.1007/s13300-015-0130-2
- [58] Tanaka K, Saisho Y, Manesso E, Tanaka M, Meguro S, Irie J, et al. Effects of liraglutide monotherapy on beta cell function and pancreatic enzymes compared with metformin in japanese overweight/obese patients with type 2 diabetes mellitus: A subpopulation analysis of the KIND-LM randomized trial. *Clinical Drug Investigation*. 2015;**35**(10):675-684. DOI: 10.1007/s40261-015-0331-5
- [59] Bhupathiraju SN, Tobias DK, Malik VS, Willett WC, Hu FB. Glycemic index, glycemic load, and risk of type 2 diabetes: Results from 3 large US cohorts and an updated meta-analysis. *American Journal of Clinical Nutrition*. 2014;**100**(1):218-232. DOI: 10.3945/ajcn.113.079533
- [60] Brouns F, Bjorck I, Frayn KN, Gibbs AL, Lang V, Slama G, Wolever TM. Glycaemic index methodology. *Nutrition Research Reviews*. 2005;**18**(1):145-171. DOI: 10.1079/NRR2005100
- [61] Wolever T, Jenkins D, Jenkins A, Josse R. The glycemic index: Methodology and clinical implications. *The American Journal of Clinical Nutrition*. 1991;**54**:846-854
- [62] American Diabetes Association. Standards of medical care in diabetes. *Diabetes Care*. 2017;**925**;38 (Suppl. 1):S1-S2. Available from: http://care.diabetesjournals.org/content/suppl/924/12/23/38.Supplement_1.DC1/January_Supplement_Combined_Final.6-99.pdf. [Accessed: Aug 30, 2017]
- [63] Diabetes and Nutrition Study Group of the European Association for the Study of Diabetes. Recommendations for the nutritional management of patients with diabetes mellitus. *European Journal of Clinical Nutrition*. 1998;**54**:353-355
- [64] Canadian Diabetes Association Clinical Practice Guidelines Expert. Nutrition therapy. *Canadian Journal of Diabetes*. 2013;**37**(Suppl 1):S45-S55. DOI: 10.1016/j.jcjd.2013.01.019

- [65] Angarita L, Miranda JL, Aparicio D, Parra K, Uzcátegui M, Céspedes V, Durán S, Reyna N. Glycemic index, glycemic load and insulin response of two formulas of isoglucose with different sweeteners and dietary fiber in healthy adults and type-2 diabetes. *Nutricion Hospitalaria*. 2017;**3**(34):532-539. DOI: 10.20960/nh.654
- [66] International Diabetes Federation (IDF). Guideline for management of postmeal glucose in diabetes. *Diabetes Research and Clinical Practice*. 2014;**103**(2):256-268
- [67] Augustin LS, Kendall CW, Jenkins DJ, Willett A, Astrup WC, Barclay AW, et al. Glycemic index, glycemic load and glycemic response: An International Scientific Consensus Summit from the International Carbohydrate Quality Consortium (ICQC). *Nutrition, Metabolism & Cardiovascular Diseases*. 2015;**25**(9):795-815. DOI: 10.1016/j.numecd.925.05.005
- [68] Pi-Sunyer FX. Glycemic index and disease. *American Journal of Clinical Nutrition*. 2002;**76**:290S-298S
- [69] Atkinson FS, Foster-Powell K, Brand Miller JC. International tables of glycemic index and glycemic load values. *Diabetes Care*. 2002;**31**(12):2281-2283. DOI: 10.2337/dc08-1239
- [70] Wolever TM, Augustin LS, Brand-Miller JC, Delport E, Livesey G, Ludwig DS, Sievenpiper JL. Glycemic index is as reliable as macronutrients on food labels. *American Journal of Clinical Nutrition*. 2017;**105**(3):768-769. DOI: 10.3945/ajcn.116.146092
- [71] Matthan NR, Ausman LM, Meng H, Tighiouart H, Lichtenstein AH. Estimating the reliability of glycemic index values and potential sources of methodological and biological variability. *American Journal of Clinical Nutrition*. 2016;**104**(4):1004-1013. DOI: 10.3945/ajcn.116.137208
- [72] Bao J, Atkinson F, Petocz P, Willett WC, Brand-Miller JC. Prediction of postprandial glycemia and insulinemia in lean, young, healthy adults: glycemic load compared with carbohydrate content alone. *American Journal of Clinical Nutrition*. 2011;**93**(5):984-996. DOI: 10.3945/ajcn.110.005033
- [73] González-Ortiz M, Martínez-Abundis E, Robles-Cervantes JA, Pérez-Rubio K. Comparison of two liquid nutritional supplements designed for patients with diabetes: Effect on glucose and insulin metabolism in healthy subjects. *Pharma Nutrition*. 2015;**3**(1):7-10. DOI: doi.org/10.1016/j.phanu.924.11.002
- [74] Gunerud UJ, Ostman EM, Bejorck BL. Effects of whey protein on glycaemia and insulinaemia to an oral glucose load in healthy adults; a dose response study. *European Journal of Clinical Nutrition*. 2013;**67**(7):749-753. DOI: 10.1038/ejcn.923.88
- [75] Livesey G, Taylor R, Livesey H, Liu S. Is there a dose-response relation of dietary glycemic load to risk of type 2 diabetes? Meta-analysis of prospective cohort studies. *American Journal of Clinical Nutrition*. 2013;**97**(3):584-596. DOI: 10.3945/ajcn.112.041467
- [76] Cheng G, Xue H, Luo J, Jia H, Zhang L, Dai J, Buyken AE. Relevance of the dietary glycemic index, glycemic load and genetic predisposition for the glucose homeostasis of

Chinese adults without diabetes. *Scientific Reports*. 2017;**7**(1):400. DOI: 10.1038/s4368-017-00453-9

- [77] Monteagudo-Mera A, Arthur JC, Jobin C, Keku T, Bruno-Barcena JM, Azcarate-Peril MA. High purity galacto-oligosaccharides enhance specific *Bifidobacterium* species and their metabolic activity in the mouse gut microbiome. *Beneficial Microbes*. 2016;**7**(2):247-264. DOI: 10.3920/BM925.0114
- [78] Latulippe ME et al. ILSI Brazil International Workshop on Functional Foods: A narrative review of the scientific evidence in the area of carbohydrates, microbiome, and health. *Food & Nutrition Research*. 2013;**57**. DOI: 10.3402/fnr.v57i0.8314
- [79] de Cossío LF, Fourrier C, Sauvant J, Everard A, Capuron L, Cani PD, Layé S, Castanon N. Impact of prebiotics on metabolic and behavioral alterations in a mouse model of metabolic syndrome. *Brain, Behavior, and Immunity*. 2017;**64**:33-49. DOI: 10.1016/j.bbi.926.12.022
- [80] Delzenne NM, Cani PD, Neyrinck AM. Modulation of glucagon-like peptide 1 and energy metabolism by inulin and oligofructose: Experimental data. *Journal of Nutrition*. 2007;**137** (11 Suppl):2547S-2551S
- [81] Dewulf EM et al. Inulin-type fructans with prebiotic properties counteract GPR43 overexpression and PPAR γ -related adipogenesis in the white adipose tissue of high-fat diet-fed mice. *Journal of Nutritional Biochemistry*. 2011;**22**(8):712-722. DOI: 10.1016/j.jnutbio.920.05.009
- [82] Chan C, Hyslop CM, Shrivastava V, Ochoa A, Reimer RA, Huang C. Oligofructose as an adjunct in treatment of diabetes in NOD mice. *Scientific Reports*. 2016;**6**:37627. DOI: 10.1038/srep37627
- [83] Cluny NL, Eller LK, Keenan CM, Reimer RA, Sharkey KA. Interactive effects of oligofructose and obesity predisposition on gut hormones and microbiota in diet-induced obese rats. *Obesity*. 2015;**23**(4):769-778. DOI: 10.1002/oby.21017
- [84] Stahel P, Kim JJ, Xiao C, Cant JP. Of the milk sugars, galactose, but not prebiotic galacto-oligosaccharide, improves insulin sensitivity in male Sprague-Dawley rats. *PLoS One*. 2017;**12**(2):e063260. DOI: 10.1371/journal.pone.063260
- [85] Fuentes Berrio L, Acevedo Correa D, Chantré CA, Gelvez Ordoñez VM. Alimentos Funcionales: Impacto Y retos Para El Desarrollo Y Bienestar De La Sociedad Colombiana. *Biotechnología en el Sector Agropecuario y Agroindustrial*. 2015;**13**(2):140. DOI: 10.7784/BSAA(13)140-149
- [86] Prieto-Hontoria P. Innovación y tendencias alimentarias Innovation and food trends. *Contribuciones Científicas y Tecnológicas*. 2016;**41**:15-20
- [87] Cornelio FS, Maldonado ZG, Ugas AR, Borjas A. Los alimentos funcionales: ¿son buenos para nuestra salud? functional foods: Are good for our health?. In *Crescendo. Ciencias de la Salud*. 2016;**3**(32):58-61

- [88] Lutz RM. Desafíos en investigación, desarrollo e innovación en alimentos y nutrición. *Revista Chilena de Nutrición*. 2013;**40**(4):404-407. DOI: 10.4067/S0717-7573013000400012
- [89] Valenzuela BA, Valenzuela R, Sanhueza J, Morales IG. Alimentos funcionales, nutraceúticos y foshu: “vamos hacia un nuevo concepto de alimentación”. *Revista Chilena de Nutrición*. 2014;**41**(2):89-204. DOI: 10.4067/S0717-757301400091011
- [90] Illanes A. Alimentos funcionales y biotecnología. *Revista Colombiana de Biotecnología*. 2015;**17**(1):5-8. DOI: 10.3146/rev.colomb.biote.v17n1.50997
- [91] Revista de Investigaciones Agropecuarias (RIA). RIA. Revista de Investigaciones Agropecuarias Vol. 42. Instituto Nacional de Tecnología Agropecuaria (INTA). Available from: http://www.scielo.org.ar/scielo.php?script=sci_arttext&pid=S589231492600091004&lng=es&nrm=iso&tlng=es
- [92] Gardner C, Wylie-Rosett J, Gidding SS, Steffen LM, Johnson RK, Reader D, et al. Nonnutritive sweeteners: Current use and health perspectives a scientific statement from the American heart association and the American diabetes association. *Diabetes Care*. 2012; **35**(8):708-808. DOI: 10.2337/dc12-9002
- [93] Palmnas MS, Cowan TE, Bomhof MR, Su J, Reimer RA, Vogel HJ, Hittel DS, Shearer J. Low-dose aspartame consumption differentially affects gut microbiota-host metabolic interactions in the diet-induced obese rat. *PloS One*. 2014;**9**(10):e109841. DOI: doi.org/10.1371/journal.pone.0109841