We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Role of Flavonoids as Wound Healing Agent

Muhammad Shahzad Aslam,
Muhammad Syarhabil Ahmad, Humayun Riaz,
Syed Atif Raza, Shahzad Hussain,
Omer Salman Qureshi, Povydysh Maria,
Zainab Hamzah and Osama Javed

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.79179>

Abstract

Flavonoids are found as the most abundant bioactive compounds all around the world. It is found in a number of medicinal plants that are used as wound healing agents in traditional medicinal uses such as *Buddleja globosa*, *Moringa oleifera*, *Lam*, *Butea monosperma*, *Parapiptadenia rigida* and *Ononis spinosa*. Flavonoids nowadays are being used in different formulation and wound healing dressings. Inflammation, proliferation and reepithelialization are involved in wound healing. Most of the wound healing medicinal plants possess multiple flavonoids that act as synergistic effect or combined effect. This chapter briefly reviews the role of flavonoids as wound healing agent in traditional and modern medicine.

Keywords: medicinal plants, flavonoids, wound healing, mechanism of flavonoids in wound healing

1. Introduction

The care of acute and chronic wound is the biggest challenges worldwide [1]. There are different kind of wounds, among them burns remain a major public health problem in developing countries [2]. There are different ways to heal the wound such as nano dressing [3], negative pressure [4], medicinal plants [5], synthetic polymers [6], gene therapy [7], stem cell [8], growth factors [9] and Functionalized Silk Biomaterials [10]. Flavonoids is one of an important bioactive source from medicinal plants that possess different pharmacological activities such as antioxidant, free radical scavenging capacity, coronary heart disease prevention,

hepatoprotective, anti-inflammatory, anticancer activities, growth regulators [11]. Wound results in edema, redness followed by pain. Inflammation is the major characteristics as a result of wound due to the release of eicosanoids, prostaglandins, leukotrienes and reactive oxygen species. By controlling all these factors will result in healing the wound faster. Recent students on Buddleja species and three Ghanaian species *Spathodea campanulata*, *Commelina diffusa* and *Secamone afzelii* showed antioxidant and anti-inflammatory properties. These anti-oxidant and anti-inflammatory properties were due to the presence of flavonoids and other bioactive compounds that works in combination to heal the wound [12]. The chapter deals with the study of role of flavonoids, mechanism of action that involves in healing the wound.

2. Wound healing medicinal plants

Healing the wound with medicinal plants is an alternative method of treatment used by traditional wound healer. The most commonly used plants all over the world is *Curcuma longa*. In Malaysia, there are few more species that act as wound healing such as *Elephantopus scaber*, *Centella asiatica*, *Clinacanthus nutans* and *Aloe barbadensis*. List of famous medicinal plants that plays an important role in wound healing is mentioned in **Table 1**.

Medicinal plants	Pharmacological uses	Reference
<i>Aloe barbadensis miller</i>	Anti-oxidant, anti-inflammatory, wound healing, antimicrobial	[13–16]
<i>Azadirachta indica</i>	Anti-malarial, wound healing, antiseptic, febrifuge, antihelminthic, anti-microbial	[17, 18]
<i>Curcuma longa</i> Linn.	Hepatoprotective, nephroprotective, anticoagulant, wound healing, anti-HIV to combat AIDS, anti-microbial	[19, 20]
<i>Clinacanthus nutans</i>	Anti-papillomavirus infectivity, anti-viral activity on varicella-zoster virus, anti-inflammatory activity, anti-herpes simplex virus type 1 and type 2 activity, anti-oxidant and protective effect against oxidative induced hemolysis	[21]
<i>Chromolaena odorata</i>	Wound healing, analgesic, anti-inflammatory and antipyretic activities	[22, 23]
<i>Centella asiatica</i>	Antimicrobial activity, anticancer activity, wound healing activity, neuroprotective activity, immunomodulatory activity, anti-inflammatory activity, hepatoprotective activity, insecticidal activity, and antioxidant activity	[24, 25]
<i>Elephantopus scaber</i>	Antimicrobial, hepatoprotective, antioxidant, antidiabetic, anti-inflammatory, analgesic, antiasthmatic, antiplatelet, and wound healing	[26]
<i>Euphorbia neriifolia</i>	Analgesic, hepatoprotective, immunostimulant, anti-inflammatory, mild CNS depressant, wound healing, radioprotective	[27, 28]
<i>Lantana camara</i>	Anticancer activity, wound healing, anti-inflammatory activity, antidiabetic activity, anthelmintic, antibacterial activity, antifungal activity, hepatoprotective activity, antioxidant activity, larvicidal activity	[29, 30]
<i>Tridax procumbens</i>	Hepatoprotective activity, anti-inflammatory, wound healing, antidiabetic activity, hypotensive effect, immunomodulating property, bronchial catarrh, dysentery, diarrhea and to prevent falling of hair promotes the growth of hair, and antimicrobial activity	[31]

Table 1. List of medicinal plants in healing the wound.

3. What is the role of flavonoids on human health?

Flavonoids are phenolic substances isolated from a wide range of vascular plants, with over 8000 individual compounds known [32]. They possess wide variety of activities such as anti-microbial [33], anti-oxidant [34], anti-cancer [35], anti-inflammatory [36] and wound healing [37]. Among all flavonoids there are some flavonoids such as flavan-3-ols and flavonols, possess a wide spectrum due to suppress a number of microbial virulence factors and show synergism with antibiotics [38]. While there has been a major focus on the antioxidant properties, there is an emerging view that flavonoids, and their in vivo metabolites, do not act as conventional hydrogen-donating antioxidants but may exert modulatory actions in cells through actions at protein kinase and lipid kinase signaling pathways. Flavonoids, and more recently their metabolites, have been reported to act at phosphoinositide 3-kinase (PI 3-kinase), Akt/protein kinase B (Akt/PKB), tyrosine kinases, protein kinase C (PKC), and mitogen activated protein kinase (MAP kinase) signaling cascades. Inhibitory or stimulatory actions at these pathways are likely to affect cellular function profoundly by altering the phosphorylation state of target molecules and by modulating gene expression [39]. In many molecular mechanisms of action for prevention against cancer, flavonoids play a major role by interacting between different types of genes and enzymes. Many mechanisms of action have been identified, including carcinogen inactivation, antiproliferation, cell cycle arrest, induction of apoptosis, inhibition of angiogenesis, antioxidation, and reversal of multidrug resistance or a combination of these mechanisms [40].

4. Flavonoids and wound healing

Catechins are one of the most widely tested classes of flavonoids for their wound healing modulation [41]. Lastly, considering that the delay in wound healing is due to insufficient or excessive fibroblast activity, some authors suggest that inhibition of fibroblast growth by flavonoids such as apigenin is beneficial for the treatment of skin injuries. Luteolin is another famous flavonoid present in medicinal plants, vegetables and fruits. It was also found as wound healing agent in different wound models [42]. Rutin which is found in many medicinal

Medicinal plants	Flavonoids isolated	Wound healing model	Reference
<i>Buddleja globosa</i>	Linarin (acacetin-7-O-rutinoside), luteolin and 6-hydroxyluteolin	Fibroblast growth stimulation	[44]
<i>Moringa oleifera</i> Lam	Vicenin-2, kaempferol and quercetin	Cell viability, proliferation, and wound scratch test assays.	[45]
<i>Butea monosperma</i>	Genistein and prunetine	Excision wound model, Incision wound model, Dead space wound model	[46]
<i>Parapiptadenia rigida</i>	Catechin	Fibroblast proliferation assay	[41]
<i>Ononis spinosa</i> L	Ononin, sativanone-7-O-glucoside	Linear incision and circular excision wound models and hydroxyproline estimation assay	[47]

Table 2. List of medicinal plants possessing flavonoids in healing the wound.

plants also possess wound healing activities [43]. List of medicinal plants possessing flavonoids in healing the wound is mentioned in **Table 2**.

5. Flavonoids in fruits and vegetables

Flavonoids found in number of fruits and vegetable. A study indicates a number of vegetable (28) and fruits (9) that possess anti-carcinogenic flavonoids [48]. Fruits such as apple, berries, and onions possess cardioprotective properties and bring a positive impact on blood pressure, vascular function and serum lipid levels [49].

6. Conclusion

Flavonoids is an essential bioactive compounds found in our daily life in the form of fruits and vegetable. It possesses wound healing properties. There are number of research that showed the importance of flavonoids as wound healing agent. For example the flowers of *Ipomoea carnea* belong to family Convolvulaceae. The isolated bioactive compounds such as Kaempferol, Kaempferol-3-O- β -D-glucoside confirm the wound healing activities on all animal models [50]. There is yet to explore its structure activity relationship on different wound models. Furthermore, there is little knowledge about mechanism involve in wound healing. Optimization use of flavonoids and their dosage as wound healing is another aspect that yet to explore.

Author details

Muhammad Shahzad Aslam^{1*}, Muhammad Syarhabil Ahmad², Humayun Riaz¹, Syed Atif Raza³, Shahzad Hussain⁴, Omer Salman Qureshi⁵, Povydysh Maria⁶, Zainab Hamzah⁷ and Osama Javed³

*Address all correspondence to: aslammuhammadshahzad@gmail.com

1 Rashid Latif College of Pharmacy, Lahore, Pakistan

2 School of Bioprocess Engineering, Universiti Malaysia Perlis, Arau, Perlis, Malaysia

3 University College of Pharmacy, University of the Punjab, Lahore, Pakistan

4 National Institute of Health, Islamabad, Pakistan

5 Department of Pharmacy, University of the Lahore, Pakistan

6 Saint-Petersburg State Chemical-Pharmaceutical Academy, Ministry of Healthcare of the Russian Federation, St. Petersburg, Russia

7 Department Chemical Engineering, Faculty of Engineering Technology (FETech), Universiti Malaysia Perlis, Padang Besar, Perlis, Malaysia

References

- [1] Serena TE. A global perspective on wound care. *Advances in Wound Care*. 2014;**3**(8): 548-552
- [2] Gupta VK, Pathak SS, Jain MK. Evaluation of burn wound healing property of ocimum sanctum by monitoring of period of re-epithelization in rabbits. *International Journal of Basic & Clinical Pharmacology Research*. 2016;**5**(1):146-148
- [3] Archana D, Dutta J, Dutta PK. Evaluation of chitosan nano dressing for wound healing: Characterization, in vitro and in vivo studies. *International Journal of Biological Macromolecules*. 2013;**57**:193-203
- [4] Huang C, Leavitt T, Bayer LR, Orgill DP. Effect of negative pressure wound therapy on wound healing. *Current Problems in Surgery*. 2014;**51**(7):301-331
- [5] Jivad N, Bahmani M, Asadi-Samani M. A review of the most important medicinal plants effective on wound healing on ethnobotany evidence of Iran. *Der Pharmacia Lettre*. 2016;**8**(2):353-357
- [6] Moura LIF, Dias AMA, Carvalho E, De Sousa HC. Recent advances on the development of wound dressings for diabetic foot ulcer treatment—A review. *Acta Biomaterialia*. 2013;**9**(7):7093-7114
- [7] Cheng CF et al. A fragment of secreted Hsp90 α carries properties that enable it to accelerate effectively both acute and diabetic wound healing in mice. *The Journal of Clinical Investigation*. 2011;**121**(11):4348-4361
- [8] Asai J et al. Topical application of ex vivo expanded endothelial progenitor cells promotes vascularisation and wound healing in diabetic mice. *International Wound Journal*. 2013;**10**(5):527-533
- [9] Barrientos S, Brem H, Stojadinovic O, Tomic-Canic M. Clinical application of growth factors and cytokines in wound healing. *Wound Repair and Regeneration*. 2014;**22**(5):569-578
- [10] Gil ES, Panilaitis B, Bellas E, Kaplan DL. Functionalized silk biomaterials for wound healing. *Advanced Healthcare Materials*. 2013;**2**(1):206-217
- [11] Kumar S, Pandey AK. Chemistry and Biological Activities of Flavonoids: An Overview. *The Scientific World Journal*. 2013;**2013**(162750):16. <https://doi.org/10.1155/2013/162750>
- [12] Houghton PJ, Hylands PJ, Mensah AY, Hensel A, Deters AM. In vitro tests and ethnopharmacological investigations: Wound healing as an example. *Journal of Ethnopharmacology*. 2005;**100**:100-107
- [13] Surjushe A, Vasani R, Saple DG. Aloe vera: A short review. *Indian Journal of Dermatology*. 2008;**53**(4):163-166
- [14] Hu Y, Xu J, Hu Q. Evaluation of antioxidant potential of Aloe vera (*Aloe barbadensis* Miller) extracts. *Journal of Agricultural and Food Chemistry*. 2003;**51**(26):7788-7791

- [15] Hutter JA et al. Antiinflammatory C-glucosyl chromone from *Aloe barbadensis*. *Journal of Natural Products*. 1996;**59**(5):541-543
- [16] Irshad S, Butt M, Younus H. In-vitro antibacterial activity of *Aloe barbadensis* Miller (*Aloe vera*). *International Research Journal*. 2011;**1**(2):59-64
- [17] Hashmat I, Azad H, Ahmed A. Neem (*Azadirachta indica* A. Juss)—A Nature's drugstore : An overview. *International Research Journal of Biological Sciences*. 2012;**1**(6):76-79
- [18] Koul O, Isman MB, Ketkar CM. Properties and uses of neem, *Azadirachta indica*. *Canadian Journal of Botany*. 1990;**68**(1):1-11
- [19] Krup V, Prakash LH, Harini A. Pharmacological activities of turmeric (*Curcuma longa* Linn): A review. *Journal of Homeopathy & Ayurvedic Medicine*. 2013;**2**(4):133. DOI: 10.4172/2167-1206.1000133
- [20] Rafieian-Kopaei M, Nasri H, Sahinfard N, Rafieian M, Rafieian S, Shirzad M. Turmeric: A spice with multifunctional medicinal properties. *Journal of Herbmmed Pharmacology*. 2014;**3**(1):5-8
- [21] Aslam MS, Ahmad MS, Mamat AS. A review on phytochemical constituents and pharmacological activities of *Clinacanthus nutans*. *International Journal of Pharmacy and Pharmaceutical Sciences*. 2015;**7**(2):30-33
- [22] Ayyanar M, Ignacimuthu S. Herbal medicines for wound healing among tribal people in southern India: Ethnobotanical and scientific evidences. *International Journal of Applied Research in Natural Products*. 2009;**2**(3):29-42
- [23] Owoyele BV, Oguntoye SO, Dare K, Ogunbiyi BA, Aruboula EA, Soladoye AO. Analgesic, anti-inflammatory and antipyretic activities from flavonoid fractions of *Chromolaena odorata*. *Journal of Medicinal Plants Research*. 2008;**2**(9):219-225
- [24] Orhan IE. *Centella asiatica* (L.) Urban: From traditional medicine to modern medicine with neuroprotective potential. *Evidence-based Complementary and Alternative Medicine*. 2012;**2012**(946259):8. <https://doi.org/10.1155/2012/946259>
- [25] Dipankar CR, Barman SK, Shaik MM. Current updates on *Centella asiatica*: Phytochemistry, pharmacology and traditional uses. *Journal of Medicinal Plants Research*. 2013;**3**(4):20-36
- [26] Hiradeve SM, Rangari VD. *Elephantopus scaber* Linn.: A review on its ethnomedical, phytochemical and pharmacological profile. *Journal of Applied Biomedicine*. 2014;**12**(2):49-61
- [27] Papiya B, Rana C. A comprehensive phyto-pharmacological review of *Euphorbia neriifolia* Linn. *Pharmacognosy Reviews*. 2008;**2**(4):57-66
- [28] Ahmed SA, Nazim S, Siraj S, Siddik PM, Wahid CA. *Euphorbia neriifolia* Linn: A phytopharmacological review. *International Research Journal of Pharmacy*. 2011;**2**(5):41-48
- [29] Saxena J, Khare S. A brief review on: Therapeutical values of *Lantana camara* plant. *International Journal of Pharmacy and Life Sciences*. 2012;**3**(3):1551-1554
- [30] Kalita S, Kumar G, Karthik L, Rao KVB. A review on medicinal properties of *Lantana camara* Linn. *Research Journal of Pharmacy and Technology*. 2012;**5**(6):711-715

- [31] Mundada S, Shivhare R. Pharmacology of *Tridax procumbens* a weed: Review. International Journal of PharmTech Research. 2010;**2**(2):1391-1394
- [32] Pietta PG. Flavonoids as antioxidants. Journal of Natural Products. 2000;**63**(7):1035-1042
- [33] Cushnie TPT, Lamb AJ. Antimicrobial activity of flavonoids. International Journal of Antimicrobial Agents. 2005;**26**(5):343-356
- [34] Procházková D, Boušová I, Wilhelmová N. Antioxidant and prooxidant properties of flavonoids. Fitoterapia. 2011;**82**(4):513-523
- [35] Ren W, Qiao Z, Wang H, Zhu L, Zhang L. Flavonoids: Promising anticancer agents. Medicinal Research Reviews. 2003;**23**(4):519-534
- [36] Kim HP, Son KH, Chang HW, Kang SS. Anti-inflammatory plant flavonoids and cellular action mechanisms. Journal of Pharmacological Sciences. 2004;**96**(3):229-245
- [37] Chirumbolo S. Flavonoids in propolis acting on mast cell-mediated wound healing. Inflammopharmacology. 2012;**20**(2):99-102
- [38] Daglia M. Polyphenols as antimicrobial agents. Current Opinion in Biotechnology. 2012;**23**(2):174-181
- [39] Williams RJ, Spencer JP, Rice-Evans C. Flavonoids: Antioxidants or signalling molecules? Free Radical Biology & Medicine. Apr. 2004;**36**(7):838-849
- [40] Sharma N, Dobhal M, Joshi Y, Chahar M. Flavonoids: A versatile source of anticancer drugs. Pharmacognosy Reviews. 2011;**5**(9):1
- [41] Schmidt CA et al. Catechin derivatives from *Parapiptadenia rigida* with in vitro wound-healing properties. Journal of Natural Products. 2010;**73**(12):2035-2041
- [42] Ozay Y et al. Evaluation of the wound healing properties of luteolin ointments on excision and incision wound models in diabetic and non-diabetic rats. Records of Natural Products. 2018;**12**(4):350-366
- [43] Tran NQ, Joung YK, Lih E, Park KD. In situ forming and rutin-releasing chitosan hydrogels as injectable dressings for dermal wound healing. Biomacromolecules. 2011;**12**(8):2872-2880
- [44] Mensah AY et al. Effects of *Buddleja globosa* leaf and its constituents relevant to wound healing. Journal of Ethnopharmacology. 2001;**77**(2-3):219-226
- [45] Muhammad AA, Pauzi NAS, Arulselvan P, Abas F, Fakurazi S. In vitro wound healing potential and identification of bioactive compounds from *Moringa oleifera* Lam. BioMed Research International. 2013;**2013**
- [46] Muralidhar A, Sudhakar Babu K, Sankar TR, Reddanna P, Latha J. Wound healing activity of flavonoid fraction isolated from the stem bark of *Butea monosperma* (Lam) in albino wistar rats. European Journal of Experimental Biology. 2013;**3**(6):1-6
- [47] Ergene Öz B, Saltan İçsan G, Küpeli Akkol E, Süntar İ, Bahadır Acıkara Ö. Isoflavonoids as wound healing agents from *Ononidis Radix*. Journal of Ethnopharmacology. 2018;**211**:384-393

- [48] Hertog MGL, Hollman PCH, Hertog MGL, Katan MB. Content of potentially anticarcinogenic flavonoids of 28 vegetables and 9 fruits commonly consumed in the Netherlands. *Journal of Agricultural and Food Chemistry*. 1992;**40**(12):2379-2383
- [49] Toh JY, Tan VMH, Lim PCY, Lim ST, Chong MFF. Flavonoids from fruit and vegetables: A focus on cardiovascular risk factors. *Current Atherosclerosis Reports*. Dec. 2013;**15**(12):368
- [50] Ambiga S, Narayanan R, Gowri D, Sukumar D, Madhavan S. Evaluation of wound healing activity of flavonoids from *Ipomoea carnea* Jacq. *Ancient Science of Life*. Jan. 2007; **26**(3):45-51