We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Chiral Mono- and α -Diimines and Their Pd(II) Complexes with Anticancer Activity

Guadalupe Hernández, Daniela Gutiérrez, Gloria E. Moreno, Oscar Portillo, René Gutiérrez and Eduardo Brambila

Abstract

The aim of this review is to provide mainly an outlook of the synthesis and characterization of chiral mono- and α -diimines ligands and their Pd(II) complexes carried out in our group in the last few years. Some other issues with simple chiral imines synthesized in our lab are also outlined. The report includes details about their versatile coordination patterns, biological activity in cancer cell lines, and engaging properties in different fields, such as materials science.

Keywords: chiral imines, Pd complexes, solvent-free reactions, anticancer activity

1. Introduction

The importance of Schiff bases resides in their structural variety as well as their ability to form a wide range of appealing structural arrangements depending of the constituents parent molecules with transition metals by acting as *N*-donor ligands, affording mono-, bi- and polynuclear complexes [1–3]. Accordingly, Schiff bases display a broad range of useful biological activities such as, *inter alia*, antibacterial, antifungal, antidiabetic, anti-inflammatory, and anticancer agents generating a huge interest in the medicine field [4–8]. The proper choice of the ligands in metal complex synthesis is essential for the activity that they could present since they determine some aspects like reactivity and lipophilicity.

We have focused our attention on the synthesis of chiral compounds since chirality is almost omnipresent in a broad range of organic molecules in the human body such as proteins, enzymes, amino acids, carbohydrates, and nucleosides. The body acts like a chiral selector metabolizing enantiomers by separate pathways and generating different pharmacological activities. For that reason, the current approach is to target specific molecules by designing more selective drugs, especially in chemotherapy where the distinction between cancerous and normal cells is essential for the success of the treatment and the reduction of the toxicity.

Likewise, the search of more eco-friendly procedures in the synthesis of organic molecules is one of the goals of our research group. Green Chemistry techniques like the use of microwave irradiation and solvent-free reactions display numerous advantages such as shorter reaction times, minimum waste, operational simplicity as well as reduction of thermal degradative byproducts along with cleaner work-up and generally higher yields [9, 10].

On the other hand, the discovery of anticancer activity of the cisplatin was a key event for the introduction of metal-based compounds to medicine, and the interest on these kind of compounds increased significantly in the last decades due to their ability to coordinate ligands in a three-dimensional configuration and bind to specific cell targets. Platinum-based drugs, particularly cisplatin, are widely used in the treatment of different types of cancer, but the toxicity and high resistance that they present limits their use. Therefore, the major challenge for chemists is the design of new drugs with less side effects. Efforts have been made to consider other metal-based complexes with cytotoxic properties, such as palladium complexes. They are known to show structural and thermodynamic analogy in regard to Pt(II) complexes, and display versatile coordination behavior and interesting properties. Palladium complexes of various donor-atom ligands have been found to possess engaging anti-tumor activity, as well as anti-inflammatory, anti-microbial, antiviral and antifungal properties [11, 12].

2. Chiral Pd(II) complexes

The incorporation of optically pure aromatic amines into α -dicarbonylic compounds bearing aromatic rings such as benzil in a 1/1 ratio generating enantiopure α -ketoimines was the first step for our investigations, considering that a flexible $X=C-C=N$ ($X = O, N$) skeleton could lead to diverse coordination modes [13]. Then, the chiral mono-imine derived from (*S*)-(-)-1-phenylethylamine and benzyl under microwave radiation in solvent-free conditions led to the formation of the *N*-donor ligand (*S*)-(-)-(1-phenylethylimino)benzylphenylketone **1** which was allowed to react with K_2PdCl_4 giving two Pd complexes: a mono- **2** and a dinuclear Pd(II) **3** complexes (Figure 1).

On the other hand, *in vitro* assays are essential to determine the capacity of the compounds to modify basic cellular functions on different cancer cells. We have

Figure 1.
Mono- and dinuclear chiral Pd(II) complexes **2** and **3** with their respective IC_{50} values.

employed sulforhodamine B staining to determine the cytotoxicity of our complexes, given the ease and high reproducibility of this method. By keeping constant the panel of human cancer cell lines (U251: central nervous system, PC-3: prostate cancer, K562: leukemia, HCT: colon cancer and MCF-7: breast cancer) we are able to compare the effects of the compounds in each cell and determine how the variations on the structure affect the activity. Those cell lines represent the most common types of cancer [14].

Complex 2 presented a common square planar geometry at the metal center with two Cl atoms *trans* and two ligands bonded through the N atoms N8 and N58 in a *trans* configuration while complex 3 is a dinuclear Pd(II) complex with one molecule in the asymmetric unit. For the binuclear complex, the coordination is carried through the N atoms (N8 and N58), like complex 2, and the C atoms of the phenyl rings of the imino functions (C14 and C64). The low level of electronic delocalization in the ligand induced a high level of flexibility in the formation of complex 3, producing a major distortion, due to the bite angles Cl-Pd-Cl and N-Pd-Cl.

The complexes 2 and 3 were tested by sulforhodamine B assays against U251, PC-3, K562, HCT and MCF-7 human cancer cell lines. Both compounds displayed cytotoxic activity, especially toward K562 (IC_{50} : 26.5 ± 0.4 and 14.8 ± 1.1 μ M for complex 2 and 3, respectively) and MCF-7 (IC_{50} : 34.5 ± 2.5 and 13.1 ± 1.0 μ M for complex 2 and 3, respectively). In general, the binuclear complex was slightly better for all cell lines exhibiting lower IC_{50} values, while complex 2 surpassed the dose of 100 μ M in U251 and HCT-15 cell lines [14].

Also, we have reported the synthesis of cyclopalladated compounds. Considering that our previous compounds displayed attractive properties, we decided to vary the substituents, replacing the aromatic rings in the α -dicarbonylic compounds by aliphatic substituents, such as two methyl groups and attaching also two chiral entities, i.e., to prepare α -diimines, as such kind of compounds have also a flexible N=C—C=N skeleton, displaying outstanding electron donor and acceptor properties and can potentially act in a variety of coordination modes. Then, the chiral diimines 4–5 were synthesized under solvent-free conditions starting from (S)-(-)-1-phenylethylamine and (S)-(-)-1-(4-methylphenyl) ethylamine with 2,3-butanedione, respectively. The reaction between Na_2PdCl_4 and each of the ligands 4–5 in a MeOH solution at ambient temperature led to the formation of the complexes 6 and 7 (Figure 2) [15]. In this case, the complex 6 is mononuclear

Figure 2.
 Synthesis of chiral Pd complexes 6–7.

with the Pd(II) center adopting a distorted square planar Pd[N₂CCl] coordination geometry where one benzene group is bonded to the metal center while the other is free of coordination. The steric hindrance produced by the benzene group and the formation of a Pd-C bond apparently blocked the dimerization of the complex. The solid obtained from complex **7** was not able to crystallize.

Both complexes **6** and **7** exhibited cytotoxic activity toward the panel of cultured cell lines previously mentioned, mainly against U251 and K562 cancer cells with IC₅₀ values of 19.8 and 22.5 μM for complex **7**, and 23.6 and 25.44 μM for **6**, respectively. According to the data, **6** cannot be considered a good candidate as an anticancer agent since its IC₅₀ values are too high for PC-3 and HCT-15, exceeding the dose of 100 μM. These compounds offer a better activity against U251 cell line compounds than the α-ketoimine complexes previously mentioned.

Thereafter, we carried out the synthesis of new unsymmetrical α-diimines by replacing one methyl group with a hydrogen atom and enlarging the number of chiral amines. A different method was used with the aim to improve the yields. Then, methylglyoxal and optically active aromatic and alicyclic primary amines were stirred in diethyl ether with Na₂SO₄ for 24 hours at room temperature leading to the formation of the ligands **8–11** (Figure 3). Solutions of the ligands **8–11** in benzene were treated this time with dichloro(1,5-cyclooctadiene) palladium (II) and stirred at room temperature under argon atmosphere to form complexes **12–15** (Figure 4). Worth-mentioning is that the coordination of the ligands took place in two different modes: chelating (σ, σ, N, N') and monodentate (σ-N) [16].

Complexes **12** and **13** expose a *s-cis* chelate system and although they are chemically similar they crystallize in different way, in two distinct space groups. We believe that the crystal symmetry modification is a consequence of the crystallization rather than small conformational variations. The complex **15** displays two diimine ligands which are coordinated to the metal center in a *trans* square planar geometry, and the same behavior is observed in complex **14**. The importance of the *trans*-geometry around the Pd center has been attributed to the comparatively higher cytotoxicity values as those for *cis*-isomers.

It seems that the small substituents on the imine N atoms facilitates the orientation toward σ, σ, N, N' coordination mode, stabilizing the complex through the chelate effect, while the monodentate (σ-N) coordination mode is favored by sterically hindered systems.

The results of the cytotoxic assay showed that Pd complexes with monodentate (σ-N) coordination mode (**14** and **15**) displayed IC₅₀ values >100 μM; these complexes were dismissed for further assays because the doses required to inhibit cell

Figure 3.
Synthesis of chiral α-diimine ligands **8–11**.

Figure 4.
 Synthesis of chiral Pd(II) complexes 12–15.

growth were too high. Complexes 12 and 13 also possessed cytotoxic activity against U251, PC-3, K562, HCT and MCF-7 cell lines, where IC_{50} ranged from 66 to 91 μM .

As such results were unpromising, we reconsidered the α -dicarbonylic compounds bearing aromatic rings, but this time with heterocyclic entities. By using the method previously used (microwave irradiation in solvent-free conditions), the chiral α -ketoimines 16–17 were synthesized from (S)-(-)-1-phenylethylamine and (S)-(-)-1-(4-methylphenyl) ethylamine with 2,2'-pyridil, respectively (Figure 5).

Complexes 18–19 (Figure 6) were synthesized by the reaction between $Pd(COD)Cl_2$ and each ligand 16–17 in a solution of benzene. It was not possible to obtain a monocystal of complex 19, however the crystal data of 18 showed that α -ketoimine 16 is a bidentate ligand and Pd(II) displayed a square-planar coordination geometry. In the case of 16, the conjugation of imine and carbonyl double bonds with the aromatic systems and the substitution of vicinal C1 and C2 by pyridil rings implied that the ligand adopted a *gauche* conformation [17].

The data from the sulforhodamine B assay evidenced that none of the compounds possess cytotoxicity toward K562, however they are able to inhibit cell

Figure 5.
 Synthesis of chiral imines 16–17.

PC-3: 29 μ M, U251: 37 μ M,
HCT-15: 31 μ M, MCF-7: 22 μ M and
K562: >100 μ M

PC-3: 45 μ M, U251: 48 μ M,
HCT-15: 36 μ M, MCF-7: 22 μ M and
K562: >100 μ M

Figure 6.
Chiral Pd(II) complexes **18–19**.

growth in U251, PC-3, HCT-15 and MCF-7, being **18** slightly better than **19** for all cell lines. The studies suggest that the nature of the aromatic rings have an impact in the cytotoxicity and the coordination mode.

Such results were not particularly impressive (at least a factor of 10 poorer than cisplatin), but they certainly do show variations in activity as well in the other cases.

It must be pointed out that even when the Pd-Schiff Base-complexes displayed cell growth inhibition against different classes of cancer, the IC₅₀ that they have showed are not comparable with cisplatin. In general, Pd(II) complexes are kinetically less stable than those of Pt(II), by losing their structural integrity in biological fluids in a short period of time due to their rapid exchange. More specific studies *in vitro* and *in vivo* need to be done to determine their toxicity and to understand in a better way the mechanisms of action since it will aid the development of more efficient palladium-based drugs.

On the other hand, considering other alternatives to the flexible X=C—C=N (X = O, N) skeleton, for example as a heterodiene, we have also reported the microwave-assisted Diels-Alder [4+2] cycloaddition reaction of the optically pure α -ketoimines **20–21** and α -diimines **22–23**, with fullerene C₆₀. The chiral α -ketoimines **20–21** were readily synthesized in quantitative yield under solvent-free conditions starting from (S)-(-)-1-phenylethylamine and (S)-(-)-1-(4-methylphenyl) ethylamine with pyruvaldehyde, respectively, and upon reaction of C₆₀ under focused-microwave irradiation in benzene, after 20 min the formation of the adducts **24–25** was observed (**Figure 7**) [18].

With the chiral α -diimines **22–23**, which were also readily prepared from (S)-(-)-1-phenylethylamine and (S)-(-)-1-(4-methylphenyl) ethylamine with pyruvaldehyde, respectively, the adducts **26–27** were obtained (**Figure 8**).

In addition, extending our studies to include some other transition metals, we have reported the preparation of chiral Hg(II) complexes with simpler chiral imines **28–30** as they present some relevant crystallographic features along with antimicrobial activity [19]. Thus, the solvent-free reaction of 2-pyridylcarboxaldehyde with optically active aromatic and alicyclic primary amines afforded the chiral imines **28–30** in almost quantitative yields (see **Figure 9**).

Solutions of the chiral imines **28–30** in methanol were treated with HgCl₂ with stirring at room temperature for 1 h, leading to the formation of complexes **31–33** (**Figure 10**).

Figure 7.
 Adducts **24–25** synthesized.

Figure 8.
 Adducts **26–27** synthesized.

Figure 9.
 Synthesis of chiral imines **28–30**.

Likewise, preliminary data have revealed that chiral imines **34–37** derived from 2-pyridylcarboxaldehyde and the optically active aromatic amines (*S*)-(-)-1-(4-methylphenyl) ethylamine, (*S*)-(-)-1-(4-methoxyphenyl) ethylamine, (*S*)-(-)-1-(4-chlorophenyl) ethylamine and (*R*)-(+)-1-(4-fluorophenyl) ethylamine under solvent-free conditions (**Figure 11**) were allowed to react with Zn(CLO)₄ affording

Figure 10.
Chiral Hg(II) complexes 31–33.

Figure 11.
Chiral imines 34–37.

Zn complexes 38–41 (**Figure 12**) with cytotoxic activity against the aforementioned human cancer cell lines as well as low toxicity in brine shrimps, along with antibacterial activity against *P. aeruginosa*, *E. coli* and *S. aureus*. Such results will be reported in due time.

On the other hand, simpler chiral imines have triggered interest in some other fields, especially in materials science; where by changing the substituents in the chiral moiety can afford morphological, optical and structural changes resulting in photoluminescent properties, which are extremely interesting since the viewpoint of physicists. In this context, we have recently reported a series of halogenated

Figure 12.
 Chiral Zn(II) complexes 38–41.

Figure 13.
 Chiral halogenated imines 42–44.

Figure 14.
 Chiral imines 45–48.

imines (**Figure 13**) derived from 2-naphtaldehyde and optically pure halogenated amines, under solvent-free conditions. As a result, imines **42–44** with a lamellar morphology exhibited photoluminescent properties. By changing the halogen atoms in the chiral moiety of the imines, the crystalline packing was modified. The bands observed in the visible region are caused by interstitial defects, vacancies,

grain boundaries and stacking faults in the crystals. The intensity of the bands increased in the following order: $\text{—F} < \text{—Cl} < \text{—Br}$, according to the increase in atomic radii. These features result attractive because of their possible applications in organic electroluminescent devices, organic light- emitting diodes, etc. [20, 21].

Likewise, in a series of chiral imines derived from 2-naphtaldehyde but with the halogen atoms in the *para*-position of the benzene ring of the amines replaced by other functional groups, such as —CH_3 , —OCH_3 and naphthyl groups, imines **45–48** (**Figure 14**) exhibited green luminescence. As the previous results, the variations on the functional group as well as the molecular packing determined the morphological changes and consequently the luminescent properties of the imines [22].

3. Conclusion

The chemistry of Schiff bases and their transition metal complexes, especially Pd, is a field that is being noticed not only for their remarkable biological properties but also for their extensive applications in other fields. This area requires further studies to be carried out, and improvements in the permeability and transport are some of the factors to take into account in the design of these metal-based complexes.

Acknowledgements

Support from VIEP-UAP is acknowledged.

Conflict of interest

The authors declare no conflict of interest.

Author details

Guadalupe Hernández^{1*}, Daniela Gutiérrez¹, Gloria E. Moreno¹, Oscar Portillo¹, René Gutiérrez¹ and Eduardo Brambila²

¹ Laboratorio de Síntesis de Complejos, Facultad de Ciencias Químicas, Universidad Autónoma de Puebla, Ciudad Universitaria, Puebla, Mexico

² Laboratorio de Química Clínica, Facultad de Ciencias Químicas, Universidad Autónoma de Puebla, Ciudad Universitaria, Puebla, Mexico

*Address all correspondence to: guadalupe.hernandez@correo.buap.mx

IntechOpen

© 2018 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] Qin W, Long S, Panunzio M, Biondi S. Schiff bases: A short survey on an evergreen chemistry tool. *Molecules*. 2013;**18**:12264-12289. DOI: 10.3390/molecules181012264
- [2] Vigato PA, Tamburini S. The challenge of cyclic and acyclic Schiff bases and related derivatives. *Coordination Chemistry Reviews*. 2004;**248**:1717-2128. DOI: 10.1016/j.ccr.2003.09.003
- [3] Martin SF. Recent applications of imines as key intermediates in the synthesis of alkaloids and novel nitrogen heterocycles. *Pure and Applied Chemistry*. 2009;**81**:195-204. DOI: 10.1351/PAC-CON-08-07-03
- [4] Kajal A, Bala S, Kamboj S, Sharma N, Saini V. Schiff bases: A versatile pharmacophore. *Journal of Catalysis*. 2013;**2013**:1-12. Art.ID 893512. DOI: 10.1155/2013/893512
- [5] Zoubi WA, Al-Hamdani AAS, Kaseem M. Synthesis and antioxidant activities of Schiff bases and their complexes: A review. *Applied Organometallic Chemistry*. 2016;**30**:810-817. DOI: 10.1002/aoc.3506
- [6] da Silva CM, da Silva DL, Modolo LV, Alves RB, de Resende MA, Martins CVB, et al. Schiff bases: A short review of their antimicrobial activities. *Journal of Advanced Research*. 2011;**2**:1-8. DOI: 10.1016/j.jare.2010.05.004
- [7] Przybylski P, Huczynski A, Pyta K, Brzezinski B, Bartl F. Biological properties of Schiff bases and azo derivatives of phenols. *Current Organic Chemistry*. 2009;**13**:124-148. DOI: 10.2174/138527209787193774
- [8] Abd El-Halim HF, Omar MM, Mohamed GG, El-Ela Sayed MA. Spectroscopic and biological activity studies on tridentate Schiff bases and their transition metal complexes. *European Journal of Chemistry*. 2011;**2**:178-188. DOI: 10.5155/eurjchem.2.2.178.188.240
- [9] Andrade CKZ, Dar AR. Applying green processes and techniques to simplify reaction work-ups. *Tetrahedron*. 2016;**72**:7375-7391. DOI: 10.1016/j.tet.2016.09.055
- [10] Sarkar A, Santra S, Kundu SK, Hajra A, Zyryanov GV, Chupakin ON, et al. A decade update on solvent and catalyst-free neat organic reactions: A step forward towards sustainability. *Green Chemistry*. 2016;**18**:4475-4525. DOI: 10.1039/c6gc01279e
- [11] Kapdi AR, Fairlamb JS. Anti-cancer palladium complexes: A focus on PdX_2L_2 palladacycles and related complexes. *Chemical Society Reviews*. 2014;**43**:4751-4777. DOI: 10.1039/c4cs00063c
- [12] Lazarevic T, Rilak A, Bugarcic ZD. Platinum, palladium, gold and ruthenium complexes as anticancer agents: Current clinical uses, cytotoxicity studies and future perspectives. *European Journal of Medicinal Chemistry*. 2017;**142**:8-31. DOI: 10.1016/ejmech.2017.04.007
- [13] Peláez MA, Ramírez T, Martínez M, Sharma P, Álvarez C, Gutiérrez R. Synthesis, crystal structures and anticancer activity of the new chiral mono- and dinuclear palladium (II) complexes derived from (S)-(-)-(1-phenylethylimino)benzylphenylketone. *Zeitschrift für Anorganische und Allgemeine Chemie*. 2004;**630**:1489-1494. DOI: 10.1002/zaac.200400140
- [14] Monks A, Scudiero D, Skehan P, Shoemaker R, Paul K, Vistica D, et al. Feasibility of a high-flux anticancer drug screen using a diverse panel of cultured human tumor cell lines.

Journal of the National Cancer Institute. 1991;**83**:757-766. DOI: 10.1093/jnci/83.11.757

[15] Cruz S, Bernès S, Sharma P, Vazquez R, Hernández T, Portillo R, et al. Synthesis, characterization and anticancer activity of new palladacycles derived from chiral α -diimines. *Applied Organometallic Chemistry*. 2010;**24**:8-11. DOI: 10.1002/aoc.1570

[16] Vázquez J, Bernès S, Sharma P, Pérez J, Hernández G, Tovar A, et al. Synthesis, characterization and anticancer activity of new chiral Pd (II)-complexes derived from unsymmetrical α -diimines ligands. *Polyhedron*. 2011;**30**:2514-2522. DOI: 10.1016/j.poly.2011.06.020

[17] Gutiérrez D, Bernès S, Hernández G, Portillo O, Moreno GE, Sharma M, et al. New chiral α -ketoimine-Pd(II) complexes and their anticancer activity. *Journal of Coordination Chemistry*. 2015;**68**:3805-3813. DOI: 10.1080/00958972.2015.1084618

[18] Tovar A, Peña U, Hernández G, Portillo R, Gutiérrez R. Microwave-assisted synthesis of new adducts from the Diels-Alder [4+2]-cycloaddition reaction of chiral α -oxo imines and α -diimines with fullerene C60. *Synthesis*. 2007:22-24. DOI: 10.1055/s-2006-958932

[19] Hernández G, Bernes S, Portillo O, Ruíz A, Moreno GE, Gutiérrez R. Crystal structures of three mercury(II) complexes [HgCL₂L] where L is a bidentate chiral imine ligand. *Acta Crystallographica Section E*. 2015;**E71**:1462-1466. DOI: 10.1107/S2056989015020368

[20] Hernández-Téllez G, Moreno GE, Bernes S, Mendoza A, Portillo O, Sharma P, et al. Crystal structures of ten enantiopure Schiff bases bearing a naphthyl group. *Acta Crystallographica*

Section E. 2016;**E72**:583-589. DOI: 10.1107/S2056989016004692

[21] Sosa A, Portillo O, Hernández G, Moreno GE, Juárez G, Chávez M, et al. Optical and morphological changes in the solid state of chiral imines bearing halogen substituents (F, Cl and Br). *Material Letters*. 2017;**189**:313-316. DOI: 10.1016/j.matlet.2016.11.053

[22] Portillo Moreno O, Meléndez Bustamante FJ, Chávez Portillo M, Moreno Morales GE, Hernández Téllez G, Sosa Sánchez A, et al. Photoluminescent green emission band induced by the systematic change of $-\text{CH}_3$, $-\text{OCH}_3$ and naphthyl groups in chiral imines. *Crystal Growth & Design*. 2018;**18**:660-668. DOI: 10.1021/acs.cgd.7b00851