We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Tea and Oral Health

Aswini Y. Balappanavar

Abstract

Tea consumption as a beverage is very common in various parts of the world. It has attained a worldwide liking and measure of social status in many parts. Tea contains various chemicals which have positive effects on health from heart to skin. It has been associated with the cure of aging to potent anticancer agent also. Considering these facts an attempt was made to establish a relation between tea and oral health. Tea has its effects on oral microorganisms, anticariogenic properties, and reduction of gingivitis as well as periodontitis. A cup of tea immediately after lunch had reduced dental caries in children and rinsing with 0.2% Chinese green tea decreased plaque and the gingival index significantly. Tea has been found to be effective against oral cancer, precancerous lesions and conditions as well. Hence tea has been rightly said as a functional food for health. Green tea has shown to have bactericidal effects on *Porphyromonas gingivalis* and *Prevotella* species. The gingival inflammation is reduced and a marked reduction in pocket size has been noticed. Tea selectively induces p57 and apoptosis as well as inhibits the growth and invasion of oral carcinoma.

Keywords: tea, oral cancer, antimicrobial properties, catechins, EGCG, gingivitis, periodontitis

1. Introduction

Ancient Chinese and Japanese medicines have emphasized the fact that green tea consumption could heal wounds and cure diseases. In 2737 BC Chinese had a belief on tea for its healing powers. Lu Yu who was a scholar in China, who had written a treatise in AD 780, entitled *Cha Ching*, states that 'tea tempers the spirits, harmonizes the mind, dispels the lassitude, relieves fatigues, awakens thought, prevents drowsiness, refreshes the body and clears the perspective faculties' [1]. In the ancient system of Medicine, Ayurveda had listed tea in the group of medicaments as 'rasayanas' that bring about positive health, resistance to diseases and assured full lifespan of quality living, unlike drugs that cure after disease has struck [2].

Tea has been considered a desired drink worldwide. The world consumed 2.9 million tones of tea in 2016 which was more when compared to 1.6 in 2002 and this may shoot to 3.3 million tones in 2021 (Euromonitor data). More than half of tea is contributed by Asia and the top three markets by per capita consumption of tea are Turkey, Ireland and UAE [3]. Different forms of tea are obtainable in the market claiming varied health benefits. These categories are based on the oxidation process. There are six different types of tea produced [4]:

- White: wilted and unoxidized
- Yellow: unwilted and unoxidized but allowed to yellow
- Green: unwilted and unoxidized
- Oolong: wilted, bruised and partially oxidized
- Black: wilted, sometime crushed, and fully oxidized which is called red tea in Chinese culture
- Post fermented: green tea that has been allowed to ferment/compost “black tea in Chinese culture”

Tea has shown to have various benefits on Health and oral health. Tea reduces the risk of several major life style related diseases which include cancer, arteriosclerosis and cardiovascular diseases, neural and obesity problems, diabetes, diseases of the kidneys and liver, pulmonary ailments, flu, SARS and even AIDS. The impact on oral health though less researched has been beneficial.

2. Tea and oral health

Oral/dental diseases are a costly burden to health care services, accounting for between 5 and 10% of total health care expenditures and exceeding the cost of treating cardiovascular disease, cancer and osteoporosis in industrialized countries [5]. In low-income countries, the cost of traditional restorative treatment of dental disease would probably exceed the available resources for health care. Dental health promotion and preventive strategies are clearly more affordable and sustainable. Although not life-threatening, dental diseases have a detrimental effect on quality of life in childhood through to old age, having an impact on self-esteem, eating ability, nutrition and health.

Oral health is related to diet in many ways, for example, through nutritional influences on craniofacial development, oral cancer and oral infectious diseases [6]. Both animal studies and experimental investigations in humans have shown that black tea extract increases plaque fluoride concentration and reduces the cariogenicity of a sugar rich diet. The protection against the causative organisms of dental caries is well documented by Banerjee in 1990 [7]. Sava et al. showed that the melanin-like pigment from black tea has immunostimulant activity [8].

2.1 Anticariogenic effects

Dental caries is a transmissible microbial disease affecting the hard tissues of teeth caused by acids from bacterial metabolism leading to demineralization and dissolution of enamel and dentin. The bacteria responsible of producing organic acids as a by-product of their metabolism of fermentable carbohydrates. The caries process is a continuum in the oral cavity resulting from many cycles of demineralization and remineralization [9].

The studies of cariostatic effects of tea were started in the 1940s and 1950s showing fluoride to be the active component [10]. Reports by many researchers have showed that the tea consumption leads to reduction in dental caries in humans and experimental animals, that tannins and fluoride were the reason for this inhibitory effect [11–14]. Despite the positive animal data supporting the positive relationship between tea and dental caries prevention, relatively little attention has

been given to this field of research. Green tea extracts, or polyphenols, have been reported to inhibit *in vitro* growth, acid production and water insoluble glucan synthesis by glucosyltransferase enzyme of *Streptococcus mutans* [12, 15–18]. Similar findings have been reported for oolong tea by Nakahara et al., in 1993 and Ooshima et al., in 1993 [18, 19]. Taiwanese green, black and oolong teas have also been shown to inhibit *in vitro* growth of selected cariogenic and periodontal pathogens [20, 21]. In an adult human study by Wu et al. [22] rinsing with black tea ten times a day for 7 days resulted in significantly less pronounced pH fall, a lower plaque index ($P < 0.05$) and lower numbers of mutans streptococci and total oral streptococci in plaque but not in saliva. Fluoride concentrations in plaque and saliva increased, reaching a maximum at day 7. Black tea and its polyphenols may benefit human oral health by inhibition of dental plaque, acidity and cariogenic microflora.

Animal studies have shown that specific pathogen-free (SPF) rats infected with *S. mutans* and then fed a cariogenic diet containing green tea polyphenols demonstrated significantly lowered caries scores [17]. Supplementing drinking water of rats with 0.1% green tea polyphenol along with a cariogenic diet also significantly reduced total fissure caries lesions [23]. Animal studies using oolong tea gave similar results and it was suggested that active substances may affect bacterial virulence factors other than the glucosyltransferase enzymes [24]. Caries were found to be significantly lower among children who drank a cup of tea immediately after lunch and the tea polyphenols, rather than fluoride, were found to be responsible for the anticariogenic effects [25]. Another study reported that rinsing with 0.2% Chinese green tea while brushing decreased plaque and the gingival index significantly [26].

Tea drinking has been attributed as one of the factors in the declining prevalence of caries in Tunisia [27]. Tea extracts have also been shown to inhibit human salivary amylase and tea consumption may reduce the cariogenic potential of starch-containing foods, such as biscuits and cakes, because tea may reduce the tendency for these foods to serve as slow-release sources of fermentable carbohydrate [28]. It is likely that cariogenic challenge in a cariogenic diet may be overcome by the simultaneous presence of green tea in the diet. An anticariogenic potential of black tea has been suggested in various *in vitro* studies. [20, 29, 30]. Black tea and its polyphenols inhibited growth, acid production, metabolism and glucosyltransferase enzyme activity of Mutans Streptococci and dental plaque bacteria.

Tea is a source of fluoride (F) as well as many other dietary trace elements. The caries-preventive effect of teas was first believed to be due to its fluoride content. More recent studies, however, have pointed out that the polyphenol contents of tea may affect plaque formation and metabolism as well [31]. The commercial tea plant, *Camellia sinensis*, takes up F from the soil by passive diffusion and concentrates it in the leaves by transpiration [32]. Due to differing soils, types of tea leaves, infusion times and methods of analysis, a great deal of variation in tea content has been found. Coupled with the various drinking habits among different people, it is very difficult to calculate the contribution of tea to total fluoride intake. A recent animal study showed that rats consuming black tea (prepared from fluoride-free water) over a 2-week period had a significantly lower rate of caries than those consuming non-fluoridated water. Furthermore, the caries scores in the group receiving tea were significantly greater than those in the group receiving fluoridated water. The authors suggested that black tea consumption attenuates the development of caries in young, caries-prone rats [14]. Wei et al. found a 15-min infusion to result in a mean fluoride concentration of 1.75 p.p.m. for 15 Chinese teas, 1.24 p.p.m. for 11 Ceylon/Indian teas and a negligible F amount for six herbal teas [32]. The bioavailability of fluoride in tea has been said to be approximately 85%. A review by Kavanagh and Renehan lists over 10 papers that measured the fluoride content of various teas [33]. Several studies have also calculated the total F intake from tea consumption.

When using the assumptions of 2.5 cups/day, 150 ml per cup and 2.2 p.p.m. F diluted in half with milk for children, an ingested range of 0.1 mg to 1.08 mg was found. Wei et al. estimated a daily F intake at a level of 1.05 mg [32]. As well as teas are unlikely to cause fluorosis by themselves, but they may be significant contributors to the total fluoride intake of children.

2.2 Tea and gingivitis/periodontal disease

Periodontal disease (PD) is one of the most omnipresent diseases of mankind, which is also second most common oral disease worldwide, after dental caries. This is a chronic condition in which a multiple and complex group of inflammatory diseases are affecting the periodontal complex i.e., tissue that surround and support the teeth (Periodontium, bone, gingival fibers). Negligence towards this condition may show further deterioration of periodontium leading to progressive loss of the alveolar bone around the teeth and subsequent loss of teeth. In fact, PD remains the most common cause of tooth loss in the world today; in the United States, it has a prevalence of 30–50% of the population and can affect up to 90% of the population worldwide [34].

Researchers have observed that for every one cup of green tea consumed per day, there was a decrease in the indicators of gingival inflammation, in turn reducing the periodontal disease. Green tea catechin has been shown to be bactericidal against *Porphyromonas gingivalis* and *Prevotella* species *in vitro*. A local slow delivery system of green catechin along with mechanical treatment was found to be effective in improving periodontal status. There was a reduction pocket size and the suppression of peptidase activities in the gingival crevicular fluid [35]. Tea catechins containing the galloyl radical (Epicatechin Gallate [ECG] and Epigallocatechin Gallate [EGCG]) possess the ability to inhibit both eukaryotic and prokaryotic cell-derived collagenase, an enzyme that plays an important role in the disruption of the collagen component in the gingival tissues of patients with periodontal disease [36–38]. Catechin derivatives have been reported to inhibit certain proteases and toxic metabolites of *P. gingivalis* and may reduce periodontal breakdown [39, 40]. Green tea catechins EGCG have also been shown to inhibit protein tyrosine phosphatase in *Prevotella intermedia* [41]. Zhu et al. have shown that purified tea polyphenols inhibited *in vitro* growth and H₂S production of *P. gingivalis* and *Fusobacterium nucleatum* associated with human halitosis [42].

The molecular and cellular effects of green tea on oral cells of smokers were researched [43]. A recent human study investigated the effect of tea polyphenols in the form of chew candies on gingival inflammation over a 4-week period [44]. The approximal plaque index (API) and sulcus bleeding index (SBI) were determined at the end of day 7 and day 28. These authors suggested that tea polyphenols might exert a positive influence on gingival inflammation.

2.3 Oral cancer

Oral cancer is a global public health problem and relevant to dentists due to proximity of this area to the work carried out by them. It is located within the top 10 ranking incidence of cancers and despite the progress in research and therapy, survival has not improved significantly in the last years, representing a continuing challenge for biomedical science. Oral cancer is a malignant neoplasia, which arises on the lip or oral cavity. It was traditionally defined as a squamous cell carcinoma (OSCC), as 90% of cancers are histologically originated in the squamous cells in the oral cavity [45].

The influence of tea on oral cancer as well as various studies has been summarized in **Tables 1–3** [44, 46–75]. Green tea polyphenols are found to induce

Sl no.	Author	Type of study	Sample	Country	Intervention	Outcome
1.	The Indian-US Head and Neck Cancer Cooperative Group, 1997	Population based chemo preventive experimental trial—6 months	64	India	Green—3.6 g per day and 5.4 g per day	Feasible
2.	Khafif A et al., 1998	<i>In vitro</i> experimental trial			(–)-epigallocatechin-3-gallate (EGCG) from green tea	A reduction of 4.4–8.5-fold in cell cycle of cancer cells
3.	Khafif A, et al., 1998	<i>In vitro</i> experimental trial—mice			EGCG [(–)-epigallocatechin-3-gallate]	Inhibited cancer cells growth
4.	Yang CS, Lee MJ, Chen L., 1999	<i>In vivo</i> experimental trial—human	6		Green tea	Increase in (–)-epigallo-catechin (EGC; 11.7–43.9 microg/ml), EGC-3-gallate (EGCG; 4.8–22 microg/ml), and (–)-epicatechin (EC; 1.8–7.5 microg/ml) levels in saliva
5.	Li N et al., 1999	Double blind randomized controlled human clinical trial— 6 months	64	China	Tea—3 g/day	Reduction in size in 37.9% of leekoplakic patients
6.	Li N, Han C, Chen J, 1999	<i>In vivo</i> animal experimental trial—hamsters			1.5% green tea, 0.1% tea pigments, and 0.5% mixed tea	Reduced buccal pouch tumor burden and the incidence of dysplasia and oral carcinoma (P < 0.01)
7.	Li N et al., 1998	Randomized control trial—human— 6 months	36		Mixed tea—3 g/day oral and 0.1% topical	Significant decrease in micronuclei (P < 0.001)
8.	Masuda M, Suzui M, Weinstein IB, 2001	<i>In vitro</i> experimental trial			Egcg—10 microg/ml	growth inhibition (P < 0.001)
9.	Hsu SD et al., 2002	<i>In vitro</i> experimental trial			Tea extracts, green tea polyphenols, (–)-epigallo-catechin-3-gallate (EGCG)	Selectively induce apoptosis only in oral carcinoma cells, EGCG inhibit the growth and invasion of oral carcinoma cells
10.	Li N et al., 2002	<i>In vivo</i> experimental trial—hamsters 18 months			0.6% green tea powder, 0.6% green tea powder + 10 mumol curcumin	Decreased the number of visible tumors and the tumor volume. Suppression of cell proliferation, induction of apoptosis, and inhibition of angiogenesis
11.	Masuda M et al., 2003	<i>In vitro</i> experimental trial			10 or 30 microg of EGCG	50% inhibition of growth of carcinoma cells
12.	Lee MJ et al., 2004	<i>In vivo</i> cross over Clinical trial—human			2 g of black tea, 2 g of green tea	Concen-trations of catechins (C(max) = 131.0–2.2 micro M) and theaflavins (C(max) = 1.8–0.6 micro M) were observed in saliva in the 1st hour

Sl no.	Author	Type of study	Sample	Country	Intervention	Outcome
13.	Srinivasan P, Sabitha KE, Shyamaladevi CS. 2004	<i>In vivo</i> experimental trial—rats 1 month			Green tea polyphenols (GTP) (200 mg/kg)	Enhances the cellular thiol status thereby mitigate oral cancer
14.	Babich H et al., 2005	<i>In vitro</i> experimental trial			Catechin gallate (CG), epigallocatechin gallate (EGCG), epigallocatechin (EGC), catechin (C) and epicatechin (EC)	Reduced carcinoma HSC-2 cells of oral cavity
15.	Gonzalez de Mejia E et al., 2005	<i>In vitro</i> controlled experimental study			Yerba mate tea products	Inhibition of topoisomerase II (cancer cell proliferation)
16.	Hsu S et al., 2005	<i>In vitro</i> experimental study			Egcg	p21WAF1 is involved in EGCG-induced growth arrest of OSC2 cells
17.	Halder A et al., 2005	<i>In vivo</i> human experimental trial—1 year	82	India	Black tea	Significant decrease in the micronuclei frequency and chromosomal aberrations, which correlated with the clinical improvement
18.	Hua Y et al., 2006	<i>In vitro</i> experimental trial			Tea polyphenols	The human telomerase reverse transcriptase (hTERT) gene in the Tca8113 cancerous cell line was less (0.1 g/l, TP 0.05 g/l) when compared to controls (0.32 ± 0.05, 0.41 ± 0.04 and 0.72 ± 0.05, respectively) (P < 0.05)
19.	Ko SY et al., 2007	<i>In vivo</i> experimental trial—hamsters			Green tea	Amyloid precursor protein (APP) expression was also significantly increased in MBN-induced HBP carcinomas but was significantly reduced by tea intake (P < 0.0001)
20.	Tsao AS et al., 2009	Randomized control human trial—12 weeks			Green tea extracts at 500, 750, or 1000 mg/m ² or placebo thrice daily	The OPL clinical response rate was higher in all GTE arms (n = 28; 50%) versus placebo (n = 11; 18.2%; P = 0.09) However, the two higher-dose GTE arms [58.8% (750 and 1000 mg/m ²), 36.4% (500 mg/m ²), and 18.2% (placebo); P = 0.03] had higher responses, improved histology (21.4% versus 9.1%; P = 0.65)

Table 1.
Experimental studies for tea and oral cancer.

Sl no.	Author	Type of study	Risk	Sample	Country	Outcome
1.	Bundgaard T, et al., 1995	Case control study	OR = 0.45	561 Cases—161 Controls—400	Denmark	Squamous cell carcinoma occurrence
2.	Ariyawardana A et al., 2007	Cross sectional		12,716	Srilanka	46.1 per 1000 for leukoplakia and 16.4 per 1000 for oral submucous fibrosis
3.	Ide R et al., 2007	Longitudinal study—10.3 years	HR—0.51 0.60 0.31	50,221	Japan	37 oral cancer cases. Did not suggest a prominent inverse association of green tea consumption with oral cancer, although there was a tendency for a reduced risk in women

Table 2.
Descriptive studies of tea and oral cancer.

apoptosis (programmed cell death) in many types of tumor cells, including oral cancer cells. However, how the normal cells escape the apoptotic effect has not still been understood by the researchers. The effect of extracts and polyphenols of green tea as well as (–)-epigallocatechin-3-gallate (EGCG) which is the most potent green tea polyphenol on normal human keratinocytes and oral carcinoma cells were assessed through assays for cell growth, invasion, combined with apoptosis. It was shown that the green tea and its constituents selectively induce apoptosis, whereas EGCG usually inhibits the growth and invasion of oral carcinoma cells. This difference in the identification of normal cells and malignant cells by green tea and its constituents was attributed to the induction of p57, a cell cycle regulator.

Sl no.	Author	Type of study	Input	Outcome
1.	Weisburger JH, Chung FL, 2002	Review	Tea	Chemo preventive effects of tea and mechanisms
2.	Chung FL, Schwartz J, Herzog CR, Yang YM, 2003	Review	Tea	Cancer
3.	Lodi G et al., 2004	Review	Tea	Leukoplakia
4.	Wu CD, Wei GX, 2007	Review	Tea	Reduced oral cancer
5.	Klass CM, Shin DM	Review	Green tea	Premalignant lesions
6.	Boehm K et al., 2009	Systematic review	Tea	Reduced oral cancer
7.	Yang CS et al., 2008	Review	Tea and tea polyphenols	Reduced carcinogenesis

Table 3.
Review.

Figure 1.
The effect of tea on cancer.

p57 mediated survival pathway in normal epithelial cells is the reason for the chemopreventive effects of green tea polyphenols in normal cells, while oral carcinoma cells undergo an apoptotic pathway. Regular consumption of green tea has favorable effects in the prevention of oral cancer. The oxidative stress and inflammation in the oral cavity may be reduced in the presence of green tea polyphenols (**Figure 1**). Green tea prevents the transformation of healthy cells to malignant cells and locally facilitates the induction of apoptosis in oral cancer cells.

3. Conclusions: tea—a multi edged sword

Tea is rich in the beneficial polyphenols and similar components that can supplement the recommended 5–10 vegetables and fruits per day. Tea is not a drug. It is a health food, akin to *Rasayanas* known to the ancient Indians. It has shown its effect on various diseases and oral diseases.

The above review also leads us to conclude that this popular beverage can regress tumors directly by inhibiting tumor angiogenesis, blocking metastasis and inducing apoptosis in cancer cells. Thus, tea, previously considered only as a popular beverage, can now emerge as a ‘multiedged sword’ against the various diseases.

Although there has been a substantial amount of research related to the study of teas and their health benefits, it has been difficult to compare data between laboratories due to the lack of standardization in experimental procedures. Teas used in studies often differed in their types, sources, method of manufacture and procedures for extraction. Analytical data of tea preparations were often not specified or provided, making the comparison of *in vitro* or *in vivo* data difficult among laboratories. Improvement in this aspect and encouragement in designing new multidisciplinary research approaches will strengthen our knowledge concerning this ancient beverage with its many health attributes.

At present, the use of tea in clinical application is still a long way from reality, and further controlled clinical trials in humans are warranted. Furthermore, consumption of tea may have added oral health benefits by controlling ‘through prevention’ the most prevalent infectious disease of humankind, namely caries considering the Indian scenario. With the added dental health implication among many other bioregulatory functions, tea can be considered as a functional food for oral health.

Acknowledgements

I acknowledge the various resources from which I have gathered the information.

Conflict of interest

No conflict of interest.

Author details

Aswini Y. Balappanavar
Department of Public Health Dentistry, Maulana Azad Institute of Dental Sciences,
New Delhi, India

*Address all correspondence to: phdaswini@gmail.com

IntechOpen

© 2020 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] Shukla Y. Tea and cancer chemoprevention. A comprehensive review. *Asian Pacific Journal of Cancer Prevention*. 2007;**8**:155-166
- [2] Hara Y. *Green Tea: Health Benefits and Applications*. 1st ed. Marcel Dekker, Inc. New York, Basil: CRC Press; 2001. 167 p
- [3] Quick Facts: Global Tea Consumption [Internet]. 2018. Available from: <https://blog.euromonitor.com/2017/08/quick-facts-global-tea-consumption.html> [Accessed: 27 July 2018]
- [4] Tong L. *Chinese Tea*. 1st ed. Beijing: China Intercontinental Press; 2005. 137 p. ISBN 7-5085-0835-1
- [5] Sheiham A. Oral health, general health and quality of life. *Bulletin of World Health Organization*. 2009;**83**(9):641-720
- [6] Daly B, Newton T, Batchelor P, Jones K. Oral health care needs and oral health-related quality of life (OHIP-14) in homeless people. *Community Dentistry and Oral Epidemiology*. 2010;**38**(2):136-144
- [7] Bannerji B. Tea: Its pharmacological significance. In: Bannerji B, editor. *Tea Research Global Perspective*. 1st ed. Calcutta, India: Tea Research Association India; 1990. pp. 212-214
- [8] Sava VM, Galkin VM, Hong Meng Y, Yan P, Huang GWS, Hong MY, et al. A novel melanin-like pigment derived from black tea leaves with immunomodulator activity. *Food Research International*. 2001;**34**:337-343
- [9] Featherstone JD. Dental caries: A dynamic disease process. *Australian Dental Journal*. 2008;**53**(3):286-291. DOI: 10.1111/j.1834-7819.2008.00064.x
- [10] Gershon-Cohen J. Fluoride in tea and caries in rats. *Nature*. 1954;**173**:304-365
- [11] Elvin-Lewis M, Steelman R. The anticariogenic effects of tea drinking among Dallas children. *Journal of Dental Research*. 1968;**65**:198
- [12] Rosen S, Elvin-Lewis M, Beck FM, Beck EX. Anticariogenic effects of tea in rats. *Journal of Dental Research*. 1984;**63**:658-660
- [13] Smullen J, Koutsou GA, Foster HA, Zumbé A, Storey DM. The antibacterial activity of plant extracts containing polyphenols against *Streptococcus mutans*. *Caries Research*. 2007;**41**(5):342-349
- [14] Touyz LZ, Amsel R. Anticariogenic effects of black tea (*Camellia sinensis*) in caries prone-rats. *Quintessence International*. 2001;**32**(8):647-650
- [15] Kashket S, Paolino VJ, Lewis DA, van Houte J. In-vitro inhibition of glucosyltransferase from the dental plaque bacterium *Streptococcus mutans* by common beverages and food extracts. *Archives of Oral Biology*. 1985;**30**:821-826
- [16] Hattori M, Kusumoto IT, Namba T, Ishigami T, Hara Y. Effect of tea polyphenols on glucan synthesis by glucosyltransferase from *Streptococcus mutans*. *Chemical Pharmaceutical Bulletin (Tokyo)*. 1990;**38**:717-720
- [17] Otake S, Makimura M, Kuroki T, Nishihara Y, Hirasawa M. Anticaries effects of polyphenolic compounds from Japanese green tea. *Caries Research*. 1991;**25**:438-443
- [18] Nakahara K, Kawabata S, Ono H, Ogura K, Tanaka T, Ooshima T, et al. Inhibitory effect of oolong tea polyphenols on glycosyltransferases of mutans streptococci. *Applied and Environmental Microbiology*. 1993;**59**:968-973

- [19] Ooshima T, Minami T, Aono W, Izumitani A, Sobue S, Fujiwara T, et al. Oolong tea polyphenols inhibit experimental dental caries in SPF rats infected with mutans streptococci. *Caries Research*. 1993;**27**:124-129
- [20] Bhalla P, Srikantha R, Grigsby W, Wu-Yuan CD. Effect of green tea extracts on regrowth and metabolism of human plaque. *Journal of Dental Research*. 1997;**76**:267
- [21] Hamilton-Miller JM. Anti-cariogenic properties of tea (*Camellia sinensis*). *Journal of Medical Microbiology*. 2001;**50**(4):299-302
- [22] Wu CD, Wefel JS, Lingstrom P. Anticariogenic potential of black tea. In: *Proceedings, American Society for Microbiology 101 Annual Meeting*; Orlando, Florida. 2001. 327 p
- [23] Sakanaka S, Aizawa M, Kim M, Yamamoto T. Inhibitory effects of green tea polyphenols on growth and cellular adherence of an oral bacterium, *Porphyromonas gingivalis*. *Bioscience, Biotechnology and Biochemistry*. 1996;**60**:745-749
- [24] Ooshima T, Minami T, Matsumoto M, Fujiwara T, Sobue S, Hamada S. Comparison of the cariostatic effects between regimens to administer oolong tea polyphenols in SPF rats. *Caries Research*. 1998;**32**:75-80
- [25] Sakanaka S, Kim M, Taniguchi M, Yamamoto T. Antibacterial substance in Japanese green tea extract against *Streptococcus mutans*, a cariogenic bacterium. *Agricultural and Biological Chemistry*. 1989;**53**:2307-2311
- [26] You SQ. Study on feasibility of Chinese green tea polyphenols (CTP) for preventing dental caries. *Chung-Hua Kou Chiang i Hsueh Tsa Chih Chinese Journal of Stomatology*. 1993;**28**:197-254
- [27] Abid A. Oral health in Tunisia. *International Dental Journal*. 2004;**54**(6 Suppl. 1):389-394
- [28] Zhang J, Kashet S. Inhibition of salivary amylase by black and green teas and their effects on the intraoral hydrolysis of starch. *Caries Research*. 1998;**32**:233-238
- [29] Wei GX, Wu CD, Nickelsen J, Wefel JS, Lingstrom P. Effects of black tea extract on oral pathogens. *Journal of Dental Research*. 1999;**78**:414
- [30] Sarkar S, Sett P, Chowdhury T, Ganguly DK. Effect of black tea on teeth. *Journal of the Indian Society of Pedodontics and Preventive Dentistry*. 2000;**18**:139-140
- [31] Yu H, Oho T, Xu LX. Effects of several tea components on acid resistance of human tooth enamel. *Journal of Dentistry*. 1995;**13**:s101-s105
- [32] Wei SHY, Hattab FN, Mellberg JR. Concentration of fluoride and selected other elements in teas. *Nutrition*. 1989;**5**:237-240
- [33] Kavanagh D, Renehan J. Fluoride in tea and its dental significance: A review. *Journal of the Irish Dental Association*. 1998;**44**:100-105
- [34] Khan SA, Kong EF, Meiller TF, Jabra-Rizk MA. Periodontal diseases: Bug induced, host promoted. *PLoS Pathogens*. 2015;**11**(7):e1004952
- [35] Hiasawa M, Takada K, Makimura M, Otake S. Improvement of periodontal status by green tea catechin using a local delivery system: A clinical pilot study. *Journal of Periodontal Research*. 2002;**37**(6):433-438
- [36] Osawa K, Matsumoto T, Yasuda H, Kato T, Naito Y, Okuda K. The inhibitory effect of plant extracts on the collagenolytic activity and cytotoxicity of human gingival fibroblasts by *Porphyromonas gingivalis* crude enzyme. *Bulletin of the Tokyo Dental College*. 1991;**32**:1-7

- [37] Demeule M, Brossard M, Page M, Gingras D, Beliveau R. Matrix metalloproteinase inhibition by green tea catechins. *Biochimica et Biophysica Acta*. 2000;**1478**(1):51-60
- [38] Maeda-Yamamoto M, Suzuki N, Sawai Y, Miyase T, Sano M, Hashimoto-Ohta A, et al. Association of suppression of extracellular signal-regulated kinase phosphorylation by epigallocatechin gallate with the reduction of matrix metalloproteinase activities in human fibrosarcoma HT1080 cells. *Journal of Agricultural and Food Chemistry*. 2003;**51**:1858-1863
- [39] Okamoto M, Sugimoto A, Leung KP, Nakayama K, Maeda N. Inhibitory effect of green tea catechins on cysteine proteinases in *Porphyromonas gingivalis*. *Oral Microbiology and Immunology*. 2004;**19**:118-120
- [40] Sakanaka S, Okada Y. Inhibitory effects of green tea polyphenols on the production of a virulence factor of the periodontal-disease-causing anaerobic bacterium *Porphyromonas gingivalis*. *Journal of Agricultural and Food Chemistry*. 2004;**52**:1688-1692
- [41] Okamoto M, Leung KP, Ansai T, Sugimoto A, Maeda N. Inhibitory effects of green tea catechins on protein tyrosine phosphatase in *Prevotella intermedia*. *Oral Microbiology and Immunology*. 2003;**18**:192-195
- [42] Zhu M, Wei GX, Wu CD. Effect of tea polyphenols on growth and H₂S production of halitosis causing bacteria. In: Presented at American Society for Microbiology 103rd General Meeting; Washington, DC. 2003
- [43] Schwartz JL, Baker V, Larios E, Chung FL. Molecular and cellular effects of green tea on oral cells of smokers: A pilot study. *Molecular Nutrition Food Research*. 2005;**49**:43-51
- [44] Krahwinkel T, Willershausen B. The effect of sugar-free green tea chew candies on the degree of inflammation of the gingiva. *European Journal of Medical Research*. 2000;**5**:463-467
- [45] Rivera C. Essentials of oral cancer. *International Journal of Clinical and Experimental Pathology*. 2015;**8**(9):11884-11894
- [46] Bundgaard T, Wildt J, Frydenberg M, Elbrønd O. Case-control study of squamous cell cancer of the oral cavity in Denmark. *Cancer Causes & Control*. 1995;**6**(1):57-67
- [47] Ariyawardana A, Sitheequ MA, Ranasinghe AW, Perera I, Tilakaratne WM, Amaratunga EA, et al. Prevalence of oral cancer and pre-cancer and associated risk factors among tea estate workers in the central Sri Lanka. *Journal of Oral Pathology & Medicine*. 2007;**36**(10):581-587
- [48] Ide R, Fujino Y, Hoshiyama Y, Mizoue T, Kubo T, Pham TM, et al. A prospective study of green tea consumption and oral cancer incidence in Japan. *Annals of Epidemiology*. 2007;**17**(10):821-826 Epub 2007 Jul 2
- [49] Lodi G, Sardella A, Bez C, Demarosi F, Carrassi A. Interventions for treating oral leukoplakia. *Cochrane Database of Systematic Reviews*. 2004;**3**:CD001829
- [50] Wu CD, Wei GX. Tea as a functional food for oral health. *Nutrition*. 2002;**18**(5):443-444
- [51] Green tea and leukoplakia. The Indian-US Head and Neck Cancer Cooperative Group. *American Journal of Surgery*. 1997;**174**(5):552-555
- [52] Khafif A, Schantz SP, Chou TC, Edelstein D, Sacks PG. Quantitation of chemopreventive synergism between (–)-epigallocatechin-3-gallate and in normal, premalignant and

malignant human oral epithelial cells. Carcinogenesis. 1998;**19**(3):419-424

[53] Khafif A, Schantz SP, Al-Rawi M, Edelstein D, Sacks PG. Green tea regulates cell cycle progression in oral leukoplakia. Head & Neck. 1998;**20**(6):528-534

[54] Yang CS, Lee MJ, Chen L. Human salivary tea catechin levels and catechin esterase activities: Implication in human cancer prevention studies. Cancer Epidemiology, Biomarkers & Prevention. 1999;**8**(1):83-89

[55] Li N, Sun Z, Han C, Chen J. The chemopreventive effects of tea on human oral precancerous mucosa lesions. Proceedings of the Society for Experimental Biology and Medicine. 1999;**220**(4):218-224

[56] Li N, Han C, Chen J. Tea preparations protect against DMBA-induced oral carcinogenesis in hamsters. Nutrition and Cancer. 1999;**35**(1):73-79

[57] Li N, Sun Z, Liu Z, Han C. Study on the preventive effect of tea on DNA damage of the buccal mucosa cells in oral leukoplakias induce by cigarette smoking. Wei Sheng Yan Jiu. 1998;**27**(3):173-174

[58] Masuda M, Suzui M, Weinstein IB. Effects of epigallocatechin-3-gallate on growth, epidermal growth factor receptor signaling pathways, gene expression, and chemosensitivity in human head and neck squamous cell carcinoma cell lines. Clinical Cancer Research. 2001;**7**(12):4220-4229

[59] Hsu SD, Singh BB, Lewis JB, Borke JL, Dickinson DP, Drake L, et al. Chemoprevention of oral cancer by green tea. General Dentistry. 2002;**50**(2):140-146

[60] Weisburger JH, Chung FL. Mechanisms of chronic disease

causation by nutritional factors and tobacco products and their prevention by tea polyphenols. Food and Chemical Toxicology. 2002;**40**(8):1145-1154

[61] Li N, Chen X, Han C, Chen J. Chemopreventive effect of tea and curcumin on DMBA-induced oral carcinogenesis in hamsters. Wei Sheng Yan Jiu. 2002;**31**(5):354-357

[62] Masuda M, Suzui M, Lim JT, Weinstein IB. Epigallocatechin-3-gallate inhibits activation of HER-2/neu and downstream signaling pathways in human head and neck and breast carcinoma cells. Clinical Cancer Research. 2003;**15**(9):3486-3491

[63] Chung FL, Schwartz J, Herzog CR, Yang YM. Tea and cancer prevention: Studies in animals and humans. The Journal of Nutrition. 2003;**133**(10):3268S-3274S

[64] Lee MJ, Lambert JD, Prabhu S, Meng X, Lu H, Maliakal P, et al. Delivery of tea polyphenols to the oral cavity by green tea leaves and black tea extract. Cancer Epidemiology, Biomarkers & Prevention. 2004;**13**(1):132-137

[65] Srinivasan P, Sabitha KE, Shyamaladevi CS. Therapeutic efficacy of green tea polyphenols on cellular thiols in 4-nitroquinoline 1-oxide-induced oral carcinogenesis. Chemico-Biological Interactions. 2004;**149**(2-3):81-87

[66] Babich H, Krupka ME, Nissim HA, Zuckerbraun HL. Differential in vitro cytotoxicity of (–)-epicatechin gallate (EPICATECHIN GALLATE (ECG)) to cancer and normal cells from the human oral cavity. Toxicology In Vitro. 2005;**19**(2):231-242

[67] Gonzalez de Mejia E, Song YS, Ramirez-Mares MV, Kobayashi H. Effect of yerba mate (*Ilex paraguariensis*) tea

on topoisomerase inhibition and oral carcinoma cell proliferation. *Journal of Agricultural and Food Chemistry*. 2005;53(6):1966-1973

[68] Hsu S et al. Role of p21WAF1 in green tea polyphenol-induced growth arrest and apoptosis of oral carcinoma cells. *Anticancer Research*. 2005;25(1A):63-67

[69] Halder A, Raychowdhury R, Ghosh A, De M. Black tea (*Camellia sinensis*) as a chemopreventive agent in oral precancerous lesions. *Journal of Environmental Pathology, Toxicology and Oncology*. 2005;24(2):141-144

[70] Hua Y, Jianhua L, Qiuliang W, Jun F, Zhi C. Effects of tea polyphenols on telomerase activity of a tongue cancer cell line: A preliminary study. *International Journal of Oral and Maxillofacial Surgery*. 2006;35(4):352-355. Epub 2005 Nov 8

[71] Ko SY, Chang KW, Lin SC, Hsu HC, Liu TY. The repressive effect of green tea ingredients on amyloid precursor protein (APP) expression in oral carcinoma cells in vitro and in vivo. *Cancer Letters*. 2007;245(1-2):81-89. Epub 2006 Feb 3

[72] Klass CM, Shin DM. Current status and future perspectives of chemoprevention in head and neck cancer. *Current Cancer Drug Targets*. 2007;7(7):623-632

[73] Yang CS, Ju J, Lu G, Xiao H, Hao X, Sang S, et al. Cancer prevention by tea and tea polyphenols. *Asia Pacific Journal of Clinical Nutrition*. 2008;17(Suppl 1):245-248

[74] Tsao AS, Liu D, Martin J, Tang XM, Lee JJ, El-Naggar AK, et al. Phase II randomized, placebo-controlled trial of green tea extract in patients with high-risk oral premalignant lesions. *Cancer Prevention Research (Philadelphia, Pa.)*. 2009;2(11):931-941

[75] Boehm K, Borrelli F, Ernst E, Habacher G, Hung SK, Milazzo S, et al. Green tea (*Camellia sinensis*) for the prevention of cancer. *Cochrane Database of Systematic Reviews*. 2009;3:CD005004