We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Radiation Protective and Immunopotentiating Effect of Lymphocytes by β -Glucan

---

Yeun-Hwa Gu

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.80496>

Abstract

Various intractable diseases have been cured by the development of many new medicines. However, cancer is still a major cause of death. We focused on β -glucan which heat treated *Enterococcus faecalis* in human intestine, and investigated antioxidant activity, antitumor activity, immune response, etc. In this study, we studied the immunostimulatory effect of β -glucan and its application as a radioprotective agent. As a result, β -D-glucan is known to possess antioxidant activity. Its action is thought to be due to radical scavenger reaction. In our study, β -glucan increased the amount of killer cells and suppressor T cells. Therefore, administration of β -glucan is thought to have an anticancer action. In the future, we hypothesize that β -glucan could be used as an immunopotentiating agent which is effective for the prevention of cancer. From now on, further research will be necessary on the search for cancer cell receptors and to conduct fundamental research on the synergistic effects of radiation therapy. β -glucan showed strong antitumor activity against two kinds of solid carcinoma. Taken altogether, this strongly suggests that β -glucan enhances innate functions of macrophages and NK cells, and as a result, secondarily enhances the immune reaction and suppresses tumor growth.

Keywords: radioprotection, β -glucan, antitumor activity, immunoresponse, IFN- γ , NK cells, TNF- α , lymphocytes

---

1. Introduction

Previously, various intractable diseases have been cured by the development of many new or alternative medicines approaches. Although there is considerable progress in the development of anticancer drugs, cancer is still the leading cause of death in developed countries. Carcinogenesis induced by various environmental factors has different mechanisms.

Depending on the type of cancer, the therapeutic effect varies and individual differences are also variably large. Therefore, early detection, early treatment and early treatment are urgent [1].

The primary cause of death in Japan is malignant tumors. The incidence of malignant tumors is increasing year by year and one out of three patients die from malignant tumors [2]. Japanese longevity is considered a lifestyle such as ingestion of a lot of fish, ingestion of enzyme foodstuffs and diligence [3, 4]. In any case, malignant tumors have to be destroyed and, at present, there are three main forms of treatment that directly kill the lesion or malignant cells [5].

These three major forms of therapy have severe mental and physical burdens as well as the side effects and weakening of the immune system are also serious problems. In addition, there is no possibility of cessation of treatment, and the motivation for life is also considerably reduced [6–9].

Therefore, research on treatments that exclude the burden on the body as much as possible, such as surgery using a robot and immunotherapy, is underway in recent cancer treatment methods [10, 11].

At the same time, research has also been made to improve disease from the root cause from the viewpoint of oriental medicine. However, it seems that the three major therapies will continue to establish immovable status in the treatment of malignant tumors, and side effects caused by them as well as the reduction of immunity are main problems [3, 12–14].

Radiotherapy is especially important because it is a noninvasive and function-preserving method. And, if the side effects on normal tissues related to treatment can be overcome, it would be one of the most important methods in the future. When hematopoietic tissue is affected by radiation, bacterial and viral infections due to immune system weakness, genetic changes due to cell necrosis and DNA damage, mutations, and cancer occur as side effects [15–17].

For that reason, various radioprotective agents have also been studied. In fact, a radiation protection agent called WR-2721 was developed at the Walter Reed Army Hospital in the United States and clinical trials were conducted, but it is no longer commonly used due to side effects such as vomiting and rash [18–20].

Therefore, it is urgent to develop a radiation protection agent with fewer side effects, and approaches focusing on natural substances, which are said to have relatively few side effects.

In recent years, approaches for reducing the side effects of anticancer drugs with natural substances in Japan and Korea are prosperous, among which propolis is drawing attention as a natural anticancer agent and an immunopotentiating agent [19, 21].

Various medicinal ingredients and physiological effects have been reported on this substance. As components thereof, carbohydrates, amino acids, minerals, terpenoids, amino acids and the like are contained and components contained therein are changed by the extraction method.

In the case of general alcohol extraction, numerous lipophilic components such as flavonoids, which are said to be effective ingredients, are extracted, but in the case of the water extraction method devised by Suzuki et al. the levels of flavonoids decrease. However, there are reports that water extraction has higher antitumor effect than alcohol extraction. Antitumor effects are reported for β -(1-3) D glucan, β -(1-6) D glucan [22, 23].

Based on the above information, in this study, we examined the protective effect of β -glucan on lymphocytes exposed to radiation as well as its ability in reducing side effects caused by radiation therapy [24–27]. In case of radiation therapy, it is applied when the radiation sensitivity is high under the condition that the cancer condition is local.

We also examined the effect on tumor growth when combined with radiation therapy. Therefore, this study aims to propose β -glucan as the first step as a new radioprotective agent that can be used at the time of radiotherapy and to contribute data for future development.

2. Materials and methods

2.1. Animals

Mice were purchased from a Japanese SLC company and adjusted to the experimental conditions of this study for 1 week prior to the start of the experiment. The mouse used in this study is a male C3H/HeJ mouse. In order to synchronize the purchased mouse, the light was adjusted (12 L:12 D) and kept. Food and water were allowed *ad libitum*.

2.2. Test material

Enterococcus Faecalis 2001®, a glucan product, composed of yeast extract, dextrin and gelatin was supplied by Nihon BRM Co. Ltd. (Tokyo, Japan) and β -1,3 glucan as an active ingredient was contained at the ratio of 6.5 mg/g of the product. *Enterococcus Faecalis* 2001® is hereafter described as β -glucan throughout this paper. Glucan was suspended in physiological saline at concentrations of 2, 4 and 8% (w/v).

2.3. Radio-protective effect

Each mouse was intraperitoneally injected with β -glucan at a dose of 200, 400, 2800 mg/kg/day at 1 day intervals for 2 weeks. The sham control group of mice received an equal volume of normal saline. After the final injection, the mice were irradiated with 2 Gy of X-ray. Whole body irradiation was performed with a dose of 8 Gy (dose rate 1.12 Gy/min) using an X-ray irradiation device (MG226/4.5, Phillips, Inc., Tokyo). Body weight and number of surviving animals were monitored daily.

2.4. Antitumor effect

SSC-7 carcinoma cells were inoculated subcutaneously (5×10^5) into the right femur of mice.

After the average long diameter of the tumors reached 5 mm, mice were intraperitoneally injected with β -glucan suspended in physiological saline at a dose of 200, 400 and 800 mg/kg/day for 5 consecutive days. Control mice received an equal volume of normal saline. At the given time point, the major and minor axes of the tumor were measured with a caliper and the tumor volume was calculated by the following equation; $V = (\pi/6) ab^2$, a: major axis length, b: minor axis length.

2.5. Leukocyte and lymphocyte counts

The β -glucan suspended in physiological saline was intraperitoneally injected into mice at doses of 200, 400, 800 mg/kg. Control mice received an equal volume of normal saline. After the injection, in order to measure the number of white blood cells and lymphocytes using an automated hemocytometer (Celltac- α , MEK-6318, Nihon kouden Co. Ltd., Tokyo), a blood sample was transferred from the tail vein into a heparinized tube.

2.6. NK cell activity

The β -glucan suspended in physiological saline was intraperitoneally injected at a dose of 400 and 800 mg/kg for 2 weeks at intervals of 1 day. Vehicle control mice received an equal volume of normal saline. Spleen cells were prepared to measure NK cell mediated cytotoxicity by ^{51}Cr -release from labeled AC-1 cells (2×10^4 cells) after the final injection. Briefly, ^{51}Cr labeled YAC-1 cells (2×10^4 cells) were added to various dilutions of splenocyte suspensions in flat bottom microplates. The mixture was incubated for 4 h at 37°C in a CO_2 -incubator. The radioactivity released to the supernatant was counted with a γ -counter and the magnitude of cell lysis calculated based on the average radioactivity of the control group was defined as NK cell activity.

2.7. Statistical analysis

Significance of the difference in each parameter among groups was assessed by Tukey's multiple comparison tests following analysis of variance. Values of $p < 0.05$ were considered significant.

3. Results

3.1. Radio-protective effect and leukocyte and lymphocyte counts

The survival of irradiated mice is summarized in **Figures 1–5**. All of the animals in the irradiated control group died from the 5th day to the 7th day following irradiation. Glucan injected intraperitoneally prolonged the survival of mice 80% on the 7th day at the doses of 200 mg/kg.

The number of leukocytes increased with time at least up to 24 h after each repeated dose of glucan in a dose-dependent manner. The lymphocyte counts also showed a similar tendency to increase as in the leukocyte counts.

3.2. Antitumor effect

Figure 6 shows anticancer effect of various β -glucan concentrations. The tumor of the control group grew with time, whereas glucan injected intraperitoneally made tumor growth delay significantly in a dose-dependent manner **Figure 3**.

3.3. NK activity

Figure 7 shows NK activity of various β -glucan concentrations. NK cell activities in mice are shown in **Figure 7**. The NK activity increased significantly about twofold to threefold after each repeated dose of glucan (400 and 800 mg/kg).

3.4. Discussion

In the present study, we have found that β -glucan showed antioxidant activity. Previously, Ekaterini et al. have reported that β -glucan shows antioxidant activity [28], suggesting that β -glucan may be responsible for the activity that we found in our study. β -Glucan is well known for its radioprotective and antitumor effect *in vivo* [29–32], and these effects were reproduced in this study.

Figure 1. Leukocyte counts on different days after irradiation in mice of different groups. The number of leukocyte was calculated from the pre-irradiation values taken as 100%. The bars represent standard deviation. *Statistically significant ($p < 0.05$) from the control group.

Figure 2. Lymphocyte counts on different days after irradiation in mice of different groups. The number of lymphocyte was calculated from the pre-irradiation values taken as 100%. The bars represent standard deviation. *Statistically significant ($p < 0.05$) from the control group.

Figure 3. The increased percentage of CD4+, CD8+ and CD16+ T-lymphocytes in PBLs compared to the experimental data baselines of the groups. The unit is in percentage (%). Significantly different from * $p < 0.05$ control group vs. β -glucan groups by Dunnett's test.

Figure 4. Survival after irradiation. Surviving fraction was increased after injection of β -glucan.

Figure 5. Radiation protection effect of mucosal damage of small intestine to β -glucan of both doses of 12 and 24 mg/kg concentration.

Figure 6. Effect of β -glucan on the tumor growth in mice inoculated with SSC-7 carcinoma cells. Groups of 10 mice were subjected to different treatment. Results represent means $\pm$ S.D. *Statistically significant ($p < 0.05$) from the control group.

Figure 7. Repeated dose effect of β -glucan on the NK activity in mice. Groups of 10 mice each were subjected to each treatment. Results represent means $\pm$ S.D. *Statistically significant ($p < 0.05$) from the control group. **Statistically significant ($p < 0.01$) from the control group.

To confirm the elucidation mechanism by which β -glucan exerts these effects, the number of white blood cells and lymphocytes was monitored as a hematopoietic effect. Furthermore, NK cell activity was measured as an immunological parameter. The results of these parameters demonstrated that the radioprotective effect of β -glucan is mediated, at least in part, by hematopoietic effects in irradiated mice. This is because the leukocyte and lymphocyte counts

Figure 8. This slide shows a mechanism of radiation protection effect of various β -glucan concentrations.

Figure 9. This slide demonstrates a mechanism of activity of NK and LAK cells of lymphocytes by various β -glucan concentrations.

increased with a single administration of β -glucan. In addition, the enhanced immunological activity as seen by the increase in NK cell activity by β -glucan appears to play a role in the prevention of secondary infections associated with radiation.

Natural killer (NK) and lymphokine-activated killer (LAK) cells are well known to be associated with cytotoxic effects on various kinds of tumor cells [33, 34]. The reason why the immature immune response of young mice was activated to the same level as observed in adult mice is that the various components in the sample used in this experiment most probably activated IL-2 and IL-6 of the immune system. IFN- γ , cytokines and the like are multifunctional factors showing antiviral effect, inhibition of cell proliferation, antitumor effect, macrophage activation, enhancement of NK cell activation, regulation of immune response, and differentiation induction.

Specifically, high levels of IFN- γ were measured in mice bearing the S-180 carcinoma, after administration of β -glucan [35–38].

IL-4 is mainly associated with a class switch of IgM to IgG. IFN- γ and IL-4 are related to differentiation of Th1 and TH2, respectively, and these cytokines suppress each other [37–39].

When antitumor action was examined using two kinds of sarcoma (Ehrlich solid carcinoma and Sarcoma-180 carcinoma), tumor-suppressive ratios after treatment with β -glucan was 83%. When Sarcoma-180 solid carcinoma was used, tumor-suppressive ratios was 60%. Thus, β -glucan showed strong antitumor activity against two kinds of solid carcinoma [40, 41].

Therefore, there is the possibility that after treatment with β -glucan, in which no antioxidant activity is found, the balance between NF- κ B and I- κ B is lost, and the apparent tumor-suppressive ratio is low due to suppression of apoptosis of tumor cells. Thus, it is possible that β -glucan restores the balance between NF- κ B and I- κ B, and induces apoptosis and tumor suppression secondarily [42].

Figure 10. This slide shows a mechanism of activity of antitumor effect by β -glucan.

Therefore, increased activity of NK by β -glucan contributes most probably to attenuated tumor growth in tumor-bearing mice. Based on these data, glucan is expected to be a promising agent for the treatment of cancer patients receiving radiotherapy (**Figures 8–10**).

4. Conclusion

In this study we have evaluated the anticancer properties of β -glucan using a mouse model. β -glucan was found to upregulate CD8⁺ lymphocytes and leukocytes as well as reducing tumor volume and mass. We believe two phenomena may be correlated but further research is required to elucidate the exact role of β -glucan. Nonetheless, it may be possible to suggest that β -glucan could be used as a potential anticancer agent. β -Glucan was also found to improve recovery rate of lymphocytes and leukocytes following irradiation. We believe this may be due to antioxidant properties of β -glucan suggesting its possible use as an alternative radioprotective agent with reduced side effects.

Author details

Yeun-Hwa Gu

Address all correspondence to: gu.y@junshin-u.ac.jp

Graduate School of Health Science, Department of Radiological Science, Faculty of Health Science, Junshin Gakuen University, Minami-ku, Fukuoka, Japan

References

- [1] Sapi E, Alvero AB, Chen W, O'Malley D, Hao XY, Dwipoyono B, et al. Resistance of ovarian carcinoma cells to docetaxel is XIAP dependent and reversible by phenoxodiol. *Oncology Research*. 2004;**14**(11–12):567-578
- [2] Cockerham LG, Walden TL Jr, Landauer MR, Dallas CE, Mickley GA Jr. Ionizing radiation. In: Hayes AW, editor. *Principles and Methods of Toxicology*. Philadelphia, PA: Taylor and Francis; 2001. pp. 699-671
- [3] Nair CKK, Parida DK, Nomura T. Radioprotectors in radiotherapy. *Journal of Radiation Research*. 2001;**42**:21-37
- [4] Xu XR, Zhou J, Wen JS, Tao ZQ, Kang AL, Huang JX, et al. The relationship between radioprotection and *Lactobacillus*-estrogenic action of nonsteroidal estrogens and related compounds (author's transl). *Yao Xue Xue Bao*. 1980;**15**:648-655

- [5] Landauer MR, Davis HD, Kumar KS, Weiss JF. Behavioral toxicity of selected radio-protectors. *Advances in Space Research*. 1992;**12**:273-283
- [6] Nakamura T, Itokawa Y, Tajima M, Ukawa Y, Cho KH, Choi JS, et al. Radioprotective effect of *Lyophyllum decastes* and the effect on immunological functions in irradiated mice. *Journal of Traditional Chinese Medicine*. 2007;**27**(1):70-75
- [7] Power KA, Thompson LU. Ligand-induced regulation of ERalpha and ERbeta is indicative of human breast cancer cell proliferation. *Breast Cancer Research and Treatment*. 2003;**81**:209-221
- [8] Fang YC, Chen BH, Huang RF, Lu YF. Effect of *Lactobacillus* supplementation on tissue lipid peroxidation of serum, liver and low-density lipoprotein in hamsters. *The Journal of Nutritional Biochemistry*. 2004;**5**:142-148
- [9] Wei H, Zhang X, Wang Y, Lebwahl M. Inhibition of ultraviolet light-induced oxidative events in the skin and internal organs of hairless mice by isoflavone. *Cancer Letters*. 2002;**185**:21-29
- [10] Shimoi K, Masuda S, Furugori M, Esaki S, Kinae N. Radioprotective effect of antioxidative flavonoids in gamma ray irradiated mice. *Carcinogenesis*. 1994;**15**:2669-2672
- [11] Yanagisawa T, Gu YH, Tsuchihashi E, Umekawa M, Yamamoto H, Iwasa T, et al. Analgesic and anti-neoplastic effects of immunization-active fraction of *Enterococcus faecalis*. *Journal of Oriental Medicine*. 2001;**5**(1):97-102
- [12] Blackett NM, Marsh JC, Gordon MY, Okell SF, Aguado M. Simultaneous assay by six methods of the effect on hemopoietic precursor cells of adriamycin, methyl-CCNU, ⁶⁰Co rays, vinblastine, and cytosine arabinoside. *Experimental Hematology*. 1978;**6**:2-12
- [13] Lohrmann HP, Schreml W. *Cytotoxic Drugs and the Granulopoietic System*. Berlin: Springer Verlag; 1982
- [14] Umadevi P. Normal tissue protection in cancer therapy: Progress and prospects. *Acta Oncologica*. 1998;**37**:247-251
- [15] Weiss JF. Pharmacological approaches to protection against radiation induced lethality and other damages. *Environmental Health Perspectives*. 1997;**105**:1473-1478
- [16] Kasai S, Fujimoto S, Nitta K, Baba H, Kunimoto T. Antitumor activity of polymorphonuclear leukocytes activated by a β -1,3-D-glucan. *Journal of Pharmacobio-Dynamics*. 1991;**14**:519-525
- [17] Patchen ML, MacVittie TJ. Comparative effects of soluble and particulate glucans on survival in irradiated mice. *Journal of Biological Response Modifiers*. 1986;**5**:45-60
- [18] Wools WR, DiLuzio NR. The phagocytic and proliferative response of the RES following glucan administration. *Journal of the Reticuloendothelial Society*. 1964;**1**:160-169
- [19] Diluzio NR, Pisano JC, Saba TM. Evaluation of the mechanism of glucan-induced stimulation of the reticuloendothelial system. *Journal of the Reticuloendothelial Society*. 1970;**7**: 731-742

- [20] Wooles WR, DiLuzio NR. Influence of RE hyperfunction on bone marrow transplantation. *The American Journal of Physiology*. 1962;**203**:404-408
- [21] Burgaleta C, Golde DW. Effect of glucan on granulopoiesis and macrophage genesis in mice. *Cancer Research*. 1977;**37**:1739-1742
- [22] Patchen ML, Lotzova E. Modulation of murine hemopoiesis by glucan. *Experimental Hematology*. 1980;**8**:409-422
- [23] Patchen ML, MacVittie TJ. Dose-dependent responses of murine pluripotent stem cells and myeloid and erythroid progenitor cells following administration of the immunomodulating agent glucan. *Immunopharmacology*. 1983;**5**:303-313
- [24] Vetvicka V, Thornton BP, Ross GD. Soluble beta-glucan polysaccharide binding to the lectin site of neutrophil or natural killer cell complement receptor type 3 (CD11b/CD18) generates a primed state of the receptor capable of mediating cytotoxicity of iC3b-opsonized target cells. *The Journal of Clinical Investigation*. 1996;**98**:50-61
- [25] Xia Y, Vetvicka V, Yan J, Hanikyrova M, Mayadas T, Ross GD. The β -glucan-binding lectin site of mouse CR3 (CD11b/CD18) and its function in generating a primed state of the receptor that mediates cytotoxic activation in response to iC3b-opsonized target cells. *Journal of Immunology*. 1999;**162**:2281-2290
- [26] Yan J, Vetvicka V, Xia Y, Coxon A, Carroll MC, Mayadas TN, et al. β -Glucan, a "specific" biologic response modifier that uses antibodies to target tumors for cytotoxic recognition by leukocyte complement receptor type 3 (CD11b/CD18). *Journal of Immunology*. 1999;**163**:3045-3052
- [27] Cheung NK, Modak S, Vickers A, Knuckles B. Orally administered β -glucans enhance anti-tumor effects of monoclonal antibodies. *Cancer Immunology, Immunotherapy*. 2003;**51**:557-564
- [28] Gu YH, Takagi Y, Nakamura T, Hasegawa T, Suzuki I, Oshima M, et al. Enhancement of radioprotection and anti-tumor immunity by yeast-derived beta-glucan in mice. *Journal of Medicinal Food*. 2005;**8**(2):154-158
- [29] Kang YN, Itokawa Y, Maenaka T, Yamashita T, Nakamura T, Hasegawa T, et al. Effects of *Ganoderma lucidum* on immunocompetence in mice. *Medicine of Biology*. 2006;**150**:90-96
- [30] Terai K, Ryu MS, Itokawa Y, Maenaka T, Nakamura T, Hasegawa T, et al. Damage of radioprotection and antitumor effects of water-soluble *propolis*. *Oriental Pharmacy and Experimental Medicine*. 2006;**6**:12-20
- [31] Gu YH, Oshima M, Hasegawa T. Dose dependence of the severity of radiation-induced thymic lymphoma in mice. *Mutagens & Carcinogens*. 2002;**22**:266-273
- [32] Choi IS, Itokawa Y, Maenaka T, Yamashita T, Mitsumoto N, Tano K, et al. Antioxidant activity and anti-tumor immunity by *Propolis* in mice. *Oriental Pharmacy and Experimental Medicine*. 2005;**5**:100-109

- [33] Nair CKK, Uma-dvi P, Shimanskaya R, Kunugita N, Murase H, Gu Y, et al. Water soluble vitamin E (TMG) as a radioprotector. *Indian Journal of Experimental Biology*. 2003;**41**: 1365-1371
- [34] Gu YH, Maenaka T, Saito K, Yamashita T, Choi IS, Terai K, et al. Antioxidant and immuno-enhancing effects of *Echinacea purpurea* (American herb) *in vivo*. *Oriental Pharmacy and Experimental Medicine*. 2005;**5**:48-56
- [35] Gu YH, Choi HJ, Yamashita T, Kang KM, Iwasa M, Lee MJ, et al. Pharmaceutical production of anti-tumor and immune-potentiating *Enterococcus faecalis*-2001 β -glucans: Enhanced activity of macrophage and lymphocytes in tumor-implanted mice. *Current Pharmaceutical Biotechnology*. 2017;**18**:653-661
- [36] Ukawa Y, Gu YH, Ohtsuki M, Suzuki I, Hisamatsu MJ. Antitumor effects of trehalose on salcoma 180 in ICR mice. *Journal of Applied Glycoscience*. 2005;**52**:367-368
- [37] Gu YH, Fujimiya Y, Itokwa Y, Oshima M, Choi JS, Miura M, et al. Tumoricidal effects of beta-glucans: Mechanisms include both antioxidant activity plus enhanced systemic and topical immunity. *Nutrition and Cancer*. 2008;**60**:685-691
- [38] Tsiapali E, Whaley S, Kalbfleisch J, Ensley HE, Browder IW, Williams DL. Glucans exhibit weak antioxidant activity, but stimulate macrophage free radical activity. *Free Radical Biology & Medicine*. 2001;**30**(4):393-402
- [39] Shigeo K, Kazuyo M. Role of innate lymphocytes in infection and inflammation. *Frontiers in Immunology*. 2012;**3**(101):1-13
- [40] Di J, Bi S, Zhang M. Third-generation superoxide anion sensor based on superoxide dismutase directly immobilized by sol-gel thin film on gold electrode. *Biosensors & Bioelectronics*. 2004;**19**(11):1479-1486
- [41] Steinbeck MJ, Khan AU, Karnovsky MJ. Extracellular production of singlet oxygen by stimulated macrophages quantified using 9,10-diphenylanthracene and perylene in a polystyrene film. *The Journal of Biological Chemistry*. 1993;**268**(21):15649-15654
- [42] Schneider-Stock R, Diab-Assef M, Rohrbeck A, Foltzer-Jourdainne C, Boltze C, Hartig R, et al. Gali-Muhtasib H.5-Aza-cytidine is a potent inhibitor of DNA methyltransferase 3a and induces apoptosis in HCT-116 colon cancer cells via Gadd45- and p53-dependent mechanisms. *The Journal of Pharmacology and Experimental Therapeutics*. 2005;**312**(2): 525-536