We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

A Genomics Perspective on Pig Domestication

Mirte Bosse

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.82646>

Abstract

Land animal domestication has typically led to remarkable phenotypic diversity, stemming from a broad genetic background. The process of land animal domestication turns out to be a complex, long-term event with extensive gene-flow between wild and captive populations. Using pig as model, this chapter provides an in-depth overview of domestication-related events leading towards the genetic diversity in extant pig breeds. Five events in the evolutionary history and domestication of pigs can be recognized that are important for the genetic variation in modern pig genomes: (1) Speciation of *Sus* species in Island South-East Asia (ISEA); (2) Divergence between European and Asian lineages; (3) Independent domestication leading to separate domesticated clades in Europe and Asia; (4) Hybridization between domesticated pigs from Asia and Europe; and (5) Breed formation. Remarkably, the extensive mixture of genetic material leading towards the current European commercial pigs has resulted in domestic breeds that are genetically more diverse than their wild ancestors. Nowadays, commercial breeding and genomics go hand in hand. Genomics has not only proven useful to provide understanding about the domestication history of pigs but also about the molecular mechanisms underlying traits of interest. Moreover, genomic selection is an important tool integral to modern commercial breeding.

Keywords: genomics, pig, hybridization, selection, domestication

1. The process of livestock domestication

Domestication of land animals has typically led to a wide variety of domestic forms, with remarkable phenotypic diversity not seen in the wild. However, the underlying molecular variation resulting in a specific phenotype often stems from mutations predating domestication. Although domestication generally leads towards a reduction in effective population size, land animal domestication cannot be seen as a simple split of a subset of individuals from

their wild progenitors. The meaning of the word domestication is poorly defined and lacks consistency across different scientific disciplines [1]. From a population genetics perspective, domestication results in a deliberate separation of the captive, and then domesticated population from its parent population. Domestication is, therefore, initially indistinguishable from any other event that results in reduction of gene-flow between populations, and creating opportunity to respond to new selective pressures [2]. The simplest definition of domestication considers a domestic population as a subset of the wild population with cessation of gene-flow [3]. Therefore, one can expect that domestication results in a reduction of genetic variation in the domesticated population. The onset of domestication occurred in multiple geographically distinct areas during the late Pleistocene to early Holocene transition (12,000–8200 B.P, [4]). The process of land animal domestication, however, turns out to be a complex, long-term event initiated by cultural transitions related to food production [5, 6]. The definition of an animal to be considered domesticated varies, however, some common characteristics emerge from literature. Teletchea and Fontaine propose that a domesticated animal should be selectively bred in captivity and modified from its wild ancestors [7]. It is important to realize that those early considered domestic populations were genetically and phenotypically hardly distinguishable from wild types, and therefore geographical location was a better predictor of local characteristics than domestication status [3]. The general assumption that multiple centers of domestication exist has important implications for the source of genetic and phenotypic variation in domesticated species. In cattle, for example, two distinct cattle lineages that separated ~300,000 ya, contributed to two major lineages of extant cattle, that is, taurine cattle (originating from *Bos taurus*) and indicine cattle (originating from *Bos indicus*) [8]. It is not unlikely that multiple populations of wild land animals that are now extinct contributed to the genetic diversity that is observed in modern breeds [9]. The domestic animal populations accompanying human settlements did not necessarily remain at their original location of domestication. Rather, they moved along with early farmers spreading in Asia and from Eastern Anatolia throughout Europe [10]. During this process, the connection of domestic animals and farmers was relatively loose, enabling animals to hybridize with local wild populations [11]. Only centuries later, animals were actually kept in strict enclosures and intentionally bred for specific purposes, leading towards the best-known characteristic of domestic animals: docility [12]. This controlled environment drastically reduced the opportunity of domestic herds to interbreed with local wild populations, which enabled strong divergence between domestic and wild forms. We should realize the genetic basis of the modifications leading towards morphological differences in domestic animals compared to their wild ancestors is mostly provided by standing genetic variation, that is, mutations that were already present before the onset of domestication and selection. Therefore, indicating the genetic underpinnings of domestication remain challenging [13, 14]. Arguably, we can speak about a domestic population if not only the gene pool is distinct from the wild variety, but also (artificially) selected variants leading to desired phenotypes are at high(er) frequency in the domestic population [15–18]. In this chapter, an in-depth overview is provided for the complex process of domestication, admixture, and selection leading towards the genetic diversity in extant breeds, using pig as model.

2. Genomic insight in pig domestication

Domesticated species are good models to study genomic and phenotypic consequences of demography and selection [19]. The use of higher DNA marker densities has enabled researchers to reveal the complexity of livestock domestication, which was shown to be far more complex than a single sampling from the wild [20]. Genotyping and sequencing technologies have opened up many opportunities to reveal the complex history of domestication, admixture, and selection in livestock [4, 20]. Combining modern sequence technologies with extensive studies on fossil records and land animal usage now enables the reconstruction of domestication in details. Apart from a suitable history and documentation, the availability of detailed genetic information is crucial to be able to study genomic alterations due to domestication. Pig (*Sus scrofa*, Linnaeus, 1758) was the first livestock species for which a genome consortium was established with the intention to completely map the genome [21, 22]. The design of a 60k single nucleotide polymorphism (SNP) chip for pigs in 2009 greatly contributed to the applicability of genomics techniques in pig breeding, and simultaneously increased possibilities for population genomics studies [23]. The establishment of a consortium to sequence the pig genome in 2003 and publication of the pig reference genome in 2012 opened up an even greater window of opportunities to study various aspects of the genetics of pig, since the highest resolution possible became reality [21, 24]. Together with the evolutionary history of pig, these provide an unprecedented study system to demonstrate the impact of domestication from a genomics perspective. Pig genomes contain a complex composition of segments, reflecting the different backgrounds that contributed to the domestic animal it is today. Disentangling these genomic signatures provides enormous information about the complex background and history of the worlds' most consumed meat type [25].

3. Conceptual history of the pig (*Sus scrofa*)

Here I will discuss genomic variation within and between different populations of pigs, providing deeper understanding of how domestication has influenced genetic diversity of pigs. Five major events in the evolutionary history and domestication of pigs can be recognized that are of importance for the distribution of genetic variation in modern pig genomes (**Figure 1**).

3.1. Speciation of *Sus* in island South-East Asia

Knowledge about the source of the domesticated form, the origin of the species, is essential to understand genetic variation within modern breeds. The Suidae family is particularly interesting for molecular genetic studies as it is one of the few mammalian lineages that has closely related species living today. Multiple *Sus* species originated roughly ~4 million years ago on Island Southeast Asia (ISEA). The island structure in this region probably promoted speciation, since the bearded pig *Sus barbatus*, the warty pigs *S. celebensis* and *S. verrucosus* but also wild *S. scrofa* occur on separate islands. The phylogenetic structure within the genus *Sus*

Figure 1. Schematic overview of the history of the pig (*Sus scrofa*). Five main events in pig history are indicated in blue, with the approximate timing of those events in red: (1) Speciation of *Sus* species in Island South-East Asia (ISEA). (2) Divergence between European and Asian *S. scrofa* lineages. (3) Independent domestication leading to separate domesticated clades in Europe and Asia. (4) Hybridization between domesticated pigs from Asia and Europe. (5) Breed formation.

has been studied intensively and revealed a complex history of admixture [26, 27]. The past connection of landmasses at the Sunda shelf and isolation of Indonesian islands by the rapid sea level rise after the last glaciatiion period [28] created a dynamic process of (re)colonization, isolation and admixture of different *Sus* species and populations [29, 30]. The species that gave rise to the domesticated pig, *Sus scrofa*, has its origin in Southeast Asia some ~4 Mya and colonized almost the entire Eurasian mainland from there. The widespread and opportunistic nature of this species probably contributed to the fact that *Sus scrofa* is the only pig species that was successfully domesticated [25].

3.2. Divergence between European and Asian *S. scrofa*

Sus scrofa is widespread within Eurasia (**Figure 1**) and consists of many isolated wild and domesticated populations. The divergence between Western-European and Eastern-Asian populations has been estimated at about 1.2 Mya [24, 29], and has resulted in many fixed molecular differences between the two groups. This divergence not only resulted in a European and an Asian *S. scrofa* clade, but also in differences in demographic history and population size. The last glacial maximum probably reduced population sizes of both European and Asian

wild boars, but the reduction was most severe in Europe [24]. The geographic distribution of wild boar over Europe faced another severe decline starting in the middle ages and lasted until the late eighteenth century [25]. In the mid-nineteenth century, natural or human-mediated recolonization events resulted in isolated populations expanding their range. Some of these isolated populations were small in effective size for decades or longer, causing inbreeding and population differentiation. Local re-stocking of populations with geographically distinct wild boar resulted in complex genetic structures and signatures of population dynamics [31, 32]. Such complex genetic architectures have been detected in Italian and Luxembourgian wild boars. However, these mixed genomes could have been shaped due to ancient glaciation events [33] or because of recent mixture [34]. Asian *Sus scrofa* is thought to have had a larger effective population size which, together with its proximity to the origin of the species, results in higher genetic diversity compared to the European clade [24, 35, 36]. These highly distinct groups of wild boar provided the basis of the genetic background of the later domesticated pigs.

3.3. Independent domestication leading to separate clades

The demographic and geographic history of the domesticated pig may be just as complex as that of its wild counterpart. There is compelling evidence that pig domestication events occurred at multiple locations, Eastern Anatolia and China independently, some 9000–10,000 years ago [26, 37]. Domestication has not been a single event, but rather a long period with recurrent admixture with wild populations [38]. Following initial domestication, the traits selected as well as how animals were kept, strongly differed in Europe and Asia resulting in highly different domesticated pigs between Europe and Asia. Asian pigs were kept in close proximity of humans, often integrated in their settlements. By contrast, European pigs were roaming freely in forested areas in the surroundings [39, 40]. Only during the Industrial Revolution, a more strict pig farming system was adopted and implemented to fulfill the increasing demand for pork. Because of recurrent gene-flow between wild and domestic pigs, a reduction in genetic diversity cannot be observed in domesticated pigs compared to their presumed wild counterparts [35, 38, 41]. One should realize though that European and Asian domesticated pigs have been geographically isolated for over a million years ago, because they have distinct wild origins. Therefore, they genetically resemble local wild boar more than domestic pigs from different geographic origins [24, 35]. This dichotomy also underlies the fact that European pigs and wild boar are genetically less diverse than Asian wild boar and domestic pigs.

3.4. Hybridization between domesticated pigs of different origin

It is well documented that during the Industrial Revolution in Europe, European pigs have been deliberately hybridized with Asian pigs. Urbanization in Europe increased the demand for meat such as pork, but during those times, pig farmers would still have their pigs roaming in surrounding forests. Forest cover was decreasing and a different pig production system seemed inevitable [40]. Due to this changing environment, pig breeders sought a way to improve their stock in such way that pigs had to become adapted to living in small(er) enclosures, be more prolific and gain weight more rapidly. This led to selection for traits better adapted to

the changed environment. Many of these traits were already present in Asian domestic pigs. Therefore, British farmers started crossbreeding their own pigs with these Asian pigs [40]. This introgression of Asian genetic material into European populations has long been demonstrated by genetic markers [42, 43]. Moreover, the intentional crossbreeding and consecutive artificial selection on Asia-derived traits enabled adaptive loci to emerge in the genome of European domestic pigs. Genes of Asian origin have been demonstrated to contribute to increased fertility and fatness in commercial Large White pigs [44, 45]. Very recently, hybridization between wild and domesticated pigs has been reported in Western Europe, resulting in traceable Asian genetic material in local wild boar populations in Germany [31, 32, 34].

3.5. Breed formation and globalization

Due to the worldwide consumption of pork, the species is farmed at a global scale, far exceeding its original natural distribution (IUCN). The influence and contribution of commercial pig breeds to local ecology and biodiversity is however debated [31, 32, 46]. Also, escape or intentional release of local stocks have resulted in feralization of domesticated pigs, which is now a major population in the United States, although the continent is not part of the native range of the species [47]. The domesticated pig as it is used nowadays for agricultural purposes consists of many breeds that have been separated and kept isolated for decades, which has resulted in many genetic differences between these breeds. Breed and population specific genetic studies have greatly enhanced the dissection of complex traits that are economically important. Knowing and understanding the origin and distribution of variation in (domesticated) species is important for conservation of genetic resources, such as culturally important heritage breeds [48]. Local husbandry and breeding techniques have created an enormous diversification of pig breeds. Generally, European breeds can be categorized into global commercial breeds, stemming from the White type in England, and local heritage breeds, developed locally and now often endangered [39]. It is notable that many heritage breeds genetically resemble the local wild boar more than global pig breeds, most likely because they were not improved by Asian gene-flow two centuries ago [35, 49–51]. The globalization of pig breeding and consumption has swamped local pig breeds with common commercial breeds from British heritage background, such as Large White, Landrace, Pietrain and Duroc [39]. Also, extensive admixture between breeds of different origin is known to occur, highly dependent on local breeding practices.

4. The hybrid nature of (pig) genomes

Increasing evidence showed that humans play an important role in stimulating hybridization in wild species, either unintentionally or on purpose. Human-induced hybridization can not only be a by-product of globalization as some species became widely distributed due to human mobility, but it can also be intentional such as in domesticated species [40, 52]. It is becoming apparent that many livestock species/breeds are actually a mixture of highly divergent populations with a mixed demographic history, combined in one genome. The formation of livestock breeds provides a good example of how man has influenced the

genomic architecture of a species. In cattle, for example, exchange of genetic material between different species promoted the uptake of beneficial traits from closely related species [53]. In pig, domestication does not seem to have left a clear population bottleneck, as demonstrated by the high level of genetic variation in European pigs [38]. This suggests that the majority of the genetic variation that is present in European wild boar is also present in domestic breeds, even though modern pigs are phenotypically clearly different from their wild counterparts. Moreover, the gene-flow with wild populations as well as between different domestic lineages enabled pig breeders to select for locally and globally preferred traits, using a broad genetic background [44, 45]. Remarkably, the extensive mixture of genetic material leading towards the current European commercial pigs has resulted in domestic breeds that are genetically more diverse than their wild ancestors in Europe [24, 35, 36, 41]. This counter-intuitive characteristic of commercial pigs is mainly driven by the influx of Asian genes during the Industrial Revolution [45]; local heritage breeds that do not display signs of Asian gene-flow tend to have lower genetic diversity [50, 51]. Nowadays, many breeds and definitions are used to describe the origin of (local) stock, with some being a complex mixture of Asian and European heritage, depending on the geographical region and the breeding practice of pig farmers.

5. Breeding and genomics go hand in hand

Pig farming has drastically changed since first domestication. Today's elaborate pig breeding industry has only few characteristics in common with early pig farmers, and has resulted in a highly professional large-scale pork production system, making use of latest technologies in animal breeding. Selection for particular traits not only improved due to more precise phenotyping and better defined traits such as carcass quality, growth rate and fertility [54], but also because of crossing breeds with desirable traits of different origins [45]. The use of pedigree information and large-scale tracking of animal relatedness has speeded up the improvement of pig breeds. In other livestock, especially cattle, the implementation of the use of genetic markers on top of pedigree information resulted in even more efficient selection [55]. The recent and rapid genetic progress can be achieved due to the implementation of genomic selection, in which animals are selected based on their performance predicted from their genotypes, rather than phenotypes [56]. This way, animals can be selected at an earlier stage, and predicted phenotypes for typically female traits can also be implemented using genotype information from males [57, 58].

Genomics has not only proven useful as a tool in genomic selection, but also has provided more understanding about the molecular mechanisms that underlie traits of interest. Knowledge about the link between genes and trait enables more accurate breeding [54]. Moreover, if the function of a specific gene is known, it can provide insight into the selection history of a breed. Numerous studies have successfully identified selection for genes linked to specific commercially important traits (**Table 1**). Interestingly, some of these genes under selection in European breeds have an Asian origin [59–61]. Also, genome-wide scans for detrimental variants have identified mutations in commercial populations with negative effects [62, 63].

Gene	Trait	Study
<i>KIT</i>	Coat color	Andersson and Plastow [66]
<i>KITLG</i>	Coat color	Okumura et al., [61]
<i>MC1R</i>	Coat color	Kijas et al., [67]; Fang et al., [68]
<i>EDNRB</i>	Coat color	Ai et al., [59]; Wilkinson et al., [69]
<i>IGF2</i>	Lean growth	van Laere et al., [70]
<i>RYR1</i>	Lean growth	Fujii et al., [71]
<i>PRKAG3</i>	Lean growth	Milan et al., [72]
<i>NR6A1</i>	Body size	Rubin et al., [73]
<i>PLAG1</i>	Body size	Rubin et al., [73]
<i>LCORL</i>	Body size	Rubin et al., [73]
<i>OSTN</i>	Body composition	Rubin et al., [73]
<i>CLDN1</i>	Fertility	Choi et al., [74]
<i>AHR</i>	Fertility	Bosse et al., [44]
<i>TWIST1</i>	Fatness	Choi et al., [74]
<i>LEMD3</i>	Ear morphology	Wilkinson et al., [59]

Table 1. Non-exhaustive list of genes associated with commercially important traits in pigs.

Recent work demonstrates that some variants that cause lethality in homozygous state are present at relatively high frequency in commercial pig lines [64, 65]. Knowing these recessive lethal mutations can aid in avoiding matings between carriers of such mutations within the breeding scheme. Overall, genomics has provided valuable insight into variation in pigs: what its origin is, how is it maintained, reduced and increased. This turned out to be a complex interplay of molecular processes, selection, demographic history, gene-flow and human interference. Moreover, genomics is an important tool in the pig industry nowadays and is integral to modern commercial breeding.

Author details

Mirte Bosse^{1,2*}

*Address all correspondence to: mirte.bosse@wur.nl

1 Wageningen University & Research – Animal Breeding and Genomics, Wageningen, The Netherlands

2 VU University Amsterdam – Animal Ecology, Amsterdam, The Netherlands

References

- [1] Zeder MA. The domestication of animals. *Journal of Anthropological Research*. 2012; **68**:161-190
- [2] Zeder MA. Domestication as a model system for the extended evolutionary synthesis. *Interface Focus*. 2017;**7**:20160133. DOI: 10.1098/rsfs.2016.0133
- [3] Larson G, Burger J. A population genetics view of animal domestication. *Trends in Genetics*. 2013;**29**(4):197-205
- [4] Larson G, Piperno DR, Allaby RG, Purugganan MD, Andersson L, Arroyo-Kalin M, et al. Current perspectives and the future of domestication studies. *Proceedings of the National Academy of Sciences of the United States of America*. 2014;**111**(17):6139-6146
- [5] Gepts P et al., editors. *Biodiversity in Agriculture: Domestication, Evolution, and Sustainability*. Cambridge: Cambridge Univ Press; 2012
- [6] Price D, Bar-Yosef O, editors. *The Beginnings of Agriculture: New Data, New Ideas*. *Curr Anthropol* 52 (Supp 4). Chicago: Wenner-Gren Symposium Series; Univ of Chicago Press; 2011
- [7] Teletchea F, Fontaine P. Levels of domestication in fish: Implications for the sustainable future of aquaculture. *Fish and Fisheries*. 2014;**15**:181-195
- [8] Pitt D, Sevane N, Nicolasi E, MacHugh D, Colli L, Martinez R, et al. Domestication of cattle: Two or three events? *Evolutionary Applications*. 2018;**1**-14. DOI: 10.1111/eva.12674
- [9] Scheu A, Powell A, Bollongino R, Vigne J-D, Tresset A, Çakırlar C, et al. The genetic prehistory of domesticated cattle from their origin to the spread across Europe. *BMC Genetics*. 2015;**16**:54. DOI: 10.1186/s12863-015-0203-2
- [10] Driscoll CA, Macdonald DW, O'Brien SJ. From wild animals to domestic pets, an evolutionary view of domestication. *Proceedings of the National Academy of Sciences of the United States of America*. 2009;**106**(suppl. 1):9971. DOI: 10.1073/pnas.0901586106
- [11] Zeder MA, Emshwiller E, Smith BD, Bradley DG. Documenting domestication: The intersection of genetics and archaeology. *Trends in Genetics*. 2006;**22**:139-155
- [12] Zeder MA. Central questions in the domestication of plants and animals. *Proceedings of the National Academy of Sciences of the United States of America*. 2006;**15**:105-117
- [13] Christie MR, Marine ML, French RA, Blouin MS. Genetic adaptation to captivity can occur in a single generation. *Proceedings of the National Academy of Sciences of the United States of America*. 2012;**109**:238-242
- [14] Christie MR, Marine ML, Fox SE, French RA, Blouin MS. A single generation of domestication heritably alters the expression of hundreds of genes. *Nature Communications*. 2016;**7**:10676. DOI: 10.1038/ncomms10676

- [15] Diamond J. Evolution, consequences and future of plant and animal domestication. *Nature*. 2002;**418**:700-707
- [16] Mignon-Grasteau S, Boissy A, Bouix J, et al. Genetics of adaptation and domestication in livestock. *Livestock Science*. 2005;**93**:3-14
- [17] Mirkena T, Duguma G, Haile A, Tibbo M, Okeyo AM, Wurzinger M, et al. Genetics of adaptation in domestic farm animals: A review. *Livestock Science*. 2010;**132**:1-12
- [18] Teletchea F. Domestication and genetics: What a comparison between land and aquatic species can bring. In: Pontarotti P, editor. *Evolutionary Biology. Biodiversification from Genotype to Phenotype*. Cham, Switzerland: Springer; 2015. pp. 389-402
- [19] Megens HJ, Groenen MAM. Domestication: A long-term genetic experiment domesticated species form a treasure-trove for molecular characterization of Mendelian traits by exploiting the specific genetic structure of these species in across-breed genome wide association studies. *Heredity*. 2012;**109**(1):1-3
- [20] Bruford MW, Bradley DG, Luikart G. DNA markers reveal the complexity of livestock domestication. *Nature Reviews. Genetics*. 2003;**4**(11):900-910
- [21] Schook LB, Beever JE, Rogers J, Humphray S, Archibald A, Chardon P, et al. Swine genome sequencing consortium (SGSC): A strategic roadmap for sequencing the pig genome. *Comparative and Functional Genomics*. 2005;**6**:251-255
- [22] Archibald AL et al. Pig genome sequence – Analysis and publication strategy. *BMC Genomics*. 2010;**11**:438
- [23] Ramos AM, Crooijmans RP, Affara NA, Amaral AJ, Archibald AL, Beever JE, et al. Design of a high density SNP genotyping assay in the pig using SNPs identified and characterized by next generation sequencing technology. *PLoS One*. 2009;**4**(8):e6524
- [24] Groenen MA, Archibald AL, Uenishi H, Tuggle CK, Takeuchi Y, Rothschild MF, Rogel-Gaillard C, Park C, Milan D, Megens HJ, et al. Analyses of pig genomes provide insight into porcine demography and evolution. *Nature*. 2012;**491**(7424):393-398
- [25] Groenen MAM. A decade of pig genome sequencing: A window on pig domestication and evolution. *Genetics Selection Evolution*. 2016;**48**:23. DOI: 10.1186/s12711-016-0204-2
- [26] Larson G, Dobney K, Albarella U, Fang M, Matisoo-Smith E, Robins J, et al. Worldwide phylogeography of wild boar reveals multiple centers of pig domestication. *Science*. 2005;**307**:1618-1621
- [27] Mona S, Randi E, Tommaseo-Ponzetta M. Evolutionary history of the genus *Sus* inferred from cytochrome b sequences. *Molecular Phylogenetics and Evolution*. 2007;**45**(2):757-762
- [28] Hanebuth T, Stattegger K, Grootes PM. Rapid flooding of the Sunda shelf: A late-glacial sea-level record. *Science*. 2000;**288**(5468):1033-1035
- [29] Frantz LA, Schraiber JG, Madsen O, Megens HJ, Bosse M, Paudel Y, et al. Genome sequencing reveals fine scale diversification and reticulation history during speciation in *Sus*. *Genome Biology*. 2013;**14**(9):R107

- [30] Frantz LAF, Madsen O, Megens H-J, Groenen MAM, Lohse K. Testing models of speciation from genome sequences: Divergence and asymmetric admixture in island south-east Asian *Sus* species during the Plio-Pleistocene climatic fluctuations. *Molecular Ecology*. 2014;**23**:5566-5574
- [31] Goedbloed DJ, Megens HJ, van Hooft P, Herrero-Medrano JM, Lutz W, Alexandri P, et al. Genome-wide single nucleotide polymorphism analysis reveals recent genetic introgression from domestic pigs into northwest European wild boar populations. *Molecular Ecology*. 2013;**22**(3):856-866
- [32] Goedbloed DJ, van Hooft P, Megens HJ, Langenbeck K, Lutz W, Crooijmans RPMA, et al. Reintroductions and genetic introgression from domestic pigs have shaped the genetic population structure of northwest European wild boar. *BMC Genetics*. 2013;**14**:43
- [33] Scandura M, Iacolina L, Crestanello B, Pecchioli E, Di Benedetto MF, Russo V, et al. Ancient vs. recent processes as factors shaping the genetic variation of the European wild boar: Are the effects of the last glaciation still detectable? *Molecular Ecology*. 2008;**17**(7):1745-1762
- [34] Frantz AC, Zachos FE, Kirschning J, Cellina S, Bertouille S, Mamuris Z, et al. Genetic evidence for introgression between domestic pigs and wild boars (*Sus scrofa*) in Belgium and Luxembourg: A comparative approach with multiple marker systems. *Biological Journal of the Linnean Society*. 2013;**110**(1):104-115
- [35] Bosse M, Megens HJ, Madsen O, Frantz LAF, Paudel Y, Crooijmans RPMA, et al. Untangling the hybrid nature of modern pig genomes: A mosaic derived from biogeographically distinct and highly divergent *Sus scrofa* populations. *Molecular Ecology*. 2014;**23**(16):4089-4102
- [36] Bosse M, Madsen O, Megens HJ, Frantz LA, Paudel Y, Crooijmans RP, et al. Hybrid origin of European commercial pigs examined by an in-depth haplotype analysis on chromosome 1. *Frontiers in Genetics*. 2014;**5**:442
- [37] Megens HJ, Crooijmans RPMA, Cristobal MS, Hui X, Li N, Groenen MAM. Biodiversity of pig breeds from China and Europe estimated from pooled DNA samples: Differences in microsatellite variation between two areas of domestication. *Genetics Selection Evolution*. 2008;**40**(1):103-128
- [38] Frantz LA, Schraiber JG, Madsen O, Megens HJ, Cagan A, Bosse M, et al. Evidence of long-term gene flow and selection during domestication from analyses of Eurasian wild and domestic pig genomes. *Nature Genetics*. 2015;**47**:1141-1148
- [39] Porter V. *Pigs – A Handbook to the Breeds of the World*. Mountfield, East Sussex: Helm Information; 1993
- [40] White S. From globalized pig breeds to capitalist pigs: A study in animal cultures and evolutionary history. *Environmental History*. 2011;**16**(1):94-120
- [41] Bosse M, Megens HJ, Madsen O, Paudel Y, Frantz LA, Schook LB, et al. Regions of homozygosity in the porcine genome: Consequence of demography and the recombination landscape. *PLoS Genetics*. 2012;**8**(11):e1003100

- [42] Giuffra E, Kijas JMH, Amarger V, Carlborg O, Jeon JT, Andersson L. The origin of the domestic pig: Independent domestication and subsequent introgression. *Genetics*. 2000; **154**(4):1785-1791
- [43] Amaral AJ, Ferretti L, Megens HJ, Crooijmans RPMA, Nie HS, Ramos-Onsins SE, et al. Genome-wide footprints of pig domestication and selection revealed through massive parallel sequencing of pooled DNA. *PLoS One*. 2011; **6**(4):e14782
- [44] Bosse M, Megens HJ, Frantz LAF, Madsen O, Larson G, Paudel Y, et al. Genomic analysis reveals selection for Asian genes in European pigs following human-mediated introgression. *Nature Communications*. 2014; **5**:4392
- [45] Bosse M, Lopes MS, Madsen O, Megens HJ, Crooijmans RP, Frantz LA, et al. Artificial selection on introduced Asian haplotypes shaped the genetic architecture in European commercial pigs. *Proceedings of the Biological Sciences*. 2015; **282**:pii:20152019. DOI: 10.1098/rspb.2015.2019
- [46] Hall SJG, Bradley DG. Conserving livestock breed biodiversity. *Trends in Ecology & Evolution*. 1995; **10**(7):267-270
- [47] Mayer JJ, Brisbin IL Jr. *Wild Pigs in the United States: Their History, Comparative Morphology, and Current Status*. 2nd ed. Athens, GA: The University of Georgia Press; 2008
- [48] Groeneveld LF, Lenstra JA, Eding H, Toro MA, Scherf B, Pilling D, et al. Genetic diversity in farm animals - a review. *Animal Genetics*. 2010; **41**:6-31
- [49] Herrero-Medrano JM, Megens HJ, Crooijmans RP, Abellaneda JM, Ramis G. Farm-by-farm analysis of microsatellite, mtDNA and SNP genotype data reveals inbreeding and crossbreeding as threats to the survival of a native Spanish pig breed. *Animal Genetics*. 2013; **44**(3):259-266
- [50] Herrero-Medrano JM, Megens HJ, Groenen MAM, Bosse M, Perez-Enciso M, Crooijmans RPMA. Whole-genome sequence analysis reveals differences in population management and selection of European low-input pig breeds. *BMC Genomics*. 2014; **15**:601
- [51] Herrero-Medrano JM, Megens HJ, Groenen MAM, Ramis G, Bosse M, Perez-Enciso M, et al. Conservation genomic analysis of domestic and wild pig populations from the Iberian Peninsula. *BMC Genetics*. 2013; **14**:106
- [52] Kijas JW, Lenstra JA, Hayes B, Boitard S, Porto Neto LR, et al. Genome-wide analysis of the world's sheep breeds reveals high levels of historic mixture and strong recent selection. *PLoS Biology*. 2012; **10**:e1001258
- [53] Wu D-D, Ding X-D, Wang S, Wojcik JM, Zhang Y, Tokarska M, et al. Pervasive introgression facilitated domestication and adaptation in the *Bos* species complex. *Nature Ecology & Evolution*. 2018; **2**:1139-1145
- [54] Merks J, Mathur P, Knol E. New phenotypes for new breeding goals in pigs. *Animal*. 2012; **6**(04):535-543

- [55] Doekes HP, Veerkamp RF, Bijma P, Hiemstra SJ, Windig JJ. Trends in genome-wide and region-specific genetic diversity in the Dutch-Flemish Holstein–friesian breeding program from 1986 to 2015. *Genetics, Selection, Evolution*. 2018;**50**:15. DOI: 10.1186/s12711-018-0385-y
- [56] Meuwissen T, Hayes B, Goddard M. Prediction of total genetic value using genome-wide dense marker maps. *Genetics*. 2001;**157**(4):1819-1829
- [57] Hayes BJ, Bowman PJ, Chamberlain AC, Goddard ME. Genomic selection in dairy cattle: Progress and challenges. *Journal of Dairy Science*. 2008;**92**:433-443
- [58] Hayes BJ, Visscher PM, Goddard ME. Increased accuracy of selection by using the realised relationship matrix. *Genetical Research*. 2009;**91**:47-60
- [59] Wilkinson S, Lu ZH, Megens HJ, Archibald AL, Haley C, Jackson IJ, et al. Signatures of diversifying selection in European pig breeds. *PLoS Genetics*. 2013;**9**(4):e1003453
- [60] Ojeda A, Huang LS, Ren J, Angiolillo A, Cho IC, Soto H, et al. Selection in the making: A worldwide survey of haplotypic diversity around a causative mutation in porcine IGF2. *Genetics*. 2008;**178**(3):1639-1652
- [61] Okumura N, Matsumoto T, Hamasima N, Awata T. Single nucleotide polymorphisms of the KIT and KITLG genes in pigs. *Animal Science Journal*. 2008;**79**(3):303-313
- [62] Bosse M, Megens HJ, Derks MFL, de Cara MAR, Groenen MAM. Deleterious alleles in the context of domestication, inbreeding and selection. *Evolutionary Applications*. 2018:1-12. doi.org/10.1111/eva.12691
- [63] Makino T, Rubin CJ, Carneiro M, Axelsson E, Andersson L, Webster MT. Elevated proportions of deleterious genetic variation in domestic animals and plants. *Genome Biology and Evolution*. 2018;**10**(1):276-290
- [64] Derks MFL, Megens HJ, Bosse M, Lopes MS, Harlizius B, Groenen MAM. A systematic survey to identify lethal recessive variation in highly managed pig populations. *BMC Genomics*. 2017;**18**:858
- [65] Derks MFL, Lopes MS, Bosse M, Madsen O, Dibbits B, Harlizius B, et al. Balancing selection on a recessive lethal deletion with pleiotropic effects on two neighboring genes in the porcine genome. *PLoS Genetics*. 2018;**14**(9):e1007661. DOI: 10.1371/journal.pgen.1007661
- [66] Andersson L, Plastow G. Molecular genetics of coat colour variation. In: Rothschild MF, Ruvinsky A, editors. *The Genetics of the Pig*. 2nd ed. Wallingford: CABI; 2011. pp. 38-50
- [67] Kijas JMH, Wales R, Tornsten A, Chardon P, Moller M, Andersson L. Melanocortin receptor 1 (MC1R) mutations and coat color in pigs. *Genetics*. 1998;**150**(3):1177-1185
- [68] Fang M, Larson G, Ribeiro HS, Li N, Andersson L. Contrasting mode of evolution at a coat color locus in wild and domestic pigs. *PLoS Genetics*. 2009;**5**(1):e1000341

- [69] Ai H, Fang X, Yang B, Huang Z, Chen H, Mao L, et al. Adaptation and possible ancient interspecies introgression in pigs identified by whole-genome sequencing. *Nature Genetics*. 2015;**47**:217-225
- [70] Van Laere AS, Nguyen M, Braunschweig M, Nezer C, Collette C, Moreau L, et al. A regulatory mutation in IGF2 causes a major QTL effect on muscle growth in the pig. *Nature*. 2003;**425**:832-836
- [71] Fujii J, Otsu K, Zorzato F, de Leon S, Khanna VK, Weiler JE, et al. Identification of a mutation in porcine ryanodine receptor associated with malignant hyperthermia. *Science*. 1991;**253**:448-451
- [72] Milan D, Jeon JT, Looft C, Amarger V, Robic A, Thelander M, et al. A mutation in PRKAG3 associated with excess glycogen content in pig skeletal muscle. *Science*. 2000;**288**:1248-1251
- [73] Rubin CJ, Megens HJ, Barrio AM, Maqbool K, Sayyab S, Schwochow D, et al. Strong signatures of selection in the domestic pig genome. *Proceedings of the National Academy of Sciences of the United States of America*. 2012;**109**(48):19529-19536
- [74] Choi JW, Chung WH, Lee KT, Cho ES, Lee SW, Choi BH, et al. Whole-genome resequencing analyses of five pig breeds, including Korean wild and native, and three European origin breeds. *DNA Research*. 2015;**22**:259-267