We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Brown Adipose Tissue Energy Metabolism

Yuan Lu

Abstract

The brown adipose tissue (BAT) evolved as a specialized thermogenic organ in mammals. Nutrients (i.e., fatty acids and glucose) from the intracellular storage and peripheral tissues are critical to the BAT thermogenic function. The BAT converts the chemical energy stored in nutrients to thermo energy through UCP1-mediated nonshivering thermogenesis (NST). Activated BAT contributes significantly to the whole body energy substrate homeostasis. It is now well-recognized that adult humans possess BAT with functional thermoactivity. Thus, BAT energy metabolism has a significant therapeutic potential in the management of metabolic disorders, such as obesity, insulin resistance, type 2 diabetes, and lipid abnormality in humans.

Keywords: brown adipose tissue, brown adipocyte, metabolism, fatty acid, glucose, metabolic disorders

1. Introduction

Brown adipose tissue (BAT) evolved as a specialized thermogenic organ in the modern eutherian mammals, including *Homo sapiens* [1–7]. The main function of BAT is mediating adaptive thermogenesis or nonshivering thermogenesis (NST), when mammals are challenged in the cold environment. The NST function is a critical adaptation, which helps to maintain the homeothermy in mammals and gives them the evolutionary advantage to survive in the cold habitat, and to thrive from the Arctic to the Antarctic region [3, 4]. During the cold challenge, BAT metabolizes nutrients (i.e., fatty acids and glucose) and converts the chemical energy to thermo energy through NST. In addition to its thermogenic function, activated BAT also contributes significantly to whole body energy substrate homeostasis. When activated, BAT presents physiologically significant benefit in fatty acids and glucose homeostasis as well as insulin sensitivity in mammals [8–24]. It is now well-recognized that adult humans possess BAT, which has functional thermoactivity [8–10, 25]. Cold challenge-activated BAT is detected mainly in the supraclavicular, paravertebral, and cervical regions in adult humans [9, 10, 25, 26]. Accumulating evidences indicate that BAT function is inversely associated with age, body mass index (BMI), and diabetic status in adult humans, which indicates that the activation of BAT has potential translational implication in the management of metabolic disorders, such as diabetes and obesity in humans [8–14, 16, 17, 21, 27].

Brown adipocyte, which is endowed with mitochondria, is the most important thermogenic functional unit of BAT [4, 28]. In the mitochondrion of brown

adipocyte, energy generated from nutrients is initially stored as proton gradient membrane potential across the mitochondrial inner membrane, and then converted directly to thermo energy by the uncoupling protein 1 (UCP1)-mediated proton flow [4, 29]. In BAT, brown adipocytes are surrounded by abundant capillaries. The heat generated in the brown adipocyte mitochondria can be quickly distributed by the blood flow to maintain the steady core body temperature in mammals [4]. In addition to classical brown adipocytes, beige/brite adipocytes can be induced from the white adipocytes to conduct thermogenesis upon the cold challenge or catecholamine stimulation [26, 30, 31]. This process is termed as “browning” [26, 30, 31]. Thermogenesis in beige/brite adipocytes can also contribute to mammals’ body temperature maintenance and whole body metabolism [26, 30, 31]. Beige/brite adipose tissue generates heat through both UCP1-independent thermogenesis, including calcium cycling in and out of endoplasmic reticulum, futile cycle between creatine and phosphocreatine, and UCP1-dependent thermogenesis in mitochondria [26, 30–33]. The energy metabolism is essential for the optimal UCP1-mediated mitochondrial thermogenic function of brown and beige/brite adipocytes [4, 34, 35]. In this chapter, we discuss the importance of energy metabolism in maintaining brown adipocyte thermogenic function and the proceeding of targeting metabolic disorder through BAT activation in human studies.

2. Fatty acid metabolism in brown adipocytes

Thermogenic brown adipocyte possesses a high capacity for fatty acid β -oxidation that has been reported in both rodent and humans [4, 12, 14, 36]. Fatty acids serve as the activator for UCP1, which is a fatty acid/ H^+ symporter directly mediating proton flow and thermogenesis [37]. Fatty acids also serve as the main energy substrate mediating the uncoupling and thermogenic function in brown adipocytes [2, 36–41]. In addition, fatty acids can enhance brown adipocyte thermogenic capacity through the nuclear receptor peroxisome proliferator-activated receptors (PPARs), which are the master transcription regulators for the expression of genes involved in lipid metabolism, oxidative phosphorylation, and the key thermogenic protein UCP1 in brown adipocytes [42, 43].

Intracellular fatty acids are stored in the format of triglyceride in the heterogeneous multilocular lipid droplets in the brown adipocyte [4]. Upon the cold challenge, triglycerides stored in the brown adipocyte lipid droplet are lipolysed. Triglyceride lipolysis is a sequential process that involves different enzymes, resulting in the liberation of glycerol and fatty acids for heat production [44]. The lipid droplet is composed of triglycerides and cholesterol esters, which are surrounded by a monolayer of phospholipids [44]. Important proteins with regulatory and enzymatic functions, including perilipin and CGI58, coexist on the phospholipid monolayer to regulate the lipid trafficking and other functions of the lipid droplet [44–47]. Perilipin stabilizes the lipid droplet and prevents it from lipolysis under basal condition. Upon cold challenge, β -adrenergic stimulation leads to the activation of G-protein-coupled receptor (GPCR) and adenylate cyclase, which subsequently increases the cAMP level in brown adipocyte [48]. cAMP then activates protein kinase A (PKA), which phosphorylates perilipin at its serine residues [49–52]. The phosphorylated perilipin releases CGI-58, an adipose triglyceride lipase (ATGL)-activating protein. CGI-58, subsequently, binds and activates ATGL. Activated ATGL hydrolyzes triglycerides and generates free fatty acids and diglycerides [50, 53–55]. Upon the cold challenge, PKA also phosphorylates serine residues on another key lipolysis enzyme hormone sensitive lipase (HSL) [56]. Although HSL is capable of hydrolyzing triglycerides, diglycerides, monoglycerides,

and a broad array of other lipid substrate, it is the rate-limiting enzyme for hydrolyzing the diglycerides *in vivo* [56]. The phosphorylated HSL catalyzes the diglycerides and generates free fatty acids and monoglycerides [56]. As the last step of lipolysis, monoglycerides is hydrolyzed by monoacylglycerol lipase (MGL) to form free fatty acids and glycerol [57].

Given the importance of fatty acid in brown adipocyte thermogenic function, it is reasonable to predict that the deficiency of the key lipolysis enzymes and fatty acid transportation proteins in brown adipocytes can lead to a defected BAT thermogenic function. Human and rodent *in vivo* studies using both pharmacological and genetic approaches to manipulate triglyceride lipolysis processes were reported [36, 39, 58–60]. In the initial studies, nicotinic acid (NiAc) was used to inhibit intracellular triglyceride lipolysis through acting on metabolite-sensing Gi-protein-coupled GPR109A and subsequent PKA activation [36, 58, 61]. A study in rats suggested the brown adipocyte intracellular lipid lipolysis played a major role in thermogenesis in rodents [36]. It showed that the intracellular triglyceride lipolysis contributes to 84% of thermogenesis during an acute cold challenge (10°C, 2–6 hours) and 74% of thermogenesis during a chronic cold challenge (10°C, 21 days) [36]. The importance of the brown adipocyte intracellular triglyceride-derived fatty acids was also reported in a human study [58]. In this study, administering intracellular triglyceride lipolysis inhibitor NiAc significantly blunted BAT oxidative metabolism in cold-challenged young healthy humans (average 30 years of age with an average BMI of 24.5 kg/m²). During a 3-hour cold challenge at 10°C, NiAc administration suppressed BAT intracellular triglyceride lipolysis by about 50% and BAT oxidative metabolism by 70%, despite of the increased blood flow in the BAT [58].

One caveat from both the aforementioned human and rodent *in vivo* studies is that NiAc-mediated lipolysis inhibition took effect in both brown and white adipocytes in addition to other tissues, which makes it hard to delineate the contribution of lipolysis from each cell type. An *in vitro* study nicely confirmed the importance of intracellular lipolysis in brown adipocytes [2]. In cultured primary mouse brown adipocytes, the inhibition of both of the key lipolysis enzymes adipose triglyceride lipase (ATGL) and hormone-sensitive lipase (HSL) led to a 97% decrease in isopropanol-induced brown adipocyte respiration, indicating the imperative requirement of intracellular lipolysis in activated brown adipocytes [2]. These studies suggested that fatty acids liberated from the intracellular triglyceride storage serve as a critical fuel resource and contribute significantly to BAT thermogenesis in brown adipocytes upon activation.

Further studies in genetically manipulated mice showed similar results. The ATGL-knockout mouse presented defective triglyceride lipolysis function with increased BAT weight (8.5-fold), enlarged lipid droplet (20-fold), and decreased BAT explant lipid hydrolysis activity (–85%) [38]. The impaired triglyceride lipolysis activity in the *ATGL-knockout* mouse led to a defective thermogenic function. Upon a 5-hour 4°C challenge, mouse body temperature dropped to a critical low point at around 25°C [38]. Another study using mice with CGI-58 deficiency in both white and brown adipocytes also showed decreased BAT thermogenic function. The *adipose-CGI-58-knockout* mice were cold-sensitive, but only under fasted state [60]. The *HSL-null* mouse adipose tissue also presented defected triglyceride lipolysis function. Upon catecholamine-stimulation, *HSL-null* adipose tissue explants exhibited significantly reduced fatty acid and almost blunted glycerol release into the culture medium, parallel with diglycerides accumulation in both white and brown adipose tissue [56, 62]. However, it seems that HSL-mediated lipolysis function is not as critical as ATGL-mediated lipolysis function in brown adipocytes. An *in vitro* study showed that isopropanol-induced UCP1 activity was

largely dependent on ATGL function (80%) compared to HSL function (35%) in cultured primary brown adipocytes, and the combined inhibition of both ATGL and HSL functions led to an almost complete block of UCP1-mediated thermogenic function (97%) [2].

These studies highlight the cardinal role of intracellular triglyceride liberation in the brown adipocyte thermogenic function, but raise the question if the brown adipocyte intracellular lipolysis is essential for BAT to maintain the adaptive thermogenesis *in vivo*. Interestingly, recent studies using *UCP1-Cre*-mediated knockout of either ATGL or CGI-58 gene in mice showed an unexpected insignificant impact of brown adipocyte intracellular lipolysis and suggested that the circulating fatty acids mobilized from peripheral tissues may play more important roles in BAT thermogenesis in mice [59, 60]. These studies demonstrated that the loss of either ATGL or CGI58-mediated triglycerides lipolysis in brown adipocytes did not compromise the cold challenge-induced thermogenic response in mice [59, 60]. The phenotypic discrepancy between the systemic and BAT-specific *ATGL* and *CGI-58* knockout mice brings very important insights to *in vivo* BAT fatty acid metabolism, and indicates that the brown adipocyte intracellular lipolysis is not the only energy resource for *in vivo* adaptive thermogenesis during cold challenge. Although it is not clear about the exact partition of intracellular and circulating fatty acids contribution to the BAT thermogenic function, it is reasonable to speculate that the core body temperature maintenance is vital for mammals to maintain their optimal function during the cold challenge, when the intracellular lipolysis function is impaired or insufficient, circulating energy substrates from other metabolic tissues, that is, white adipose tissue and liver, are mobilized to provide energy substrates for BAT-mediated adaptive thermogenesis to maintain the adequate core body temperature and ensure the optimal functionality of mammals.

Long-chain fatty acids (LCFAs) are the most abundant format of fatty acid energy substrate in mammals [63–66]. The liberated LCFAs from intracellular lipid droplet are facilitated and transported to mitochondrion and nucleus by fatty acid-binding proteins (FABPs) to conduct their functions [67]. Of the six FABP isoforms, FABP4 (also termed adipocyte p2) and FABP5 are the major FABP isoforms in brown adipocytes [68–74]. Mice mutated in both *FABP4* and *FABP5* gene developed severe hypothermia during fasting after an acute cold challenge (1–3 hours), indicating that fatty acids transportation plays an indispensable role in BAT thermogenic function [74].

The LCFAs-mediated mitochondrial oxidation and UCP-1 activation function require sequential carnitine acyltransferases in order to translocate the LCFA into mitochondrial matrix. Carnitine palmitoyltransferase 1 (CPT1), located on the mitochondrial outer membrane, is the rate-limiting enzyme that mediates LCFAs inward translocation into mitochondrial matrix [63–66]. CPT1 exists in tissue-specific isoforms, and CPT1b is the major isoform expressed in brown adipocytes [75–77]. Mouse embryos-carried homozygous knockout of *Cpt1b* gene were lethal before embryonic day 9.5–11.5 and a normal percentage of *CPT1b*^{+/-} mice was born from the *CPT1b*^{+/-} and wild type breeding pairs [78]. However, more than 50% of the *CPT1b*^{+/-} pups were lost before weaning [78]. The detailed experiment showed that ~7% *CPT1b*^{+/-} mice developed fatal hypothermia following a 3-hour cold challenge (4°C) and ~52% *CPT1b*^{+/-} mice developed fatal hypothermia following a 6-hour cold challenge (4°C), indicating the critical contribution of CPT1b-mediated LCFAs transportation and thermogenesis during the cold challenge in infant/young mice [78]. Carnitine palmitoyltransferase 2 (CPT2) is located on mitochondrial inner membrane to further mediate LCFAs translocation into mitochondrial matrix. In line with the study using the *CPT1b*-deficient mice, an *adipose-specific CPT2 knockout* mice also presented hypothermic phenotype after 3 hours cold

challenge (4°C) and decreased LCFAs oxidation in isolated MEF cells [39, 79]. These studies indicate that fatty acid transportation is critical for BAT thermogenic function during the cold challenge.

The intramitochondrial LCFAs contribute to the thermogenesis through UCP-1 activation and β -oxidation. A detailed study confirmed that mice with impaired fatty acid β -oxidation are cold-intolerant [80]. The acyl-CoA dehydrogenases, which catalyze the initial steps of fatty acid β -oxidation, are composed of a group of enzymes, including very long-chain acyl CoA dehydrogenase (VLCAD), long-chain acyl CoA dehydrogenase (LCAD), and short-chain acyl CoA dehydrogenase (SCAD) [81, 82]. A detailed experiment showed that fetal hypothermia was presented in all of the homozygous knockout mice ($VLCAD^{-/-}$, $LCAD^{-/-}$, and $SCAD^{-/-}$) after 1-hour 4°C challenge [80]. Although the mice with single heterozygous of VLCAD, LCAD, and SCAD genes were cold tolerate; more than 30% of mice with double heterozygous $VLCAD^{+/-}/LCAD^{+/-}$ or $LCAD^{+/-}/SCAD^{+/-}$ and triple heterozygous $VLCAD^{+/-}/LCAD^{+/-}/SCAD^{+/-}$ combinations developed hypothermia upon the cold challenge [80], indicating the essential role of the LCFAs-mediated thermogenic function in mitochondria.

In summary, these studies highlight the importance of the brown adipocyte lipolysis and the liberated intracellular fatty acid transportation and oxidation during thermogenesis. In addition, these studies indicate that brown adipocytes not only use the intracellular lipid storage, but also the circulating energy substrate to maintain the critical thermogenic function for the organismal survival and optimal function in mammals [2, 12, 38, 44, 56, 59, 60, 80]. In modern days, BAT's ability to metabolize fatty acids mobilized from other peripheral storages, including white adipose tissue and liver, makes it a good potential therapeutic target in humans. Studies have shown that BAT mediates significant plasma lipid clearance during the cold challenge in rodent and humans under both physiological and pathological conditions [83–85].

One study showed that the activated BAT is involved in the basal and post-prandial triglyceride metabolism in rodents [83]. In this study, compared to mice kept at 22°C, mice kept at 4°C had significantly lower triglyceride-rich lipoproteins (TRLs)-triglyceride concentration [83]. The study also showed that the activated BAT was involved in the post-prandial triglyceride metabolic process, evidenced by an oral ^3H -triolein tolerance test. Under cold challenge condition, the BAT ^3H -triolein uptake was significantly higher than that of either liver or skeletal muscle, suggesting the significant triglyceride/triglyceride-rich lipoprotein metabolism in activated BAT. For the triglyceride clearance, circulating triglycerides rose to a peak at 2 hours postprandial and declined subsequently in mice kept at 22°C. In contrast, the triglyceride level remained persistently low in mice kept at 4°C, suggesting that the BAT possesses a high postprandial triglyceride clearance function [83]. Most interestingly, the cold challenge increased the uptake of radio-labeled lipoprotein in BAT and reduced the uptake in liver [83]. The cold challenge-induced lipid clearance shift suggests that BAT can be targeted for lipid metabolism *in vivo*. In a pathological setting, the genetically manipulated $Apoa5^{-/-}$ mice with severe hyperlipidemia were studied. A 4–24 hours 4°C challenge corrected $Apoa5^{-/-}$ mouse plasma lipid concentration to the values comparable to wild-type mice, indicating the significant impact of BAT on whole body lipid metabolism in the rodent under the pathological condition [83, 86, 87]. Other studies also showed that the BAT preferentially uptook plasma triglycerides through peripheral tissue lipolysis, and the selective fatty acids uptake from triglyceride-rich lipoprotein ameliorated hyperlipidemia in rodents [84, 85].

Although the studies in rodent strongly support the importance of BAT in lipid clearance, the significance of BAT in human lipid metabolism is still unclear. The contribution of activated BAT in human body was studied using a dietary radio-labeled LCFAs tracer 14(R,S)-[^{18}F]-fluoro-6-thia-heptadecanoic acid (^{18}F THA) [88].

This study showed that a 4-hour mild cold-challenge at 18°C significantly increased dietary fatty acids distribution in BAT in humans [88]. However, given the relative small volume of BAT tissue, the dietary fatty acids clearance contribution from the BAT is less significant compared to other organs including heart, liver, white adipose tissue, or skeletal muscles, and only contributed to ~0.3% of total body dietary fatty acids clearance upon the mild cold challenge [88]. Similarly, another study using fatty acid tracer ^{18}F -fluoro-thiaheptadecanoic acid (^{18}F THA) showed that a 3-hour cold challenge at 18°C led to four times higher radio-labeled tracer uptake and ~80% metabolic rate increase in the BAT, contributing <1% of total fatty acids clearance rate in human subjects [12]. Nevertheless, these experiments suggest that BAT not only exists, but also is functionally active and can contribute to systemic metabolism in humans. One possible reason for the relatively low BAT contribution in fatty acids metabolism in humans could be due to the short cold challenge time and mild cold challenge conditions. It is possible that during the acute cold challenge, intracellular lipid lipolysis serves as the main energy resource so that it ameliorates the clearance of circulating TRLs or fatty acids.

The circulating TRLs or fatty acids are transported into brown adipocyte by a group of proteins, including lipoprotein lipase (LPL) and fatty acid transport proteins [83, 89–93]. LPL is a multifunctional protein produced by many tissues, including the adipose tissue [94]. LPL serves as a rate-limiting enzyme mediating extracellular lipolysis [94]. It hydrolyzes triglycerides into lipid-rich proteins into fatty acids and monoacylglycerol. It also mediates the cellular uptake of triglyceride and other lipids [94]. Studies have shown that cold challenge or catecholamine-stimulation induce the expression and activity of LPL in brown adipocytes through a cAMP-mediated mechanism [89–91]. After activation, LPL is released from the brown adipocyte, transferred to the capillary endothelium lumen, and serves as the anchor between the endothelium cell surface and the TRLs [95–97]. Next, the LCFAs liberated from LPL channel into brown adipocyte for thermogenic function [98, 99]. A study indicated that the local LPL activity is required for TRLs uptake into the BAT [83]. In this study, it is shown that either LPL-specific inhibitor tetrahydrolipstatin pretreatment or releasing LPL from endothelium by heparin significantly blocked the uptake of TRL and fatty acids in BAT [83].

The liberated LCFAs in circulation can be transported into cells and activated by both transmembrane fatty acid transporter proteins (FATPs) and scavenger receptor CD36 [92, 93]. FATPs are composed by a family of six proteins mediating circulating LCFAs uptake and distribution in cells [83, 92, 93, 100]. Among the six isoforms, FATP1 (SLC27A1) is the major isoform in brown and white adipose tissue [83, 92, 93, 100]. Studies showed that postprandial lipid uptake is highly dependent on the adipose tissue FATP1 [101]. The FATP1-knockout mice have decreased lipid uptake in adipose tissue and a compensatory lipid redistribution in liver and heart, where FATP1-mediated lipid uptake function is not required under normal conditions [101]. The cold challenge can induce FATP1 expression in BAT [100]. In line with this, the isolated FATP1-null brown adipocytes showed significantly less fatty acids uptake upon catecholamine stimulation [100]. *In vivo* studies showed that in cold challenged FATP1-knockout mice, BAT triglyceride storage was decreased and circulating serum free fatty acids was increased, indicating the importance of FATP1-mediated fatty acid uptake in BAT [100]. The LCFAs uptake can also be mediated by the transmembrane class B scavenger receptor CD36 [83, 102, 103]. Upon the cold challenge, CD36 is significantly upregulated in adipocytes [83, 102]. Other studies using $CD36^{-/-}$ mice clearly indicated the importance of CD36-mediated LCFAs transportation during thermogenesis. Around 60% of $Cd36^{-/-}$ mice died during a 24-hour cold challenge, which is paralleled with drastically increased plasma free fatty acids concentration [83, 102]. A recent

study indicated that CD36-mediated coenzyme Q (CoQ) uptake is required for the BAT mitochondrial thermogenic function [104].

In summary, fatty acid is indispensable for the brown adipocyte thermogenic function. Fatty acid mediates brown adipocyte thermogenesis through mitochondrial β -oxidation, UCP1-activation, and fatty acid-mediated thermogenic gene regulation. Both the intracellular fatty acids from brown adipocyte lipolysis and the liberated fatty acids from peripheral tissues play essential roles mediating the critical BAT adaptive thermogenic function to maintain the adequate core body temperature in mammals. In modern days, activating BAT thermogenic function to increase fatty acids uptake and utilization may offer new therapeutics to treat human metabolic disorders.

3. Glucose metabolism in BAT

The radio-labeled glucose analog ^{18}F -fluorodeoxyglucose (^{18}F -FDG), in combination with positron emission tomography (PET) and computed tomography (CT), provides a reliable method for the *in vivo* tissue glucose uptake study [105, 106]. Based on studies using this method, it is now well-recognized that BAT in both rodents and humans possesses a significant glucose uptake capacity upon the cold challenge [4, 9–12, 25, 26, 107–112].

Glucose uptake is regulated in brown adipocytes. *In vivo* studies showed that cold challenge significantly enhanced insulin sensitivity and subsequent glucose uptake in the BAT [9, 10, 25, 26, 107, 112, 113]. Interestingly, other studies showed that starved rats with low insulin level also had increased BAT glucose uptake during the cold challenge, indicating that the enhanced glucose uptake is not completely insulin-dependent [114]. Additional studies also indicated that the enhanced glucose uptake in brown adipocytes can be mediated by different pathways in addition to insulin stimulation, including β -adrenergic receptor activation, AMP-activated protein kinase (AMPK) activation, and mTOR activation [115–117].

The importance of glucose metabolism in BAT is a long-standing question. Glucose transporters, which facilitate glucose across the cell plasma membrane is the first rate-limiting step of the glucose metabolism [34, 118, 119]. Intracellular glucose is subsequently phosphorylated to glucose-6-phosphate (G6P) by the enzyme hexokinase (HK). Glucose-6-phosphate serves as the substrate into different pathways, including glycolysis, glycogen synthesis, and the pentose phosphate pathway (PPP) [34, 118, 119]. Glycolysis breaks down glucose to pyruvate to generate small amount of ATP and NADH [34, 118, 119]. The pyruvate is then transported into mitochondria for oxidation and energy production. Under hypoxia condition, pyruvate is disposed in the form of lactate [34, 118, 119].

Early study indicates that glucose and its metabolites contribute to the brown adipocyte thermogenesis by showing that catecholamine-induced glucose uptake was decreased when mitochondrial β -oxidation was inhibited in brown adipocytes [120]. Other studies in brown adipocyte glucose transportation also indicate the importance of the glucose in brown adipocyte metabolism [6, 112, 121, 122]. Both glucose transporter 1 (Glut1) and glucose transporter 4 (Glut4) are abundantly expressed in brown adipocytes and the insulin sensitive Glut4 is the major isoform [112, 122]. The *in vitro* study showed that knock-down of Glut1 and/or Glut4 gene in cultured brown adipocytes impaired the catecholamine-induced cell oxygen consumption by 30–50% [6, 121]. Other studies indicate the importance of glycolysis in brown adipocyte thermogenesis [115, 123, 124]. An *in vitro* study showed that the knockdown of two glycolysis enzymes, HK2 or pyruvate kinase M (PKM), decreased glucose uptake and catecholamine-induced cell oxygen consumption by 67% or

34% respectively, in brown adipocytes [6]. These studies suggested the importance of glucose metabolism in brown adipocyte function. However, another detailed study indicated that glucose oxidation does not play a major role in brown adipocyte metabolism and thermogenesis, by showing that the rate of $^{14}\text{CO}_2$ formation from the ^{14}C glucose was relatively small compared with the maximum rate of oxygen consumption in activated brown adipocytes [125]. Studies using radio-labeled glucose also suggested that glucose uptake was only sufficient to fuel maximally ~15% of the thermogenic capacity in activated rodent brown adipocytes, suggesting that the significantly upregulated glucose uptake in activated brown adipocytes is disassociated with the relative low glucose-mediated thermogenic capacity [123, 125, 126]. A more recent study indicates that the brown adipocyte energy production from glucose depends on the state of the cell: glucose and fatty acid contribute equally to brown adipocyte oxygen consumption under the basal condition; upon catecholamine-activation, oxygen consumption is mainly fueled by fatty acids [6].

The dissociation between high glucose uptake and low glucose-mediated thermogenesis in activated BAT raises an important question: what is the function of the intracellular glucose in brown adipocytes? The importance of glucose in the *de novo* lipogenesis in brown adipocytes has been reported [127–129]. Studies indicate that glucose uptake is an independent process of thermogenesis in both cold-challenged and catecholamine-activated BAT, which further supports that glucose uptake might play other role as energy substrate in activated brown adipocytes [116, 117, 130–132]. One detailed *in vitro* study using rat brown adipocytes showed that norepinephrine significantly enhanced glucose oxidation by sevenfold, while it also inhibited lipogenesis at the same time. On the other hand, insulin stimulation increased the lipogenesis by sevenfold in brown adipocytes whereas glucose oxidation remained very low. Most interestingly, the addition of insulin to the norepinephrine only potentiated the enhanced glucose oxidation, but do not enhance the lipogenesis [115]. On the other hand, another study showed that brown adipocytes converted a greater proportion of metabolized glucose into lactate and pyruvate, but only a smaller proportion into fatty acids through insulin-mediated pathway [125]. These studies suggest that glucose metabolism is involved in two different states in brown adipocytes, the thermogenic state and nonthermogenic state. It is plausible that during the thermogenic state, glucose contributes to both the energy production and lipogenesis; and during the nonthermogenic state, glucose contributes mainly to the lipogenesis and energy storage process in brown adipocytes, which explains the disassociation of glucose uptake and glucose metabolism in brown adipocytes. In addition to glucose, other energy substrates, including glycerol and amino acid (glutamate), might also contribute to BAT metabolism and thermogenesis in human and rodent brown adipocytes. However, the relative contribution and partition of these energy substrates are unclear [132–135].

4. Energy storage in BAT

The cold challenge not only enhances catabolic processes mediating energy substrate metabolism and heat generation, but also induces anabolic processes mediating fatty acid synthesis and lipogenesis, as well as glycogenesis [36, 111, 136–139].

Glucose can be stored as glycogen in brown adipocytes [36, 139]. Studies showed that glycogen synthases (GStot) and uridine diphosphate glucose pyrophosphorylase (Udgp), which mediates glycogenesis, were upregulated upon the cold challenge in BAT [36, 139]. Interestingly, glycogen hydrolysis enzyme, glycogen phosphorylase (Pygl), was also upregulated after cold challenge [36]. Although cold-challenge upregulates both glycogen synthesis and degradation, it is reported

that the stored glycogen is used up shortly after the cold challenge [139]. Indicating that glycogen is not an efficient format for energy storage and a sustainable energy resource in brown adipocytes.

The glucose uptaken by brown adipocyte can also be converted to fatty acid through the *de novo* lipogenesis. Carbohydrate response element-binding protein (ChREBP) is the major transcription factor mediating fatty acid synthesis in adipocytes. There are two isoforms of ChREBP, ChREBP α and ChREBP β . ChREBP α is abundantly expressed in BAT from rodents and humans [137, 140, 141]. *De novo* lipogenesis involves a series of enzymes mediating the sequential reactions converting glucose-derived citrate into fatty acids, including ATP-citrate lyase (ACLY), acetyl-CoA carboxylases 1 (ACC1), fatty acid synthase (FASN), and stearoyl-CoA desaturase-1 (SCD1). ChREBP α , coordinates with another transcription factor Max-like protein (MLX), directly binds to the carbohydrate response elements (ChoRE) and upregulates these *de novo* lipogenic genes expressions. ChREBP α can also increase the expression of ChREBP β to further activate the *de novo* lipogenesis enzymes [142, 143]. It has been reported that lipogenic genes and AKT2-ChREBP pathways are upregulated to optimize fuel storage and thermogenesis upon cold stimulation in BAT [136]. In accordance with these studies, *ChREBP*^{-/-} mice presented less BAT weight [142], and adipose-specific *ChREBP-knockout* mice had decreased carbohydrate-induced lipogenesis in BAT [144]. These studies indicate that ChREBP plays important roles in brown adipocyte's *de novo* lipogenesis and energy storage.

Sterol regulatory element-binding protein-1 (SREBP-1) is another transcription factor mediating the *de novo* lipogenesis [145]. Of the three different SREBP isoforms, SREBP1c is more abundant and SREBP1a is less abundant in the adipose tissue [137, 145, 146]. *In vitro* study showed that SREBP1c is sufficient to regulate lipogenic enzymes in cultured adipocytes [147]. In the adipose-specific *ap2-SREBP1c* transgenic mice, lipogenic enzyme *Acc1*, *FASN* and *Scd1* expression as well as fatty acids synthesis rate were significantly upregulated in brown adipocytes [148]. In line with this study, *ap2-SREBP1a* transgenic mice also developed adipose tissue hypertrophy in accordance with an increased lipogenic enzyme profile and enhance *de novo* lipogenesis in the BAT [148]. However, some *in vivo* studies indicated that SREBP1c's role in the *de novo* lipogenesis in adipocytes is dispensable, as evidenced by *SREBP1-knockout* mice have normal lipogenic enzymes gene expression profile and normal lipid storage in their adipose tissue [149, 150]. These studies suggest that SREBP-1 is involved in the brown adipocyte lipogenesis and triglyceride storage when the excessive energy resources are available; however, SREBP-1's function can be compensated by other factors when it is absent.

In summary, these studies suggest that in activated brown adipocytes, glycogenesis and lipogenesis are upregulated to store/restore energy substrates, which is parallel with energy substrate metabolism and thermogenesis. These coordinated anabolic and catabolic processes are important to maintain the brown adipocyte energy homeostasis.

5. The proceeding of targeting BAT in human metabolic disorders

The understating of BAT energy homeostasis and the discovery of the functional BAT in humans lead to significant interests in targeting BAT for metabolic disorders, for example, obesity, insulin resistance, type 2 diabetes, and lipid profile abnormality. The ability of BAT metabolizing fatty acid and glucose from the intracellular storage, the peripheral tissues liberation, and the dietary nutrition absorption makes it a good potential therapeutic target for combating metabolic disorders in humans. In addition to the BAT, beige/brite fat, which is coexisted in white adipose tissue,

can be recruited and activated (browning) in response to cold challenge or pharmacological stimulation and serves as a target for metabolic disorders [151–154].

It has been reported that the BAT ^{18}F -FDG uptake in humans correlates inversely with aging, adiposity, diabetic status, and BMI, indicating that the manipulation of BAT function is a possible approach for combating metabolic disorders [8–14, 16, 21–24, 155, 156]. The studies in mouse models and humans provide evidence for the metabolic benefit of BAT. Mice with genetic ablation of BAT and the *UCP1-knockout* mice under thermoneutrality, both developed obesity [18–20, 85]. In addition, BAT activation reduced hypercholesterolemia and protected mice from atherosclerosis development and liver steatosis [18–20, 85]. It is well-recognized that BAT can be activated upon seasonal temperature changes or short time period (several hours) mild cold challenge (16–19°C) in humans [7, 9, 10, 12, 21, 22, 24–27, 157]. However, the significance of BAT's contribution in human metabolism has not been clearly elucidated. Human studies indicated that BAT activity is inversely correlated with body fat deposition, suggests BAT can serve as a target for obesity [9, 10, 16, 21–24]. Studies also showed that fasting glucose was lower in the human subjects with higher BAT prevalence and the BAT activity were blunted in subjects with obesity [11, 16, 155]. More interestingly, in the same patients with multiple PET scans, BAT was more detectable when fasting glucose in the subjects were lower [16]. Other studies showed that a mild cold challenge significantly increased whole body glucose disposal, glucose oxidation, insulin sensitivity, and whole body energy expenditure in human subjects [13, 17, 158, 159]. Additional study showed that moderate cold challenge (18.06°C) significantly improved the peripheral glucose uptake and insulin sensitivity by 20%, but did not impact the pancreatic insulin secretion [17]. These studies strongly indicate the therapeutic potentials of targeting BAT for glucose metabolism in humans, but leave the question of the relative contribution of activated BAT in whole body glucose metabolism to be answered.

Cold challenge can also enhance BAT lipid metabolism. Studies showed that cold challenge led to significantly enhanced lipid mobilization, increased plasma fatty acid levels, as well as upregulated genes for lipid metabolism in human BAT [12, 58, 156, 160]. It is reported that circulating fatty acids were uptaken by BAT in cold-challenged humans by using the ^{18}F THA tracing method [88]. In addition, it has been shown that cold-activated BAT significantly contributed to whole body fatty acid utilization in healthy humans [17]. The fatty acid uptake is significantly lower in overweight human subjects compared to healthy humans [156]. Importantly, BAT activation upon mild cold challenge significantly increased systemic lipid metabolism, whole body lipolysis, triglyceride-fatty acid cycling, and fatty acid oxidation in overweight/obese subjects [14]. These studies suggest that BAT contributes to whole body lipid metabolism and homeostasis in healthy humans as well as in humans with metabolic disorders, suggesting that BAT can serve as a target for lipid abnormality in humans.

The significance of BAT/beige adipose tissue in human whole body metabolism has been studied and remarkable progress has been made in recent years. The acknowledgment that BAT can be activated and subsequently contributes to the human whole body energy expenditure is encouraging. Although the capacity and relative significance of BAT's contribution to whole body energy substrate metabolism has not been elucidated, it should be noticed that majority of the studies in humans were conducted for relative short time period with mild cold challenge conditions. Given, humans usually live under thermoneutrality, we can assume that their BAT functions are repressed under this condition. In addition, the prevalence of BAT in humans varies significantly, which depends on individual's life style, physical activity, and health conditions, which makes it harder to evaluate the contribution of BAT in whole body metabolism [7, 9–12, 21, 22, 24–27, 88, 156, 157].

Hence, a sustainable chronic mild cold challenge strategy aiming to recruit more BAT/beige adipose tissue and enhance their oxidative capacity might provide more significant therapeutic potential in humans over time, especially the human subjects with metabolic disorders. Future studies should also delineate the relative contribution from glucose and fatty acid in human BAT under physiological and pathological conditions; as well as compare the glucose and fatty acid partitioning in different tissues and organs, including skeletal muscle, heart, liver and BAT. These details will guide us to establish better strategies targeting metabolic disorders through BAT activation in the future.

6. Conclusion

The energy metabolism plays critical role in maintaining BAT thermogenic function in mammals. Through the energy metabolism, the chemical energy stored in nutrients (i.e., fatty acids and glucose) can be converted to thermoenergy and dissipated as heat in the BAT. Activated brown adipocytes not only contribute to the intracellular substrate homeostasis, but also contribute significantly to the whole body energy metabolism. BAT with functional thermoactivity is present in adult humans. Thus, BAT activation has remarkable therapeutic implications in human metabolic disorder management.

Acknowledgements

This work was supported by American Heart Association National Scientist Development grant, AHA SDG grant, 12050558 to Yuan Lu; and NIH R35HL135789 to Mukesh K. Jain.

Conflict of interest

The author has no conflict of interest to declare.

Author details

Yuan Lu^{1,2*}

1 Case Western Reserve University School of Medicine, Cleveland, Ohio, USA

2 Harrington Heart and Vascular Institute, University Hospitals Cleveland Medical Center, Cleveland, Ohio, USA

*Address all correspondence to: yuan.lu@case.edu

IntechOpen

© 2019 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] Himms-Hagen J. Nonshivering thermogenesis. *Brain Research Bulletin*. 1984;**12**(2):151-160
- [2] Li Y et al. Taking control over intracellular fatty acid levels is essential for the analysis of thermogenic function in cultured primary brown and brite/beige adipocytes. *EMBO Reports*. 2014;**15**(10):1069-1076
- [3] Oelkrug R, Polymeropoulos ET, Jastroch M. Brown adipose tissue: Physiological function and evolutionary significance. *Journal of Comparative Physiology. B*. 2015;**185**(6):587-606
- [4] Cannon B, Nedergaard J. Brown adipose tissue: Function and physiological significance. *Physiological Reviews*. 2004;**84**(1):277-359
- [5] Kajimura S, Seale P, Spiegelman BM. Transcriptional control of brown fat development. *Cell Metabolism*. 2010;**11**(4):257-262
- [6] Winther S et al. Restricting glycolysis impairs brown adipocyte glucose and oxygen consumption. *American Journal of Physiology. Endocrinology and Metabolism*. 2018;**314**(3):E214-E223
- [7] Cohade C et al. Uptake in supraclavicular area fat ("USA-fat"): Description on 18F-FDG PET/CT. *Journal of Nuclear Medicine*. 2003;**44**(2):170-176
- [8] Virtanen KA et al. Functional brown adipose tissue in healthy adults. *The New England Journal of Medicine*. 2009;**360**(15):1518-1525
- [9] Cypess AM et al. Identification and importance of brown adipose tissue in adult humans. *The New England Journal of Medicine*. 2009;**360**(15):1509-1517
- [10] van Marken Lichtenbelt WD et al. Cold-activated brown adipose tissue in healthy men. *The New England Journal of Medicine*. 2009;**360**(15):1500-1508
- [11] Orava J et al. Different metabolic responses of human brown adipose tissue to activation by cold and insulin. *Cell Metabolism*. 2011;**14**(2):272-279
- [12] Ouellet V et al. Brown adipose tissue oxidative metabolism contributes to energy expenditure during acute cold exposure in humans. *The Journal of Clinical Investigation*. 2012;**122**(2):545-552
- [13] Chondronikola M et al. Brown adipose tissue improves whole-body glucose homeostasis and insulin sensitivity in humans. *Diabetes*. 2014;**63**(12):4089-4099
- [14] Chondronikola M et al. Brown adipose tissue activation is linked to distinct systemic effects on lipid metabolism in humans. *Cell Metabolism*. 2016;**23**(6):1200-1206
- [15] Foster DO. Quantitative contribution of brown adipose tissue thermogenesis to overall metabolism. *Canadian Journal of Biochemistry and Cell Biology*. 1984;**62**(7):618-622
- [16] Lee P et al. A critical appraisal of the prevalence and metabolic significance of brown adipose tissue in adult humans. *American Journal of Physiology. Endocrinology and Metabolism*. 2010;**299**(4):E601-E606
- [17] Iwen KA et al. Cold-induced brown adipose tissue activity alters plasma fatty acids and improves glucose metabolism in men. *The Journal of Clinical Endocrinology and Metabolism*. 2017;**102**(11):4226-4234

- [18] Lowell BB et al. Development of obesity in transgenic mice after genetic ablation of brown adipose tissue. *Nature*. 1993;**366**(6457):740-742
- [19] Feldmann HM et al. UCP1 ablation induces obesity and abolishes diet-induced thermogenesis in mice exempt from thermal stress by living at thermoneutrality. *Cell Metabolism*. 2009;**9**(2):203-209
- [20] Poekes L et al. Activation of brown adipose tissue enhances the efficacy of caloric restriction for treatment of nonalcoholic steatohepatitis. *Laboratory Investigation*. 2018
- [21] Saito M et al. High incidence of metabolically active brown adipose tissue in healthy adult humans: Effects of cold exposure and adiposity. *Diabetes*. 2009;**58**(7):1526-1531
- [22] Yoneshiro T et al. Age-related decrease in cold-activated brown adipose tissue and accumulation of body fat in healthy humans. *Obesity (Silver Spring)*. 2011;**19**(9):1755-1760
- [23] Pfannenberger C et al. Impact of age on the relationships of brown adipose tissue with sex and adiposity in humans. *Diabetes*. 2010;**59**(7):1789-1793
- [24] Ouellet V et al. Outdoor temperature, age, sex, body mass index, and diabetic status determine the prevalence, mass, and glucose-uptake activity of 18F-FDG-detected BAT in humans. *The Journal of Clinical Endocrinology and Metabolism*. 2011;**96**(1):192-199
- [25] Hany TF et al. Brown adipose tissue: A factor to consider in symmetrical tracer uptake in the neck and upper chest region. *European Journal of Nuclear Medicine and Molecular Imaging*. 2002;**29**(10):1393-1398
- [26] Enerback S. The origins of brown adipose tissue. *The New England Journal of Medicine*. 2009;**360**(19):2021-2023
- [27] Carpentier AC et al. Brown adipose tissue energy metabolism in humans. *Frontiers in Endocrinology (Lausanne)*. 2018;**9**:447
- [28] Lindberg O et al. Studies of the mitochondrial energy-transfer system of brown adipose tissue. *The Journal of Cell Biology*. 1967;**34**(1):293-310
- [29] Kozak UC et al. An upstream enhancer regulating brown-fat-specific expression of the mitochondrial uncoupling protein gene. *Molecular and Cellular Biology*. 1994;**14**(1):59-67
- [30] Ishibashi J, Seale P. Medicine. Beige can be slimming. *Science*. 2010;**328**(5982):1113-1114
- [31] Petrovic N et al. Chronic peroxisome proliferator-activated receptor gamma (PPARgamma) activation of epididymally derived white adipocyte cultures reveals a population of thermogenically competent, UCP1-containing adipocytes molecularly distinct from classic brown adipocytes. *The Journal of Biological Chemistry*. 2010;**285**(10):7153-7164
- [32] Ikeda K et al. UCP1-independent signaling involving SERCA2b-mediated calcium cycling regulates beige fat thermogenesis and systemic glucose homeostasis. *Nature Medicine*. 2017;**23**(12):1454-1465
- [33] Kazak L et al. A creatine-driven substrate cycle enhances energy expenditure and thermogenesis in beige fat. *Cell*. 2015;**163**(3):643-655
- [34] Gottlieb E. p53 guards the metabolic pathway less travelled. *Nature Cell Biology*. 2011;**13**(3):195-197
- [35] Golozoubova V et al. Only UCP1 can mediate adaptive nonshivering thermogenesis in the cold. *The FASEB Journal*. 2001;**15**(11):2048-2050
- [36] Labbe SM et al. In vivo measurement of energy substrate

contribution to cold-induced brown adipose tissue thermogenesis. The FASEB Journal. 2015;**29**(5):2046-2058

[37] Fedorenko A, Lishko PV, Kirichok Y. Mechanism of fatty-acid-dependent UCP1 uncoupling in brown fat mitochondria. Cell. 2012;**151**(2):400-413

[38] Haemmerle G et al. Defective lipolysis and altered energy metabolism in mice lacking adipose triglyceride lipase. Science. 2006;**312**(5774):734-737

[39] Lee J, Ellis JM, Wolfgang MJ. Adipose fatty acid oxidation is required for thermogenesis and potentiates oxidative stress-induced inflammation. Cell Reports. 2015;**10**(2):266-279

[40] Shabalina IG et al. Within brown-fat cells, UCP1-mediated fatty acid-induced uncoupling is independent of fatty acid metabolism. Biochimica et Biophysica Acta. 2008;**1777**(7-8):642-650

[41] Osuga J et al. Targeted disruption of hormone-sensitive lipase results in male sterility and adipocyte hypertrophy, but not in obesity. Proceedings of the National Academy of Sciences of the United States of America. 2000;**97**(2):787-792

[42] Lock EA, Mitchell AM, Elcombe CR. Biochemical mechanisms of induction of hepatic peroxisome proliferation. Annual Review of Pharmacology and Toxicology. 1989;**29**:145-163

[43] Zechner R, Madeo F, Kratky D. Cytosolic lipolysis and lipophagy: Two sides of the same coin. Nature Reviews. Molecular Cell Biology. 2017;**18**(11):671-684

[44] Zimmermann R et al. Fate of fat: The role of adipose triglyceride lipase in lipolysis. Biochimica et Biophysica Acta. 2009;**1791**(6):494-500

[45] Londos C et al. Role of PAT proteins in lipid metabolism. Biochimie. 2005;**87**(1):45-49

[46] Brasaemle DL. Thematic review series: Adipocyte biology. The perilipin family of structural lipid droplet proteins: Stabilization of lipid droplets and control of lipolysis. Journal of Lipid Research. 2007;**48**(12):2547-2559

[47] Ducharme NA, Bickel PE. Lipid droplets in lipogenesis and lipolysis. Endocrinology. 2008;**149**(3):942-949

[48] Steinberg D, Huttunen JK. The role of cyclic AMP in activation of hormone-sensitive lipase of adipose tissue. Advances in Cyclic Nucleotide Research. 1972;**1**:47-62

[49] Sztalryd C et al. Perilipin A is essential for the translocation of hormone-sensitive lipase during lipolytic activation. The Journal of Cell Biology. 2003;**161**(6):1093-1103

[50] Granneman JG et al. Analysis of lipolytic protein trafficking and interactions in adipocytes. The Journal of Biological Chemistry. 2007;**282**(8):5726-5735

[51] Su CL et al. Mutational analysis of the hormone-sensitive lipase translocation reaction in adipocytes. The Journal of Biological Chemistry. 2003;**278**(44):43615-43619

[52] Miyoshi H et al. Perilipin promotes hormone-sensitive lipase-mediated adipocyte lipolysis via phosphorylation-dependent and -independent mechanisms. The Journal of Biological Chemistry. 2006;**281**(23):15837-15844

[53] Miyoshi H et al. Control of adipose triglyceride lipase action by serine 517 of perilipin A globally regulates protein kinase A-stimulated lipolysis in adipocytes. The Journal of Biological Chemistry. 2007;**282**(2):996-1002

- [54] Zimmermann R et al. Fat mobilization in adipose tissue is promoted by adipose triglyceride lipase. *Science*. 2004;**306**(5700):1383-1386
- [55] Jenkins CM et al. Identification, cloning, expression, and purification of three novel human calcium-independent phospholipase A2 family members possessing triacylglycerol lipase and acylglycerol transacylase activities. *The Journal of Biological Chemistry*. 2004;**279**(47):48968-48975
- [56] Haemmerle G et al. Hormone-sensitive lipase deficiency in mice causes diglyceride accumulation in adipose tissue, muscle, and testis. *The Journal of Biological Chemistry*. 2002;**277**(7):4806-4815
- [57] Fredrikson G, Tornqvist H, Belfrage P. Hormone-sensitive lipase and monoacylglycerol lipase are both required for complete degradation of adipocyte triacylglycerol. *Biochimica et Biophysica Acta*. 1986;**876**(2):288-293
- [58] Blondin DP et al. Inhibition of intracellular triglyceride lipolysis suppresses cold-induced Brown adipose tissue metabolism and increases shivering in humans. *Cell Metabolism*. 2017;**25**(2):438-447
- [59] Schreiber R et al. Cold-induced thermogenesis depends on ATGL-mediated lipolysis in cardiac muscle, but not Brown adipose tissue. *Cell Metabolism*. 2017;**26**(5):753 e7-763 e7
- [60] Shin H et al. Lipolysis in brown adipocytes is not essential for cold-induced thermogenesis in mice. *Cell Metabolism*. 2017;**26**(5):764 e5-777 e5
- [61] Ahmed K, Tunaru S, Offermanns S. GPR109A, GPR109B and GPR81, a family of hydroxy-carboxylic acid receptors. *Trends in Pharmacological Sciences*. 2009;**30**(11):557-562
- [62] Mulder H et al. Hormone-sensitive lipase null mice exhibit signs of impaired insulin sensitivity whereas insulin secretion is intact. *The Journal of Biological Chemistry*. 2003;**278**(38):36380-36388
- [63] Neely JR, Rovetto MJ, Oram JF. Myocardial utilization of carbohydrate and lipids. *Progress in Cardiovascular Diseases*. 1972;**15**(3):289-329
- [64] Mitchell GA et al. Medical aspects of ketone body metabolism. *Clinical and Investigative Medicine*. 1995;**18**(3):193-216
- [65] Britton CH et al. Human liver mitochondrial carnitine palmitoyltransferase I: Characterization of its cDNA and chromosomal localization and partial analysis of the gene. *Proceedings of the National Academy of Sciences of the United States of America*. 1995;**92**(6):1984-1988
- [66] McGarry JD, Brown NF. The mitochondrial carnitine palmitoyltransferase system. From concept to molecular analysis. *European Journal of Biochemistry*. 1997;**244**(1):1-14
- [67] Furuhashi M, Hotamisligil GS. Fatty acid-binding proteins: Role in metabolic diseases and potential as drug targets. *Nature Reviews. Drug Discovery*. 2008;**7**(6):489-503
- [68] Hunt CR et al. Adipocyte P2 gene: Developmental expression and homology of 5'-flanking sequences among fat cell-specific genes. *Proceedings of the National Academy of Sciences of the United States of America*. 1986;**83**(11):3786-3790
- [69] Haunerland NH, Spener F. Fatty acid-binding proteins—Insights from genetic manipulations. *Progress in Lipid Research*. 2004;**43**(4):328-349

- [70] Makowski L, Hotamisligil GS. The role of fatty acid binding proteins in metabolic syndrome and atherosclerosis. *Current Opinion in Lipidology*. 2005;**16**(5):543-548
- [71] Shen WJ et al. Interaction of rat hormone-sensitive lipase with adipocyte lipid-binding protein. *Proceedings of the National Academy of Sciences of the United States of America*. 1999;**96**(10):5528-5532
- [72] Scheja L et al. Altered insulin secretion associated with reduced lipolytic efficiency in aP2^{-/-} mice. *Diabetes*. 1999;**48**(10):1987-1994
- [73] Coe NR, Simpson MA, Bernlohr DA. Targeted disruption of the adipocyte lipid-binding protein (aP2 protein) gene impairs fat cell lipolysis and increases cellular fatty acid levels. *Journal of Lipid Research*. 1999;**40**(5):967-972
- [74] Syamsunarno MR et al. Fatty acid binding protein 4 and 5 play a crucial role in thermogenesis under the conditions of fasting and cold stress. *PLoS One*. 2014;**9**(6):e90825
- [75] Esser V et al. Expression of a cDNA isolated from rat brown adipose tissue and heart identifies the product as the muscle isoform of carnitine palmitoyltransferase I (M-CPT I). M-CPT I is the predominant CPT I isoform expressed in both white (epididymal) and brown adipocytes. *The Journal of Biological Chemistry*. 1996;**271**(12):6972-6977
- [76] Nyman LR et al. Homozygous carnitine palmitoyltransferase 1a (liver isoform) deficiency is lethal in the mouse. *Molecular Genetics and Metabolism*. 2005;**86**(1-2):179-187
- [77] Yamazaki N et al. High expression of a novel carnitine palmitoyltransferase I like protein in rat brown adipose tissue and heart: Isolation and characterization of its cDNA clone. *FEBS Letters*. 1995;**363**(1-2):41-45
- [78] Ji S et al. Homozygous carnitine palmitoyltransferase 1b (muscle isoform) deficiency is lethal in the mouse. *Molecular Genetics and Metabolism*. 2008;**93**(3):314-322
- [79] Gonzalez-Hurtado E et al. Fatty acid oxidation is required for active and quiescent brown adipose tissue maintenance and thermogenic programming. *Molecular Metabolism*. 2018;**7**:45-56
- [80] Schuler AM et al. Synergistic heterozygosity in mice with inherited enzyme deficiencies of mitochondrial fatty acid beta-oxidation. *Molecular Genetics and Metabolism*. 2005;**85**(1):7-11
- [81] Thorpe C, Kim JJ. Structure and mechanism of action of the acyl-CoA dehydrogenases. *The FASEB Journal*. 1995;**9**(9):718-725
- [82] Gregersen N et al. Mutation analysis in mitochondrial fatty acid oxidation defects: Exemplified by acyl-CoA dehydrogenase deficiencies, with special focus on genotype-phenotype relationship. *Human Mutation*. 2001;**18**(3):169-189
- [83] Bartelt A et al. Brown adipose tissue activity controls triglyceride clearance. *Nature Medicine*. 2011;**17**(2):200-205
- [84] Khedoe PP et al. Brown adipose tissue takes up plasma triglycerides mostly after lipolysis. *Journal of Lipid Research*. 2015;**56**(1):51-59
- [85] Berbee JF et al. Brown fat activation reduces hypercholesterolaemia and protects from atherosclerosis development. *Nature Communications*. 2015;**6**:6356
- [86] Merkel M et al. Apolipoprotein AV accelerates plasma hydrolysis

of triglyceride-rich lipoproteins by interaction with proteoglycan-bound lipoprotein lipase. *The Journal of Biological Chemistry*. 2005;**280**(22):21553-21560

[87] Pennacchio LA et al. An apolipoprotein influencing triglycerides in humans and mice revealed by comparative sequencing. *Science*. 2001;**294**(5540):169-173

[88] Blondin DP et al. Dietary fatty acid metabolism of brown adipose tissue in cold-acclimated men. *Nature Communications*. 2017;**8**:14146

[89] Amri EZ et al. Maturation and secretion of lipoprotein lipase in cultured adipose cells. II. Effects of tunicamycin on activation and secretion of the enzyme. *Biochimica et Biophysica Acta*. 1986;**875**(2):334-343

[90] Carneheim C, Nedergaard J, Cannon B. Cold-induced beta-adrenergic recruitment of lipoprotein lipase in brown fat is due to increased transcription. *The American Journal of Physiology*. 1988;**254**(2 Pt 1):E155-E161

[91] Ong JM, Kern PA. The role of glucose and glycosylation in the regulation of lipoprotein lipase synthesis and secretion in rat adipocytes. *The Journal of Biological Chemistry*. 1989;**264**(6):3177-3182

[92] Stahl A. A current review of fatty acid transport proteins (SLC27). *Pflügers Archiv*. 2004;**447**(5):722-727

[93] Anderson CM, Stahl A. SLC27 fatty acid transport proteins. *Molecular Aspects of Medicine*. 2013;**34**(2-3):516-528

[94] Wang H, Eckel RH. Lipoprotein lipase: From gene to obesity. *American Journal of Physiology. Endocrinology and Metabolism*. 2009;**297**(2):E271-E288

[95] Beigneux AP et al. Glycosylphosphatidylinositol-anchored high-density lipoprotein-binding protein 1 plays a critical role in the lipolytic processing of chylomicrons. *Cell Metabolism*. 2007;**5**(4):279-291

[96] Davies BS et al. GPIHBP1 is responsible for the entry of lipoprotein lipase into capillaries. *Cell Metabolism*. 2010;**12**(1):42-52

[97] Beisiegel U, Weber W, Bengtsson-Olivecrona G. Lipoprotein lipase enhances the binding of chylomicrons to low density lipoprotein receptor-related protein. *Proceedings of the National Academy of Sciences of the United States of America*. 1991;**88**(19):8342-8346

[98] Mead JR, Irvine SA, Ramji DP. Lipoprotein lipase: Structure, function, regulation, and role in disease. *Journal of Molecular Medicine (Berlin, Germany)*. 2002;**80**(12):753-769

[99] Wang CS, Hartsuck J, McConathy WJ. Structure and functional properties of lipoprotein lipase. *Biochimica et Biophysica Acta*. 1992;**1123**(1):1-17

[100] Wu Q et al. Fatty acid transport protein 1 is required for nonshivering thermogenesis in brown adipose tissue. *Diabetes*. 2006;**55**(12):3229-3237

[101] Wu Q et al. FATP1 is an insulin-sensitive fatty acid transporter involved in diet-induced obesity. *Molecular and Cellular Biology*. 2006;**26**(9):3455-3467

[102] Harmon CM, Abumrad NA. Binding of sulfosuccinimidyl fatty acids to adipocyte membrane proteins: Isolation and amino-terminal sequence of an 88-kD protein implicated in transport of long-chain fatty acids. *The Journal of Membrane Biology*. 1993;**133**(1):43-49

[103] Tao N, Wagner SJ, Lublin DM. CD36 is palmitoylated on both

N- and C-terminal cytoplasmic tails. *The Journal of Biological Chemistry*. 1996;**271**(37):22315-22320

[104] Anderson CM et al. Dependence of brown adipose tissue function on CD36-mediated coenzyme Q uptake. *Cell Reports*. 2015;**10**(4):505-515

[105] Som P et al. A fluorinated glucose analog, 2-fluoro-2-deoxy-D-glucose (F-18): Nontoxic tracer for rapid tumor detection. *Journal of Nuclear Medicine*. 1980;**21**(7):670-675

[106] Kelloff GJ et al. Progress and promise of FDG-PET imaging for cancer patient management and oncologic drug development. *Clinical Cancer Research*. 2005;**11**(8):2785-2808

[107] Greco-Perotto R et al. Stimulatory effect of cold adaptation on glucose utilization by brown adipose tissue. Relationship with changes in the glucose transporter system. *The Journal of Biological Chemistry*. 1987;**262**(16):7732-7736

[108] Vallerand AL, Perusse F, Bukowiecki LJ. Cold exposure potentiates the effect of insulin on in vivo glucose uptake. *The American Journal of Physiology*. 1987;**253**(2 Pt 1):E179-E186

[109] Vallerand AL, Perusse F, Bukowiecki LJ. Stimulatory effects of cold exposure and cold acclimation on glucose uptake in rat peripheral tissues. *The American Journal of Physiology*. 1990;**259**(5 Pt 2):R1043-R1049

[110] Shimizu Y, Nikami H, Saito M. Sympathetic activation of glucose utilization in brown adipose tissue in rats. *Journal of Biochemistry*. 1991;**110**(5):688-692

[111] Yu XX et al. Cold elicits the simultaneous induction of fatty acid synthesis and beta-oxidation in murine brown adipose tissue: Prediction

from differential gene expression and confirmation in vivo. *The FASEB Journal*. 2002;**16**(2):155-168

[112] Shimizu Y et al. Increased expression of glucose transporter GLUT-4 in brown adipose tissue of fasted rats after cold exposure. *The American Journal of Physiology*. 1993;**264**(6 Pt 1):E890-E895

[113] Wang X, Wahl R. Responses of the insulin signaling pathways in the brown adipose tissue of rats following cold exposure. *PLoS One*. 2014;**9**(6):e99772

[114] Shibata H et al. Cold exposure reverses inhibitory effects of fasting on peripheral glucose uptake in rats. *The American Journal of Physiology*. 1989;**257**(1 Pt 2):R96-R101

[115] Ebner S et al. Effects of insulin and norepinephrine on glucose transport and metabolism in rat brown adipocytes. Potentiation by insulin of norepinephrine-induced glucose oxidation. *European Journal of Biochemistry*. 1987;**170**(1-2):469-474

[116] Hutchinson DS et al. Beta-adrenoceptors, but not alpha-adrenoceptors, stimulate AMP-activated protein kinase in brown adipocytes independently of uncoupling protein-1. *Diabetologia*. 2005;**48**(11):2386-2395

[117] Olsen JM et al. Glucose uptake in brown fat cells is dependent on mTOR complex 2-promoted GLUT1 translocation. *The Journal of Cell Biology*. 2014;**207**(3):365-374

[118] Bell GI et al. Molecular biology of mammalian glucose transporters. *Diabetes Care*. 1990;**13**(3):198-208

[119] Gatenby RA, Gillies RJ. Why do cancers have high aerobic glycolysis? *Nature Reviews. Cancer*. 2004;**4**(11):891-899

[120] Marette A, Bukowiecki LJ. Noradrenaline stimulates glucose

- transport in rat brown adipocytes by activating thermogenesis. Evidence that fatty acid activation of mitochondrial respiration enhances glucose transport. *The Biochemical Journal*. 1991;**277**(Pt 1): 119-124
- [121] Lee P et al. Brown adipose tissue exhibits a glucose-responsive thermogenic biorhythm in humans. *Cell Metabolism*. 2016;**23**(4):602-609
- [122] Shimizu Y et al. Noradrenaline increases glucose transport into brown adipocytes in culture by a mechanism different from that of insulin. *The Biochemical Journal*. 1996;**314**(Pt 2):485-490
- [123] Ma SW, Foster DO. Uptake of glucose and release of fatty acids and glycerol by rat brown adipose tissue in vivo. *Canadian Journal of Physiology and Pharmacology*. 1986;**64**(5):609-614
- [124] Jeong JH, Chang JS, Jo YH. Intracellular glycolysis in brown adipose tissue is essential for optogenetically induced nonshivering thermogenesis in mice. *Scientific Reports*. 2018;**8**(1):6672
- [125] Saggerson ED, McAllister TW, Baht HS. Lipogenesis in rat brown adipocytes. Effects of insulin and noradrenaline, contributions from glucose and lactate as precursors and comparisons with white adipocytes. *The Biochemical Journal*. 1988;**251**(3):701-709
- [126] Isler D, Hill HP, Meier MK. Glucose metabolism in isolated brown adipocytes under beta-adrenergic stimulation. Quantitative contribution of glucose to total thermogenesis. *The Biochemical Journal*. 1987;**245**(3):789-793
- [127] Nikami H et al. Cold exposure increases glucose utilization and glucose transporter expression in brown adipose tissue. *Biochemical and Biophysical Research Communications*. 1992;**185**(3):1078-1082
- [128] Smith DM et al. Glucose transporter expression and glucose utilization in skeletal muscle and brown adipose tissue during starvation and re-feeding. *The Biochemical Journal*. 1992;**282**(Pt 1):231-235
- [129] Sugden MC, Holness MJ. Physiological modulation of the uptake and fate of glucose in brown adipose tissue. *The Biochemical Journal*. 1993;**295**(Pt 1):171-176
- [130] Hankir MK et al. Dissociation between brown adipose tissue (18) F-FDG uptake and thermogenesis in uncoupling protein 1-deficient mice. *Journal of Nuclear Medicine*. 2017;**58**(7):1100-1103
- [131] Olsen JM et al. beta3-Adrenergically induced glucose uptake in brown adipose tissue is independent of UCP1 presence or activity: Mediation through the mTOR pathway. *Molecular Metabolism*. 2017;**6**(6):611-619
- [132] Weir G et al. Substantial metabolic activity of human brown adipose tissue during warm conditions and cold-induced lipolysis of local triglycerides. *Cell Metabolism*. 2018;**27**(6):1348 e4-1355 e4
- [133] Chakrabarty K, Chaudhuri B, Jeffay H. Glycerokinase activity in human brown adipose tissue. *Journal of Lipid Research*. 1983;**24**(4):381-390
- [134] Kawashita NH et al. Glycerokinase activity in brown adipose tissue: A sympathetic regulation? *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology*. 2002;**282**(4):R1185-R1190
- [135] Festuccia WT et al. Control of glyceroneogenic activity in rat brown adipose tissue. *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology*. 2003;**285**(1):R177-R182

- [136] Sanchez-Gurmaches J et al. Brown fat AKT2 is a cold-induced kinase that stimulates ChREBP-mediated *De novo* lipogenesis to optimize fuel storage and thermogenesis. *Cell Metabolism*. 2018;**27**(1):195 e6-209 e6
- [137] Song Z, Xiaoli AM, Yang F. Regulation and metabolic significance of *De novo* lipogenesis in adipose tissues. *Nutrients*. 2018;**10**(10):1383
- [138] Guilherme A et al. Adipocyte lipid synthesis coupled to neuronal control of thermogenic programming. *Molecular Metabolism*. 2017;**6**(8):781-796
- [139] Farkas V, Kelenyi G, Sandor A. A dramatic accumulation of glycogen in the brown adipose tissue of rats following recovery from cold exposure. *Archives of Biochemistry and Biophysics*. 1999;**365**(1):54-61
- [140] Hurtado del Pozo C et al. ChREBP expression in the liver, adipose tissue and differentiated preadipocytes in human obesity. *Biochimica et Biophysica Acta*. 2011;**1811**(12):1194-1200
- [141] Witte N et al. The glucose sensor ChREBP links *De novo* lipogenesis to PPARgamma activity and adipocyte differentiation. *Endocrinology*. 2015;**156**(11):4008-4019
- [142] Iizuka K et al. Deficiency of carbohydrate response element-binding protein (ChREBP) reduces lipogenesis as well as glycolysis. *Proceedings of the National Academy of Sciences of the United States of America*. 2004;**101**(19):7281-7286
- [143] Herman MA et al. A novel ChREBP isoform in adipose tissue regulates systemic glucose metabolism. *Nature*. 2012;**484**(7394):333-338
- [144] Vijayakumar A et al. Absence of carbohydrate response element binding protein in adipocytes causes systemic insulin resistance and impairs glucose transport. *Cell Reports*. 2017;**21**(4):1021-1035
- [145] Eberle D et al. SREBP transcription factors: Master regulators of lipid homeostasis. *Biochimie*. 2004;**86**(11):839-848
- [146] Osborne TF. Sterol regulatory element-binding proteins (SREBPs): Key regulators of nutritional homeostasis and insulin action. *The Journal of Biological Chemistry*. 2000;**275**(42):32379-32382
- [147] Kim JB, Spiegelman BM. ADD1/SREBP1 promotes adipocyte differentiation and gene expression linked to fatty acid metabolism. *Genes & Development*. 1996;**10**(9):1096-1107
- [148] Horton JD et al. Overexpression of sterol regulatory element-binding protein-1a in mouse adipose tissue produces adipocyte hypertrophy, increased fatty acid secretion, and fatty liver. *The Journal of Biological Chemistry*. 2003;**278**(38):36652-36660
- [149] Shimano H et al. Elevated levels of SREBP-2 and cholesterol synthesis in livers of mice homozygous for a targeted disruption of the SREBP-1 gene. *The Journal of Clinical Investigation*. 1997;**100**(8):2115-2124
- [150] Yahagi N et al. Absence of sterol regulatory element-binding protein-1 (SREBP-1) ameliorates fatty livers but not obesity or insulin resistance in *Lep(ob)/Lep(ob)* mice. *The Journal of Biological Chemistry*. 2002;**277**(22):19353-19357
- [151] Guerra C et al. Emergence of brown adipocytes in white fat in mice is under genetic control. Effects on body weight and adiposity. *The Journal of Clinical Investigation*. 1998;**102**(2):412-420
- [152] Cederberg A et al. FOXC2 is a winged helix gene that counteracts obesity, hypertriglyceridemia, and

diet-induced insulin resistance. *Cell*. 2001;**106**(5):563-573

[153] Kiefer FW et al. Retinaldehyde dehydrogenase 1 regulates a thermogenic program in white adipose tissue. *Nature Medicine*. 2012;**18**(6):918-925

[154] Schulz TJ et al. Brown-fat paucity due to impaired BMP signalling induces compensatory browning of white fat. *Nature*. 2013;**495**(7441):379-383

[155] Orava J et al. Blunted metabolic responses to cold and insulin stimulation in brown adipose tissue of obese humans. *Obesity (Silver Spring)*. 2013;**21**(11):2279-2287

[156] Blondin DP et al. Selective impairment of glucose but not fatty acid or oxidative metabolism in brown adipose tissue of subjects with type 2 diabetes. *Diabetes*. 2015;**64**(7):2388-2397

[157] Au-Yong IT et al. Brown adipose tissue and seasonal variation in humans. *Diabetes*. 2009;**58**(11):2583-2587

[158] Leitner BP et al. Mapping of human brown adipose tissue in lean and obese young men. *Proceedings of the National Academy of Sciences of the United States of America*. 2017;**114**(32):8649-8654

[159] Hanssen MJ et al. Short-term cold acclimation improves insulin sensitivity in patients with type 2 diabetes mellitus. *Nature Medicine*. 2015;**21**(8):863-865

[160] Hoeke G et al. Short-term cooling increases serum triglycerides and small high-density lipoprotein levels in humans. *Journal of Clinical Lipidology*. 2017;**11**(4):920 e2-928 e2