We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Introductory Chapter: Molecular Docking and Molecular Dynamics Techniques to Achieve Rational Drug Design

Amalia Stefaniu

Molecular docking and molecular mechanics simulations are important approaches to achieve a rational drug design or a chemical process modeling. It goes to deep molecular insights as structures and mechanisms helping researchers to characterize various conformations and molecular interactions in terms of energy and binding affinities, giving the possibility to search among dozens, hundreds of real or imaginary compounds, the most suitable for a precise, well-defined purpose. The biochemical purpose derives from the chosen macromolecular target, protein, or enzyme. Starting from a known substance with a known mechanism of action and biological activity, we can imagine other related compounds as drug candidates with better efficacy and fewer side effects. These *in silico* methods help us to identify and select among large compound libraries the most suitable therapeutic agent before even starting its chemical synthesis. That can be called virtual chemistry before reaction tube. It is very convenient, reducing the consumption of chemical reagents, preclinical, clinical trials, and time.

The purpose of this book project is to clearly explain the principles of molecular docking and molecular dynamics, with examples of algorithms and procedures proposed by different software programs for small molecule-protein or protein-protein complexes of medical or materials sciences interest.

Molecular docking studies provide us an overview of type of interactions occurring in ligand (small molecule)-protein or protein-protein complexes and rank the candidate poses by their affinity scoring function.

The concept of molecular recognition of ligand at the protein/enzyme active site, classically named “lock and key,” has been extended at “hand and glove,” considering the protein flexibility and reciprocal adaptability between the receptor and ligand [1].

Molecular dynamics simulations explore extrinsic surface and bulk properties of various forms of pharmaceutically active molecules to aid the selection of a successful candidate. It involves accurate evaluation of binding pathways, kinetics, and thermodynamics of ligands in different solvents.

Both these computer-aided drug design (CADD) methods lead to ligand identification and optimization, favoring rapid development of pharmaceutical compounds.

1. Molecular docking approaches and challenges

Different software algorithms use various approaches such as rigid protein or flexible protein, rigid receptor, soft receptor, flexible side chains, induced fit, or multiple structure algorithms [2].

The steps for conducting molecular docking studies are:

- Ligand preparation consists in generation, optimization, and analysis of its 3D structure. Among multiple conformers, the most stable, as lowest energy, can be used for docking simulations. An aspect to be considered is the fact that in physiological media, the ligand appears ionized. The effect of solvation due to the surrounded water molecules must be solved. The presence of active site water molecules influences the docking pose of the ligand and makes questionable the accuracy of the method [2]. Three-dimensional structures of small ligand molecules are available in virtual databases such as Cambridge Structural Database (CSD), Available Chemical Directory (ACD), MDL Drug Data Report (MDDR), or National Cancer Institute Database (NCI).
- Receptor preparation. The use of a rigid target protein will conduct a single conformation of the receptor. Flexible protein involves different conformations to bind the ligand. Often the site water molecules are removed before performing a docking simulation.

Protein Data Bank (<https://www.rcsb.org/>) provides various solved 3D structures of protein, protein fragments, nucleic acids, and protein-ligand complexes. The assemblies are characterized by X-ray crystallography, nuclear magnetic resonance (NMR), infrared spectroscopy, and or/electron density and are available as PDB files format. This online tool allows us to explore and analyze the structures or to compare any protein in the PDB archive, including support for rigid-body and flexible alignments.

Also, for simulation the optimized ligand structure must be imported and used in the docking software as *.pdb or compatible file.

- Identify the binding site: This step plays a key role in structure-based drug design. It can be determined experimentally or computationally. Some software are created to identifying and analyzing binding sites and predicting receptor druggability [3].
- Dock ligands: Different algorithms are used, fragment-based algorithms, genetic algorithm, Monte Carlo algorithms, and molecular dynamics protocols.
- Docking validation and results analysis: For validation, the software must reproduce the real binding site that was founded and characterized by X-ray crystallography or NMR techniques. To dock ligand similar derivative structures, the same binding site is used, and different conformation dues to rotations around flexible bond are performed for each new structure. The results conduct to predicting preferential binding orientation and the strength of binding affinity, interactions (type, strength, bond length); the conformations are ranked by mean of scoring functions [or root-mean-score deviation (RMSD)]. Furthermore, the stability of receptor-ligand complexes is assessed, and ligand/pharmaceutical small compound druggability is evaluated.

Pharmaceutical applications

- Exploring DNA binding properties of some malignant tumor chemotherapeutic agents [4–7] (to identify the DNA binding site, to predict interactions between potential therapeutic compound and DNA, to assess the stability of DNA-complexes, and to establish correlations between structure and cytotoxicity).

- In silico modeling as attempts to find new efficient therapeutic compounds against pathogens, causative agents of infectious disorders, antitubercular drugs [8–11], antibiotics agents against *Escherichia coli* [12–14], *Pseudomonas aeruginosa* [14–16], *Staphylococcus aureus* [13, 14, 16–19], *Bacillus cereus* [13, 16], *Klebsiella pneumoniae* [13], or others.

The computational findings must be completed and confirmed by biological assays to determine in vitro activity, by measuring minimum inhibitory concentration against tested microorganisms.

Molecular dynamics (MD) simulations are useful approaches when analysis of thermodynamic and kinetic properties of ligand-binding events is required to consider. Besides, MD has become effective tools used to modeling chemical processes and to evaluating different parameters of materials in different media (water or gas): velocity direction of removal of material electrical discharge machining (EDM) [20], indentation [21], wear and friction [22], nano-cutting [23], and laser machining [24].

Author details

Amalia Stefaniu
National Institute for Chemical, Pharmaceutical Research and Development,
Romania

*Address all correspondence to: astefaniu@gmail.com

IntechOpen

© 2019 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] Sledz P, Caflisch A. Protein structure-based drug design: From docking to molecular dynamics. *Current Opinion in Structural Biology*. 2018;**48**:93-102
- [2] Elokely KM, Doerksen RJ. Docking challenge: Protein sampling and molecular docking performance. *Journal of Chemical Information and Modeling*. 2013;**53**(8):1934-1945. DOI: 10.1021/ci400040d
- [3] Halgren TA. Identifying and characterizing binding sites and assessing druggability. *Journal of Chemical Information and Modeling*. 2009;**49**(2):377-389. DOI: 10.1021/ci800324m
- [4] Agarwal S, Chadha D, Mehrotra R. Molecular modeling and spectroscopic studies of semustine binding with DNA and its comparison with lomustine–DNA adduct formation. *Journal of Biomolecular Structure & Dynamics*. 2015;**33**(8):1653-1668
- [5] Subhani S, Jamil K. Molecular docking of chemotherapeutic agents to CYP3A4 in non-small cell lung cancer. *Biomedicine & Pharmacotherapy*. 2015;**73**:65-74. DOI: 10.1016/j.biopha.2015.05.018
- [6] Cathcart J, Pulkoski-Grossa A, Cao J. Targeting matrix metalloproteinases in cancer: Bringing new life to old ideas. *Genes and Diseases*. 2015;**2**(1):26-34
- [7] Pirvu L, Stefaniu A, Neagu G, Albu B, Pintilie L. In vitro cytotoxic and antiproliferative activity of *Cydonia oblonga* flower petals, leaf and fruit pellet ethanolic extracts. Docking simulation of the active flavonoids on anti-apoptotic protein Bcl-2. *Open Chemistry*. 2018;**16**(1):591-604
- [8] Nunn CM, Djordjevic S, Hillas PJ, Nishida C, Ortiz de Montellano PR. The crystal structure of Mycobacterium tuberculosis alkylhydroperoxidase Ahpd, a potential target for antitubercular drug design. *The Journal of Biological Chemistry*. 2002;**277**:20033-20040. DOI: 10.1074/jbc.M200864200
- [9] Salunke SB, Azad AK, Kapuriya NP, Balada-Llasat JM, Pancholi P, Schlesinger LS, et al. Design and synthesis of novel anti-tuberculosis agents from the celecoxib pharmacophore. *Bioorganic & Medicinal Chemistry*. 2015;**23**(9):1935-1943. DOI: 10.1016/j.bmc.2015.03.041
- [10] Tiwari R, Möllmann U, Cho S, Franzblau SG, Miller PA, Miller MJ. Design and syntheses of anti-tuberculosis agents inspired by BTZ043 using a scaffold simplification strategy. *ACS Medicinal Chemistry Letters*. 2014;**5**(5):587-591. DOI: 10.1021/ml500039g
- [11] Umesiri FE, Lick A, Fricke C, Nathaniel TI. Boronic-aurone derivatives as anti-tubercular agents: Design, synthesis and biological evaluation. *Medicinal Chemistry*. 2015;**5**:437-441. DOI: 10.4172/2161-0444.1000297
- [12] Cheng K, Zheng Q-Z, Qian Y, Shi L, Zhao J, Zhu H-L. Synthesis, antibacterial activities and molecular docking studies of peptide and Schiff bases as targeted antibiotics. *Bioorganic & Medicinal Chemistry*. 2009;**17**(23):7861-7871
- [13] Gullapelli K, Brahmeshwari G, Ravichander M, Kusuma U. Synthesis, antibacterial and molecular docking studies of new benzimidazole derivative. *EJBAS*. 2017;**4**(4):303-309
- [14] Pintilie L, Stefaniu A, Nicu AI, Maganu M, Caproiu MT. Design, synthesis and docking studies of some novel fluoroquinolone compounds with antibacterial activity. *Revista de Chimie*. 2018;**69**(4):815-822

- [15] El-Attar MAZ, Elbayaa RY, Shaaban OG, Habib NS, Wahab AEA, Abdelwahab IA, et al. Design, synthesis, antibacterial evaluation and molecular docking studies of some new quinoxaline derivatives targeting dihydropyrimidinase synthase enzyme. *Bioorganic Chemistry*. 2018;**76**:437-448
- [16] Srivastava R, Gupta SK, Naaz F, Singh A, Singh VK, Verma R, et al. Synthesis, antibacterial activity, synergistic effect, cytotoxicity, docking and molecular dynamics of benzimidazole analogues. *Computational Biology and Chemistry*. 2018;**76**:1-16. DOI: 10.1016/j.compbiolchem.2018.05.021
- [17] Barakat A, Al-Majid AM, Al-Qahtany BM, Ali M, Teleb M, Al-Agamy MH, et al. Synthesis, antimicrobial activity, pharmacophore modeling and molecular docking studies of new pyrazole-dimedone hybrid architectures. *Chemistry Central Journal*. 2018;**12**(1):29-42. DOI: 10.1186/s13065-018-0399-0
- [18] Pintilie L, Stefaniu A, Nicu AI, Caproiu MT, Maganu M. Synthesis, antimicrobial activity and docking studies of novel 8-chloro-quinolones. *Revista de Chimie*. 2016;**67**(3):438-445
- [19] Pintilie L, Stefaniu A, Nicu AI, Negut C, Tanase C, Caproiu MT. Design, synthesis and molecular docking of some oxazolidinone compounds. *Revista de Chimie*. 2018;**69**(11):2981-2986
- [20] Yue X, Yang X. Study on the distribution of removal material of EDM in deionized water and gas with molecular dynamics simulation. 18th CIRP Conference on Electro Physical and Chemical Machining (ISEM XVIII) *Procedia CIRP* 42. 2016. 691-696
- [21] Cheong WCD, Zhang LC. Molecular dynamics simulation of phase transformations in silicon monocrystals due to nano-indentation. *Nanotechnology*. 2000;**11**:173-180
- [22] Maekawa K, Itoh A. Friction and tool wear in nano-scale machining-a molecular dynamics approach. *Wear*. 1995;**188**(1-2):115-122
- [23] Zhang L, Zhao HW, Dai YYH, Du XC, Tang PY, et al. Molecular dynamics simulation of deformation accumulation in repeated nanometric cutting on single-crystal copper. *RSC Advances*. 2015;**5**:12678-12685
- [24] Shih C-Y, Shugaev MV, Wu C, Zhigilei LV. Generation of subsurface voids, incubation effect, and formation of nanoparticles in short pulse laser interactions with bulk metal targets in liquid: Molecular dynamics study. *Journal of Physical Chemistry C*. 2017;**121**(30):16549-16567. DOI: 10.1021/acs.jpcc.7b02301