We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Antioxidant and Biological Activity of *Cissus sicyoides* and *Rosmarinus officinalis* Extracts

Marielba de los Angeles Rodriguez Salazar,
Glides Rafael Olivo Urbina, Priscila do Nascimento Bezerra,
Vânia Maria Borges Cunha, Marcilene Paiva da Silva,
Flávia Cristina Seabra Pires, Ana Paula de Souza e Silva,
Sérgio Henrique Brabo de Sousa
and Raul Nunes de Carvalho Jr

Abstract

This chapter will describe the antioxidant and biological activity of *Cissus sicyoides* and *Rosmarinus officinalis* leaf extracts, which represent an important natural source of antioxidants. These plants contain several bioactive compounds with high antioxidant activity, such as phenolic compounds, which are compounds that prevent or delay oxidative stress, acting as free radical scavengers (FRSs), and thus reduce the onset of cardiovascular disease, cancer, diabetes, epilepsy, stroke, among other diseases. The supercritical fluid extraction (SFE) has been studied to obtain antioxidant compounds from natural sources, without the drawbacks associated with conventional extraction processes, such as the use of organic solvents, which present toxicity and contaminate the extracts, is proposed.

Keywords: *C. sicyoides*, *R. officinalis*, antioxidant activity, biological activity, supercritical extraction

1. Introduction

The Amazonian biodiversity presents a great source of foods and medicinal plants rich in antioxidant compounds whose study and conscious exploration contribute to the region sustainable development [1, 2]. The plants have a great importance due to their medicinal and nutritional properties. About 70–90% of the world population prefers the use of medicinal plants or plant extracts to treat common diseases [3, 4]. Plants have been extensively studied in recent years for their antioxidant activity. The main classes of plant chemicals are phenolic compounds, tocopherols, carotenoids, and alkaloids. Among these compounds, phenolic compounds are the most important. They prevent or delay oxidative stress, acting as free radical scavengers (FRSs), and thus reduce the onset of different chronic diseases [5–8].

Antioxidants are a set of substances that can delay or inhibit oxidation reactions and act as a defense mechanism to neutralize the harmful effects of oxidation in biological systems and foods [6, 9, 10]. Oxidative stress is considered a state of imbalance where excessive amounts of reactive oxygen and nitrogen species (ROS/RNS, for example, superoxide anion, hydrogen peroxide, hydroxyl radical, peroxynitrite) exceed the capacity of endogenous antioxidants (uric acid, superoxide dismutase, catalase, glutathione peroxidase), leading to the oxidation of a biomacromolecule variety such as enzymes, proteins, DNA, and lipids. Exogenous antioxidants (phenolic compounds, carotenoids, tocopherols, and ascorbates) are consumed in the diet mainly of fruits, leaves, seeds, vegetables, and cereals, they have the function of increasing or protecting the antioxidant defense in biological systems and, therefore, they are important for endogenous oxidative stability [11–13].

It is conflicting that oxygen and nitrogen, considered essential for biological processes, are also cofactors for toxic and degenerative processes. In this sense, the antioxidant compounds act through different chemical mechanisms in order to minimize or maintain redox balance in vivo [9, 14]. There are several mechanisms by which oxidation can be inhibited. In general, the mechanisms involved include FRSs, ester bond enzymatic hydrolysis, transition metal ion sequestration, and enzyme-catalyzed peroxide reduction. The last three mechanisms mentioned do not cease reactive species action, but prevent the formation of molecules capable of promoting free radical chain reactions [15].

There is a growing interest in new sources of natural antioxidant compounds due to synthetic antioxidants such as butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), and tert-butylhydroquinone (TBHQ) in the food industry being severely restricted, since they may show carcinogenic effects on living organisms [16–18]. In this sense, the scientific community and consumers are looking for new bioactive compounds of natural origin that can be used to develop new treatments against diseases. In addition, they may be employed in the food industry as functional food ingredients.

Cissus sicyoides L., which belongs to the Vitaceae family, is also known as vegetal insulin, anil-trepador, bejuco-caro, cipó-pucá, or puci. It is considered a plant from the Neotropical region and is usually found in the Amazon region [19, 20]. According to studies on *C. sicyoides* extract composition, the presence of bioactive compounds with high antioxidant activity as carotenoids and phenolic compounds (flavonoid, resveratrol, coumarins, and tannins) was found [21–23]. Therefore, it is a plant traditionally used by Brazilian popular medicine to treat rheumatism, epilepsy, stroke, abscesses, arthritis, and diabetes [23, 24].

Rosmarinus officinalis is an aromatic plant of the Lamiaceae family, native to the Mediterranean region. Today, it has been grown in many parts of the world and is known as rosemary [25, 26]. It has been recognized as one of the plants with great antioxidant activity. Among the most effective antioxidant constituents, cyclic diterpene diphenols, carnosic acid, rosmarinic acid, and carnosol have been identified. *R. officinalis* extracts have been used in the treatment and/or prevention of diseases such as cancer, Alzheimer's disease, urinary and gastrointestinal infections, diabetes, ischemia, and atherosclerosis [17, 25–32]. *R. officinalis* extract has been commercially exploited as a natural antioxidant [5, 16].

Supercritical fluid extraction (SFE) has already been studied to obtain bioactive compounds from natural sources. Salazar et al. and Carvalho et al. showed that the application of SFE technology is successful in obtaining extracts from *C. sicyoides* and *R. officinalis*, respectively, with high antioxidant capacity [24, 33]. SFE is based on the use of solvents with temperatures and pressures above their critical points, which have a high solvency power. One of the most commonly used solvents in SFE is carbon dioxide (CO₂) since its critical points are moderate, nontoxic,

non-flammable, affordable, chemically inert, and apolar and has an ideal behavior for thermosensitive compound extraction. In SFE it is possible to obtain totally solvent-free extracts without compound thermal degradation due to the low operating temperatures applied [34–37].

Supercritical CO₂ (Sc-CO₂) has a limitation in dissolving polar molecules. However, this disadvantage can be solved by polar solvent addition, called modifiers or cosolvents, which modify the supercritical fluid polarity and, consequently, improve polar fraction extraction rich in bioactive substances, such as phenolic compounds related to high antioxidant activity [37, 38]. The aim of this chapter is to describe the antioxidant and biological activity of *Cissus sicyoides* and *Rosmarinus officinalis* leaf extracts that represent an important natural source of antioxidants. In addition to providing an overview of the SFE that is currently presented as a modern and environmentally safe extraction technology for antioxidant compound extraction.

2. *Cissus sicyoides*

2.1 Botanical description

Cissus sicyoides L. or *Cissus verticillata* L., which belongs to the Vitaceae family, is also known as vegetal insulin, cipó-pucá, or puci. It is considered a plant of the Neotropical region and is usually found in the Amazon region. It is a climbing plant, which can reach up to 6 m in length, and presents fleshy articulated branch, alternating leaf of ovate format, pale or yellowish-green flowers, and round fruit, with variations of color from violet to black (**Figure 1**) [19, 20].

2.2 Chemical composition

The bioactive compounds present in the leaf and stem are represented by carotenoids (α -carotene and β -carotene) [39] and phenolic compounds such as

Figure 1.
 Description of *C. sicyoides* parts and the main chemical structure of the antioxidant compounds.

flavonoids (quercetin 3-O-rhamnoside and kaempferol 3-O-rhamnoside) [21]. But also, three new flavonoid glycosides were found, denominate cissosides I, II, and III (kaempferol 3-O- α -L-(5''-O-acetyl)-arabinofuranosyl-7-O- α -L-rhamnopyranoside, quercetin 3-O- α -L-arabinofuranosyl-7-O- α -L-rhamnopyranoside, and quercetin 3-O- α -L-(5''-O-acetyl)-arabinofuranosyl-7-O- α -L-rhamnopyranoside) [22].

Recently, three different flavonoids were identified (quercetin-3-O-hexoside, quercetin-3-O-deoxyhexoside, and kaempferol-3-O-deoxyhexoside) [23]. In addition, resveratrol (3,5,4'-trihydroxystilbene) [40] and a new benzofuran-type stilbene (cissusin) [22], tannins, coumarins (glycoside 5,6,7,8-tetrahydroxycoumarin-5 β -xylopyranoside and sabandin), and steroids (β -sitosterol and 3 β -O- β -D-glucopyranosyl sitosterol) were found [21]. The presence of essential oils was also detected [41]. In the supercritical extracts of *C. sicyoides* phytochemical screening obtained from leaves and stems by high-performance thin-layer chromatography (HPTLC), the presence of terpenes, phenolic compounds, and flavonoids was evidenced [24]. In the fruit composition analysis, three anthocyanins were found (delphinidin-3-rutinoside, cyanidin-3-rhamnosyl-arabinoside, and delphinidin-3-rhamnoside) [42]. Therefore, the fruit of this plant may have potential use as a food coloring. **Figure 1** shows the chemical structures of the main antioxidant compounds found in *C. sicyoides*.

2.3 Antioxidant and biological activity

In *C. sicyoides* polyphenolic profile, we can find flavonoids: the quercetin and kaempferol as the majority and its various isomers. The mechanism of quercetin antioxidant action has been associated mainly with the reduction of ROS/RNS, which is a compound that prevents or retards oxidative stress, which enables the prevention of various chronic diseases [6, 43]. In the study conducted by Crespo et al., treatment with quercetin and kaempferol prevented the production of ROS such as peroxides, superoxide anion, and nitric oxide. These results confirmed the differential protective effect of these flavonoids in the diet against oxidative stress induced by pro-inflammatory stimuli in parenchymal liver cells [44]. Resveratrol plays an important antioxidant role in reducing hydroxyl radicals, superoxide, and metal-induced radicals, as well as showing antioxidant abilities in ROS-producing cells. Also, it has a protective effect against lipid peroxidation in cell membranes and DNA damage caused by ROS/RNS [45]. Benzofuran is a potent radical scavenger capable of inhibiting lipid peroxidation, its FRS capacity being greater than α -tocopherol [46].

Figure 2 shows the results of the qualitative analysis of antioxidant activity by HPTLC of *C. sicyoides* extracts obtained by supercritical extraction (essays 1–15), hexane extract (HE), and ethanolic extract (EE) obtained with conventional extraction by Soxhlet. The plaque was derivatized with DPPH (2,2-diphenyl-1-picrylhydrazyl), and it was possible to detect the presence of yellow spots on the plaque purple bottom resulting from the reduction of the DPPH $\bullet$; in the presence of antioxidant substances, 2,2-diphenyl-picryl-hydrazine is reduced, losing its purple coloration. The study results confirmed the presence of chemical constituent characteristic of this plant, with antioxidant activity. In the same study, a quantitative determination of the antioxidant activity by the DPPH method was carried out. It was demonstrated that with the extractive methodologies (SFE and Soxhlet) used it was possible to extract with low EC₅₀ values, related to a high antioxidant activity; for EE, the value of EC₅₀ (325.67 g of extract/g of DPPH) is similar to the value of EC₅₀ (404.81 g of extract/g of DPPH) obtained with SFE [24].

In the in vitro antioxidant activity determination by the ABTS method of the *C. sicyoides* aqueous extract obtained by decoction, it was evidenced that the extract has an antioxidant activity of IC₅₀ = 13.0 $\pm$ 0.2 μ g/ml. These results indicate that

Figure 2.
 Qualitative analysis of antioxidant activity by HPTLC of *C. sicyoides* extracts obtained by supercritical extraction and conventional extraction (Soxhlet) [29].

the extract is a potential source of natural antioxidant and may be useful in the prevention of diseases associated with oxidative stress [47]. Thus, the antioxidant activity results of *C. sicyoides* extracts are related to the extraction methods and to the solvent used.

Due to antioxidant properties, *C. sicyoides* has been used by folk medicine to treat rheumatism, epilepsy, stroke, abscesses, arthritis, and diabetes; it has also been used to treat respiratory diseases. Some biological activities are attributed to the plant as anti-inflammatory, antirheumatic, antiepileptic, antihypertensive, antimicrobial, antipyretic, antioxidant, antiallergic, anticancer, and antidiabetic activities [23, 40, 48–50].

Several studies point to the application of *C. sicyoides* in the treatment of various diseases, in order to demonstrate the cytotoxic activity of the *C. sicyoides* aqueous extract obtained by decoction against cells human epidermoid carcinoma no. 2 (HEp-2 cells), showing complete inhibition of cell division after 24 h of treatment [51]. Also, the antitumor activity of *C. sicyoides* hydroalcoholic extract obtained by maceration, in animals at doses of 300 and 600 mg/kg in weight, was investigated, being demonstrated that the extract showed an inhibition of tumor activity in sarcoma-180 of 49 and 62% and Ehrlich carcinoma of 69 and 84% [52].

Regarding the plant anti-inflammatory activity, it has been demonstrated that the oral administration of 300 and 500 mg/kg of the *C. sicyoides* stem's aqueous extract obtained by decoction in mice with induced edema has a potent anti-inflammatory activity, and administration of the extract produced an approximately 50% reduction of the induced edema [53]. Resveratrol was indicated as one of the constituents responsible for the anti-inflammatory and antiallergic properties presented by *C. sicyoides* alcoholic extract obtained by maceration [40]. However, modern ethnopharmacological use reports that the *C. sicyoides* hydroalcoholic extract by percolation has anti-inflammatory and antidiarrheal actions, due to the abundant presence of flavonol-O-glycoside derivatives of quercetin and kaempferol, which are mainly responsible for the plant pharmacological effects [23].

In addition, pharmacological effects were detected in the treatment and/or prevention of dysfunctions such as hypertension and vasoconstriction of arteries, veins, and capillaries with aqueous extract of *C. sicyoides*. These compounds act at the membrane level, increasing the calcium entry through the membrane as well as acting on the internal calcium deposits [54].

In the evaluation of the antidiabetic potential of *C. sicyoides*, the effects of leaf tea from the plant were studied; the *in vivo* experimental model chosen proved to be an appropriate treatment, reducing blood and urine glucose levels [48]. Later, it

was demonstrated that treatment of diabetic rats with *C. sicyoides* aqueous extract obtained by decoction, for 7 days (100 and 200 mg/kg), reduced blood glucose levels by 22 and 25%, respectively [49]. However, *C. sicyoides* leaf tea was used to investigate the plant therapeutic efficacy in volunteers who are diabetic and intolerant to glucose. A single dose of tea (1 g of dried leaf powder in 150 ml of water) was used for a period of 7 days. It was observed in people intolerant to glucose that the tea had antidiabetic activity [55].

C. sicyoides has antibacterial activity, showing inhibitory capacity against bacteria that cause food poisoning [56], which causes acute effects in the gastrointestinal tract and, in some cases, a high severity that patients come to death (*Bacillus cereus*, *Bacillus subtilis*, *Bacillus megaterium*, *Staphylococcus aureus*, and *Escherichia coli*). In addition, the antifungal activity of plant leaf and stem alcoholic extracts was demonstrated, inhibiting the growth of fungi *Cladosporium sphaerospermum* and *Cladosporium cladosporioides* [57].

Recently, Salazar et al. carried out the biological activity determination of *C. sicyoides* supercritical extract; an in vivo test using a focal cerebral ischemia model was performed, and the extract had shown to have a neuroprotective and anti-inflammatory effect, justifying the use in traditional folk medicine for central nervous system diseases. These effects were associated to the presence of phenolic compounds in the extract [24]. Therefore, the results of these studies justify the traditional use of *C. sicyoides*, pointing to the plant extract potential benefit as a possible alternative medicine in disease treatment.

3. *Rosmarinus officinalis*

3.1 Botanical description

Rosmarinus officinalis is an aromatic plant of the Lamiaceae family, native to the Mediterranean region, and is also cultivated in Central Asia, India, Southeast Asia, South Africa, Australia, the United States, and Brazil. Today, it has been grown in many parts of the world and is commonly known as rosemary. The plant is a bush that reaches from 0.50 to 1.50 m in height, with very pungent aroma leaves and blue, violet, and white flowers (**Figure 3**) [25, 26, 58].

3.2 Chemical composition

R. officinalis chemical composition varies greatly, due to some factors that directly influence the quality, amount of oil, and extract produced. However, it is possible to verify, through the literature, that its main chemical constituents are flavones, diterpenes, steroids, and triterpenes [17, 26, 27, 31]. The phenolic compounds present in *R. officinalis* were grouped into three classes: (i) phenolic acids (vanillic, caffeic, ferulic, and rosmarinic acids), (ii) diterpenes (carnosol, rosmadial, carnosic acid, methyl carbonate, rosmanol, epirosmanol, epiisorosmanol, epirosmanol methyl ether, and epiisorosmanol ethyl ether), and (iii) flavonoids (hesperetin, apigenin, genkwanin, 4'-methoxytecto-chrysin, cirsimaritin, scutellarein, 4'',5,7,8-tetrahydroxyflavone, homoplantagin, and 6-hydroxyluteolin 7-glucoside) [27]. Recently, a *R. officinalis* chromatographic analysis was carried out, which revealed two large groups: oxygenated monoterpenes and hydrocarbonated monoterpenes. The main constituents of these groups were 1,8-cineole followed by camphor, borneol, and α - and β -pinene. The oxygenated and hydrocarbonated sesquiterpenes were composed of caryophyllene and caryophyllene oxide [26]. In the supercritical extracts of *R. officinalis* leaves, chemical analysis confirmed the

Figure 3.
 Description of *R. officinalis* parts and the main chemical structure of the antioxidant compounds.

presence of 1,8-cineole, camphor, carnosic acid, and rosmarinic acid [17, 33, 59].

Figure 3 shows the chemical structures of the main antioxidant compounds found in *R. officinalis*.

3.3 Antioxidant and biological activity

R. officinalis has been recognized as one of the plants with high antioxidant activity [28–30]. Its antioxidant effect is due to the phenolic compounds present in the leaves and stems [60]. Among the most effective antioxidant constituents, cyclic diterpene diphenols, carnosic acid, and carnosol were identified. In addition, its extract contains epirosmanol, rosmanol, metilcarnosato, isorosmanol, and other caffeic acid derivatives [61, 62].

The action mechanism of these compounds has been widely discussed in several studies. The carnosic acid and carnosol act as potent sequestrants of peroxy radicals and are responsible for 90% of the antioxidant properties, where both are inhibitors of lipid peroxidation in liposomal and microsomal systems, besides being good sequestrants of hydroxyl radicals. Specifically, carnosic acid removes hydrogen peroxide but may also act as a substrate for its ability to increase or maintain the superoxide dismutase and glutathione peroxidase activities. The most important elements in the *R. officinalis* structure are the diterpenes containing the aromatic ring (C₁₁–C₁₂) in the catechol group, with the conjugation of three basic rings. The

catechol group is responsible for eliminating the radical electrons formed as an oxidation result. Lactone carnosol, rosmarinic acid, and hesperetin were cited in the literature as important FRSs [31, 63–65]. Rosmarinic acid has two aromatic rings, each with two OH groups that are capable of donating hydrogen and chelating metals [66]. Caffeic acid derivatives may act as metal ion chelators (Fe^{2+}), thus reducing the formation of ROS [67].

The antioxidant of *R. officinalis* extracts obtained in SFE was confirmed. Carvalho et al. analyzed the plant antioxidant activity through a coupled reaction of β -carotene and linoleic acid; the results indicated that the extracts obtained at high pressures and low temperature (300 bar/40°C) exhibited the highest antioxidant activities, in comparison to extracts obtained in low pressures (150 bar/30°C). In any case, antioxidant activities were always above the control used (β -carotene and linolenic acid) as shown in **Figure 4**. The authors state that the antioxidant action remained approximately constant for the 3-h reaction for all extracts tested. The major compounds detected in the extracts were camphor (0.6% d.b.) and 1,8-cineol (0.043% d.b.). The extract obtained by hydrodistillation showed the highest yield of camphor (1.22% d.b.) and 1,8 cineol (0.23% d.b.) compared to other extraction methods (SFE and Soxhlet) [33].

Thus, the antioxidant capacity of *R. officinalis* leaf and stem extract obtained with supercritical CO_2 was also evaluated by the ORAC method. The extract showed a high antioxidant capacity (1.9 mol Trolox/mg) similar to that of BHT and vitamin E (2.8–3.0 mol Trolox/mg). In addition, the extract presented a high percentage of lipid oxidation inhibition (88%) of fatty acids present in an analyzed cosmetic foundation. The extract volatile fraction was characterized by compounds such as camphor, 1,8-cineol, and trans- β -caryophyllene present in relative amounts of less than 25% [59].

Due to antioxidant properties, *R. officinalis* has been used in food preservation and in disease treatment. In food preservation, components such as rosmanol and carnosol prevent oxidation and microbial contamination and also can be up to four times more effective than BHA and equal to BHT as an antioxidant [27, 68, 69]. Different studies have demonstrated the potent activity of *R. officinalis* in inhibiting the formation of hydroperoxides, reducing carotenoid color loss, and retarding lipid oxidation in corn [78] and hazelnut oils [70].

The *R. officinalis* extract has been successfully commercially exploited as a natural antioxidant, for its use as synthetic antioxidants such as BHA, BHT, and TBHQ, in the food industry and is severely restricted as they may have carcinogenic effects on living organisms [5, 16, 17]. In this way, *R. officinalis* extract may be useful to

Figure 4. Antioxidant activity of *R. officinalis* extracts obtained with supercritical CO_2 [38].

replace or even decrease the synthetic antioxidants in foods. As preservatives, the extracts offer several technological advantages and benefits to consumers [71].

Health problems derived from lipid oxidation have attracted consumers' and researchers' attention, since numerous diseases are linked to dietary and biological lipid oxidation products. Therefore, *R. officinalis* extracts have been related to several biological activities, such as anticancer, antidiuretic, anti-inflammatory, antibacterial, antidiabetic, antiangiogenic, antioxidant, and hepatoprotective [28, 71–74], besides allowing the use of the plant in the treatment and/or prevention of Alzheimer's disease, urinary and gastrointestinal infections, diabetes, aging, ischemia, and atherosclerosis [25, 32].

Among the most important groups of compounds isolated from the plant, phenolic diterpenes account most of their biological activity. These compounds have been indicated in recent years as inhibitors of neuronal cell-induced death by a variety of agents both in vitro and in vivo, confirming the therapeutic potential of these compounds for Alzheimer's disease, due to the compounds multifunctional nature in the neuronal protection mediated by the plant antioxidant activity [32].

Several studies show that *R. officinalis* has pharmacological activity for chemoprevention and cancer therapy. In the extract antiproliferative activity evaluation against human ovarian cancer cells, it was corroborated that the extract inhibited the proliferation of cancer cell lines, affecting the cell cycle in multiple phases. In addition, it induced apoptosis by modifying the multiple gene expression that regulates apoptosis. Thus, the extract can be considered as an adjuvant to chemotherapy [62]. Also, the antiangiogenic effect of the carnosic acid present in *R. officinalis* extract was corroborated in angiogenesis models using human umbilical vein endothelial cells in relation to the tube formation in the reconstituted basement membrane, chemotaxis, and proliferation. Carnosic acid from the extract may be useful in preventing disorders due to angiogenesis, and its antiangiogenic effect may contribute to a neuroprotective effect [72].

The actions of *R. officinalis* leaf ethanolic extract obtained with Soxhlet extraction were tested in glucose homeostasis and antioxidant defense in rabbits. Serum levels, glucose levels and insulin levels were studied in diabetic rabbits (alloxan was used to induce diabetes); it was shown that at a dose of 200 mg/kg was possible to reduce the blood glucose level and to increase the serum insulin concentration. In addition, during 1 week of animal treatment with the extract, it was demonstrated that it had an ability to inhibit lipid peroxidation and to activate the antioxidant enzymes. Due to its potent antioxidant properties, the plant extract has a remarkable antidiabetogenic effect [75]. Recently, the antidiabetic and antihypercholesterolemic action of flavonoid-rich fractions of *R. officinalis* (fractions obtained with n-butanol and diethyl ether) in diabetic mice induced by streptozotocin was evaluated. Both fractions showed a decrease in the glucose level at a dose of 400 mg/kg, especially the fraction obtained with diethyl ether; plasma glucose levels decreased up to 60.38%. The pancreas histopathological study showed that both fractions regenerated the pancreatic β cells and increased the mass of islets. *R. officinalis* fractions exhibited a potent antidiabetic effect, while the fraction obtained with n-butanol showed a high anti-hypercholesterolemic activity [76].

The anti-inflammatory activity of *R. officinalis* supercritical extracts was studied. Absorptions of extract fractions were tested on monolayers of Caco-2 cells (2–12 h of incubation). Human macrophages were treated with basolateral fractions, and TNF- α , IL-1 β , IL-6, and IL-10 secretions were measured by ELISA. Fractions obtained after 8 and 12 h in absorption experiments caused a considerable reduction in the excretion of pro-inflammatory cytokines. This reduction in cytokine secretion levels was associated with the amounts of carnosol and carnosic acid. Thus, the *R. officinalis* supercritical extract can be used in formulations to inflammatory disease prevention [77].

In relation to the antibacterial activity, *R. officinalis* essential oils obtained by hydrodistillation exhibited antibacterial activity against *Escherichia coli*, *Salmonella typhi*, *S. enteritidis*, and *Shigella sonnei*; this activity was associated with the oil ability to reduce DPPH radical formation ($CI_{50} = 3.82 \mu\text{g/ml}$) [61]. However, the antibacterial and antifungal activities of *R. officinalis* leaf extracts obtained by SFE extraction were confirmed, and the extracts showed antibacterial activity against Gram-positive bacteria (*Staphylococcus aureus* and *Bacillus cereus*) and Gram-negative bacteria (*Escherichia coli* and *Pseudomonas aeruginosa*) and antifungals against *Candida albicans*. Obtaining *R. officinalis* extracts by SFE has been shown to be a promising extraction with respect to its incorporation into various foods, cosmetics, and pharmaceuticals products that a natural aroma, color, and antioxidant/antimicrobial additive are desired. These properties are also necessary for the food industry in order to find possible alternatives to synthetic preservatives [17].

4. Supercritical fluid extraction (SFE) of antioxidant compounds from plant matrices

When a new extract from a natural source is tested, the most important aspects to take into account are the extraction method and the type of solvent used, as this will affect the antioxidant properties. Several extraction methods for the selective extraction from plant matrices such as *R. officinalis* were identified in the scientific literature [71].

Thus, the bioactive compound extraction has been considered one of the most important steps in the approach of obtaining or recovering bioactive compounds. Conventional extractions have been the most used technology for these compound recovery. It is based on the extraction power of different solvents and the application of high temperatures, promoting mass transfer. However, there are drawbacks associated with conventional extraction processes such as the use of large amounts of organic solvents, toxic to human health and the environment, extraction time, and the use of high temperatures that can degrade the thermosensitive compounds [6, 8]. They motivated the search for environmentally safe extraction techniques such as microwave-assisted extraction (MAE), ultrasonic-assisted extraction (UAE), pressurized liquid extraction (PLE), and supercritical fluid extraction (SFE) [36, 78]. SFE has already been studied to obtain antioxidant compounds from natural sources [24, 33, 77, 79]. **Table 1** presents the antioxidant activity values of different plant extracts obtained with SFE, involving the plants under study (*C. sicyoides* and *R. officinalis*).

4.1 SFE procedure

A solvent is considered a supercritical fluid when the pressure and temperature of the system are above its critical point. This point is defined as the highest temperature and pressure at which a substance can exist in equilibrium between the liquid and vapor phases. Above its critical temperature (T_c) and critical pressure (P_c), the supercritical fluid can be considered as an expanded liquid or as a compressed gas, whose density (ρ) is relatively high and consequently has a high solvency power. This effect gives the solvent a certain degree of selectivity, in addition to allowing easy separation of the solvent from the solute, which can be achieved by a simple system depressurizing, resulting in products totally solvent-free and without thermal degradation of the compounds of interest, due to low operating temperatures [35, 38].

Plants	Extraction conditions	Solvents	Method of determination	Antioxidant capacity	Biological Activity	Refs.
<i>C. sicyoides</i>	40°C/400 bar	CO ₂ + 10% of ethanol	DPPH	404.81 ± 2.78 EC ₅₀ : g of extract/g of DPPH	Neuroprotective and anti-inflammatory effect	[29]
<i>R. officinalis</i>	40°C/300 bar	CO ₂	DPPH	12.85 ± 0.10 IC ₅₀ : µg.ml ⁻¹	Antioxidant, antibacterial, and antifungal	[21]
	100°C/350 bar	CO ₂	DPPH	0.23 ± 0.01 IC ₅₀ : mg/ml	Antioxidant	[84]
	50°C/300 bar	CO ₂	ORAC	1.9 ± 0.10 µmol Trolox/mg extract	Antioxidant	[59]
<i>Mangifera indica</i> L.	55°C/100 bar	CO ₂ + 20% of ethanol	DPPH	2.13 ± 0.24 EC ₅₀ : DPPH µg/µg dry extract	Antioxidant	[85]
<i>Eugenia uniflora</i> L.	60°C/400 bar	CO ₂ + ethanol	DPPH	>200 EC ₅₀ (µg/ml)	Antioxidant	[86]
<i>Raphanus sativus</i> L.	35°C/400 bar	CO ₂	DPPH	359 mg TE/100 g dry extract	Antioxidant and anti-inflammatory	[37]
<i>Piper nigrum</i> L	40°C/300 bar	CO ₂	DPPH	103.28 EC ₅₀ : of µg.ml ⁻¹	Antioxidant	[87]

Table 1.
Presentation of the antioxidant activity values of different plant extracts obtained with SFE, involving the plants under study (*C. sicyoides* and *R. officinalis*).

One of the most commonly used solvents in SFE is carbon dioxide (CO₂) because its critical points are moderate (T_c = 31.1°C, P_c = 73.8 bar, and critical density (ρ_c) = 0.468 g/cm³), nontoxic, non-flammable, affordable, chemically inert, and apolar and has an ideal behavior for thermosensitive compound extraction [34, 37]. In addition to the supercritical CO₂ (Sc-CO₂), there are other substances that are also used as supercritical fluids, as shown in **Table 2**.

Due to its low polarity, Sc-CO₂ presents a limitation to dissolve polar molecules. However, this disadvantage can be solved by the addition of polar solvents, called modifiers or cosolvents, which modify the supercritical fluid polarity and, consequently, improve the extraction of polar fractions rich in bioactive substances, such as phenol compounds related to high antioxidant activity [37, 38]. Methanol is the solvent most used as a modifier for various plant matrices, but it is toxic and

Fluid	T _c (°C)	P _c (bar)	ρ _c (g/cm ³)
Nitrous oxide (N ₂ O)	36.5	71.0	0.457
Ethane (C ₂ H ₆)	32.2	48.8	0.203
Propane (C ₃ H ₈)	96.7	42.5	0.220
Propylene (C ₃ H ₆)	91.9	46.2	0.230
Benzene (C ₆ H ₆)	289.0	48.9	0.302
Toluene (C ₇ H ₈)	318.6	41.1	0.290
Ammonia (NH ₃)	132.5	112.8	0.240
Water (H ₂ O)	374.2	220.5	0.272

Table 2.
Critical properties of some substances used as solvents in supercritical extraction processes.

Figure 5.
Scheme of the SFE procedure of plant matrices.

different from ethanol, which is an environmentally safe solvent being a good choice for SFE processes, and can be used in the extraction of natural products [80, 81]. Water is also a very attractive cosolvent for natural product extraction due to its high polarity, which considerably increases the polarity of Sc-CO₂ [79].

For antioxidant compound extraction and recovery by SFE, several vegetable matrices were used, such as seeds, fruits, leaves, flowers, rhizomes, roots, fruit peels, and tree branches. The SFE process consists basically in the extraction of soluble compounds present in the solid matrix by a supercritical solvent and then separates these compounds from the solvent after depressurizing the system. In order to achieve an efficient and adequate extraction, several factors must be taken into account, having a careful control of the operating conditions and process step optimization [35, 36, 82].

Initially, the raw material must pass through a pretreatment stage before being fed into the fixed bed extractor; this procedure is performed to prepare the solid particles, allowing a greater efficiency to be achieved in the extraction process [83]. As shown in **Figure 5**, after the raw material is collected, one of the first stages of its pretreatment is the solid matrix moisture reduction, for example, drying leaves in an oven with air circulation. Generally, the plant matrix moisture should not exceed 14% (wet basis). Another important step is the moisture content determination by the distillation method of the Jacobs immiscible solvent, with the purpose of knowing if the quantity of water in the sample is adequate for the supercritical extraction process. The sieving stage is applied to standardize and determine the average particle size of the solid particles. The real and apparent density and bed porosity determination is also very important as they affect the particles packaging in the extraction vessel and consequently the solvent flow and the mass and heat transfer processes [35, 82].

After a suitable pretreatment, the solid matrix is placed in an extraction vessel forming a fixed bed. Depending on the compounds of interest, the supercritical solvent (Sc-CO₂) or solvent + cosolvent is fed by the solvent pump and/or cosolvent into the extraction vessel, where it continuously flows through the fixed bed and dissolves the extractable components from the solid matrix. The mixture of solutes that is removed from the solid matrix is called extract. In the separation step, the mixture formed by the solvent extraction + extract leaves the vessel and feeds the separator (collection flask) where the mixture is separated by rapid reduction of

pressure (ambient pressure). The extract precipitates in the separator, and the solvent is removed from the system [38, 83].

5. Conclusion

The identification of new natural antioxidant compounds is of great interest to the food, pharmaceutical, and cosmetic industry in order to find possible alternatives to synthetic antioxidants. In this way, plants such as *C. sicyoides* and *R. officinalis* have been extensively studied for their antioxidant activity. The *C. sicyoides* extract obtained by SFE has a neuroprotective and anti-inflammatory effect; these effects are associated with the presence of phenolic compounds and the high antioxidant activity in the extract. *R. officinalis* extract is antibacterial, antifungal, anti-inflammatory, and effective, associated with the presence of carnosic acid, carnosol, rosmarinic acids, and hesperetin. It has been corroborated that these plants contain chemical compounds that exhibit the capacity of FRSs and reduce the onset of different diseases. Finally, obtaining extracts from plant matrices using environmentally safe extraction technology such as SFE represents a great opportunity to obtain bioactive compounds.

Acknowledgements

Mar Salazar (Process Number: 1777277) thanks CAPES for the doctorate scholarship.

Conflict of interest

The authors have no conflict of interest to declare and are responsible for the content and writing of the manuscript.

Author details

Marielba de los Angeles Rodriguez Salazar*, Glides Rafael Olivo Urbina, Priscila do Nascimento Bezerra, Vânia Maria Borges Cunha, Marcilene Paiva da Silva, Flávia Cristina Seabra Pires, Ana Paula de Souza e Silva, Sérgio Henrique Brabo de Sousa and Raul Nunes de Carvalho Jr
LABEX/FEA (Faculty of Food Engineering), Federal University of Para, Belém, Pará, Brazil

*Address all correspondence to: marielba434@hotmail.com

IntechOpen

© 2019 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] Souza JNS, Silva EM, Loir A, Rees JF, Rogez H, Larondelle Y. Antioxidant capacity of four polyphenol-rich Amazonian plant extracts: A correlation study using chemical and biological in vitro assays. *Food Chemistry*. 2008;**106**(1):331-339. DOI: 10.1016/j.foodchem.2007.05.011
- [2] Virgolin LB, Seixas FRF, Janzantti NS. Composition, content of bioactive compounds, and antioxidant activity of fruit pulps from the Brazilian Amazon biome. *Pesquisa Agropecuaria Brasileira*. 2017;**52**(10):933-941. DOI: 10.1590/S0100-204X2017001000013
- [3] Ekor M. The growing use of herbal medicines: Issues relating to adverse reactions and challenges in monitoring safety. *Frontiers in Neurology*. 2014;**4**: 1-10. DOI: 10.3389/fphar.2013.00177
- [4] Alcantara RGL, Joaquim RHVT, Sampaio SF. Plantas medicinais: O conhecimento e uso popular. *Revista de APS*. 2015;**18**(4):470-482
- [5] Damodaran S, Parkin K, Fennema OR. *Fennema's Food Chemistry*. 4th ed. New York, Boca Raton: CRC Press/Taylor & Francis; 2008. 1144 p
- [6] Dai J, Mumper RJ. Plant phenolics: Extraction, analysis and their antioxidant and anticancer properties. *Molecules*. 2010;**15**:7313-7352. DOI: 10.3390/molecules15107313
- [7] Khoddami A, Wilkes MA, Roberts TH. Techniques for analysis of plant phenolic compounds. *Molecules*. 2013;**18**:2328-2375. DOI: 10.3390/molecules18022328
- [8] Khaw K, Parat M, Shaw PN, Falconer JR. Solvent supercritical fluid technologies to extract bioactive compounds from natural sources: A review. *Molecules*. 2017;**22**:2-22. DOI: 10.3390/molecules22071186
- [9] Halliwell B. The definition and measurement of antioxidants in biological systems. *Free Radical Biology & Medicine*. 1995;**18**(1):125-126
- [10] Droge W. Free radicals in the physiological control of cell function. *Physiological Reviews*. 2002;**82**:47-95
- [11] Sies H. Physiological society symposium: Impaired endothelial and smooth muscle cell function in oxidative stress oxidative stress: Oxidants and antioxidants. *Experimental Physiology*. 1996;**82**:291-295
- [12] Maleki D, Rad AH, Khalili L, Alipour B. Antioxidants and natural compounds. In: Gowder SJT, editor. *Basic Principles and Clinical Significance of Oxidative Stress*. London: IntechOpen; 2015. DOI: 10.5772/59293
- [13] Xu D, Li Y, Meng X, Zhou T, Zhou Y, Zheng J, et al. Natural antioxidants in foods and medicinal plants: Extraction, assessment and resources. *International Journal of Molecular Sciences*. 2017;**18**:20-32. DOI: 10.3390/ijms18010096
- [14] Halliwell B, Gutteridge JMC. The importance of free radicals and catalytic metal ions in human diseases. *Molecular Aspects of Medicine*. 1985;**8**:89-193
- [15] Thomas MJ. The role of free radicals and antioxidants. *Nutrition*. 2000;**16**:716-718
- [16] Gachkar L, Yadegari D, Rezaei MB, Taghizadeh M, Astaneh SA, Rasoli I. Chemical and biological characteristics of *Cuminum cyminum* and *Rosmarinus officinalis* essential oils. *Food Chemistry*. 2007;**102**:898-904. DOI: 10.1016/j.foodchem.2006.06.035
- [17] Genena AK, Hense H, Smania Junior A, Souza SM. Rosemary

(*Rosmarinus officinalis*)—A study of the composition, antioxidant and antimicrobial activities of extracts obtained with supercritical carbon dioxide. *Ciência e Tecnologia de Alimentos*. 2008;**28**(2):463-469

[18] Shahidi F. Antioxidants: Principles and applications. In: Shahidi F, editor. *Handbook of Antioxidants for Food Preservation*. 1st ed. United Kingdom: Elsevier Ltd; 2015. p. 514. DOI: 10.1016/B978-1-78242-089-7.00001-4

[19] Drobnik J, de Oliveira AB. *Cissus verticillata* (L.) Nicolson and C.E. Jarvis (Vitaceae): Its identification and usage in the sources from 16th to 19th century. *Journal of Ethnopharmacology*. 2015;**171**:317-329. DOI: 10.1016/j.jep.2015.06.003

[20] Pires FCS, Almeida MM, Silva APS, Salazar MAR, Urbina GRO, Silva PS. Therapeutic potential of extracts from Amazonian plants with antimicrobial activity. In: Méndez-Vilas A, editor. *Antimicrobial Research: Novel Bioknowledge and Educational Programs*. 6th ed. Badajoz: Formatex Research Center; 2017. pp. 504-515

[21] Beltrame F, Ferreira A, Cortez D. Coumarin glycoside from *Cissus sicyoides*. *Natural Product Letters*. 2002;**16**(4):213-216. DOI: 10.1080/10.575630290015736

[22] Xu F, Matsuda H, Hata H, Sugawara K, Nakamura S, Yoshikawa M. Structures of new flavonoids and benzofuran-type stilbene and degranulation inhibitors of rat basophilic leukemia cells from the Brazilian herbal medicine *Cissus sicyoides*. *Chemical and Pharmaceutical Bulletin*. 2009;**57**(10):1089-1095. DOI: 10.1248/cpb.57.1089

[23] Beserra FP, de Cássia Santos R, Périco LL, Rodrigues VP, de Almeida Kiguti LR, Saldanha LL, et al. *Cissus sicyoides*: Pharmacological mechanisms

involved in the anti-inflammatory and antidiarrheal activities. *International Journal of Molecular Sciences*. 2016;**17**(2):1-15. DOI: 10.3390/ijms17020149

[24] Salazar MAR, Costa JV, Olivo GRU, Cunha VMB, Silva MP, Bezerra PN, et al. Chemical composition, antioxidant activity, neuroprotective and anti-inflammatory effects of cipó-pucá (*Cissus sicyoides* L.) extracts obtained from supercritical extraction. *The Journal of Supercritical Fluids*. 2018;**138**:36-45. DOI: 10.1016/j.supflu.2018.03.022

[25] Machado DG, Cunha MP, Neis VB, Balen GO, Colla AR, Grando J, et al. *Rosmarinus officinalis* L. Hydroalcoholic extract, similar to fluoxetine, reverses depressive-like behavior without altering learning deficit in olfactory bulbectomized mice. *Journal of Ethnopharmacology*. 2012;**143**:158-169. DOI: 10.1016/j.jep.2012.06.017

[26] Hannour K, Boughdad A, Maataoui A, Bouchelta A. Chemical composition of *Rosmarinus officinalis* (Lamiaceae) essential oils and evaluation of their toxicity against *Bruchus rufimanus* (Coleoptera: Chrysomelidae - Bruchinae) in Morocco. *International Journal of Tropical Insect Sciences*. 2018;**38**:192-204. DOI: 10.1017/S1742758418000012

[27] Cuvelier M, Richard H, Berset C. Antioxidative activity and phenolic composition of pilot-plant and commercial extracts of sage and rosemary. *Journal of the American Oil Chemists' Society*. 1996;**73**:645-652

[28] Haloui M, Louedec L, Michel J, Lyoussi B. Experimental diuretic effects of *Rosmarinus officinalis* and *Centaureum erythraea*. *Journal of Ethnopharmacology*. 2000;**71**:465-472

[29] Zheng W, Wang SY. Antioxidant activity and phenolic compounds in

- p>selected herbs antioxidant activity and phenolic compounds in selected herbs.
- Journal of Agricultura and Food Chemistry*
- . 2001;
- 49**
- (11):5165-5170. DOI: 10.1021/jf010697n
- [30] Moreno S, Scheyer T, Romano CS, Vojnov AA. Antioxidant and antimicrobial activities of rosemary extracts linked to their polyphenol composition. *Free Radical Research*. 2006;**40**:223-231. DOI: 10.1080/10715760500473834
- [31] Kontogianni VG, Tomic G, Nikolic I, Nerantzaki AA, Sayyad N, Stosic-grujicic S, et al. Phytochemical profile of *Rosmarinus officinalis* and *Salvia officinalis* extracts and correlation to their antioxidant and anti-proliferative activity. *Food Chemistry*. 2013;**136**(1):120-129. DOI: 10.1016/j.foodchem.2012.07.091
- [32] Habtemariam S. The therapeutic potential of rosemary (*Rosmarinus officinalis*) diterpenes for alzheimer's disease. *Evidence-Based Complementary and Alternative Medicine*. 2016;**2016**:1-14. DOI: 10.1155/2016/2680409
- [33] Carvalho RN Jr, Moura LS, Rosa PTV, Meireles MAA. Supercritical fluid extraction from rosemary (*Rosmarinus officinalis*): Kinetic data, extract's global yield, composition, and antioxidant activity. *The Journal of Supercritical Fluids*. 2005;**35**(3):197-204. DOI: 10.1016/j.supflu.2005.01.009
- [34] Brunner G. Supercritical fluids: Technology and application to food processing. *Journal Food Engineering*. 2005;**67**(1-2):21-33. DOI: 10.1016/j.jfoodeng.2004.05.060
- [35] Pereira CG, Meireles MAA. Supercritical fluid extraction of bioactive compounds: Fundamentals, applications and economic perspectives. *Food and Bioprocess Technology*. 2010;**3**(3):340-372. DOI: 10.1007/s11947-009-0263-2
- [36] Herrero M, Castro-Puyana M, Mendiola JA, Ibañez E. Compressed fluids for the extraction of bioactive compounds. *Trends in Analytical Chemistry*. 2013;**43**(2):67-83. DOI: 10.1016/j.trac.2012.12.008
- [37] Goyeneche R, Fanovich A, Rodrigues CR, Nicolao MC, Discala K. Supercritical CO2 extraction of bioactive compounds from radish leaves: Yield, antioxidant capacity and cytotoxicity. *The Journal of Supercritical Fluids*. 2018;**135**:78-83. DOI: 10.1016/j.supflu.2018.01.004
- [38] de Melo MMR, Silvestre AJD, Silva CM. Supercritical fluid extraction of vegetable matrices: Applications, trends and future perspectives of a convincing green technology. *The Journal of Supercritical Fluids*. 2014;**92**:115-176. DOI: 10.1016/j.supflu.2014.04.007
- [39] da Silva GA, de Almeida-Muradian LB, Akisue G, Ferro VO. Padronização do extrato de *Cissus sicyoides* L. (insulina vegetal) e identificação de carotenos. *Revista Brasileira de Farmacognosia*. 1996;**1**:96-112
- [40] Quílez AM, Saenz MT, García MD, de la Puerta R. Phytochemical analysis and anti-allergic study of *Agave intermixta* Trel. and *Cissus sicyoides* L. *Journal Pharmacy Pharmacology*. 2004;**56**(9):1185-1189. DOI: 10.1211/0022357044102
- [41] de Oliveira AB, de Mendonça MS, Azevedo AA, Meira RMSA. Anatomy and histochemistry of the vegetative organs of *Cissus verticillata*—A native medicinal plant of the Brazilian Amazon. *Brazilian Journal of Pharmacognosy*. 2012;**22**(6):1201-1211. DOI: 10.1590/S0102-695X2012005000092
- [42] Toledo MCF, Reyes FGR, Iaderoza M, Francis FJ, Draetta S. Anthocyanins from anil trepador (*Cissus sicyoides*, Linn.). *Journal of Food Science*. 1983;**48**:1368

- [43] El-shafey MM, Abd-allah GM, Mohamadin AM, Harisa GI, Mariee AD. Quercetin protects against acetaminophen-induced hepatorenal toxicity by reducing reactive oxygen and nitrogen species. *Pathophysiology*. 2015;**22**(1):49-55. DOI: 10.1016/j.pathophys.2014.12.002
- [44] Crespo I, García-Mediavilla MV, Almar M, González P, Tunón MJ, Sánchez-Campos S, et al. Differential effects of dietary flavonoids on reactive oxygen and nitrogen species generation and changes in antioxidant enzyme expression induced by proinflammatory cytokines in Chang liver cells. *Food and Chemical Toxicology*. 2008;**46**:1555-1569. DOI: 10.1016/j.fct.2007.12.014
- [45] Perrone D, Fuggetta MP, Ardito F, Cottarelli A, Filippis ADE, Ravagnan G, et al. Resveratrol (3,5,4'-trihydroxystilbene) and its properties in oral diseases (review). *Experimental and Therapeutic Medicine*. 2017;**14**:3-9. DOI: 10.3892/etm.2017.4472
- [46] Bolkenius FN, Verne-misner J, Wagner J, Grisar JM. Amphiphilic cY-tocopherol analogues as inhibitors of brain lipid peroxidation. *European Journal of Pharmacology*. 1996;**298**:37-43
- [47] Khalil NM, Pepato MT, Brunetti IL. Free radical scavenging profile and myeloperoxidase inhibition of extracts from antidiabetic plants: *Bauhinia forficata* and *Cissus sicyoides*. *Biological Research*. 2008;**41**(2):165-171. DOI: 10.4067/S0716-97602008000200006
- [48] Pepato MT, Baviera AM, Vendramini RC, da Silva Perez MDP, Kettelhut ID. *Cissus sicyoides* (princess vine) in the long-term treatment of streptozotocin-diabetic rats. *Biotechnology and Applied Biochemistry*. 2003;**37**(Part 1):15-20. DOI: 10.1042/BA20020065
- [49] Viana GSB, Medeiros ACC, Lacerda AMR, Leal LKAM, Vale TG, Matos FJDA. Hypoglycemic and anti-lipemic effects of the aqueous extract from *Cissus sicyoides*. *BMC Pharmacology*. 2004;**4**:1-7. DOI: 10.1186/1471-2210-4-9
- [50] Fernandes G, Banu J. Medicinal properties of plants from the genus *Cissus*: A review. *Journal of Medicinal Plants Research*. 2012;**6**(16):3080-3086. DOI: 10.5897/JMPR11.1637
- [51] Sáenz MT, García MD, Quílez A, Ahumada MC. Cytotoxic activity of *Agave intermixta* L. (Agavaceae) and *Cissus sicyoides* L. (Vitaceae). *Phytotherapy Research*. 2000;**14**:552-554
- [52] Lucena FRS, Almeida ER, Aguiar JS, Silva TG, Souza VMO, Nascimento SC. Cytotoxic, antitumor and leukocyte migration activities of resveratrol and sitosterol present in the hidroalcoholic extract of *Cissus sicyoides* L., Vitaceae, leaves. *Brazilian Journal of Pharmacognosy*. 2010;**20**(5):729-733. DOI: 10.1590/S0102-695X2010005000002
- [53] García MD, Quílez AM, Sáenz MT, Martínez-Domínguez ME, De La Puerta R. Anti-inflammatory activity of *Agave intermixta* Trel. and *Cissus sicyoides* L., species used in the caribbean traditional medicine. *Journal of Ethnopharmacology*. 2000;**71**(3):395-400. DOI: 10.1016/S0378-8741(00)00160-4
- [54] García X, Cartas-Heredia L, Lorenzana-Jímenez M, Gijón E. Vasoconstrictor effect of *Cissus sicyoides* on Guinea-pig aortic rings. *General Pharmacology*. 1997;**29**(3):457-462. DOI: 10.1016/S0306-3623(96)00478-8
- [55] Santos HB, Modesto-filho J, Diniz M de FF, Vasconcelos THC, Pereira FS de B, Ramalho J do A, et al. Artigo avaliação do efeito hipoglicemiante de *Cissus sicyoides* em estudos clínicos fase

II. Brazilian Journal of Pharmacognosy. 2008;**18**(1):70-76

[56] Garcia M, Saenz MT, Puerta R, Quilez A, Fernandez MA. Antibacterial activity of *Agave intermixta* and *Cissus sicyoides*. Fitoterapia. 1999;**70**:71-73. DOI: 10.1016/S0367-326X(98)00009-4

[57] Silva L, Carreira RC, Oniki GH, Agripino DG, Young MCM, Ladeira AM. Crescimento e análise do potencial antifúngico em plantas de *Cissus verticillata* (L.) Nicolson & Jarvis (Vitaceae). Revista Brasileira de Plantas Mediciniais. 2007;**9**(1):73-79

[58] Brender T, Gruenwald J, Jaenicke C. Herbal Remedies. 2nd ed. Berlin: Schaper & Brümmer GmbH & Co., Salzgitter; 1997

[59] Rodríguez EA, Árias AJ, Vásquez EG, Martínez JR, Stashenko EE. Rendimiento y capacidad antioxidante de extractos de *Rosmarinus officinalis*, *Salvia officinalis* y *Psidium guajava* obtenidos con CO₂ supercrítico. Revista de la Academia Colombiana de Universidad del Zulia. 2012;**XXXVI**:305-315

[60] Govaris A, Florou-paneri P, Botsoglou E, Giannenas I, Amvrosiadis I, Botsoglou N. The inhibitory potential of feed supplementation with rosemary and/or α -tocopheryl acetate on microbial growth and lipid oxidation of Turkey breast during refrigerated storage. Lebensmittel-Wissenschaft & Technologie. 2007;**40**:331-337. DOI: 10.1016/j.lwt.2005.10.006

[61] Bozin B, Mimica-Dukic N, Samojlik I, Jovin E. Antimicrobial and antioxidant properties of rosemary and sage (*Rosmarinus officinalis* L. and *Salvia officinalis* L., Lamiaceae) essential oils. Journal of Agricultura and Food Chemistry. 2007;**55**:7879-7885. DOI: 10.1021/jf0715323

[62] Tai J, Cheung S, Wu M, Hasman D. Antiproliferation effect of rosemary (*Rosmarinus officinalis*) on human ovarian cancer cells in vitro. Phytomedicine. 2012;**19**(5):436-443. DOI: 10.1016/j.phymed.2011.12.012

[63] Munne-Bosch S, Alegre L. Subcellular compartmentation of the diterpene carnosic acid and its derivatives in the leaves of rosemary. Plant Physiology. 2001;**125**(2):1094-1102. DOI: 10.1104/pp.125.2.1094

[64] Wijeratne SSK, Cuppett SL. Potential of rosemary (*Rosmarinus officinalis* L.) diterpenes in preventing lipid hydroperoxide-mediated oxidative stress in Caco-2 cells. Journal of Agricultural and Food Chemistry. 2007;**55**(4):1193-1199. DOI: 10.1021/jf063089m

[65] Cui L, Kim MO, Seo JH, Kim IS, Kim NY, Lee SH, et al. Abietane diterpenoids of *Rosmarinus officinalis* and their diacylglycerol acyltransferase-inhibitory activity. Food Chemistry. 2012;**132**(4):1775-1780. DOI: 10.1016/j.foodchem.2011.11.138

[66] Hras AR, Hadolin M, Knez Z, Bauman D. Comparison of antioxidative and synergistic effects of rosemary extract with α -tocopherol, ascorbyl palmitate and citric acid in sunflower oil. Food Chemistry. 2000;**71**:229-233

[67] Fang X, Wada S. Enhancing the antioxidant effect of α -tocopherol with rosemary in inhibiting catalyzed oxidation caused by Fe²⁺ and hemoprotein. Food Research International. 1993;**26**(6):405-411. DOI: 10.1016/0963-9969(93)90086-X

[68] Nakatani N, Inatani R. Two antioxidative diterpenes from rosemary (*Rosmarinus officinalis* L.) and a revised structure for rosmannol. Agricultural and Biological Chemistry. 1984;**48**:2081-2085

- [69] Houlihan CM, Ho CT, Chang SS. The structure of rosmariquinone—A new antioxidant isolated from *Rosmarinus officinalis* L. Journal of the American Oil Chemists' Society. 1985;**62**(1):96-98. DOI: 10.1007/BF02541500
- [70] Tohma S, Turan S. Rosemary plant (*Rosmarinus officinalis* L.), solvent extract and essential oil can be used to extend the usage life of hazelnut oil during deep frying. European Journal of Lipid Science and Technology. 2015;**117**:1978-1990. DOI: 10.1002/ejlt.201400382
- [71] Nieto G, Ros G, Castillo J. Antioxidant and antimicrobial properties of rosemary (*Rosmarinus officinalis* L.): A review. Medicine. 2018;**5**(3):98. DOI: 10.3390/medicines5030098
- [72] Kayashima TK, Matsubara KM. Antiangiogenic effect of carnosic acid and carnosol, neuroprotective compounds in rosemary leaves. Bioscience Biotechnology Biochemistry. 2012;**76**(1):115-119. DOI: 10.1271/bbb.110584
- [73] Al-attar AM, Shawush NA. Influence of olive and rosemary leaves extracts on chemically induced liver cirrhosis in male rats. Saudi Journal of Biological Sciences. 2015;**22**(2):157-163. DOI: 10.1016/j.sjbs.2014.08.005
- [74] Nieto G. Biological activities of three essential oils of the Lamiaceae family. Medicine. 2017;**4**(63):1-10. DOI: 10.3390/medicines4030063
- [75] Bakirel T, Bakirel U, Keleş OÜ, Ülgen SG, Yardibi H. In vivo assessment of antidiabetic and antioxidant activities of rosemary (*Rosmarinus officinalis*) in alloxan-diabetic rabbits. Journal of Ethnopharmacology. 2008;**116**(1):64-73. DOI: 10.1016/j.jep.2007.10.039
- [76] Belmouhoub M, Chebout I, Iguer-ouada M. Antidiabetic and anti-hypercholesterolemic effects of flavonoid-rich fractions of *Rosmarinus officinalis* in streptozotocin-induced diabetes in mice. Phytothérapie. 2017;1-7. DOI: 10.1007/s10298-017-1103-6
- [77] Arranz E, Mes J, Wichers HJ, Jaime L, Mendiola JA, Reglero G, et al. Anti-inflammatory activity of the basolateral fraction of Caco-2 cells exposed to a rosemary supercritical extract. Journal of Functional Foods. 2015;**13**:384-390. DOI: 10.1016/j.jff.2015.01.015
- [78] Garcia-Salas P, Morales-Soto A, Segura-Carretero A, Fernández-Gutiérrez A. Phenolic-compound-extraction systems for fruit and vegetable samples. Molecules. 2010;**15**(12):8813-8826. DOI: 10.3390/molecules15128813
- [79] Jiao G, Pour KA. Extraction of anthocyanins from haskap berry pulp using supercritical carbon dioxide: Influence of co-solvent composition and pretreatment. LWT—Food Science and Technology. 2018;**98**:237-244. DOI: 10.1016/j.lwt.2018.08.042
- [80] Moura PM, Prado GHC, Meireles MAA, Pereira CG. Supercritical fluid extraction from guava (*Psidium guajava*) leaves: Global yield, composition and kinetic data. The Journal of Supercritical Fluids. 2012;**62**:116-122. DOI: 10.1016/j.supflu.2011.11.014
- [81] Solana M, Boschiero I, Acqua SD, Bertucco A. Extraction of bioactive enriched fractions from *Eruca sativa* leaves by supercritical CO₂ technology using different co-solvents. The Journal of Supercritical Fluids. 2014;**94**:245-251. DOI: 10.1016/j.supflu.2014.08.022
- [82] da Silva RFF, Rocha-santos TAP, Duarte AC. Supercritical fluid extraction of bioactive compounds. Trends in Analytical Chemistry. 2016;**76**:40-51. DOI: 10.1016/j.trac.2015.11.013

[83] Jesus SP, Meireles MAA. Supercritical fluid extraction: A global perspective of the fundamental concepts of this eco-friendly extraction technique. In: Chemat F, Vian MA, editors. *Alternative Solvents for Natural Products Extraction*. 1st ed. London: Springer-Verlag Berlin Heidelberg; 2014. pp. 39-72. DOI: 10.1007/978-3-662-43628-8

[84] Babovic N, Djilas S, Jadranin M, Vajs V, Ivanovic J, Petrovic S, Zizovic I. Supercritical carbon dioxide extraction of antioxidant fractions from selected Lamiaceae herbs and their antioxidant capacity. *Innovative Food Science and Emerging Technologies*. 2010;**11**(1):98-107. DOI: 10.1016/j.ifset.2009.08.013

[85] Fernández-ponce MT, Casas L, Mantell C, Rodríguez M, De la Ossa EM. Extraction of antioxidant compounds from different varieties of *Mangifera indica* leaves using green technologies. *The Journal of Supercritical Fluids*. 2012;**72**:168-175. DOI: 10.1016/j.supflu.2012.07.016

[86] Martinez-Correa HA, Magalhães PM, Queiroga CL, Peixoto CA, Oliveira AL, Cabral FA. Extracts from pitanga (*Eugenia uniflora* L.) leaves: Influence of extraction process on antioxidant properties and yield of phenolic compounds. *The Journal of Supercritical Fluids*. 2011;**55**(3):998-1006. DOI: 10.1016/j.supflu.2010.09.001

[87] Bagheri H, Abdul Manap MY Bin, Solati Z. Antioxidant activity of *Piper nigrum* L. essential oil extracted by supercritical CO₂ extraction and hydro-distillation. *Talanta*. 2014;**121**:220-228. DOI: 10.1016/j.talanta.2014.01.007